

Adolescent Behaviors and Experiences Survey								
Alcohol and Other Drug Use	Grade 9		Grade 10		Grade 11		Grade 12	
	Percentage	Confidence Interval						
Had their first drink of alcohol before age 13 years (other than a few sips)	14.3	11.9 - 17.1	10.0	8.1 - 12.3	10.8	8.6 - 13.5	8.9	7.2 - 11.0
Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	12.9	9.8 - 16.6	17.1	14.0 - 20.8	21.7	18.2 - 25.7	27.2	23.8 - 30.9
Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey)	4.6	2.5 - 8.2	5.3	3.5 - 7.8	9.6	7.5 - 12.1	12.1	9.5 - 15.2
Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	1.1	0.5 - 2.2	1.5	0.9 - 2.7	2.7	1.7 - 4.4	3.6	2.2 - 5.8
Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	32.5	24.3 - 42.0	40.0	34.0 - 46.2	40.7	34.2 - 47.6	38.7	33.0 - 44.8
Ever used marijuana (one or more times during their life)	15.9	12.5 - 19.9	20.8	18.0 - 24.0	28.6	24.8 - 32.8	36.0	33.6 - 38.3
Tried marijuana for the first time before age 13 years	4.5	3.2 - 6.3	3.5	2.3 - 5.2	3.1	2.1 - 4.4	2.6	1.7 - 3.8
Currently used marijuana (one or more times during the 30 days before the survey)	9.4	7.0 - 12.7	10.5	8.2 - 13.2	13.4	11.2 - 15.9	18.4	15.8 - 21.4
Ever used synthetic marijuana (one or more times during their life)	5.0	3.5 - 7.0	5.5	4.2 - 7.1	6.4	5.0 - 8.3	6.3	4.7 - 8.4
Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	10.3	8.5 - 12.4	9.6	7.6 - 12.0	9.7	7.9 - 11.8	8.6	7.1 - 10.4
Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)	5.3	4.2 - 6.7	4.2	3.2 - 5.7	4.0	2.8 - 5.7	3.6	2.6 - 4.9
Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	1.0	0.6 - 1.8	1.3	0.6 - 2.8	1.4	0.8 - 2.3	2.1	1.3 - 3.4
Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	7.9	5.9 - 10.5	5.7	4.4 - 7.5	5.9	4.3 - 7.9	3.2	2.2 - 4.6
Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.7	0.3 - 1.4	0.8	0.3 - 2.2	0.5	0.2 - 0.9	0.7	0.4 - 1.3
Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	1.1	0.6 - 2.0	0.7	0.3 - 1.4	0.6	0.3 - 1.4	1.0	0.6 - 1.7
Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)	1.5	0.9 - 2.6	2.0	1.2 - 3.3	1.5	0.9 - 2.5	2.8	1.8 - 4.4
Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)	2.3	1.4 - 3.8	2.9	1.9 - 4.4	3.5	2.4 - 4.9	6.6	4.6 - 9.2
Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times during their life)	9.8	7.7 - 12.5	8.4	6.4 - 11.0	8.9	6.9 - 11.3	9.3	7.1 - 12.1
Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	0.6	0.2 - 1.5	0.6	0.2 - 1.5	0.1	0.1 - 0.3	0.8	0.4 - 1.5
Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended school in-person during the 12 months before the survey)	9.7	7.0 - 13.3	10.1	7.0 - 14.4	8.6	6.2 - 11.9	8.9	6.8 - 11.5