

Adolescent Behaviors and Experiences Survey

	Grade 9		Grade 10		Grade 11		Grade 12	
Mental Health	Percentage	Confidence Interval						
Who felt sad or hopeless (almost every day for two or more weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	43.3	38.6 - 48.1	44.6	40.8 - 48.4	44.7	40.6 - 48.8	44.4	40.7 - 48.2
Seriously considered attempting suicide (during the 12 months before the survey)	22.1	18.3 - 26.4	19.6	17.0 - 22.5	18.5	15.9 - 21.3	19.2	16.3 - 22.6
Made a plan about how they would attempt suicide (during the 12 months before the survey)	15.6	12.7 - 18.9	15.9	13.7 - 18.3	15.0	12.2 - 18.3	14.8	11.4 - 18.9
Actually attempted suicide (one or more times during the 12 months before the survey)	10.3	7.8 - 13.6	9.9	8.1 - 12.2	8.4	6.4 - 10.9	7.2	4.9 - 10.4
Had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.0	1.2 - 3.3	1.7	1.0 - 2.7	2.0	1.2 - 3.3	1.8	1.1 - 3.0
Reported that their mental health was most of the time or always not good (including stress, anxiety, and depression, during the 30 days before the survey)	30.2	26.7 - 33.9	31.3	27.8 - 35.0	30.2	26.7 - 34.0	32.8	29.0 - 36.8