

Adolescent Behaviors and Experiences Survey								
Sexual Behaviors	Grade 9		Grade 10		Grade 11		Grade 12	
	Percentage	Confidence Interval						
Ever had sexual intercourse	13.6	10.8 - 16.9	21.9	18.1 - 26.3	33.2	29.1 - 37.6	47.0	43.4 - 50.6
Had sexual intercourse for the first time before age 13 years	2.2	1.4 - 3.5	2.0	1.3 - 3.1	1.9	1.1 - 3.0	2.4	1.5 - 3.9
Had sexual intercourse with four or more persons during their life	1.9	1.3 - 3.0	4.1	3.0 - 5.6	5.0	3.5 - 7.0	12.3	9.4 - 15.9
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	7.3	5.2 - 10.1	13.8	10.7 - 17.6	21.6	18.3 - 25.3	33.6	30.1 - 37.3
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	29.3	20.7 - 39.5	12.3	8.9 - 16.6	17.3	11.6 - 25.1	14.9	11.3 - 19.4
Used a condom during last sexual intercourse (among students who were currently sexually active)	50.9	39.7 - 62.0	57.8	50.6 - 64.7	56.9	50.1 - 63.4	51.5	46.3 - 56.7
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	8.6	4.8 - 15.1	15.8	11.1 - 22.0	25.2	19.5 - 32.0	25.4	20.5 - 30.9
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	4.0	3.0 - 5.4	4.7	3.3 - 6.8	5.7	4.3 - 7.5	8.9	7.3 - 10.8
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	1.9	1.2 - 3.0	3.8	2.8 - 5.2	5.1	3.9 - 6.6	8.0	6.2 - 10.3
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	1.4	0.2 - 8.6	6.3	3.4 - 11.4	5.7	3.1 - 10.3	6.8	4.9 - 9.4
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	15.4	9.4 - 24.3	27.8	21.0 - 35.8	38.3	31.3 - 45.7	37.4	32.2 - 42.8
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	6.3	3.1 - 12.3	9.3	6.4 - 13.5	17.4	12.7 - 23.4	10.6	7.9 - 13.9
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	17.9	10.5 - 28.7	17.8	12.1 - 25.3	13.2	9.3 - 18.3	10.8	7.6 - 15.2