Centers for Disease Control and Prevention National Center for Emerging and Zoonotic Infectious Diseases



Infection Prevention and Control (IPC) for Marburg Virus Disease (MVD):

Healthcare Worker and Inpatient Monitoring

Healthcare Settings with Limited to Intermediate Resources

Updated: March 2023

Learning Objectives

After this presentation, participants will be able to

- Explain why healthcare worker monitoring is important in the context of MVD.
- Explain why inpatient monitoring is important in the context of MVD.
- Describe steps for monitoring inpatients.

Screening for MVD involves 2 parts. What are they?

_____ and _____

Screening for MVD involves 2 parts:

Temperature check and questionnaire

Why Monitor Healthcare Workers?

- Healthcare workers are at high risk for exposure to MVD.
- If they are ill, they can infect co-workers or patients.
- Screening healthcare workers regularly helps quickly identify if they are ill, provide early care, and isolate them from others.
- This helps protect you, your patients and co-workers, and your community.

Healthcare Worker Monitoring

- Screen all healthcare workers prior to each shift.
 - Temperature check
 - Questionnaire: signs and symptoms + risk factors in past 21 days
- Encourage reporting symptoms don't penalize
 - Sick healthcare workers should be evaluated for possible Marburg virus disease
- Maintain a record of healthcare workers entering isolation areas because of high-risk for potential exposure

Why Monitor Inpatients?

The incubation period for MVD is between 2-21 days.

- Some patients may be admitted for other reasons and may not have signs or symptoms of MVD upon admission.
- They may develop signs and symptoms during their stay in your facility.
- Quickly identifying them as potentially having MVD and isolating them from others helps protect you, your patients and co-workers, and your community.

Inpatient Monitoring Process

- Apply standard precautions for all patients.
- Take temperature at least twice a day for all inpatients
- Assess patients for signs and symptoms at least once a day.
 Assess immediately in the presence of fever (> 38°C).
- Complete the Inpatient Monitoring Form.



Inpatient Monitoring Form

- Temperature 2 times/day
- Signs and symptoms 1 time/day
- Optional: Exposures in the last 21 days
 - This may be asked to increase recall of patient, if feasible

| Date | Name of patient | Age in month / <u>year</u> | Sex | Temp. 1 | Temp. 2 | New symptoms (including temperature) | Other (new contact information) |
|------|-----------------|----------------------------------|-----|---------|------------|--|---------------------------------|
| | | | | | | | |
| | | | | | | | |
| | | | | | • | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

Knowledge Check: Monitoring

You notice that a healthcare worker monitoring patients in the maternity ward of the hospital is walking around to take temperatures of patients and ask about symptoms. They record the temperature but do not record symptoms.

When you ask about a particular patient's symptoms the healthcare worker responds: "The patient has felt nauseous and had some diarrhea and a headache. The patient has had a fever since yesterday, but she does not show signs of MVD because she is not vomiting and does not look hemorrhagic."

- 1. What is the risk of not keeping track of symptoms?
- 2. What is the correct process for in-patient monitoring?

Feedback: Monitoring

1. What is the risk of not keeping track of symptoms?

- If symptoms not recorded, it can be difficult to track any worsening symptoms
- Change in nursing staff between shifts means information might not be transferred between shifts

2. What is the correct process for in-patient monitoring?

- Health workers should complete in-patient identification form COMPLETELY for each patient and not skip over any steps
- Complete in-patient identification as agreed upon in country or facility policy
- If a patient meets the case definition for MVD, isolate them immediately by separating them from other patients.

Key Takeaways

 Monitoring healthcare workers and inpatients helps identify and isolate people who potentially have MVD from others in the facility. This helps protect you, your patients and co-workers, and your community.

When monitoring inpatients, always fill out inpatient monitoring forms
 completely and as agreed upon in your facility to ensure that patients
 who become ill while in your facility can be identified and isolated
 quickly.

Thank you!

For more information, contact CDC 1-800-CDC-INFO (232-4636)
TTY: 1-888-232-6348 www.cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

