

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,423,712	100.0	803.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	616,067	25.4	204.3
2	Malignant neoplasms (C00-C97)	562,875	23.2	186.6
3	Cerebrovascular diseases (I60-I69)	135,952	5.6	45.1
4	Chronic lower respiratory diseases (J40-J47)	127,924	5.3	42.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123,706	5.1	41.0
6	Alzheimer's disease (G30)	74,632	3.1	24.7
7	Diabetes mellitus (E10-E14)	71,382	2.9	23.7
8	Influenza and pneumonia (J09-J18)	52,717	2.2	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46,448	1.9	15.4
10	Septicemia (A40-A41)	34,828	1.4	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,598	1.4	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,165	1.2	9.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,965	1.0	7.9
14	Parkinson's disease (G20-G21)	20,058	0.8	6.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,361	0.8	6.1
...	All other causes (Residual)	451,034	18.6	149.5
All races, both sexes, 1-4 years				
...	All causes	4,703	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,588	33.8	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	546	11.6	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	398	8.5	2.4
4	Malignant neoplasms (C00-C97)	364	7.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	3.7	1.1
6	Influenza and pneumonia (J09-J18)	109	2.3	0.7
7	Septicemia (A40-A41)	78	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	70	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	57	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	52	1.1	0.3
12	Complications of medical and surgical care (Y40-Y84,Y88)	23	0.5	0.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.1
14	Anemias (D50-D64)	17	0.4	*
15	Meningitis (G00,G03)	16	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	16	0.3	*
...	All other causes (Residual)	1,115	23.7	6.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,147	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,194	35.7	5.5
2	Malignant neoplasms (C00-C97)	959	15.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	374	6.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	346	5.6	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	3.9	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	3.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	118	1.9	0.3
8	Influenza and pneumonia (J09-J18)	103	1.7	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	1.4	0.2
10	Cerebrovascular diseases (I60-I69)	83	1.4	0.2
11	Septicemia (A40-A41)	74	1.2	0.2
12	Anemias (D50-D64)	32	0.5	0.1
13	Meningitis (G00,G03)	25	0.4	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.4	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	22	0.4	0.1
...	All other causes (Residual)	1,284	20.9	3.2
All races, both sexes, 15-24 years				
...	All causes	33,982	100.0	79.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,897	46.8	37.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,551	16.3	13.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,140	12.2	9.7
4	Malignant neoplasms (C00-C97)	1,653	4.9	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,084	3.2	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	402	1.2	0.9
7	Cerebrovascular diseases (I60-I69)	195	0.6	0.5
8	Diabetes mellitus (E10-E14)	168	0.5	0.4
9	Influenza and pneumonia (J09-J18)	163	0.5	0.4
10	Septicemia (A40-A41)	160	0.5	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	160	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	160	0.5	0.4
13	Chronic lower respiratory diseases (J40-J47)	149	0.4	0.4
14	Legal intervention (Y35,Y89.0)	97	0.3	0.2
15	Anemias (D50-D64)	90	0.3	0.2
...	All other causes (Residual)	3,913	11.5	9.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	42,572	100.0	104.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,977	35.2	36.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,278	12.4	13.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,758	11.2	11.7
4	Malignant neoplasms (C00-C97)	3,463	8.1	8.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,223	7.6	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,091	2.6	2.7
7	Diabetes mellitus (E10-E14)	610	1.4	1.5
8	Cerebrovascular diseases (I60-I69)	505	1.2	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	417	1.0	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	384	0.9	0.9
11	Influenza and pneumonia (J09-J18)	331	0.8	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	326	0.8	0.8
13	Septicemia (A40-A41)	297	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	263	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	261	0.6	0.6
...	All other causes (Residual)	6,388	15.0	15.7
All races, both sexes, 35-44 years				
...	All causes	79,606	100.0	184.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,931	21.3	39.2
2	Malignant neoplasms (C00-C97)	13,288	16.7	30.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,839	14.9	27.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,722	8.4	15.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,572	4.5	8.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,052	3.8	7.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,570	3.2	6.0
8	Cerebrovascular diseases (I60-I69)	2,133	2.7	4.9
9	Diabetes mellitus (E10-E14)	1,984	2.5	4.6
10	Septicemia (A40-A41)	910	1.1	2.1
11	Chronic lower respiratory diseases (J40-J47)	796	1.0	1.8
12	Influenza and pneumonia (J09-J18)	784	1.0	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	754	0.9	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	513	0.6	1.2
15	Viral hepatitis (B15-B19)	489	0.6	1.1
...	All other causes (Residual)	13,269	16.7	30.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	184,686	100.0	420.9
1	Malignant neoplasms (C00-C97)	50,167	27.2	114.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,434	20.3	85.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,315	11.0	46.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,212	4.4	18.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,778	4.2	17.7
6	Cerebrovascular diseases (I60-I69)	6,385	3.5	14.6
7	Diabetes mellitus (E10-E14)	5,753	3.1	13.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,156	2.3	9.5
9	Chronic lower respiratory diseases (J40-J47)	4,153	2.2	9.5
10	Viral hepatitis (B15-B19)	2,815	1.5	6.4
11	Septicemia (A40-A41)	2,431	1.3	5.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,233	1.2	5.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,140	1.2	4.9
14	Influenza and pneumonia (J09-J18)	1,909	1.0	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,235	0.7	2.8
...	All other causes (Residual)	27,570	14.9	62.8
All races, both sexes, 55-64 years				
...	All causes	287,110	100.0	877.7
1	Malignant neoplasms (C00-C97)	103,171	35.9	315.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,527	22.8	200.3
3	Chronic lower respiratory diseases (J40-J47)	12,777	4.5	39.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,193	4.2	37.3
5	Diabetes mellitus (E10-E14)	11,304	3.9	34.6
6	Cerebrovascular diseases (I60-I69)	10,500	3.7	32.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	8,004	2.8	24.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,069	1.8	15.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,440	1.5	13.6
10	Septicemia (A40-A41)	4,231	1.5	12.9
11	Influenza and pneumonia (J09-J18)	3,152	1.1	9.6
12	Viral hepatitis (B15-B19)	2,413	0.8	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,124	0.7	6.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,721	0.6	5.3
15	Aortic aneurysm and dissection (I71)	1,483	0.5	4.5
...	All other causes (Residual)	39,001	13.6	119.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	389,238	100.0	2,011.3
1	Malignant neoplasms (C00-C97)	138,466	35.6	715.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,589	23.0	462.9
3	Chronic lower respiratory diseases (J40-J47)	28,664	7.4	148.1
4	Cerebrovascular diseases (I60-I69)	18,007	4.6	93.0
5	Diabetes mellitus (E10-E14)	15,112	3.9	78.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,753	2.2	45.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,752	2.0	40.1
8	Septicemia (A40-A41)	6,345	1.6	32.8
9	Influenza and pneumonia (J09-J18)	5,547	1.4	28.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,167	1.3	26.7
11	Alzheimer's disease (G30)	3,984	1.0	20.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,133	0.8	16.2
13	Aortic aneurysm and dissection (I71)	2,616	0.7	13.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,444	0.6	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,402	0.6	12.4
...	All other causes (Residual)	51,257	13.2	264.9
All races, both sexes, 75-84 years				
...	All causes	652,682	100.0	5,011.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	171,257	26.2	1,315.0
2	Malignant neoplasms (C00-C97)	163,608	25.1	1,256.3
3	Chronic lower respiratory diseases (J40-J47)	48,041	7.4	368.9
4	Cerebrovascular diseases (I60-I69)	41,979	6.4	322.3
5	Alzheimer's disease (G30)	23,009	3.5	176.7
6	Diabetes mellitus (E10-E14)	21,189	3.2	162.7
7	Influenza and pneumonia (J09-J18)	14,859	2.3	114.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,711	2.3	113.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,736	2.1	105.5
10	Septicemia (A40-A41)	10,403	1.6	79.9
11	Parkinson's disease (G20-G21)	9,363	1.4	71.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,442	1.0	49.5
13	Pneumonitis due to solids and liquids (J69)	5,187	0.8	39.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,717	0.7	36.2
15	Aortic aneurysm and dissection (I71)	4,346	0.7	33.4
...	All other causes (Residual)	99,835	15.3	766.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	713,647	100.0	12,946.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235,249	33.0	4,267.7
2	Malignant neoplasms (C00-C97)	87,656	12.3	1,590.2
3	Cerebrovascular diseases (I60-I69)	55,975	7.8	1,015.5
4	Alzheimer's disease (G30)	46,804	6.6	849.1
5	Chronic lower respiratory diseases (J40-J47)	32,857	4.6	596.1
6	Influenza and pneumonia (J09-J18)	25,535	3.6	463.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,021	2.2	290.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,803	2.2	286.7
9	Diabetes mellitus (E10-E14)	15,227	2.1	276.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,536	1.5	191.1
11	Septicemia (A40-A41)	9,614	1.3	174.4
12	Pneumonitis due to solids and liquids (J69)	8,451	1.2	153.3
13	Parkinson's disease (G20-G21)	7,911	1.1	143.5
14	Atherosclerosis (I70)	4,590	0.6	83.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,209	0.6	76.4
...	All other causes (Residual)	137,209	19.2	2,489.1
All races, male, all ages				
...	All causes	1,203,968	100.0	809.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	309,821	25.7	208.4
2	Malignant neoplasms (C00-C97)	292,857	24.3	197.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,827	6.6	53.7
4	Chronic lower respiratory diseases (J40-J47)	61,235	5.1	41.2
5	Cerebrovascular diseases (I60-I69)	54,111	4.5	36.4
6	Diabetes mellitus (E10-E14)	35,478	2.9	23.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,269	2.3	18.3
8	Influenza and pneumonia (J09-J18)	24,071	2.0	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,616	1.9	15.2
10	Alzheimer's disease (G30)	21,800	1.8	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19,151	1.6	12.9
12	Septicemia (A40-A41)	15,839	1.3	10.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,538	1.2	9.8
14	Parkinson's disease (G20-G21)	11,554	1.0	7.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,417	0.8	6.3
...	All other causes (Residual)	204,384	17.0	137.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,634	100.0	31.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	987	37.5	11.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	286	10.9	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	8.1	2.5
4	Malignant neoplasms (C00-C97)	191	7.3	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	3.1	1.0
6	Influenza and pneumonia (J09-J18)	55	2.1	0.7
7	Septicemia (A40-A41)	44	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	34	1.3	0.4
9	Chronic lower respiratory diseases (J40-J47)	30	1.1	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	20	0.8	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.3	*
...	All other causes (Residual)	626	23.8	7.4
All races, male, 5-14 years				
...	All causes	3,585	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,346	37.5	6.6
2	Malignant neoplasms (C00-C97)	499	13.9	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	210	5.9	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	5.4	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	3.7	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	3.7	0.6
7	Chronic lower respiratory diseases (J40-J47)	78	2.2	0.4
8	Influenza and pneumonia (J09-J18)	52	1.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.3	0.2
10	Septicemia (A40-A41)	45	1.3	0.2
11	Cerebrovascular diseases (I60-I69)	37	1.0	0.2
12	Anemias (D50-D64)	21	0.6	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	14	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	Meningitis (G00,G03)	12	0.3	*
15	Pneumonitis due to solids and liquids (J69)	12	0.3	*
...	All other causes (Residual)	741	20.7	3.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	25,316	100.0	115.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,905	47.0	54.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,829	19.1	22.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,481	13.8	15.9
4	Malignant neoplasms (C00-C97)	987	3.9	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	700	2.8	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	241	1.0	1.1
7	Cerebrovascular diseases (I60-I69)	113	0.4	0.5
8	Diabetes mellitus (E10-E14)	98	0.4	0.4
9	Chronic lower respiratory diseases (J40-J47)	96	0.4	0.4
10	Legal intervention (Y35,Y89.0)	94	0.4	0.4
11	Influenza and pneumonia (J09-J18)	91	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	88	0.3	0.4
13	Septicemia (A40-A41)	85	0.3	0.4
14	Anemias (D50-D64)	49	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.2	0.2
...	All other causes (Residual)	2,416	9.5	11.1
All races, male, 25-34 years				
...	All causes	29,792	100.0	144.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,384	38.2	55.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,281	14.4	20.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,019	13.5	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,177	7.3	10.5
5	Malignant neoplasms (C00-C97)	1,692	5.7	8.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	653	2.2	3.2
7	Diabetes mellitus (E10-E14)	359	1.2	1.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	261	0.9	1.3
9	Cerebrovascular diseases (I60-I69)	252	0.8	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	243	0.8	1.2
11	Influenza and pneumonia (J09-J18)	187	0.6	0.9
12	Septicemia (A40-A41)	153	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	150	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	0.5	0.7
15	Legal intervention (Y35,Y89.0)	126	0.4	0.6
...	All other causes (Residual)	3,714	12.5	18.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	50,105	100.0	231.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,022	24.0	55.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,345	16.7	38.6
3	Malignant neoplasms (C00-C97)	5,701	11.4	26.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,152	10.3	23.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,506	5.0	11.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,289	4.6	10.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,722	3.4	8.0
8	Diabetes mellitus (E10-E14)	1,228	2.5	5.7
9	Cerebrovascular diseases (I60-I69)	1,143	2.3	5.3
10	Septicemia (A40-A41)	463	0.9	2.1
11	Influenza and pneumonia (J09-J18)	443	0.9	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	0.9	2.0
13	Chronic lower respiratory diseases (J40-J47)	392	0.8	1.8
14	Viral hepatitis (B15-B19)	326	0.7	1.5
15	Aortic aneurysm and dissection (I71)	278	0.6	1.3
...	All other causes (Residual)	7,667	15.3	35.5
All races, male, 45-54 years				
...	All causes	114,456	100.0	530.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,911	23.5	124.6
2	Malignant neoplasms (C00-C97)	25,370	22.2	117.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,017	12.2	64.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,824	5.1	27.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,788	5.1	26.8
6	Diabetes mellitus (E10-E14)	3,513	3.1	16.3
7	Cerebrovascular diseases (I60-I69)	3,507	3.1	16.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3,028	2.6	14.0
9	Chronic lower respiratory diseases (J40-J47)	2,125	1.9	9.8
10	Viral hepatitis (B15-B19)	1,997	1.7	9.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,572	1.4	7.3
12	Septicemia (A40-A41)	1,305	1.1	6.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,260	1.1	5.8
14	Influenza and pneumonia (J09-J18)	1,126	1.0	5.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	760	0.7	3.5
...	All other causes (Residual)	16,353	14.3	75.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	173,618	100.0	1,100.6
1	Malignant neoplasms (C00-C97)	56,559	32.6	358.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,558	26.2	288.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,357	4.8	53.0
4	Chronic lower respiratory diseases (J40-J47)	6,733	3.9	42.7
5	Diabetes mellitus (E10-E14)	6,644	3.8	42.1
6	Cerebrovascular diseases (I60-I69)	5,992	3.5	38.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,732	3.3	36.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,826	2.2	24.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,478	1.4	15.7
10	Septicemia (A40-A41)	2,203	1.3	14.0
11	Influenza and pneumonia (J09-J18)	1,881	1.1	11.9
12	Viral hepatitis (B15-B19)	1,761	1.0	11.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,344	0.8	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,273	0.7	8.1
15	Aortic aneurysm and dissection (I71)	1,093	0.6	6.9
...	All other causes (Residual)	22,184	12.8	140.6
All races, male, 65-74 years				
...	All causes	218,344	100.0	2,456.9
1	Malignant neoplasms (C00-C97)	75,924	34.8	854.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,532	25.4	624.9
3	Chronic lower respiratory diseases (J40-J47)	14,679	6.7	165.2
4	Cerebrovascular diseases (I60-I69)	9,351	4.3	105.2
5	Diabetes mellitus (E10-E14)	8,279	3.8	93.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,593	2.6	62.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,044	1.9	45.5
8	Septicemia (A40-A41)	3,256	1.5	36.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3,219	1.5	36.2
10	Influenza and pneumonia (J09-J18)	3,044	1.4	34.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,003	0.9	22.5
12	Aortic aneurysm and dissection (I71)	1,698	0.8	19.1
13	Alzheimer's disease (G30)	1,694	0.8	19.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,601	0.7	18.0
15	Parkinson's disease (G20-G21)	1,507	0.7	17.0
...	All other causes (Residual)	26,920	12.3	302.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	320,803	100.0	6,038.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,005	27.4	1,656.5
2	Malignant neoplasms (C00-C97)	85,927	26.8	1,617.4
3	Chronic lower respiratory diseases (J40-J47)	23,439	7.3	441.2
4	Cerebrovascular diseases (I60-I69)	17,700	5.5	333.2
5	Diabetes mellitus (E10-E14)	9,874	3.1	185.9
6	Alzheimer's disease (G30)	8,250	2.6	155.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,435	2.3	139.9
8	Influenza and pneumonia (J09-J18)	7,412	2.3	139.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,268	2.3	136.8
10	Parkinson's disease (G20-G21)	5,694	1.8	107.2
11	Septicemia (A40-A41)	4,768	1.5	89.7
12	Pneumonitis due to solids and liquids (J69)	2,926	0.9	55.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,667	0.8	50.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,570	0.8	48.4
15	Aortic aneurysm and dissection (I71)	2,341	0.7	44.1
...	All other causes (Residual)	44,527	13.9	838.1
All races, male, 85 years and over				
...	All causes	248,866	100.0	14,006.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,120	33.0	4,621.8
2	Malignant neoplasms (C00-C97)	39,964	16.1	2,249.2
3	Cerebrovascular diseases (I60-I69)	15,915	6.4	895.7
4	Chronic lower respiratory diseases (J40-J47)	13,480	5.4	758.7
5	Alzheimer's disease (G30)	11,479	4.6	646.0
6	Influenza and pneumonia (J09-J18)	9,642	3.9	542.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,686	2.7	376.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,201	2.5	349.0
9	Diabetes mellitus (E10-E14)	5,462	2.2	307.4
10	Parkinson's disease (G20-G21)	4,051	1.6	228.0
11	Pneumonitis due to solids and liquids (J69)	3,944	1.6	222.0
12	Septicemia (A40-A41)	3,364	1.4	189.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,805	1.1	157.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,700	0.7	95.7
15	Aortic aneurysm and dissection (I71)	1,453	0.6	81.8
...	All other causes (Residual)	40,600	16.3	2,285.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,219,744	100.0	797.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	306,246	25.1	200.2
2	Malignant neoplasms (C00-C97)	270,018	22.1	176.5
3	Cerebrovascular diseases (I60-I69)	81,841	6.7	53.5
4	Chronic lower respiratory diseases (J40-J47)	66,689	5.5	43.6
5	Alzheimer's disease (G30)	52,832	4.3	34.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,879	3.6	28.7
7	Diabetes mellitus (E10-E14)	35,904	2.9	23.5
8	Influenza and pneumonia (J09-J18)	28,646	2.3	18.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,832	2.0	15.6
10	Septicemia (A40-A41)	18,989	1.6	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,548	1.2	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,014	0.8	6.5
13	Parkinson's disease (G20-G21)	8,504	0.7	5.6
14	Pneumonitis due to solids and liquids (J69)	8,054	0.7	5.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,329	0.6	4.8
...	All other causes (Residual)	242,419	19.9	158.5
All races, female, 1-4 years				
...	All causes	2,069	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	601	29.0	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	260	12.6	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	8.9	2.3
4	Malignant neoplasms (C00-C97)	173	8.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	4.4	1.1
6	Influenza and pneumonia (J09-J18)	54	2.6	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	36	1.7	0.4
8	Septicemia (A40-A41)	34	1.6	0.4
9	Cerebrovascular diseases (I60-I69)	32	1.5	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.5	0.4
11	Chronic lower respiratory diseases (J40-J47)	27	1.3	0.3
12	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.7	*
13	Anemias (D50-D64)	11	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	501	24.2	6.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,562	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	848	33.1	4.3
2	Malignant neoplasms (C00-C97)	460	18.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	7.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.3	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	4.2	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	2.1	0.3
7	Influenza and pneumonia (J09-J18)	51	2.0	0.3
8	Cerebrovascular diseases (I60-I69)	46	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	40	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.4	0.2
11	Septicemia (A40-A41)	29	1.1	0.1
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	11	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
...	All other causes (Residual)	528	20.6	2.7
All races, female, 15-24 years				
...	All causes	8,666	100.0	42.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,992	46.1	19.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	722	8.3	3.5
3	Malignant neoplasms (C00-C97)	666	7.7	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	659	7.6	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	384	4.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	160	1.8	0.8
8	Cerebrovascular diseases (I60-I69)	82	0.9	0.4
9	Septicemia (A40-A41)	75	0.9	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.8	0.3
10	Influenza and pneumonia (J09-J18)	72	0.8	0.3
12	Diabetes mellitus (E10-E14)	70	0.8	0.3
13	Chronic lower respiratory diseases (J40-J47)	53	0.6	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.5	0.2
15	Anemias (D50-D64)	41	0.5	0.2
...	All other causes (Residual)	1,413	16.3	6.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,780	100.0	64.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,593	28.1	18.0
2	Malignant neoplasms (C00-C97)	1,771	13.9	8.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,046	8.2	5.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	997	7.8	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	739	5.8	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	438	3.4	2.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	326	2.6	1.6
8	Cerebrovascular diseases (I60-I69)	253	2.0	1.3
9	Diabetes mellitus (E10-E14)	251	2.0	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	1.4	0.9
11	Septicemia (A40-A41)	144	1.1	0.7
11	Influenza and pneumonia (J09-J18)	144	1.1	0.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	123	1.0	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	0.9	0.6
15	Chronic lower respiratory diseases (J40-J47)	113	0.9	0.6
...	All other causes (Residual)	2,548	19.9	12.8
All races, female, 35-44 years				
...	All causes	29,501	100.0	136.9
1	Malignant neoplasms (C00-C97)	7,587	25.7	35.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,909	16.6	22.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,494	11.8	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,570	5.3	7.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,066	3.6	4.9
6	Cerebrovascular diseases (I60-I69)	990	3.4	4.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	848	2.9	3.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	763	2.6	3.5
9	Diabetes mellitus (E10-E14)	756	2.6	3.5
10	Septicemia (A40-A41)	447	1.5	2.1
11	Chronic lower respiratory diseases (J40-J47)	404	1.4	1.9
12	Influenza and pneumonia (J09-J18)	341	1.2	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	326	1.1	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	256	0.9	1.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	220	0.7	1.0
...	All other causes (Residual)	5,524	18.7	25.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	70,230	100.0	315.2
1	Malignant neoplasms (C00-C97)	24,797	35.3	111.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,523	15.0	47.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,298	9.0	28.3
4	Cerebrovascular diseases (I60-I69)	2,878	4.1	12.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,424	3.5	10.9
6	Diabetes mellitus (E10-E14)	2,240	3.2	10.1
7	Chronic lower respiratory diseases (J40-J47)	2,028	2.9	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,954	2.8	8.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,128	1.6	5.1
10	Septicemia (A40-A41)	1,126	1.6	5.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	973	1.4	4.4
12	Viral hepatitis (B15-B19)	818	1.2	3.7
13	Influenza and pneumonia (J09-J18)	783	1.1	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	568	0.8	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	0.7	2.1
...	All other causes (Residual)	11,217	16.0	50.3
All races, female, 55-64 years				
...	All causes	113,492	100.0	670.1
1	Malignant neoplasms (C00-C97)	46,612	41.1	275.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,969	17.6	117.9
3	Chronic lower respiratory diseases (J40-J47)	6,044	5.3	35.7
4	Diabetes mellitus (E10-E14)	4,660	4.1	27.5
5	Cerebrovascular diseases (I60-I69)	4,508	4.0	26.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,836	3.4	22.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,272	2.0	13.4
8	Septicemia (A40-A41)	2,028	1.8	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,962	1.7	11.6
10	Influenza and pneumonia (J09-J18)	1,271	1.1	7.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,243	1.1	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	851	0.7	5.0
13	Viral hepatitis (B15-B19)	652	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	578	0.5	3.4
15	Alzheimer's disease (G30)	398	0.4	2.3
...	All other causes (Residual)	16,608	14.6	98.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	170,894	100.0	1,633.0
1	Malignant neoplasms (C00-C97)	62,542	36.6	597.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,057	19.9	325.4
3	Chronic lower respiratory diseases (J40-J47)	13,985	8.2	133.6
4	Cerebrovascular diseases (I60-I69)	8,656	5.1	82.7
5	Diabetes mellitus (E10-E14)	6,833	4.0	65.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,708	2.2	35.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,160	1.8	30.2
8	Septicemia (A40-A41)	3,089	1.8	29.5
9	Influenza and pneumonia (J09-J18)	2,503	1.5	23.9
10	Alzheimer's disease (G30)	2,290	1.3	21.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,948	1.1	18.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,532	0.9	14.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,048	0.6	10.0
14	Aortic aneurysm and dissection (I71)	918	0.5	8.8
15	Parkinson's disease (G20-G21)	803	0.5	7.7
...	All other causes (Residual)	23,822	13.9	227.6
All races, female, 75-84 years				
...	All causes	331,879	100.0	4,304.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,252	25.1	1,079.7
2	Malignant neoplasms (C00-C97)	77,681	23.4	1,007.4
3	Chronic lower respiratory diseases (J40-J47)	24,602	7.4	319.1
4	Cerebrovascular diseases (I60-I69)	24,279	7.3	314.9
5	Alzheimer's disease (G30)	14,759	4.4	191.4
6	Diabetes mellitus (E10-E14)	11,315	3.4	146.7
7	Influenza and pneumonia (J09-J18)	7,447	2.2	96.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,276	2.2	94.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,468	1.9	83.9
10	Septicemia (A40-A41)	5,635	1.7	73.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,775	1.1	49.0
12	Parkinson's disease (G20-G21)	3,669	1.1	47.6
13	Pneumonitis due to solids and liquids (J69)	2,261	0.7	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,147	0.6	27.8
15	Aortic aneurysm and dissection (I71)	2,005	0.6	26.0
...	All other causes (Residual)	55,308	16.7	717.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	464,781	100.0	12,442.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	153,129	32.9	4,099.3
2	Malignant neoplasms (C00-C97)	47,692	10.3	1,276.7
3	Cerebrovascular diseases (I60-I69)	40,060	8.6	1,072.4
4	Alzheimer's disease (G30)	35,325	7.6	945.7
5	Chronic lower respiratory diseases (J40-J47)	19,377	4.2	518.7
6	Influenza and pneumonia (J09-J18)	15,893	3.4	425.5
7	Diabetes mellitus (E10-E14)	9,765	2.1	261.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,602	2.1	257.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,335	2.0	249.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,731	1.7	207.0
11	Septicemia (A40-A41)	6,250	1.3	167.3
12	Pneumonitis due to solids and liquids (J69)	4,507	1.0	120.7
13	Parkinson's disease (G20-G21)	3,860	0.8	103.3
14	Atherosclerosis (I70)	3,230	0.7	86.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,509	0.5	67.2
...	All other causes (Residual)	96,516	20.8	2,583.8
White, both sexes, all ages				
...	All causes	2,074,151	100.0	851.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	531,636	25.6	218.3
2	Malignant neoplasms (C00-C97)	483,939	23.3	198.7
3	Chronic lower respiratory diseases (J40-J47)	118,081	5.7	48.5
4	Cerebrovascular diseases (I60-I69)	114,695	5.5	47.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106,252	5.1	43.6
6	Alzheimer's disease (G30)	68,933	3.3	28.3
7	Diabetes mellitus (E10-E14)	56,390	2.7	23.2
8	Influenza and pneumonia (J09-J18)	45,947	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,871	1.8	15.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,348	1.5	12.9
11	Septicemia (A40-A41)	27,750	1.3	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25,490	1.2	10.5
13	Parkinson's disease (G20-G21)	18,864	0.9	7.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,583	0.9	7.6
15	Pneumonitis due to solids and liquids (J69)	15,229	0.7	6.3
...	All other causes (Residual)	374,143	18.0	153.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,287	100.0	25.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,169	35.6	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	396	12.0	3.1
3	Malignant neoplasms (C00-C97)	287	8.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	6.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	3.4	0.9
6	Influenza and pneumonia (J09-J18)	73	2.2	0.6
7	Septicemia (A40-A41)	60	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	40	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	34	1.0	0.3
11	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
14	Meningitis (G00,G03)	12	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
...	All other causes (Residual)	777	23.6	6.1
White, both sexes, 5-14 years				
...	All causes	4,480	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,625	36.3	5.2
2	Malignant neoplasms (C00-C97)	776	17.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	274	6.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	4.4	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	3.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	2.8	0.4
7	Influenza and pneumonia (J09-J18)	77	1.7	0.2
8	Cerebrovascular diseases (I60-I69)	63	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	1.3	0.2
10	Septicemia (A40-A41)	59	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	54	1.2	0.2
12	Meningitis (G00,G03)	19	0.4	*
13	Anemias (D50-D64)	18	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	16	0.4	*
...	All other causes (Residual)	934	20.8	3.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,137	100.0	75.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,481	53.6	40.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,442	13.7	10.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,193	8.7	6.6
4	Malignant neoplasms (C00-C97)	1,296	5.2	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	706	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	299	1.2	0.9
7	Cerebrovascular diseases (I60-I69)	138	0.5	0.4
8	Influenza and pneumonia (J09-J18)	129	0.5	0.4
9	Diabetes mellitus (E10-E14)	109	0.4	0.3
10	Septicemia (A40-A41)	106	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	86	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	79	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.2	0.2
...	All other causes (Residual)	2,906	11.6	8.8
White, both sexes, 25-34 years				
...	All causes	31,181	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,646	40.6	39.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,524	14.5	14.2
3	Malignant neoplasms (C00-C97)	2,660	8.5	8.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,073	6.6	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,050	6.6	6.4
6	Diabetes mellitus (E10-E14)	379	1.2	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	367	1.2	1.2
8	Cerebrovascular diseases (I60-I69)	332	1.1	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	322	1.0	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	310	1.0	1.0
11	Influenza and pneumonia (J09-J18)	225	0.7	0.7
12	Septicemia (A40-A41)	194	0.6	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	193	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	159	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	0.5	0.4
...	All other causes (Residual)	4,606	14.8	14.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	59,571	100.0	172.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,211	23.9	41.2
2	Malignant neoplasms (C00-C97)	10,289	17.3	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,406	14.1	24.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,033	10.1	17.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,184	3.7	6.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,577	2.6	4.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,522	2.6	4.4
8	Cerebrovascular diseases (I60-I69)	1,356	2.3	3.9
9	Diabetes mellitus (E10-E14)	1,344	2.3	3.9
10	Septicemia (A40-A41)	608	1.0	1.8
11	Influenza and pneumonia (J09-J18)	580	1.0	1.7
12	Chronic lower respiratory diseases (J40-J47)	578	1.0	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	421	0.7	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	0.7	1.2
15	Viral hepatitis (B15-B19)	387	0.6	1.1
...	All other causes (Residual)	9,669	16.2	28.1
White, both sexes, 45-54 years				
...	All causes	142,233	100.0	394.0
1	Malignant neoplasms (C00-C97)	39,557	27.8	109.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,135	19.8	77.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,018	12.0	47.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,259	5.1	20.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,007	4.9	19.4
6	Cerebrovascular diseases (I60-I69)	4,139	2.9	11.5
7	Diabetes mellitus (E10-E14)	4,123	2.9	11.4
8	Chronic lower respiratory diseases (J40-J47)	3,395	2.4	9.4
9	Viral hepatitis (B15-B19)	2,345	1.6	6.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,747	1.2	4.8
11	Septicemia (A40-A41)	1,628	1.1	4.5
12	Influenza and pneumonia (J09-J18)	1,403	1.0	3.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,346	0.9	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,271	0.9	3.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	0.5	1.9
...	All other causes (Residual)	21,191	14.9	58.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	231,287	100.0	835.4
1	Malignant neoplasms (C00-C97)	85,373	36.9	308.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,585	22.3	186.3
3	Chronic lower respiratory diseases (J40-J47)	11,412	4.9	41.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,224	4.4	36.9
5	Diabetes mellitus (E10-E14)	8,378	3.6	30.3
6	Cerebrovascular diseases (I60-I69)	7,438	3.2	26.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,879	3.0	24.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,792	2.1	17.3
9	Septicemia (A40-A41)	3,120	1.3	11.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,892	1.3	10.4
11	Influenza and pneumonia (J09-J18)	2,483	1.1	9.0
12	Viral hepatitis (B15-B19)	1,887	0.8	6.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,313	0.6	4.7
14	Aortic aneurysm and dissection (I71)	1,192	0.5	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,190	0.5	4.3
...	All other causes (Residual)	31,129	13.5	112.4
White, both sexes, 65-74 years				
...	All causes	327,377	100.0	1,970.3
1	Malignant neoplasms (C00-C97)	118,826	36.3	715.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,600	22.5	443.0
3	Chronic lower respiratory diseases (J40-J47)	26,389	8.1	158.8
4	Cerebrovascular diseases (I60-I69)	14,023	4.3	84.4
5	Diabetes mellitus (E10-E14)	11,652	3.6	70.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,529	2.3	45.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,655	1.7	34.0
8	Septicemia (A40-A41)	4,852	1.5	29.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	4,624	1.4	27.8
10	Influenza and pneumonia (J09-J18)	4,551	1.4	27.4
11	Alzheimer's disease (G30)	3,563	1.1	21.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,317	0.7	13.9
13	Aortic aneurysm and dissection (I71)	2,295	0.7	13.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,190	0.7	13.2
15	Parkinson's disease (G20-G21)	2,115	0.6	12.7
...	All other causes (Residual)	43,196	13.2	260.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	579,122	100.0	5,030.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	151,041	26.1	1,312.1
2	Malignant neoplasms (C00-C97)	145,585	25.1	1,264.7
3	Chronic lower respiratory diseases (J40-J47)	44,986	7.8	390.8
4	Cerebrovascular diseases (I60-I69)	36,340	6.3	315.7
5	Alzheimer's disease (G30)	21,115	3.6	183.4
6	Diabetes mellitus (E10-E14)	17,411	3.0	151.2
7	Influenza and pneumonia (J09-J18)	13,074	2.3	113.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,584	2.2	109.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,282	2.1	106.7
10	Parkinson's disease (G20-G21)	8,832	1.5	76.7
11	Septicemia (A40-A41)	8,684	1.5	75.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,113	0.9	44.4
13	Pneumonitis due to solids and liquids (J69)	4,647	0.8	40.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,301	0.7	37.4
15	Aortic aneurysm and dissection (I71)	3,959	0.7	34.4
...	All other causes (Residual)	89,168	15.4	774.6
White, both sexes, 85 years and over				
...	All causes	651,521	100.0	13,176.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215,539	33.1	4,359.1
2	Malignant neoplasms (C00-C97)	79,226	12.2	1,602.3
3	Cerebrovascular diseases (I60-I69)	50,734	7.8	1,026.1
4	Alzheimer's disease (G30)	43,484	6.7	879.4
5	Chronic lower respiratory diseases (J40-J47)	30,976	4.8	626.5
6	Influenza and pneumonia (J09-J18)	23,217	3.6	469.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,933	2.3	302.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,966	2.1	282.5
9	Diabetes mellitus (E10-E14)	12,968	2.0	262.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,048	1.4	183.0
11	Septicemia (A40-A41)	8,280	1.3	167.5
12	Pneumonitis due to solids and liquids (J69)	7,792	1.2	157.6
13	Parkinson's disease (G20-G21)	7,497	1.2	151.6
14	Atherosclerosis (I70)	4,253	0.7	86.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,928	0.6	79.4
...	All other causes (Residual)	125,680	19.3	2,541.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,023,951	100.0	848.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	266,908	26.1	221.1
2	Malignant neoplasms (C00-C97)	252,049	24.6	208.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,059	6.6	56.4
4	Chronic lower respiratory diseases (J40-J47)	55,934	5.5	46.3
5	Cerebrovascular diseases (I60-I69)	44,714	4.4	37.0
6	Diabetes mellitus (E10-E14)	28,744	2.8	23.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,725	2.4	20.5
8	Influenza and pneumonia (J09-J18)	20,720	2.0	17.2
9	Alzheimer's disease (G30)	20,185	2.0	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,242	1.8	15.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16,801	1.6	13.9
12	Septicemia (A40-A41)	12,600	1.2	10.4
13	Parkinson's disease (G20-G21)	10,895	1.1	9.0
14	Pneumonitis due to solids and liquids (J69)	8,010	0.8	6.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,106	0.7	5.9
...	All other causes (Residual)	168,259	16.4	139.4
White, male, 1-4 years				
...	All causes	1,847	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	738	40.0	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	11.4	3.2
3	Malignant neoplasms (C00-C97)	146	7.9	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	116	6.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.5	0.7
6	Septicemia (A40-A41)	33	1.8	0.5
7	Influenza and pneumonia (J09-J18)	32	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	19	1.0	*
10	Cerebrovascular diseases (I60-I69)	16	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.7	*
12	Meningitis (G00,G03)	7	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	427	23.1	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,590	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,000	38.6	6.2
2	Malignant neoplasms (C00-C97)	404	15.6	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	5.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	4.1	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	3.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.5	0.6
7	Septicemia (A40-A41)	38	1.5	0.2
8	Influenza and pneumonia (J09-J18)	36	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	34	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	32	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.2	0.2
12	Anemias (D50-D64)	13	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
14	Diabetes mellitus (E10-E14)	9	0.3	*
14	Pneumonitis due to solids and liquids (J69)	9	0.3	*
...	All other causes (Residual)	541	20.9	3.4
White, male, 15-24 years				
...	All causes	18,513	100.0	108.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,121	54.7	59.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,896	15.6	16.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,792	9.7	10.5
4	Malignant neoplasms (C00-C97)	781	4.2	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	445	2.4	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	0.9	1.0
7	Cerebrovascular diseases (I60-I69)	76	0.4	0.4
8	Influenza and pneumonia (J09-J18)	74	0.4	0.4
9	Diabetes mellitus (E10-E14)	73	0.4	0.4
10	Septicemia (A40-A41)	59	0.3	0.3
11	Chronic lower respiratory diseases (J40-J47)	49	0.3	0.3
12	Legal intervention (Y35,Y89.0)	46	0.2	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.1	0.2
...	All other causes (Residual)	1,837	9.9	10.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	21,998	100.0	134.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,624	43.7	58.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,676	16.7	22.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,625	7.4	9.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,433	6.5	8.7
5	Malignant neoplasms (C00-C97)	1,329	6.0	8.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	263	1.2	1.6
7	Diabetes mellitus (E10-E14)	234	1.1	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	219	1.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	0.8	1.1
10	Cerebrovascular diseases (I60-I69)	167	0.8	1.0
11	Influenza and pneumonia (J09-J18)	127	0.6	0.8
12	Septicemia (A40-A41)	103	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	86	0.4	0.5
14	Legal intervention (Y35,Y89.0)	81	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.3	0.5
...	All other causes (Residual)	2,770	12.6	16.9
White, male, 35-44 years				
...	All causes	38,121	100.0	218.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,074	26.4	57.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,058	15.9	34.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,628	12.1	26.5
4	Malignant neoplasms (C00-C97)	4,522	11.9	25.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,477	3.9	8.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,249	3.3	7.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,117	2.9	6.4
8	Diabetes mellitus (E10-E14)	861	2.3	4.9
9	Cerebrovascular diseases (I60-I69)	746	2.0	4.3
10	Influenza and pneumonia (J09-J18)	323	0.8	1.8
11	Septicemia (A40-A41)	319	0.8	1.8
12	Chronic lower respiratory diseases (J40-J47)	278	0.7	1.6
13	Viral hepatitis (B15-B19)	260	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	242	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	214	0.6	1.2
...	All other causes (Residual)	5,753	15.1	32.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	89,556	100.0	498.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,876	23.3	116.2
2	Malignant neoplasms (C00-C97)	20,133	22.5	112.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,725	13.1	65.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,436	6.1	30.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,007	5.6	27.9
6	Diabetes mellitus (E10-E14)	2,591	2.9	14.4
7	Cerebrovascular diseases (I60-I69)	2,330	2.6	13.0
8	Chronic lower respiratory diseases (J40-J47)	1,777	2.0	9.9
9	Viral hepatitis (B15-B19)	1,686	1.9	9.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,422	1.6	7.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	903	1.0	5.0
12	Septicemia (A40-A41)	878	1.0	4.9
13	Influenza and pneumonia (J09-J18)	810	0.9	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	788	0.9	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	0.5	2.4
...	All other causes (Residual)	12,762	14.3	71.0
White, male, 55-64 years				
...	All causes	140,785	100.0	1,042.7
1	Malignant neoplasms (C00-C97)	46,817	33.3	346.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,646	26.0	271.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,984	5.0	51.7
4	Chronic lower respiratory diseases (J40-J47)	5,956	4.2	44.1
5	Diabetes mellitus (E10-E14)	5,091	3.6	37.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,963	3.5	36.8
7	Cerebrovascular diseases (I60-I69)	4,242	3.0	31.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,618	2.6	26.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,655	1.2	12.3
10	Septicemia (A40-A41)	1,616	1.1	12.0
11	Influenza and pneumonia (J09-J18)	1,486	1.1	11.0
12	Viral hepatitis (B15-B19)	1,395	1.0	10.3
13	Aortic aneurysm and dissection (I71)	909	0.6	6.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	828	0.6	6.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	705	0.5	5.2
...	All other causes (Residual)	17,874	12.7	132.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	184,828	100.0	2,396.7
1	Malignant neoplasms (C00-C97)	65,198	35.3	845.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,500	25.2	603.0
3	Chronic lower respiratory diseases (J40-J47)	13,399	7.2	173.7
4	Cerebrovascular diseases (I60-I69)	7,272	3.9	94.3
5	Diabetes mellitus (E10-E14)	6,653	3.6	86.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,798	2.6	62.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,025	1.6	39.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,874	1.6	37.3
9	Septicemia (A40-A41)	2,494	1.3	32.3
10	Influenza and pneumonia (J09-J18)	2,475	1.3	32.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,901	1.0	24.7
12	Alzheimer's disease (G30)	1,522	0.8	19.7
13	Aortic aneurysm and dissection (I71)	1,514	0.8	19.6
14	Parkinson's disease (G20-G21)	1,374	0.7	17.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,177	0.6	15.3
...	All other causes (Residual)	22,652	12.3	293.7
White, male, 75-84 years				
...	All causes	286,824	100.0	6,049.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,676	27.4	1,659.3
2	Malignant neoplasms (C00-C97)	76,691	26.7	1,617.4
3	Chronic lower respiratory diseases (J40-J47)	21,735	7.6	458.4
4	Cerebrovascular diseases (I60-I69)	15,322	5.3	323.1
5	Diabetes mellitus (E10-E14)	8,425	2.9	177.7
6	Alzheimer's disease (G30)	7,608	2.7	160.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,663	2.3	140.5
8	Influenza and pneumonia (J09-J18)	6,517	2.3	137.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,400	2.2	135.0
10	Parkinson's disease (G20-G21)	5,399	1.9	113.9
11	Septicemia (A40-A41)	4,068	1.4	85.8
12	Pneumonitis due to solids and liquids (J69)	2,624	0.9	55.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,370	0.8	50.0
14	Aortic aneurysm and dissection (I71)	2,158	0.8	45.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,137	0.7	45.1
...	All other causes (Residual)	40,031	14.0	844.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	228,227	100.0	14,286.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,979	33.3	4,756.1
2	Malignant neoplasms (C00-C97)	35,995	15.8	2,253.2
3	Cerebrovascular diseases (I60-I69)	14,457	6.3	905.0
4	Chronic lower respiratory diseases (J40-J47)	12,589	5.5	788.0
5	Alzheimer's disease (G30)	10,710	4.7	670.4
6	Influenza and pneumonia (J09-J18)	8,752	3.8	547.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,967	2.6	373.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,848	2.6	366.1
9	Diabetes mellitus (E10-E14)	4,799	2.1	300.4
10	Parkinson's disease (G20-G21)	3,848	1.7	240.9
11	Pneumonitis due to solids and liquids (J69)	3,664	1.6	229.4
12	Septicemia (A40-A41)	2,906	1.3	181.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,402	1.1	150.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,588	0.7	99.4
15	Aortic aneurysm and dissection (I71)	1,371	0.6	85.8
...	All other causes (Residual)	37,352	16.4	2,338.1
White, female, all ages				
...	All causes	1,050,200	100.0	854.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,728	25.2	215.5
2	Malignant neoplasms (C00-C97)	231,890	22.1	188.8
3	Cerebrovascular diseases (I60-I69)	69,981	6.7	57.0
4	Chronic lower respiratory diseases (J40-J47)	62,147	5.9	50.6
5	Alzheimer's disease (G30)	48,748	4.6	39.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,193	3.6	31.1
7	Diabetes mellitus (E10-E14)	27,646	2.6	22.5
8	Influenza and pneumonia (J09-J18)	25,227	2.4	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,629	1.8	15.2
10	Septicemia (A40-A41)	15,150	1.4	12.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,477	1.1	9.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,689	0.8	7.1
13	Parkinson's disease (G20-G21)	7,969	0.8	6.5
14	Pneumonitis due to solids and liquids (J69)	7,219	0.7	5.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,623	0.6	5.4
...	All other causes (Residual)	205,884	19.6	167.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,440	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	29.9	6.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.8	3.0
3	Malignant neoplasms (C00-C97)	141	9.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	6.9	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	4.5	1.0
6	Influenza and pneumonia (J09-J18)	41	2.8	0.7
7	Septicemia (A40-A41)	27	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.7	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	21	1.5	0.3
10	Cerebrovascular diseases (I60-I69)	18	1.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.6	*
14	Meningitis (G00,G03)	5	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (Residual)	349	24.2	5.6
White, female, 5-14 years				
...	All causes	1,890	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	625	33.1	4.1
2	Malignant neoplasms (C00-C97)	372	19.7	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	6.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	4.0	0.5
6	Influenza and pneumonia (J09-J18)	41	2.2	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.8	0.2
8	Cerebrovascular diseases (I60-I69)	31	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.5	0.2
10	Septicemia (A40-A41)	21	1.1	0.1
11	Chronic lower respiratory diseases (J40-J47)	20	1.1	0.1
12	Meningitis (G00,G03)	12	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.5	*
14	Diabetes mellitus (E10-E14)	6	0.3	*
15	Anemias (D50-D64)	5	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	379	20.1	2.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,624	100.0	41.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,360	50.7	20.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	546	8.2	3.4
3	Malignant neoplasms (C00-C97)	515	7.8	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	6.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	3.9	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	86	1.3	0.5
8	Cerebrovascular diseases (I60-I69)	62	0.9	0.4
9	Influenza and pneumonia (J09-J18)	55	0.8	0.3
10	Septicemia (A40-A41)	47	0.7	0.3
11	Diabetes mellitus (E10-E14)	36	0.5	0.2
12	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.5	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.4	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.3	*
...	All other causes (Residual)	1,023	15.4	6.4
White, female, 25-34 years				
...	All causes	9,183	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,022	32.9	19.6
2	Malignant neoplasms (C00-C97)	1,331	14.5	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	848	9.2	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	640	7.0	4.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	425	4.6	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	193	2.1	1.3
7	Cerebrovascular diseases (I60-I69)	165	1.8	1.1
8	Diabetes mellitus (E10-E14)	145	1.6	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	1.5	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	104	1.1	0.7
11	Influenza and pneumonia (J09-J18)	98	1.1	0.6
12	Septicemia (A40-A41)	91	1.0	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	73	0.8	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.7	0.4
...	All other causes (Residual)	1,755	19.1	11.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	21,450	100.0	126.2
1	Malignant neoplasms (C00-C97)	5,767	26.9	33.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,137	19.3	24.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,348	10.9	13.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,405	6.6	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	707	3.3	4.2
6	Cerebrovascular diseases (I60-I69)	610	2.8	3.6
7	Diabetes mellitus (E10-E14)	483	2.3	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	460	2.1	2.7
9	Chronic lower respiratory diseases (J40-J47)	300	1.4	1.8
10	Septicemia (A40-A41)	289	1.3	1.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	273	1.3	1.6
12	Influenza and pneumonia (J09-J18)	257	1.2	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	1.0	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	0.8	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	137	0.6	0.8
...	All other causes (Residual)	3,906	18.2	23.0
White, female, 45-54 years				
...	All causes	52,677	100.0	290.5
1	Malignant neoplasms (C00-C97)	19,424	36.9	107.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,259	13.8	40.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,293	10.0	29.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,000	3.8	11.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,823	3.5	10.1
6	Cerebrovascular diseases (I60-I69)	1,809	3.4	10.0
7	Chronic lower respiratory diseases (J40-J47)	1,618	3.1	8.9
8	Diabetes mellitus (E10-E14)	1,532	2.9	8.4
9	Septicemia (A40-A41)	750	1.4	4.1
10	Viral hepatitis (B15-B19)	659	1.3	3.6
11	Influenza and pneumonia (J09-J18)	593	1.1	3.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	558	1.1	3.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	368	0.7	2.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	325	0.6	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	0.5	1.5
...	All other causes (Residual)	8,395	15.9	46.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	90,502	100.0	638.0
1	Malignant neoplasms (C00-C97)	38,556	42.6	271.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,939	16.5	105.3
3	Chronic lower respiratory diseases (J40-J47)	5,456	6.0	38.5
4	Diabetes mellitus (E10-E14)	3,287	3.6	23.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,240	3.6	22.8
6	Cerebrovascular diseases (I60-I69)	3,196	3.5	22.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,916	2.1	13.5
8	Septicemia (A40-A41)	1,504	1.7	10.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,237	1.4	8.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,174	1.3	8.3
11	Influenza and pneumonia (J09-J18)	997	1.1	7.0
12	Viral hepatitis (B15-B19)	492	0.5	3.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	485	0.5	3.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	485	0.5	3.4
15	Alzheimer's disease (G30)	374	0.4	2.6
...	All other causes (Residual)	13,164	14.5	92.8
White, female, 65-74 years				
...	All causes	142,549	100.0	1,600.9
1	Malignant neoplasms (C00-C97)	53,628	37.6	602.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,100	19.0	304.4
3	Chronic lower respiratory diseases (J40-J47)	12,990	9.1	145.9
4	Cerebrovascular diseases (I60-I69)	6,751	4.7	75.8
5	Diabetes mellitus (E10-E14)	4,999	3.5	56.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,731	1.9	30.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,630	1.8	29.5
8	Septicemia (A40-A41)	2,358	1.7	26.5
9	Influenza and pneumonia (J09-J18)	2,076	1.5	23.3
10	Alzheimer's disease (G30)	2,041	1.4	22.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,750	1.2	19.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,059	0.7	11.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	908	0.6	10.2
14	Aortic aneurysm and dissection (I71)	781	0.5	8.8
15	Parkinson's disease (G20-G21)	741	0.5	8.3
...	All other causes (Residual)	20,006	14.0	224.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	292,298	100.0	4,317.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,365	24.8	1,068.9
2	Malignant neoplasms (C00-C97)	68,894	23.6	1,017.7
3	Chronic lower respiratory diseases (J40-J47)	23,251	8.0	343.4
4	Cerebrovascular diseases (I60-I69)	21,018	7.2	310.5
5	Alzheimer's disease (G30)	13,507	4.6	199.5
6	Diabetes mellitus (E10-E14)	8,986	3.1	132.7
7	Influenza and pneumonia (J09-J18)	6,557	2.2	96.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,921	2.0	87.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,882	2.0	86.9
10	Septicemia (A40-A41)	4,616	1.6	68.2
11	Parkinson's disease (G20-G21)	3,433	1.2	50.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,976	1.0	44.0
13	Pneumonitis due to solids and liquids (J69)	2,023	0.7	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,931	0.7	28.5
15	Aortic aneurysm and dissection (I71)	1,801	0.6	26.6
...	All other causes (Residual)	49,137	16.8	725.8
White, female, 85 years and over				
...	All causes	423,294	100.0	12,646.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	139,560	33.0	4,169.6
2	Malignant neoplasms (C00-C97)	43,231	10.2	1,291.6
3	Cerebrovascular diseases (I60-I69)	36,277	8.6	1,083.8
4	Alzheimer's disease (G30)	32,774	7.7	979.2
5	Chronic lower respiratory diseases (J40-J47)	18,387	4.3	549.3
6	Influenza and pneumonia (J09-J18)	14,465	3.4	432.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,085	2.1	271.4
8	Diabetes mellitus (E10-E14)	8,169	1.9	244.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,999	1.9	239.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,646	1.6	198.6
11	Septicemia (A40-A41)	5,374	1.3	160.6
12	Pneumonitis due to solids and liquids (J69)	4,128	1.0	123.3
13	Parkinson's disease (G20-G21)	3,649	0.9	109.0
14	Atherosclerosis (I70)	2,987	0.7	89.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,340	0.6	69.9
...	All other causes (Residual)	88,223	20.8	2,635.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	289,585	100.0	723.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,209	24.6	177.9
2	Malignant neoplasms (C00-C97)	64,049	22.1	160.0
3	Cerebrovascular diseases (I60-I69)	17,085	5.9	42.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,559	4.7	33.9
5	Diabetes mellitus (E10-E14)	12,459	4.3	31.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,870	3.1	22.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,392	2.9	21.0
8	Chronic lower respiratory diseases (J40-J47)	7,901	2.7	19.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,470	2.2	16.2
10	Septicemia (A40-A41)	6,297	2.2	15.7
11	Influenza and pneumonia (J09-J18)	5,155	1.8	12.9
12	Certain conditions originating in the perinatal period (P00-P96)	5,052	1.7	12.6
13	Alzheimer's disease (G30)	4,760	1.6	11.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,640	1.6	11.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,558	0.9	6.4
...	All other causes (Residual)	51,129	17.7	127.7
Black, both sexes, 1-4 years				
...	All causes	1,137	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	28.6	12.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	14.8	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	10.5	4.4
4	Malignant neoplasms (C00-C97)	55	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.8	1.6
6	Chronic lower respiratory diseases (J40-J47)	30	2.6	1.1
7	Influenza and pneumonia (J09-J18)	29	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	27	2.4	1.0
9	Septicemia (A40-A41)	16	1.4	*
9	Cerebrovascular diseases (I60-I69)	16	1.4	*
11	Anemias (D50-D64)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
...	All other causes (Residual)	271	23.8	10.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,363	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	459	33.7	7.0
2	Malignant neoplasms (C00-C97)	143	10.5	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	10.3	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	5.6	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.6	1.0
6	Chronic lower respiratory diseases (J40-J47)	60	4.4	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.1	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.5	0.3
8	Influenza and pneumonia (J09-J18)	21	1.5	0.3
10	Anemias (D50-D64)	13	1.0	*
10	Cerebrovascular diseases (I60-I69)	13	1.0	*
12	Septicemia (A40-A41)	11	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
14	Diabetes mellitus (E10-E14)	6	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	274	20.1	4.2
Black, both sexes, 15-24 years				
...	All causes	7,380	100.0	109.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,213	43.5	47.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,749	23.7	25.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	406	5.5	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	4.7	5.2
5	Malignant neoplasms (C00-C97)	261	3.5	3.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.5	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.1	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	69	0.9	1.0
9	Anemias (D50-D64)	67	0.9	1.0
10	Chronic lower respiratory diseases (J40-J47)	65	0.9	1.0
11	Diabetes mellitus (E10-E14)	56	0.8	0.8
12	Legal intervention (Y35,Y89.0)	48	0.7	0.7
13	Septicemia (A40-A41)	47	0.6	0.7
14	Cerebrovascular diseases (I60-I69)	46	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.4	0.4
...	All other causes (Residual)	785	10.6	11.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,662	100.0	168.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,560	26.5	44.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,779	18.4	31.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	997	10.3	17.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	705	7.3	12.3
5	Malignant neoplasms (C00-C97)	618	6.4	10.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	481	5.0	8.4
7	Diabetes mellitus (E10-E14)	213	2.2	3.7
8	Cerebrovascular diseases (I60-I69)	148	1.5	2.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	1.2	2.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	108	1.1	1.9
11	Anemias (D50-D64)	102	1.1	1.8
12	Chronic lower respiratory diseases (J40-J47)	96	1.0	1.7
13	Influenza and pneumonia (J09-J18)	94	1.0	1.6
14	Septicemia (A40-A41)	90	0.9	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	0.9	1.5
...	All other causes (Residual)	1,475	15.3	25.8
Black, both sexes, 35-44 years				
...	All causes	16,990	100.0	299.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,992	17.6	52.8
2	Malignant neoplasms (C00-C97)	2,365	13.9	41.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,158	12.7	38.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,002	11.8	35.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,366	8.0	24.1
6	Cerebrovascular diseases (I60-I69)	655	3.9	11.5
7	Diabetes mellitus (E10-E14)	562	3.3	9.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	2.5	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	1.9	5.8
10	Septicemia (A40-A41)	274	1.6	4.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	203	1.2	3.6
12	Chronic lower respiratory diseases (J40-J47)	193	1.1	3.4
13	Influenza and pneumonia (J09-J18)	178	1.0	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.0	3.1
15	Anemias (D50-D64)	142	0.8	2.5
...	All other causes (Residual)	2,970	17.5	52.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	36,818	100.0	694.2
1	Malignant neoplasms (C00-C97)	8,818	24.0	166.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,357	22.7	157.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,766	7.5	52.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,351	6.4	44.3
5	Cerebrovascular diseases (I60-I69)	1,946	5.3	36.7
6	Diabetes mellitus (E10-E14)	1,433	3.9	27.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	853	2.3	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	823	2.2	15.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	802	2.2	15.1
10	Septicemia (A40-A41)	721	2.0	13.6
11	Chronic lower respiratory diseases (J40-J47)	672	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	535	1.5	10.1
13	Influenza and pneumonia (J09-J18)	439	1.2	8.3
14	Viral hepatitis (B15-B19)	382	1.0	7.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	321	0.9	6.1
...	All other causes (Residual)	5,599	15.2	105.6
Black, both sexes, 55-64 years				
...	All causes	47,682	100.0	1,414.7
1	Malignant neoplasms (C00-C97)	14,762	31.0	438.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,289	25.8	364.6
3	Cerebrovascular diseases (I60-I69)	2,591	5.4	76.9
4	Diabetes mellitus (E10-E14)	2,486	5.2	73.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,599	3.4	47.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,391	2.9	41.3
7	Chronic lower respiratory diseases (J40-J47)	1,178	2.5	35.0
8	Septicemia (A40-A41)	1,004	2.1	29.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	995	2.1	29.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	867	1.8	25.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	752	1.6	22.3
12	Influenza and pneumonia (J09-J18)	573	1.2	17.0
13	Viral hepatitis (B15-B19)	409	0.9	12.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	319	0.7	9.5
15	Aortic aneurysm and dissection (I71)	232	0.5	6.9
...	All other causes (Residual)	6,235	13.1	185.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	51,384	100.0	2,791.4
1	Malignant neoplasms (C00-C97)	16,055	31.2	872.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,642	26.5	741.1
3	Cerebrovascular diseases (I60-I69)	3,257	6.3	176.9
4	Diabetes mellitus (E10-E14)	2,861	5.6	155.4
5	Chronic lower respiratory diseases (J40-J47)	1,897	3.7	103.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,841	3.6	100.0
7	Septicemia (A40-A41)	1,320	2.6	71.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	911	1.8	49.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	830	1.6	45.1
10	Influenza and pneumonia (J09-J18)	800	1.6	43.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	404	0.8	21.9
12	Alzheimer's disease (G30)	360	0.7	19.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	251	0.5	13.6
14	Aortic aneurysm and dissection (I71)	238	0.5	12.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	231	0.4	12.5
...	All other causes (Residual)	6,486	12.6	352.3
Black, both sexes, 75-84 years				
...	All causes	58,986	100.0	5,725.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,559	28.1	1,607.3
2	Malignant neoplasms (C00-C97)	14,277	24.2	1,385.8
3	Cerebrovascular diseases (I60-I69)	4,385	7.4	425.6
4	Diabetes mellitus (E10-E14)	3,068	5.2	297.8
5	Chronic lower respiratory diseases (J40-J47)	2,340	4.0	227.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,059	3.5	199.9
7	Alzheimer's disease (G30)	1,627	2.8	157.9
8	Septicemia (A40-A41)	1,519	2.6	147.4
9	Influenza and pneumonia (J09-J18)	1,312	2.2	127.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,095	1.9	106.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	764	1.3	74.2
12	Pneumonitis due to solids and liquids (J69)	438	0.7	42.5
13	Parkinson's disease (G20-G21)	361	0.6	35.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	310	0.5	30.1
15	Aortic aneurysm and dissection (I71)	271	0.5	26.3
...	All other causes (Residual)	8,601	14.6	834.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	49,194	100.0	12,281.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,773	32.1	3,937.8
2	Malignant neoplasms (C00-C97)	6,682	13.6	1,668.2
3	Cerebrovascular diseases (I60-I69)	3,992	8.1	996.6
4	Alzheimer's disease (G30)	2,724	5.5	680.1
5	Diabetes mellitus (E10-E14)	1,771	3.6	442.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,751	3.6	437.1
7	Influenza and pneumonia (J09-J18)	1,611	3.3	402.2
8	Chronic lower respiratory diseases (J40-J47)	1,355	2.8	338.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,192	2.4	297.6
10	Septicemia (A40-A41)	1,179	2.4	294.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	1.3	154.0
12	Pneumonitis due to solids and liquids (J69)	502	1.0	125.3
13	Atherosclerosis (I70)	280	0.6	69.9
14	Parkinson's disease (G20-G21)	264	0.5	65.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.4	49.4
...	All other causes (Residual)	9,303	18.9	2,322.5
Black, male, all ages				
...	All causes	148,309	100.0	775.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,669	24.1	186.5
2	Malignant neoplasms (C00-C97)	33,069	22.3	172.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,268	6.2	48.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,584	5.1	39.7
5	Cerebrovascular diseases (I60-I69)	7,549	5.1	39.5
6	Diabetes mellitus (E10-E14)	5,493	3.7	28.7
7	Chronic lower respiratory diseases (J40-J47)	4,207	2.8	22.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,186	2.8	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,772	2.5	19.7
10	Certain conditions originating in the perinatal period (P00-P96)	2,846	1.9	14.9
11	Septicemia (A40-A41)	2,835	1.9	14.8
12	Influenza and pneumonia (J09-J18)	2,498	1.7	13.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,979	1.3	10.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,682	1.1	8.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,606	1.1	8.4
...	All other causes (Residual)	24,066	16.2	125.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	621	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	30.4	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	14.3	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	9.3	4.2
4	Malignant neoplasms (C00-C97)	33	5.3	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.5	2.0
6	Influenza and pneumonia (J09-J18)	16	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.9	*
9	Septicemia (A40-A41)	11	1.8	*
10	Anemias (D50-D64)	4	0.6	*
10	Cerebrovascular diseases (I60-I69)	4	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	148	23.8	10.8
Black, male, 5-14 years				
...	All causes	816	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	34.2	8.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	11.8	2.9
3	Malignant neoplasms (C00-C97)	72	8.8	2.2
4	Chronic lower respiratory diseases (J40-J47)	41	5.0	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.3	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.7	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.8	*
9	Influenza and pneumonia (J09-J18)	13	1.6	*
10	Anemias (D50-D64)	8	1.0	*
11	Septicemia (A40-A41)	5	0.6	*
12	Meningitis (G00,G03)	4	0.5	*
13	Cerebrovascular diseases (I60-I69)	3	0.4	*
13	Diseases of appendix (K35-K38)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	172	21.1	5.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,751	100.0	168.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,918	50.7	85.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	22.8	38.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	6.1	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	4.1	6.9
5	Malignant neoplasms (C00-C97)	145	2.5	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	56	1.0	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.9	1.5
8	Legal intervention (Y35,Y89.0)	46	0.8	1.3
9	Chronic lower respiratory diseases (J40-J47)	44	0.8	1.3
10	Anemias (D50-D64)	37	0.6	1.1
11	Cerebrovascular diseases (I60-I69)	30	0.5	0.9
12	Septicemia (A40-A41)	22	0.4	0.6
12	Diabetes mellitus (E10-E14)	22	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
15	Influenza and pneumonia (J09-J18)	12	0.2	*
...	All other causes (Residual)	456	7.9	13.3
Black, male, 25-34 years				
...	All causes	6,648	100.0	240.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,283	34.3	82.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,369	20.6	49.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	632	9.5	22.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	409	6.2	14.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	376	5.7	13.6
6	Malignant neoplasms (C00-C97)	263	4.0	9.5
7	Diabetes mellitus (E10-E14)	116	1.7	4.2
8	Cerebrovascular diseases (I60-I69)	71	1.1	2.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.9	2.2
10	Chronic lower respiratory diseases (J40-J47)	58	0.9	2.1
11	Anemias (D50-D64)	56	0.8	2.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.8	1.9
13	Influenza and pneumonia (J09-J18)	51	0.8	1.8
14	Legal intervention (Y35,Y89.0)	43	0.6	1.6
15	Septicemia (A40-A41)	41	0.6	1.5
...	All other causes (Residual)	767	11.5	27.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	10,106	100.0	378.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,951	19.3	73.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,563	15.5	58.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,223	12.1	45.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,100	10.9	41.2
5	Malignant neoplasms (C00-C97)	907	9.0	34.0
6	Cerebrovascular diseases (I60-I69)	335	3.3	12.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	334	3.3	12.5
8	Diabetes mellitus (E10-E14)	314	3.1	11.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	1.8	6.6
10	Septicemia (A40-A41)	127	1.3	4.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	124	1.2	4.6
12	Influenza and pneumonia (J09-J18)	103	1.0	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.0	3.8
14	Chronic lower respiratory diseases (J40-J47)	99	1.0	3.7
15	Aortic aneurysm and dissection (I71)	87	0.9	3.3
...	All other causes (Residual)	1,561	15.4	58.5
Black, male, 45-54 years				
...	All causes	21,495	100.0	876.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,304	24.7	216.3
2	Malignant neoplasms (C00-C97)	4,365	20.3	178.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,949	9.1	79.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,563	7.3	63.8
5	Cerebrovascular diseases (I60-I69)	1,006	4.7	41.0
6	Diabetes mellitus (E10-E14)	795	3.7	32.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	623	2.9	25.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	560	2.6	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	2.0	17.7
10	Septicemia (A40-A41)	380	1.8	15.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	307	1.4	12.5
12	Chronic lower respiratory diseases (J40-J47)	304	1.4	12.4
13	Influenza and pneumonia (J09-J18)	271	1.3	11.1
14	Viral hepatitis (B15-B19)	254	1.2	10.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	245	1.1	10.0
...	All other causes (Residual)	3,134	14.6	127.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	28,127	100.0	1,870.8
1	Malignant neoplasms (C00-C97)	8,181	29.1	544.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,762	27.6	516.3
3	Cerebrovascular diseases (I60-I69)	1,500	5.3	99.8
4	Diabetes mellitus (E10-E14)	1,293	4.6	86.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,151	4.1	76.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	742	2.6	49.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	730	2.6	48.6
8	Chronic lower respiratory diseases (J40-J47)	672	2.4	44.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	606	2.2	40.3
10	Septicemia (A40-A41)	528	1.9	35.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	406	1.4	27.0
12	Influenza and pneumonia (J09-J18)	332	1.2	22.1
13	Viral hepatitis (B15-B19)	296	1.1	19.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	268	1.0	17.8
15	Aortic aneurysm and dissection (I71)	149	0.5	9.9
...	All other causes (Residual)	3,511	12.5	233.5
Black, male, 65-74 years				
...	All causes	27,672	100.0	3,604.9
1	Malignant neoplasms (C00-C97)	8,747	31.6	1,139.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,595	27.4	989.4
3	Cerebrovascular diseases (I60-I69)	1,714	6.2	223.3
4	Diabetes mellitus (E10-E14)	1,317	4.8	171.6
5	Chronic lower respiratory diseases (J40-J47)	1,061	3.8	138.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	877	3.2	114.2
7	Septicemia (A40-A41)	659	2.4	85.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	2.2	78.7
9	Influenza and pneumonia (J09-J18)	456	1.6	59.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	411	1.5	53.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	272	1.0	35.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	175	0.6	22.8
13	Alzheimer's disease (G30)	152	0.5	19.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.5	17.7
15	Aortic aneurysm and dissection (I71)	129	0.5	16.8
15	Pneumonitis due to solids and liquids (J69)	129	0.5	16.8
...	All other causes (Residual)	3,238	11.7	421.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	26,815	100.0	7,169.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,478	27.9	1,999.2
2	Malignant neoplasms (C00-C97)	7,245	27.0	1,936.9
3	Cerebrovascular diseases (I60-I69)	1,840	6.9	491.9
4	Chronic lower respiratory diseases (J40-J47)	1,296	4.8	346.5
5	Diabetes mellitus (E10-E14)	1,169	4.4	312.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	863	3.2	230.7
7	Influenza and pneumonia (J09-J18)	642	2.4	171.6
8	Septicemia (A40-A41)	609	2.3	162.8
9	Alzheimer's disease (G30)	539	2.0	144.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	436	1.6	116.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	1.5	106.4
12	Pneumonitis due to solids and liquids (J69)	244	0.9	65.2
13	Parkinson's disease (G20-G21)	201	0.7	53.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	142	0.5	38.0
15	Aortic aneurysm and dissection (I71)	118	0.4	31.5
...	All other causes (Residual)	3,595	13.4	961.1
Black, male, 85 years and over				
...	All causes	15,255	100.0	12,964.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,565	29.9	3,879.6
2	Malignant neoplasms (C00-C97)	3,103	20.3	2,637.1
3	Cerebrovascular diseases (I60-I69)	1,020	6.7	866.9
4	Chronic lower respiratory diseases (J40-J47)	606	4.0	515.0
5	Alzheimer's disease (G30)	589	3.9	500.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	581	3.8	493.8
7	Influenza and pneumonia (J09-J18)	560	3.7	475.9
8	Diabetes mellitus (E10-E14)	463	3.0	393.5
9	Septicemia (A40-A41)	390	2.6	331.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	289	1.9	245.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	1.5	198.0
12	Pneumonitis due to solids and liquids (J69)	189	1.2	160.6
13	Parkinson's disease (G20-G21)	117	0.8	99.4
14	Atherosclerosis (I70)	74	0.5	62.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	62.0
...	All other causes (Residual)	2,403	15.8	2,042.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,276	100.0	675.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,540	25.2	170.0
2	Malignant neoplasms (C00-C97)	30,980	21.9	148.2
3	Cerebrovascular diseases (I60-I69)	9,536	6.7	45.6
4	Diabetes mellitus (E10-E14)	6,966	4.9	33.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,620	3.3	22.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,291	3.0	20.5
7	Chronic lower respiratory diseases (J40-J47)	3,694	2.6	17.7
8	Septicemia (A40-A41)	3,462	2.5	16.6
9	Alzheimer's disease (G30)	3,459	2.4	16.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,661	1.9	12.7
11	Influenza and pneumonia (J09-J18)	2,657	1.9	12.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,284	1.6	10.9
13	Certain conditions originating in the perinatal period (P00-P96)	2,206	1.6	10.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,286	0.9	6.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	883	0.6	4.2
...	All other causes (Residual)	26,751	18.9	127.9
Black, female, 1-4 years				
...	All causes	516	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	26.4	10.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	15.3	6.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	11.8	4.6
4	Malignant neoplasms (C00-C97)	22	4.3	1.7
5	Chronic lower respiratory diseases (J40-J47)	17	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	15	2.9	*
8	Influenza and pneumonia (J09-J18)	13	2.5	*
9	Cerebrovascular diseases (I60-I69)	12	2.3	*
10	Anemias (D50-D64)	8	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	120	23.3	9.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	547	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	32.9	5.6
2	Malignant neoplasms (C00-C97)	71	13.0	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	8.0	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.3	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.1	0.9
6	Chronic lower respiratory diseases (J40-J47)	19	3.5	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.2	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Influenza and pneumonia (J09-J18)	8	1.5	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
13	Anemias (D50-D64)	5	0.9	*
14	Diabetes mellitus (E10-E14)	4	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	104	19.0	3.2
Black, female, 15-24 years				
...	All causes	1,629	100.0	48.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	440	27.0	13.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	295	18.1	8.9
3	Malignant neoplasms (C00-C97)	116	7.1	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	6.9	3.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	69	4.2	2.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	3.3	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	53	3.3	1.6
8	Diabetes mellitus (E10-E14)	34	2.1	1.0
9	Anemias (D50-D64)	30	1.8	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.8	0.9
11	Septicemia (A40-A41)	25	1.5	0.8
12	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.6
13	Cerebrovascular diseases (I60-I69)	16	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.9	*
15	Influenza and pneumonia (J09-J18)	11	0.7	*
...	All other causes (Residual)	308	18.9	9.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,014	100.0	102.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	13.6	13.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	12.1	12.4
3	Malignant neoplasms (C00-C97)	355	11.8	12.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	329	10.9	11.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	9.2	9.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	108	3.6	3.7
7	Diabetes mellitus (E10-E14)	97	3.2	3.3
8	Cerebrovascular diseases (I60-I69)	77	2.6	2.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	2.4	2.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.8	1.8
11	Septicemia (A40-A41)	49	1.6	1.7
12	Anemias (D50-D64)	46	1.5	1.6
13	Influenza and pneumonia (J09-J18)	43	1.4	1.5
14	Chronic lower respiratory diseases (J40-J47)	38	1.3	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	1.0
...	All other causes (Residual)	665	22.1	22.5
Black, female, 35-44 years				
...	All causes	6,884	100.0	229.1
1	Malignant neoplasms (C00-C97)	1,458	21.2	48.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,041	15.1	34.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	779	11.3	25.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	595	8.6	19.8
5	Cerebrovascular diseases (I60-I69)	320	4.6	10.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	3.9	8.9
7	Diabetes mellitus (E10-E14)	248	3.6	8.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.2	5.1
9	Septicemia (A40-A41)	147	2.1	4.9
10	Chronic lower respiratory diseases (J40-J47)	94	1.4	3.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	1.3	3.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	79	1.1	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.1	2.5
13	Influenza and pneumonia (J09-J18)	75	1.1	2.5
15	Anemias (D50-D64)	64	0.9	2.1
...	All other causes (Residual)	1,400	20.3	46.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	15,323	100.0	537.2
1	Malignant neoplasms (C00-C97)	4,453	29.1	156.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,053	19.9	107.0
3	Cerebrovascular diseases (I60-I69)	940	6.1	33.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	817	5.3	28.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	788	5.1	27.6
6	Diabetes mellitus (E10-E14)	638	4.2	22.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	388	2.5	13.6
8	Chronic lower respiratory diseases (J40-J47)	368	2.4	12.9
9	Septicemia (A40-A41)	341	2.2	12.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	293	1.9	10.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	228	1.5	8.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	1.2	6.3
13	Influenza and pneumonia (J09-J18)	168	1.1	5.9
14	Viral hepatitis (B15-B19)	128	0.8	4.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	0.5	2.7
...	All other causes (Residual)	2,465	16.1	86.4
Black, female, 55-64 years				
...	All causes	19,555	100.0	1,047.4
1	Malignant neoplasms (C00-C97)	6,581	33.7	352.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,527	23.2	242.5
3	Diabetes mellitus (E10-E14)	1,193	6.1	63.9
4	Cerebrovascular diseases (I60-I69)	1,091	5.6	58.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	661	3.4	35.4
6	Chronic lower respiratory diseases (J40-J47)	506	2.6	27.1
7	Septicemia (A40-A41)	476	2.4	25.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	2.3	24.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	346	1.8	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	261	1.3	14.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	253	1.3	13.6
12	Influenza and pneumonia (J09-J18)	241	1.2	12.9
13	Viral hepatitis (B15-B19)	113	0.6	6.1
14	Aortic aneurysm and dissection (I71)	83	0.4	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	3.7
...	All other causes (Residual)	2,706	13.8	144.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	23,712	100.0	2,209.5
1	Malignant neoplasms (C00-C97)	7,308	30.8	681.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,047	25.5	563.5
3	Diabetes mellitus (E10-E14)	1,544	6.5	143.9
4	Cerebrovascular diseases (I60-I69)	1,543	6.5	143.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	964	4.1	89.8
6	Chronic lower respiratory diseases (J40-J47)	836	3.5	77.9
7	Septicemia (A40-A41)	661	2.8	61.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	419	1.8	39.0
9	Influenza and pneumonia (J09-J18)	344	1.5	32.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	1.3	28.6
11	Alzheimer's disease (G30)	208	0.9	19.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	132	0.6	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	115	0.5	10.7
14	Aortic aneurysm and dissection (I71)	109	0.5	10.2
15	Pneumonitis due to solids and liquids (J69)	76	0.3	7.1
...	All other causes (Residual)	3,099	13.1	288.8
Black, female, 75-84 years				
...	All causes	32,171	100.0	4,902.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,081	28.2	1,384.0
2	Malignant neoplasms (C00-C97)	7,032	21.9	1,071.7
3	Cerebrovascular diseases (I60-I69)	2,545	7.9	387.9
4	Diabetes mellitus (E10-E14)	1,899	5.9	289.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,196	3.7	182.3
6	Alzheimer's disease (G30)	1,088	3.4	165.8
7	Chronic lower respiratory diseases (J40-J47)	1,044	3.2	159.1
8	Septicemia (A40-A41)	910	2.8	138.7
9	Influenza and pneumonia (J09-J18)	670	2.1	102.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	659	2.0	100.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	1.1	55.8
12	Pneumonitis due to solids and liquids (J69)	194	0.6	29.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.5	25.6
14	Parkinson's disease (G20-G21)	160	0.5	24.4
15	Aortic aneurysm and dissection (I71)	153	0.5	23.3
...	All other causes (Residual)	5,006	15.6	762.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	33,939	100.0	11,997.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,208	33.0	3,962.0
2	Malignant neoplasms (C00-C97)	3,579	10.5	1,265.2
3	Cerebrovascular diseases (I60-I69)	2,972	8.8	1,050.6
4	Alzheimer's disease (G30)	2,135	6.3	754.7
5	Diabetes mellitus (E10-E14)	1,308	3.9	462.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,170	3.4	413.6
7	Influenza and pneumonia (J09-J18)	1,051	3.1	371.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	903	2.7	319.2
9	Septicemia (A40-A41)	789	2.3	278.9
10	Chronic lower respiratory diseases (J40-J47)	749	2.2	264.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	1.1	135.7
12	Pneumonitis due to solids and liquids (J69)	313	0.9	110.6
13	Atherosclerosis (I70)	206	0.6	72.8
14	Parkinson's disease (G20-G21)	147	0.4	52.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.4	44.2
...	All other causes (Residual)	6,900	20.3	2,439.1
American Indian, both sexes, all ages				
...	All causes	14,367	100.0	444.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,648	18.4	81.8
2	Malignant neoplasms (C00-C97)	2,561	17.8	79.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	11.8	52.6
4	Diabetes mellitus (E10-E14)	790	5.5	24.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	709	4.9	21.9
6	Chronic lower respiratory diseases (J40-J47)	611	4.3	18.9
7	Cerebrovascular diseases (I60-I69)	586	4.1	18.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.7	12.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	2.0	9.0
10	Influenza and pneumonia (J09-J18)	280	1.9	8.7
11	Septicemia (A40-A41)	230	1.6	7.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	1.5	6.8
13	Alzheimer's disease (G30)	191	1.3	5.9
14	Certain conditions originating in the perinatal period (P00-P96)	158	1.1	4.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	0.9	3.8
...	All other causes (Residual)	2,874	20.0	88.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	97	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	44.3	24.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	7.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
5	Influenza and pneumonia (J09-J18)	2	2.1	*
5	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
7	Malignant neoplasms (C00-C97)	1	1.0	*
7	Cerebrovascular diseases (I60-I69)	1	1.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.0	*
...	All other causes (Residual)	24	24.7	13.6
American Indian, both sexes, 5-14 years				
...	All causes	94	100.0	18.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	48.9	8.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	9.6	*
3	Malignant neoplasms (C00-C97)	6	6.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.2	*
6	Septicemia (A40-A41)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
...	All other causes (Residual)	17	18.1	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	622	100.0	103.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	51.3	53.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	19.6	20.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	9.2	9.5
4	Malignant neoplasms (C00-C97)	22	3.5	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.0	*
7	Influenza and pneumonia (J09-J18)	4	0.6	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.6	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
10	Septicemia (A40-A41)	2	0.3	*
10	Cerebrovascular diseases (I60-I69)	2	0.3	*
10	Aortic aneurysm and dissection (I71)	2	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
10	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Anemias (D50-D64)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	60	9.6	10.0
American Indian, both sexes, 25-34 years				
...	All causes	731	100.0	146.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	42.3	61.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	13.4	19.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	8.9	13.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	8.2	12.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	37	5.1	7.4
6	Malignant neoplasms (C00-C97)	25	3.4	5.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.0	*
8	Septicemia (A40-A41)	6	0.8	*
8	Diabetes mellitus (E10-E14)	6	0.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
11	Influenza and pneumonia (J09-J18)	4	0.5	*
12	Cerebrovascular diseases (I60-I69)	3	0.4	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	97	13.3	19.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,213	100.0	264.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	26.4	69.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	145	12.0	31.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	11.3	29.9
4	Malignant neoplasms (C00-C97)	83	6.8	18.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	6.1	16.2
6	Diabetes mellitus (E10-E14)	45	3.7	9.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	3.5	9.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	31	2.6	6.8
9	Cerebrovascular diseases (I60-I69)	28	2.3	6.1
10	Viral hepatitis (B15-B19)	19	1.6	*
11	Influenza and pneumonia (J09-J18)	18	1.5	*
12	Septicemia (A40-A41)	12	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.6	*
...	All other causes (Residual)	230	19.0	50.2
American Indian, both sexes, 45-54 years				
...	All causes	1,958	100.0	456.4
1	Malignant neoplasms (C00-C97)	330	16.9	76.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	16.1	73.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	14.3	65.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	259	13.2	60.4
5	Diabetes mellitus (E10-E14)	92	4.7	21.4
6	Cerebrovascular diseases (I60-I69)	51	2.6	11.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.6	11.7
8	Chronic lower respiratory diseases (J40-J47)	48	2.5	11.2
9	Septicemia (A40-A41)	46	2.3	10.7
10	Viral hepatitis (B15-B19)	33	1.7	7.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.6	7.5
12	Influenza and pneumonia (J09-J18)	28	1.4	6.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.2	5.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	1.2	5.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.4	*
...	All other causes (Residual)	332	17.0	77.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,367	100.0	858.3
1	Malignant neoplasms (C00-C97)	623	26.3	225.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	468	19.8	169.7
3	Diabetes mellitus (E10-E14)	167	7.1	60.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	167	7.1	60.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	6.3	54.4
6	Chronic lower respiratory diseases (J40-J47)	100	4.2	36.3
7	Cerebrovascular diseases (I60-I69)	83	3.5	30.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	2.4	20.7
9	Septicemia (A40-A41)	44	1.9	16.0
10	Viral hepatitis (B15-B19)	39	1.6	14.1
11	Influenza and pneumonia (J09-J18)	35	1.5	12.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	1.0	8.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.3	*
...	All other causes (Residual)	378	16.0	137.1
American Indian, both sexes, 65-74 years				
...	All causes	2,536	100.0	1,853.8
1	Malignant neoplasms (C00-C97)	685	27.0	500.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	543	21.4	396.9
3	Diabetes mellitus (E10-E14)	208	8.2	152.0
4	Chronic lower respiratory diseases (J40-J47)	173	6.8	126.5
5	Cerebrovascular diseases (I60-I69)	114	4.5	83.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	3.7	68.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.9	54.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	66	2.6	48.2
9	Septicemia (A40-A41)	49	1.9	35.8
10	Influenza and pneumonia (J09-J18)	39	1.5	28.5
11	Alzheimer's disease (G30)	24	0.9	17.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.8	14.6
13	Aortic aneurysm and dissection (I71)	15	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.6	*
15	Pneumonitis due to solids and liquids (J69)	13	0.5	*
...	All other causes (Residual)	406	16.0	296.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,546	100.0	3,760.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	24.7	929.0
2	Malignant neoplasms (C00-C97)	561	22.0	828.6
3	Chronic lower respiratory diseases (J40-J47)	185	7.3	273.2
4	Diabetes mellitus (E10-E14)	173	6.8	255.5
5	Cerebrovascular diseases (I60-I69)	154	6.0	227.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	3.0	113.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	2.7	100.4
8	Influenza and pneumonia (J09-J18)	67	2.6	99.0
9	Alzheimer's disease (G30)	65	2.6	96.0
10	Septicemia (A40-A41)	41	1.6	60.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.5	54.6
12	Pneumonitis due to solids and liquids (J69)	21	0.8	31.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.8	29.5
14	Parkinson's disease (G20-G21)	17	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.6	*
...	All other causes (Residual)	416	16.3	614.4
American Indian, both sexes, 85 years and over				
...	All causes	1,771	100.0	6,708.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	466	26.3	1,765.1
2	Malignant neoplasms (C00-C97)	224	12.6	848.5
3	Cerebrovascular diseases (I60-I69)	148	8.4	560.6
4	Diabetes mellitus (E10-E14)	99	5.6	375.0
5	Alzheimer's disease (G30)	96	5.4	363.6
6	Chronic lower respiratory diseases (J40-J47)	86	4.9	325.7
7	Influenza and pneumonia (J09-J18)	75	4.2	284.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	3.0	204.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	1.9	125.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.5	98.5
11	Pneumonitis due to solids and liquids (J69)	25	1.4	94.7
12	Septicemia (A40-A41)	24	1.4	90.9
12	Parkinson's disease (G20-G21)	24	1.4	90.9
14	Aortic aneurysm and dissection (I71)	12	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.6	*
...	All other causes (Residual)	368	20.8	1,393.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,885	100.0	488.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,520	19.3	94.1
2	Malignant neoplasms (C00-C97)	1,345	17.1	83.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,129	14.3	69.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	415	5.3	25.7
5	Diabetes mellitus (E10-E14)	381	4.8	23.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	310	3.9	19.2
7	Chronic lower respiratory diseases (J40-J47)	299	3.8	18.5
8	Cerebrovascular diseases (I60-I69)	267	3.4	16.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	2.1	10.1
10	Influenza and pneumonia (J09-J18)	150	1.9	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.6	7.8
12	Septicemia (A40-A41)	102	1.3	6.3
13	Certain conditions originating in the perinatal period (P00-P96)	83	1.1	5.1
14	Alzheimer's disease (G30)	69	0.9	4.3
15	Viral hepatitis (B15-B19)	66	0.8	4.1
...	All other causes (Residual)	1,460	18.5	90.4
American Indian, male, 1-4 years				
...	All causes	57	100.0	63.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	42.1	26.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	8.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	8.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.3	*
5	Influenza and pneumonia (J09-J18)	2	3.5	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.8	*
...	All other causes (Residual)	14	24.6	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	61	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	49.2	11.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	9.8	*
3	Malignant neoplasms (C00-C97)	4	6.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.9	*
6	Septicemia (A40-A41)	1	1.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.6	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	*
...	All other causes (Residual)	11	18.0	*
American Indian, male, 15-24 years				
...	All causes	441	100.0	143.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	50.8	73.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	22.4	32.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	10.2	14.7
4	Malignant neoplasms (C00-C97)	16	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.9	*
7	Aortic aneurysm and dissection (I71)	2	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
9	Septicemia (A40-A41)	1	0.2	*
9	Anemias (D50-D64)	1	0.2	*
9	Cerebrovascular diseases (I60-I69)	1	0.2	*
9	Influenza and pneumonia (J09-J18)	1	0.2	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
9	Pneumonitis due to solids and liquids (J69)	1	0.2	*
9	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	37	8.4	12.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	514	100.0	198.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	41.6	82.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	15.8	31.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	10.1	20.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	8.4	16.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	22	4.3	8.5
6	Malignant neoplasms (C00-C97)	18	3.5	*
7	Septicemia (A40-A41)	4	0.8	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
9	Diabetes mellitus (E10-E14)	3	0.6	*
9	Influenza and pneumonia (J09-J18)	3	0.6	*
11	Cerebrovascular diseases (I60-I69)	2	0.4	*
11	Aortic aneurysm and dissection (I71)	2	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
14	Salmonella infections (A01-A02)	1	0.2	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	59	11.5	22.8
American Indian, male, 35-44 years				
...	All causes	768	100.0	332.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	28.1	93.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	11.7	39.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	90	11.7	39.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	7.7	25.5
5	Malignant neoplasms (C00-C97)	37	4.8	16.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	4.2	13.9
7	Diabetes mellitus (E10-E14)	29	3.8	12.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	21	2.7	9.1
9	Cerebrovascular diseases (I60-I69)	14	1.8	*
10	Viral hepatitis (B15-B19)	12	1.6	*
10	Influenza and pneumonia (J09-J18)	12	1.6	*
12	Septicemia (A40-A41)	7	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
...	All other causes (Residual)	133	17.3	57.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,193	100.0	573.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	19.6	112.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	14.8	85.0
3	Malignant neoplasms (C00-C97)	163	13.7	78.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	153	12.8	73.5
5	Diabetes mellitus (E10-E14)	60	5.0	28.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.1	17.8
7	Cerebrovascular diseases (I60-I69)	29	2.4	13.9
8	Septicemia (A40-A41)	24	2.0	11.5
8	Chronic lower respiratory diseases (J40-J47)	24	2.0	11.5
10	Influenza and pneumonia (J09-J18)	23	1.9	11.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.8	10.6
12	Viral hepatitis (B15-B19)	20	1.7	9.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.4	*
...	All other causes (Residual)	190	15.9	91.3
American Indian, male, 55-64 years				
...	All causes	1,372	100.0	1,037.0
1	Malignant neoplasms (C00-C97)	350	25.5	264.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	312	22.7	235.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	6.9	71.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	93	6.8	70.3
5	Diabetes mellitus (E10-E14)	92	6.7	69.5
6	Cerebrovascular diseases (I60-I69)	49	3.6	37.0
6	Chronic lower respiratory diseases (J40-J47)	49	3.6	37.0
8	Influenza and pneumonia (J09-J18)	27	2.0	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.9	19.7
10	Viral hepatitis (B15-B19)	23	1.7	17.4
11	Septicemia (A40-A41)	19	1.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.4	*
...	All other causes (Residual)	196	14.3	148.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,357	100.0	2,131.7
1	Malignant neoplasms (C00-C97)	360	26.5	565.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	24.5	521.5
3	Diabetes mellitus (E10-E14)	104	7.7	163.4
4	Chronic lower respiratory diseases (J40-J47)	92	6.8	144.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	4.6	97.4
6	Cerebrovascular diseases (I60-I69)	53	3.9	83.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	2.8	59.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.6	55.0
9	Septicemia (A40-A41)	22	1.6	34.6
9	Influenza and pneumonia (J09-J18)	22	1.6	34.6
11	Aortic aneurysm and dissection (I71)	12	0.9	*
12	Parkinson's disease (G20-G21)	9	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.7	*
14	Viral hepatitis (B15-B19)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
15	Atherosclerosis (I70)	7	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
...	All other causes (Residual)	178	13.1	279.6
American Indian, male, 75-84 years				
...	All causes	1,214	100.0	4,193.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	26.9	1,129.5
2	Malignant neoplasms (C00-C97)	285	23.5	984.5
3	Chronic lower respiratory diseases (J40-J47)	86	7.1	297.1
4	Cerebrovascular diseases (I60-I69)	77	6.3	266.0
5	Diabetes mellitus (E10-E14)	59	4.9	203.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	4.1	172.7
7	Influenza and pneumonia (J09-J18)	32	2.6	110.5
8	Alzheimer's disease (G30)	25	2.1	86.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.9	79.4
10	Septicemia (A40-A41)	16	1.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.0	*
13	Parkinson's disease (G20-G21)	11	0.9	*
14	Pneumonitis due to solids and liquids (J69)	10	0.8	*
15	Aortic aneurysm and dissection (I71)	8	0.7	*
...	All other causes (Residual)	180	14.8	621.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	667	100.0	7,638.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	24.9	1,901.1
2	Malignant neoplasms (C00-C97)	111	16.6	1,271.2
3	Cerebrovascular diseases (I60-I69)	42	6.3	481.0
4	Chronic lower respiratory diseases (J40-J47)	37	5.5	423.7
5	Diabetes mellitus (E10-E14)	34	5.1	389.4
5	Alzheimer's disease (G30)	34	5.1	389.4
7	Influenza and pneumonia (J09-J18)	23	3.4	263.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	3.3	251.9
9	Pneumonitis due to solids and liquids (J69)	15	2.2	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.1	*
11	Parkinson's disease (G20-G21)	11	1.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.2	*
13	Septicemia (A40-A41)	6	0.9	*
14	Aortic aneurysm and dissection (I71)	5	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
...	All other causes (Residual)	135	20.2	1,546.0
American Indian, female, all ages				
...	All causes	6,482	100.0	400.0
1	Malignant neoplasms (C00-C97)	1,216	18.8	75.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,128	17.4	69.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	8.8	35.3
4	Diabetes mellitus (E10-E14)	409	6.3	25.2
5	Cerebrovascular diseases (I60-I69)	319	4.9	19.7
6	Chronic lower respiratory diseases (J40-J47)	312	4.8	19.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	294	4.5	18.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.6	10.2
9	Influenza and pneumonia (J09-J18)	130	2.0	8.0
10	Septicemia (A40-A41)	128	2.0	7.9
11	Alzheimer's disease (G30)	122	1.9	7.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	1.3	5.1
13	Certain conditions originating in the perinatal period (P00-P96)	75	1.2	4.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.0	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.0	3.8
...	All other causes (Residual)	1,405	21.7	86.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	40	100.0	46.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	47.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.0	*
5	Malignant neoplasms (C00-C97)	1	2.5	*
5	Cerebrovascular diseases (I60-I69)	1	2.5	*
5	Acute bronchitis and bronchiolitis (J20-J21)	1	2.5	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.5	*
...	All other causes (Residual)	10	25.0	*
American Indian, female, 5-14 years				
...	All causes	33	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	48.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	9.1	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	9.1	*
4	Malignant neoplasms (C00-C97)	2	6.1	*
5	Septicemia (A40-A41)	1	3.0	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.0	*
5	Chronic lower respiratory diseases (J40-J47)	1	3.0	*
...	All other causes (Residual)	6	18.2	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	181	100.0	61.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	52.5	32.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	12.7	7.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	6.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	3.9	*
5	Malignant neoplasms (C00-C97)	6	3.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.2	*
7	Influenza and pneumonia (J09-J18)	3	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
9	Septicemia (A40-A41)	1	0.6	*
9	Cerebrovascular diseases (I60-I69)	1	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
9	Pneumonitis due to solids and liquids (J69)	1	0.6	*
9	Diseases of appendix (K35-K38)	1	0.6	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
...	All other causes (Residual)	23	12.7	7.8
American Indian, female, 25-34 years				
...	All causes	217	100.0	90.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	43.8	39.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	7.8	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	7.8	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	15	6.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	6.0	*
6	Malignant neoplasms (C00-C97)	7	3.2	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	7	3.2	*
8	Diabetes mellitus (E10-E14)	3	1.4	*
9	Septicemia (A40-A41)	2	0.9	*
9	Pneumonitis due to solids and liquids (J69)	2	0.9	*
11	Scarlet fever and erysipelas (A38,A46)	1	0.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
11	Cerebrovascular diseases (I60-I69)	1	0.5	*
11	Influenza and pneumonia (J09-J18)	1	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	*
...	All other causes (Residual)	32	14.7	13.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	445	100.0	196.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	23.4	45.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	55	12.4	24.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	10.6	20.7
4	Malignant neoplasms (C00-C97)	46	10.3	20.3
5	Diabetes mellitus (E10-E14)	16	3.6	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.4	*
7	Cerebrovascular diseases (I60-I69)	14	3.1	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	2.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	2.2	*
10	Viral hepatitis (B15-B19)	7	1.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.6	*
12	Influenza and pneumonia (J09-J18)	6	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.3	*
14	Septicemia (A40-A41)	5	1.1	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
...	All other causes (Residual)	90	20.2	39.6
American Indian, female, 45-54 years				
...	All causes	765	100.0	346.4
1	Malignant neoplasms (C00-C97)	167	21.8	75.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	106	13.9	48.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	13.5	46.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	10.6	36.7
5	Diabetes mellitus (E10-E14)	32	4.2	14.5
6	Chronic lower respiratory diseases (J40-J47)	24	3.1	10.9
7	Septicemia (A40-A41)	22	2.9	10.0
7	Cerebrovascular diseases (I60-I69)	22	2.9	10.0
9	Viral hepatitis (B15-B19)	13	1.7	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.3	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.9	*
14	Influenza and pneumonia (J09-J18)	5	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	139	18.2	62.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	995	100.0	693.5
1	Malignant neoplasms (C00-C97)	273	27.4	190.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	15.7	108.7
3	Diabetes mellitus (E10-E14)	75	7.5	52.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	74	7.4	51.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	5.5	38.3
6	Chronic lower respiratory diseases (J40-J47)	51	5.1	35.5
7	Cerebrovascular diseases (I60-I69)	34	3.4	23.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	3.1	21.6
9	Septicemia (A40-A41)	25	2.5	17.4
10	Viral hepatitis (B15-B19)	16	1.6	*
11	Influenza and pneumonia (J09-J18)	8	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.6	*
14	Nutritional deficiencies (E40-E64)	4	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.3	*
15	Anemias (D50-D64)	3	0.3	*
15	Aortic aneurysm and dissection (I71)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	166	16.7	115.7
American Indian, female, 65-74 years				
...	All causes	1,179	100.0	1,611.9
1	Malignant neoplasms (C00-C97)	325	27.6	444.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	17.9	288.5
3	Diabetes mellitus (E10-E14)	104	8.8	142.2
4	Chronic lower respiratory diseases (J40-J47)	81	6.9	110.7
5	Cerebrovascular diseases (I60-I69)	61	5.2	83.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	3.3	53.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.6	42.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.4	38.3
9	Septicemia (A40-A41)	27	2.3	36.9
10	Alzheimer's disease (G30)	19	1.6	*
11	Influenza and pneumonia (J09-J18)	17	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.9	*
13	Pneumonitis due to solids and liquids (J69)	10	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
...	All other causes (Residual)	204	17.3	278.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,332	100.0	3,436.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	22.7	779.2
2	Malignant neoplasms (C00-C97)	276	20.7	712.1
3	Diabetes mellitus (E10-E14)	114	8.6	294.1
4	Chronic lower respiratory diseases (J40-J47)	99	7.4	255.4
5	Cerebrovascular diseases (I60-I69)	77	5.8	198.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	3.4	116.1
7	Alzheimer's disease (G30)	40	3.0	103.2
8	Influenza and pneumonia (J09-J18)	35	2.6	90.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.0	69.7
10	Septicemia (A40-A41)	25	1.9	64.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.8	61.9
12	Pneumonitis due to solids and liquids (J69)	11	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.6	*
15	Parkinson's disease (G20-G21)	6	0.5	*
15	Atherosclerosis (I70)	6	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
...	All other causes (Residual)	223	16.7	575.4
American Indian, female, 85 years and over				
...	All causes	1,104	100.0	6,248.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	300	27.2	1,697.9
2	Malignant neoplasms (C00-C97)	113	10.2	639.5
3	Cerebrovascular diseases (I60-I69)	106	9.6	599.9
4	Diabetes mellitus (E10-E14)	65	5.9	367.9
5	Alzheimer's disease (G30)	62	5.6	350.9
6	Influenza and pneumonia (J09-J18)	52	4.7	294.3
7	Chronic lower respiratory diseases (J40-J47)	49	4.4	277.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.9	181.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.7	*
10	Septicemia (A40-A41)	18	1.6	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.6	*
12	Parkinson's disease (G20-G21)	13	1.2	*
13	Pneumonitis due to solids and liquids (J69)	10	0.9	*
14	Aortic aneurysm and dissection (I71)	7	0.6	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.6	*
...	All other causes (Residual)	233	21.1	1,318.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	45,609	100.0	308.7
1	Malignant neoplasms (C00-C97)	12,326	27.0	83.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,574	23.2	71.6
3	Cerebrovascular diseases (I60-I69)	3,586	7.9	24.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,194	4.8	14.9
5	Diabetes mellitus (E10-E14)	1,743	3.8	11.8
6	Influenza and pneumonia (J09-J18)	1,335	2.9	9.0
7	Chronic lower respiratory diseases (J40-J47)	1,331	2.9	9.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	900	2.0	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	893	2.0	6.0
10	Alzheimer's disease (G30)	748	1.6	5.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	634	1.4	4.3
12	Septicemia (A40-A41)	551	1.2	3.7
13	Certain conditions originating in the perinatal period (P00-P96)	499	1.1	3.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	408	0.9	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	0.8	2.4
...	All other causes (Residual)	7,530	16.5	51.0
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	182	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	28.0	6.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	12.6	2.7
3	Malignant neoplasms (C00-C97)	21	11.5	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	6.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.9	*
6	Influenza and pneumonia (J09-J18)	5	2.7	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
8	Septicemia (A40-A41)	2	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
10	Salmonella infections (A01-A02)	1	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
10	Meningitis (G00,G03)	1	0.5	*
10	Cerebrovascular diseases (I60-I69)	1	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	49	26.9	5.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	210	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	30.5	3.4
2	Malignant neoplasms (C00-C97)	34	16.2	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	8.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.2	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.8	*
6	Cerebrovascular diseases (I60-I69)	7	3.3	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.3	*
8	Influenza and pneumonia (J09-J18)	5	2.4	*
9	Septicemia (A40-A41)	2	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.0	*
14	Meningococcal infection (A39)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Meningitis (G00,G03)	1	0.5	*
14	Pneumonitis due to solids and liquids (J69)	1	0.5	*
14	Hernia (K40-K46)	1	0.5	*
...	All other causes (Residual)	42	20.0	2.2
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	843	100.0	43.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	41.3	17.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	20.2	8.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	10.4	4.5
4	Malignant neoplasms (C00-C97)	74	8.8	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.7	*
7	Cerebrovascular diseases (I60-I69)	9	1.1	*
8	Influenza and pneumonia (J09-J18)	7	0.8	*
9	Septicemia (A40-A41)	5	0.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
10	Diabetes mellitus (E10-E14)	3	0.4	*
10	Meningitis (G00,G03)	3	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
10	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	2	0.2	*
...	All other causes (Residual)	93	11.0	4.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	998	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	24.3	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	17.5	6.8
3	Malignant neoplasms (C00-C97)	160	16.0	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	9.3	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	8.3	3.2
6	Cerebrovascular diseases (I60-I69)	22	2.2	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	18	1.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.4	*
9	Diabetes mellitus (E10-E14)	12	1.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.9	*
12	Influenza and pneumonia (J09-J18)	8	0.8	*
13	Septicemia (A40-A41)	7	0.7	*
14	Aortic aneurysm and dissection (I71)	6	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	127	12.7	5.0
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,832	100.0	71.5
1	Malignant neoplasms (C00-C97)	551	30.1	21.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	16.6	11.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	13.2	9.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	10.4	7.5
5	Cerebrovascular diseases (I60-I69)	94	5.1	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	3.6	2.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	2.1	1.5
8	Diabetes mellitus (E10-E14)	33	1.8	1.3
9	Viral hepatitis (B15-B19)	18	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.9	*
11	Septicemia (A40-A41)	16	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	15	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.5	*
...	All other causes (Residual)	212	11.6	8.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,677	100.0	180.2
1	Malignant neoplasms (C00-C97)	1,462	39.8	71.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	627	17.1	30.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	6.8	12.3
4	Cerebrovascular diseases (I60-I69)	249	6.8	12.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	4.0	7.3
6	Diabetes mellitus (E10-E14)	105	2.9	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	93	2.5	4.6
8	Viral hepatitis (B15-B19)	55	1.5	2.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	1.2	2.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.1	2.0
11	Influenza and pneumonia (J09-J18)	39	1.1	1.9
12	Chronic lower respiratory diseases (J40-J47)	38	1.0	1.9
13	Septicemia (A40-A41)	36	1.0	1.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.7	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.6	1.1
15	Aortic aneurysm and dissection (I71)	23	0.6	1.1
...	All other causes (Residual)	417	11.3	20.4
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	5,774	100.0	418.8
1	Malignant neoplasms (C00-C97)	2,413	41.8	175.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	20.5	86.0
3	Cerebrovascular diseases (I60-I69)	388	6.7	28.1
4	Diabetes mellitus (E10-E14)	273	4.7	19.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	3.8	16.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.7	7.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	1.7	7.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.6	6.6
9	Chronic lower respiratory diseases (J40-J47)	87	1.5	6.3
10	Viral hepatitis (B15-B19)	78	1.4	5.7
11	Septicemia (A40-A41)	63	1.1	4.6
12	Influenza and pneumonia (J09-J18)	61	1.1	4.4
13	Aortic aneurysm and dissection (I71)	52	0.9	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.7	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	0.7	3.1
...	All other causes (Residual)	581	10.1	42.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,941	100.0	1,046.6
1	Malignant neoplasms (C00-C97)	2,900	36.5	382.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,804	22.7	237.8
3	Cerebrovascular diseases (I60-I69)	613	7.7	80.8
4	Diabetes mellitus (E10-E14)	391	4.9	51.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	2.8	29.0
6	Chronic lower respiratory diseases (J40-J47)	205	2.6	27.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	2.3	24.0
8	Influenza and pneumonia (J09-J18)	157	2.0	20.7
9	Septicemia (A40-A41)	124	1.6	16.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	1.2	12.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.9	9.6
12	Viral hepatitis (B15-B19)	71	0.9	9.4
13	Aortic aneurysm and dissection (I71)	68	0.9	9.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.7	6.9
15	Parkinson's disease (G20-G21)	48	0.6	6.3
...	All other causes (Residual)	940	11.8	123.9
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	12,028	100.0	2,904.2
1	Malignant neoplasms (C00-C97)	3,185	26.5	769.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,028	25.2	731.1
3	Cerebrovascular diseases (I60-I69)	1,100	9.1	265.6
4	Diabetes mellitus (E10-E14)	537	4.5	129.7
5	Chronic lower respiratory diseases (J40-J47)	530	4.4	128.0
6	Influenza and pneumonia (J09-J18)	406	3.4	98.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	311	2.6	75.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	302	2.5	72.9
9	Alzheimer's disease (G30)	202	1.7	48.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	1.6	47.6
11	Septicemia (A40-A41)	159	1.3	38.4
12	Parkinson's disease (G20-G21)	153	1.3	36.9
13	Aortic aneurysm and dissection (I71)	104	0.9	25.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.8	22.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	82	0.7	19.8
...	All other causes (Residual)	1,641	13.6	396.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	11,161	100.0	7,929.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,471	31.1	2,465.9
2	Malignant neoplasms (C00-C97)	1,524	13.7	1,082.7
3	Cerebrovascular diseases (I60-I69)	1,101	9.9	782.2
4	Influenza and pneumonia (J09-J18)	632	5.7	449.0
5	Alzheimer's disease (G30)	500	4.5	355.2
6	Chronic lower respiratory diseases (J40-J47)	440	3.9	312.6
7	Diabetes mellitus (E10-E14)	389	3.5	276.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	270	2.4	191.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	2.2	177.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	2.0	156.3
11	Pneumonitis due to solids and liquids (J69)	132	1.2	93.8
12	Septicemia (A40-A41)	131	1.2	93.1
13	Parkinson's disease (G20-G21)	126	1.1	89.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.7	52.6
15	Aortic aneurysm and dissection (I71)	58	0.5	41.2
...	All other causes (Residual)	1,843	16.5	1,309.3
Asian or Pacific Islander, male, all ages				
...	All causes	23,823	100.0	331.4
1	Malignant neoplasms (C00-C97)	6,394	26.8	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,724	24.0	79.6
3	Cerebrovascular diseases (I60-I69)	1,581	6.6	22.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,371	5.8	19.1
5	Diabetes mellitus (E10-E14)	860	3.6	12.0
6	Chronic lower respiratory diseases (J40-J47)	795	3.3	11.1
7	Influenza and pneumonia (J09-J18)	703	3.0	9.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	628	2.6	8.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	2.0	6.6
10	Septicemia (A40-A41)	302	1.3	4.2
11	Certain conditions originating in the perinatal period (P00-P96)	290	1.2	4.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	1.2	4.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	253	1.1	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	250	1.0	3.5
15	Alzheimer's disease (G30)	245	1.0	3.4
...	All other causes (Residual)	3,665	15.4	51.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	109	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	33.0	8.3
2	Malignant neoplasms (C00-C97)	12	11.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	11.0	*
4	Influenza and pneumonia (J09-J18)	5	4.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.6	*
6	Chronic lower respiratory diseases (J40-J47)	3	2.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	1.8	*
9	Salmonella infections (A01-A02)	1	0.9	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	0.9	*
...	All other causes (Residual)	30	27.5	7.0
Asian or Pacific Islander, male, 5-14 years				
...	All causes	118	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	31.4	3.8
2	Malignant neoplasms (C00-C97)	19	16.1	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	7.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	5.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.2	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	3.4	*
7	Influenza and pneumonia (J09-J18)	3	2.5	*
8	Cerebrovascular diseases (I60-I69)	2	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.7	*
11	Meningococcal infection (A39)	1	0.8	*
11	Septicemia (A40-A41)	1	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Meningitis (G00,G03)	1	0.8	*
11	Pneumonitis due to solids and liquids (J69)	1	0.8	*
...	All other causes (Residual)	23	19.5	2.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	611	100.0	61.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	41.1	25.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	134	21.9	13.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	12.1	7.4
4	Malignant neoplasms (C00-C97)	45	7.4	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.5	*
6	Congenital malformations, deformations and chromosomal abnormalities	9	1.5	*
7	Cerebrovascular diseases (I60-I69)	6	1.0	*
8	Influenza and pneumonia (J09-J18)	4	0.7	*
9	Septicemia (A40-A41)	3	0.5	*
9	Diabetes mellitus (E10-E14)	3	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
11	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Aortic aneurysm and dissection (I71)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	53	8.7	5.3
Asian or Pacific Islander, male, 25-34 years				
...	All causes	632	100.0	50.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	28.0	14.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	18.2	9.1
3	Malignant neoplasms (C00-C97)	82	13.0	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	10.9	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	9.3	4.7
6	Cerebrovascular diseases (I60-I69)	12	1.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.6	*
8	Diabetes mellitus (E10-E14)	6	0.9	*
8	Influenza and pneumonia (J09-J18)	6	0.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.9	*
11	Septicemia (A40-A41)	5	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
13	Aortic aneurysm and dissection (I71)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities	3	0.5	*
...	All other causes (Residual)	66	10.4	5.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,110	100.0	88.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	22.2	19.7
2	Malignant neoplasms (C00-C97)	235	21.2	18.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	15.2	13.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	11.8	10.5
5	Cerebrovascular diseases (I60-I69)	48	4.3	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.6	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	31	2.8	2.5
8	Diabetes mellitus (E10-E14)	24	2.2	1.9
9	Viral hepatitis (B15-B19)	13	1.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.2	*
11	Septicemia (A40-A41)	10	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
12	Aortic aneurysm and dissection (I71)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Influenza and pneumonia (J09-J18)	5	0.5	*
...	All other causes (Residual)	121	10.9	9.7
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,212	100.0	229.1
1	Malignant neoplasms (C00-C97)	709	32.1	73.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	497	22.5	51.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	7.5	17.2
4	Cerebrovascular diseases (I60-I69)	142	6.4	14.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	4.8	11.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	68	3.1	7.0
7	Diabetes mellitus (E10-E14)	67	3.0	6.9
8	Viral hepatitis (B15-B19)	37	1.7	3.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	1.4	3.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.2	2.7
11	Septicemia (A40-A41)	23	1.0	2.4
12	Influenza and pneumonia (J09-J18)	22	1.0	2.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.9	2.2
14	Chronic lower respiratory diseases (J40-J47)	20	0.9	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
...	All other causes (Residual)	262	11.8	27.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	3,334	100.0	523.1
1	Malignant neoplasms (C00-C97)	1,211	36.3	190.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	838	25.1	131.5
3	Cerebrovascular diseases (I60-I69)	201	6.0	31.5
4	Diabetes mellitus (E10-E14)	168	5.0	26.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	3.8	19.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	70	2.1	11.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.0	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	1.9	10.2
9	Chronic lower respiratory diseases (J40-J47)	56	1.7	8.8
10	Viral hepatitis (B15-B19)	47	1.4	7.4
11	Septicemia (A40-A41)	40	1.2	6.3
12	Influenza and pneumonia (J09-J18)	36	1.1	5.6
13	Aortic aneurysm and dissection (I71)	31	0.9	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.9	4.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.7	3.6
...	All other causes (Residual)	325	9.7	51.0
Asian or Pacific Islander, male, 65-74 years				
...	All causes	4,487	100.0	1,304.7
1	Malignant neoplasms (C00-C97)	1,619	36.1	470.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,105	24.6	321.3
3	Cerebrovascular diseases (I60-I69)	312	7.0	90.7
4	Diabetes mellitus (E10-E14)	205	4.6	59.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	2.9	37.5
6	Chronic lower respiratory diseases (J40-J47)	127	2.8	36.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.4	31.1
8	Influenza and pneumonia (J09-J18)	91	2.0	26.5
9	Septicemia (A40-A41)	81	1.8	23.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.1	14.5
11	Aortic aneurysm and dissection (I71)	43	1.0	12.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.8	10.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.8	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	0.7	9.0
15	Parkinson's disease (G20-G21)	29	0.6	8.4
...	All other causes (Residual)	489	10.9	142.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,950	100.0	3,538.4
1	Malignant neoplasms (C00-C97)	1,706	28.7	1,014.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,524	25.6	906.3
3	Cerebrovascular diseases (I60-I69)	461	7.7	274.2
4	Chronic lower respiratory diseases (J40-J47)	322	5.4	191.5
5	Diabetes mellitus (E10-E14)	221	3.7	131.4
5	Influenza and pneumonia (J09-J18)	221	3.7	131.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	2.6	93.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	2.5	88.6
9	Parkinson's disease (G20-G21)	83	1.4	49.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.4	48.2
11	Alzheimer's disease (G30)	78	1.3	46.4
12	Septicemia (A40-A41)	75	1.3	44.6
13	Aortic aneurysm and dissection (I71)	57	1.0	33.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.9	30.3
15	Pneumonitis due to solids and liquids (J69)	48	0.8	28.5
...	All other causes (Residual)	716	12.0	425.8
Asian or Pacific Islander, male, 85 years and over				
...	All causes	4,717	100.0	8,918.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,410	29.9	2,665.8
2	Malignant neoplasms (C00-C97)	755	16.0	1,427.4
3	Cerebrovascular diseases (I60-I69)	396	8.4	748.7
4	Influenza and pneumonia (J09-J18)	307	6.5	580.4
5	Chronic lower respiratory diseases (J40-J47)	248	5.3	468.9
6	Diabetes mellitus (E10-E14)	166	3.5	313.8
7	Alzheimer's disease (G30)	146	3.1	276.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	2.5	219.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	106	2.2	200.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	2.2	200.4
11	Pneumonitis due to solids and liquids (J69)	76	1.6	143.7
12	Parkinson's disease (G20-G21)	75	1.6	141.8
13	Septicemia (A40-A41)	62	1.3	117.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.8	68.1
15	Aortic aneurysm and dissection (I71)	28	0.6	52.9
...	All other causes (Residual)	684	14.5	1,293.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	21,786	100.0	287.2
1	Malignant neoplasms (C00-C97)	5,932	27.2	78.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,850	22.3	63.9
3	Cerebrovascular diseases (I60-I69)	2,005	9.2	26.4
4	Diabetes mellitus (E10-E14)	883	4.1	11.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	823	3.8	10.8
6	Influenza and pneumonia (J09-J18)	632	2.9	8.3
7	Chronic lower respiratory diseases (J40-J47)	536	2.5	7.1
8	Alzheimer's disease (G30)	503	2.3	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	1.9	5.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	348	1.6	4.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	1.2	3.6
12	Septicemia (A40-A41)	249	1.1	3.3
13	Certain conditions originating in the perinatal period (P00-P96)	209	1.0	2.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.7	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	155	0.7	2.0
...	All other causes (Residual)	3,809	17.5	50.2
Asian or Pacific Islander, female, 1-4 years				
...	All causes	73	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	20.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.1	*
3	Malignant neoplasms (C00-C97)	9	12.3	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	12.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.5	*
6	Septicemia (A40-A41)	2	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
7	Meningitis (G00,G03)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.4	*
...	All other causes (Residual)	19	26.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	92	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	29.3	2.9
2	Malignant neoplasms (C00-C97)	15	16.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.7	*
4	Cerebrovascular diseases (I60-I69)	5	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.3	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	4.3	*
7	Influenza and pneumonia (J09-J18)	2	2.2	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	2	2.2	*
10	Septicemia (A40-A41)	1	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
10	Anemias (D50-D64)	1	1.1	*
10	Hernia (K40-K46)	1	1.1	*
...	All other causes (Residual)	19	20.7	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	232	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	41.8	10.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	15.5	3.8
3	Malignant neoplasms (C00-C97)	29	12.5	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.3	*
6	Cerebrovascular diseases (I60-I69)	3	1.3	*
6	Influenza and pneumonia (J09-J18)	3	1.3	*
9	Septicemia (A40-A41)	2	0.9	*
9	Meningitis (G00,G03)	2	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
11	Aortic aneurysm and dissection (I71)	1	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
11	Pneumonitis due to solids and liquids (J69)	1	0.4	*
11	Diseases of appendix (K35-K38)	1	0.4	*
11	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
...	All other causes (Residual)	31	13.4	3.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	366	100.0	28.1
1	Malignant neoplasms (C00-C97)	78	21.3	6.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	18.0	5.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	16.4	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	6.6	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.6	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	18	4.9	*
7	Cerebrovascular diseases (I60-I69)	10	2.7	*
8	Diabetes mellitus (E10-E14)	6	1.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.1	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Septicemia (A40-A41)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Aortic aneurysm and dissection (I71)	2	0.5	*
12	Influenza and pneumonia (J09-J18)	2	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	55	15.0	4.2
Asian or Pacific Islander, female, 35-44 years				
...	All causes	722	100.0	54.9
1	Malignant neoplasms (C00-C97)	316	43.8	24.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	10.1	5.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	8.3	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	8.0	4.4
5	Cerebrovascular diseases (I60-I69)	46	6.4	3.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.6	2.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	15	2.1	*
8	Diabetes mellitus (E10-E14)	9	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
11	Septicemia (A40-A41)	6	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
13	Viral hepatitis (B15-B19)	5	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
15	Influenza and pneumonia (J09-J18)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	72	10.0	5.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,465	100.0	136.2
1	Malignant neoplasms (C00-C97)	753	51.4	70.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	8.9	12.1
3	Cerebrovascular diseases (I60-I69)	107	7.3	9.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	5.8	7.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	2.9	3.9
6	Diabetes mellitus (E10-E14)	38	2.6	3.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.7	2.3
8	Viral hepatitis (B15-B19)	18	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
10	Influenza and pneumonia (J09-J18)	17	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	1.0	*
13	Septicemia (A40-A41)	13	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.8	*
...	All other causes (Residual)	166	11.3	15.4
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,440	100.0	329.2
1	Malignant neoplasms (C00-C97)	1,202	49.3	162.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	14.2	46.8
3	Cerebrovascular diseases (I60-I69)	187	7.7	25.2
4	Diabetes mellitus (E10-E14)	105	4.3	14.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	3.8	12.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.4	4.5
7	Viral hepatitis (B15-B19)	31	1.3	4.2
7	Chronic lower respiratory diseases (J40-J47)	31	1.3	4.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.3	4.2
10	Influenza and pneumonia (J09-J18)	25	1.0	3.4
11	Septicemia (A40-A41)	23	0.9	3.1
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.9	3.1
13	Aortic aneurysm and dissection (I71)	21	0.9	2.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	21	0.9	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.6	*
...	All other causes (Residual)	239	9.8	32.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,454	100.0	832.7
1	Malignant neoplasms (C00-C97)	1,281	37.1	308.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	699	20.2	168.5
3	Cerebrovascular diseases (I60-I69)	301	8.7	72.6
4	Diabetes mellitus (E10-E14)	186	5.4	44.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	2.6	21.9
6	Chronic lower respiratory diseases (J40-J47)	78	2.3	18.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.2	18.1
8	Influenza and pneumonia (J09-J18)	66	1.9	15.9
9	Septicemia (A40-A41)	43	1.2	10.4
9	Viral hepatitis (B15-B19)	43	1.2	10.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.2	10.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.1	9.2
13	Aortic aneurysm and dissection (I71)	25	0.7	6.0
14	Alzheimer's disease (G30)	22	0.6	5.3
15	Parkinson's disease (G20-G21)	19	0.6	*
...	All other causes (Residual)	444	12.9	107.0
Asian or Pacific Islander, female, 75-84 years				
...	All causes	6,078	100.0	2,470.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,504	24.7	611.4
2	Malignant neoplasms (C00-C97)	1,479	24.3	601.2
3	Cerebrovascular diseases (I60-I69)	639	10.5	259.7
4	Diabetes mellitus (E10-E14)	316	5.2	128.5
5	Chronic lower respiratory diseases (J40-J47)	208	3.4	84.5
6	Influenza and pneumonia (J09-J18)	185	3.0	75.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	2.5	62.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.5	62.2
9	Alzheimer's disease (G30)	124	2.0	50.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.9	47.2
11	Septicemia (A40-A41)	84	1.4	34.1
12	Parkinson's disease (G20-G21)	70	1.2	28.5
13	Aortic aneurysm and dissection (I71)	47	0.8	19.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	47	0.8	19.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	16.3
...	All other causes (Residual)	912	15.0	370.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85 years and over				
...	All causes	6,444	100.0	7,334.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,061	32.0	2,345.6
2	Malignant neoplasms (C00-C97)	769	11.9	875.2
3	Cerebrovascular diseases (I60-I69)	705	10.9	802.4
4	Alzheimer's disease (G30)	354	5.5	402.9
5	Influenza and pneumonia (J09-J18)	325	5.0	369.9
6	Diabetes mellitus (E10-E14)	223	3.5	253.8
7	Chronic lower respiratory diseases (J40-J47)	192	3.0	218.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	164	2.5	186.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	2.1	152.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	1.8	129.7
11	Septicemia (A40-A41)	69	1.1	78.5
12	Pneumonitis due to solids and liquids (J69)	56	0.9	63.7
13	Parkinson's disease (G20-G21)	51	0.8	58.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.6	43.2
15	Atherosclerosis (I70)	31	0.5	35.3
...	All other causes (Residual)	1,158	18.0	1,317.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable