

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,515,458	100.0	807.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	596,577	23.7	191.5
2	Malignant neoplasms (C00-C97)	576,691	22.9	185.1
3	Chronic lower respiratory diseases (J40-J47)	142,943	5.7	45.9
4	Cerebrovascular diseases (I60-I69)	128,932	5.1	41.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126,438	5.0	40.6
6	Alzheimer's disease (G30)	84,974	3.4	27.3
7	Diabetes mellitus (E10-E14)	73,831	2.9	23.7
8	Influenza and pneumonia (J09-J18)	53,826	2.1	17.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,591	1.8	14.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39,518	1.6	12.7
11	Septicemia (A40-A41)	35,748	1.4	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33,642	1.3	10.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27,853	1.1	8.9
14	Parkinson's disease (G20-G21)	23,111	0.9	7.4
15	Pneumonitis due to solids and liquids (J69)	18,195	0.7	5.8
...	All other causes (Residual)	507,588	20.2	162.9
All races, both sexes, 1-4 years				
...	All causes	4,246	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,377	32.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	493	11.6	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	9.7	2.5
4	Malignant neoplasms (C00-C97)	353	8.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.9	1.0
6	Influenza and pneumonia (J09-J18)	112	2.6	0.7
7	Septicemia (A40-A41)	61	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	53	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	42	1.0	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	34	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.5	0.1
13	Anemias (D50-D64)	14	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,036	24.4	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,401	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,635	30.3	4.0
2	Malignant neoplasms (C00-C97)	860	15.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	358	6.6	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	5.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	3.8	0.5
7	Chronic lower respiratory diseases (J40-J47)	136	2.5	0.3
8	Influenza and pneumonia (J09-J18)	118	2.2	0.3
9	Cerebrovascular diseases (I60-I69)	87	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	1.3	0.2
11	Septicemia (A40-A41)	69	1.3	0.2
12	Anemias (D50-D64)	43	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	30	0.6	0.1
14	Diabetes mellitus (E10-E14)	26	0.5	0.1
15	Pneumonitis due to solids and liquids (J69)	18	0.3	*
...	All other causes (Residual)	1,178	21.8	2.9
All races, both sexes, 15-24 years				
...	All causes	29,667	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,330	41.6	28.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,822	16.3	11.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,554	15.4	10.4
4	Malignant neoplasms (C00-C97)	1,611	5.4	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	998	3.4	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	432	1.5	1.0
7	Influenza and pneumonia (J09-J18)	220	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	186	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	172	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	170	0.6	0.4
11	Diabetes mellitus (E10-E14)	167	0.6	0.4
12	Septicemia (A40-A41)	134	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	134	0.5	0.3
14	Legal intervention (Y35,Y89.0)	102	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.3	0.2
...	All other causes (Residual)	3,536	11.9	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	113,641	100.0	137.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,748	27.1	37.3
2	Malignant neoplasms (C00-C97)	15,216	13.4	18.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,936	12.3	16.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,699	11.2	15.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,704	5.9	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,954	2.6	3.6
7	Diabetes mellitus (E10-E14)	2,528	2.2	3.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,285	2.0	2.8
9	Cerebrovascular diseases (I60-I69)	2,248	2.0	2.7
10	Influenza and pneumonia (J09-J18)	1,374	1.2	1.7
11	Septicemia (A40-A41)	1,145	1.0	1.4
12	Chronic lower respiratory diseases (J40-J47)	1,002	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	880	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	829	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	646	0.6	0.8
...	All other causes (Residual)	18,447	16.2	22.4
All races, both sexes, 45-64 years				
...	All causes	506,562	100.0	611.9
1	Malignant neoplasms (C00-C97)	161,469	31.9	195.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	105,842	20.9	127.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,907	7.1	43.4
4	Chronic lower respiratory diseases (J40-J47)	19,678	3.9	23.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	19,613	3.9	23.7
6	Diabetes mellitus (E10-E14)	18,700	3.7	22.6
7	Cerebrovascular diseases (I60-I69)	16,910	3.3	20.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15,379	3.0	18.6
9	Septicemia (A40-A41)	7,414	1.5	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,738	1.3	8.1
11	Influenza and pneumonia (J09-J18)	6,409	1.3	7.7
12	Viral hepatitis (B15-B19)	5,832	1.2	7.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,577	0.9	5.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,054	0.8	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,140	0.6	3.8
...	All other causes (Residual)	74,900	14.8	90.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,831,822	100.0	4,425.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	475,097	25.9	1,147.7
2	Malignant neoplasms (C00-C97)	397,106	21.7	959.3
3	Chronic lower respiratory diseases (J40-J47)	121,869	6.7	294.4
4	Cerebrovascular diseases (I60-I69)	109,323	6.0	264.1
5	Alzheimer's disease (G30)	84,032	4.6	203.0
6	Diabetes mellitus (E10-E14)	52,402	2.9	126.6
7	Influenza and pneumonia (J09-J18)	45,386	2.5	109.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,258	2.4	104.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,796	2.1	91.3
10	Septicemia (A40-A41)	26,746	1.5	64.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,280	1.3	56.2
12	Parkinson's disease (G20-G21)	22,535	1.2	54.4
13	Pneumonitis due to solids and liquids (J69)	16,218	0.9	39.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12,166	0.7	29.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11,030	0.6	26.6
...	All other causes (Residual)	353,578	19.3	854.2
All races, male, all ages				
...	All causes	1,254,978	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	308,398	24.6	201.2
2	Malignant neoplasms (C00-C97)	302,231	24.1	197.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,257	6.3	51.7
4	Chronic lower respiratory diseases (J40-J47)	67,521	5.4	44.0
5	Cerebrovascular diseases (I60-I69)	52,335	4.2	34.1
6	Diabetes mellitus (E10-E14)	38,324	3.1	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,003	2.5	20.2
8	Alzheimer's disease (G30)	25,677	2.0	16.8
9	Influenza and pneumonia (J09-J18)	25,401	2.0	16.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,649	1.8	14.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	21,621	1.7	14.1
12	Septicemia (A40-A41)	16,484	1.3	10.8
13	Parkinson's disease (G20-G21)	13,546	1.1	8.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,745	1.0	8.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,303	0.9	7.4
...	All other causes (Residual)	226,483	18.0	147.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,405	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	839	34.9	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.1	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	9.9	2.9
4	Malignant neoplasms (C00-C97)	183	7.6	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.5	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	37	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	29	1.2	0.4
9	Septicemia (A40-A41)	26	1.1	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
13	Anemias (D50-D64)	9	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	8	0.3	*
15	Meningitis (G00,G03)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	557	23.2	6.7
All races, male, 5-14 years				
...	All causes	3,179	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	32.5	4.9
2	Malignant neoplasms (C00-C97)	483	15.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	6.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	6.2	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	3.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	87	2.7	0.4
8	Influenza and pneumonia (J09-J18)	63	2.0	0.3
9	Cerebrovascular diseases (I60-I69)	46	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.2
11	Septicemia (A40-A41)	34	1.1	0.2
12	Anemias (D50-D64)	27	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Pneumonitis due to solids and liquids (J69)	14	0.4	*
15	Diabetes mellitus (E10-E14)	11	0.3	*
...	All other causes (Residual)	656	20.6	3.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,924	100.0	97.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,172	41.8	40.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,959	18.1	17.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,941	18.0	17.6
4	Malignant neoplasms (C00-C97)	966	4.4	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	684	3.1	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	1.3	1.2
7	Influenza and pneumonia (J09-J18)	123	0.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	117	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	104	0.5	0.5
10	Diabetes mellitus (E10-E14)	101	0.5	0.5
11	Legal intervention (Y35,Y89.0)	100	0.5	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	80	0.4	0.4
13	Septicemia (A40-A41)	72	0.3	0.3
14	Anemias (D50-D64)	60	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.2	0.2
...	All other causes (Residual)	2,125	9.7	9.5
All races, male, 25-44 years				
...	All causes	73,300	100.0	177.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,885	29.9	53.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,906	13.5	24.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,603	13.1	23.3
4	Malignant neoplasms (C00-C97)	6,747	9.2	16.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,439	7.4	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,878	2.6	4.6
7	Diabetes mellitus (E10-E14)	1,508	2.1	3.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,488	2.0	3.6
9	Cerebrovascular diseases (I60-I69)	1,190	1.6	2.9
10	Influenza and pneumonia (J09-J18)	746	1.0	1.8
11	Septicemia (A40-A41)	587	0.8	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	515	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	497	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	473	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	309	0.4	0.7
...	All other causes (Residual)	10,529	14.4	25.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	308,350	100.0	763.7
1	Malignant neoplasms (C00-C97)	86,871	28.2	215.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,079	24.0	183.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,358	7.9	60.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	13,457	4.4	33.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,655	3.8	28.9
6	Diabetes mellitus (E10-E14)	11,367	3.7	28.2
7	Chronic lower respiratory diseases (J40-J47)	10,063	3.3	24.9
8	Cerebrovascular diseases (I60-I69)	9,606	3.1	23.8
9	Viral hepatitis (B15-B19)	4,208	1.4	10.4
10	Septicemia (A40-A41)	3,944	1.3	9.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,854	1.2	9.5
12	Influenza and pneumonia (J09-J18)	3,718	1.2	9.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,412	1.1	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,536	0.8	6.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,295	0.7	5.7
...	All other causes (Residual)	42,927	13.9	106.3
All races, male, 65 years and over				
...	All causes	832,394	100.0	4,639.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	223,669	26.9	1,246.5
2	Malignant neoplasms (C00-C97)	206,939	24.9	1,153.3
3	Chronic lower respiratory diseases (J40-J47)	56,697	6.8	316.0
4	Cerebrovascular diseases (I60-I69)	41,284	5.0	230.1
5	Diabetes mellitus (E10-E14)	25,333	3.0	141.2
6	Alzheimer's disease (G30)	25,260	3.0	140.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,304	2.6	118.7
8	Influenza and pneumonia (J09-J18)	20,572	2.5	114.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,183	2.2	101.3
10	Parkinson's disease (G20-G21)	13,187	1.6	73.5
11	Septicemia (A40-A41)	11,723	1.4	65.3
12	Pneumonitis due to solids and liquids (J69)	8,471	1.0	47.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,445	1.0	47.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,257	0.8	34.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,221	0.7	34.7
...	All other causes (Residual)	138,849	16.7	773.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,260,480	100.0	796.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	288,179	22.9	182.0
2	Malignant neoplasms (C00-C97)	274,460	21.8	173.4
3	Cerebrovascular diseases (I60-I69)	76,597	6.1	48.4
4	Chronic lower respiratory diseases (J40-J47)	75,422	6.0	47.6
5	Alzheimer's disease (G30)	59,297	4.7	37.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,181	3.7	29.8
7	Diabetes mellitus (E10-E14)	35,507	2.8	22.4
8	Influenza and pneumonia (J09-J18)	28,425	2.3	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,942	1.8	14.5
10	Septicemia (A40-A41)	19,264	1.5	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,550	1.3	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,021	1.0	7.6
13	Parkinson's disease (G20-G21)	9,565	0.8	6.0
14	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,515	0.7	5.4
...	All other causes (Residual)	278,036	22.1	175.6
All races, female, 1-4 years				
...	All causes	1,841	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	29.2	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.3	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	9.4	2.2
4	Malignant neoplasms (C00-C97)	170	9.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	4.4	1.0
6	Influenza and pneumonia (J09-J18)	57	3.1	0.7
7	Septicemia (A40-A41)	35	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.8	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	472	25.6	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,222	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	27.1	3.0
2	Malignant neoplasms (C00-C97)	377	17.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	7.2	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	4.9	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	4.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	3.9	0.4
7	Influenza and pneumonia (J09-J18)	55	2.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	49	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.8	0.2
10	Septicemia (A40-A41)	35	1.6	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.5	0.2
12	Anemias (D50-D64)	16	0.7	*
13	Diabetes mellitus (E10-E14)	15	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
15	Meningitis (G00,G03)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
...	All other causes (Residual)	510	23.0	2.5
All races, female, 15-24 years				
...	All causes	7,743	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,158	40.8	14.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	863	11.1	4.0
3	Malignant neoplasms (C00-C97)	645	8.3	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	613	7.9	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	4.1	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	172	2.2	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	2.0	0.7
8	Influenza and pneumonia (J09-J18)	97	1.3	0.5
9	Cerebrovascular diseases (I60-I69)	82	1.1	0.4
10	Diabetes mellitus (E10-E14)	66	0.9	0.3
11	Septicemia (A40-A41)	62	0.8	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	0.3
14	Chronic lower respiratory diseases (J40-J47)	53	0.7	0.2
15	Anemias (D50-D64)	37	0.5	0.2
...	All other causes (Residual)	1,316	17.0	6.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	40,341	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,863	22.0	21.5
2	Malignant neoplasms (C00-C97)	8,469	21.0	20.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,333	10.7	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,793	6.9	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,265	3.1	3.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,076	2.7	2.6
7	Cerebrovascular diseases (I60-I69)	1,058	2.6	2.6
8	Diabetes mellitus (E10-E14)	1,020	2.5	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	797	2.0	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	646	1.6	1.6
11	Influenza and pneumonia (J09-J18)	628	1.6	1.5
12	Septicemia (A40-A41)	558	1.4	1.4
13	Chronic lower respiratory diseases (J40-J47)	505	1.3	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	0.9	0.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	356	0.9	0.9
...	All other causes (Residual)	7,609	18.9	18.5
All races, female, 45-64 years				
...	All causes	198,212	100.0	467.4
1	Malignant neoplasms (C00-C97)	74,598	37.6	175.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,763	16.0	74.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,549	5.8	27.2
4	Chronic lower respiratory diseases (J40-J47)	9,615	4.9	22.7
5	Diabetes mellitus (E10-E14)	7,333	3.7	17.3
6	Cerebrovascular diseases (I60-I69)	7,304	3.7	17.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,156	3.1	14.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,724	1.9	8.8
9	Septicemia (A40-A41)	3,470	1.8	8.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,884	1.5	6.8
11	Influenza and pneumonia (J09-J18)	2,691	1.4	6.3
12	Viral hepatitis (B15-B19)	1,624	0.8	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,518	0.8	3.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,165	0.6	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	975	0.5	2.3
...	All other causes (Residual)	31,843	16.1	75.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	999,428	100.0	4,261.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,428	25.2	1,072.2
2	Malignant neoplasms (C00-C97)	190,167	19.0	810.9
3	Cerebrovascular diseases (I60-I69)	68,039	6.8	290.1
4	Chronic lower respiratory diseases (J40-J47)	65,172	6.5	277.9
5	Alzheimer's disease (G30)	58,772	5.9	250.6
6	Diabetes mellitus (E10-E14)	27,069	2.7	115.4
7	Influenza and pneumonia (J09-J18)	24,814	2.5	105.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,954	2.2	93.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,613	2.0	83.6
10	Septicemia (A40-A41)	15,023	1.5	64.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,835	1.5	63.3
12	Parkinson's disease (G20-G21)	9,348	0.9	39.9
13	Pneumonitis due to solids and liquids (J69)	7,747	0.8	33.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,945	0.6	25.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,773	0.5	20.4
...	All other causes (Residual)	214,729	21.5	915.7
White, both sexes, all ages				
...	All causes	2,156,077	100.0	872.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	513,701	23.8	207.9
2	Malignant neoplasms (C00-C97)	492,348	22.8	199.3
3	Chronic lower respiratory diseases (J40-J47)	131,495	6.1	53.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,751	5.1	44.4
5	Cerebrovascular diseases (I60-I69)	108,542	5.0	43.9
6	Alzheimer's disease (G30)	78,055	3.6	31.6
7	Diabetes mellitus (E10-E14)	57,974	2.7	23.5
8	Influenza and pneumonia (J09-J18)	46,658	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,268	1.7	14.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,775	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	29,507	1.4	11.9
12	Septicemia (A40-A41)	28,838	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,845	1.0	8.8
14	Parkinson's disease (G20-G21)	21,688	1.0	8.8
15	Pneumonitis due to solids and liquids (J69)	16,263	0.8	6.6
...	All other causes (Residual)	427,369	19.8	173.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,980	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	34.5	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	360	12.1	3.0
3	Malignant neoplasms (C00-C97)	275	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	0.9
6	Influenza and pneumonia (J09-J18)	66	2.2	0.5
7	Septicemia (A40-A41)	45	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.3
9	Cerebrovascular diseases (I60-I69)	23	0.8	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	21	0.7	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	721	24.2	5.9
White, both sexes, 5-14 years				
...	All causes	3,927	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,242	31.6	4.0
2	Malignant neoplasms (C00-C97)	676	17.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	6.6	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	4.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	3.6	0.5
7	Influenza and pneumonia (J09-J18)	83	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	59	1.5	0.2
9	Septicemia (A40-A41)	55	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	49	1.2	0.2
12	Diabetes mellitus (E10-E14)	21	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	19	0.5	*
14	Anemias (D50-D64)	15	0.4	*
15	Pneumonitis due to solids and liquids (J69)	11	0.3	*
...	All other causes (Residual)	847	21.6	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,654	100.0	65.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,391	48.0	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,023	18.6	12.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,611	7.4	4.8
4	Malignant neoplasms (C00-C97)	1,267	5.9	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	1.4	0.9
7	Influenza and pneumonia (J09-J18)	165	0.8	0.5
8	Cerebrovascular diseases (I60-I69)	143	0.7	0.4
9	Diabetes mellitus (E10-E14)	115	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	102	0.5	0.3
11	Septicemia (A40-A41)	92	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	81	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.3	0.2
14	Legal intervention (Y35,Y89.0)	49	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,554	11.8	7.7
White, both sexes, 25-44 years				
...	All causes	85,713	100.0	134.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,249	30.6	41.1
2	Malignant neoplasms (C00-C97)	11,607	13.5	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,102	13.0	17.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,703	11.3	15.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,975	3.5	4.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,449	2.9	3.8
7	Diabetes mellitus (E10-E14)	1,689	2.0	2.6
8	Cerebrovascular diseases (I60-I69)	1,440	1.7	2.3
9	Influenza and pneumonia (J09-J18)	1,018	1.2	1.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	922	1.1	1.4
11	Septicemia (A40-A41)	780	0.9	1.2
12	Chronic lower respiratory diseases (J40-J47)	692	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	679	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	478	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-O99)	385	0.4	0.6
...	All other causes (Residual)	13,545	15.8	21.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	403,116	100.0	595.8
1	Malignant neoplasms (C00-C97)	130,368	32.3	192.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,892	20.3	121.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,416	7.5	45.0
4	Chronic lower respiratory diseases (J40-J47)	17,174	4.3	25.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	16,957	4.2	25.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,419	3.6	21.3
7	Diabetes mellitus (E10-E14)	13,727	3.4	20.3
8	Cerebrovascular diseases (I60-I69)	11,681	2.9	17.3
9	Septicemia (A40-A41)	5,423	1.3	8.0
10	Influenza and pneumonia (J09-J18)	5,068	1.3	7.5
11	Viral hepatitis (B15-B19)	4,724	1.2	7.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,389	1.1	6.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,576	0.6	3.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,068	0.5	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,943	0.5	2.9
...	All other causes (Residual)	60,291	15.0	89.1
White, both sexes, 65 years and over				
...	All causes	1,623,128	100.0	4,530.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	421,005	25.9	1,175.1
2	Malignant neoplasms (C00-C97)	348,101	21.4	971.6
3	Chronic lower respiratory diseases (J40-J47)	113,457	7.0	316.7
4	Cerebrovascular diseases (I60-I69)	95,112	5.9	265.5
5	Alzheimer's disease (G30)	77,205	4.8	215.5
6	Diabetes mellitus (E10-E14)	42,415	2.6	118.4
7	Influenza and pneumonia (J09-J18)	40,141	2.5	112.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,689	2.4	110.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,293	1.9	87.3
10	Septicemia (A40-A41)	22,344	1.4	62.4
11	Parkinson's disease (G20-G21)	21,178	1.3	59.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,012	1.2	53.1
13	Pneumonitis due to solids and liquids (J69)	14,652	0.9	40.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,013	0.7	30.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10,070	0.6	28.1
...	All other causes (Residual)	316,441	19.5	883.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,071,966	100.0	876.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	265,596	24.8	217.1
2	Malignant neoplasms (C00-C97)	259,323	24.2	212.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,123	6.4	55.7
4	Chronic lower respiratory diseases (J40-J47)	61,507	5.7	50.3
5	Cerebrovascular diseases (I60-I69)	43,264	4.0	35.4
6	Diabetes mellitus (E10-E14)	30,783	2.9	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,103	2.6	23.0
8	Alzheimer's disease (G30)	23,640	2.2	19.3
9	Influenza and pneumonia (J09-J18)	21,953	2.0	17.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,067	1.8	15.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,275	1.7	14.9
12	Septicemia (A40-A41)	13,371	1.2	10.9
13	Parkinson's disease (G20-G21)	12,746	1.2	10.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,698	0.8	7.1
15	Pneumonitis due to solids and liquids (J69)	8,653	0.8	7.1
...	All other causes (Residual)	188,864	17.6	154.4
White, male, 1-4 years				
...	All causes	1,689	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	634	37.5	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	11.4	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	8.5	2.3
4	Malignant neoplasms (C00-C97)	136	8.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.3	0.9
6	Influenza and pneumonia (J09-J18)	32	1.9	0.5
7	Septicemia (A40-A41)	20	1.2	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
10	Cerebrovascular diseases (I60-I69)	14	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	389	23.0	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,321	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	791	34.1	4.9
2	Malignant neoplasms (C00-C97)	379	16.3	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	6.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	6.1	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	4.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.8	0.4
7	Influenza and pneumonia (J09-J18)	43	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	33	1.4	0.2
9	Septicemia (A40-A41)	29	1.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	29	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.2	0.2
12	Anemias (D50-D64)	11	0.5	*
13	Pneumonitis due to solids and liquids (J69)	8	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.3	*
15	Diabetes mellitus (E10-E14)	7	0.3	*
...	All other causes (Residual)	486	20.9	3.0
White, male, 15-24 years				
...	All causes	15,839	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,724	48.8	45.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,307	20.9	19.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,308	8.3	7.6
4	Malignant neoplasms (C00-C97)	765	4.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	452	2.9	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	1.3	1.2
7	Influenza and pneumonia (J09-J18)	98	0.6	0.6
8	Cerebrovascular diseases (I60-I69)	83	0.5	0.5
9	Diabetes mellitus (E10-E14)	76	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	60	0.4	0.4
11	Septicemia (A40-A41)	55	0.3	0.3
12	Legal intervention (Y35,Y89.0)	48	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.1	0.1
15	Pneumonitis due to solids and liquids (J69)	21	0.1	0.1
...	All other causes (Residual)	1,590	10.0	9.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	56,071	100.0	173.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,660	33.3	57.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,672	15.5	26.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,875	12.3	21.2
4	Malignant neoplasms (C00-C97)	5,370	9.6	16.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,199	3.9	6.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,603	2.9	4.9
7	Diabetes mellitus (E10-E14)	1,042	1.9	3.2
8	Cerebrovascular diseases (I60-I69)	777	1.4	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	703	1.3	2.2
10	Influenza and pneumonia (J09-J18)	561	1.0	1.7
11	Septicemia (A40-A41)	407	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	391	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	335	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	0.5	0.9
15	Aortic aneurysm and dissection (I71)	205	0.4	0.6
...	All other causes (Residual)	7,979	14.2	24.6
White, male, 45-64 years				
...	All causes	248,146	100.0	744.0
1	Malignant neoplasms (C00-C97)	70,746	28.5	212.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,681	23.6	175.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,536	8.3	61.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	11,728	4.7	35.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,931	4.4	32.8
6	Chronic lower respiratory diseases (J40-J47)	8,767	3.5	26.3
7	Diabetes mellitus (E10-E14)	8,572	3.5	25.7
8	Cerebrovascular diseases (I60-I69)	6,626	2.7	19.9
9	Viral hepatitis (B15-B19)	3,458	1.4	10.4
10	Influenza and pneumonia (J09-J18)	2,961	1.2	8.9
11	Septicemia (A40-A41)	2,895	1.2	8.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,558	1.0	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,676	0.7	5.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,667	0.7	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,351	0.5	4.1
...	All other causes (Residual)	34,993	14.1	104.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	739,242	100.0	4,717.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	199,351	27.0	1,272.3
2	Malignant neoplasms (C00-C97)	181,896	24.6	1,160.9
3	Chronic lower respiratory diseases (J40-J47)	52,288	7.1	333.7
4	Cerebrovascular diseases (I60-I69)	35,683	4.8	227.7
5	Alzheimer's disease (G30)	23,267	3.1	148.5
6	Diabetes mellitus (E10-E14)	21,083	2.9	134.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,360	2.6	123.6
8	Influenza and pneumonia (J09-J18)	18,183	2.5	116.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,368	2.1	98.1
10	Parkinson's disease (G20-G21)	12,421	1.7	79.3
11	Septicemia (A40-A41)	9,916	1.3	63.3
12	Pneumonitis due to solids and liquids (J69)	7,673	1.0	49.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,863	0.9	43.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,732	0.8	36.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,714	0.8	36.5
...	All other causes (Residual)	124,444	16.8	794.2
White, female, all ages				
...	All causes	1,084,111	100.0	868.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,105	22.9	198.8
2	Malignant neoplasms (C00-C97)	233,025	21.5	186.8
3	Chronic lower respiratory diseases (J40-J47)	69,988	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	65,278	6.0	52.3
5	Alzheimer's disease (G30)	54,415	5.0	43.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,628	3.8	33.4
7	Diabetes mellitus (E10-E14)	27,191	2.5	21.8
8	Influenza and pneumonia (J09-J18)	24,705	2.3	19.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,993	1.7	14.4
10	Septicemia (A40-A41)	15,467	1.4	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,147	1.2	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,440	1.0	8.4
13	Parkinson's disease (G20-G21)	8,942	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,672	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	7,610	0.7	6.1
...	All other causes (Residual)	238,505	22.0	191.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,291	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	30.4	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	12.9	2.8
3	Malignant neoplasms (C00-C97)	139	10.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	4.0	0.9
6	Influenza and pneumonia (J09-J18)	34	2.6	0.6
7	Septicemia (A40-A41)	25	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
10	Cerebrovascular diseases (I60-I69)	9	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
...	All other causes (Residual)	330	25.6	5.6
White, female, 5-14 years				
...	All causes	1,606	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	451	28.1	3.0
2	Malignant neoplasms (C00-C97)	297	18.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	7.3	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	4.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	4.4	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	4.0	0.4
7	Influenza and pneumonia (J09-J18)	40	2.5	0.3
8	Septicemia (A40-A41)	26	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	26	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.4	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.1
12	Diabetes mellitus (E10-E14)	14	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
14	Meningitis (G00,G03)	5	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	358	22.3	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,815	100.0	36.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,667	45.9	16.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	12.3	4.4
3	Malignant neoplasms (C00-C97)	502	8.6	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	5.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	3.3	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	1.8	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	102	1.8	0.6
8	Influenza and pneumonia (J09-J18)	67	1.2	0.4
9	Cerebrovascular diseases (I60-I69)	60	1.0	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	0.2
11	Diabetes mellitus (E10-E14)	39	0.7	0.2
12	Septicemia (A40-A41)	37	0.6	0.2
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.3	*
...	All other causes (Residual)	926	15.9	5.7
White, female, 25-44 years				
...	All causes	29,642	100.0	94.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,589	25.6	24.1
2	Malignant neoplasms (C00-C97)	6,237	21.0	19.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,828	9.5	9.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,430	8.2	7.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	846	2.9	2.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	776	2.6	2.5
7	Cerebrovascular diseases (I60-I69)	663	2.2	2.1
8	Diabetes mellitus (E10-E14)	647	2.2	2.1
9	Influenza and pneumonia (J09-J18)	457	1.5	1.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	385	1.3	1.2
11	Septicemia (A40-A41)	373	1.3	1.2
12	Chronic lower respiratory diseases (J40-J47)	357	1.2	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	288	1.0	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	219	0.7	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	0.6	0.6
...	All other causes (Residual)	5,361	18.1	17.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	154,970	100.0	451.7
1	Malignant neoplasms (C00-C97)	59,622	38.5	173.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,211	15.0	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,880	6.4	28.8
4	Chronic lower respiratory diseases (J40-J47)	8,407	5.4	24.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,229	3.4	15.2
6	Diabetes mellitus (E10-E14)	5,155	3.3	15.0
7	Cerebrovascular diseases (I60-I69)	5,055	3.3	14.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,488	2.3	10.2
9	Septicemia (A40-A41)	2,528	1.6	7.4
10	Influenza and pneumonia (J09-J18)	2,107	1.4	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,831	1.2	5.3
12	Viral hepatitis (B15-B19)	1,266	0.8	3.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	909	0.6	2.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	782	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	647	0.4	1.9
...	All other causes (Residual)	24,853	16.0	72.4
White, female, 65 years and over				
...	All causes	883,886	100.0	4,384.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221,654	25.1	1,099.6
2	Malignant neoplasms (C00-C97)	166,205	18.8	824.5
3	Chronic lower respiratory diseases (J40-J47)	61,169	6.9	303.4
4	Cerebrovascular diseases (I60-I69)	59,429	6.7	294.8
5	Alzheimer's disease (G30)	53,938	6.1	267.6
6	Influenza and pneumonia (J09-J18)	21,958	2.5	108.9
7	Diabetes mellitus (E10-E14)	21,332	2.4	105.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,329	2.3	100.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,925	1.8	79.0
10	Septicemia (A40-A41)	12,428	1.4	61.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,149	1.4	60.3
12	Parkinson's disease (G20-G21)	8,757	1.0	43.4
13	Pneumonitis due to solids and liquids (J69)	6,979	0.8	34.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,281	0.6	26.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,356	0.5	21.6
...	All other causes (Residual)	191,997	21.7	952.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	290,100	100.0	679.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,372	23.6	160.2
2	Malignant neoplasms (C00-C97)	66,820	23.0	156.6
3	Cerebrovascular diseases (I60-I69)	15,853	5.5	37.1
4	Diabetes mellitus (E10-E14)	12,895	4.4	30.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,531	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,086	3.1	21.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,087	2.8	18.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,858	2.7	18.4
9	Septicemia (A40-A41)	6,028	2.1	14.1
10	Alzheimer's disease (G30)	5,482	1.9	12.8
11	Influenza and pneumonia (J09-J18)	5,167	1.8	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,006	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,215	1.5	9.9
14	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.4	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,816	1.0	6.6
...	All other causes (Residual)	55,829	19.2	130.8
Black, both sexes, 1-4 years				
...	All causes	1,052	100.0	38.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	28.3	10.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	15.1	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.3	3.9
4	Malignant neoplasms (C00-C97)	55	5.2	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.1	2.0
6	Influenza and pneumonia (J09-J18)	32	3.0	1.2
7	Chronic lower respiratory diseases (J40-J47)	25	2.4	0.9
8	Cerebrovascular diseases (I60-I69)	17	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.2	*
10	Septicemia (A40-A41)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Meningitis (G00,G03)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	248	23.6	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,181	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.2	4.8
2	Malignant neoplasms (C00-C97)	133	11.3	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	7.6	1.3
4	Chronic lower respiratory diseases (J40-J47)	84	7.1	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	6.8	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.2	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	3.8	0.7
8	Influenza and pneumonia (J09-J18)	26	2.2	0.4
9	Anemias (D50-D64)	22	1.9	0.3
10	Cerebrovascular diseases (I60-I69)	20	1.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
12	Septicemia (A40-A41)	12	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
14	Pneumonitis due to solids and liquids (J69)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (Residual)	259	21.9	3.9
Black, both sexes, 15-24 years				
...	All causes	6,690	100.0	92.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,826	42.2	39.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,387	20.7	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	7.5	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	4.6	4.3
5	Malignant neoplasms (C00-C97)	260	3.9	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	1.6	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	103	1.5	1.4
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.1
9	Anemias (D50-D64)	73	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.0	0.9
11	Diabetes mellitus (E10-E14)	48	0.7	0.7
12	Legal intervention (Y35,Y89.0)	47	0.7	0.6
13	Influenza and pneumonia (J09-J18)	44	0.7	0.6
14	Septicemia (A40-A41)	36	0.5	0.5
15	Cerebrovascular diseases (I60-I69)	32	0.5	0.4
...	All other causes (Residual)	771	11.5	10.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	22,857	100.0	198.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,675	16.1	31.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,463	15.2	30.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,352	14.7	29.0
4	Malignant neoplasms (C00-C97)	2,762	12.1	23.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,322	5.8	11.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	978	4.3	8.5
7	Diabetes mellitus (E10-E14)	719	3.1	6.2
8	Cerebrovascular diseases (I60-I69)	648	2.8	5.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	1.6	3.2
10	Septicemia (A40-A41)	314	1.4	2.7
11	Influenza and pneumonia (J09-J18)	289	1.3	2.5
12	Chronic lower respiratory diseases (J40-J47)	283	1.2	2.5
13	Anemias (D50-D64)	243	1.1	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	1.0	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	227	1.0	2.0
...	All other causes (Residual)	3,981	17.4	34.5
Black, both sexes, 45-64 years				
...	All causes	87,330	100.0	868.0
1	Malignant neoplasms (C00-C97)	25,348	29.0	251.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,964	24.0	208.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,439	5.1	44.1
4	Cerebrovascular diseases (I60-I69)	4,352	5.0	43.3
5	Diabetes mellitus (E10-E14)	4,221	4.8	42.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,454	2.8	24.4
7	Chronic lower respiratory diseases (J40-J47)	2,186	2.5	21.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,149	2.5	21.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,962	2.2	19.5
10	Septicemia (A40-A41)	1,766	2.0	17.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,355	1.6	13.5
12	Influenza and pneumonia (J09-J18)	1,110	1.3	11.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,069	1.2	10.6
14	Viral hepatitis (B15-B19)	879	1.0	8.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	551	0.6	5.5
...	All other causes (Residual)	12,525	14.3	124.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	163,679	100.0	4,446.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,214	26.4	1,174.0
2	Malignant neoplasms (C00-C97)	38,246	23.4	1,039.1
3	Cerebrovascular diseases (I60-I69)	10,735	6.6	291.7
4	Diabetes mellitus (E10-E14)	7,903	4.8	214.7
5	Chronic lower respiratory diseases (J40-J47)	6,415	3.9	174.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,510	3.4	149.7
7	Alzheimer's disease (G30)	5,405	3.3	146.8
8	Septicemia (A40-A41)	3,823	2.3	103.9
9	Influenza and pneumonia (J09-J18)	3,583	2.2	97.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,404	2.1	92.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,352	1.4	63.9
12	Pneumonitis due to solids and liquids (J69)	1,174	0.7	31.9
13	Parkinson's disease (G20-G21)	841	0.5	22.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	798	0.5	21.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	619	0.4	16.8
...	All other causes (Residual)	29,657	18.1	805.7
Black, male, all ages				
...	All causes	146,884	100.0	719.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,913	23.8	171.0
2	Malignant neoplasms (C00-C97)	33,978	23.1	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,387	5.7	41.1
4	Cerebrovascular diseases (I60-I69)	7,039	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,739	4.6	33.0
6	Diabetes mellitus (E10-E14)	6,048	4.1	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,702	3.2	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,756	2.6	18.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,744	1.9	13.4
10	Septicemia (A40-A41)	2,700	1.8	13.2
11	Influenza and pneumonia (J09-J18)	2,417	1.6	11.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,252	1.5	11.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,200	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,828	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,775	1.2	8.7
...	All other causes (Residual)	25,406	17.3	124.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	596	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	29.0	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	14.6	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	9.9	4.2
4	Malignant neoplasms (C00-C97)	31	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.7	2.0
6	Influenza and pneumonia (J09-J18)	18	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.7	*
8	Cerebrovascular diseases (I60-I69)	14	2.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	135	22.7	9.7
Black, male, 5-14 years				
...	All causes	701	100.0	20.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	28.8	5.9
2	Malignant neoplasms (C00-C97)	77	11.0	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	8.8	1.8
4	Chronic lower respiratory diseases (J40-J47)	57	8.1	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	6.3	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	4.9	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.7	0.8
8	Influenza and pneumonia (J09-J18)	16	2.3	*
9	Anemias (D50-D64)	14	2.0	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
12	Pneumonitis due to solids and liquids (J69)	5	0.7	*
13	Septicemia (A40-A41)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
...	All other causes (Residual)	131	18.7	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,124	100.0	140.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,536	49.5	69.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,037	20.2	28.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	8.2	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	3.9	5.5
5	Malignant neoplasms (C00-C97)	151	2.9	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.2	1.8
8	Chronic lower respiratory diseases (J40-J47)	54	1.1	1.5
9	Legal intervention (Y35,Y89.0)	46	0.9	1.3
10	Anemias (D50-D64)	45	0.9	1.2
11	Diabetes mellitus (E10-E14)	23	0.4	0.6
12	Influenza and pneumonia (J09-J18)	20	0.4	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
14	Cerebrovascular diseases (I60-I69)	16	0.3	*
15	Septicemia (A40-A41)	13	0.3	*
...	All other causes (Residual)	417	8.1	11.4
Black, male, 25-44 years				
...	All causes	14,065	100.0	256.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,039	21.6	55.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,394	17.0	43.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,311	16.4	42.1
4	Malignant neoplasms (C00-C97)	1,018	7.2	18.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	792	5.6	14.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	755	5.4	13.8
7	Diabetes mellitus (E10-E14)	389	2.8	7.1
8	Cerebrovascular diseases (I60-I69)	333	2.4	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.4	3.7
10	Septicemia (A40-A41)	156	1.1	2.8
11	Influenza and pneumonia (J09-J18)	150	1.1	2.7
12	Chronic lower respiratory diseases (J40-J47)	145	1.0	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.0	2.5
14	Anemias (D50-D64)	129	0.9	2.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	121	0.9	2.2
...	All other causes (Residual)	1,995	14.2	36.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	50,792	100.0	1,089.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,278	26.1	284.7
2	Malignant neoplasms (C00-C97)	13,268	26.1	284.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,093	6.1	66.3
4	Cerebrovascular diseases (I60-I69)	2,504	4.9	53.7
5	Diabetes mellitus (E10-E14)	2,349	4.6	50.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,689	3.3	36.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,283	2.5	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,187	2.3	25.5
9	Chronic lower respiratory diseases (J40-J47)	1,132	2.2	24.3
10	Septicemia (A40-A41)	938	1.8	20.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	856	1.7	18.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	785	1.5	16.8
13	Influenza and pneumonia (J09-J18)	619	1.2	13.3
14	Viral hepatitis (B15-B19)	592	1.2	12.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	445	0.9	9.5
...	All other causes (Residual)	6,774	13.3	145.2
Black, male, 65 years and over				
...	All causes	71,529	100.0	4,920.0
1	Malignant neoplasms (C00-C97)	19,425	27.2	1,336.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,014	26.6	1,307.8
3	Cerebrovascular diseases (I60-I69)	4,134	5.8	284.4
4	Chronic lower respiratory diseases (J40-J47)	3,289	4.6	226.2
5	Diabetes mellitus (E10-E14)	3,284	4.6	225.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,331	3.3	160.3
7	Influenza and pneumonia (J09-J18)	1,549	2.2	106.5
8	Septicemia (A40-A41)	1,543	2.2	106.1
9	Alzheimer's disease (G30)	1,529	2.1	105.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,274	1.8	87.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,271	1.8	87.4
12	Pneumonitis due to solids and liquids (J69)	575	0.8	39.6
13	Parkinson's disease (G20-G21)	473	0.7	32.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	368	0.5	25.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	310	0.4	21.3
...	All other causes (Residual)	11,160	15.6	767.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,216	100.0	643.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,459	23.4	150.3
2	Malignant neoplasms (C00-C97)	32,842	22.9	147.5
3	Cerebrovascular diseases (I60-I69)	8,814	6.2	39.6
4	Diabetes mellitus (E10-E14)	6,847	4.8	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,384	3.1	19.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,331	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,144	2.9	18.6
8	Alzheimer's disease (G30)	3,914	2.7	17.6
9	Septicemia (A40-A41)	3,328	2.3	15.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,806	2.0	12.6
11	Influenza and pneumonia (J09-J18)	2,750	1.9	12.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,803	1.3	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,471	1.0	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,119	0.8	5.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,041	0.7	4.7
...	All other causes (Residual)	30,163	21.1	135.5
Black, female, 1-4 years				
...	All causes	456	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	27.4	9.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	15.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	10.7	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	1.9
5	Malignant neoplasms (C00-C97)	24	5.3	1.8
6	Influenza and pneumonia (J09-J18)	14	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	6	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Meningitis (G00,G03)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Atherosclerosis (I70)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	106	23.2	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	480	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.8	3.6
2	Malignant neoplasms (C00-C97)	56	11.7	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	7.5	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.8	0.8
5	Chronic lower respiratory diseases (J40-J47)	27	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.0	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.3	*
8	Influenza and pneumonia (J09-J18)	10	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	9	1.9	*
11	Septicemia (A40-A41)	8	1.7	*
11	Anemias (D50-D64)	8	1.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
14	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	125	26.0	3.8
Black, female, 15-24 years				
...	All causes	1,566	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	22.3	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	18.5	8.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	7.0	3.1
4	Malignant neoplasms (C00-C97)	109	7.0	3.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	5.3	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	64	4.1	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.6	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	39	2.5	1.1
9	Anemias (D50-D64)	28	1.8	0.8
10	Chronic lower respiratory diseases (J40-J47)	27	1.7	0.8
11	Diabetes mellitus (E10-E14)	25	1.6	0.7
12	Influenza and pneumonia (J09-J18)	24	1.5	0.7
13	Septicemia (A40-A41)	23	1.5	0.6
14	Cerebrovascular diseases (I60-I69)	16	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	327	20.9	9.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	8,792	100.0	145.1
1	Malignant neoplasms (C00-C97)	1,744	19.8	28.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,364	15.5	22.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	958	10.9	15.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	567	6.4	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	424	4.8	7.0
6	Diabetes mellitus (E10-E14)	330	3.8	5.4
7	Cerebrovascular diseases (I60-I69)	315	3.6	5.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	211	2.4	3.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	186	2.1	3.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.9	2.7
11	Septicemia (A40-A41)	158	1.8	2.6
12	Influenza and pneumonia (J09-J18)	139	1.6	2.3
13	Chronic lower respiratory diseases (J40-J47)	138	1.6	2.3
14	Anemias (D50-D64)	114	1.3	1.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.2	1.7
...	All other causes (Residual)	1,875	21.3	31.0
Black, female, 45-64 years				
...	All causes	36,538	100.0	676.9
1	Malignant neoplasms (C00-C97)	12,080	33.1	223.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,686	21.0	142.4
3	Diabetes mellitus (E10-E14)	1,872	5.1	34.7
4	Cerebrovascular diseases (I60-I69)	1,848	5.1	34.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,346	3.7	24.9
6	Chronic lower respiratory diseases (J40-J47)	1,054	2.9	19.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	962	2.6	17.8
8	Septicemia (A40-A41)	828	2.3	15.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	765	2.1	14.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	679	1.9	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	570	1.6	10.6
12	Influenza and pneumonia (J09-J18)	491	1.3	9.1
13	Viral hepatitis (B15-B19)	287	0.8	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	0.6	3.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.4	2.9
...	All other causes (Residual)	5,699	15.6	105.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	92,150	100.0	4,138.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,200	26.3	1,086.7
2	Malignant neoplasms (C00-C97)	18,821	20.4	845.2
3	Cerebrovascular diseases (I60-I69)	6,601	7.2	296.4
4	Diabetes mellitus (E10-E14)	4,619	5.0	207.4
5	Alzheimer's disease (G30)	3,876	4.2	174.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,179	3.4	142.8
7	Chronic lower respiratory diseases (J40-J47)	3,126	3.4	140.4
8	Septicemia (A40-A41)	2,280	2.5	102.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,133	2.3	95.8
10	Influenza and pneumonia (J09-J18)	2,034	2.2	91.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,078	1.2	48.4
12	Pneumonitis due to solids and liquids (J69)	599	0.7	26.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	488	0.5	21.9
14	Parkinson's disease (G20-G21)	368	0.4	16.5
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	301	0.3	13.5
...	All other causes (Residual)	18,447	20.0	828.4
American Indian, both sexes, all ages				
...	All causes	15,945	100.0	366.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,918	18.3	67.1
2	Malignant neoplasms (C00-C97)	2,810	17.6	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,829	11.5	42.1
4	Diabetes mellitus (E10-E14)	927	5.8	21.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	821	5.1	18.9
6	Chronic lower respiratory diseases (J40-J47)	657	4.1	15.1
7	Cerebrovascular diseases (I60-I69)	600	3.8	13.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	459	2.9	10.6
9	Influenza and pneumonia (J09-J18)	349	2.2	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	1.8	6.6
11	Septicemia (A40-A41)	269	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.6	5.8
13	Alzheimer's disease (G30)	226	1.4	5.2
14	Certain conditions originating in the perinatal period (P00-P96)	126	0.8	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	0.8	2.9
...	All other causes (Residual)	3,289	20.6	75.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	85	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	36.5	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	10.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	8.2	*
4	Malignant neoplasms (C00-C97)	5	5.9	*
5	Influenza and pneumonia (J09-J18)	3	3.5	*
6	Septicemia (A40-A41)	2	2.4	*
6	Cerebrovascular diseases (I60-I69)	2	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
9	Meningococcal infection (A39)	1	1.2	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	21	24.7	6.7
American Indian, both sexes, 5-14 years				
...	All causes	96	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	33.3	4.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.5	*
3	Malignant neoplasms (C00-C97)	9	9.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
6	Cerebrovascular diseases (I60-I69)	4	4.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
8	Septicemia (A40-A41)	1	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Influenza and pneumonia (J09-J18)	1	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
8	Pneumonitis due to solids and liquids (J69)	1	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.0	*
...	All other causes (Residual)	17	17.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	589	100.0	77.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	47.4	36.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	22.9	17.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.0	6.9
4	Malignant neoplasms (C00-C97)	14	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Influenza and pneumonia (J09-J18)	5	0.8	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
10	Cerebrovascular diseases (I60-I69)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	57	9.7	7.5
American Indian, both sexes, 25-44 years				
...	All causes	2,042	100.0	164.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	643	31.5	51.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	229	11.2	18.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	9.8	16.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	8.6	14.1
5	Malignant neoplasms (C00-C97)	113	5.5	9.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	5.2	8.6
7	Diabetes mellitus (E10-E14)	79	3.9	6.4
8	Cerebrovascular diseases (I60-I69)	31	1.5	2.5
9	Influenza and pneumonia (J09-J18)	28	1.4	2.3
10	Septicemia (A40-A41)	21	1.0	1.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.5	*
15	Viral hepatitis (B15-B19)	9	0.4	*
...	All other causes (Residual)	353	17.3	28.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	5,059	100.0	551.6
1	Malignant neoplasms (C00-C97)	1,042	20.6	113.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	929	18.4	101.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	562	11.1	61.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	464	9.2	50.6
5	Diabetes mellitus (E10-E14)	333	6.6	36.3
6	Chronic lower respiratory diseases (J40-J47)	167	3.3	18.2
7	Cerebrovascular diseases (I60-I69)	148	2.9	16.1
8	Septicemia (A40-A41)	107	2.1	11.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	1.9	10.2
10	Influenza and pneumonia (J09-J18)	92	1.8	10.0
11	Viral hepatitis (B15-B19)	83	1.6	9.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.5	8.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	1.2	6.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.6	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	2.5
...	All other causes (Residual)	852	16.8	92.9
American Indian, both sexes, 65 years and over				
...	All causes	7,708	100.0	2,803.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,797	23.3	653.5
2	Malignant neoplasms (C00-C97)	1,626	21.1	591.4
3	Diabetes mellitus (E10-E14)	513	6.7	186.6
4	Chronic lower respiratory diseases (J40-J47)	475	6.2	172.8
5	Cerebrovascular diseases (I60-I69)	411	5.3	149.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	3.1	88.0
7	Alzheimer's disease (G30)	222	2.9	80.7
8	Influenza and pneumonia (J09-J18)	219	2.8	79.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.6	72.4
10	Septicemia (A40-A41)	133	1.7	48.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	122	1.6	44.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.2	32.7
13	Parkinson's disease (G20-G21)	72	0.9	26.2
14	Pneumonitis due to solids and liquids (J69)	65	0.8	23.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.7	18.9
...	All other causes (Residual)	1,470	19.1	534.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,638	100.0	395.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	19.4	76.9
2	Malignant neoplasms (C00-C97)	1,486	17.2	68.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,227	14.2	56.1
4	Diabetes mellitus (E10-E14)	453	5.2	20.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	453	5.2	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	4.1	16.1
7	Chronic lower respiratory diseases (J40-J47)	304	3.5	13.9
8	Cerebrovascular diseases (I60-I69)	257	3.0	11.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	2.3	9.0
10	Influenza and pneumonia (J09-J18)	177	2.0	8.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.5	5.9
12	Septicemia (A40-A41)	109	1.3	5.0
13	Certain conditions originating in the perinatal period (P00-P96)	73	0.8	3.3
14	Viral hepatitis (B15-B19)	71	0.8	3.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.8	3.0
...	All other causes (Residual)	1,605	18.6	73.4
American Indian, male, 1-4 years				
...	All causes	46	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	45.7	13.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Malignant neoplasms (C00-C97)	3	6.5	*
5	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
6	Cerebrovascular diseases (I60-I69)	1	2.2	*
6	Influenza and pneumonia (J09-J18)	1	2.2	*
...	All other causes (Residual)	9	19.6	*
American Indian, male, 5-14 years				
...	All causes	50	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	30.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	14.0	*
3	Malignant neoplasms (C00-C97)	6	12.0	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	10.0	*
5	Cerebrovascular diseases (I60-I69)	2	4.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.0	*
7	Influenza and pneumonia (J09-J18)	1	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
7	Pneumonitis due to solids and liquids (J69)	1	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (Residual)	8	16.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	422	100.0	107.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	45.5	48.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	25.8	27.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	10.4	11.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.1	*
5	Malignant neoplasms (C00-C97)	8	1.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
6	Legal intervention (Y35,Y89.0)	4	0.9	*
8	Influenza and pneumonia (J09-J18)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Pneumonitis due to solids and liquids (J69)	2	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	37	8.8	9.4
American Indian, male, 25-44 years				
...	All causes	1,292	100.0	202.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	34.2	69.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	11.9	24.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	9.4	19.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	115	8.9	18.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	6.8	13.8
6	Malignant neoplasms (C00-C97)	50	3.9	7.8
7	Diabetes mellitus (E10-E14)	49	3.8	7.7
8	Influenza and pneumonia (J09-J18)	15	1.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.0	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Septicemia (A40-A41)	8	0.6	*
12	Viral hepatitis (B15-B19)	6	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.5	*
15	Legal intervention (Y35,Y89.0)	5	0.4	*
...	All other causes (Residual)	202	15.6	31.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-64 years				
...	All causes	2,953	100.0	658.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	625	21.2	139.3
2	Malignant neoplasms (C00-C97)	544	18.4	121.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	13.3	87.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	264	8.9	58.9
5	Diabetes mellitus (E10-E14)	177	6.0	39.5
6	Cerebrovascular diseases (I60-I69)	80	2.7	17.8
7	Chronic lower respiratory diseases (J40-J47)	75	2.5	16.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	2.3	14.9
9	Viral hepatitis (B15-B19)	57	1.9	12.7
10	Influenza and pneumonia (J09-J18)	53	1.8	11.8
11	Septicemia (A40-A41)	49	1.7	10.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	1.7	10.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.3	8.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.7	4.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.6	*
...	All other causes (Residual)	446	15.1	99.4
American Indian, male, 65 years and over				
...	All causes	3,673	100.0	3,002.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	924	25.2	755.4
2	Malignant neoplasms (C00-C97)	875	23.8	715.4
3	Diabetes mellitus (E10-E14)	226	6.2	184.8
4	Chronic lower respiratory diseases (J40-J47)	220	6.0	179.9
5	Cerebrovascular diseases (I60-I69)	162	4.4	132.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	3.9	116.1
7	Influenza and pneumonia (J09-J18)	104	2.8	85.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	2.3	69.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	70	1.9	57.2
10	Alzheimer's disease (G30)	65	1.8	53.1
11	Septicemia (A40-A41)	49	1.3	40.1
12	Parkinson's disease (G20-G21)	39	1.1	31.9
13	Pneumonitis due to solids and liquids (J69)	31	0.8	25.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.8	22.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.7	22.1
...	All other causes (Residual)	626	17.0	511.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	7,307	100.0	338.1
1	Malignant neoplasms (C00-C97)	1,324	18.1	61.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,238	16.9	57.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	8.2	27.9
4	Diabetes mellitus (E10-E14)	474	6.5	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	368	5.0	17.0
6	Chronic lower respiratory diseases (J40-J47)	353	4.8	16.3
7	Cerebrovascular diseases (I60-I69)	343	4.7	15.9
8	Influenza and pneumonia (J09-J18)	172	2.4	8.0
9	Alzheimer's disease (G30)	161	2.2	7.4
10	Septicemia (A40-A41)	160	2.2	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.2	7.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	1.5	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.0	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.8	2.7
15	Certain conditions originating in the perinatal period (P00-P96)	53	0.7	2.5
...	All other causes (Residual)	1,664	22.8	77.0
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	25.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.7	*
4	Septicemia (A40-A41)	2	5.1	*
4	Malignant neoplasms (C00-C97)	2	5.1	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
7	Meningococcal infection (A39)	1	2.6	*
7	Cerebrovascular diseases (I60-I69)	1	2.6	*
7	Infections of kidney (N10-N12,N13.6,N15.1)	1	2.6	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	12	30.8	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 5-14 years				
...	All causes	46	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	37.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	13.0	*
3	Malignant neoplasms (C00-C97)	3	6.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
5	Cerebrovascular diseases (I60-I69)	2	4.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
8	Septicemia (A40-A41)	1	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.2	*
...	All other causes (Residual)	9	19.6	*
American Indian, female, 15-24 years				
...	All causes	167	100.0	45.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	52.1	23.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	15.6	7.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	5.4	*
4	Malignant neoplasms (C00-C97)	6	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.4	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.4	*
7	Cerebrovascular diseases (I60-I69)	3	1.8	*
8	Influenza and pneumonia (J09-J18)	2	1.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.2	*
10	Diabetes mellitus (E10-E14)	1	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
10	Peptic ulcer (K25-K28)	1	0.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	20	12.0	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-44 years				
...	All causes	750	100.0	124.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	26.8	33.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	114	15.2	18.9
3	Malignant neoplasms (C00-C97)	63	8.4	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	7.2	8.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	6.3	7.8
6	Diabetes mellitus (E10-E14)	30	4.0	5.0
7	Cerebrovascular diseases (I60-I69)	19	2.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	2.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	14	1.9	*
10	Septicemia (A40-A41)	13	1.7	*
10	Influenza and pneumonia (J09-J18)	13	1.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
...	All other causes (Residual)	140	18.7	23.2
American Indian, female, 45-64 years				
...	All causes	2,106	100.0	449.4
1	Malignant neoplasms (C00-C97)	498	23.6	106.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	14.4	64.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	200	9.5	42.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	8.1	36.3
5	Diabetes mellitus (E10-E14)	156	7.4	33.3
6	Chronic lower respiratory diseases (J40-J47)	92	4.4	19.6
7	Cerebrovascular diseases (I60-I69)	68	3.2	14.5
8	Septicemia (A40-A41)	58	2.8	12.4
9	Influenza and pneumonia (J09-J18)	39	1.9	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.8	7.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	1.3	5.8
12	Viral hepatitis (B15-B19)	26	1.2	5.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.4	*
15	Pneumonitis due to solids and liquids (J69)	8	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	8	0.4	*
...	All other causes (Residual)	395	18.8	84.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65 years and over				
...	All causes	4,035	100.0	2,643.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	873	21.6	571.9
2	Malignant neoplasms (C00-C97)	751	18.6	492.0
3	Diabetes mellitus (E10-E14)	287	7.1	188.0
4	Chronic lower respiratory diseases (J40-J47)	255	6.3	167.1
5	Cerebrovascular diseases (I60-I69)	249	6.2	163.1
6	Alzheimer's disease (G30)	157	3.9	102.9
7	Influenza and pneumonia (J09-J18)	115	2.9	75.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.8	74.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	2.5	65.5
10	Septicemia (A40-A41)	84	2.1	55.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.5	40.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	52	1.3	34.1
13	Pneumonitis due to solids and liquids (J69)	34	0.8	22.3
14	Parkinson's disease (G20-G21)	33	0.8	21.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.6	16.4
...	All other causes (Residual)	844	20.9	552.9
Asian or Pacific Islander, both sexes, all ages				
...	All causes	53,336	100.0	305.3
1	Malignant neoplasms (C00-C97)	14,713	27.6	84.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,586	21.7	66.3
3	Cerebrovascular diseases (I60-I69)	3,937	7.4	22.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,327	4.4	13.3
5	Diabetes mellitus (E10-E14)	2,035	3.8	11.6
6	Chronic lower respiratory diseases (J40-J47)	1,705	3.2	9.8
7	Influenza and pneumonia (J09-J18)	1,652	3.1	9.5
8	Alzheimer's disease (G30)	1,211	2.3	6.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,043	2.0	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	949	1.8	5.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	878	1.6	5.0
12	Septicemia (A40-A41)	613	1.1	3.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	498	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	482	0.9	2.8
15	Parkinson's disease (G20-G21)	460	0.9	2.6
...	All other causes (Residual)	9,247	17.3	52.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	129	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	16.3	2.2
2	Malignant neoplasms (C00-C97)	18	14.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	12.4	*
4	Influenza and pneumonia (J09-J18)	11	8.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	7.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.1	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.1	*
8	Septicemia (A40-A41)	2	1.6	*
8	Anemias (D50-D64)	2	1.6	*
10	Tuberculosis (A16-A19)	1	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
...	All other causes (Residual)	40	31.0	4.2
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	197	100.0	8.5
1	Malignant neoplasms (C00-C97)	42	21.3	1.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	20.3	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	5.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	5.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	4.6	*
7	Influenza and pneumonia (J09-J18)	8	4.1	*
8	Anemias (D50-D64)	6	3.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	2.5	*
10	Cerebrovascular diseases (I60-I69)	4	2.0	*
11	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.0	*
13	Salmonella infections (A01-A02)	1	0.5	*
13	Septicemia (A40-A41)	1	0.5	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (Residual)	45	22.8	1.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	734	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	37.2	10.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	21.8	6.4
3	Malignant neoplasms (C00-C97)	70	9.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	8.7	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.4	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.6	*
7	Cerebrovascular diseases (I60-I69)	8	1.1	*
8	Septicemia (A40-A41)	6	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
8	Influenza and pneumonia (J09-J18)	6	0.8	*
11	Aortic aneurysm and dissection (I71)	5	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.3	*
13	Legal intervention (Y35,Y89.0)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	68	9.3	2.7
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,029	100.0	52.8
1	Malignant neoplasms (C00-C97)	734	24.2	12.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	504	16.6	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	13.8	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	12.6	6.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	5.2	2.8
6	Cerebrovascular diseases (I60-I69)	129	4.3	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.6	0.9
8	Diabetes mellitus (E10-E14)	41	1.4	0.7
9	Influenza and pneumonia (J09-J18)	39	1.3	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	36	1.2	0.6
11	Septicemia (A40-A41)	30	1.0	0.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.9	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.8	0.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	0.4
15	Viral hepatitis (B15-B19)	21	0.7	0.4
...	All other causes (Residual)	414	13.7	7.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	11,057	100.0	267.2
1	Malignant neoplasms (C00-C97)	4,711	42.6	113.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,057	18.6	49.7
3	Cerebrovascular diseases (I60-I69)	729	6.6	17.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	490	4.4	11.8
5	Diabetes mellitus (E10-E14)	419	3.8	10.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	2.8	7.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	230	2.1	5.6
8	Chronic lower respiratory diseases (J40-J47)	151	1.4	3.6
9	Viral hepatitis (B15-B19)	146	1.3	3.5
10	Influenza and pneumonia (J09-J18)	139	1.3	3.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.1	3.0
12	Septicemia (A40-A41)	118	1.1	2.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	0.9	2.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	0.6	1.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.6	1.5
15	Aortic aneurysm and dissection (I71)	64	0.6	1.5
...	All other causes (Residual)	1,136	10.3	27.5
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	37,307	100.0	2,315.3
1	Malignant neoplasms (C00-C97)	9,133	24.5	566.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,081	24.3	563.6
3	Cerebrovascular diseases (I60-I69)	3,065	8.2	190.2
4	Diabetes mellitus (E10-E14)	1,571	4.2	97.5
5	Chronic lower respiratory diseases (J40-J47)	1,522	4.1	94.5
6	Influenza and pneumonia (J09-J18)	1,443	3.9	89.6
7	Alzheimer's disease (G30)	1,200	3.2	74.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	975	2.6	60.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	794	2.1	49.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	774	2.1	48.0
11	Septicemia (A40-A41)	446	1.2	27.7
12	Parkinson's disease (G20-G21)	444	1.2	27.6
13	Pneumonitis due to solids and liquids (J69)	327	0.9	20.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	303	0.8	18.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	219	0.6	13.6
...	All other causes (Residual)	6,010	16.1	373.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	27,490	100.0	328.6
1	Malignant neoplasms (C00-C97)	7,444	27.1	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,209	22.6	74.2
3	Cerebrovascular diseases (I60-I69)	1,775	6.5	21.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,520	5.5	18.2
5	Diabetes mellitus (E10-E14)	1,040	3.8	12.4
6	Chronic lower respiratory diseases (J40-J47)	1,008	3.7	12.0
7	Influenza and pneumonia (J09-J18)	854	3.1	10.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	2.6	8.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	1.8	5.8
10	Alzheimer's disease (G30)	404	1.5	4.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	1.3	4.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.2	3.9
13	Septicemia (A40-A41)	304	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.3
15	Parkinson's disease (G20-G21)	263	1.0	3.1
...	All other causes (Residual)	4,503	16.4	53.8
Asian or Pacific Islander, male, 1-4 years				
...	All causes	74	100.0	15.4
1	Malignant neoplasms (C00-C97)	13	17.6	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	14.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	12.2	*
4	Influenza and pneumonia (J09-J18)	4	5.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.4	*
6	Anemias (D50-D64)	2	2.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.4	*
...	All other causes (Residual)	27	36.5	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	107	100.0	9.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	23.4	2.2
2	Malignant neoplasms (C00-C97)	21	19.6	1.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.7	*
7	Influenza and pneumonia (J09-J18)	3	2.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.8	*
9	Anemias (D50-D64)	2	1.9	*
10	Salmonella infections (A01-A02)	1	0.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	Diabetes mellitus (E10-E14)	1	0.9	*
...	All other causes (Residual)	28	26.2	2.4
Asian or Pacific Islander, male, 15-24 years				
...	All causes	539	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	40.6	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	22.6	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.8	4.2
4	Malignant neoplasms (C00-C97)	42	7.8	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.3	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.9	*
7	Cerebrovascular diseases (I60-I69)	5	0.9	*
7	Aortic aneurysm and dissection (I71)	5	0.9	*
9	Septicemia (A40-A41)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Legal intervention (Y35,Y89.0)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	39	7.2	3.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-44 years				
...	All causes	1,872	100.0	68.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	20.8	14.3
2	Malignant neoplasms (C00-C97)	309	16.5	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	296	15.8	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	288	15.4	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	6.0	4.1
6	Cerebrovascular diseases (I60-I69)	68	3.6	2.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	39	2.1	1.4
8	Diabetes mellitus (E10-E14)	28	1.5	1.0
9	Influenza and pneumonia (J09-J18)	20	1.1	0.7
10	Viral hepatitis (B15-B19)	18	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.9	*
12	Septicemia (A40-A41)	16	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	14	0.7	*
...	All other causes (Residual)	226	12.1	8.3
Asian or Pacific Islander, male, 45-64 years				
...	All causes	6,459	100.0	337.9
1	Malignant neoplasms (C00-C97)	2,313	35.8	121.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,495	23.1	78.2
3	Cerebrovascular diseases (I60-I69)	396	6.1	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	337	5.2	17.6
5	Diabetes mellitus (E10-E14)	269	4.2	14.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	3.3	11.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	182	2.8	9.5
8	Viral hepatitis (B15-B19)	101	1.6	5.3
9	Chronic lower respiratory diseases (J40-J47)	89	1.4	4.7
10	Influenza and pneumonia (J09-J18)	85	1.3	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.1	3.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	1.0	3.3
13	Septicemia (A40-A41)	62	1.0	3.2
14	Aortic aneurysm and dissection (I71)	46	0.7	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	0.6	2.0
...	All other causes (Residual)	697	10.8	36.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65 years and over				
...	All causes	17,950	100.0	2,570.2
1	Malignant neoplasms (C00-C97)	4,743	26.4	679.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,380	24.4	627.1
3	Cerebrovascular diseases (I60-I69)	1,305	7.3	186.9
4	Chronic lower respiratory diseases (J40-J47)	900	5.0	128.9
5	Diabetes mellitus (E10-E14)	740	4.1	106.0
6	Influenza and pneumonia (J09-J18)	736	4.1	105.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	528	2.9	75.6
8	Alzheimer's disease (G30)	399	2.2	57.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.2	57.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	283	1.6	40.5
11	Parkinson's disease (G20-G21)	254	1.4	36.4
12	Septicemia (A40-A41)	215	1.2	30.8
13	Pneumonitis due to solids and liquids (J69)	192	1.1	27.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.8	21.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	105	0.6	15.0
...	All other causes (Residual)	2,619	14.6	375.0
Asian or Pacific Islander, female, all ages				
...	All causes	25,846	100.0	283.8
1	Malignant neoplasms (C00-C97)	7,269	28.1	79.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,377	20.8	59.1
3	Cerebrovascular diseases (I60-I69)	2,162	8.4	23.7
4	Diabetes mellitus (E10-E14)	995	3.8	10.9
5	Alzheimer's disease (G30)	807	3.1	8.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.1	8.9
7	Influenza and pneumonia (J09-J18)	798	3.1	8.8
8	Chronic lower respiratory diseases (J40-J47)	697	2.7	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	525	2.0	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	1.8	5.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	1.2	3.5
12	Septicemia (A40-A41)	309	1.2	3.4
13	Certain conditions originating in the perinatal period (P00-P96)	204	0.8	2.2
14	Parkinson's disease (G20-G21)	197	0.8	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	191	0.7	2.1
...	All other causes (Residual)	4,725	18.3	51.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	55	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	18.2	*
2	Influenza and pneumonia (J09-J18)	7	12.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	12.7	*
4	Malignant neoplasms (C00-C97)	5	9.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
7	Septicemia (A40-A41)	2	3.6	*
7	Chronic lower respiratory diseases (J40-J47)	2	3.6	*
9	Tuberculosis (A16-A19)	1	1.8	*
...	All other causes (Residual)	13	23.6	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	90	100.0	7.9
1	Malignant neoplasms (C00-C97)	21	23.3	1.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	16.7	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.6	*
3	Influenza and pneumonia (J09-J18)	5	5.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.6	*
6	Anemias (D50-D64)	4	4.4	*
6	Cerebrovascular diseases (I60-I69)	4	4.4	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	4.4	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.3	*
10	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	2	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
13	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	17	18.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	195	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	27.7	4.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	19.5	3.1
3	Malignant neoplasms (C00-C97)	28	14.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	4.6	*
7	Influenza and pneumonia (J09-J18)	4	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.5	*
9	Cerebrovascular diseases (I60-I69)	3	1.5	*
11	Septicemia (A40-A41)	2	1.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.0	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Meningitis (G00,G03)	1	0.5	*
13	Pneumonitis due to solids and liquids (J69)	1	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (Residual)	24	12.3	2.0
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,157	100.0	38.4
1	Malignant neoplasms (C00-C97)	425	36.7	14.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	11.2	4.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	9.9	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	7.5	2.9
5	Cerebrovascular diseases (I60-I69)	61	5.3	2.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	4.0	1.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	36	3.1	1.2
8	Influenza and pneumonia (J09-J18)	19	1.6	*
9	Septicemia (A40-A41)	14	1.2	*
10	Diabetes mellitus (E10-E14)	13	1.1	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.6	*
...	All other causes (Residual)	169	14.6	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-64 years				
...	All causes	4,598	100.0	206.5
1	Malignant neoplasms (C00-C97)	2,398	52.2	107.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	562	12.2	25.2
3	Cerebrovascular diseases (I60-I69)	333	7.2	15.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	3.3	6.9
5	Diabetes mellitus (E10-E14)	150	3.3	6.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	2.2	4.6
7	Chronic lower respiratory diseases (J40-J47)	62	1.3	2.8
8	Septicemia (A40-A41)	56	1.2	2.5
9	Influenza and pneumonia (J09-J18)	54	1.2	2.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.2	2.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	48	1.0	2.2
12	Viral hepatitis (B15-B19)	45	1.0	2.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	0.7	1.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	0.6	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	1.3
...	All other causes (Residual)	493	10.7	22.1
Asian or Pacific Islander, female, 65 years and over				
...	All causes	19,357	100.0	2,120.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,701	24.3	514.9
2	Malignant neoplasms (C00-C97)	4,390	22.7	480.9
3	Cerebrovascular diseases (I60-I69)	1,760	9.1	192.8
4	Diabetes mellitus (E10-E14)	831	4.3	91.0
5	Alzheimer's disease (G30)	801	4.1	87.7
6	Influenza and pneumonia (J09-J18)	707	3.7	77.4
7	Chronic lower respiratory diseases (J40-J47)	622	3.2	68.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	491	2.5	53.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	447	2.3	49.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	395	2.0	43.3
11	Septicemia (A40-A41)	231	1.2	25.3
12	Parkinson's disease (G20-G21)	190	1.0	20.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	151	0.8	16.5
14	Pneumonitis due to solids and liquids (J69)	135	0.7	14.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	114	0.6	12.5
...	All other causes (Residual)	3,391	17.5	371.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011