

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,543,279	100.0	810.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	599,711	23.6	191.0
2	Malignant neoplasms (C00-C97)	582,623	22.9	185.6
3	Chronic lower respiratory diseases (J40-J47)	143,489	5.6	45.7
4	Cerebrovascular diseases (I60-I69)	128,546	5.1	40.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,792	5.0	40.7
6	Alzheimer's disease (G30)	83,637	3.3	26.6
7	Diabetes mellitus (E10-E14)	73,932	2.9	23.6
8	Influenza and pneumonia (J09-J18)	50,636	2.0	16.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,622	1.8	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40,600	1.6	12.9
11	Septicemia (A40-A41)	35,842	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34,979	1.4	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29,115	1.1	9.3
14	Parkinson's disease (G20-G21)	23,818	0.9	7.6
15	Pneumonitis due to solids and liquids (J69)	17,897	0.7	5.7
...	All other causes (Residual)	525,040	20.6	167.3
All races, both sexes, 1-4 years				
...	All causes	4,218	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,353	32.1	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	501	11.9	3.1
3	Malignant neoplasms (C00-C97)	392	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	3.7	1.0
6	Influenza and pneumonia (J09-J18)	93	2.2	0.6
7	Septicemia (A40-A41)	62	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	42	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	17	0.4	*
13	Anemias (D50-D64)	15	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (Residual)	1,065	25.2	6.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,200	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,550	29.8	3.8
2	Malignant neoplasms (C00-C97)	912	17.5	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	327	6.3	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	311	6.0	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	311	6.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	3.4	0.4
7	Chronic lower respiratory diseases (J40-J47)	119	2.3	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	85	1.6	0.2
9	Influenza and pneumonia (J09-J18)	85	1.6	0.2
11	Septicemia (A40-A41)	52	1.0	0.1
12	Diabetes mellitus (E10-E14)	34	0.7	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	34	0.7	0.1
14	Anemias (D50-D64)	28	0.5	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
...	All other causes (Residual)	1,068	20.5	2.6
All races, both sexes, 15-24 years				
...	All causes	29,182	100.0	66.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,908	40.8	27.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,872	16.7	11.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,614	15.8	10.5
4	Malignant neoplasms (C00-C97)	1,574	5.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	956	3.3	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	423	1.4	1.0
7	Diabetes mellitus (E10-E14)	196	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	183	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	169	0.6	0.4
10	Influenza and pneumonia (J09-J18)	147	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	143	0.5	0.3
12	Septicemia (A40-A41)	120	0.4	0.3
13	Legal intervention (Y35,Y89.0)	118	0.4	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	116	0.4	0.3
15	Anemias (D50-D64)	89	0.3	0.2
...	All other causes (Residual)	3,554	12.2	8.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	113,753	100.0	137.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,885	27.2	37.3
2	Malignant neoplasms (C00-C97)	15,011	13.2	18.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,720	12.1	16.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,974	11.4	15.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,047	6.2	8.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,066	2.7	3.7
7	Diabetes mellitus (E10-E14)	2,513	2.2	3.0
8	Cerebrovascular diseases (I60-I69)	2,265	2.0	2.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,997	1.8	2.4
10	Septicemia (A40-A41)	1,082	1.0	1.3
11	Influenza and pneumonia (J09-J18)	1,041	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	1,032	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	881	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	797	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	663	0.6	0.8
...	All other causes (Residual)	18,779	16.5	22.7
All races, both sexes, 45-64 years				
...	All causes	509,069	100.0	614.4
1	Malignant neoplasms (C00-C97)	161,158	31.7	194.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,493	20.9	128.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,216	7.1	43.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	20,107	3.9	24.3
5	Chronic lower respiratory diseases (J40-J47)	19,745	3.9	23.8
6	Diabetes mellitus (E10-E14)	18,300	3.6	22.1
7	Cerebrovascular diseases (I60-I69)	16,724	3.3	20.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15,791	3.1	19.1
9	Septicemia (A40-A41)	7,322	1.4	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,829	1.3	8.2
11	Viral hepatitis (B15-B19)	5,944	1.2	7.2
12	Influenza and pneumonia (J09-J18)	5,754	1.1	6.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,375	0.9	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,341	0.9	5.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,205	0.6	3.9
...	All other causes (Residual)	76,765	15.1	92.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,858,081	100.0	4,306.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	477,840	25.7	1,107.5
2	Malignant neoplasms (C00-C97)	403,497	21.7	935.2
3	Chronic lower respiratory diseases (J40-J47)	122,375	6.6	283.6
4	Cerebrovascular diseases (I60-I69)	109,127	5.9	252.9
5	Alzheimer's disease (G30)	82,690	4.5	191.7
6	Diabetes mellitus (E10-E14)	52,881	2.8	122.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,698	2.4	103.6
8	Influenza and pneumonia (J09-J18)	43,355	2.3	100.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,740	2.0	87.5
10	Septicemia (A40-A41)	27,022	1.5	62.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,312	1.3	56.3
12	Parkinson's disease (G20-G21)	23,203	1.2	53.8
13	Pneumonitis due to solids and liquids (J69)	15,837	0.9	36.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12,246	0.7	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11,770	0.6	27.3
...	All other causes (Residual)	369,488	19.9	856.4
All races, male, all ages				
...	All causes	1,273,722	100.0	824.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	312,491	24.5	202.3
2	Malignant neoplasms (C00-C97)	305,670	24.0	197.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,010	6.3	51.8
4	Chronic lower respiratory diseases (J40-J47)	67,673	5.3	43.8
5	Cerebrovascular diseases (I60-I69)	52,638	4.1	34.1
6	Diabetes mellitus (E10-E14)	38,584	3.0	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,780	2.5	20.6
8	Alzheimer's disease (G30)	25,653	2.0	16.6
9	Influenza and pneumonia (J09-J18)	24,013	1.9	15.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,731	1.8	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22,623	1.8	14.6
12	Septicemia (A40-A41)	16,789	1.3	10.9
13	Parkinson's disease (G20-G21)	14,121	1.1	9.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,208	1.0	8.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,859	0.9	7.7
...	All other causes (Residual)	233,879	18.4	151.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,395	100.0	29.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	33.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	11.5	3.4
3	Malignant neoplasms (C00-C97)	218	9.1	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	8.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Cerebrovascular diseases (I60-I69)	36	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	35	1.5	0.4
9	Septicemia (A40-A41)	29	1.2	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	27	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.8	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	13	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (Residual)	580	24.2	7.1
All races, male, 5-14 years				
...	All causes	3,036	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	982	32.3	4.7
2	Malignant neoplasms (C00-C97)	513	16.9	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	7.4	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	6.0	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	170	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	62	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.4	0.2
10	Influenza and pneumonia (J09-J18)	40	1.3	0.2
11	Septicemia (A40-A41)	32	1.1	0.2
12	Diabetes mellitus (E10-E14)	18	0.6	*
13	Anemias (D50-D64)	16	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
...	All other causes (Residual)	588	19.4	2.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,456	100.0	95.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,849	41.2	39.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,995	18.6	17.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,918	18.3	17.4
4	Malignant neoplasms (C00-C97)	924	4.3	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	3.0	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	250	1.2	1.1
7	Legal intervention (Y35,Y89.0)	112	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	108	0.5	0.5
9	Diabetes mellitus (E10-E14)	106	0.5	0.5
10	Chronic lower respiratory diseases (J40-J47)	98	0.5	0.4
11	Influenza and pneumonia (J09-J18)	90	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.4	0.3
13	Septicemia (A40-A41)	54	0.3	0.2
14	Anemias (D50-D64)	50	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.2	0.2
...	All other causes (Residual)	2,131	9.9	9.5
All races, male, 25-44 years				
...	All causes	73,712	100.0	177.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,148	30.0	53.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,170	13.8	24.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,445	12.8	22.8
4	Malignant neoplasms (C00-C97)	6,651	9.0	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,789	7.9	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,980	2.7	4.8
7	Diabetes mellitus (E10-E14)	1,515	2.1	3.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,309	1.8	3.2
9	Cerebrovascular diseases (I60-I69)	1,259	1.7	3.0
10	Septicemia (A40-A41)	567	0.8	1.4
11	Influenza and pneumonia (J09-J18)	563	0.8	1.4
12	Chronic lower respiratory diseases (J40-J47)	511	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	498	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	466	0.6	1.1
15	Aortic aneurysm and dissection (I71)	321	0.4	0.8
...	All other causes (Residual)	10,520	14.3	25.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	310,165	100.0	767.5
1	Malignant neoplasms (C00-C97)	86,673	27.9	214.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,484	24.0	184.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,395	7.9	60.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	13,918	4.5	34.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,910	3.8	29.5
6	Diabetes mellitus (E10-E14)	11,238	3.6	27.8
7	Chronic lower respiratory diseases (J40-J47)	10,014	3.2	24.8
8	Cerebrovascular diseases (I60-I69)	9,511	3.1	23.5
9	Viral hepatitis (B15-B19)	4,226	1.4	10.5
10	Septicemia (A40-A41)	3,953	1.3	9.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,900	1.3	9.7
12	Influenza and pneumonia (J09-J18)	3,321	1.1	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,265	1.1	8.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,738	0.9	6.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,357	0.8	5.8
...	All other causes (Residual)	44,262	14.3	109.5
All races, male, 65 years and over				
...	All causes	849,719	100.0	4,516.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	227,519	26.8	1,209.2
2	Malignant neoplasms (C00-C97)	210,648	24.8	1,119.6
3	Chronic lower respiratory diseases (J40-J47)	56,938	6.7	302.6
4	Cerebrovascular diseases (I60-I69)	41,628	4.9	221.2
5	Diabetes mellitus (E10-E14)	25,702	3.0	136.6
6	Alzheimer's disease (G30)	25,235	3.0	134.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,144	2.6	117.7
8	Influenza and pneumonia (J09-J18)	19,847	2.3	105.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,242	2.1	97.0
10	Parkinson's disease (G20-G21)	13,739	1.6	73.0
11	Septicemia (A40-A41)	12,048	1.4	64.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,843	1.0	47.0
13	Pneumonitis due to solids and liquids (J69)	8,435	1.0	44.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,699	0.8	35.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,332	0.7	33.7
...	All other causes (Residual)	145,720	17.1	774.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,269,557	100.0	796.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,220	22.6	180.2
2	Malignant neoplasms (C00-C97)	276,953	21.8	173.7
3	Cerebrovascular diseases (I60-I69)	75,908	6.0	47.6
4	Chronic lower respiratory diseases (J40-J47)	75,816	6.0	47.6
5	Alzheimer's disease (G30)	57,984	4.6	36.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,782	3.8	30.0
7	Diabetes mellitus (E10-E14)	35,348	2.8	22.2
8	Influenza and pneumonia (J09-J18)	26,623	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,891	1.8	14.4
10	Septicemia (A40-A41)	19,053	1.5	12.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,256	1.4	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,356	1.0	7.8
13	Parkinson's disease (G20-G21)	9,697	0.8	6.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,820	0.7	5.5
15	Pneumonitis due to solids and liquids (J69)	8,184	0.6	5.1
...	All other causes (Residual)	287,666	22.7	180.4
All races, female, 1-4 years				
...	All causes	1,823	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	547	30.0	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	12.4	2.9
3	Malignant neoplasms (C00-C97)	174	9.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	7.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	4.1	0.9
6	Influenza and pneumonia (J09-J18)	38	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.9	0.4
8	Septicemia (A40-A41)	33	1.8	0.4
9	Cerebrovascular diseases (I60-I69)	20	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	15	0.8	*
12	Anemias (D50-D64)	10	0.5	*
13	Meningitis (G00,G03)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	474	26.0	6.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,164	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	26.2	2.8
2	Malignant neoplasms (C00-C97)	399	18.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	7.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	6.0	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	3.9	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.7	0.4
7	Chronic lower respiratory diseases (J40-J47)	57	2.6	0.3
8	Influenza and pneumonia (J09-J18)	45	2.1	0.2
9	Cerebrovascular diseases (I60-I69)	44	2.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.8	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	23	1.1	0.1
12	Septicemia (A40-A41)	20	0.9	0.1
13	Diabetes mellitus (E10-E14)	16	0.7	*
14	Anemias (D50-D64)	12	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
...	All other causes (Residual)	480	22.2	2.4
All races, female, 15-24 years				
...	All causes	7,726	100.0	36.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,059	39.6	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	954	12.3	4.5
3	Malignant neoplasms (C00-C97)	650	8.4	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	619	8.0	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	312	4.0	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	2.2	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	169	2.2	0.8
8	Diabetes mellitus (E10-E14)	90	1.2	0.4
9	Cerebrovascular diseases (I60-I69)	75	1.0	0.3
10	Septicemia (A40-A41)	66	0.9	0.3
11	Influenza and pneumonia (J09-J18)	57	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	45	0.6	0.2
13	Anemias (D50-D64)	39	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.5	0.2
...	All other causes (Residual)	1,342	17.4	6.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	40,041	100.0	96.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,737	21.8	21.1
2	Malignant neoplasms (C00-C97)	8,360	20.9	20.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,275	10.7	10.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,804	7.0	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,258	3.1	3.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,086	2.7	2.6
7	Cerebrovascular diseases (I60-I69)	1,006	2.5	2.4
8	Diabetes mellitus (E10-E14)	998	2.5	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	688	1.7	1.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	663	1.7	1.6
11	Chronic lower respiratory diseases (J40-J47)	521	1.3	1.3
12	Septicemia (A40-A41)	515	1.3	1.2
13	Influenza and pneumonia (J09-J18)	478	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	383	1.0	0.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	331	0.8	0.8
...	All other causes (Residual)	7,938	19.8	19.2
All races, female, 45-64 years				
...	All causes	198,904	100.0	468.6
1	Malignant neoplasms (C00-C97)	74,485	37.4	175.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,009	16.1	75.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,821	5.9	27.9
4	Chronic lower respiratory diseases (J40-J47)	9,731	4.9	22.9
5	Cerebrovascular diseases (I60-I69)	7,213	3.6	17.0
6	Diabetes mellitus (E10-E14)	7,062	3.6	16.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,189	3.1	14.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,881	2.0	9.1
9	Septicemia (A40-A41)	3,369	1.7	7.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,929	1.5	6.9
11	Influenza and pneumonia (J09-J18)	2,433	1.2	5.7
12	Viral hepatitis (B15-B19)	1,718	0.9	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,603	0.8	3.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,110	0.6	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	962	0.5	2.3
...	All other causes (Residual)	32,389	16.3	76.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	1,008,362	100.0	4,144.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,321	24.8	1,028.8
2	Malignant neoplasms (C00-C97)	192,849	19.1	792.6
3	Cerebrovascular diseases (I60-I69)	67,499	6.7	277.4
4	Chronic lower respiratory diseases (J40-J47)	65,437	6.5	269.0
5	Alzheimer's disease (G30)	57,455	5.7	236.1
6	Diabetes mellitus (E10-E14)	27,179	2.7	111.7
7	Influenza and pneumonia (J09-J18)	23,508	2.3	96.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,554	2.2	92.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,498	1.9	80.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,469	1.5	63.6
11	Septicemia (A40-A41)	14,974	1.5	61.5
12	Parkinson's disease (G20-G21)	9,464	0.9	38.9
13	Pneumonitis due to solids and liquids (J69)	7,402	0.7	30.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,914	0.6	24.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,071	0.5	20.8
...	All other causes (Residual)	223,768	22.2	919.7
White, both sexes, all ages				
...	All causes	2,175,178	100.0	876.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,372	23.6	207.2
2	Malignant neoplasms (C00-C97)	496,885	22.8	200.2
3	Chronic lower respiratory diseases (J40-J47)	131,782	6.1	53.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110,789	5.1	44.6
5	Cerebrovascular diseases (I60-I69)	107,964	5.0	43.5
6	Alzheimer's disease (G30)	76,590	3.5	30.9
7	Diabetes mellitus (E10-E14)	57,806	2.7	23.3
8	Influenza and pneumonia (J09-J18)	43,585	2.0	17.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,606	1.7	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,066	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,747	1.4	12.4
12	Septicemia (A40-A41)	28,945	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,908	1.1	9.2
14	Parkinson's disease (G20-G21)	22,352	1.0	9.0
15	Pneumonitis due to solids and liquids (J69)	15,938	0.7	6.4
...	All other causes (Residual)	441,843	20.3	178.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,958	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,004	33.9	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	366	12.4	3.0
3	Malignant neoplasms (C00-C97)	301	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	189	6.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.4	0.8
6	Influenza and pneumonia (J09-J18)	67	2.3	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.5	0.4
8	Septicemia (A40-A41)	37	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	35	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	0.8	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Pneumonitis due to solids and liquids (J69)	7	0.2	*
15	Anemias (D50-D64)	6	0.2	*
15	Diabetes mellitus (E10-E14)	6	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (Residual)	726	24.5	6.0
White, both sexes, 5-14 years				
...	All causes	3,711	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,087	29.3	3.5
2	Malignant neoplasms (C00-C97)	712	19.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	6.7	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	6.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	5.3	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.1	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	2.0	0.2
8	Cerebrovascular diseases (I60-I69)	67	1.8	0.2
9	Influenza and pneumonia (J09-J18)	63	1.7	0.2
10	Septicemia (A40-A41)	43	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	41	1.1	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	24	0.6	0.1
13	Diabetes mellitus (E10-E14)	17	0.5	*
14	Meningitis (G00,G03)	14	0.4	*
15	Anemias (D50-D64)	11	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.3	*
...	All other causes (Residual)	758	20.4	2.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,101	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,001	47.4	30.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,001	19.0	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,579	7.5	4.7
4	Malignant neoplasms (C00-C97)	1,218	5.8	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	613	2.9	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	321	1.5	1.0
7	Cerebrovascular diseases (I60-I69)	133	0.6	0.4
8	Diabetes mellitus (E10-E14)	131	0.6	0.4
9	Influenza and pneumonia (J09-J18)	110	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	98	0.5	0.3
11	Septicemia (A40-A41)	81	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.2
13	Legal intervention (Y35,Y89.0)	65	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,572	12.2	7.7
White, both sexes, 25-44 years				
...	All causes	85,598	100.0	133.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,256	30.7	41.1
2	Malignant neoplasms (C00-C97)	11,467	13.4	17.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,281	13.2	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,417	11.0	14.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,060	3.6	4.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,549	3.0	4.0
7	Diabetes mellitus (E10-E14)	1,692	2.0	2.6
8	Cerebrovascular diseases (I60-I69)	1,514	1.8	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	768	0.9	1.2
10	Influenza and pneumonia (J09-J18)	763	0.9	1.2
11	Septicemia (A40-A41)	729	0.9	1.1
12	Chronic lower respiratory diseases (J40-J47)	694	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	647	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	495	0.6	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	385	0.4	0.6
...	All other causes (Residual)	13,881	16.2	21.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	403,796	100.0	598.7
1	Malignant neoplasms (C00-C97)	129,833	32.2	192.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,784	20.3	121.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,607	7.6	45.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	17,548	4.3	26.0
5	Chronic lower respiratory diseases (J40-J47)	17,162	4.3	25.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,760	3.7	21.9
7	Diabetes mellitus (E10-E14)	13,372	3.3	19.8
8	Cerebrovascular diseases (I60-I69)	11,475	2.8	17.0
9	Septicemia (A40-A41)	5,393	1.3	8.0
10	Viral hepatitis (B15-B19)	4,852	1.2	7.2
11	Influenza and pneumonia (J09-J18)	4,441	1.1	6.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,373	1.1	6.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,745	0.7	4.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,048	0.5	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,956	0.5	2.9
...	All other causes (Residual)	61,447	15.2	91.1
White, both sexes, 65 years and over				
...	All causes	1,642,659	100.0	4,406.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	422,104	25.7	1,132.3
2	Malignant neoplasms (C00-C97)	353,294	21.5	947.7
3	Chronic lower respiratory diseases (J40-J47)	113,774	6.9	305.2
4	Cerebrovascular diseases (I60-I69)	94,677	5.8	254.0
5	Alzheimer's disease (G30)	75,733	4.6	203.2
6	Diabetes mellitus (E10-E14)	42,587	2.6	114.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,116	2.5	110.3
8	Influenza and pneumonia (J09-J18)	38,048	2.3	102.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,091	1.9	83.4
10	Septicemia (A40-A41)	22,565	1.4	60.5
11	Parkinson's disease (G20-G21)	21,803	1.3	58.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,933	1.2	53.5
13	Pneumonitis due to solids and liquids (J69)	14,260	0.9	38.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,134	0.7	29.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10,624	0.6	28.5
...	All other causes (Residual)	329,916	20.1	885.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,085,250	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	267,970	24.7	218.0
2	Malignant neoplasms (C00-C97)	261,975	24.1	213.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,677	6.3	55.9
4	Chronic lower respiratory diseases (J40-J47)	61,452	5.7	50.0
5	Cerebrovascular diseases (I60-I69)	43,435	4.0	35.3
6	Diabetes mellitus (E10-E14)	30,832	2.8	25.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,681	2.6	23.3
8	Alzheimer's disease (G30)	23,607	2.2	19.2
9	Influenza and pneumonia (J09-J18)	20,575	1.9	16.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,973	1.8	16.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,269	1.7	14.9
12	Septicemia (A40-A41)	13,654	1.3	11.1
13	Parkinson's disease (G20-G21)	13,281	1.2	10.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,098	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	8,662	0.8	7.0
...	All other causes (Residual)	195,109	18.0	158.7
White, male, 1-4 years				
...	All causes	1,683	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	35.1	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	12.4	3.4
3	Malignant neoplasms (C00-C97)	165	9.8	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.8	0.8
6	Influenza and pneumonia (J09-J18)	42	2.5	0.7
7	Cerebrovascular diseases (I60-I69)	25	1.5	0.4
8	Septicemia (A40-A41)	19	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.2	*
...	All other causes (Residual)	408	24.2	6.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,132	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	676	31.7	4.2
2	Malignant neoplasms (C00-C97)	399	18.7	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	8.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	5.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	5.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	2.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	33	1.5	0.2
9	Influenza and pneumonia (J09-J18)	29	1.4	0.2
10	Septicemia (A40-A41)	28	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	0.9	0.1
12	Diabetes mellitus (E10-E14)	10	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
14	Meningitis (G00,G03)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	412	19.3	2.6
White, male, 15-24 years				
...	All causes	15,362	100.0	89.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,452	48.5	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,253	21.2	19.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,279	8.3	7.5
4	Malignant neoplasms (C00-C97)	705	4.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	2.7	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	77	0.5	0.4
8	Diabetes mellitus (E10-E14)	76	0.5	0.4
9	Influenza and pneumonia (J09-J18)	68	0.4	0.4
10	Legal intervention (Y35,Y89.0)	62	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	49	0.3	0.3
12	Septicemia (A40-A41)	38	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	23	0.1	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.1	0.1
...	All other causes (Residual)	1,626	10.6	9.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	56,016	100.0	172.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,827	33.6	58.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,830	15.8	27.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,620	11.8	20.4
4	Malignant neoplasms (C00-C97)	5,252	9.4	16.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,293	4.1	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,670	3.0	5.1
7	Diabetes mellitus (E10-E14)	1,039	1.9	3.2
8	Cerebrovascular diseases (I60-I69)	850	1.5	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	612	1.1	1.9
10	Influenza and pneumonia (J09-J18)	418	0.7	1.3
11	Septicemia (A40-A41)	397	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	373	0.7	1.1
13	Chronic lower respiratory diseases (J40-J47)	328	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	290	0.5	0.9
15	Aortic aneurysm and dissection (I71)	218	0.4	0.7
...	All other causes (Residual)	7,999	14.3	24.6
White, male, 45-64 years				
...	All causes	248,826	100.0	748.3
1	Malignant neoplasms (C00-C97)	70,365	28.3	211.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,542	23.5	176.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,539	8.3	61.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	12,246	4.9	36.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,128	4.5	33.5
6	Chronic lower respiratory diseases (J40-J47)	8,664	3.5	26.1
7	Diabetes mellitus (E10-E14)	8,426	3.4	25.3
8	Cerebrovascular diseases (I60-I69)	6,589	2.6	19.8
9	Viral hepatitis (B15-B19)	3,500	1.4	10.5
10	Septicemia (A40-A41)	2,954	1.2	8.9
11	Influenza and pneumonia (J09-J18)	2,566	1.0	7.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,557	1.0	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,795	0.7	5.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,692	0.7	5.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,365	0.5	4.1
...	All other causes (Residual)	35,898	14.4	108.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	752,719	100.0	4,587.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	202,145	26.9	1,231.9
2	Malignant neoplasms (C00-C97)	185,057	24.6	1,127.8
3	Chronic lower respiratory diseases (J40-J47)	52,370	7.0	319.2
4	Cerebrovascular diseases (I60-I69)	35,829	4.8	218.3
5	Alzheimer's disease (G30)	23,230	3.1	141.6
6	Diabetes mellitus (E10-E14)	21,277	2.8	129.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,181	2.7	123.0
8	Influenza and pneumonia (J09-J18)	17,396	2.3	106.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,359	2.0	93.6
10	Parkinson's disease (G20-G21)	12,936	1.7	78.8
11	Septicemia (A40-A41)	10,165	1.4	61.9
12	Pneumonitis due to solids and liquids (J69)	7,614	1.0	46.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,155	1.0	43.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,039	0.8	36.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,826	0.8	35.5
...	All other causes (Residual)	130,140	17.3	793.1
White, female, all ages				
...	All causes	1,089,928	100.0	869.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246,402	22.6	196.7
2	Malignant neoplasms (C00-C97)	234,910	21.6	187.5
3	Chronic lower respiratory diseases (J40-J47)	70,330	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	64,529	5.9	51.5
5	Alzheimer's disease (G30)	52,983	4.9	42.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,112	3.9	33.6
7	Diabetes mellitus (E10-E14)	26,974	2.5	21.5
8	Influenza and pneumonia (J09-J18)	23,010	2.1	18.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,797	1.6	14.2
10	Septicemia (A40-A41)	15,291	1.4	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,810	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,774	1.0	8.6
13	Parkinson's disease (G20-G21)	9,071	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,925	0.7	6.3
15	Pneumonitis due to solids and liquids (J69)	7,276	0.7	5.8
...	All other causes (Residual)	246,734	22.6	196.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,275	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	32.4	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	12.4	2.7
3	Malignant neoplasms (C00-C97)	136	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	6.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.2	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.2	0.5
7	Influenza and pneumonia (J09-J18)	25	2.0	0.4
8	Septicemia (A40-A41)	18	1.4	*
9	Cerebrovascular diseases (I60-I69)	10	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
13	Anemias (D50-D64)	3	0.2	*
13	Meningitis (G00,G03)	3	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
13	Pneumonitis due to solids and liquids (J69)	3	0.2	*
...	All other causes (Residual)	319	25.0	5.4
White, female, 5-14 years				
...	All causes	1,579	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	411	26.0	2.7
2	Malignant neoplasms (C00-C97)	313	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	7.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	5.7	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	4.4	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.5	0.4
7	Cerebrovascular diseases (I60-I69)	34	2.2	0.2
7	Influenza and pneumonia (J09-J18)	34	2.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	2.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
12	Septicemia (A40-A41)	15	0.9	*
13	Diabetes mellitus (E10-E14)	7	0.4	*
13	Meningitis (G00,G03)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
...	All other causes (Residual)	351	22.2	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,739	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,549	44.4	15.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	748	13.0	4.6
3	Malignant neoplasms (C00-C97)	513	8.9	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	300	5.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	2.4	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	98	1.7	0.6
8	Cerebrovascular diseases (I60-I69)	56	1.0	0.3
9	Diabetes mellitus (E10-E14)	55	1.0	0.3
10	Septicemia (A40-A41)	43	0.7	0.3
11	Influenza and pneumonia (J09-J18)	42	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	0.2
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.2	*
...	All other causes (Residual)	914	15.9	5.7
White, female, 25-44 years				
...	All causes	29,582	100.0	94.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,429	25.1	23.6
2	Malignant neoplasms (C00-C97)	6,215	21.0	19.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,797	9.5	8.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,451	8.3	7.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	879	3.0	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	767	2.6	2.4
7	Cerebrovascular diseases (I60-I69)	664	2.2	2.1
8	Diabetes mellitus (E10-E14)	653	2.2	2.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	385	1.3	1.2
10	Chronic lower respiratory diseases (J40-J47)	366	1.2	1.2
11	Influenza and pneumonia (J09-J18)	345	1.2	1.1
12	Septicemia (A40-A41)	332	1.1	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	274	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	205	0.7	0.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	156	0.5	0.5
...	All other causes (Residual)	5,664	19.1	18.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	154,970	100.0	453.2
1	Malignant neoplasms (C00-C97)	59,468	38.4	173.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,242	15.0	68.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,068	6.5	29.4
4	Chronic lower respiratory diseases (J40-J47)	8,498	5.5	24.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,302	3.4	15.5
6	Diabetes mellitus (E10-E14)	4,946	3.2	14.5
7	Cerebrovascular diseases (I60-I69)	4,886	3.2	14.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,632	2.3	10.6
9	Septicemia (A40-A41)	2,439	1.6	7.1
10	Influenza and pneumonia (J09-J18)	1,875	1.2	5.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,816	1.2	5.3
12	Viral hepatitis (B15-B19)	1,352	0.9	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	950	0.6	2.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	787	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	612	0.4	1.8
...	All other causes (Residual)	25,097	16.2	73.4
White, female, 65 years and over				
...	All causes	889,940	100.0	4,264.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	219,959	24.7	1,054.0
2	Malignant neoplasms (C00-C97)	168,237	18.9	806.2
3	Chronic lower respiratory diseases (J40-J47)	61,404	6.9	294.2
4	Cerebrovascular diseases (I60-I69)	58,848	6.6	282.0
5	Alzheimer's disease (G30)	52,503	5.9	251.6
6	Diabetes mellitus (E10-E14)	21,310	2.4	102.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,935	2.4	100.3
8	Influenza and pneumonia (J09-J18)	20,652	2.3	99.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,732	1.8	75.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,778	1.4	61.2
11	Septicemia (A40-A41)	12,400	1.4	59.4
12	Parkinson's disease (G20-G21)	8,867	1.0	42.5
13	Pneumonitis due to solids and liquids (J69)	6,646	0.7	31.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,308	0.6	25.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,585	0.5	22.0
...	All other causes (Residual)	199,776	22.4	957.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	295,222	100.0	683.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,123	23.8	162.3
2	Malignant neoplasms (C00-C97)	67,379	22.8	156.0
3	Cerebrovascular diseases (I60-I69)	15,894	5.4	36.8
4	Diabetes mellitus (E10-E14)	12,983	4.4	30.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,709	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,375	3.2	21.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,241	2.8	19.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,207	2.8	19.0
9	Septicemia (A40-A41)	5,983	2.0	13.8
10	Alzheimer's disease (G30)	5,451	1.8	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,200	1.8	12.0
12	Influenza and pneumonia (J09-J18)	5,000	1.7	11.6
13	Certain conditions originating in the perinatal period (P00-P96)	3,958	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,921	1.3	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,791	0.9	6.5
...	All other causes (Residual)	58,007	19.6	134.3
Black, both sexes, 1-4 years				
...	All causes	1,021	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	27.4	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.0	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.6	3.9
4	Malignant neoplasms (C00-C97)	65	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.3	1.6
6	Chronic lower respiratory diseases (J40-J47)	30	2.9	1.1
7	Influenza and pneumonia (J09-J18)	21	2.1	0.8
8	Septicemia (A40-A41)	18	1.8	*
8	Cerebrovascular diseases (I60-I69)	18	1.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	261	25.6	9.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,196	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	30.0	5.3
2	Malignant neoplasms (C00-C97)	148	12.4	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	9.2	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	6.4	1.1
5	Chronic lower respiratory diseases (J40-J47)	75	6.3	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.7	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.6	0.6
8	Diabetes mellitus (E10-E14)	17	1.4	*
9	Influenza and pneumonia (J09-J18)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.3	*
11	Anemias (D50-D64)	13	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
15	Septicemia (A40-A41)	7	0.6	*
...	All other causes (Residual)	244	20.4	3.6
Black, both sexes, 15-24 years				
...	All causes	6,705	100.0	91.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,906	43.3	39.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,396	20.8	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	7.8	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	289	4.3	3.9
5	Malignant neoplasms (C00-C97)	250	3.7	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	1.3	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.3	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	68	1.0	0.9
9	Anemias (D50-D64)	65	1.0	0.9
10	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.8
11	Diabetes mellitus (E10-E14)	58	0.9	0.8
12	Legal intervention (Y35,Y89.0)	45	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	43	0.6	0.6
14	Septicemia (A40-A41)	33	0.5	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.4
...	All other causes (Residual)	762	11.4	10.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	22,798	100.0	195.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,723	16.3	31.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,696	16.2	31.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,419	15.0	29.3
4	Malignant neoplasms (C00-C97)	2,601	11.4	22.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,183	5.2	10.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,040	4.6	8.9
7	Diabetes mellitus (E10-E14)	712	3.1	6.1
8	Cerebrovascular diseases (I60-I69)	594	2.6	5.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	1.5	3.0
10	Chronic lower respiratory diseases (J40-J47)	309	1.4	2.6
11	Septicemia (A40-A41)	306	1.3	2.6
12	Anemias (D50-D64)	259	1.1	2.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	235	1.0	2.0
14	Influenza and pneumonia (J09-J18)	233	1.0	2.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	230	1.0	2.0
...	All other causes (Residual)	3,910	17.2	33.5
Black, both sexes, 45-64 years				
...	All causes	88,613	100.0	868.4
1	Malignant neoplasms (C00-C97)	25,451	28.7	249.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,687	24.5	212.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,476	5.1	43.9
4	Cerebrovascular diseases (I60-I69)	4,385	4.9	43.0
5	Diabetes mellitus (E10-E14)	4,102	4.6	40.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,273	2.6	22.3
7	Chronic lower respiratory diseases (J40-J47)	2,258	2.5	22.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,244	2.5	22.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,830	2.1	17.9
10	Septicemia (A40-A41)	1,732	2.0	17.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,447	1.6	14.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,102	1.2	10.8
13	Influenza and pneumonia (J09-J18)	1,101	1.2	10.8
14	Viral hepatitis (B15-B19)	876	1.0	8.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	583	0.7	5.7
...	All other causes (Residual)	13,066	14.7	128.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	167,751	100.0	4,363.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,247	26.4	1,151.0
2	Malignant neoplasms (C00-C97)	38,851	23.2	1,010.6
3	Cerebrovascular diseases (I60-I69)	10,802	6.4	281.0
4	Diabetes mellitus (E10-E14)	8,093	4.8	210.5
5	Chronic lower respiratory diseases (J40-J47)	6,634	4.0	172.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,558	3.3	144.6
7	Alzheimer's disease (G30)	5,380	3.2	140.0
8	Septicemia (A40-A41)	3,813	2.3	99.2
9	Influenza and pneumonia (J09-J18)	3,541	2.1	92.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,537	2.1	92.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,378	1.4	61.9
12	Pneumonitis due to solids and liquids (J69)	1,165	0.7	30.3
13	Parkinson's disease (G20-G21)	875	0.5	22.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	813	0.5	21.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	722	0.4	18.8
...	All other causes (Residual)	31,342	18.7	815.3
Black, male, all ages				
...	All causes	150,586	100.0	728.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,140	24.0	174.7
2	Malignant neoplasms (C00-C97)	34,427	22.9	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,583	5.7	41.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,129	4.7	34.5
5	Cerebrovascular diseases (I60-I69)	7,061	4.7	34.1
6	Diabetes mellitus (E10-E14)	6,145	4.1	29.7
7	Chronic lower respiratory diseases (J40-J47)	4,885	3.2	23.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,818	2.5	18.5
9	Septicemia (A40-A41)	2,709	1.8	13.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,546	1.7	12.3
11	Influenza and pneumonia (J09-J18)	2,404	1.6	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,310	1.5	11.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,235	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,908	1.3	9.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,807	1.2	8.7
...	All other causes (Residual)	26,479	17.6	128.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	578	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	29.2	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	8.8	3.7
4	Malignant neoplasms (C00-C97)	39	6.7	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	24	4.2	1.7
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.9	*
9	Septicemia (A40-A41)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	9	1.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	130	22.5	9.3
Black, male, 5-14 years				
...	All causes	731	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	32.6	6.9
2	Malignant neoplasms (C00-C97)	83	11.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	10.0	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.3	1.3
5	Chronic lower respiratory diseases (J40-J47)	40	5.5	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	4.5	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.3	0.7
8	Anemias (D50-D64)	8	1.1	*
8	Diabetes mellitus (E10-E14)	8	1.1	*
8	Influenza and pneumonia (J09-J18)	8	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.1	*
12	Cerebrovascular diseases (I60-I69)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Septicemia (A40-A41)	4	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	143	19.6	4.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,127	100.0	138.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,617	51.0	70.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,020	19.9	27.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	423	8.3	11.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	3.8	5.2
5	Malignant neoplasms (C00-C97)	148	2.9	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.1	1.5
8	Chronic lower respiratory diseases (J40-J47)	46	0.9	1.2
9	Legal intervention (Y35,Y89.0)	43	0.8	1.2
10	Anemias (D50-D64)	34	0.7	0.9
11	Diabetes mellitus (E10-E14)	27	0.5	0.7
12	Cerebrovascular diseases (I60-I69)	24	0.5	0.6
13	Influenza and pneumonia (J09-J18)	20	0.4	0.5
14	Septicemia (A40-A41)	13	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.1	*
...	All other causes (Residual)	396	7.7	10.7
Black, male, 25-44 years				
...	All causes	14,286	100.0	256.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,292	23.0	59.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,461	17.2	44.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,363	16.5	42.5
4	Malignant neoplasms (C00-C97)	965	6.8	17.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	850	5.9	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	662	4.6	11.9
7	Diabetes mellitus (E10-E14)	403	2.8	7.2
8	Cerebrovascular diseases (I60-I69)	330	2.3	5.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	1.3	3.3
10	Chronic lower respiratory diseases (J40-J47)	170	1.2	3.1
11	Septicemia (A40-A41)	147	1.0	2.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	133	0.9	2.4
13	Anemias (D50-D64)	124	0.9	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	0.8	2.1
14	Influenza and pneumonia (J09-J18)	115	0.8	2.1
...	All other causes (Residual)	1,974	13.8	35.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	51,514	100.0	1,089.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,750	26.7	290.8
2	Malignant neoplasms (C00-C97)	13,403	26.0	283.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,111	6.0	65.8
4	Cerebrovascular diseases (I60-I69)	2,440	4.7	51.6
5	Diabetes mellitus (E10-E14)	2,306	4.5	48.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,531	3.0	32.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,225	2.4	25.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,222	2.4	25.8
9	Chronic lower respiratory diseases (J40-J47)	1,168	2.3	24.7
10	Septicemia (A40-A41)	893	1.7	18.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	891	1.7	18.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	844	1.6	17.8
13	Influenza and pneumonia (J09-J18)	638	1.2	13.5
14	Viral hepatitis (B15-B19)	592	1.1	12.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	453	0.9	9.6
...	All other causes (Residual)	7,047	13.7	149.0
Black, male, 65 years and over				
...	All causes	74,347	100.0	4,871.5
1	Malignant neoplasms (C00-C97)	19,781	26.6	1,296.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,718	26.5	1,292.0
3	Cerebrovascular diseases (I60-I69)	4,231	5.7	277.2
4	Chronic lower respiratory diseases (J40-J47)	3,433	4.6	224.9
5	Diabetes mellitus (E10-E14)	3,400	4.6	222.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,387	3.2	156.4
7	Septicemia (A40-A41)	1,593	2.1	104.4
8	Influenza and pneumonia (J09-J18)	1,580	2.1	103.5
9	Alzheimer's disease (G30)	1,520	2.0	99.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,348	1.8	88.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,346	1.8	88.2
12	Pneumonitis due to solids and liquids (J69)	598	0.8	39.2
13	Parkinson's disease (G20-G21)	522	0.7	34.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	449	0.6	29.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	364	0.5	23.9
...	All other causes (Residual)	12,077	16.2	791.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	144,636	100.0	642.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,983	23.5	150.9
2	Malignant neoplasms (C00-C97)	32,952	22.8	146.3
3	Cerebrovascular diseases (I60-I69)	8,833	6.1	39.2
4	Diabetes mellitus (E10-E14)	6,838	4.7	30.4
5	Chronic lower respiratory diseases (J40-J47)	4,490	3.1	19.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,389	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,126	2.9	18.3
8	Alzheimer's disease (G30)	3,899	2.7	17.3
9	Septicemia (A40-A41)	3,274	2.3	14.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,890	2.0	12.8
11	Influenza and pneumonia (J09-J18)	2,596	1.8	11.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,723	1.2	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,375	1.0	6.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,112	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	984	0.7	4.4
...	All other causes (Residual)	31,172	21.6	138.4
Black, female, 1-4 years				
...	All causes	443	100.0	33.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	25.1	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	12.2	4.0
4	Malignant neoplasms (C00-C97)	26	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Septicemia (A40-A41)	9	2.0	*
6	Cerebrovascular diseases (I60-I69)	9	2.0	*
6	Influenza and pneumonia (J09-J18)	9	2.0	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	122	27.5	9.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	465	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	26.0	3.7
2	Malignant neoplasms (C00-C97)	65	14.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	8.0	1.1
4	Chronic lower respiratory diseases (J40-J47)	35	7.5	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	6.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.3	0.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	2.2	*
8	Diabetes mellitus (E10-E14)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.7	*
9	Influenza and pneumonia (J09-J18)	8	1.7	*
11	Anemias (D50-D64)	5	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Septicemia (A40-A41)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	101	21.7	3.0
Black, female, 15-24 years				
...	All causes	1,578	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	376	23.8	10.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	18.3	8.0
3	Malignant neoplasms (C00-C97)	102	6.5	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.4	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.1	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	68	4.3	1.9
7	Anemias (D50-D64)	31	2.0	0.9
7	Diabetes mellitus (E10-E14)	31	2.0	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.9	0.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.6	0.7
11	Septicemia (A40-A41)	20	1.3	0.6
12	Cerebrovascular diseases (I60-I69)	19	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
15	Influenza and pneumonia (J09-J18)	12	0.8	*
...	All other causes (Residual)	342	21.7	9.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	8,512	100.0	139.3
1	Malignant neoplasms (C00-C97)	1,636	19.2	26.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,333	15.7	21.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	958	11.3	15.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	521	6.1	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	431	5.1	7.1
6	Diabetes mellitus (E10-E14)	309	3.6	5.1
7	Cerebrovascular diseases (I60-I69)	264	3.1	4.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	230	2.7	3.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	2.2	3.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.0	2.7
11	Septicemia (A40-A41)	159	1.9	2.6
12	Chronic lower respiratory diseases (J40-J47)	139	1.6	2.3
13	Anemias (D50-D64)	135	1.6	2.2
14	Influenza and pneumonia (J09-J18)	118	1.4	1.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.2	1.7
...	All other causes (Residual)	1,821	21.4	29.8
Black, female, 45-64 years				
...	All causes	37,099	100.0	677.5
1	Malignant neoplasms (C00-C97)	12,048	32.5	220.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,937	21.4	145.0
3	Cerebrovascular diseases (I60-I69)	1,945	5.2	35.5
4	Diabetes mellitus (E10-E14)	1,796	4.8	32.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,365	3.7	24.9
6	Chronic lower respiratory diseases (J40-J47)	1,090	2.9	19.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,019	2.7	18.6
8	Septicemia (A40-A41)	839	2.3	15.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	742	2.0	13.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	608	1.6	11.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	603	1.6	11.0
12	Influenza and pneumonia (J09-J18)	463	1.2	8.5
13	Viral hepatitis (B15-B19)	284	0.8	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	0.6	3.9
15	Anemias (D50-D64)	156	0.4	2.8
...	All other causes (Residual)	5,993	16.2	109.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	93,404	100.0	4,029.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,529	26.3	1,058.2
2	Malignant neoplasms (C00-C97)	19,070	20.4	822.7
3	Cerebrovascular diseases (I60-I69)	6,571	7.0	283.5
4	Diabetes mellitus (E10-E14)	4,693	5.0	202.5
5	Alzheimer's disease (G30)	3,860	4.1	166.5
6	Chronic lower respiratory diseases (J40-J47)	3,201	3.4	138.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,171	3.4	136.8
8	Septicemia (A40-A41)	2,220	2.4	95.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,189	2.3	94.4
10	Influenza and pneumonia (J09-J18)	1,961	2.1	84.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	1.1	44.5
12	Pneumonitis due to solids and liquids (J69)	567	0.6	24.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	449	0.5	19.4
14	Parkinson's disease (G20-G21)	353	0.4	15.2
15	Anemias (D50-D64)	287	0.3	12.4
...	All other causes (Residual)	19,251	20.6	830.5
American Indian, both sexes, all ages				
...	All causes	16,527	100.0	375.7
1	Malignant neoplasms (C00-C97)	3,019	18.3	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,950	17.8	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,922	11.6	43.7
4	Diabetes mellitus (E10-E14)	985	6.0	22.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	918	5.6	20.9
6	Chronic lower respiratory diseases (J40-J47)	708	4.3	16.1
7	Cerebrovascular diseases (I60-I69)	580	3.5	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	485	2.9	11.0
9	Influenza and pneumonia (J09-J18)	306	1.9	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	1.8	6.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.6	6.0
12	Septicemia (A40-A41)	255	1.5	5.8
13	Alzheimer's disease (G30)	217	1.3	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	0.9	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	129	0.8	2.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.8	2.9
...	All other causes (Residual)	3,216	19.5	73.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	89	100.0	28.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	43.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	12.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.7	*
4	Malignant neoplasms (C00-C97)	5	5.6	*
5	Septicemia (A40-A41)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
6	Cerebrovascular diseases (I60-I69)	1	1.1	*
6	Influenza and pneumonia (J09-J18)	1	1.1	*
6	Hernia (K40-K46)	1	1.1	*
...	All other causes (Residual)	21	23.6	6.7
American Indian, both sexes, 5-14 years				
...	All causes	101	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	50.5	6.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	9.9	*
3	Malignant neoplasms (C00-C97)	6	5.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.0	*
7	Influenza and pneumonia (J09-J18)	2	2.0	*
9	Anemias (D50-D64)	1	1.0	*
9	Cerebrovascular diseases (I60-I69)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	15	14.9	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	604	100.0	78.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	44.5	35.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	25.2	19.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	10.6	8.3
4	Malignant neoplasms (C00-C97)	15	2.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.3	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
8	Diabetes mellitus (E10-E14)	4	0.7	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
10	Septicemia (A40-A41)	3	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
12	Pneumonitis due to solids and liquids (J69)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.3	*
...	All other causes (Residual)	57	9.4	7.4
American Indian, both sexes, 25-44 years				
...	All causes	2,132	100.0	169.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	684	32.1	54.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	235	11.0	18.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	9.3	15.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	9.2	15.7
5	Malignant neoplasms (C00-C97)	122	5.7	9.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	5.7	9.6
7	Diabetes mellitus (E10-E14)	53	2.5	4.2
8	Septicemia (A40-A41)	27	1.3	2.2
9	Cerebrovascular diseases (I60-I69)	22	1.0	1.8
10	Influenza and pneumonia (J09-J18)	21	1.0	1.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.8	*
14	Viral hepatitis (B15-B19)	14	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.5	*
...	All other causes (Residual)	376	17.6	29.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	5,186	100.0	555.9
1	Malignant neoplasms (C00-C97)	1,106	21.3	118.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	882	17.0	94.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	611	11.8	65.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	516	9.9	55.3
5	Diabetes mellitus (E10-E14)	363	7.0	38.9
6	Chronic lower respiratory diseases (J40-J47)	181	3.5	19.4
7	Cerebrovascular diseases (I60-I69)	139	2.7	14.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	2.0	11.4
9	Viral hepatitis (B15-B19)	82	1.6	8.8
10	Influenza and pneumonia (J09-J18)	81	1.6	8.7
11	Septicemia (A40-A41)	80	1.5	8.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.4	7.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	1.0	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.9	5.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.4	2.5
...	All other causes (Residual)	844	16.3	90.5
American Indian, both sexes, 65 years and over				
...	All causes	8,045	100.0	2,749.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,845	22.9	630.4
2	Malignant neoplasms (C00-C97)	1,763	21.9	602.4
3	Diabetes mellitus (E10-E14)	565	7.0	193.1
4	Chronic lower respiratory diseases (J40-J47)	516	6.4	176.3
5	Cerebrovascular diseases (I60-I69)	415	5.2	141.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	3.0	82.0
7	Alzheimer's disease (G30)	212	2.6	72.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	2.6	70.4
9	Influenza and pneumonia (J09-J18)	196	2.4	67.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	161	2.0	55.0
11	Septicemia (A40-A41)	139	1.7	47.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.2	33.1
13	Pneumonitis due to solids and liquids (J69)	65	0.8	22.2
14	Parkinson's disease (G20-G21)	63	0.8	21.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	12.6
...	All other causes (Residual)	1,525	19.0	521.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	9,067	100.0	410.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,731	19.1	78.3
2	Malignant neoplasms (C00-C97)	1,559	17.2	70.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	14.1	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	517	5.7	23.4
5	Diabetes mellitus (E10-E14)	514	5.7	23.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	4.3	17.5
7	Chronic lower respiratory diseases (J40-J47)	370	4.1	16.7
8	Cerebrovascular diseases (I60-I69)	266	2.9	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	2.3	9.4
10	Influenza and pneumonia (J09-J18)	144	1.6	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.6	6.5
12	Septicemia (A40-A41)	114	1.3	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	0.8	3.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.8	3.3
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.8	3.3
...	All other causes (Residual)	1,619	17.9	73.2
American Indian, male, 1-4 years				
...	All causes	54	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	51.9	17.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	11.1	*
3	Malignant neoplasms (C00-C97)	4	7.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.6	*
5	Cerebrovascular diseases (I60-I69)	1	1.9	*
...	All other causes (Residual)	12	22.2	*
American Indian, male, 5-14 years				
...	All causes	65	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	56.9	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	12.3	*
3	Malignant neoplasms (C00-C97)	4	6.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.5	*
6	Influenza and pneumonia (J09-J18)	1	1.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.5	*
...	All other causes (Residual)	7	10.8	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	427	100.0	107.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	43.8	47.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	27.2	29.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	11.5	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.3	*
5	Malignant neoplasms (C00-C97)	9	2.1	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
7	Legal intervention (Y35,Y89.0)	3	0.7	*
10	Diabetes mellitus (E10-E14)	2	0.5	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Septicemia (A40-A41)	1	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Cerebrovascular diseases (I60-I69)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	31	7.3	7.8
American Indian, male, 25-44 years				
...	All causes	1,348	100.0	208.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	460	34.1	71.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	11.6	24.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	138	10.2	21.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	9.9	20.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	7.3	15.3
6	Malignant neoplasms (C00-C97)	60	4.5	9.3
7	Diabetes mellitus (E10-E14)	34	2.5	5.3
8	Influenza and pneumonia (J09-J18)	13	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.9	*
10	Septicemia (A40-A41)	11	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
12	Cerebrovascular diseases (I60-I69)	8	0.6	*
13	Legal intervention (Y35,Y89.0)	7	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
...	All other causes (Residual)	196	14.5	30.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-64 years				
...	All causes	3,098	100.0	679.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	605	19.5	132.7
2	Malignant neoplasms (C00-C97)	580	18.7	127.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	13.4	90.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	296	9.6	64.9
5	Diabetes mellitus (E10-E14)	210	6.8	46.1
6	Chronic lower respiratory diseases (J40-J47)	95	3.1	20.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	2.8	19.1
8	Cerebrovascular diseases (I60-I69)	79	2.6	17.3
9	Viral hepatitis (B15-B19)	49	1.6	10.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.5	9.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	1.4	9.7
12	Influenza and pneumonia (J09-J18)	43	1.4	9.4
13	Septicemia (A40-A41)	40	1.3	8.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.0	6.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.6	*
...	All other causes (Residual)	463	14.9	101.6
American Indian, male, 65 years and over				
...	All causes	3,867	100.0	2,960.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	976	25.2	747.3
2	Malignant neoplasms (C00-C97)	901	23.3	689.8
3	Chronic lower respiratory diseases (J40-J47)	270	7.0	206.7
4	Diabetes mellitus (E10-E14)	268	6.9	205.2
5	Cerebrovascular diseases (I60-I69)	176	4.6	134.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	3.5	104.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	2.2	66.6
8	Influenza and pneumonia (J09-J18)	83	2.1	63.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	78	2.0	59.7
10	Septicemia (A40-A41)	62	1.6	47.5
11	Alzheimer's disease (G30)	53	1.4	40.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	1.0	29.9
13	Parkinson's disease (G20-G21)	34	0.9	26.0
14	Pneumonitis due to solids and liquids (J69)	32	0.8	24.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	0.5	*
...	All other causes (Residual)	653	16.9	500.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	7,460	100.0	340.9
1	Malignant neoplasms (C00-C97)	1,460	19.6	66.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,219	16.3	55.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	8.6	29.4
4	Diabetes mellitus (E10-E14)	471	6.3	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	401	5.4	18.3
6	Chronic lower respiratory diseases (J40-J47)	338	4.5	15.4
7	Cerebrovascular diseases (I60-I69)	314	4.2	14.4
8	Alzheimer's disease (G30)	162	2.2	7.4
8	Influenza and pneumonia (J09-J18)	162	2.2	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.0	6.9
11	Septicemia (A40-A41)	141	1.9	6.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	1.3	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.0	3.5
14	Certain conditions originating in the perinatal period (P00-P96)	57	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.8	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.8	2.6
...	All other causes (Residual)	1,652	22.1	75.5
American Indian, female, 1-4 years				
...	All causes	35	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	31.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	14.3	*
3	Septicemia (A40-A41)	3	8.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.6	*
5	Malignant neoplasms (C00-C97)	1	2.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.9	*
5	Influenza and pneumonia (J09-J18)	1	2.9	*
5	Hernia (K40-K46)	1	2.9	*
...	All other causes (Residual)	9	25.7	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 5-14 years				
...	All causes	36	100.0	9.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	38.9	*
2	Malignant neoplasms (C00-C97)	2	5.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	5.6	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.8	*
6	Anemias (D50-D64)	1	2.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.8	*
6	Cerebrovascular diseases (I60-I69)	1	2.8	*
6	Influenza and pneumonia (J09-J18)	1	2.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.8	*
...	All other causes (Residual)	8	22.2	*
American Indian, female, 15-24 years				
...	All causes	177	100.0	47.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	46.3	22.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	20.3	9.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	8.5	*
4	Malignant neoplasms (C00-C97)	6	3.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.3	*
6	Septicemia (A40-A41)	2	1.1	*
6	Diabetes mellitus (E10-E14)	2	1.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
10	Pneumonitis due to solids and liquids (J69)	1	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
10	Legal intervention (Y35,Y89.0)	1	0.6	*
...	All other causes (Residual)	22	12.4	5.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-44 years				
...	All causes	784	100.0	128.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	28.6	36.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	97	12.4	15.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	8.3	10.7
4	Malignant neoplasms (C00-C97)	62	7.9	10.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.1	6.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	2.8	3.6
7	Diabetes mellitus (E10-E14)	19	2.4	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	17	2.2	*
9	Septicemia (A40-A41)	16	2.0	*
10	Cerebrovascular diseases (I60-I69)	14	1.8	*
11	Viral hepatitis (B15-B19)	10	1.3	*
12	Influenza and pneumonia (J09-J18)	8	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (Residual)	167	21.3	27.4
American Indian, female, 45-64 years				
...	All causes	2,088	100.0	437.7
1	Malignant neoplasms (C00-C97)	526	25.2	110.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	13.3	58.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	220	10.5	46.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	9.4	41.3
5	Diabetes mellitus (E10-E14)	153	7.3	32.1
6	Chronic lower respiratory diseases (J40-J47)	86	4.1	18.0
7	Cerebrovascular diseases (I60-I69)	60	2.9	12.6
8	Septicemia (A40-A41)	40	1.9	8.4
9	Influenza and pneumonia (J09-J18)	38	1.8	8.0
10	Viral hepatitis (B15-B19)	33	1.6	6.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.2	5.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.4	*
15	Pneumonitis due to solids and liquids (J69)	8	0.4	*
...	All other causes (Residual)	378	18.1	79.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65 years and over				
...	All causes	4,178	100.0	2,578.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	869	20.8	536.3
2	Malignant neoplasms (C00-C97)	862	20.6	532.0
3	Diabetes mellitus (E10-E14)	297	7.1	183.3
4	Chronic lower respiratory diseases (J40-J47)	246	5.9	151.8
5	Cerebrovascular diseases (I60-I69)	239	5.7	147.5
6	Alzheimer's disease (G30)	159	3.8	98.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	2.8	73.4
8	Influenza and pneumonia (J09-J18)	113	2.7	69.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	2.5	63.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	83	2.0	51.2
11	Septicemia (A40-A41)	77	1.8	47.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.4	35.8
13	Pneumonitis due to solids and liquids (J69)	33	0.8	20.4
14	Parkinson's disease (G20-G21)	29	0.7	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	13.0
...	All other causes (Residual)	870	20.8	536.9
Asian or Pacific Islander, both sexes, all ages				
...	All causes	56,352	100.0	311.5
1	Malignant neoplasms (C00-C97)	15,340	27.2	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,266	21.8	67.8
3	Cerebrovascular diseases (I60-I69)	4,108	7.3	22.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	4.2	13.1
5	Diabetes mellitus (E10-E14)	2,158	3.8	11.9
6	Influenza and pneumonia (J09-J18)	1,745	3.1	9.6
7	Chronic lower respiratory diseases (J40-J47)	1,624	2.9	9.0
8	Alzheimer's disease (G30)	1,379	2.4	7.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,152	2.0	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,054	1.9	5.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	857	1.5	4.7
12	Septicemia (A40-A41)	659	1.2	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	523	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	476	0.8	2.6
15	Parkinson's disease (G20-G21)	472	0.8	2.6
...	All other causes (Residual)	10,167	18.0	56.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	150	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	20.0	3.1
2	Malignant neoplasms (C00-C97)	21	14.0	2.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.0	2.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
7	Septicemia (A40-A41)	4	2.7	*
7	Influenza and pneumonia (J09-J18)	4	2.7	*
9	Anemias (D50-D64)	2	1.3	*
9	Cerebrovascular diseases (I60-I69)	2	1.3	*
9	Acute bronchitis and bronchiolitis (J20-J21)	2	1.3	*
12	Tuberculosis (A16-A19)	1	0.7	*
12	Meningitis (G00,G03)	1	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (Residual)	40	26.7	4.1
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	192	100.0	8.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	27.6	2.2
2	Malignant neoplasms (C00-C97)	46	24.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	8.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	5.2	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	5.2	*
6	Influenza and pneumonia (J09-J18)	4	2.1	*
7	Anemias (D50-D64)	3	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
9	Septicemia (A40-A41)	2	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
9	Cerebrovascular diseases (I60-I69)	2	1.0	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (Residual)	37	19.3	1.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	772	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	31.3	9.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	25.3	7.6
3	Malignant neoplasms (C00-C97)	91	11.8	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	8.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.2	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
7	Cerebrovascular diseases (I60-I69)	6	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
10	Influenza and pneumonia (J09-J18)	4	0.5	*
10	Legal intervention (Y35,Y89.0)	4	0.5	*
12	Septicemia (A40-A41)	3	0.4	*
12	Diabetes mellitus (E10-E14)	3	0.4	*
14	Tuberculosis (A16-A19)	2	0.3	*
14	Anemias (D50-D64)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	88	11.4	3.4
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,225	100.0	54.2
1	Malignant neoplasms (C00-C97)	821	25.5	13.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	526	16.3	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	456	14.1	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	409	12.7	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	4.4	2.4
6	Cerebrovascular diseases (I60-I69)	135	4.2	2.3
7	Diabetes mellitus (E10-E14)	56	1.7	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	47	1.5	0.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	31	1.0	0.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.9	0.5
12	Influenza and pneumonia (J09-J18)	24	0.7	0.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.7	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.7	0.4
15	Septicemia (A40-A41)	20	0.6	0.3
15	Aortic aneurysm and dissection (I71)	20	0.6	0.3
...	All other causes (Residual)	436	13.5	7.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	11,474	100.0	268.4
1	Malignant neoplasms (C00-C97)	4,768	41.6	111.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,140	18.7	50.1
3	Cerebrovascular diseases (I60-I69)	725	6.3	17.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	522	4.5	12.2
5	Diabetes mellitus (E10-E14)	463	4.0	10.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	3.0	8.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	213	1.9	5.0
8	Chronic lower respiratory diseases (J40-J47)	144	1.3	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.2	3.3
10	Viral hepatitis (B15-B19)	134	1.2	3.1
11	Influenza and pneumonia (J09-J18)	131	1.1	3.1
12	Septicemia (A40-A41)	117	1.0	2.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	0.9	2.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	0.8	2.2
15	Aortic aneurysm and dissection (I71)	79	0.7	1.8
...	All other causes (Residual)	1,360	11.9	31.8
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	39,626	100.0	2,289.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,644	24.3	557.3
2	Malignant neoplasms (C00-C97)	9,589	24.2	554.1
3	Cerebrovascular diseases (I60-I69)	3,233	8.2	186.8
4	Diabetes mellitus (E10-E14)	1,636	4.1	94.5
5	Influenza and pneumonia (J09-J18)	1,570	4.0	90.7
6	Chronic lower respiratory diseases (J40-J47)	1,451	3.7	83.8
7	Alzheimer's disease (G30)	1,365	3.4	78.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	964	2.4	55.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	885	2.2	51.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	745	1.9	43.1
11	Septicemia (A40-A41)	505	1.3	29.2
12	Parkinson's disease (G20-G21)	462	1.2	26.7
13	Pneumonitis due to solids and liquids (J69)	347	0.9	20.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	263	0.7	15.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	262	0.7	15.1
...	All other causes (Residual)	6,705	16.9	387.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	28,819	100.0	332.8
1	Malignant neoplasms (C00-C97)	7,709	26.7	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,650	23.1	76.8
3	Cerebrovascular diseases (I60-I69)	1,876	6.5	21.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,472	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,093	3.8	12.6
6	Chronic lower respiratory diseases (J40-J47)	966	3.4	11.2
7	Influenza and pneumonia (J09-J18)	890	3.1	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	805	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	1.7	5.8
10	Alzheimer's disease (G30)	439	1.5	5.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.3	4.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.1	3.8
13	Septicemia (A40-A41)	312	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	262	0.9	3.0
15	Parkinson's disease (G20-G21)	252	0.9	2.9
...	All other causes (Residual)	4,889	17.0	56.5
Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	22.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	16.3	*
3	Malignant neoplasms (C00-C97)	10	12.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
7	Tuberculosis (A16-A19)	1	1.3	*
7	Septicemia (A40-A41)	1	1.3	*
7	Anemias (D50-D64)	1	1.3	*
7	Cerebrovascular diseases (I60-I69)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.3	*
...	All other causes (Residual)	23	28.8	4.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	108	100.0	9.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	28.7	2.6
2	Malignant neoplasms (C00-C97)	27	25.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	7.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.5	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	6.5	*
6	Anemias (D50-D64)	3	2.8	*
7	Influenza and pneumonia (J09-J18)	2	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
9	Cerebrovascular diseases (I60-I69)	1	0.9	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	0.9	*
...	All other causes (Residual)	18	16.7	*
Asian or Pacific Islander, male, 15-24 years				
...	All causes	540	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	35.2	14.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	23.3	9.7
3	Malignant neoplasms (C00-C97)	62	11.5	4.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	9.3	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	5.7	2.4
6	Cerebrovascular diseases (I60-I69)	6	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.1	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
11	Septicemia (A40-A41)	2	0.4	*
12	Anemias (D50-D64)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Influenza and pneumonia (J09-J18)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	52	9.6	4.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-44 years				
...	All causes	2,062	100.0	72.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	19.4	14.1
2	Malignant neoplasms (C00-C97)	374	18.1	13.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	333	16.1	11.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	329	16.0	11.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	5.1	3.7
6	Cerebrovascular diseases (I60-I69)	71	3.4	2.5
7	Diabetes mellitus (E10-E14)	39	1.9	1.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	39	1.9	1.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.1	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.9	*
11	Influenza and pneumonia (J09-J18)	17	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.8	*
13	Viral hepatitis (B15-B19)	16	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	*
14	Aortic aneurysm and dissection (I71)	14	0.7	*
...	All other causes (Residual)	252	12.2	8.9
Asian or Pacific Islander, male, 45-64 years				
...	All causes	6,727	100.0	340.6
1	Malignant neoplasms (C00-C97)	2,325	34.6	117.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,587	23.6	80.4
3	Cerebrovascular diseases (I60-I69)	403	6.0	20.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	4.9	16.8
5	Diabetes mellitus (E10-E14)	296	4.4	15.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	3.6	12.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	154	2.3	7.8
8	Chronic lower respiratory diseases (J40-J47)	87	1.3	4.4
9	Viral hepatitis (B15-B19)	85	1.3	4.3
10	Influenza and pneumonia (J09-J18)	74	1.1	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.1	3.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.0	3.5
13	Septicemia (A40-A41)	66	1.0	3.3
14	Aortic aneurysm and dissection (I71)	57	0.8	2.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	0.8	2.9
...	All other causes (Residual)	821	12.2	41.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65 years and over				
...	All causes	18,786	100.0	2,507.2
1	Malignant neoplasms (C00-C97)	4,909	26.1	655.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,680	24.9	624.6
3	Cerebrovascular diseases (I60-I69)	1,392	7.4	185.8
4	Chronic lower respiratory diseases (J40-J47)	865	4.6	115.4
5	Influenza and pneumonia (J09-J18)	788	4.2	105.2
6	Diabetes mellitus (E10-E14)	757	4.0	101.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	480	2.6	64.1
8	Alzheimer's disease (G30)	432	2.3	57.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	409	2.2	54.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	301	1.6	40.2
11	Parkinson's disease (G20-G21)	247	1.3	33.0
12	Septicemia (A40-A41)	228	1.2	30.4
13	Pneumonitis due to solids and liquids (J69)	191	1.0	25.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	133	0.7	17.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.7	16.8
...	All other causes (Residual)	2,848	15.2	380.1
Asian or Pacific Islander, female, all ages				
...	All causes	27,533	100.0	292.0
1	Malignant neoplasms (C00-C97)	7,631	27.7	80.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,616	20.4	59.6
3	Cerebrovascular diseases (I60-I69)	2,232	8.1	23.7
4	Diabetes mellitus (E10-E14)	1,065	3.9	11.3
5	Alzheimer's disease (G30)	940	3.4	10.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	3.3	9.5
7	Influenza and pneumonia (J09-J18)	855	3.1	9.1
8	Chronic lower respiratory diseases (J40-J47)	658	2.4	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	553	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	480	1.7	5.1
11	Septicemia (A40-A41)	347	1.3	3.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	1.3	3.7
13	Parkinson's disease (G20-G21)	220	0.8	2.3
14	Certain conditions originating in the perinatal period (P00-P96)	214	0.8	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	197	0.7	2.1
...	All other causes (Residual)	5,278	19.2	56.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	70	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.1	*
2	Malignant neoplasms (C00-C97)	11	15.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	11.4	*
4	Septicemia (A40-A41)	3	4.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.3	*
4	Influenza and pneumonia (J09-J18)	3	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.3	*
9	Anemias (D50-D64)	1	1.4	*
9	Meningitis (G00,G03)	1	1.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.4	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.4	*
...	All other causes (Residual)	17	24.3	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	84	100.0	7.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	26.2	1.9
2	Malignant neoplasms (C00-C97)	19	22.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.6	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	3.6	*
6	Septicemia (A40-A41)	2	2.4	*
6	Influenza and pneumonia (J09-J18)	2	2.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.2	*
8	Cerebrovascular diseases (I60-I69)	1	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.2	*
...	All other causes (Residual)	19	22.6	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	232	100.0	18.4
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	29.7	5.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	22.4	4.1
3	Malignant neoplasms (C00-C97)	29	12.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
7	Influenza and pneumonia (J09-J18)	3	1.3	*
9	Tuberculosis (A16-A19)	2	0.9	*
9	Diabetes mellitus (E10-E14)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
12	Septicemia (A40-A41)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Aortic aneurysm and dissection (I71)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	34	14.7	2.7
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,163	100.0	37.2
1	Malignant neoplasms (C00-C97)	447	38.4	14.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	10.8	4.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	10.6	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	6.9	2.6
5	Cerebrovascular diseases (I60-I69)	64	5.5	2.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	3.3	1.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.7	1.0
8	Diabetes mellitus (E10-E14)	17	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.9	*
11	Septicemia (A40-A41)	8	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.7	*
14	Influenza and pneumonia (J09-J18)	7	0.6	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	179	15.4	5.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-64 years				
...	All causes	4,747	100.0	206.5
1	Malignant neoplasms (C00-C97)	2,443	51.5	106.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	11.6	24.1
3	Cerebrovascular diseases (I60-I69)	322	6.8	14.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	4.0	8.3
5	Diabetes mellitus (E10-E14)	167	3.5	7.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	2.1	4.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.4	3.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.2	2.6
9	Influenza and pneumonia (J09-J18)	57	1.2	2.5
9	Chronic lower respiratory diseases (J40-J47)	57	1.2	2.5
11	Septicemia (A40-A41)	51	1.1	2.2
12	Viral hepatitis (B15-B19)	49	1.0	2.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	0.8	1.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.7	1.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	1.1
...	All other causes (Residual)	535	11.3	23.3
Asian or Pacific Islander, female, 65 years and over				
...	All causes	20,840	100.0	2,123.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,964	23.8	505.9
2	Malignant neoplasms (C00-C97)	4,680	22.5	476.9
3	Cerebrovascular diseases (I60-I69)	1,841	8.8	187.6
4	Alzheimer's disease (G30)	933	4.5	95.1
5	Diabetes mellitus (E10-E14)	879	4.2	89.6
6	Influenza and pneumonia (J09-J18)	782	3.8	79.7
7	Chronic lower respiratory diseases (J40-J47)	586	2.8	59.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	484	2.3	49.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	2.3	48.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	444	2.1	45.2
11	Septicemia (A40-A41)	277	1.3	28.2
12	Parkinson's disease (G20-G21)	215	1.0	21.9
13	Pneumonitis due to solids and liquids (J69)	156	0.7	15.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.7	13.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	130	0.6	13.2
...	All other causes (Residual)	3,857	18.5	393.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012