

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,596,993	100.0	821.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	611,105	23.5	193.3
2	Malignant neoplasms (C00-C97)	584,881	22.5	185.0
3	Chronic lower respiratory diseases (J40-J47)	149,205	5.7	47.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130,557	5.0	41.3
5	Cerebrovascular diseases (I60-I69)	128,978	5.0	40.8
6	Alzheimer's disease (G30)	84,767	3.3	26.8
7	Diabetes mellitus (E10-E14)	75,578	2.9	23.9
8	Influenza and pneumonia (J09-J18)	56,979	2.2	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47,112	1.8	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41,149	1.6	13.0
11	Septicemia (A40-A41)	38,156	1.5	12.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	36,427	1.4	11.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30,770	1.2	9.7
14	Parkinson's disease (G20-G21)	25,196	1.0	8.0
15	Pneumonitis due to solids and liquids (J69)	18,579	0.7	5.9
...	All other causes (Residual)	537,554	20.7	170.0
All races, both sexes, 1-4 years				
...	All causes	4,068	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,316	32.4	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	476	11.7	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	8.3	2.1
4	Malignant neoplasms (C00-C97)	328	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	4.2	1.1
6	Influenza and pneumonia (J09-J18)	102	2.5	0.6
7	Chronic lower respiratory diseases (J40-J47)	64	1.6	0.4
8	Septicemia (A40-A41)	53	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	45	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	38	0.9	0.2
12	Meningitis (G00,G03)	23	0.6	0.1
13	Anemias (D50-D64)	20	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
...	All other causes (Residual)	1,021	25.1	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,340	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,521	28.5	3.7
2	Malignant neoplasms (C00-C97)	895	16.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	7.4	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	6.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	3.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	155	2.9	0.4
8	Influenza and pneumonia (J09-J18)	128	2.4	0.3
9	Cerebrovascular diseases (I60-I69)	89	1.7	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	1.2	0.2
11	Septicemia (A40-A41)	55	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	33	0.6	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	26	0.5	0.1
14	Diabetes mellitus (E10-E14)	24	0.4	0.1
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	1,147	21.5	2.8
All races, both sexes, 15-24 years				
...	All causes	28,486	100.0	64.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,619	40.8	26.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,878	17.1	11.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,329	15.2	9.8
4	Malignant neoplasms (C00-C97)	1,496	5.3	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	941	3.3	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	362	1.3	0.8
7	Influenza and pneumonia (J09-J18)	197	0.7	0.4
8	Diabetes mellitus (E10-E14)	193	0.7	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	178	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	155	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	153	0.5	0.3
12	Septicemia (A40-A41)	138	0.5	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	109	0.4	0.2
14	Legal intervention (Y35,Y89.0)	99	0.3	0.2
15	Anemias (D50-D64)	93	0.3	0.2
...	All other causes (Residual)	3,546	12.4	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	115,036	100.0	138.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,563	27.4	37.9
2	Malignant neoplasms (C00-C97)	15,022	13.1	18.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,599	11.8	16.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,899	11.2	15.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,817	5.9	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,167	2.8	3.8
7	Diabetes mellitus (E10-E14)	2,636	2.3	3.2
8	Cerebrovascular diseases (I60-I69)	2,195	1.9	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,877	1.6	2.3
10	Influenza and pneumonia (J09-J18)	1,330	1.2	1.6
11	Septicemia (A40-A41)	1,153	1.0	1.4
12	Chronic lower respiratory diseases (J40-J47)	1,051	0.9	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	893	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	864	0.8	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	742	0.6	0.9
...	All other causes (Residual)	19,228	16.7	23.1
All races, both sexes, 45-64 years				
...	All causes	515,851	100.0	620.9
1	Malignant neoplasms (C00-C97)	159,509	30.9	192.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	107,735	20.9	129.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,414	7.3	45.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	20,736	4.0	25.0
5	Chronic lower respiratory diseases (J40-J47)	20,561	4.0	24.7
6	Diabetes mellitus (E10-E14)	18,960	3.7	22.8
7	Cerebrovascular diseases (I60-I69)	16,789	3.3	20.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15,756	3.1	19.0
9	Septicemia (A40-A41)	7,790	1.5	9.4
10	Influenza and pneumonia (J09-J18)	7,012	1.4	8.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,972	1.4	8.4
12	Viral hepatitis (B15-B19)	5,847	1.1	7.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,650	0.9	5.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4,198	0.8	5.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,164	0.6	3.8
...	All other causes (Residual)	78,758	15.3	94.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,904,640	100.0	4,260.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	488,156	25.6	1,092.0
2	Malignant neoplasms (C00-C97)	407,558	21.4	911.7
3	Chronic lower respiratory diseases (J40-J47)	127,194	6.7	284.5
4	Cerebrovascular diseases (I60-I69)	109,602	5.8	245.2
5	Alzheimer's disease (G30)	83,786	4.4	187.4
6	Diabetes mellitus (E10-E14)	53,751	2.8	120.2
7	Influenza and pneumonia (J09-J18)	48,031	2.5	107.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,942	2.4	102.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39,080	2.1	87.4
10	Septicemia (A40-A41)	28,815	1.5	64.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25,564	1.3	57.2
12	Parkinson's disease (G20-G21)	24,526	1.3	54.9
13	Pneumonitis due to solids and liquids (J69)	16,425	0.9	36.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	12,488	0.7	27.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12,433	0.7	27.8
...	All other causes (Residual)	381,289	20.0	852.9
All races, male, all ages				
...	All causes	1,306,034	100.0	839.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	321,347	24.6	206.5
2	Malignant neoplasms (C00-C97)	307,559	23.5	197.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81,916	6.3	52.6
4	Chronic lower respiratory diseases (J40-J47)	70,317	5.4	45.2
5	Cerebrovascular diseases (I60-I69)	53,691	4.1	34.5
6	Diabetes mellitus (E10-E14)	39,841	3.1	25.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,055	2.5	20.6
8	Influenza and pneumonia (J09-J18)	26,804	2.1	17.2
9	Alzheimer's disease (G30)	25,836	2.0	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23,709	1.8	15.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,493	1.8	15.1
12	Septicemia (A40-A41)	17,994	1.4	11.6
13	Parkinson's disease (G20-G21)	15,088	1.2	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,963	1.0	8.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,726	1.0	8.2
...	All other causes (Residual)	240,695	18.4	154.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,323	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	35.8	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	10.0	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	7.9	2.3
4	Malignant neoplasms (C00-C97)	183	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	4.1	1.2
6	Influenza and pneumonia (J09-J18)	53	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	38	1.6	0.5
8	Septicemia (A40-A41)	32	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.4
10	Cerebrovascular diseases (I60-I69)	22	0.9	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	566	24.4	7.0
All races, male, 5-14 years				
...	All causes	3,077	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	950	30.9	4.5
2	Malignant neoplasms (C00-C97)	471	15.3	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	252	8.2	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	5.5	0.8
6	Chronic lower respiratory diseases (J40-J47)	100	3.2	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.0	0.4
8	Influenza and pneumonia (J09-J18)	70	2.3	0.3
9	Cerebrovascular diseases (I60-I69)	44	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.3	0.2
11	Septicemia (A40-A41)	32	1.0	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	17	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
15	Meningitis (G00,G03)	11	0.4	*
...	All other causes (Residual)	626	20.3	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	20,864	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,545	41.0	37.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,903	18.7	17.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,765	18.0	16.7
4	Malignant neoplasms (C00-C97)	859	4.1	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	628	3.0	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	223	1.1	1.0
7	Diabetes mellitus (E10-E14)	120	0.6	0.5
8	Influenza and pneumonia (J09-J18)	114	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	101	0.5	0.4
10	Legal intervention (Y35,Y89.0)	95	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	85	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	81	0.4	0.4
13	Septicemia (A40-A41)	74	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	0.3
15	Anemias (D50-D64)	54	0.3	0.2
...	All other causes (Residual)	2,160	10.4	9.6
All races, male, 25-44 years				
...	All causes	74,534	100.0	178.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,591	30.3	54.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,066	13.5	24.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,366	12.6	22.4
4	Malignant neoplasms (C00-C97)	6,684	9.0	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,580	7.5	13.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,055	2.8	4.9
7	Diabetes mellitus (E10-E14)	1,614	2.2	3.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,299	1.7	3.1
9	Cerebrovascular diseases (I60-I69)	1,217	1.6	2.9
10	Influenza and pneumonia (J09-J18)	717	1.0	1.7
11	Septicemia (A40-A41)	575	0.8	1.4
12	Chronic lower respiratory diseases (J40-J47)	525	0.7	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	491	0.7	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	482	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	338	0.5	0.8
...	All other causes (Residual)	10,934	14.7	26.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	314,322	100.0	775.6
1	Malignant neoplasms (C00-C97)	85,776	27.3	211.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,510	24.0	186.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,249	8.0	62.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	14,343	4.6	35.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,766	3.7	29.0
6	Diabetes mellitus (E10-E14)	11,566	3.7	28.5
7	Chronic lower respiratory diseases (J40-J47)	10,429	3.3	25.7
8	Cerebrovascular diseases (I60-I69)	9,718	3.1	24.0
9	Septicemia (A40-A41)	4,196	1.3	10.4
10	Viral hepatitis (B15-B19)	4,120	1.3	10.2
11	Influenza and pneumonia (J09-J18)	4,022	1.3	9.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,990	1.3	9.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,129	1.0	7.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,980	0.9	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,302	0.7	5.7
...	All other causes (Residual)	45,226	14.4	111.6
All races, male, 65 years and over				
...	All causes	877,700	100.0	4,478.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235,460	26.8	1,201.3
2	Malignant neoplasms (C00-C97)	213,550	24.3	1,089.5
3	Chronic lower respiratory diseases (J40-J47)	59,109	6.7	301.6
4	Cerebrovascular diseases (I60-I69)	42,544	4.8	217.1
5	Diabetes mellitus (E10-E14)	26,520	3.0	135.3
6	Alzheimer's disease (G30)	25,420	2.9	129.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,069	2.6	117.7
8	Influenza and pneumonia (J09-J18)	21,717	2.5	110.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,925	2.2	96.6
10	Parkinson's disease (G20-G21)	14,640	1.7	74.7
11	Septicemia (A40-A41)	12,997	1.5	66.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,625	1.1	49.1
13	Pneumonitis due to solids and liquids (J69)	8,855	1.0	45.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7,287	0.8	37.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,505	0.7	33.2
...	All other causes (Residual)	151,477	17.3	772.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,290,959	100.0	804.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,758	22.4	180.6
2	Malignant neoplasms (C00-C97)	277,322	21.5	172.8
3	Chronic lower respiratory diseases (J40-J47)	78,888	6.1	49.2
4	Cerebrovascular diseases (I60-I69)	75,287	5.8	46.9
5	Alzheimer's disease (G30)	58,931	4.6	36.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,641	3.8	30.3
7	Diabetes mellitus (E10-E14)	35,737	2.8	22.3
8	Influenza and pneumonia (J09-J18)	30,175	2.3	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,619	1.8	14.7
10	Septicemia (A40-A41)	20,162	1.6	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,807	1.4	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,718	1.0	7.9
13	Parkinson's disease (G20-G21)	10,108	0.8	6.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,094	0.7	5.7
15	Pneumonitis due to solids and liquids (J69)	8,439	0.7	5.3
...	All other causes (Residual)	294,273	22.8	183.4
All races, female, 1-4 years				
...	All causes	1,745	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.8	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	14.0	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	8.8	2.0
4	Malignant neoplasms (C00-C97)	145	8.3	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.2	0.9
6	Influenza and pneumonia (J09-J18)	49	2.8	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	26	1.5	0.3
9	Septicemia (A40-A41)	21	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	16	0.9	*
12	Meningitis (G00,G03)	10	0.6	*
13	Anemias (D50-D64)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	455	26.1	5.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,263	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	571	25.2	2.8
2	Malignant neoplasms (C00-C97)	424	18.7	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	7.2	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	6.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	4.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.5	0.4
7	Influenza and pneumonia (J09-J18)	58	2.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	55	2.4	0.3
9	Cerebrovascular diseases (I60-I69)	45	2.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.1	0.1
11	Septicemia (A40-A41)	23	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
13	Diabetes mellitus (E10-E14)	12	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Anemias (D50-D64)	8	0.4	*
...	All other causes (Residual)	519	22.9	2.6
All races, female, 15-24 years				
...	All causes	7,622	100.0	35.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,074	40.3	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	975	12.8	4.5
3	Malignant neoplasms (C00-C97)	637	8.4	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	564	7.4	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	4.1	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	178	2.3	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	1.8	0.6
8	Influenza and pneumonia (J09-J18)	83	1.1	0.4
9	Diabetes mellitus (E10-E14)	73	1.0	0.3
10	Cerebrovascular diseases (I60-I69)	68	0.9	0.3
11	Septicemia (A40-A41)	64	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.7	0.3
13	Anemias (D50-D64)	39	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.4	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.3	0.1
...	All other causes (Residual)	1,307	17.1	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	40,502	100.0	97.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,972	22.2	21.6
2	Malignant neoplasms (C00-C97)	8,338	20.6	20.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,233	10.5	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,833	7.0	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,237	3.1	3.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,112	2.7	2.7
7	Diabetes mellitus (E10-E14)	1,022	2.5	2.5
8	Cerebrovascular diseases (I60-I69)	978	2.4	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	742	1.8	1.8
10	Influenza and pneumonia (J09-J18)	613	1.5	1.5
11	Septicemia (A40-A41)	578	1.4	1.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	578	1.4	1.4
13	Chronic lower respiratory diseases (J40-J47)	526	1.3	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	1.0	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	373	0.9	0.9
...	All other causes (Residual)	7,956	19.6	19.2
All races, female, 45-64 years				
...	All causes	201,529	100.0	473.5
1	Malignant neoplasms (C00-C97)	73,733	36.6	173.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,225	16.0	75.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,165	6.0	28.6
4	Chronic lower respiratory diseases (J40-J47)	10,132	5.0	23.8
5	Diabetes mellitus (E10-E14)	7,394	3.7	17.4
6	Cerebrovascular diseases (I60-I69)	7,071	3.5	16.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,393	3.2	15.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,990	2.0	9.4
9	Septicemia (A40-A41)	3,594	1.8	8.4
10	Influenza and pneumonia (J09-J18)	2,990	1.5	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,982	1.5	7.0
12	Viral hepatitis (B15-B19)	1,727	0.9	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,670	0.8	3.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,069	0.5	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	925	0.5	2.2
...	All other causes (Residual)	33,469	16.6	78.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	1,026,940	100.0	4,090.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252,696	24.6	1,006.6
2	Malignant neoplasms (C00-C97)	194,008	18.9	772.8
3	Chronic lower respiratory diseases (J40-J47)	68,085	6.6	271.2
4	Cerebrovascular diseases (I60-I69)	67,058	6.5	267.1
5	Alzheimer's disease (G30)	58,366	5.7	232.5
6	Diabetes mellitus (E10-E14)	27,231	2.7	108.5
7	Influenza and pneumonia (J09-J18)	26,314	2.6	104.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,873	2.2	91.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,155	2.0	80.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,939	1.6	63.5
11	Septicemia (A40-A41)	15,818	1.5	63.0
12	Parkinson's disease (G20-G21)	9,886	1.0	39.4
13	Pneumonitis due to solids and liquids (J69)	7,570	0.7	30.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,928	0.6	23.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,201	0.5	20.7
...	All other causes (Residual)	229,812	22.4	915.4
White, both sexes, all ages				
...	All causes	2,217,103	100.0	889.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	522,645	23.6	209.6
2	Malignant neoplasms (C00-C97)	498,116	22.5	199.8
3	Chronic lower respiratory diseases (J40-J47)	136,682	6.2	54.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112,803	5.1	45.2
5	Cerebrovascular diseases (I60-I69)	107,909	4.9	43.3
6	Alzheimer's disease (G30)	77,387	3.5	31.0
7	Diabetes mellitus (E10-E14)	58,925	2.7	23.6
8	Influenza and pneumonia (J09-J18)	49,013	2.2	19.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,270	1.7	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37,154	1.7	14.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	31,871	1.4	12.8
12	Septicemia (A40-A41)	30,889	1.4	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,149	1.1	9.7
14	Parkinson's disease (G20-G21)	23,553	1.1	9.4
15	Pneumonitis due to solids and liquids (J69)	16,458	0.7	6.6
...	All other causes (Residual)	452,279	20.4	181.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,774	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	33.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	345	12.4	2.9
3	Malignant neoplasms (C00-C97)	246	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	3.7	0.9
6	Influenza and pneumonia (J09-J18)	70	2.5	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.3	0.3
8	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	30	1.1	0.3
10	Septicemia (A40-A41)	29	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	20	0.7	0.2
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	11	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	709	25.6	5.9
White, both sexes, 5-14 years				
...	All causes	3,841	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,105	28.8	3.5
2	Malignant neoplasms (C00-C97)	717	18.7	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	8.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	6.3	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	4.4	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	2.7	0.3
7	Influenza and pneumonia (J09-J18)	92	2.4	0.3
8	Cerebrovascular diseases (I60-I69)	66	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	59	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.2	0.1
11	Septicemia (A40-A41)	38	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	24	0.6	0.1
13	Diabetes mellitus (E10-E14)	20	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
15	Meningitis (G00,G03)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	811	21.1	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	20,618	100.0	62.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,718	47.1	29.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,003	19.4	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,503	7.3	4.5
4	Malignant neoplasms (C00-C97)	1,138	5.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	602	2.9	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	1.4	0.8
7	Influenza and pneumonia (J09-J18)	159	0.8	0.5
8	Diabetes mellitus (E10-E14)	113	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	107	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	103	0.5	0.3
11	Septicemia (A40-A41)	92	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	68	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
14	Legal intervention (Y35,Y89.0)	49	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,578	12.5	7.8
White, both sexes, 25-44 years				
...	All causes	86,307	100.0	134.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,781	31.0	41.8
2	Malignant neoplasms (C00-C97)	11,487	13.3	17.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,218	13.0	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,254	10.7	14.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,855	3.3	4.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,613	3.0	4.1
7	Diabetes mellitus (E10-E14)	1,756	2.0	2.7
8	Cerebrovascular diseases (I60-I69)	1,395	1.6	2.2
9	Influenza and pneumonia (J09-J18)	1,011	1.2	1.6
10	Septicemia (A40-A41)	785	0.9	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	752	0.9	1.2
12	Chronic lower respiratory diseases (J40-J47)	710	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	690	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	506	0.6	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	455	0.5	0.7
...	All other causes (Residual)	14,039	16.3	21.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	408,371	100.0	606.0
1	Malignant neoplasms (C00-C97)	128,254	31.4	190.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,384	20.2	122.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,504	7.7	46.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	17,942	4.4	26.6
5	Chronic lower respiratory diseases (J40-J47)	17,774	4.4	26.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,698	3.6	21.8
7	Diabetes mellitus (E10-E14)	13,825	3.4	20.5
8	Cerebrovascular diseases (I60-I69)	11,623	2.8	17.2
9	Septicemia (A40-A41)	5,750	1.4	8.5
10	Influenza and pneumonia (J09-J18)	5,431	1.3	8.1
11	Viral hepatitis (B15-B19)	4,753	1.2	7.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,497	1.1	6.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,036	0.7	4.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,956	0.5	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,924	0.5	2.9
...	All other causes (Residual)	63,020	15.4	93.5
White, both sexes, 65 years and over				
...	All causes	1,679,966	100.0	4,363.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	429,985	25.6	1,116.7
2	Malignant neoplasms (C00-C97)	356,214	21.2	925.1
3	Chronic lower respiratory diseases (J40-J47)	118,026	7.0	306.5
4	Cerebrovascular diseases (I60-I69)	94,633	5.6	245.8
5	Alzheimer's disease (G30)	76,513	4.6	198.7
6	Diabetes mellitus (E10-E14)	43,203	2.6	112.2
7	Influenza and pneumonia (J09-J18)	42,149	2.5	109.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,053	2.5	109.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32,153	1.9	83.5
10	Septicemia (A40-A41)	24,103	1.4	62.6
11	Parkinson's disease (G20-G21)	22,966	1.4	59.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,824	1.2	54.1
13	Pneumonitis due to solids and liquids (J69)	14,725	0.9	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,316	0.7	29.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11,284	0.7	29.3
...	All other causes (Residual)	339,819	20.2	882.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,110,956	100.0	899.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	275,101	24.8	222.6
2	Malignant neoplasms (C00-C97)	263,167	23.7	213.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70,161	6.3	56.8
4	Chronic lower respiratory diseases (J40-J47)	63,757	5.7	51.6
5	Cerebrovascular diseases (I60-I69)	44,203	4.0	35.8
6	Diabetes mellitus (E10-E14)	31,745	2.9	25.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,943	2.6	23.4
8	Alzheimer's disease (G30)	23,648	2.1	19.1
9	Influenza and pneumonia (J09-J18)	22,907	2.1	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20,884	1.9	16.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,800	1.7	15.2
12	Septicemia (A40-A41)	14,602	1.3	11.8
13	Parkinson's disease (G20-G21)	14,127	1.3	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,940	0.9	8.0
15	Pneumonitis due to solids and liquids (J69)	8,989	0.8	7.3
...	All other causes (Residual)	199,982	18.0	161.9
White, male, 1-4 years				
...	All causes	1,601	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	37.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	10.1	2.7
3	Malignant neoplasms (C00-C97)	142	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	0.9
6	Influenza and pneumonia (J09-J18)	39	2.4	0.6
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Septicemia (A40-A41)	16	1.0	*
11	Cerebrovascular diseases (I60-I69)	9	0.6	*
12	Meningitis (G00,G03)	7	0.4	*
13	Anemias (D50-D64)	6	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	393	24.5	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,219	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	684	30.8	4.3
2	Malignant neoplasms (C00-C97)	371	16.7	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	9.1	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	5.9	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	4.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	2.4	0.3
7	Influenza and pneumonia (J09-J18)	48	2.2	0.3
8	Chronic lower respiratory diseases (J40-J47)	38	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	31	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.2	0.2
11	Septicemia (A40-A41)	21	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
15	Meningitis (G00,G03)	8	0.4	*
...	All other causes (Residual)	470	21.2	2.9
White, male, 15-24 years				
...	All causes	14,880	100.0	87.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,172	48.2	42.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,203	21.5	18.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	8.1	7.1
4	Malignant neoplasms (C00-C97)	664	4.5	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	398	2.7	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	1.2	1.0
7	Influenza and pneumonia (J09-J18)	94	0.6	0.5
8	Diabetes mellitus (E10-E14)	67	0.5	0.4
9	Cerebrovascular diseases (I60-I69)	61	0.4	0.4
10	Septicemia (A40-A41)	50	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.3
12	Legal intervention (Y35,Y89.0)	45	0.3	0.3
13	Chronic lower respiratory diseases (J40-J47)	42	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.1	0.1
...	All other causes (Residual)	1,610	10.8	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	56,319	100.0	172.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,165	34.0	58.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,731	15.5	26.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,497	11.5	19.9
4	Malignant neoplasms (C00-C97)	5,214	9.3	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,128	3.8	6.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,744	3.1	5.4
7	Diabetes mellitus (E10-E14)	1,100	2.0	3.4
8	Cerebrovascular diseases (I60-I69)	799	1.4	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	603	1.1	1.9
10	Influenza and pneumonia (J09-J18)	546	1.0	1.7
11	Septicemia (A40-A41)	401	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	393	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	330	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	0.5	0.8
15	Aortic aneurysm and dissection (I71)	210	0.4	0.6
...	All other causes (Residual)	8,185	14.5	25.1
White, male, 45-64 years				
...	All causes	251,660	100.0	757.2
1	Malignant neoplasms (C00-C97)	69,451	27.6	209.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,269	23.6	178.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,165	8.4	63.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	12,539	5.0	37.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,981	4.4	33.0
6	Chronic lower respiratory diseases (J40-J47)	9,031	3.6	27.2
7	Diabetes mellitus (E10-E14)	8,674	3.4	26.1
8	Cerebrovascular diseases (I60-I69)	6,759	2.7	20.3
9	Viral hepatitis (B15-B19)	3,400	1.4	10.2
10	Septicemia (A40-A41)	3,121	1.2	9.4
11	Influenza and pneumonia (J09-J18)	3,117	1.2	9.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,571	1.0	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,012	0.8	6.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,600	0.6	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,331	0.5	4.0
...	All other causes (Residual)	36,639	14.6	110.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	775,654	100.0	4,550.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	208,697	26.9	1,224.3
2	Malignant neoplasms (C00-C97)	187,291	24.1	1,098.7
3	Chronic lower respiratory diseases (J40-J47)	54,289	7.0	318.5
4	Cerebrovascular diseases (I60-I69)	36,509	4.7	214.2
5	Alzheimer's disease (G30)	23,293	3.0	136.6
6	Diabetes mellitus (E10-E14)	21,887	2.8	128.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,937	2.7	122.8
8	Influenza and pneumonia (J09-J18)	18,998	2.4	111.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,894	2.0	93.2
10	Parkinson's disease (G20-G21)	13,736	1.8	80.6
11	Septicemia (A40-A41)	10,940	1.4	64.2
12	Pneumonitis due to solids and liquids (J69)	7,957	1.0	46.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,729	1.0	45.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,580	0.8	38.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,958	0.8	35.0
...	All other causes (Residual)	134,959	17.4	791.7
White, female, all ages				
...	All causes	1,106,147	100.0	879.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,544	22.4	196.8
2	Malignant neoplasms (C00-C97)	234,949	21.2	186.8
3	Chronic lower respiratory diseases (J40-J47)	72,925	6.6	58.0
4	Cerebrovascular diseases (I60-I69)	63,706	5.8	50.6
5	Alzheimer's disease (G30)	53,739	4.9	42.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,642	3.9	33.9
7	Diabetes mellitus (E10-E14)	27,180	2.5	21.6
8	Influenza and pneumonia (J09-J18)	26,106	2.4	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,470	1.7	14.7
10	Septicemia (A40-A41)	16,287	1.5	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,209	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,987	1.0	8.7
13	Parkinson's disease (G20-G21)	9,426	0.9	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,211	0.7	6.5
15	Pneumonitis due to solids and liquids (J69)	7,469	0.7	5.9
...	All other causes (Residual)	252,297	22.8	200.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,173	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	27.7	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	15.6	3.1
3	Malignant neoplasms (C00-C97)	104	8.9	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.1	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
8	Septicemia (A40-A41)	13	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	11	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Meningitis (G00,G03)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	316	26.9	5.4
White, female, 5-14 years				
...	All causes	1,622	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	26.0	2.8
2	Malignant neoplasms (C00-C97)	346	21.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	6.9	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	6.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	4.1	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	3.2	0.3
7	Influenza and pneumonia (J09-J18)	44	2.7	0.3
8	Cerebrovascular diseases (I60-I69)	35	2.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.1
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	17	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	9	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	346	21.3	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,738	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,546	44.4	15.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	800	13.9	5.0
3	Malignant neoplasms (C00-C97)	474	8.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	291	5.1	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	3.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	1.9	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	103	1.8	0.6
8	Influenza and pneumonia (J09-J18)	65	1.1	0.4
9	Diabetes mellitus (E10-E14)	46	0.8	0.3
9	Cerebrovascular diseases (I60-I69)	46	0.8	0.3
11	Septicemia (A40-A41)	42	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
...	All other causes (Residual)	935	16.3	5.8
White, female, 25-44 years				
...	All causes	29,988	100.0	95.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,616	25.4	24.2
2	Malignant neoplasms (C00-C97)	6,273	20.9	19.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,757	9.2	8.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,487	8.3	7.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	869	2.9	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	727	2.4	2.3
7	Diabetes mellitus (E10-E14)	656	2.2	2.1
8	Cerebrovascular diseases (I60-I69)	596	2.0	1.9
9	Influenza and pneumonia (J09-J18)	465	1.6	1.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	455	1.5	1.4
11	Septicemia (A40-A41)	384	1.3	1.2
12	Chronic lower respiratory diseases (J40-J47)	380	1.3	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	297	1.0	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	0.8	0.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	149	0.5	0.5
...	All other causes (Residual)	5,644	18.8	17.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	156,711	100.0	458.8
1	Malignant neoplasms (C00-C97)	58,803	37.5	172.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,115	14.8	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,339	6.6	30.3
4	Chronic lower respiratory diseases (J40-J47)	8,743	5.6	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,403	3.4	15.8
6	Diabetes mellitus (E10-E14)	5,151	3.3	15.1
7	Cerebrovascular diseases (I60-I69)	4,864	3.1	14.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,717	2.4	10.9
9	Septicemia (A40-A41)	2,629	1.7	7.7
10	Influenza and pneumonia (J09-J18)	2,314	1.5	6.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,926	1.2	5.6
12	Viral hepatitis (B15-B19)	1,353	0.9	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,024	0.7	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	732	0.5	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	657	0.4	1.9
...	All other causes (Residual)	25,941	16.6	75.9
White, female, 65 years and over				
...	All causes	904,312	100.0	4,214.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221,288	24.5	1,031.2
2	Malignant neoplasms (C00-C97)	168,923	18.7	787.2
3	Chronic lower respiratory diseases (J40-J47)	63,737	7.0	297.0
4	Cerebrovascular diseases (I60-I69)	58,124	6.4	270.9
5	Alzheimer's disease (G30)	53,220	5.9	248.0
6	Influenza and pneumonia (J09-J18)	23,151	2.6	107.9
7	Diabetes mellitus (E10-E14)	21,316	2.4	99.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,116	2.3	98.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,259	1.8	75.8
10	Septicemia (A40-A41)	13,163	1.5	61.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,095	1.4	61.0
12	Parkinson's disease (G20-G21)	9,230	1.0	43.0
13	Pneumonitis due to solids and liquids (J69)	6,768	0.7	31.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,358	0.6	25.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,704	0.5	21.9
...	All other causes (Residual)	204,860	22.7	954.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	302,969	100.0	693.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,010	23.8	164.8
2	Malignant neoplasms (C00-C97)	67,953	22.4	155.5
3	Cerebrovascular diseases (I60-I69)	16,269	5.4	37.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,413	4.4	30.7
5	Diabetes mellitus (E10-E14)	13,385	4.4	30.6
6	Chronic lower respiratory diseases (J40-J47)	9,918	3.3	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,393	2.8	19.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,059	2.7	18.4
9	Septicemia (A40-A41)	6,250	2.1	14.3
10	Alzheimer's disease (G30)	5,714	1.9	13.1
11	Influenza and pneumonia (J09-J18)	5,567	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,455	1.8	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.3	9.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,742	1.2	8.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,053	1.0	7.0
...	All other causes (Residual)	59,733	19.7	136.7
Black, both sexes, 1-4 years				
...	All causes	1,006	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	29.7	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	13.5	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	9.0	3.4
4	Malignant neoplasms (C00-C97)	64	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	5.9	2.2
6	Chronic lower respiratory diseases (J40-J47)	27	2.7	1.0
7	Influenza and pneumonia (J09-J18)	23	2.3	0.8
8	Septicemia (A40-A41)	20	2.0	0.7
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	13	1.3	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Meningitis (G00,G03)	6	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	231	23.0	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,169	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	28.2	4.9
2	Malignant neoplasms (C00-C97)	122	10.4	1.8
3	Chronic lower respiratory diseases (J40-J47)	87	7.4	1.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	7.4	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	7.2	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	5.0	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.4	0.8
8	Influenza and pneumonia (J09-J18)	28	2.4	0.4
9	Cerebrovascular diseases (I60-I69)	19	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
11	Septicemia (A40-A41)	13	1.1	*
12	Anemias (D50-D64)	11	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	247	21.1	3.6
Black, both sexes, 15-24 years				
...	All causes	6,538	100.0	88.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,731	41.8	37.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,415	21.6	19.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	523	8.0	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	290	4.4	3.9
5	Malignant neoplasms (C00-C97)	241	3.7	3.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.1
8	Diabetes mellitus (E10-E14)	73	1.1	1.0
9	Anemias (D50-D64)	70	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.1	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	0.8
12	Legal intervention (Y35,Y89.0)	44	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.6	0.5
14	Septicemia (A40-A41)	36	0.6	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.4
...	All other causes (Residual)	754	11.5	10.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	23,290	100.0	197.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,773	16.2	31.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,717	16.0	31.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,576	15.4	30.2
4	Malignant neoplasms (C00-C97)	2,562	11.0	21.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,079	4.6	9.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	995	4.3	8.4
7	Diabetes mellitus (E10-E14)	760	3.3	6.4
8	Cerebrovascular diseases (I60-I69)	641	2.8	5.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	345	1.5	2.9
10	Septicemia (A40-A41)	318	1.4	2.7
11	Chronic lower respiratory diseases (J40-J47)	307	1.3	2.6
12	Influenza and pneumonia (J09-J18)	257	1.1	2.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	241	1.0	2.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	237	1.0	2.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	235	1.0	2.0
...	All other causes (Residual)	4,247	18.2	35.9
Black, both sexes, 45-64 years				
...	All causes	90,220	100.0	873.6
1	Malignant neoplasms (C00-C97)	25,371	28.1	245.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,988	24.4	212.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,774	5.3	46.2
4	Diabetes mellitus (E10-E14)	4,308	4.8	41.7
5	Cerebrovascular diseases (I60-I69)	4,296	4.8	41.6
6	Chronic lower respiratory diseases (J40-J47)	2,459	2.7	23.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,219	2.5	21.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,174	2.4	21.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,022	2.2	19.6
10	Septicemia (A40-A41)	1,817	2.0	17.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,458	1.6	14.1
12	Influenza and pneumonia (J09-J18)	1,315	1.5	12.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,119	1.2	10.8
14	Viral hepatitis (B15-B19)	857	0.9	8.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	625	0.7	6.1
...	All other causes (Residual)	13,418	14.9	129.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	173,597	100.0	4,312.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,745	26.4	1,136.4
2	Malignant neoplasms (C00-C97)	39,583	22.8	983.3
3	Cerebrovascular diseases (I60-I69)	11,220	6.5	278.7
4	Diabetes mellitus (E10-E14)	8,235	4.7	204.6
5	Chronic lower respiratory diseases (J40-J47)	6,947	4.0	172.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,784	3.3	143.7
7	Alzheimer's disease (G30)	5,626	3.2	139.8
8	Septicemia (A40-A41)	3,991	2.3	99.1
9	Influenza and pneumonia (J09-J18)	3,842	2.2	95.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,755	2.2	93.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,587	1.5	64.3
12	Pneumonitis due to solids and liquids (J69)	1,245	0.7	30.9
13	Parkinson's disease (G20-G21)	984	0.6	24.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	802	0.5	19.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	789	0.5	19.6
...	All other causes (Residual)	32,462	18.7	806.4
Black, male, all ages				
...	All causes	154,767	100.0	739.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,096	24.0	177.2
2	Malignant neoplasms (C00-C97)	34,671	22.4	165.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,017	5.8	43.1
4	Cerebrovascular diseases (I60-I69)	7,338	4.7	35.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,937	4.5	33.1
6	Diabetes mellitus (E10-E14)	6,378	4.1	30.5
7	Chronic lower respiratory diseases (J40-J47)	5,073	3.3	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,976	2.6	19.0
9	Septicemia (A40-A41)	2,882	1.9	13.8
10	Influenza and pneumonia (J09-J18)	2,696	1.7	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,496	1.6	11.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,479	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,215	1.4	10.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,955	1.3	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,891	1.2	9.0
...	All other causes (Residual)	27,667	17.9	132.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	561	100.0	40.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	31.6	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	12.5	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.9	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.4	2.6
5	Malignant neoplasms (C00-C97)	33	5.9	2.4
6	Chronic lower respiratory diseases (J40-J47)	15	2.7	*
7	Septicemia (A40-A41)	13	2.3	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
9	Cerebrovascular diseases (I60-I69)	12	2.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	122	21.7	8.8
Black, male, 5-14 years				
...	All causes	675	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	31.9	6.2
2	Malignant neoplasms (C00-C97)	66	9.8	1.9
3	Chronic lower respiratory diseases (J40-J47)	55	8.1	1.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	8.1	1.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.9	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	5.3	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.6	0.9
8	Influenza and pneumonia (J09-J18)	18	2.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.6	*
10	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Septicemia (A40-A41)	8	1.2	*
12	Anemias (D50-D64)	5	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	116	17.2	3.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,041	100.0	135.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,479	49.2	66.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,023	20.3	27.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	8.5	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	3.8	5.2
5	Malignant neoplasms (C00-C97)	129	2.6	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.6
7	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.5
8	Diabetes mellitus (E10-E14)	48	1.0	1.3
9	Anemias (D50-D64)	44	0.9	1.2
9	Legal intervention (Y35,Y89.0)	44	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.7	0.9
12	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
13	Septicemia (A40-A41)	19	0.4	*
14	Influenza and pneumonia (J09-J18)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	435	8.6	11.7
Black, male, 25-44 years				
...	All causes	14,812	100.0	262.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,272	22.1	57.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,581	17.4	45.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,444	16.5	43.3
4	Malignant neoplasms (C00-C97)	1,026	6.9	18.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	814	5.5	14.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	663	4.5	11.7
7	Diabetes mellitus (E10-E14)	442	3.0	7.8
8	Cerebrovascular diseases (I60-I69)	343	2.3	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	1.3	3.4
10	Chronic lower respiratory diseases (J40-J47)	172	1.2	3.0
11	Septicemia (A40-A41)	154	1.0	2.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.0	2.5
13	Influenza and pneumonia (J09-J18)	137	0.9	2.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	129	0.9	2.3
15	Anemias (D50-D64)	112	0.8	2.0
...	All other causes (Residual)	2,191	14.8	38.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	52,347	100.0	1,093.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,799	26.4	288.4
2	Malignant neoplasms (C00-C97)	13,275	25.4	277.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,343	6.4	69.9
4	Cerebrovascular diseases (I60-I69)	2,461	4.7	51.4
5	Diabetes mellitus (E10-E14)	2,391	4.6	50.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,476	2.8	30.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,335	2.6	27.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,282	2.4	26.8
9	Chronic lower respiratory diseases (J40-J47)	1,223	2.3	25.6
10	Septicemia (A40-A41)	941	1.8	19.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	882	1.7	18.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	864	1.7	18.1
13	Influenza and pneumonia (J09-J18)	752	1.4	15.7
14	Viral hepatitis (B15-B19)	577	1.1	12.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	477	0.9	10.0
...	All other causes (Residual)	7,269	13.9	151.9
Black, male, 65 years and over				
...	All causes	77,436	100.0	4,818.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,534	26.5	1,277.7
2	Malignant neoplasms (C00-C97)	20,140	26.0	1,253.2
3	Cerebrovascular diseases (I60-I69)	4,468	5.8	278.0
4	Chronic lower respiratory diseases (J40-J47)	3,547	4.6	220.7
5	Diabetes mellitus (E10-E14)	3,493	4.5	217.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,477	3.2	154.1
7	Influenza and pneumonia (J09-J18)	1,720	2.2	107.0
8	Septicemia (A40-A41)	1,716	2.2	106.8
9	Alzheimer's disease (G30)	1,612	2.1	100.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,486	1.9	92.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,451	1.9	90.3
12	Pneumonitis due to solids and liquids (J69)	646	0.8	40.2
13	Parkinson's disease (G20-G21)	581	0.8	36.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	490	0.6	30.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	389	0.5	24.2
...	All other causes (Residual)	12,686	16.4	789.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	148,202	100.0	651.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,914	23.6	153.4
2	Malignant neoplasms (C00-C97)	33,282	22.5	146.2
3	Cerebrovascular diseases (I60-I69)	8,931	6.0	39.2
4	Diabetes mellitus (E10-E14)	7,007	4.7	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,845	3.3	21.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,417	3.0	19.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,396	3.0	19.3
8	Alzheimer's disease (G30)	4,050	2.7	17.8
9	Septicemia (A40-A41)	3,368	2.3	14.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,959	2.0	13.0
11	Influenza and pneumonia (J09-J18)	2,871	1.9	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,840	1.2	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,263	0.9	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,122	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,098	0.7	4.8
...	All other causes (Residual)	31,839	21.5	139.9
Black, female, 1-4 years				
...	All causes	445	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	27.4	9.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	14.8	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	9.2	3.1
4	Malignant neoplasms (C00-C97)	31	7.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	5.2	1.7
6	Chronic lower respiratory diseases (J40-J47)	12	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.2	*
8	Septicemia (A40-A41)	7	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	103	23.1	7.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	494	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	23.3	3.5
2	Malignant neoplasms (C00-C97)	56	11.3	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	8.9	1.3
4	Chronic lower respiratory diseases (J40-J47)	32	6.5	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.5	1.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.5	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.0	0.6
8	Influenza and pneumonia (J09-J18)	10	2.0	*
9	Cerebrovascular diseases (I60-I69)	9	1.8	*
10	Anemias (D50-D64)	6	1.2	*
11	Septicemia (A40-A41)	5	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
15	Meningitis (G00,G03)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	124	25.1	3.7
Black, female, 15-24 years				
...	All causes	1,497	100.0	41.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	26.2	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	252	16.8	6.9
3	Malignant neoplasms (C00-C97)	112	7.5	3.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.4	2.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	6.3	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	69	4.6	1.9
7	Anemias (D50-D64)	26	1.7	0.7
8	Diabetes mellitus (E10-E14)	25	1.7	0.7
8	Chronic lower respiratory diseases (J40-J47)	25	1.7	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.5	0.6
12	Septicemia (A40-A41)	17	1.1	*
12	Cerebrovascular diseases (I60-I69)	17	1.1	*
14	Influenza and pneumonia (J09-J18)	16	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	303	20.2	8.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	8,478	100.0	137.3
1	Malignant neoplasms (C00-C97)	1,536	18.1	24.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,329	15.7	21.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	995	11.7	16.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	445	5.2	7.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	416	4.9	6.7
6	Diabetes mellitus (E10-E14)	318	3.8	5.2
7	Cerebrovascular diseases (I60-I69)	298	3.5	4.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	237	2.8	3.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	2.1	2.9
10	Septicemia (A40-A41)	164	1.9	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	155	1.8	2.5
12	Chronic lower respiratory diseases (J40-J47)	135	1.6	2.2
13	Influenza and pneumonia (J09-J18)	120	1.4	1.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	112	1.3	1.8
15	Anemias (D50-D64)	102	1.2	1.7
...	All other causes (Residual)	1,935	22.8	31.3
Black, female, 45-64 years				
...	All causes	37,873	100.0	683.4
1	Malignant neoplasms (C00-C97)	12,096	31.9	218.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,189	21.6	147.8
3	Diabetes mellitus (E10-E14)	1,917	5.1	34.6
4	Cerebrovascular diseases (I60-I69)	1,835	4.8	33.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,431	3.8	25.8
6	Chronic lower respiratory diseases (J40-J47)	1,236	3.3	22.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	937	2.5	16.9
8	Septicemia (A40-A41)	876	2.3	15.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	698	1.8	12.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	687	1.8	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	594	1.6	10.7
12	Influenza and pneumonia (J09-J18)	563	1.5	10.2
13	Viral hepatitis (B15-B19)	280	0.7	5.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	0.6	4.3
15	Anemias (D50-D64)	148	0.4	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	0.4	2.7
...	All other causes (Residual)	6,001	15.8	108.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	96,161	100.0	3,976.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,211	26.2	1,042.4
2	Malignant neoplasms (C00-C97)	19,443	20.2	803.9
3	Cerebrovascular diseases (I60-I69)	6,752	7.0	279.2
4	Diabetes mellitus (E10-E14)	4,742	4.9	196.1
5	Alzheimer's disease (G30)	4,014	4.2	166.0
6	Chronic lower respiratory diseases (J40-J47)	3,400	3.5	140.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,307	3.4	136.7
8	Septicemia (A40-A41)	2,275	2.4	94.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,269	2.4	93.8
10	Influenza and pneumonia (J09-J18)	2,122	2.2	87.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,136	1.2	47.0
12	Pneumonitis due to solids and liquids (J69)	599	0.6	24.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	413	0.4	17.1
14	Parkinson's disease (G20-G21)	403	0.4	16.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	299	0.3	12.4
...	All other causes (Residual)	19,776	20.6	817.7
American Indian, both sexes, all ages				
...	All causes	17,052	100.0	382.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,139	18.4	70.4
2	Malignant neoplasms (C00-C97)	3,109	18.2	69.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	10.7	41.1
4	Diabetes mellitus (E10-E14)	959	5.6	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	944	5.5	21.2
6	Chronic lower respiratory diseases (J40-J47)	757	4.4	17.0
7	Cerebrovascular diseases (I60-I69)	595	3.5	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	521	3.1	11.7
9	Influenza and pneumonia (J09-J18)	375	2.2	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	302	1.8	6.8
11	Septicemia (A40-A41)	288	1.7	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	1.4	5.4
13	Alzheimer's disease (G30)	238	1.4	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	183	1.1	4.1
15	Viral hepatitis (B15-B19)	133	0.8	3.0
...	All other causes (Residual)	3,435	20.1	77.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	104	100.0	33.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	41.3	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	15.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	10.6	*
4	Malignant neoplasms (C00-C97)	3	2.9	*
4	Influenza and pneumonia (J09-J18)	3	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.9	*
7	Nutritional deficiencies (E40-E64)	1	1.0	*
7	Cerebrovascular diseases (I60-I69)	1	1.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
...	All other causes (Residual)	21	20.2	6.8
American Indian, both sexes, 5-14 years				
...	All causes	88	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	39.8	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	12.5	*
3	Malignant neoplasms (C00-C97)	10	11.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.3	*
8	Viral hepatitis (B15-B19)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Pneumonitis due to solids and liquids (J69)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
...	All other causes (Residual)	15	17.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	548	100.0	71.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	44.7	31.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	27.4	19.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	7.3	5.2
4	Malignant neoplasms (C00-C97)	15	2.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.3	*
7	Diabetes mellitus (E10-E14)	5	0.9	*
8	Septicemia (A40-A41)	3	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	52	9.5	6.8
American Indian, both sexes, 25-44 years				
...	All causes	2,149	100.0	169.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	657	30.6	51.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	238	11.1	18.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	10.2	17.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	8.3	14.1
5	Malignant neoplasms (C00-C97)	136	6.3	10.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	5.8	9.7
7	Diabetes mellitus (E10-E14)	72	3.4	5.7
8	Influenza and pneumonia (J09-J18)	37	1.7	2.9
9	Septicemia (A40-A41)	28	1.3	2.2
9	Cerebrovascular diseases (I60-I69)	28	1.3	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.6	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
15	Viral hepatitis (B15-B19)	9	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.4	*
...	All other causes (Residual)	362	16.8	28.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	5,403	100.0	568.5
1	Malignant neoplasms (C00-C97)	1,135	21.0	119.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	988	18.3	103.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	573	10.6	60.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	552	10.2	58.1
5	Diabetes mellitus (E10-E14)	345	6.4	36.3
6	Chronic lower respiratory diseases (J40-J47)	159	2.9	16.7
7	Cerebrovascular diseases (I60-I69)	136	2.5	14.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	2.2	12.3
9	Viral hepatitis (B15-B19)	105	1.9	11.0
10	Septicemia (A40-A41)	103	1.9	10.8
11	Influenza and pneumonia (J09-J18)	89	1.6	9.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	1.6	9.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.0	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	0.8	4.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.6	3.7
...	All other causes (Residual)	877	16.2	92.3
American Indian, both sexes, 65 years and over				
...	All causes	8,441	100.0	2,704.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,952	23.1	625.4
2	Malignant neoplasms (C00-C97)	1,810	21.4	579.9
3	Chronic lower respiratory diseases (J40-J47)	581	6.9	186.2
4	Diabetes mellitus (E10-E14)	536	6.3	171.7
5	Cerebrovascular diseases (I60-I69)	427	5.1	136.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	3.0	81.7
7	Influenza and pneumonia (J09-J18)	237	2.8	75.9
8	Alzheimer's disease (G30)	232	2.7	74.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.3	62.5
10	Septicemia (A40-A41)	152	1.8	48.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	151	1.8	48.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.4	38.8
13	Parkinson's disease (G20-G21)	65	0.8	20.8
14	Pneumonitis due to solids and liquids (J69)	59	0.7	18.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.4	11.5
...	All other causes (Residual)	1,632	19.3	522.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	9,331	100.0	416.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,843	19.8	82.3
2	Malignant neoplasms (C00-C97)	1,650	17.7	73.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,177	12.6	52.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	511	5.5	22.8
5	Diabetes mellitus (E10-E14)	490	5.3	21.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	401	4.3	17.9
7	Chronic lower respiratory diseases (J40-J47)	370	4.0	16.5
8	Cerebrovascular diseases (I60-I69)	253	2.7	11.3
9	Influenza and pneumonia (J09-J18)	190	2.0	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	2.0	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.5	6.4
12	Septicemia (A40-A41)	139	1.5	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.0	4.2
14	Viral hepatitis (B15-B19)	80	0.9	3.6
15	Alzheimer's disease (G30)	77	0.8	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	3.4
...	All other causes (Residual)	1,649	17.7	73.6
American Indian, male, 1-4 years				
...	All causes	65	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	36.9	15.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	16.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	10.8	*
4	Malignant neoplasms (C00-C97)	2	3.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.1	*
6	Nutritional deficiencies (E40-E64)	1	1.5	*
6	Cerebrovascular diseases (I60-I69)	1	1.5	*
6	Influenza and pneumonia (J09-J18)	1	1.5	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.5	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
...	All other causes (Residual)	14	21.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	47	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	38.3	*
2	Malignant neoplasms (C00-C97)	8	17.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.9	*
4	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.1	*
7	Pneumonitis due to solids and liquids (J69)	1	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (Residual)	5	10.6	*
American Indian, male, 15-24 years				
...	All causes	388	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	44.3	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	29.6	29.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	7.2	7.1
4	Malignant neoplasms (C00-C97)	9	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.5	*
7	Diabetes mellitus (E10-E14)	4	1.0	*
8	Legal intervention (Y35,Y89.0)	3	0.8	*
9	Septicemia (A40-A41)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	32	8.2	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-44 years				
...	All causes	1,345	100.0	205.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	32.0	65.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	12.9	26.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	131	9.7	20.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	9.1	18.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	7.5	15.4
6	Malignant neoplasms (C00-C97)	61	4.5	9.3
7	Diabetes mellitus (E10-E14)	41	3.0	6.3
8	Influenza and pneumonia (J09-J18)	16	1.2	*
9	Septicemia (A40-A41)	13	1.0	*
10	Cerebrovascular diseases (I60-I69)	11	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
14	Viral hepatitis (B15-B19)	6	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.4	*
...	All other causes (Residual)	211	15.7	32.2
American Indian, male, 45-64 years				
...	All causes	3,190	100.0	685.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	675	21.2	145.1
2	Malignant neoplasms (C00-C97)	620	19.4	133.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	11.8	81.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	305	9.6	65.6
5	Diabetes mellitus (E10-E14)	193	6.1	41.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	2.7	18.3
7	Cerebrovascular diseases (I60-I69)	71	2.2	15.3
8	Chronic lower respiratory diseases (J40-J47)	67	2.1	14.4
9	Viral hepatitis (B15-B19)	63	2.0	13.5
10	Septicemia (A40-A41)	56	1.8	12.0
11	Influenza and pneumonia (J09-J18)	54	1.7	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.3	8.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.2	8.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	1.2	8.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.9	6.0
...	All other causes (Residual)	479	15.0	103.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65 years and over				
...	All causes	4,096	100.0	2,930.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,034	25.2	739.7
2	Malignant neoplasms (C00-C97)	950	23.2	679.6
3	Chronic lower respiratory diseases (J40-J47)	291	7.1	208.2
4	Diabetes mellitus (E10-E14)	252	6.2	180.3
5	Cerebrovascular diseases (I60-I69)	169	4.1	120.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	3.4	99.4
7	Influenza and pneumonia (J09-J18)	114	2.8	81.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	2.3	68.0
9	Alzheimer's disease (G30)	73	1.8	52.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.8	52.2
11	Septicemia (A40-A41)	67	1.6	47.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.2	35.8
13	Parkinson's disease (G20-G21)	38	0.9	27.2
14	Pneumonitis due to solids and liquids (J69)	31	0.8	22.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	15.7
...	All other causes (Residual)	698	17.0	499.3
American Indian, female, all ages				
...	All causes	7,721	100.0	348.2
1	Malignant neoplasms (C00-C97)	1,459	18.9	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	16.8	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	8.5	29.6
4	Diabetes mellitus (E10-E14)	469	6.1	21.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	433	5.6	19.5
6	Chronic lower respiratory diseases (J40-J47)	387	5.0	17.5
7	Cerebrovascular diseases (I60-I69)	342	4.4	15.4
8	Influenza and pneumonia (J09-J18)	185	2.4	8.3
9	Alzheimer's disease (G30)	161	2.1	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.1	7.2
11	Septicemia (A40-A41)	149	1.9	6.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	1.6	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.2	4.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.7	2.4
15	Viral hepatitis (B15-B19)	53	0.7	2.4
...	All other causes (Residual)	1,709	22.1	77.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	48.7	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	12.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
5	Malignant neoplasms (C00-C97)	1	2.6	*
5	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
...	All other causes (Residual)	7	17.9	*
American Indian, female, 5-14 years				
...	All causes	41	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	41.5	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	9.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.8	*
4	Malignant neoplasms (C00-C97)	2	4.9	*
5	Viral hepatitis (B15-B19)	1	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
5	Influenza and pneumonia (J09-J18)	1	2.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.4	*
...	All other causes (Residual)	10	24.4	*
American Indian, female, 15-24 years				
...	All causes	160	100.0	42.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	45.6	19.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	21.9	9.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	7.5	*
4	Malignant neoplasms (C00-C97)	6	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.9	*
7	Septicemia (A40-A41)	1	0.6	*
7	Diabetes mellitus (E10-E14)	1	0.6	*
7	Influenza and pneumonia (J09-J18)	1	0.6	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	20	12.5	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-44 years				
...	All causes	804	100.0	130.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	28.1	36.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	107	13.3	17.3
3	Malignant neoplasms (C00-C97)	75	9.3	12.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	7.1	9.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	5.7	7.5
6	Diabetes mellitus (E10-E14)	31	3.9	5.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.9	3.7
8	Influenza and pneumonia (J09-J18)	21	2.6	3.4
9	Cerebrovascular diseases (I60-I69)	17	2.1	*
10	Septicemia (A40-A41)	15	1.9	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
15	Viral hepatitis (B15-B19)	3	0.4	*
...	All other causes (Residual)	153	19.0	24.8
American Indian, female, 45-64 years				
...	All causes	2,213	100.0	455.9
1	Malignant neoplasms (C00-C97)	515	23.3	106.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	14.1	64.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	247	11.2	50.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	8.8	40.2
5	Diabetes mellitus (E10-E14)	152	6.9	31.3
6	Chronic lower respiratory diseases (J40-J47)	92	4.2	19.0
7	Cerebrovascular diseases (I60-I69)	65	2.9	13.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.3	10.3
9	Septicemia (A40-A41)	47	2.1	9.7
10	Viral hepatitis (B15-B19)	42	1.9	8.7
11	Influenza and pneumonia (J09-J18)	35	1.6	7.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	1.4	6.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.4	*
...	All other causes (Residual)	394	17.8	81.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65 years and over				
...	All causes	4,345	100.0	2,521.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	918	21.1	532.7
2	Malignant neoplasms (C00-C97)	860	19.8	499.1
3	Chronic lower respiratory diseases (J40-J47)	290	6.7	168.3
4	Diabetes mellitus (E10-E14)	284	6.5	164.8
5	Cerebrovascular diseases (I60-I69)	258	5.9	149.7
6	Alzheimer's disease (G30)	159	3.7	92.3
7	Influenza and pneumonia (J09-J18)	123	2.8	71.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	2.7	67.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	2.3	58.0
10	Septicemia (A40-A41)	85	2.0	49.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	78	1.8	45.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.6	41.2
13	Pneumonitis due to solids and liquids (J69)	28	0.6	16.2
14	Parkinson's disease (G20-G21)	27	0.6	15.7
15	Aortic aneurysm and dissection (I71)	16	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	16	0.4	*
...	All other causes (Residual)	916	21.1	531.6
Asian or Pacific Islander, both sexes, all ages				
...	All causes	59,869	100.0	321.4
1	Malignant neoplasms (C00-C97)	15,703	26.2	84.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,311	22.2	71.4
3	Cerebrovascular diseases (I60-I69)	4,205	7.0	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,508	4.2	13.5
5	Diabetes mellitus (E10-E14)	2,309	3.9	12.4
6	Influenza and pneumonia (J09-J18)	2,024	3.4	10.9
7	Chronic lower respiratory diseases (J40-J47)	1,848	3.1	9.9
8	Alzheimer's disease (G30)	1,428	2.4	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,147	1.9	6.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,121	1.9	6.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	983	1.6	5.3
12	Septicemia (A40-A41)	729	1.2	3.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	559	0.9	3.0
14	Parkinson's disease (G20-G21)	528	0.9	2.8
15	Certain conditions originating in the perinatal period (P00-P96)	487	0.8	2.6
...	All other causes (Residual)	10,979	18.3	58.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	184	100.0	18.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	24.5	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	15.8	3.0
3	Malignant neoplasms (C00-C97)	15	8.2	*
4	Influenza and pneumonia (J09-J18)	6	3.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.7	*
7	Septicemia (A40-A41)	4	2.2	*
7	Meningitis (G00,G03)	4	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.6	*
11	Cerebrovascular diseases (I60-I69)	2	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
13	Shigellosis and amebiasis (A03,A06)	1	0.5	*
13	Anemias (D50-D64)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
...	All other causes (Residual)	55	29.9	5.6
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	242	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	21.1	2.1
2	Malignant neoplasms (C00-C97)	46	19.0	1.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.2	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	5.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	4.5	*
7	Influenza and pneumonia (J09-J18)	7	2.9	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.9	*
9	Septicemia (A40-A41)	4	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.7	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.7	*
13	Meningitis (G00,G03)	2	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
15	Hernia (K40-K46)	1	0.4	*
...	All other causes (Residual)	56	23.1	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	782	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	30.8	9.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	25.8	7.8
3	Malignant neoplasms (C00-C97)	102	13.0	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	7.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.2	*
7	Cerebrovascular diseases (I60-I69)	8	1.0	*
8	Septicemia (A40-A41)	7	0.9	*
9	Influenza and pneumonia (J09-J18)	5	0.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
12	Legal intervention (Y35,Y89.0)	3	0.4	*
15	Tuberculosis (A16-A19)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
...	All other causes (Residual)	89	11.4	3.4
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,290	100.0	53.8
1	Malignant neoplasms (C00-C97)	837	25.4	13.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	549	16.7	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	14.2	7.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	393	11.9	6.4
5	Cerebrovascular diseases (I60-I69)	131	4.0	2.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	3.7	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	75	2.3	1.2
8	Diabetes mellitus (E10-E14)	48	1.5	0.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.2	0.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.0	0.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.8	0.4
12	Influenza and pneumonia (J09-J18)	25	0.8	0.4
13	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.4
14	Septicemia (A40-A41)	22	0.7	0.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.6	0.3
...	All other causes (Residual)	480	14.6	7.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	11,857	100.0	268.6
1	Malignant neoplasms (C00-C97)	4,749	40.1	107.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,375	20.0	53.8
3	Cerebrovascular diseases (I60-I69)	734	6.2	16.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	563	4.7	12.8
5	Diabetes mellitus (E10-E14)	482	4.1	10.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	316	2.7	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	220	1.9	5.0
8	Influenza and pneumonia (J09-J18)	177	1.5	4.0
9	Chronic lower respiratory diseases (J40-J47)	169	1.4	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	1.4	3.8
11	Viral hepatitis (B15-B19)	132	1.1	3.0
12	Septicemia (A40-A41)	120	1.0	2.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	0.8	2.3
14	Aortic aneurysm and dissection (I71)	79	0.7	1.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	0.6	1.7
...	All other causes (Residual)	1,397	11.8	31.6
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	42,636	100.0	2,290.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,474	24.6	562.6
2	Malignant neoplasms (C00-C97)	9,951	23.3	534.5
3	Cerebrovascular diseases (I60-I69)	3,322	7.8	178.4
4	Influenza and pneumonia (J09-J18)	1,803	4.2	96.9
5	Diabetes mellitus (E10-E14)	1,777	4.2	95.5
6	Chronic lower respiratory diseases (J40-J47)	1,640	3.8	88.1
7	Alzheimer's disease (G30)	1,415	3.3	76.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,047	2.5	56.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	948	2.2	50.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	864	2.0	46.4
11	Septicemia (A40-A41)	569	1.3	30.6
12	Parkinson's disease (G20-G21)	511	1.2	27.4
13	Pneumonitis due to solids and liquids (J69)	396	0.9	21.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	279	0.7	15.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	264	0.6	14.2
...	All other causes (Residual)	7,376	17.3	396.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	30,980	100.0	347.4
1	Malignant neoplasms (C00-C97)	8,071	26.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,307	23.6	81.9
3	Cerebrovascular diseases (I60-I69)	1,897	6.1	21.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,561	5.0	17.5
5	Diabetes mellitus (E10-E14)	1,228	4.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,117	3.6	12.5
7	Influenza and pneumonia (J09-J18)	1,011	3.3	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	820	2.6	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	1.9	6.4
10	Alzheimer's disease (G30)	447	1.4	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	433	1.4	4.9
12	Septicemia (A40-A41)	371	1.2	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	359	1.2	4.0
14	Parkinson's disease (G20-G21)	299	1.0	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	277	0.9	3.1
...	All other causes (Residual)	5,208	16.8	58.4
Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	13.5	*
3	Malignant neoplasms (C00-C97)	6	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
5	Septicemia (A40-A41)	3	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
5	Chronic lower respiratory diseases (J40-J47)	3	3.1	*
8	Meningitis (G00,G03)	2	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
9	Anemias (D50-D64)	1	1.0	*
9	Hernia (K40-K46)	1	1.0	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	30	31.3	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	136	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	24.3	2.7
2	Malignant neoplasms (C00-C97)	26	19.1	2.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	5.9	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.4	*
6	Chronic lower respiratory diseases (J40-J47)	5	3.7	*
7	Influenza and pneumonia (J09-J18)	4	2.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.9	*
9	Septicemia (A40-A41)	3	2.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.2	*
9	Cerebrovascular diseases (I60-I69)	3	2.2	*
12	Meningitis (G00,G03)	2	1.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.5	*
14	Hernia (K40-K46)	1	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
...	All other causes (Residual)	27	19.9	2.2
Asian or Pacific Islander, male, 15-24 years				
...	All causes	555	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	32.1	13.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	28.1	11.9
3	Malignant neoplasms (C00-C97)	57	10.3	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	8.3	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.8	*
7	Influenza and pneumonia (J09-J18)	4	0.7	*
8	Septicemia (A40-A41)	3	0.5	*
8	Cerebrovascular diseases (I60-I69)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
11	Tuberculosis (A16-A19)	2	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	57	10.3	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-44 years				
...	All causes	2,058	100.0	70.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	20.1	14.2
2	Malignant neoplasms (C00-C97)	383	18.6	13.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	16.9	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	303	14.7	10.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	3.8	2.7
6	Cerebrovascular diseases (I60-I69)	64	3.1	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	2.5	1.8
8	Diabetes mellitus (E10-E14)	31	1.5	1.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.3	0.9
10	Influenza and pneumonia (J09-J18)	18	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Viral hepatitis (B15-B19)	15	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.6	*
15	Aortic aneurysm and dissection (I71)	9	0.4	*
...	All other causes (Residual)	275	13.4	9.5
Asian or Pacific Islander, male, 45-64 years				
...	All causes	7,125	100.0	349.1
1	Malignant neoplasms (C00-C97)	2,430	34.1	119.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,767	24.8	86.6
3	Cerebrovascular diseases (I60-I69)	427	6.0	20.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	363	5.1	17.8
5	Diabetes mellitus (E10-E14)	308	4.3	15.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	223	3.1	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	164	2.3	8.0
8	Chronic lower respiratory diseases (J40-J47)	108	1.5	5.3
9	Influenza and pneumonia (J09-J18)	99	1.4	4.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.4	4.8
11	Viral hepatitis (B15-B19)	80	1.1	3.9
12	Septicemia (A40-A41)	78	1.1	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	0.9	3.1
14	Aortic aneurysm and dissection (I71)	57	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	0.7	2.5
...	All other causes (Residual)	807	11.3	39.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65 years and over				
...	All causes	20,514	100.0	2,542.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,195	25.3	643.8
2	Malignant neoplasms (C00-C97)	5,169	25.2	640.5
3	Cerebrovascular diseases (I60-I69)	1,398	6.8	173.2
4	Chronic lower respiratory diseases (J40-J47)	982	4.8	121.7
5	Diabetes mellitus (E10-E14)	888	4.3	110.0
6	Influenza and pneumonia (J09-J18)	885	4.3	109.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	542	2.6	67.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	2.2	56.9
9	Alzheimer's disease (G30)	442	2.2	54.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	360	1.8	44.6
11	Parkinson's disease (G20-G21)	285	1.4	35.3
12	Septicemia (A40-A41)	274	1.3	34.0
13	Pneumonitis due to solids and liquids (J69)	221	1.1	27.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	144	0.7	17.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.7	16.9
...	All other causes (Residual)	3,134	15.3	388.4
Asian or Pacific Islander, female, all ages				
...	All causes	28,889	100.0	297.4
1	Malignant neoplasms (C00-C97)	7,632	26.4	78.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,004	20.8	61.8
3	Cerebrovascular diseases (I60-I69)	2,308	8.0	23.8
4	Diabetes mellitus (E10-E14)	1,081	3.7	11.1
5	Influenza and pneumonia (J09-J18)	1,013	3.5	10.4
6	Alzheimer's disease (G30)	981	3.4	10.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	947	3.3	9.7
8	Chronic lower respiratory diseases (J40-J47)	731	2.5	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	1.9	5.7
11	Septicemia (A40-A41)	358	1.2	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	1.0	3.1
13	Parkinson's disease (G20-G21)	229	0.8	2.4
14	Certain conditions originating in the perinatal period (P00-P96)	210	0.7	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	200	0.7	2.1
...	All other causes (Residual)	5,771	20.0	59.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	88	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	21.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	18.2	*
3	Malignant neoplasms (C00-C97)	9	10.2	*
4	Influenza and pneumonia (J09-J18)	6	6.8	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.3	*
5	Meningitis (G00,G03)	2	2.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
5	Cerebrovascular diseases (I60-I69)	2	2.3	*
9	Shigellosis and amebiasis (A03,A06)	1	1.1	*
9	Septicemia (A40-A41)	1	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	25	28.4	5.2
Asian or Pacific Islander, female, 5-14 years				
...	All causes	106	100.0	8.8
1	Malignant neoplasms (C00-C97)	20	18.9	1.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	17.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	7.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	6.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	5.7	*
7	Influenza and pneumonia (J09-J18)	3	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	29	27.4	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	227	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	27.8	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	20.3	3.6
3	Malignant neoplasms (C00-C97)	45	19.8	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.1	*
7	Cerebrovascular diseases (I60-I69)	5	2.2	*
8	Septicemia (A40-A41)	4	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Influenza and pneumonia (J09-J18)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	29	12.8	2.3
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,232	100.0	38.4
1	Malignant neoplasms (C00-C97)	454	36.9	14.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	11.0	4.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	9.7	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	7.3	2.8
5	Cerebrovascular diseases (I60-I69)	67	5.4	2.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	3.4	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	39	3.2	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	24	1.9	0.7
9	Diabetes mellitus (E10-E14)	17	1.4	*
10	Septicemia (A40-A41)	15	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.6	*
14	Influenza and pneumonia (J09-J18)	7	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.6	*
...	All other causes (Residual)	166	13.5	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-64 years				
...	All causes	4,732	100.0	199.4
1	Malignant neoplasms (C00-C97)	2,319	49.0	97.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	608	12.8	25.6
3	Cerebrovascular diseases (I60-I69)	307	6.5	12.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	4.2	8.4
5	Diabetes mellitus (E10-E14)	174	3.7	7.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	2.0	3.9
7	Influenza and pneumonia (J09-J18)	78	1.6	3.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.5	2.9
9	Chronic lower respiratory diseases (J40-J47)	61	1.3	2.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.2	2.4
11	Viral hepatitis (B15-B19)	52	1.1	2.2
12	Septicemia (A40-A41)	42	0.9	1.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.8	1.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	0.8	1.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.5	1.1
...	All other causes (Residual)	575	12.2	24.2
Asian or Pacific Islander, female, 65 years and over				
...	All causes	22,122	100.0	2,097.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,279	23.9	500.5
2	Malignant neoplasms (C00-C97)	4,782	21.6	453.4
3	Cerebrovascular diseases (I60-I69)	1,924	8.7	182.4
4	Alzheimer's disease (G30)	973	4.4	92.3
5	Influenza and pneumonia (J09-J18)	918	4.1	87.0
6	Diabetes mellitus (E10-E14)	889	4.0	84.3
7	Chronic lower respiratory diseases (J40-J47)	658	3.0	62.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	2.3	47.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	504	2.3	47.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	2.2	46.4
11	Septicemia (A40-A41)	295	1.3	28.0
12	Parkinson's disease (G20-G21)	226	1.0	21.4
13	Pneumonitis due to solids and liquids (J69)	175	0.8	16.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	143	0.6	13.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	120	0.5	11.4
...	All other causes (Residual)	4,242	19.2	402.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013