

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	156,419	100.0	295.0
1	Malignant neoplasms (C00-C97)	34,147	21.8	64.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,595	20.2	59.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,447	7.3	21.6
4	Cerebrovascular diseases (I60-I69)	7,767	5.0	14.6
5	Diabetes mellitus (E10-E14)	7,363	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,988	3.2	9.4
7	Chronic lower respiratory diseases (J40-J47)	4,477	2.9	8.4
8	Alzheimer's disease (G30)	3,772	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,049	1.9	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,964	1.9	5.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,837	1.8	5.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,740	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,429	1.6	4.6
14	Septicemia (A40-A41)	2,304	1.5	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,859	1.2	3.5
...	All other causes (Residual)	32,681	20.9	61.6
Hispanic, both sexes, 1-4 years				
...	All causes	899	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	29.9	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	14.0	3.1
3	Malignant neoplasms (C00-C97)	99	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
6	Influenza and pneumonia (J09-J18)	22	2.4	0.5
7	Septicemia (A40-A41)	14	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.3	*
10	Cerebrovascular diseases (I60-I69)	11	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.6	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
...	All other causes (Residual)	216	24.0	5.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	510	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	28.8	2.9
2	Malignant neoplasms (C00-C97)	116	22.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	4.1	0.4
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	3.1	*
6	Influenza and pneumonia (J09-J18)	14	2.7	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.2	*
9	Septicemia (A40-A41)	8	1.6	*
9	Cerebrovascular diseases (I60-I69)	8	1.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
12	Meningitis (G00,G03)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	107	21.0	2.1
Hispanic, both sexes, 10-14 years				
...	All causes	573	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	27.9	3.4
2	Malignant neoplasms (C00-C97)	103	18.0	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	7.9	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.0	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.5	0.4
7	Cerebrovascular diseases (I60-I69)	16	2.8	*
8	Septicemia (A40-A41)	8	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
10	Influenza and pneumonia (J09-J18)	4	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Anemias (D50-D64)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	117	20.4	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,780	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	38.4	14.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	20.1	7.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	263	14.8	5.7
4	Malignant neoplasms (C00-C97)	153	8.6	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.6	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.0	0.8
7	Cerebrovascular diseases (I60-I69)	10	0.6	*
8	Influenza and pneumonia (J09-J18)	9	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.4	*
9	Legal intervention (Y35,Y89.0)	8	0.4	*
11	Diabetes mellitus (E10-E14)	7	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
13	Septicemia (A40-A41)	5	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
15	Anemias (D50-D64)	4	0.2	*
...	All other causes (Residual)	181	10.2	4.0
Hispanic, both sexes, 20-24 years				
...	All causes	2,993	100.0	65.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,188	39.7	25.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	558	18.6	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	412	13.8	9.0
4	Malignant neoplasms (C00-C97)	217	7.3	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.1	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.1	0.7
7	Cerebrovascular diseases (I60-I69)	26	0.9	0.6
8	Legal intervention (Y35,Y89.0)	22	0.7	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	18	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.6	*
10	Influenza and pneumonia (J09-J18)	17	0.6	*
12	Diabetes mellitus (E10-E14)	16	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
14	Septicemia (A40-A41)	10	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
...	All other causes (Residual)	343	11.5	7.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,960	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,112	37.6	25.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	455	15.4	10.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	316	10.7	7.2
4	Malignant neoplasms (C00-C97)	272	9.2	6.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	4.4	3.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	41	1.4	0.9
7	Diabetes mellitus (E10-E14)	39	1.3	0.9
8	Cerebrovascular diseases (I60-I69)	32	1.1	0.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.1	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.9	0.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.9	0.6
12	Septicemia (A40-A41)	21	0.7	0.5
13	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Influenza and pneumonia (J09-J18)	17	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
...	All other causes (Residual)	401	13.5	9.2
Hispanic, both sexes, 30-34 years				
...	All causes	3,256	100.0	75.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,048	32.2	24.4
2	Malignant neoplasms (C00-C97)	407	12.5	9.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	326	10.0	7.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	316	9.7	7.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	6.9	5.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	89	2.7	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	66	2.0	1.5
7	Cerebrovascular diseases (I60-I69)	66	2.0	1.5
9	Diabetes mellitus (E10-E14)	47	1.4	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.9	0.7
11	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.9	0.7
12	Septicemia (A40-A41)	26	0.8	0.6
12	Influenza and pneumonia (J09-J18)	26	0.8	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.7	0.5
15	Legal intervention (Y35,Y89.0)	17	0.5	*
...	All other causes (Residual)	517	15.9	12.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,519	100.0	88.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	924	26.3	23.2
2	Malignant neoplasms (C00-C97)	555	15.8	13.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	338	9.6	8.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	274	7.8	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	240	6.8	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	164	4.7	4.1
7	Cerebrovascular diseases (I60-I69)	109	3.1	2.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	79	2.2	2.0
9	Diabetes mellitus (E10-E14)	77	2.2	1.9
10	Septicemia (A40-A41)	42	1.2	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.1	1.0
12	Influenza and pneumonia (J09-J18)	38	1.1	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	34	1.0	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.7	0.7
15	Chronic lower respiratory diseases (J40-J47)	22	0.6	0.6
...	All other causes (Residual)	558	15.9	14.0
Hispanic, both sexes, 40-44 years				
...	All causes	4,819	100.0	131.4
1	Malignant neoplasms (C00-C97)	1,035	21.5	28.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	888	18.4	24.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	602	12.5	16.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	309	6.4	8.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	273	5.7	7.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	4.4	5.8
7	Cerebrovascular diseases (I60-I69)	179	3.7	4.9
8	Diabetes mellitus (E10-E14)	155	3.2	4.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	105	2.2	2.9
10	Influenza and pneumonia (J09-J18)	65	1.3	1.8
11	Septicemia (A40-A41)	51	1.1	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.0	1.3
13	Viral hepatitis (B15-B19)	48	1.0	1.3
14	Aortic aneurysm and dissection (I71)	25	0.5	0.7
14	Chronic lower respiratory diseases (J40-J47)	25	0.5	0.7
...	All other causes (Residual)	797	16.5	21.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,767	100.0	211.0
1	Malignant neoplasms (C00-C97)	1,650	24.4	51.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	975	14.4	30.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	903	13.3	28.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	573	8.5	17.9
5	Cerebrovascular diseases (I60-I69)	296	4.4	9.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	3.8	8.0
7	Diabetes mellitus (E10-E14)	238	3.5	7.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	179	2.6	5.6
9	Viral hepatitis (B15-B19)	143	2.1	4.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	2.1	4.5
11	Septicemia (A40-A41)	106	1.6	3.3
12	Influenza and pneumonia (J09-J18)	77	1.1	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	1.1	2.4
14	Chronic lower respiratory diseases (J40-J47)	47	0.7	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	0.6	1.3
...	All other causes (Residual)	1,063	15.7	33.2
Hispanic, both sexes, 50-54 years				
...	All causes	9,011	100.0	335.9
1	Malignant neoplasms (C00-C97)	2,495	27.7	93.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,498	16.6	55.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	842	9.3	31.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	789	8.8	29.4
5	Diabetes mellitus (E10-E14)	442	4.9	16.5
6	Cerebrovascular diseases (I60-I69)	362	4.0	13.5
7	Viral hepatitis (B15-B19)	231	2.6	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	2.2	7.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	157	1.7	5.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	1.6	5.4
11	Septicemia (A40-A41)	126	1.4	4.7
12	Chronic lower respiratory diseases (J40-J47)	113	1.3	4.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	1.1	3.8
14	Influenza and pneumonia (J09-J18)	86	1.0	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	0.7	2.5
...	All other causes (Residual)	1,355	15.0	50.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	10,806	100.0	517.4
1	Malignant neoplasms (C00-C97)	3,250	30.1	155.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,020	18.7	96.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	806	7.5	38.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	674	6.2	32.3
5	Diabetes mellitus (E10-E14)	593	5.5	28.4
6	Cerebrovascular diseases (I60-I69)	449	4.2	21.5
7	Viral hepatitis (B15-B19)	286	2.6	13.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	2.1	10.9
9	Chronic lower respiratory diseases (J40-J47)	178	1.6	8.5
10	Septicemia (A40-A41)	174	1.6	8.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	1.5	7.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	134	1.2	6.4
13	Influenza and pneumonia (J09-J18)	129	1.2	6.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	103	1.0	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	0.8	4.1
...	All other causes (Residual)	1,538	14.2	73.6
Hispanic, both sexes, 60-64 years				
...	All causes	11,814	100.0	764.0
1	Malignant neoplasms (C00-C97)	3,795	32.1	245.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,439	20.6	157.7
3	Diabetes mellitus (E10-E14)	771	6.5	49.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	681	5.8	44.0
5	Cerebrovascular diseases (I60-I69)	508	4.3	32.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	449	3.8	29.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	2.1	15.8
8	Chronic lower respiratory diseases (J40-J47)	229	1.9	14.8
9	Viral hepatitis (B15-B19)	213	1.8	13.8
10	Septicemia (A40-A41)	208	1.8	13.5
11	Influenza and pneumonia (J09-J18)	187	1.6	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.0	7.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	0.9	6.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	73	0.6	4.7
15	Pneumonitis due to solids and liquids (J69)	48	0.4	3.1
...	All other causes (Residual)	1,753	14.8	113.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	12,022	100.0	1,092.8
1	Malignant neoplasms (C00-C97)	3,936	32.7	357.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,537	21.1	230.6
3	Diabetes mellitus (E10-E14)	904	7.5	82.2
4	Cerebrovascular diseases (I60-I69)	565	4.7	51.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	498	4.1	45.3
6	Chronic lower respiratory diseases (J40-J47)	320	2.7	29.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	2.7	29.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	2.3	25.3
9	Septicemia (A40-A41)	220	1.8	20.0
10	Influenza and pneumonia (J09-J18)	196	1.6	17.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.1	12.5
12	Viral hepatitis (B15-B19)	97	0.8	8.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.6	6.7
14	Alzheimer's disease (G30)	69	0.6	6.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	0.6	6.1
...	All other causes (Residual)	1,804	15.0	164.0
Hispanic, both sexes, 70-74 years				
...	All causes	12,846	100.0	1,671.9
1	Malignant neoplasms (C00-C97)	3,993	31.1	519.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,852	22.2	371.2
3	Diabetes mellitus (E10-E14)	833	6.5	108.4
4	Cerebrovascular diseases (I60-I69)	679	5.3	88.4
5	Chronic lower respiratory diseases (J40-J47)	466	3.6	60.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	382	3.0	49.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	2.6	42.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	2.2	36.2
9	Influenza and pneumonia (J09-J18)	260	2.0	33.8
10	Septicemia (A40-A41)	229	1.8	29.8
11	Alzheimer's disease (G30)	160	1.2	20.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	1.2	20.2
13	Parkinson's disease (G20-G21)	93	0.7	12.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.6	9.8
15	Viral hepatitis (B15-B19)	64	0.5	8.3
...	All other causes (Residual)	1,998	15.6	260.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	15,347	100.0	2,759.3
1	Malignant neoplasms (C00-C97)	4,081	26.6	733.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,501	22.8	629.4
3	Diabetes mellitus (E10-E14)	981	6.4	176.4
4	Cerebrovascular diseases (I60-I69)	934	6.1	167.9
5	Chronic lower respiratory diseases (J40-J47)	649	4.2	116.7
6	Alzheimer's disease (G30)	404	2.6	72.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	396	2.6	71.2
8	Influenza and pneumonia (J09-J18)	343	2.2	61.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	2.0	56.5
10	Septicemia (A40-A41)	294	1.9	52.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	287	1.9	51.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	211	1.4	37.9
13	Parkinson's disease (G20-G21)	210	1.4	37.8
14	Pneumonitis due to solids and liquids (J69)	91	0.6	16.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.6	15.6
...	All other causes (Residual)	2,564	16.7	461.0
Hispanic, both sexes, 80-84 years				
...	All causes	18,267	100.0	4,682.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,627	25.3	1,186.1
2	Malignant neoplasms (C00-C97)	3,770	20.6	966.4
3	Cerebrovascular diseases (I60-I69)	1,244	6.8	318.9
4	Diabetes mellitus (E10-E14)	1,004	5.5	257.4
5	Chronic lower respiratory diseases (J40-J47)	838	4.6	214.8
6	Alzheimer's disease (G30)	781	4.3	200.2
7	Influenza and pneumonia (J09-J18)	490	2.7	125.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	2.4	113.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	387	2.1	99.2
10	Parkinson's disease (G20-G21)	316	1.7	81.0
11	Septicemia (A40-A41)	285	1.6	73.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	280	1.5	71.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	215	1.2	55.1
14	Pneumonitis due to solids and liquids (J69)	135	0.7	34.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	118	0.6	30.2
...	All other causes (Residual)	3,336	18.3	855.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	17,380	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,607	26.5	---
2	Malignant neoplasms (C00-C97)	2,672	15.4	---
3	Cerebrovascular diseases (I60-I69)	1,219	7.0	---
4	Alzheimer's disease (G30)	1,040	6.0	---
5	Chronic lower respiratory diseases (J40-J47)	831	4.8	---
6	Diabetes mellitus (E10-E14)	759	4.4	---
7	Influenza and pneumonia (J09-J18)	495	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	405	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	345	2.0	---
11	Parkinson's disease (G20-G21)	259	1.5	---
12	Septicemia (A40-A41)	248	1.4	---
13	Pneumonitis due to solids and liquids (J69)	175	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	106	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.6	---
...	All other causes (Residual)	3,750	21.6	---
Hispanic, both sexes, 90-94 years				
...	All causes	10,954	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,284	30.0	---
2	Malignant neoplasms (C00-C97)	1,158	10.6	---
3	Alzheimer's disease (G30)	842	7.7	---
4	Cerebrovascular diseases (I60-I69)	715	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	502	4.6	---
6	Influenza and pneumonia (J09-J18)	380	3.5	---
7	Diabetes mellitus (E10-E14)	351	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	205	1.9	---
11	Pneumonitis due to solids and liquids (J69)	144	1.3	---
12	Septicemia (A40-A41)	142	1.3	---
13	Parkinson's disease (G20-G21)	128	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	---
15	Atherosclerosis (I70)	43	0.4	---
...	All other causes (Residual)	2,538	23.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	4,015	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,323	33.0	---
2	Alzheimer's disease (G30)	335	8.3	---
3	Malignant neoplasms (C00-C97)	319	7.9	---
4	Cerebrovascular diseases (I60-I69)	261	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	144	3.6	---
6	Influenza and pneumonia (J09-J18)	125	3.1	---
7	Diabetes mellitus (E10-E14)	108	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.5	---
11	Pneumonitis due to solids and liquids (J69)	50	1.2	---
12	Septicemia (A40-A41)	45	1.1	---
13	Parkinson's disease (G20-G21)	29	0.7	---
14	Atherosclerosis (I70)	22	0.5	---
15	Anemias (D50-D64)	16	0.4	---
...	All other causes (Residual)	1,003	25.0	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,061	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	33.9	---
2	Alzheimer's disease (G30)	74	7.0	---
3	Malignant neoplasms (C00-C97)	56	5.3	---
3	Cerebrovascular diseases (I60-I69)	56	5.3	---
5	Influenza and pneumonia (J09-J18)	41	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	34	3.2	---
7	Diabetes mellitus (E10-E14)	32	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.6	---
10	Septicemia (A40-A41)	15	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	---
12	Pneumonitis due to solids and liquids (J69)	10	0.9	---
13	Atherosclerosis (I70)	9	0.8	---
14	Parkinson's disease (G20-G21)	7	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.3	---
15	Anemias (D50-D64)	3	0.3	---
15	Nutritional deficiencies (E40-E64)	3	0.3	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.3	---
...	All other causes (Residual)	294	27.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	85,238	100.0	316.5
1	Malignant neoplasms (C00-C97)	17,849	20.9	66.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,329	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,255	9.7	30.7
4	Diabetes mellitus (E10-E14)	3,763	4.4	14.0
5	Cerebrovascular diseases (I60-I69)	3,636	4.3	13.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,422	4.0	12.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,292	2.7	8.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,277	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,268	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,557	1.8	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,552	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,373	1.6	5.1
13	Alzheimer's disease (G30)	1,193	1.4	4.4
14	Septicemia (A40-A41)	1,175	1.4	4.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	952	1.1	3.5
...	All other causes (Residual)	16,345	19.2	60.7
Hispanic, male, 1-4 years				
...	All causes	494	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	31.2	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	15.0	3.5
3	Malignant neoplasms (C00-C97)	56	11.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.6	*
6	Influenza and pneumonia (J09-J18)	11	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
8	Septicemia (A40-A41)	7	1.4	*
8	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
14	Anemias (D50-D64)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	114	23.1	5.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	282	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	32.3	3.6
2	Malignant neoplasms (C00-C97)	64	22.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	4.3	*
5	Influenza and pneumonia (J09-J18)	9	3.2	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.8	*
6	Chronic lower respiratory diseases (J40-J47)	8	2.8	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.1	*
9	Septicemia (A40-A41)	5	1.8	*
9	Cerebrovascular diseases (I60-I69)	5	1.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
12	Anemias (D50-D64)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	50	17.7	2.0
Hispanic, male, 10-14 years				
...	All causes	326	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	31.9	4.3
2	Malignant neoplasms (C00-C97)	51	15.6	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	8.6	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	8.0	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	7.1	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.7	*
7	Cerebrovascular diseases (I60-I69)	9	2.8	*
8	Septicemia (A40-A41)	4	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.2	*
10	Diseases of appendix (K35-K38)	2	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
11	Anemias (D50-D64)	1	0.3	*
11	Diabetes mellitus (E10-E14)	1	0.3	*
11	Meningitis (G00,G03)	1	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
11	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	55	16.9	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,328	100.0	56.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	495	37.3	21.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	24.1	13.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	15.6	8.8
4	Malignant neoplasms (C00-C97)	96	7.2	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.5	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.6	0.9
7	Legal intervention (Y35,Y89.0)	8	0.6	*
8	Cerebrovascular diseases (I60-I69)	7	0.5	*
9	Influenza and pneumonia (J09-J18)	6	0.5	*
10	Septicemia (A40-A41)	2	0.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
10	Diabetes mellitus (E10-E14)	2	0.2	*
10	Aortic aneurysm and dissection (I71)	2	0.2	*
10	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.2	*
10	Pneumonitis due to solids and liquids (J69)	2	0.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	119	9.0	5.0
Hispanic, male, 20-24 years				
...	All causes	2,327	100.0	96.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	40.5	38.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	498	21.4	20.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	341	14.7	14.1
4	Malignant neoplasms (C00-C97)	132	5.7	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.8	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.9	0.9
6	Legal intervention (Y35,Y89.0)	21	0.9	0.9
8	Cerebrovascular diseases (I60-I69)	16	0.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
9	Influenza and pneumonia (J09-J18)	11	0.5	*
11	Diabetes mellitus (E10-E14)	10	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
14	Septicemia (A40-A41)	5	0.2	*
14	Anemias (D50-D64)	5	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	227	9.8	9.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,233	100.0	96.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	889	39.8	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	396	17.7	17.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	11.9	11.4
4	Malignant neoplasms (C00-C97)	156	7.0	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	4.7	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	37	1.7	1.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.0	0.9
8	Diabetes mellitus (E10-E14)	21	0.9	0.9
9	Cerebrovascular diseases (I60-I69)	20	0.9	0.9
10	Legal intervention (Y35,Y89.0)	18	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.7	*
12	Septicemia (A40-A41)	11	0.5	*
12	Influenza and pneumonia (J09-J18)	11	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.3	*
...	All other causes (Residual)	251	11.2	10.8
Hispanic, male, 30-34 years				
...	All causes	2,345	100.0	104.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	862	36.8	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	11.8	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	10.9	11.3
4	Malignant neoplasms (C00-C97)	190	8.1	8.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	7.4	7.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	73	3.1	3.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	53	2.3	2.4
8	Cerebrovascular diseases (I60-I69)	39	1.7	1.7
9	Diabetes mellitus (E10-E14)	31	1.3	1.4
10	Influenza and pneumonia (J09-J18)	19	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
12	Legal intervention (Y35,Y89.0)	16	0.7	*
13	Septicemia (A40-A41)	15	0.6	*
14	Aortic aneurysm and dissection (I71)	12	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
...	All other causes (Residual)	303	12.9	13.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,414	100.0	118.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	30.2	35.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	247	10.2	12.1
3	Malignant neoplasms (C00-C97)	242	10.0	11.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	9.3	10.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	8.2	9.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	125	5.2	6.1
7	Cerebrovascular diseases (I60-I69)	70	2.9	3.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	63	2.6	3.1
9	Diabetes mellitus (E10-E14)	48	2.0	2.3
10	Septicemia (A40-A41)	28	1.2	1.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.1	1.3
12	Influenza and pneumonia (J09-J18)	26	1.1	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
14	Aortic aneurysm and dissection (I71)	16	0.7	*
15	Viral hepatitis (B15-B19)	15	0.6	*
...	All other causes (Residual)	339	14.0	16.6
Hispanic, male, 40-44 years				
...	All causes	3,179	100.0	169.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	671	21.1	35.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	14.4	24.5
3	Malignant neoplasms (C00-C97)	437	13.7	23.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	235	7.4	12.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	223	7.0	11.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	5.5	9.3
7	Cerebrovascular diseases (I60-I69)	121	3.8	6.5
8	Diabetes mellitus (E10-E14)	106	3.3	5.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	73	2.3	3.9
10	Influenza and pneumonia (J09-J18)	44	1.4	2.3
11	Viral hepatitis (B15-B19)	33	1.0	1.8
12	Septicemia (A40-A41)	29	0.9	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.9	1.5
14	Aortic aneurysm and dissection (I71)	23	0.7	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.5	*
15	Legal intervention (Y35,Y89.0)	17	0.5	*
...	All other causes (Residual)	487	15.3	26.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,383	100.0	269.1
1	Malignant neoplasms (C00-C97)	740	16.9	45.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	740	16.9	45.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	689	15.7	42.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	442	10.1	27.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	4.3	11.7
6	Cerebrovascular diseases (I60-I69)	179	4.1	11.0
7	Diabetes mellitus (E10-E14)	162	3.7	9.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	134	3.1	8.2
9	Viral hepatitis (B15-B19)	118	2.7	7.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	2.5	6.8
11	Septicemia (A40-A41)	59	1.3	3.6
12	Influenza and pneumonia (J09-J18)	50	1.1	3.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.1	2.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.7	1.8
15	Chronic lower respiratory diseases (J40-J47)	24	0.5	1.5
...	All other causes (Residual)	669	15.3	41.1
Hispanic, male, 50-54 years				
...	All causes	5,786	100.0	432.2
1	Malignant neoplasms (C00-C97)	1,265	21.9	94.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,092	18.9	81.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	603	10.4	45.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	603	10.4	45.0
5	Diabetes mellitus (E10-E14)	282	4.9	21.1
6	Cerebrovascular diseases (I60-I69)	221	3.8	16.5
7	Viral hepatitis (B15-B19)	172	3.0	12.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	2.8	12.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	122	2.1	9.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	1.6	6.9
11	Septicemia (A40-A41)	86	1.5	6.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	1.3	5.8
13	Chronic lower respiratory diseases (J40-J47)	58	1.0	4.3
14	Influenza and pneumonia (J09-J18)	54	0.9	4.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	0.9	3.7
...	All other causes (Residual)	847	14.6	63.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	6,980	100.0	688.0
1	Malignant neoplasms (C00-C97)	1,790	25.6	176.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,460	20.9	143.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	622	8.9	61.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	7.3	50.1
5	Diabetes mellitus (E10-E14)	362	5.2	35.7
6	Cerebrovascular diseases (I60-I69)	275	3.9	27.1
7	Viral hepatitis (B15-B19)	207	3.0	20.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	2.1	14.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	1.8	12.5
10	Septicemia (A40-A41)	111	1.6	10.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	110	1.6	10.8
12	Chronic lower respiratory diseases (J40-J47)	103	1.5	10.2
13	Influenza and pneumonia (J09-J18)	73	1.0	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	0.9	6.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	0.9	6.1
...	All other causes (Residual)	962	13.8	94.8
Hispanic, male, 60-64 years				
...	All causes	7,278	100.0	994.1
1	Malignant neoplasms (C00-C97)	2,137	29.4	291.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,641	22.5	224.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	485	6.7	66.2
4	Diabetes mellitus (E10-E14)	434	6.0	59.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	4.8	47.5
6	Cerebrovascular diseases (I60-I69)	307	4.2	41.9
7	Viral hepatitis (B15-B19)	164	2.3	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	1.9	18.6
9	Chronic lower respiratory diseases (J40-J47)	135	1.9	18.4
10	Influenza and pneumonia (J09-J18)	119	1.6	16.3
11	Septicemia (A40-A41)	115	1.6	15.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	1.2	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.0	10.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	55	0.8	7.5
15	Pneumonitis due to solids and liquids (J69)	35	0.5	4.8
...	All other causes (Residual)	1,006	13.8	137.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	6,968	100.0	1,386.0
1	Malignant neoplasms (C00-C97)	2,177	31.2	433.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,664	23.9	331.0
3	Diabetes mellitus (E10-E14)	487	7.0	96.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	326	4.7	64.8
5	Cerebrovascular diseases (I60-I69)	287	4.1	57.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	3.1	43.4
7	Chronic lower respiratory diseases (J40-J47)	184	2.6	36.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.4	33.0
9	Influenza and pneumonia (J09-J18)	113	1.6	22.5
10	Septicemia (A40-A41)	105	1.5	20.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.0	13.9
12	Viral hepatitis (B15-B19)	56	0.8	11.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	0.7	10.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	8.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.6	7.8
...	All other causes (Residual)	983	14.1	195.5
Hispanic, male, 70-74 years				
...	All causes	7,134	100.0	2,116.2
1	Malignant neoplasms (C00-C97)	2,226	31.2	660.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,747	24.5	518.2
3	Diabetes mellitus (E10-E14)	435	6.1	129.0
4	Cerebrovascular diseases (I60-I69)	347	4.9	102.9
5	Chronic lower respiratory diseases (J40-J47)	276	3.9	81.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	197	2.8	58.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	2.5	52.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	2.3	48.4
9	Influenza and pneumonia (J09-J18)	135	1.9	40.0
10	Septicemia (A40-A41)	130	1.8	38.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.1	22.2
12	Alzheimer's disease (G30)	66	0.9	19.6
13	Parkinson's disease (G20-G21)	54	0.8	16.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.6	12.5
15	Aortic aneurysm and dissection (I71)	37	0.5	11.0
15	Pneumonitis due to solids and liquids (J69)	37	0.5	11.0
...	All other causes (Residual)	990	13.9	293.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,918	100.0	3,389.3
1	Malignant neoplasms (C00-C97)	2,222	28.1	951.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,886	23.8	807.3
3	Diabetes mellitus (E10-E14)	483	6.1	206.7
4	Cerebrovascular diseases (I60-I69)	447	5.6	191.3
5	Chronic lower respiratory diseases (J40-J47)	360	4.5	154.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.6	89.0
7	Influenza and pneumonia (J09-J18)	184	2.3	78.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	2.2	75.8
9	Alzheimer's disease (G30)	164	2.1	70.2
10	Septicemia (A40-A41)	147	1.9	62.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	137	1.7	58.6
12	Parkinson's disease (G20-G21)	112	1.4	47.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	1.1	39.0
14	Pneumonitis due to solids and liquids (J69)	57	0.7	24.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.5	17.1
...	All other causes (Residual)	1,203	15.2	514.9
Hispanic, male, 80-84 years				
...	All causes	8,634	100.0	5,543.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,277	26.4	1,461.9
2	Malignant neoplasms (C00-C97)	1,937	22.4	1,243.6
3	Cerebrovascular diseases (I60-I69)	519	6.0	333.2
4	Diabetes mellitus (E10-E14)	438	5.1	281.2
5	Chronic lower respiratory diseases (J40-J47)	419	4.9	269.0
6	Alzheimer's disease (G30)	276	3.2	177.2
7	Influenza and pneumonia (J09-J18)	245	2.8	157.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.5	136.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	2.5	136.1
10	Parkinson's disease (G20-G21)	183	2.1	117.5
11	Septicemia (A40-A41)	130	1.5	83.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	1.4	75.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.1	59.7
14	Pneumonitis due to solids and liquids (J69)	64	0.7	41.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.7	37.2
...	All other causes (Residual)	1,453	16.8	932.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	7,316	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,002	27.4	---
2	Malignant neoplasms (C00-C97)	1,278	17.5	---
3	Cerebrovascular diseases (I60-I69)	452	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	376	5.1	---
5	Alzheimer's disease (G30)	336	4.6	---
6	Diabetes mellitus (E10-E14)	298	4.1	---
7	Influenza and pneumonia (J09-J18)	225	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	2.4	---
10	Parkinson's disease (G20-G21)	137	1.9	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	1.9	---
12	Septicemia (A40-A41)	97	1.3	---
13	Pneumonitis due to solids and liquids (J69)	94	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	39	0.5	---
...	All other causes (Residual)	1,427	19.5	---
Hispanic, male, 90-94 years				
...	All causes	3,874	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,191	30.7	---
2	Malignant neoplasms (C00-C97)	511	13.2	---
3	Alzheimer's disease (G30)	215	5.5	---
4	Cerebrovascular diseases (I60-I69)	213	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	203	5.2	---
6	Influenza and pneumonia (J09-J18)	154	4.0	---
7	Diabetes mellitus (E10-E14)	128	3.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	2.0	---
11	Pneumonitis due to solids and liquids (J69)	72	1.9	---
12	Parkinson's disease (G20-G21)	62	1.6	---
13	Septicemia (A40-A41)	57	1.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	---
15	Atherosclerosis (I70)	17	0.4	---
...	All other causes (Residual)	773	20.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,140	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	32.9	---
2	Malignant neoplasms (C00-C97)	116	10.2	---
3	Cerebrovascular diseases (I60-I69)	78	6.8	---
4	Alzheimer's disease (G30)	67	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	46	4.0	---
6	Influenza and pneumonia (J09-J18)	41	3.6	---
7	Diabetes mellitus (E10-E14)	27	2.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.6	---
11	Pneumonitis due to solids and liquids (J69)	15	1.3	---
12	Septicemia (A40-A41)	14	1.2	---
13	Parkinson's disease (G20-G21)	9	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	---
15	Anemias (D50-D64)	4	0.4	---
15	Aortic aneurysm and dissection (I71)	4	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.4	---
...	All other causes (Residual)	262	23.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 100 years and over				
...	All causes	230	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	40.0	---
2	Malignant neoplasms (C00-C97)	17	7.4	---
3	Alzheimer's disease (G30)	11	4.8	---
3	Chronic lower respiratory diseases (J40-J47)	11	4.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	4.3	---
5	Influenza and pneumonia (J09-J18)	10	4.3	---
7	Pneumonitis due to solids and liquids (J69)	8	3.5	---
8	Diabetes mellitus (E10-E14)	7	3.0	---
8	Cerebrovascular diseases (I60-I69)	7	3.0	---
10	Septicemia (A40-A41)	3	1.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.3	---
12	Parkinson's disease (G20-G21)	2	0.9	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
14	Atherosclerosis (I70)	1	0.4	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	---
14	Hyperplasia of prostate (N40)	1	0.4	---
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	---
...	All other causes (Residual)	40	17.4	---
Hispanic, female, all ages				
...	All causes	71,181	100.0	272.7
1	Malignant neoplasms (C00-C97)	16,298	22.9	62.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,266	20.0	54.7
3	Cerebrovascular diseases (I60-I69)	4,131	5.8	15.8
4	Diabetes mellitus (E10-E14)	3,600	5.1	13.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,192	4.5	12.2
6	Alzheimer's disease (G30)	2,579	3.6	9.9
7	Chronic lower respiratory diseases (J40-J47)	2,209	3.1	8.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,566	2.2	6.0
9	Influenza and pneumonia (J09-J18)	1,492	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,412	2.0	5.4
11	Septicemia (A40-A41)	1,129	1.6	4.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,056	1.5	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,017	1.4	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	838	1.2	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	545	0.8	2.1
...	All other causes (Residual)	15,851	22.3	60.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	405	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	28.4	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	12.8	2.6
3	Malignant neoplasms (C00-C97)	43	10.6	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.7	*
6	Influenza and pneumonia (J09-J18)	11	2.7	*
8	Septicemia (A40-A41)	7	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
15	Anemias (D50-D64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	98	24.2	4.8
Hispanic, female, 5-9 years				
...	All causes	228	100.0	9.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	24.6	2.3
2	Malignant neoplasms (C00-C97)	52	22.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	9.2	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	3.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	3.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.2	*
6	Influenza and pneumonia (J09-J18)	5	2.2	*
8	Septicemia (A40-A41)	3	1.3	*
8	Meningitis (G00,G03)	3	1.3	*
8	Cerebrovascular diseases (I60-I69)	3	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
14	Diabetes mellitus (E10-E14)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	52	22.8	2.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 10-14 years				
...	All causes	247	100.0	10.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	22.7	2.4
2	Malignant neoplasms (C00-C97)	52	21.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	6.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	5.7	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.2	*
7	Cerebrovascular diseases (I60-I69)	7	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	6	2.4	*
9	Septicemia (A40-A41)	4	1.6	*
9	Influenza and pneumonia (J09-J18)	4	1.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	1.2	*
12	Anemias (D50-D64)	2	0.8	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Meningitis (G00,G03)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	58	23.5	2.5
Hispanic, female, 15-19 years				
...	All causes	452	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	41.6	8.5
2	Malignant neoplasms (C00-C97)	57	12.6	2.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	12.4	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	8.2	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	3.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.8	*
8	Diabetes mellitus (E10-E14)	5	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
10	Septicemia (A40-A41)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Anemias (D50-D64)	3	0.7	*
10	Cerebrovascular diseases (I60-I69)	3	0.7	*
10	Influenza and pneumonia (J09-J18)	3	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	49	10.8	2.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 20-24 years				
...	All causes	666	100.0	30.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	36.8	11.3
2	Malignant neoplasms (C00-C97)	85	12.8	3.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	10.7	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	9.0	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.2	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	18	2.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.8	*
8	Cerebrovascular diseases (I60-I69)	10	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
9	Diabetes mellitus (E10-E14)	6	0.9	*
9	Influenza and pneumonia (J09-J18)	6	0.9	*
12	Septicemia (A40-A41)	5	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.3	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	97	14.6	4.5
Hispanic, female, 25-29 years				
...	All causes	727	100.0	35.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	30.7	10.9
2	Malignant neoplasms (C00-C97)	116	16.0	5.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	8.1	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	7.0	2.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	28	3.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.6	1.3
7	Diabetes mellitus (E10-E14)	18	2.5	*
8	Cerebrovascular diseases (I60-I69)	12	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.7	*
10	Septicemia (A40-A41)	10	1.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
13	Influenza and pneumonia (J09-J18)	6	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (Residual)	135	18.6	6.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 30-34 years				
...	All causes	911	100.0	44.5
1	Malignant neoplasms (C00-C97)	217	23.8	10.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	20.4	9.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	6.7	3.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.7	2.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	5.5	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	28	3.1	1.4
7	Cerebrovascular diseases (I60-I69)	27	3.0	1.3
8	Diabetes mellitus (E10-E14)	16	1.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.4	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.4	*
12	Septicemia (A40-A41)	11	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.2	*
14	Influenza and pneumonia (J09-J18)	7	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
...	All other causes (Residual)	197	21.6	9.6
Hispanic, female, 35-39 years				
...	All causes	1,105	100.0	57.1
1	Malignant neoplasms (C00-C97)	313	28.3	16.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	17.6	10.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	8.2	4.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	4.5	2.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	3.9	2.2
6	Cerebrovascular diseases (I60-I69)	39	3.5	2.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	39	3.5	2.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	34	3.1	1.8
9	Diabetes mellitus (E10-E14)	29	2.6	1.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.4	*
11	Septicemia (A40-A41)	14	1.3	*
12	Influenza and pneumonia (J09-J18)	12	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
...	All other causes (Residual)	200	18.1	10.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 40-44 years				
...	All causes	1,640	100.0	91.5
1	Malignant neoplasms (C00-C97)	598	36.5	33.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	13.2	12.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	8.7	8.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	74	4.5	4.1
5	Cerebrovascular diseases (I60-I69)	58	3.5	3.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	3.0	2.8
7	Diabetes mellitus (E10-E14)	49	3.0	2.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	2.3	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	32	2.0	1.8
10	Septicemia (A40-A41)	22	1.3	1.2
11	Influenza and pneumonia (J09-J18)	21	1.3	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.2	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	20	1.2	1.1
14	Viral hepatitis (B15-B19)	15	0.9	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
...	All other causes (Residual)	273	16.6	15.2
Hispanic, female, 45-49 years				
...	All causes	2,384	100.0	151.1
1	Malignant neoplasms (C00-C97)	910	38.2	57.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	9.9	14.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	9.0	13.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	131	5.5	8.3
5	Cerebrovascular diseases (I60-I69)	117	4.9	7.4
6	Diabetes mellitus (E10-E14)	76	3.2	4.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	2.7	4.1
8	Septicemia (A40-A41)	47	2.0	3.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	45	1.9	2.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	1.3	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.2	1.8
12	Influenza and pneumonia (J09-J18)	27	1.1	1.7
13	Viral hepatitis (B15-B19)	25	1.0	1.6
14	Chronic lower respiratory diseases (J40-J47)	23	1.0	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.6	*
...	All other causes (Residual)	380	15.9	24.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 50-54 years				
...	All causes	3,225	100.0	239.9
1	Malignant neoplasms (C00-C97)	1,230	38.1	91.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	406	12.6	30.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	7.4	17.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	186	5.8	13.8
5	Diabetes mellitus (E10-E14)	160	5.0	11.9
6	Cerebrovascular diseases (I60-I69)	141	4.4	10.5
7	Viral hepatitis (B15-B19)	59	1.8	4.4
8	Chronic lower respiratory diseases (J40-J47)	55	1.7	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.6	3.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.3	3.1
11	Septicemia (A40-A41)	40	1.2	3.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.1	2.6
13	Influenza and pneumonia (J09-J18)	32	1.0	2.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.5	*
...	All other causes (Residual)	508	15.8	37.8
Hispanic, female, 55-59 years				
...	All causes	3,826	100.0	356.2
1	Malignant neoplasms (C00-C97)	1,460	38.2	135.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	560	14.6	52.1
3	Diabetes mellitus (E10-E14)	231	6.0	21.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	184	4.8	17.1
5	Cerebrovascular diseases (I60-I69)	174	4.5	16.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	4.3	15.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	2.2	7.7
8	Viral hepatitis (B15-B19)	79	2.1	7.4
9	Chronic lower respiratory diseases (J40-J47)	75	2.0	7.0
10	Septicemia (A40-A41)	63	1.6	5.9
11	Influenza and pneumonia (J09-J18)	56	1.5	5.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.1	3.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	0.8	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.6	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.6	2.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	0.6	2.0
...	All other causes (Residual)	554	14.5	51.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 60-64 years				
...	All causes	4,536	100.0	557.1
1	Malignant neoplasms (C00-C97)	1,658	36.6	203.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	798	17.6	98.0
3	Diabetes mellitus (E10-E14)	337	7.4	41.4
4	Cerebrovascular diseases (I60-I69)	201	4.4	24.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	196	4.3	24.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	2.4	13.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	2.2	12.4
8	Chronic lower respiratory diseases (J40-J47)	94	2.1	11.5
9	Septicemia (A40-A41)	93	2.1	11.4
10	Influenza and pneumonia (J09-J18)	68	1.5	8.4
11	Viral hepatitis (B15-B19)	49	1.1	6.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	0.9	5.0
13	Alzheimer's disease (G30)	20	0.4	2.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	0.3	*
...	All other causes (Residual)	740	16.3	90.9
Hispanic, female, 65-69 years				
...	All causes	5,054	100.0	846.0
1	Malignant neoplasms (C00-C97)	1,759	34.8	294.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	873	17.3	146.1
3	Diabetes mellitus (E10-E14)	417	8.3	69.8
4	Cerebrovascular diseases (I60-I69)	278	5.5	46.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	172	3.4	28.8
6	Chronic lower respiratory diseases (J40-J47)	136	2.7	22.8
7	Septicemia (A40-A41)	115	2.3	19.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	2.2	18.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.0	17.1
10	Influenza and pneumonia (J09-J18)	83	1.6	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.3	11.2
12	Alzheimer's disease (G30)	43	0.9	7.2
13	Viral hepatitis (B15-B19)	41	0.8	6.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.7	5.5
15	Parkinson's disease (G20-G21)	18	0.4	*
...	All other causes (Residual)	805	15.9	134.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 70-74 years				
...	All causes	5,712	100.0	1,324.6
1	Malignant neoplasms (C00-C97)	1,767	30.9	409.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,105	19.3	256.2
3	Diabetes mellitus (E10-E14)	398	7.0	92.3
4	Cerebrovascular diseases (I60-I69)	332	5.8	77.0
5	Chronic lower respiratory diseases (J40-J47)	190	3.3	44.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	185	3.2	42.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.9	38.5
8	Influenza and pneumonia (J09-J18)	125	2.2	29.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	1.8	23.4
10	Septicemia (A40-A41)	99	1.7	23.0
11	Alzheimer's disease (G30)	94	1.6	21.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	1.4	18.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.7	9.0
13	Parkinson's disease (G20-G21)	39	0.7	9.0
15	Viral hepatitis (B15-B19)	37	0.6	8.6
...	All other causes (Residual)	955	16.7	221.5
Hispanic, female, 75-79 years				
...	All causes	7,429	100.0	2,303.0
1	Malignant neoplasms (C00-C97)	1,859	25.0	576.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,615	21.7	500.6
3	Diabetes mellitus (E10-E14)	498	6.7	154.4
4	Cerebrovascular diseases (I60-I69)	487	6.6	151.0
5	Chronic lower respiratory diseases (J40-J47)	289	3.9	89.6
6	Alzheimer's disease (G30)	240	3.2	74.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	2.5	58.3
8	Influenza and pneumonia (J09-J18)	159	2.1	49.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.0	46.5
10	Septicemia (A40-A41)	147	2.0	45.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	1.8	42.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.6	37.2
13	Parkinson's disease (G20-G21)	98	1.3	30.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	14.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	40	0.5	12.4
...	All other causes (Residual)	1,355	18.2	420.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 80-84 years				
...	All causes	9,633	100.0	4,110.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,350	24.4	1,002.7
2	Malignant neoplasms (C00-C97)	1,833	19.0	782.1
3	Cerebrovascular diseases (I60-I69)	725	7.5	309.4
4	Diabetes mellitus (E10-E14)	566	5.9	241.5
5	Alzheimer's disease (G30)	505	5.2	215.5
6	Chronic lower respiratory diseases (J40-J47)	419	4.3	178.8
7	Influenza and pneumonia (J09-J18)	245	2.5	104.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	2.4	97.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	1.8	74.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	1.7	69.1
11	Septicemia (A40-A41)	155	1.6	66.1
12	Parkinson's disease (G20-G21)	133	1.4	56.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	122	1.3	52.1
14	Pneumonitis due to solids and liquids (J69)	71	0.7	30.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.6	25.6
...	All other causes (Residual)	1,883	19.5	803.5
Hispanic, female, 85-89 years				
...	All causes	10,064	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,605	25.9	---
2	Malignant neoplasms (C00-C97)	1,394	13.9	---
3	Cerebrovascular diseases (I60-I69)	767	7.6	---
4	Alzheimer's disease (G30)	704	7.0	---
5	Diabetes mellitus (E10-E14)	461	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	455	4.5	---
7	Influenza and pneumonia (J09-J18)	270	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	209	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	1.9	---
11	Septicemia (A40-A41)	151	1.5	---
12	Parkinson's disease (G20-G21)	122	1.2	---
13	Pneumonitis due to solids and liquids (J69)	81	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.5	---
...	All other causes (Residual)	2,323	23.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 90-94 years				
...	All causes	7,080	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,093	29.6	---
2	Malignant neoplasms (C00-C97)	647	9.1	---
3	Alzheimer's disease (G30)	627	8.9	---
4	Cerebrovascular diseases (I60-I69)	502	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	299	4.2	---
6	Influenza and pneumonia (J09-J18)	226	3.2	---
7	Diabetes mellitus (E10-E14)	223	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	153	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	1.7	---
11	Septicemia (A40-A41)	85	1.2	---
12	Pneumonitis due to solids and liquids (J69)	72	1.0	---
13	Parkinson's disease (G20-G21)	66	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.5	---
15	Atherosclerosis (I70)	26	0.4	---
...	All other causes (Residual)	1,765	24.9	---
Hispanic, female, 95-99 years				
...	All causes	2,875	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	948	33.0	---
2	Alzheimer's disease (G30)	268	9.3	---
3	Malignant neoplasms (C00-C97)	203	7.1	---
4	Cerebrovascular diseases (I60-I69)	183	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	98	3.4	---
6	Influenza and pneumonia (J09-J18)	84	2.9	---
7	Diabetes mellitus (E10-E14)	81	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	1.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.5	---
11	Pneumonitis due to solids and liquids (J69)	35	1.2	---
12	Septicemia (A40-A41)	31	1.1	---
13	Atherosclerosis (I70)	21	0.7	---
14	Parkinson's disease (G20-G21)	20	0.7	---
15	Anemias (D50-D64)	12	0.4	---
...	All other causes (Residual)	724	25.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 100 years and over				
...	All causes	831	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	268	32.3	---
2	Alzheimer's disease (G30)	63	7.6	---
3	Cerebrovascular diseases (I60-I69)	49	5.9	---
4	Malignant neoplasms (C00-C97)	39	4.7	---
5	Influenza and pneumonia (J09-J18)	31	3.7	---
6	Diabetes mellitus (E10-E14)	25	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	23	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.7	---
10	Septicemia (A40-A41)	12	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	---
12	Atherosclerosis (I70)	8	1.0	---
13	Parkinson's disease (G20-G21)	5	0.6	---
14	Anemias (D50-D64)	3	0.4	---
14	Nutritional deficiencies (E40-E64)	3	0.4	---
...	All other causes (Residual)	259	31.2	---
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,016,896	100.0	1,004.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	481,991	23.9	240.2
2	Malignant neoplasms (C00-C97)	462,499	22.9	230.4
3	Chronic lower respiratory diseases (J40-J47)	127,116	6.3	63.3
4	Cerebrovascular diseases (I60-I69)	100,154	5.0	49.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99,288	4.9	49.5
6	Alzheimer's disease (G30)	72,772	3.6	36.3
7	Diabetes mellitus (E10-E14)	50,443	2.5	25.1
8	Influenza and pneumonia (J09-J18)	40,460	2.0	20.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,727	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,105	1.6	16.5
11	Septicemia (A40-A41)	26,592	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25,720	1.3	12.8
13	Parkinson's disease (G20-G21)	21,201	1.1	10.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,056	1.0	10.5
15	Pneumonitis due to solids and liquids (J69)	15,073	0.7	7.5
...	All other causes (Residual)	405,699	20.1	202.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,094	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	35.6	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	11.7	2.9
3	Malignant neoplasms (C00-C97)	202	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	6.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.5	0.9
6	Influenza and pneumonia (J09-J18)	46	2.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	522	24.9	6.2
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,150	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	352	30.6	3.2
2	Malignant neoplasms (C00-C97)	239	20.8	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	6.5	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	6.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.9	0.3
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.4	0.3
7	Influenza and pneumonia (J09-J18)	22	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	19	1.7	*
9	Septicemia (A40-A41)	13	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
12	Meningitis (G00,G03)	6	0.5	*
13	Anemias (D50-D64)	4	0.3	*
13	Diabetes mellitus (E10-E14)	4	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	255	22.2	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,517	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	29.4	3.9
2	Malignant neoplasms (C00-C97)	262	17.3	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	13.2	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	5.1	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	4.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.7	0.2
8	Cerebrovascular diseases (I60-I69)	25	1.6	0.2
9	Influenza and pneumonia (J09-J18)	23	1.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
11	Septicemia (A40-A41)	14	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
...	All other causes (Residual)	279	18.4	2.4
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,580	100.0	45.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	47.8	22.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,246	22.3	10.3
3	Malignant neoplasms (C00-C97)	341	6.1	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	202	3.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	2.6	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	2.0	0.9
7	Cerebrovascular diseases (I60-I69)	36	0.6	0.3
8	Influenza and pneumonia (J09-J18)	32	0.6	0.3
9	Septicemia (A40-A41)	26	0.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.2
11	Diabetes mellitus (E10-E14)	21	0.4	0.2
13	Legal intervention (Y35,Y89.0)	12	0.2	*
14	Pneumonitis due to solids and liquids (J69)	11	0.2	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.2	*
...	All other causes (Residual)	670	12.0	5.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	10,895	100.0	83.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,503	50.5	42.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,101	19.3	16.2
3	Malignant neoplasms (C00-C97)	517	4.7	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	504	4.6	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	3.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.3	1.1
7	Diabetes mellitus (E10-E14)	88	0.8	0.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	62	0.6	0.5
9	Cerebrovascular diseases (I60-I69)	61	0.6	0.5
10	Influenza and pneumonia (J09-J18)	53	0.5	0.4
11	Septicemia (A40-A41)	40	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	34	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.2	0.2
14	Legal intervention (Y35,Y89.0)	24	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.2
...	All other causes (Residual)	1,383	12.7	10.6
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,777	100.0	102.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,666	44.3	45.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,378	18.6	19.0
3	Malignant neoplasms (C00-C97)	817	6.4	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	5.2	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	504	3.9	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	1.0	1.0
7	Diabetes mellitus (E10-E14)	124	1.0	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	99	0.8	0.8
9	Cerebrovascular diseases (I60-I69)	98	0.8	0.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	85	0.7	0.7
11	Influenza and pneumonia (J09-J18)	84	0.7	0.7
12	Septicemia (A40-A41)	79	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	68	0.5	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.3	0.3
...	All other causes (Residual)	1,904	14.9	15.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	14,489	100.0	118.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,692	39.3	46.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,303	15.9	18.8
3	Malignant neoplasms (C00-C97)	1,303	9.0	10.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,159	8.0	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	480	3.3	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	267	1.8	2.2
7	Diabetes mellitus (E10-E14)	210	1.4	1.7
8	Cerebrovascular diseases (I60-I69)	180	1.2	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	1.0	1.1
10	Influenza and pneumonia (J09-J18)	132	0.9	1.1
11	Septicemia (A40-A41)	91	0.6	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	90	0.6	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	72	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	0.5	0.6
...	All other causes (Residual)	2,230	15.4	18.3
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	16,936	100.0	148.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,085	30.0	44.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,462	14.5	21.6
3	Malignant neoplasms (C00-C97)	2,261	13.4	19.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,075	12.3	18.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	465	2.7	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	2.5	3.7
7	Diabetes mellitus (E10-E14)	360	2.1	3.2
8	Cerebrovascular diseases (I60-I69)	272	1.6	2.4
9	Septicemia (A40-A41)	146	0.9	1.3
10	Influenza and pneumonia (J09-J18)	135	0.8	1.2
11	Chronic lower respiratory diseases (J40-J47)	129	0.8	1.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.7	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	113	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	64	0.4	0.6
...	All other causes (Residual)	2,715	16.0	23.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	27,004	100.0	205.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,878	21.8	44.7
2	Malignant neoplasms (C00-C97)	4,845	17.9	36.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,238	15.7	32.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,970	11.0	22.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,133	4.2	8.6
6	Diabetes mellitus (E10-E14)	682	2.5	5.2
7	Cerebrovascular diseases (I60-I69)	586	2.2	4.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	445	1.6	3.4
9	Chronic lower respiratory diseases (J40-J47)	350	1.3	2.7
10	Septicemia (A40-A41)	276	1.0	2.1
11	Influenza and pneumonia (J09-J18)	267	1.0	2.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	254	0.9	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	0.6	1.1
15	Viral hepatitis (B15-B19)	131	0.5	1.0
...	All other causes (Residual)	4,628	17.1	35.2
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	46,264	100.0	322.4
1	Malignant neoplasms (C00-C97)	10,873	23.5	75.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,462	18.3	59.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,354	15.9	51.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,586	7.8	25.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,431	5.3	16.9
6	Diabetes mellitus (E10-E14)	1,269	2.7	8.8
7	Cerebrovascular diseases (I60-I69)	1,115	2.4	7.8
8	Chronic lower respiratory diseases (J40-J47)	950	2.1	6.6
9	Septicemia (A40-A41)	527	1.1	3.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	462	1.0	3.2
11	Influenza and pneumonia (J09-J18)	444	1.0	3.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	434	0.9	3.0
13	Viral hepatitis (B15-B19)	393	0.8	2.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	0.7	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	217	0.5	1.5
...	All other causes (Residual)	7,405	16.0	51.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	77,878	100.0	491.0
1	Malignant neoplasms (C00-C97)	22,799	29.3	143.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,443	19.8	97.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,198	10.5	51.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,167	5.4	26.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,949	5.1	24.9
6	Chronic lower respiratory diseases (J40-J47)	2,651	3.4	16.7
7	Diabetes mellitus (E10-E14)	2,108	2.7	13.3
8	Cerebrovascular diseases (I60-I69)	1,904	2.4	12.0
9	Viral hepatitis (B15-B19)	976	1.3	6.2
10	Septicemia (A40-A41)	919	1.2	5.8
11	Influenza and pneumonia (J09-J18)	759	1.0	4.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	668	0.9	4.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	515	0.7	3.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	471	0.6	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	0.6	2.7
...	All other causes (Residual)	11,919	15.3	75.1
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	106,760	100.0	705.1
1	Malignant neoplasms (C00-C97)	35,972	33.7	237.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,073	20.7	145.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,829	6.4	45.1
4	Chronic lower respiratory diseases (J40-J47)	4,944	4.6	32.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,457	4.2	29.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,651	3.4	24.1
7	Diabetes mellitus (E10-E14)	3,332	3.1	22.0
8	Cerebrovascular diseases (I60-I69)	2,866	2.7	18.9
9	Viral hepatitis (B15-B19)	1,541	1.4	10.2
10	Septicemia (A40-A41)	1,360	1.3	9.0
11	Influenza and pneumonia (J09-J18)	1,200	1.1	7.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,031	1.0	6.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	725	0.7	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	514	0.5	3.4
15	Aortic aneurysm and dissection (I71)	409	0.4	2.7
...	All other causes (Residual)	15,856	14.9	104.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	133,830	100.0	996.5
1	Malignant neoplasms (C00-C97)	48,975	36.6	364.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,683	21.4	213.6
3	Chronic lower respiratory diseases (J40-J47)	7,999	6.0	59.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,243	3.9	39.0
5	Diabetes mellitus (E10-E14)	4,598	3.4	34.2
6	Cerebrovascular diseases (I60-I69)	3,979	3.0	29.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,814	2.8	28.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,582	1.9	19.2
9	Septicemia (A40-A41)	1,958	1.5	14.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,641	1.2	12.2
11	Influenza and pneumonia (J09-J18)	1,546	1.2	11.5
12	Viral hepatitis (B15-B19)	1,058	0.8	7.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,047	0.8	7.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	673	0.5	5.0
15	Pneumonitis due to solids and liquids (J69)	534	0.4	4.0
...	All other causes (Residual)	19,500	14.6	145.2
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	160,009	100.0	1,467.1
1	Malignant neoplasms (C00-C97)	59,544	37.2	545.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,667	21.0	308.7
3	Chronic lower respiratory diseases (J40-J47)	13,019	8.1	119.4
4	Cerebrovascular diseases (I60-I69)	5,515	3.4	50.6
5	Diabetes mellitus (E10-E14)	5,477	3.4	50.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,475	2.8	41.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,914	1.8	26.7
8	Septicemia (A40-A41)	2,382	1.5	21.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,309	1.4	21.2
10	Influenza and pneumonia (J09-J18)	2,055	1.3	18.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,786	1.1	16.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,167	0.7	10.7
13	Alzheimer's disease (G30)	1,106	0.7	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	895	0.6	8.2
15	Parkinson's disease (G20-G21)	763	0.5	7.0
...	All other causes (Residual)	22,935	14.3	210.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	179,383	100.0	2,287.1
1	Malignant neoplasms (C00-C97)	62,137	34.6	792.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,603	21.0	479.4
3	Chronic lower respiratory diseases (J40-J47)	16,862	9.4	215.0
4	Cerebrovascular diseases (I60-I69)	7,517	4.2	95.8
5	Diabetes mellitus (E10-E14)	5,650	3.1	72.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,114	2.3	52.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,027	1.7	38.6
8	Septicemia (A40-A41)	2,706	1.5	34.5
9	Influenza and pneumonia (J09-J18)	2,632	1.5	33.6
10	Alzheimer's disease (G30)	2,473	1.4	31.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,102	1.2	26.8
12	Parkinson's disease (G20-G21)	1,783	1.0	22.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,323	0.7	16.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,276	0.7	16.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,257	0.7	16.0
...	All other causes (Residual)	26,921	15.0	343.2
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	219,714	100.0	3,699.1
1	Malignant neoplasms (C00-C97)	63,415	28.9	1,067.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,648	22.1	819.0
3	Chronic lower respiratory diseases (J40-J47)	20,186	9.2	339.8
4	Cerebrovascular diseases (I60-I69)	11,161	5.1	187.9
5	Alzheimer's disease (G30)	6,379	2.9	107.4
6	Diabetes mellitus (E10-E14)	6,181	2.8	104.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,038	2.3	84.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,027	1.8	67.8
9	Influenza and pneumonia (J09-J18)	3,925	1.8	66.1
10	Parkinson's disease (G20-G21)	3,493	1.6	58.8
11	Septicemia (A40-A41)	3,343	1.5	56.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,897	0.9	31.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,745	0.8	29.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,704	0.8	28.7
15	Pneumonitis due to solids and liquids (J69)	1,508	0.7	25.4
...	All other causes (Residual)	37,064	16.9	624.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	291,625	100.0	6,165.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,767	24.3	1,496.1
2	Malignant neoplasms (C00-C97)	64,203	22.0	1,357.4
3	Chronic lower respiratory diseases (J40-J47)	23,336	8.0	493.4
4	Cerebrovascular diseases (I60-I69)	17,586	6.0	371.8
5	Alzheimer's disease (G30)	13,581	4.7	287.1
6	Diabetes mellitus (E10-E14)	7,362	2.5	155.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,308	2.5	154.5
8	Influenza and pneumonia (J09-J18)	6,532	2.2	138.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,957	2.0	125.9
10	Parkinson's disease (G20-G21)	5,467	1.9	115.6
11	Septicemia (A40-A41)	4,235	1.5	89.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,063	1.1	64.8
13	Pneumonitis due to solids and liquids (J69)	2,565	0.9	54.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,287	0.8	48.4
15	Aortic aneurysm and dissection (I71)	1,411	0.5	29.8
...	All other causes (Residual)	55,965	19.2	1,183.2
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	330,594	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	91,187	27.6	---
2	Malignant neoplasms (C00-C97)	51,387	15.5	---
3	Cerebrovascular diseases (I60-I69)	21,959	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	20,848	6.3	---
5	Alzheimer's disease (G30)	20,723	6.3	---
6	Influenza and pneumonia (J09-J18)	8,630	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,479	2.6	---
8	Diabetes mellitus (E10-E14)	7,162	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,703	2.0	---
10	Parkinson's disease (G20-G21)	5,577	1.7	---
11	Septicemia (A40-A41)	4,505	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,400	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,461	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,477	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,585	0.5	---
...	All other causes (Residual)	71,511	21.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	249,204	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,005	30.9	---
2	Malignant neoplasms (C00-C97)	25,134	10.1	---
3	Alzheimer's disease (G30)	18,481	7.4	---
4	Cerebrovascular diseases (I60-I69)	17,462	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	11,566	4.6	---
6	Influenza and pneumonia (J09-J18)	7,568	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,550	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,888	2.0	---
9	Diabetes mellitus (E10-E14)	4,304	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,074	1.6	---
11	Pneumonitis due to solids and liquids (J69)	3,000	1.2	---
12	Parkinson's disease (G20-G21)	2,908	1.2	---
13	Septicemia (A40-A41)	2,801	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,432	0.6	---
15	Atherosclerosis (I70)	1,346	0.5	---
...	All other causes (Residual)	60,685	24.4	---
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	98,585	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,613	33.1	---
2	Alzheimer's disease (G30)	7,761	7.9	---
3	Cerebrovascular diseases (I60-I69)	6,579	6.7	---
4	Malignant neoplasms (C00-C97)	6,455	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	3,523	3.6	---
6	Influenza and pneumonia (J09-J18)	3,453	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,610	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,015	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,741	1.8	---
10	Diabetes mellitus (E10-E14)	1,315	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,183	1.2	---
12	Septicemia (A40-A41)	924	0.9	---
13	Atherosclerosis (I70)	703	0.7	---
14	Parkinson's disease (G20-G21)	630	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	464	0.5	---
...	All other causes (Residual)	26,616	27.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	19,941	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,906	34.6	---
2	Alzheimer's disease (G30)	1,479	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,168	5.9	---
4	Influenza and pneumonia (J09-J18)	853	4.3	---
5	Malignant neoplasms (C00-C97)	775	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	509	2.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	508	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	487	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	312	1.6	---
10	Pneumonitis due to solids and liquids (J69)	229	1.1	---
11	Atherosclerosis (I70)	217	1.1	---
12	Diabetes mellitus (E10-E14)	183	0.9	---
13	Septicemia (A40-A41)	154	0.8	---
14	Anemias (D50-D64)	79	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.3	---
...	All other causes (Residual)	6,014	30.2	---
Non-Hispanic, white, male, all ages				
...	All causes	998,832	100.0	1,011.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,116	25.0	253.2
2	Malignant neoplasms (C00-C97)	243,925	24.4	247.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,388	6.0	61.1
4	Chronic lower respiratory diseases (J40-J47)	59,084	5.9	59.8
5	Cerebrovascular diseases (I60-I69)	39,786	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,043	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,356	2.6	26.7
8	Alzheimer's disease (G30)	22,403	2.2	22.7
9	Influenza and pneumonia (J09-J18)	18,960	1.9	19.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,720	1.7	16.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16,518	1.7	16.7
12	Parkinson's disease (G20-G21)	12,654	1.3	12.8
13	Septicemia (A40-A41)	12,446	1.2	12.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,263	0.8	8.4
15	Pneumonitis due to solids and liquids (J69)	8,172	0.8	8.3
...	All other causes (Residual)	175,998	17.6	178.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,202	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	36.8	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	11.0	3.1
3	Malignant neoplasms (C00-C97)	108	9.0	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.8	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
9	Septicemia (A40-A41)	11	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
15	Meningococcal infection (A39)	3	0.2	*
...	All other causes (Residual)	302	25.1	7.0
Non-Hispanic, white, male, 5-9 years				
...	All causes	645	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	32.1	3.6
2	Malignant neoplasms (C00-C97)	141	21.9	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.6	0.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.4	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.2	*
7	Septicemia (A40-A41)	10	1.6	*
8	Influenza and pneumonia (J09-J18)	9	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
11	Meningitis (G00,G03)	4	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Anemias (D50-D64)	2	0.3	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Hernia (K40-K46)	2	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
...	All other causes (Residual)	145	22.5	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 10-14 years				
...	All causes	903	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	287	31.8	4.8
2	Malignant neoplasms (C00-C97)	147	16.3	2.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	16.3	2.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	4.0	0.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	4.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.0	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.7	*
8	Cerebrovascular diseases (I60-I69)	13	1.4	*
9	Influenza and pneumonia (J09-J18)	11	1.2	*
10	Septicemia (A40-A41)	9	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
12	Diabetes mellitus (E10-E14)	6	0.7	*
13	Meningitis (G00,G03)	2	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	156	17.3	2.6
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,848	100.0	61.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,846	48.0	29.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	968	25.2	15.5
3	Malignant neoplasms (C00-C97)	197	5.1	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	3.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	2.4	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	1.7	1.1
7	Cerebrovascular diseases (I60-I69)	18	0.5	*
7	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
9	Influenza and pneumonia (J09-J18)	17	0.4	*
10	Septicemia (A40-A41)	12	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.3	*
10	Legal intervention (Y35,Y89.0)	12	0.3	*
13	Diabetes mellitus (E10-E14)	11	0.3	*
14	Pneumonitis due to solids and liquids (J69)	7	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (Residual)	424	11.0	6.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 20-24 years				
...	All causes	7,978	100.0	120.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,204	52.7	63.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,755	22.0	26.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	4.5	5.4
4	Malignant neoplasms (C00-C97)	288	3.6	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	2.8	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.0	1.2
7	Diabetes mellitus (E10-E14)	54	0.7	0.8
8	Cerebrovascular diseases (I60-I69)	36	0.5	0.5
9	Influenza and pneumonia (J09-J18)	35	0.4	0.5
10	Legal intervention (Y35,Y89.0)	21	0.3	0.3
11	Septicemia (A40-A41)	19	0.2	*
11	Chronic lower respiratory diseases (J40-J47)	19	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.2	*
14	Pneumonitis due to solids and liquids (J69)	14	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.1	*
...	All other causes (Residual)	845	10.6	12.8
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,993	100.0	142.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,191	46.6	66.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,928	21.4	30.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	458	5.1	7.2
4	Malignant neoplasms (C00-C97)	451	5.0	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	362	4.0	5.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	0.9	1.3
7	Diabetes mellitus (E10-E14)	71	0.8	1.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.7	1.0
9	Cerebrovascular diseases (I60-I69)	51	0.6	0.8
10	Septicemia (A40-A41)	49	0.5	0.8
11	Influenza and pneumonia (J09-J18)	46	0.5	0.7
12	Chronic lower respiratory diseases (J40-J47)	38	0.4	0.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.4	0.6
14	Legal intervention (Y35,Y89.0)	30	0.3	0.5
15	Aortic aneurysm and dissection (I71)	22	0.2	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.2	0.3
...	All other causes (Residual)	1,093	12.2	17.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 30-34 years				
...	All causes	9,613	100.0	156.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,017	41.8	65.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,801	18.7	29.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	796	8.3	12.9
4	Malignant neoplasms (C00-C97)	616	6.4	10.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	3.4	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	162	1.7	2.6
7	Diabetes mellitus (E10-E14)	126	1.3	2.0
8	Cerebrovascular diseases (I60-I69)	99	1.0	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	1.3
10	Influenza and pneumonia (J09-J18)	76	0.8	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	56	0.6	0.9
12	Septicemia (A40-A41)	49	0.5	0.8
13	Chronic lower respiratory diseases (J40-J47)	38	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.4	0.6
15	Legal intervention (Y35,Y89.0)	29	0.3	0.5
...	All other causes (Residual)	1,306	13.6	21.2
Non-Hispanic, white, male, 35-39 years				
...	All causes	10,764	100.0	187.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,529	32.8	61.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,888	17.5	32.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,437	13.4	25.1
4	Malignant neoplasms (C00-C97)	1,021	9.5	17.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	287	2.7	5.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	286	2.7	5.0
7	Diabetes mellitus (E10-E14)	219	2.0	3.8
8	Cerebrovascular diseases (I60-I69)	149	1.4	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	86	0.8	1.5
10	Septicemia (A40-A41)	81	0.8	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.6	1.1
12	Influenza and pneumonia (J09-J18)	61	0.6	1.1
13	Chronic lower respiratory diseases (J40-J47)	55	0.5	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.5	0.9
15	Aortic aneurysm and dissection (I71)	45	0.4	0.8
...	All other causes (Residual)	1,504	14.0	26.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,572	100.0	251.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,964	23.9	60.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,949	17.8	44.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,254	13.6	34.2
4	Malignant neoplasms (C00-C97)	2,145	12.9	32.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	705	4.3	10.7
6	Diabetes mellitus (E10-E14)	417	2.5	6.3
7	Cerebrovascular diseases (I60-I69)	306	1.8	4.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	1.8	4.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	212	1.3	3.2
10	Chronic lower respiratory diseases (J40-J47)	152	0.9	2.3
11	Influenza and pneumonia (J09-J18)	136	0.8	2.1
12	Septicemia (A40-A41)	135	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	0.6	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	0.5	1.3
15	Viral hepatitis (B15-B19)	79	0.5	1.2
...	All other causes (Residual)	2,622	15.8	39.8
Non-Hispanic, white, male, 45-49 years				
...	All causes	28,166	100.0	394.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,015	21.4	84.1
2	Malignant neoplasms (C00-C97)	5,243	18.6	73.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,838	17.2	67.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,638	9.4	36.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,572	5.6	22.0
6	Diabetes mellitus (E10-E14)	816	2.9	11.4
7	Cerebrovascular diseases (I60-I69)	577	2.0	8.1
8	Chronic lower respiratory diseases (J40-J47)	421	1.5	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	362	1.3	5.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	1.1	4.3
11	Septicemia (A40-A41)	290	1.0	4.1
12	Viral hepatitis (B15-B19)	264	0.9	3.7
13	Influenza and pneumonia (J09-J18)	229	0.8	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	211	0.7	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	0.5	2.0
...	All other causes (Residual)	4,245	15.1	59.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 50-54 years				
...	All causes	48,041	100.0	611.8
1	Malignant neoplasms (C00-C97)	11,889	24.7	151.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,270	23.5	143.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,378	11.2	68.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,103	6.5	39.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,682	5.6	34.2
6	Diabetes mellitus (E10-E14)	1,345	2.8	17.1
7	Chronic lower respiratory diseases (J40-J47)	1,248	2.6	15.9
8	Cerebrovascular diseases (I60-I69)	1,043	2.2	13.3
9	Viral hepatitis (B15-B19)	694	1.4	8.8
10	Septicemia (A40-A41)	476	1.0	6.1
11	Influenza and pneumonia (J09-J18)	433	0.9	5.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	404	0.8	5.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	403	0.8	5.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	0.8	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	312	0.6	4.0
...	All other causes (Residual)	6,986	14.5	89.0
Non-Hispanic, white, male, 55-59 years				
...	All causes	66,224	100.0	891.8
1	Malignant neoplasms (C00-C97)	19,916	30.1	268.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,968	24.1	215.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,524	6.8	60.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,145	4.7	42.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,774	4.2	37.4
6	Chronic lower respiratory diseases (J40-J47)	2,482	3.7	33.4
7	Diabetes mellitus (E10-E14)	2,140	3.2	28.8
8	Cerebrovascular diseases (I60-I69)	1,683	2.5	22.7
9	Viral hepatitis (B15-B19)	1,103	1.7	14.9
10	Septicemia (A40-A41)	750	1.1	10.1
11	Influenza and pneumonia (J09-J18)	689	1.0	9.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	604	0.9	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	463	0.7	6.2
14	Aortic aneurysm and dissection (I71)	320	0.5	4.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	305	0.5	4.1
...	All other causes (Residual)	9,358	14.1	126.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 60-64 years				
...	All causes	81,410	100.0	1,247.2
1	Malignant neoplasms (C00-C97)	27,334	33.6	418.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,192	24.8	309.3
3	Chronic lower respiratory diseases (J40-J47)	4,172	5.1	63.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,563	4.4	54.6
5	Diabetes mellitus (E10-E14)	2,862	3.5	43.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,656	3.3	40.7
7	Cerebrovascular diseases (I60-I69)	2,299	2.8	35.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,003	2.5	30.7
9	Septicemia (A40-A41)	1,061	1.3	16.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	923	1.1	14.1
11	Influenza and pneumonia (J09-J18)	909	1.1	13.9
12	Viral hepatitis (B15-B19)	770	0.9	11.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	0.8	10.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	399	0.5	6.1
15	Aortic aneurysm and dissection (I71)	364	0.4	5.6
...	All other causes (Residual)	11,234	13.8	172.1
Non-Hispanic, white, male, 65-69 years				
...	All causes	93,047	100.0	1,778.2
1	Malignant neoplasms (C00-C97)	33,037	35.5	631.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,514	24.2	430.3
3	Chronic lower respiratory diseases (J40-J47)	6,538	7.0	124.9
4	Diabetes mellitus (E10-E14)	3,313	3.6	63.3
5	Cerebrovascular diseases (I60-I69)	3,062	3.3	58.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,904	3.1	55.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,864	2.0	35.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,388	1.5	26.5
9	Septicemia (A40-A41)	1,266	1.4	24.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,262	1.4	24.1
11	Influenza and pneumonia (J09-J18)	1,170	1.3	22.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	647	0.7	12.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	527	0.6	10.1
14	Parkinson's disease (G20-G21)	520	0.6	9.9
15	Aortic aneurysm and dissection (I71)	492	0.5	9.4
...	All other causes (Residual)	12,543	13.5	239.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 70-74 years				
...	All causes	100,405	100.0	2,744.5
1	Malignant neoplasms (C00-C97)	34,306	34.2	937.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,566	23.5	644.2
3	Chronic lower respiratory diseases (J40-J47)	8,259	8.2	225.8
4	Cerebrovascular diseases (I60-I69)	3,845	3.8	105.1
5	Diabetes mellitus (E10-E14)	3,261	3.2	89.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,532	2.5	69.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,676	1.7	45.8
8	Influenza and pneumonia (J09-J18)	1,463	1.5	40.0
9	Septicemia (A40-A41)	1,347	1.3	36.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,264	1.3	34.6
11	Parkinson's disease (G20-G21)	1,189	1.2	32.5
12	Alzheimer's disease (G30)	1,069	1.1	29.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,063	1.1	29.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	806	0.8	22.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	712	0.7	19.5
...	All other causes (Residual)	14,047	14.0	384.0
Non-Hispanic, white, male, 75-79 years				
...	All causes	115,961	100.0	4,367.6
1	Malignant neoplasms (C00-C97)	34,362	29.6	1,294.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,045	24.2	1,056.3
3	Chronic lower respiratory diseases (J40-J47)	9,825	8.5	370.1
4	Cerebrovascular diseases (I60-I69)	5,200	4.5	195.9
5	Diabetes mellitus (E10-E14)	3,370	2.9	126.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,779	2.4	104.7
7	Alzheimer's disease (G30)	2,505	2.2	94.4
8	Parkinson's disease (G20-G21)	2,263	2.0	85.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,175	1.9	81.9
10	Influenza and pneumonia (J09-J18)	2,050	1.8	77.2
11	Septicemia (A40-A41)	1,639	1.4	61.7
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	986	0.9	37.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	975	0.8	36.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	946	0.8	35.6
15	Pneumonitis due to solids and liquids (J69)	874	0.8	32.9
...	All other causes (Residual)	17,967	15.5	676.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 80-84 years				
...	All causes	142,312	100.0	7,269.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,974	26.0	1,888.7
2	Malignant neoplasms (C00-C97)	33,717	23.7	1,722.3
3	Chronic lower respiratory diseases (J40-J47)	10,963	7.7	560.0
4	Cerebrovascular diseases (I60-I69)	7,311	5.1	373.4
5	Alzheimer's disease (G30)	4,965	3.5	253.6
6	Diabetes mellitus (E10-E14)	3,756	2.6	191.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,721	2.6	190.1
8	Parkinson's disease (G20-G21)	3,480	2.4	177.8
9	Influenza and pneumonia (J09-J18)	3,315	2.3	169.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,149	2.2	160.9
11	Septicemia (A40-A41)	1,937	1.4	98.9
12	Pneumonitis due to solids and liquids (J69)	1,480	1.0	75.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,264	0.9	64.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,249	0.9	63.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	830	0.6	42.4
...	All other causes (Residual)	24,201	17.0	1,236.2
Non-Hispanic, white, male, 85-89 years				
...	All causes	140,129	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,845	29.1	---
2	Malignant neoplasms (C00-C97)	25,091	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,136	6.5	---
4	Cerebrovascular diseases (I60-I69)	7,587	5.4	---
5	Alzheimer's disease (G30)	6,569	4.7	---
6	Influenza and pneumonia (J09-J18)	3,910	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,799	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,248	2.3	---
9	Diabetes mellitus (E10-E14)	3,217	2.3	---
10	Parkinson's disease (G20-G21)	3,128	2.2	---
11	Septicemia (A40-A41)	1,909	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,840	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,474	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,212	0.9	---
15	Aortic aneurysm and dissection (I71)	645	0.5	---
...	All other causes (Residual)	26,519	18.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 90-94 years				
...	All causes	87,581	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,409	32.4	---
2	Malignant neoplasms (C00-C97)	11,257	12.9	---
3	Alzheimer's disease (G30)	4,861	5.6	---
4	Cerebrovascular diseases (I60-I69)	4,856	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	4,441	5.1	---
6	Influenza and pneumonia (J09-J18)	3,054	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,505	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,113	2.4	---
9	Diabetes mellitus (E10-E14)	1,648	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,524	1.7	---
11	Parkinson's disease (G20-G21)	1,487	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,123	1.3	---
13	Septicemia (A40-A41)	1,070	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	589	0.7	---
15	Atherosclerosis (I70)	460	0.5	---
...	All other causes (Residual)	18,184	20.8	---
Non-Hispanic, white, male, 95-99 years				
...	All causes	25,774	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,931	34.7	---
2	Malignant neoplasms (C00-C97)	2,431	9.4	---
3	Alzheimer's disease (G30)	1,449	5.6	---
4	Cerebrovascular diseases (I60-I69)	1,434	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,150	4.5	---
6	Influenza and pneumonia (J09-J18)	1,096	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	737	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	618	2.4	---
9	Pneumonitis due to solids and liquids (J69)	496	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	449	1.7	---
11	Diabetes mellitus (E10-E14)	377	1.5	---
12	Septicemia (A40-A41)	255	1.0	---
13	Parkinson's disease (G20-G21)	242	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	171	0.7	---
15	Atherosclerosis (I70)	166	0.6	---
...	All other causes (Residual)	5,772	22.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 100 years and over				
...	All causes	3,337	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,259	37.7	---
2	Malignant neoplasms (C00-C97)	207	6.2	---
3	Influenza and pneumonia (J09-J18)	180	5.4	---
4	Cerebrovascular diseases (I60-I69)	169	5.1	---
5	Alzheimer's disease (G30)	153	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	103	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	2.1	---
10	Pneumonitis due to solids and liquids (J69)	68	2.0	---
11	Atherosclerosis (I70)	38	1.1	---
12	Septicemia (A40-A41)	33	1.0	---
13	Diabetes mellitus (E10-E14)	28	0.8	---
14	Parkinson's disease (G20-G21)	19	0.6	---
15	Anemias (D50-D64)	17	0.5	---
...	All other causes (Residual)	831	24.9	---
Non-Hispanic, white, female, all ages				
...	All causes	1,018,064	100.0	998.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,875	22.8	227.5
2	Malignant neoplasms (C00-C97)	218,574	21.5	214.4
3	Chronic lower respiratory diseases (J40-J47)	68,032	6.7	66.7
4	Cerebrovascular diseases (I60-I69)	60,368	5.9	59.2
5	Alzheimer's disease (G30)	50,369	4.9	49.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,900	3.8	38.2
7	Diabetes mellitus (E10-E14)	23,400	2.3	23.0
8	Influenza and pneumonia (J09-J18)	21,500	2.1	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,385	1.6	16.1
10	Septicemia (A40-A41)	14,146	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,793	1.3	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,202	0.9	9.0
13	Parkinson's disease (G20-G21)	8,547	0.8	8.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,371	0.7	7.2
15	Pneumonitis due to solids and liquids (J69)	6,901	0.7	6.8
...	All other causes (Residual)	229,701	22.6	225.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	892	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	34.0	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.6	2.7
3	Malignant neoplasms (C00-C97)	94	10.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	15	1.7	*
8	Septicemia (A40-A41)	11	1.2	*
9	Cerebrovascular diseases (I60-I69)	6	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Anemias (D50-D64)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	218	24.4	5.3
Non-Hispanic, white, female, 5-9 years				
...	All causes	505	100.0	9.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	28.7	2.7
2	Malignant neoplasms (C00-C97)	98	19.4	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	7.7	0.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	7.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.8	*
7	Influenza and pneumonia (J09-J18)	13	2.6	*
8	Cerebrovascular diseases (I60-I69)	12	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Septicemia (A40-A41)	3	0.6	*
12	Anemias (D50-D64)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	106	21.0	2.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 10-14 years				
...	All causes	614	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	25.9	2.8
2	Malignant neoplasms (C00-C97)	115	18.7	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	8.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.4	0.5
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
7	Influenza and pneumonia (J09-J18)	12	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.3	*
11	Septicemia (A40-A41)	5	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
13	Anemias (D50-D64)	2	0.3	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	125	20.4	2.2
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,732	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	47.6	14.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	278	16.1	4.7
3	Malignant neoplasms (C00-C97)	144	8.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.5	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.1	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	2.7	0.8
7	Cerebrovascular diseases (I60-I69)	18	1.0	*
8	Influenza and pneumonia (J09-J18)	15	0.9	*
9	Septicemia (A40-A41)	14	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.6	*
11	Diabetes mellitus (E10-E14)	10	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
...	All other causes (Residual)	236	13.6	4.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,917	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,299	44.5	20.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	346	11.9	5.4
3	Malignant neoplasms (C00-C97)	229	7.9	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	5.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.8	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	2.2	1.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	62	2.1	1.0
8	Diabetes mellitus (E10-E14)	34	1.2	0.5
9	Cerebrovascular diseases (I60-I69)	25	0.9	0.4
10	Septicemia (A40-A41)	21	0.7	0.3
11	Influenza and pneumonia (J09-J18)	18	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	517	17.7	8.1
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,784	100.0	61.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,475	39.0	23.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	450	11.9	7.3
3	Malignant neoplasms (C00-C97)	366	9.7	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	5.4	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	3.8	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	85	2.2	1.4
7	Diabetes mellitus (E10-E14)	53	1.4	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	1.3	0.8
9	Cerebrovascular diseases (I60-I69)	47	1.2	0.8
10	Influenza and pneumonia (J09-J18)	38	1.0	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.9	0.5
12	Septicemia (A40-A41)	30	0.8	0.5
12	Chronic lower respiratory diseases (J40-J47)	30	0.8	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.3	*
...	All other causes (Residual)	751	19.8	12.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,876	100.0	80.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,675	34.4	27.6
2	Malignant neoplasms (C00-C97)	687	14.1	11.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	502	10.3	8.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	363	7.4	6.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	3.1	2.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	105	2.2	1.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	90	1.8	1.5
8	Diabetes mellitus (E10-E14)	84	1.7	1.4
9	Cerebrovascular diseases (I60-I69)	81	1.7	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.3	1.0
11	Influenza and pneumonia (J09-J18)	56	1.1	0.9
12	Septicemia (A40-A41)	42	0.9	0.7
13	Chronic lower respiratory diseases (J40-J47)	34	0.7	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.6	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.4
...	All other causes (Residual)	888	18.2	14.7
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,172	100.0	108.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,556	25.2	27.5
2	Malignant neoplasms (C00-C97)	1,240	20.1	21.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	638	10.3	11.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	574	9.3	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	178	2.9	3.1
6	Diabetes mellitus (E10-E14)	141	2.3	2.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	2.3	2.5
8	Cerebrovascular diseases (I60-I69)	123	2.0	2.2
9	Influenza and pneumonia (J09-J18)	74	1.2	1.3
9	Chronic lower respiratory diseases (J40-J47)	74	1.2	1.3
11	Septicemia (A40-A41)	65	1.1	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.0	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.9	1.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.4	0.5
...	All other causes (Residual)	1,166	18.9	20.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,432	100.0	159.4
1	Malignant neoplasms (C00-C97)	2,700	25.9	41.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,914	18.3	29.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,289	12.4	19.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	6.9	10.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	428	4.1	6.5
6	Cerebrovascular diseases (I60-I69)	280	2.7	4.3
7	Diabetes mellitus (E10-E14)	265	2.5	4.0
8	Chronic lower respiratory diseases (J40-J47)	198	1.9	3.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	1.4	2.2
10	Septicemia (A40-A41)	141	1.4	2.2
11	Influenza and pneumonia (J09-J18)	131	1.3	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.6	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.6	1.0
14	Viral hepatitis (B15-B19)	52	0.5	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.4	0.6
...	All other causes (Residual)	1,964	18.8	30.0
Non-Hispanic, white, female, 45-49 years				
...	All causes	18,098	100.0	251.4
1	Malignant neoplasms (C00-C97)	5,630	31.1	78.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,516	13.9	34.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,447	13.5	34.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	948	5.2	13.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	859	4.7	11.9
6	Cerebrovascular diseases (I60-I69)	538	3.0	7.5
7	Chronic lower respiratory diseases (J40-J47)	529	2.9	7.3
8	Diabetes mellitus (E10-E14)	453	2.5	6.3
9	Septicemia (A40-A41)	237	1.3	3.3
10	Influenza and pneumonia (J09-J18)	215	1.2	3.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	0.9	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	0.7	1.8
13	Viral hepatitis (B15-B19)	129	0.7	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	0.5	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.4	1.1
...	All other causes (Residual)	3,138	17.3	43.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 50-54 years				
...	All causes	29,837	100.0	372.5
1	Malignant neoplasms (C00-C97)	10,910	36.6	136.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,173	14.0	52.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,820	9.5	35.2
4	Chronic lower respiratory diseases (J40-J47)	1,403	4.7	17.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,267	4.2	15.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,064	3.6	13.3
7	Cerebrovascular diseases (I60-I69)	861	2.9	10.8
8	Diabetes mellitus (E10-E14)	763	2.6	9.5
9	Septicemia (A40-A41)	443	1.5	5.5
10	Influenza and pneumonia (J09-J18)	326	1.1	4.1
11	Viral hepatitis (B15-B19)	282	0.9	3.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	0.9	3.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	0.5	1.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.5	1.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	0.5	1.7
...	All other causes (Residual)	4,838	16.2	60.4
Non-Hispanic, white, female, 55-59 years				
...	All causes	40,536	100.0	525.4
1	Malignant neoplasms (C00-C97)	16,056	39.6	208.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,105	15.1	79.1
3	Chronic lower respiratory diseases (J40-J47)	2,462	6.1	31.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,305	5.7	29.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,312	3.2	17.0
6	Diabetes mellitus (E10-E14)	1,192	2.9	15.5
7	Cerebrovascular diseases (I60-I69)	1,183	2.9	15.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	877	2.2	11.4
9	Septicemia (A40-A41)	610	1.5	7.9
10	Influenza and pneumonia (J09-J18)	511	1.3	6.6
11	Viral hepatitis (B15-B19)	438	1.1	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	1.1	5.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	262	0.6	3.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	238	0.6	3.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	0.4	2.0
...	All other causes (Residual)	6,401	15.8	83.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 60-64 years				
...	All causes	52,420	100.0	759.4
1	Malignant neoplasms (C00-C97)	21,641	41.3	313.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,491	16.2	123.0
3	Chronic lower respiratory diseases (J40-J47)	3,827	7.3	55.4
4	Diabetes mellitus (E10-E14)	1,736	3.3	25.2
5	Cerebrovascular diseases (I60-I69)	1,680	3.2	24.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,680	3.2	24.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,158	2.2	16.8
8	Septicemia (A40-A41)	897	1.7	13.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	718	1.4	10.4
10	Influenza and pneumonia (J09-J18)	637	1.2	9.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	579	1.1	8.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	378	0.7	5.5
13	Viral hepatitis (B15-B19)	288	0.5	4.2
14	Alzheimer's disease (G30)	279	0.5	4.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	274	0.5	4.0
...	All other causes (Residual)	8,157	15.6	118.2
Non-Hispanic, white, female, 65-69 years				
...	All causes	66,962	100.0	1,180.1
1	Malignant neoplasms (C00-C97)	26,507	39.6	467.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,153	16.7	196.6
3	Chronic lower respiratory diseases (J40-J47)	6,481	9.7	114.2
4	Cerebrovascular diseases (I60-I69)	2,453	3.7	43.2
5	Diabetes mellitus (E10-E14)	2,164	3.2	38.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,571	2.3	27.7
7	Septicemia (A40-A41)	1,116	1.7	19.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,050	1.6	18.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,047	1.6	18.5
10	Influenza and pneumonia (J09-J18)	885	1.3	15.6
11	Alzheimer's disease (G30)	619	0.9	10.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	520	0.8	9.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	398	0.6	7.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	368	0.5	6.5
15	Pneumonitis due to solids and liquids (J69)	268	0.4	4.7
...	All other causes (Residual)	10,362	15.5	182.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 70-74 years				
...	All causes	78,978	100.0	1,887.2
1	Malignant neoplasms (C00-C97)	27,831	35.2	665.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,037	17.8	335.4
3	Chronic lower respiratory diseases (J40-J47)	8,603	10.9	205.6
4	Cerebrovascular diseases (I60-I69)	3,672	4.6	87.7
5	Diabetes mellitus (E10-E14)	2,389	3.0	57.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,582	2.0	37.8
7	Alzheimer's disease (G30)	1,404	1.8	33.5
8	Septicemia (A40-A41)	1,359	1.7	32.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,351	1.7	32.3
10	Influenza and pneumonia (J09-J18)	1,169	1.5	27.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	838	1.1	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	611	0.8	14.6
13	Parkinson's disease (G20-G21)	594	0.8	14.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	470	0.6	11.2
15	Pneumonitis due to solids and liquids (J69)	380	0.5	9.1
...	All other causes (Residual)	12,688	16.1	303.2
Non-Hispanic, white, female, 75-79 years				
...	All causes	103,753	100.0	3,158.6
1	Malignant neoplasms (C00-C97)	29,053	28.0	884.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,603	19.9	627.2
3	Chronic lower respiratory diseases (J40-J47)	10,361	10.0	315.4
4	Cerebrovascular diseases (I60-I69)	5,961	5.7	181.5
5	Alzheimer's disease (G30)	3,874	3.7	117.9
6	Diabetes mellitus (E10-E14)	2,811	2.7	85.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,259	2.2	68.8
8	Influenza and pneumonia (J09-J18)	1,875	1.8	57.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,852	1.8	56.4
10	Septicemia (A40-A41)	1,704	1.6	51.9
11	Parkinson's disease (G20-G21)	1,230	1.2	37.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,063	1.0	32.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	770	0.7	23.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	718	0.7	21.9
15	Pneumonitis due to solids and liquids (J69)	634	0.6	19.3
...	All other causes (Residual)	18,985	18.3	578.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 80-84 years				
...	All causes	149,313	100.0	5,385.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,793	22.6	1,219.0
2	Malignant neoplasms (C00-C97)	30,486	20.4	1,099.7
3	Chronic lower respiratory diseases (J40-J47)	12,373	8.3	446.3
4	Cerebrovascular diseases (I60-I69)	10,275	6.9	370.6
5	Alzheimer's disease (G30)	8,616	5.8	310.8
6	Diabetes mellitus (E10-E14)	3,606	2.4	130.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,587	2.4	129.4
8	Influenza and pneumonia (J09-J18)	3,217	2.2	116.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,808	1.9	101.3
10	Septicemia (A40-A41)	2,298	1.5	82.9
11	Parkinson's disease (G20-G21)	1,987	1.3	71.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,814	1.2	65.4
13	Pneumonitis due to solids and liquids (J69)	1,085	0.7	39.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,023	0.7	36.9
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	770	0.5	27.8
...	All other causes (Residual)	31,575	21.1	1,138.9
Non-Hispanic, white, female, 85-89 years				
...	All causes	190,465	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,342	26.4	---
2	Malignant neoplasms (C00-C97)	26,296	13.8	---
3	Cerebrovascular diseases (I60-I69)	14,372	7.5	---
4	Alzheimer's disease (G30)	14,154	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	11,712	6.1	---
6	Influenza and pneumonia (J09-J18)	4,720	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,680	2.5	---
8	Diabetes mellitus (E10-E14)	3,945	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,455	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,926	1.5	---
11	Septicemia (A40-A41)	2,596	1.4	---
12	Parkinson's disease (G20-G21)	2,449	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,621	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,265	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,023	0.5	---
...	All other causes (Residual)	44,909	23.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 90-94 years				
...	All causes	161,623	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,596	30.1	---
2	Malignant neoplasms (C00-C97)	13,877	8.6	---
3	Alzheimer's disease (G30)	13,620	8.4	---
4	Cerebrovascular diseases (I60-I69)	12,606	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	7,125	4.4	---
6	Influenza and pneumonia (J09-J18)	4,514	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,045	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,951	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,775	1.7	---
10	Diabetes mellitus (E10-E14)	2,656	1.6	---
11	Septicemia (A40-A41)	1,731	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,476	0.9	---
13	Parkinson's disease (G20-G21)	1,421	0.9	---
14	Atherosclerosis (I70)	886	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	843	0.5	---
...	All other causes (Residual)	42,501	26.3	---
Non-Hispanic, white, female, 95-99 years				
...	All causes	72,811	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,682	32.5	---
2	Alzheimer's disease (G30)	6,312	8.7	---
3	Cerebrovascular diseases (I60-I69)	5,145	7.1	---
4	Malignant neoplasms (C00-C97)	4,024	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	2,373	3.3	---
6	Influenza and pneumonia (J09-J18)	2,357	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,873	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,566	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,123	1.5	---
10	Diabetes mellitus (E10-E14)	938	1.3	---
11	Pneumonitis due to solids and liquids (J69)	687	0.9	---
12	Septicemia (A40-A41)	669	0.9	---
13	Atherosclerosis (I70)	537	0.7	---
14	Parkinson's disease (G20-G21)	388	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	293	0.4	---
...	All other causes (Residual)	20,844	28.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 100 years and over				
...	All causes	16,604	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,647	34.0	---
2	Alzheimer's disease (G30)	1,326	8.0	---
3	Cerebrovascular diseases (I60-I69)	999	6.0	---
4	Influenza and pneumonia (J09-J18)	673	4.1	---
5	Malignant neoplasms (C00-C97)	568	3.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	438	2.6	---
7	Chronic lower respiratory diseases (J40-J47)	406	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	1.4	---
10	Atherosclerosis (I70)	179	1.1	---
11	Pneumonitis due to solids and liquids (J69)	161	1.0	---
12	Diabetes mellitus (E10-E14)	155	0.9	---
13	Septicemia (A40-A41)	121	0.7	---
14	Anemias (D50-D64)	62	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.3	---
...	All other causes (Residual)	5,176	31.2	---
Non-Hispanic, black, both sexes, all ages				
...	All causes	291,179	100.0	720.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,147	23.7	171.2
2	Malignant neoplasms (C00-C97)	66,565	22.9	164.8
3	Cerebrovascular diseases (I60-I69)	15,714	5.4	38.9
4	Diabetes mellitus (E10-E14)	12,836	4.4	31.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,451	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	9,264	3.2	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,129	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,123	2.8	20.1
9	Septicemia (A40-A41)	5,918	2.0	14.7
10	Alzheimer's disease (G30)	5,392	1.9	13.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,140	1.8	12.7
12	Influenza and pneumonia (J09-J18)	4,923	1.7	12.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,848	1.3	9.5
14	Certain conditions originating in the perinatal period (P00-P96)	3,767	1.3	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,739	0.9	6.8
...	All other causes (Residual)	57,223	19.7	141.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	989	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	271	27.4	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	13.0	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	10.5	4.2
4	Malignant neoplasms (C00-C97)	64	6.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.3	1.7
6	Chronic lower respiratory diseases (J40-J47)	30	3.0	1.2
7	Influenza and pneumonia (J09-J18)	20	2.0	0.8
8	Septicemia (A40-A41)	17	1.7	*
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	252	25.5	10.2
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	535	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	34.4	6.0
2	Malignant neoplasms (C00-C97)	60	11.2	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	8.4	1.5
4	Chronic lower respiratory diseases (J40-J47)	40	7.5	1.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	7.1	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.4	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Influenza and pneumonia (J09-J18)	6	1.1	*
10	Diabetes mellitus (E10-E14)	5	0.9	*
11	Septicemia (A40-A41)	4	0.7	*
12	Anemias (D50-D64)	3	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
...	All other causes (Residual)	109	20.4	3.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	620	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	26.1	5.2
2	Malignant neoplasms (C00-C97)	80	12.9	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	10.6	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	6.5	1.3
5	Chronic lower respiratory diseases (J40-J47)	34	5.5	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	5.2	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.7	0.7
8	Diabetes mellitus (E10-E14)	12	1.9	*
9	Anemias (D50-D64)	10	1.6	*
10	Influenza and pneumonia (J09-J18)	9	1.5	*
11	Cerebrovascular diseases (I60-I69)	8	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
14	Septicemia (A40-A41)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	126	20.3	4.0
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,264	100.0	68.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,022	45.1	30.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	20.6	14.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	7.0	4.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	4.1	2.8
5	Malignant neoplasms (C00-C97)	90	4.0	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.9	1.3
7	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.7
8	Anemias (D50-D64)	18	0.8	*
9	Diabetes mellitus (E10-E14)	16	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.7	*
11	Legal intervention (Y35,Y89.0)	14	0.6	*
12	Cerebrovascular diseases (I60-I69)	13	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.4	*
14	Septicemia (A40-A41)	8	0.4	*
15	Influenza and pneumonia (J09-J18)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	263	11.6	7.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,317	100.0	124.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,847	42.8	53.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	891	20.6	25.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	8.1	10.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	4.4	5.4
5	Malignant neoplasms (C00-C97)	154	3.6	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	78	1.8	2.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.2	1.5
8	Anemias (D50-D64)	47	1.1	1.4
9	Diabetes mellitus (E10-E14)	40	0.9	1.2
10	Chronic lower respiratory diseases (J40-J47)	39	0.9	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.1
12	Legal intervention (Y35,Y89.0)	30	0.7	0.9
13	Cerebrovascular diseases (I60-I69)	29	0.7	0.8
14	Influenza and pneumonia (J09-J18)	27	0.6	0.8
15	Septicemia (A40-A41)	25	0.6	0.7
...	All other causes (Residual)	480	11.1	13.8
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,183	100.0	145.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,335	31.9	46.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	845	20.2	29.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	7.5	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	6.9	10.0
5	Malignant neoplasms (C00-C97)	210	5.0	7.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	158	3.8	5.5
7	Diabetes mellitus (E10-E14)	73	1.7	2.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	68	1.6	2.4
9	Anemias (D50-D64)	58	1.4	2.0
10	Cerebrovascular diseases (I60-I69)	43	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	42	1.0	1.5
12	Septicemia (A40-A41)	41	1.0	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.9	1.3
14	Influenza and pneumonia (J09-J18)	35	0.8	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.8	1.1
...	All other causes (Residual)	604	14.4	20.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,765	100.0	172.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,071	22.5	38.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	847	17.8	30.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	567	11.9	20.5
4	Malignant neoplasms (C00-C97)	431	9.0	15.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	259	5.4	9.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	5.4	9.4
7	Diabetes mellitus (E10-E14)	127	2.7	4.6
8	Cerebrovascular diseases (I60-I69)	78	1.6	2.8
9	Chronic lower respiratory diseases (J40-J47)	67	1.4	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	65	1.4	2.3
11	Anemias (D50-D64)	55	1.2	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.1	2.0
13	Septicemia (A40-A41)	48	1.0	1.7
14	Influenza and pneumonia (J09-J18)	45	0.9	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.6	1.0
...	All other causes (Residual)	764	16.0	27.6
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	5,557	100.0	219.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,058	19.0	41.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	786	14.1	31.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	733	13.2	29.0
4	Malignant neoplasms (C00-C97)	651	11.7	25.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	313	5.6	12.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	3.8	8.4
7	Diabetes mellitus (E10-E14)	195	3.5	7.7
8	Cerebrovascular diseases (I60-I69)	161	2.9	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	1.5	3.3
10	Anemias (D50-D64)	69	1.2	2.7
11	Influenza and pneumonia (J09-J18)	66	1.2	2.6
12	Chronic lower respiratory diseases (J40-J47)	65	1.2	2.6
13	Septicemia (A40-A41)	61	1.1	2.4
14	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.0	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	52	0.9	2.1
...	All other causes (Residual)	996	17.9	39.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	7,945	100.0	296.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,707	21.5	63.7
2	Malignant neoplasms (C00-C97)	1,265	15.9	47.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	872	11.0	32.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	6.9	20.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	431	5.4	16.1
6	Diabetes mellitus (E10-E14)	310	3.9	11.6
7	Cerebrovascular diseases (I60-I69)	303	3.8	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	252	3.2	9.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.2	6.5
10	Septicemia (A40-A41)	151	1.9	5.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.7	5.2
12	Chronic lower respiratory diseases (J40-J47)	129	1.6	4.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.4	4.0
14	Influenza and pneumonia (J09-J18)	85	1.1	3.2
15	Anemias (D50-D64)	74	0.9	2.8
...	All other causes (Residual)	1,401	17.6	52.3
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	12,571	100.0	457.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,958	23.5	107.7
2	Malignant neoplasms (C00-C97)	2,700	21.5	98.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,088	8.7	39.6
4	Cerebrovascular diseases (I60-I69)	646	5.1	23.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	602	4.8	21.9
6	Diabetes mellitus (E10-E14)	562	4.5	20.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	400	3.2	14.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	289	2.3	10.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	258	2.1	9.4
10	Chronic lower respiratory diseases (J40-J47)	241	1.9	8.8
11	Septicemia (A40-A41)	220	1.8	8.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	1.6	7.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	189	1.5	6.9
14	Influenza and pneumonia (J09-J18)	145	1.2	5.3
15	Viral hepatitis (B15-B19)	69	0.5	2.5
...	All other causes (Residual)	2,004	15.9	73.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,951	100.0	726.2
1	Malignant neoplasms (C00-C97)	5,398	27.1	196.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,724	23.7	171.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,247	6.3	45.4
4	Cerebrovascular diseases (I60-I69)	1,001	5.0	36.4
5	Diabetes mellitus (E10-E14)	833	4.2	30.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	680	3.4	24.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	464	2.3	16.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	452	2.3	16.5
9	Chronic lower respiratory diseases (J40-J47)	440	2.2	16.0
10	Septicemia (A40-A41)	362	1.8	13.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	1.7	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	332	1.7	12.1
13	Influenza and pneumonia (J09-J18)	243	1.2	8.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	0.9	6.5
15	Viral hepatitis (B15-B19)	169	0.8	6.2
...	All other causes (Residual)	3,085	15.5	112.3
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	26,160	100.0	1,093.3
1	Malignant neoplasms (C00-C97)	7,893	30.2	329.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,388	24.4	267.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,253	4.8	52.4
4	Cerebrovascular diseases (I60-I69)	1,238	4.7	51.7
5	Diabetes mellitus (E10-E14)	1,208	4.6	50.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	658	2.5	27.5
7	Chronic lower respiratory diseases (J40-J47)	646	2.5	27.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	569	2.2	23.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	555	2.1	23.2
10	Septicemia (A40-A41)	516	2.0	21.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	418	1.6	17.5
12	Viral hepatitis (B15-B19)	331	1.3	13.8
13	Influenza and pneumonia (J09-J18)	328	1.3	13.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.8	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	0.5	5.7
...	All other causes (Residual)	3,805	14.5	159.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	28,823	100.0	1,529.0
1	Malignant neoplasms (C00-C97)	9,137	31.7	484.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,361	25.5	390.5
3	Diabetes mellitus (E10-E14)	1,454	5.0	77.1
4	Cerebrovascular diseases (I60-I69)	1,452	5.0	77.0
5	Chronic lower respiratory diseases (J40-J47)	908	3.2	48.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	813	2.8	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	813	2.8	43.1
8	Septicemia (A40-A41)	613	2.1	32.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	539	1.9	28.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	483	1.7	25.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	380	1.3	20.2
12	Influenza and pneumonia (J09-J18)	368	1.3	19.5
13	Viral hepatitis (B15-B19)	300	1.0	15.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	0.4	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.3	5.1
...	All other causes (Residual)	3,983	13.8	211.3
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	26,993	100.0	2,087.1
1	Malignant neoplasms (C00-C97)	8,840	32.7	683.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,766	25.1	523.1
3	Diabetes mellitus (E10-E14)	1,533	5.7	118.5
4	Cerebrovascular diseases (I60-I69)	1,395	5.2	107.9
5	Chronic lower respiratory diseases (J40-J47)	1,050	3.9	81.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	847	3.1	65.5
7	Septicemia (A40-A41)	614	2.3	47.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	526	1.9	40.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	1.7	36.4
10	Influenza and pneumonia (J09-J18)	378	1.4	29.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	305	1.1	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	194	0.7	15.0
13	Viral hepatitis (B15-B19)	118	0.4	9.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.4	8.6
15	Pneumonitis due to solids and liquids (J69)	106	0.4	8.2
...	All other causes (Residual)	3,739	13.9	289.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	27,488	100.0	3,012.9
1	Malignant neoplasms (C00-C97)	8,522	31.0	934.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,783	24.7	743.5
3	Cerebrovascular diseases (I60-I69)	1,639	6.0	179.6
4	Diabetes mellitus (E10-E14)	1,551	5.6	170.0
5	Chronic lower respiratory diseases (J40-J47)	1,309	4.8	143.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	920	3.3	100.8
7	Septicemia (A40-A41)	665	2.4	72.9
8	Influenza and pneumonia (J09-J18)	478	1.7	52.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	460	1.7	50.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	1.6	49.1
11	Alzheimer's disease (G30)	299	1.1	32.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	170	0.6	18.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.5	14.7
14	Pneumonitis due to solids and liquids (J69)	123	0.4	13.5
15	Parkinson's disease (G20-G21)	118	0.4	12.9
...	All other causes (Residual)	3,869	14.1	424.1
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	28,584	100.0	4,358.3
1	Malignant neoplasms (C00-C97)	7,586	26.5	1,156.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,281	25.5	1,110.2
3	Cerebrovascular diseases (I60-I69)	1,884	6.6	287.3
4	Diabetes mellitus (E10-E14)	1,506	5.3	229.6
5	Chronic lower respiratory diseases (J40-J47)	1,281	4.5	195.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	3.5	153.7
7	Septicemia (A40-A41)	692	2.4	105.5
8	Alzheimer's disease (G30)	638	2.2	97.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	555	1.9	84.6
10	Influenza and pneumonia (J09-J18)	549	1.9	83.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	1.4	61.8
12	Pneumonitis due to solids and liquids (J69)	181	0.6	27.6
13	Parkinson's disease (G20-G21)	168	0.6	25.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	148	0.5	22.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	115	0.4	17.5
...	All other causes (Residual)	4,587	16.0	699.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	28,981	100.0	6,613.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,487	25.8	1,708.4
2	Malignant neoplasms (C00-C97)	6,376	22.0	1,454.9
3	Cerebrovascular diseases (I60-I69)	1,984	6.8	452.7
4	Diabetes mellitus (E10-E14)	1,439	5.0	328.4
5	Chronic lower respiratory diseases (J40-J47)	1,188	4.1	271.1
6	Alzheimer's disease (G30)	1,044	3.6	238.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,019	3.5	232.5
8	Septicemia (A40-A41)	695	2.4	158.6
9	Influenza and pneumonia (J09-J18)	622	2.1	141.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	609	2.1	139.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	1.4	90.1
12	Pneumonitis due to solids and liquids (J69)	237	0.8	54.1
13	Parkinson's disease (G20-G21)	222	0.8	50.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.6	37.9
15	Atherosclerosis (I70)	90	0.3	20.5
...	All other causes (Residual)	5,408	18.7	1,234.0
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	26,541	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,302	27.5	---
2	Malignant neoplasms (C00-C97)	4,415	16.6	---
3	Cerebrovascular diseases (I60-I69)	1,901	7.2	---
4	Alzheimer's disease (G30)	1,386	5.2	---
5	Diabetes mellitus (E10-E14)	1,104	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	976	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	908	3.4	---
8	Influenza and pneumonia (J09-J18)	666	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	606	2.3	---
10	Septicemia (A40-A41)	599	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	1.1	---
12	Pneumonitis due to solids and liquids (J69)	254	1.0	---
13	Parkinson's disease (G20-G21)	172	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	146	0.6	---
15	Atherosclerosis (I70)	96	0.4	---
...	All other causes (Residual)	5,708	21.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	17,356	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,008	28.9	---
2	Malignant neoplasms (C00-C97)	2,018	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,224	7.1	---
4	Alzheimer's disease (G30)	1,092	6.3	---
5	Diabetes mellitus (E10-E14)	620	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	540	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	521	3.0	---
8	Influenza and pneumonia (J09-J18)	473	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	462	2.7	---
10	Septicemia (A40-A41)	343	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	1.1	---
12	Pneumonitis due to solids and liquids (J69)	163	0.9	---
13	Parkinson's disease (G20-G21)	95	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.4	---
15	Atherosclerosis (I70)	65	0.4	---
...	All other causes (Residual)	4,477	25.8	---
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	7,482	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,198	29.4	---
2	Alzheimer's disease (G30)	576	7.7	---
3	Malignant neoplasms (C00-C97)	552	7.4	---
4	Cerebrovascular diseases (I60-I69)	537	7.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	3.4	---
6	Influenza and pneumonia (J09-J18)	241	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	2.9	---
8	Diabetes mellitus (E10-E14)	198	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	185	2.5	---
10	Septicemia (A40-A41)	138	1.8	---
11	Pneumonitis due to solids and liquids (J69)	69	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	0.9	---
13	Anemias (D50-D64)	30	0.4	---
14	Parkinson's disease (G20-G21)	28	0.4	---
14	Atherosclerosis (I70)	28	0.4	---
...	All other causes (Residual)	2,169	29.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,278	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	771	33.8	---
2	Alzheimer's disease (G30)	182	8.0	---
3	Cerebrovascular diseases (I60-I69)	119	5.2	---
4	Malignant neoplasms (C00-C97)	112	4.9	---
5	Influenza and pneumonia (J09-J18)	80	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	3.5	---
7	Diabetes mellitus (E10-E14)	49	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.0	---
9	Chronic lower respiratory diseases (J40-J47)	43	1.9	---
10	Septicemia (A40-A41)	29	1.3	---
11	Pneumonitis due to solids and liquids (J69)	26	1.1	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	0.8	---
13	Atherosclerosis (I70)	9	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	8	0.4	---
15	Anemias (D50-D64)	6	0.3	---
...	All other causes (Residual)	701	30.8	---
Non-Hispanic, black, male, all ages				
...	All causes	148,344	100.0	768.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,586	24.0	184.3
2	Malignant neoplasms (C00-C97)	34,005	22.9	176.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,402	5.7	43.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,039	4.7	36.5
5	Cerebrovascular diseases (I60-I69)	6,975	4.7	36.1
6	Diabetes mellitus (E10-E14)	6,072	4.1	31.5
7	Chronic lower respiratory diseases (J40-J47)	4,826	3.3	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,776	2.5	19.6
9	Septicemia (A40-A41)	2,677	1.8	13.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,491	1.7	12.9
11	Influenza and pneumonia (J09-J18)	2,371	1.6	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,288	1.5	11.9
13	Certain conditions originating in the perinatal period (P00-P96)	2,131	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,861	1.3	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,770	1.2	9.2
...	All other causes (Residual)	26,074	17.6	135.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	563	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	29.1	13.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	9.1	4.1
4	Malignant neoplasms (C00-C97)	39	6.9	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.0	2.2
6	Chronic lower respiratory diseases (J40-J47)	24	4.3	1.9
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	2.0	*
9	Septicemia (A40-A41)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.9	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	124	22.0	9.9
Non-Hispanic, black, male, 5-9 years				
...	All causes	314	100.0	20.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	36.3	7.4
2	Malignant neoplasms (C00-C97)	35	11.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	9.6	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.0	1.6
5	Chronic lower respiratory diseases (J40-J47)	17	5.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.2	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.9	*
8	Anemias (D50-D64)	3	1.0	*
8	Diabetes mellitus (E10-E14)	3	1.0	*
10	Septicemia (A40-A41)	2	0.6	*
10	Cerebrovascular diseases (I60-I69)	2	0.6	*
10	Influenza and pneumonia (J09-J18)	2	0.6	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
14	Hernia (K40-K46)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	62	19.7	4.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 10-14 years				
...	All causes	393	100.0	24.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	29.0	7.2
2	Malignant neoplasms (C00-C97)	45	11.5	2.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	11.5	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	7.9	1.9
5	Chronic lower respiratory diseases (J40-J47)	23	5.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	4.1	*
8	Anemias (D50-D64)	5	1.3	*
8	Diabetes mellitus (E10-E14)	5	1.3	*
8	Influenza and pneumonia (J09-J18)	5	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
11	Cerebrovascular diseases (I60-I69)	4	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.8	*
14	Septicemia (A40-A41)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	73	18.6	4.6
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,731	100.0	102.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	917	53.0	54.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	347	20.0	20.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	6.9	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	3.6	3.7
5	Malignant neoplasms (C00-C97)	53	3.1	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.4	1.5
7	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
7	Legal intervention (Y35,Y89.0)	14	0.8	*
9	Anemias (D50-D64)	11	0.6	*
10	Diabetes mellitus (E10-E14)	7	0.4	*
11	Cerebrovascular diseases (I60-I69)	6	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.2	*
12	Aortic aneurysm and dissection (I71)	4	0.2	*
14	Septicemia (A40-A41)	3	0.2	*
14	Influenza and pneumonia (J09-J18)	3	0.2	*
...	All other causes (Residual)	141	8.1	8.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,297	100.0	189.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,669	50.6	96.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	639	19.4	36.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	292	8.9	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	3.8	7.2
5	Malignant neoplasms (C00-C97)	89	2.7	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.7	3.3
7	Chronic lower respiratory diseases (J40-J47)	32	1.0	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	1.7
9	Legal intervention (Y35,Y89.0)	28	0.8	1.6
10	Anemias (D50-D64)	23	0.7	1.3
11	Diabetes mellitus (E10-E14)	19	0.6	*
12	Cerebrovascular diseases (I60-I69)	17	0.5	*
12	Influenza and pneumonia (J09-J18)	17	0.5	*
14	Septicemia (A40-A41)	10	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	241	7.3	13.9
Non-Hispanic, black, male, 25-29 years				
...	All causes	2,985	100.0	213.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,204	40.3	86.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	620	20.8	44.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	246	8.2	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	6.3	13.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	108	3.6	7.7
6	Malignant neoplasms (C00-C97)	95	3.2	6.8
7	Diabetes mellitus (E10-E14)	40	1.3	2.9
8	Anemias (D50-D64)	34	1.1	2.4
9	Chronic lower respiratory diseases (J40-J47)	27	0.9	1.9
10	Legal intervention (Y35,Y89.0)	26	0.9	1.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.8	1.7
12	Cerebrovascular diseases (I60-I69)	22	0.7	1.6
13	Influenza and pneumonia (J09-J18)	20	0.7	1.4
14	Septicemia (A40-A41)	19	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.5	*
...	All other causes (Residual)	280	9.4	20.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,170	100.0	240.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	949	29.9	72.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	626	19.7	47.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	11.5	27.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	221	7.0	16.8
5	Malignant neoplasms (C00-C97)	168	5.3	12.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	156	4.9	11.9
7	Diabetes mellitus (E10-E14)	60	1.9	4.6
8	Cerebrovascular diseases (I60-I69)	48	1.5	3.6
9	Chronic lower respiratory diseases (J40-J47)	41	1.3	3.1
10	Septicemia (A40-A41)	29	0.9	2.2
11	Anemias (D50-D64)	25	0.8	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.8	1.9
13	Influenza and pneumonia (J09-J18)	21	0.7	1.6
14	Aortic aneurysm and dissection (I71)	18	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
...	All other causes (Residual)	400	12.6	30.4
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,339	100.0	280.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	655	19.6	55.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	654	19.6	55.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	16.9	47.4
4	Malignant neoplasms (C00-C97)	221	6.6	18.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	5.0	14.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	4.3	12.1
7	Diabetes mellitus (E10-E14)	112	3.4	9.4
8	Cerebrovascular diseases (I60-I69)	93	2.8	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.3	3.6
10	Influenza and pneumonia (J09-J18)	33	1.0	2.8
11	Chronic lower respiratory diseases (J40-J47)	31	0.9	2.6
12	Septicemia (A40-A41)	29	0.9	2.4
12	Anemias (D50-D64)	29	0.9	2.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.9	2.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.8	2.4
...	All other causes (Residual)	507	15.2	42.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,566	100.0	361.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,122	24.6	88.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	599	13.1	47.5
3	Malignant neoplasms (C00-C97)	467	10.2	37.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	452	9.9	35.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	241	5.3	19.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	189	4.1	15.0
7	Diabetes mellitus (E10-E14)	188	4.1	14.9
8	Cerebrovascular diseases (I60-I69)	162	3.5	12.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.1	7.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	88	1.9	7.0
11	Septicemia (A40-A41)	68	1.5	5.4
11	Chronic lower respiratory diseases (J40-J47)	68	1.5	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.3	4.6
14	Aortic aneurysm and dissection (I71)	41	0.9	3.2
15	Influenza and pneumonia (J09-J18)	40	0.9	3.2
...	All other causes (Residual)	685	15.0	54.3
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,090	100.0	548.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,856	26.2	143.5
2	Malignant neoplasms (C00-C97)	1,180	16.6	91.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	719	10.1	55.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	374	5.3	28.9
5	Cerebrovascular diseases (I60-I69)	357	5.0	27.6
6	Diabetes mellitus (E10-E14)	322	4.5	24.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	318	4.5	24.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	2.3	12.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	144	2.0	11.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	2.0	11.1
11	Septicemia (A40-A41)	116	1.6	9.0
12	Chronic lower respiratory diseases (J40-J47)	113	1.6	8.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.6	8.5
14	Influenza and pneumonia (J09-J18)	82	1.2	6.3
15	Viral hepatitis (B15-B19)	50	0.7	3.9
...	All other causes (Residual)	1,045	14.7	80.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,487	100.0	892.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,024	26.3	235.0
2	Malignant neoplasms (C00-C97)	2,687	23.4	208.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	862	7.5	67.0
4	Cerebrovascular diseases (I60-I69)	533	4.6	41.4
5	Diabetes mellitus (E10-E14)	484	4.2	37.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	455	4.0	35.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	287	2.5	22.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	2.4	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.1	19.1
10	Chronic lower respiratory diseases (J40-J47)	222	1.9	17.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.7	15.2
12	Septicemia (A40-A41)	185	1.6	14.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	137	1.2	10.6
14	Influenza and pneumonia (J09-J18)	136	1.2	10.6
15	Viral hepatitis (B15-B19)	117	1.0	9.1
...	All other causes (Residual)	1,645	14.3	127.8
Non-Hispanic, black, male, 55-59 years				
...	All causes	15,358	100.0	1,395.3
1	Malignant neoplasms (C00-C97)	4,230	27.5	384.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,093	26.7	371.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	5.8	81.6
4	Cerebrovascular diseases (I60-I69)	717	4.7	65.1
5	Diabetes mellitus (E10-E14)	665	4.3	60.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	403	2.6	36.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	379	2.5	34.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	358	2.3	32.5
9	Chronic lower respiratory diseases (J40-J47)	326	2.1	29.6
10	Septicemia (A40-A41)	260	1.7	23.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	257	1.7	23.3
12	Viral hepatitis (B15-B19)	213	1.4	19.4
13	Influenza and pneumonia (J09-J18)	197	1.3	17.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	186	1.2	16.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	0.7	9.4
...	All other causes (Residual)	2,072	13.5	188.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 60-64 years				
...	All causes	16,885	100.0	1,998.5
1	Malignant neoplasms (C00-C97)	5,124	30.3	606.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,609	27.3	545.5
3	Diabetes mellitus (E10-E14)	812	4.8	96.1
4	Cerebrovascular diseases (I60-I69)	804	4.8	95.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	3.5	69.1
6	Chronic lower respiratory diseases (J40-J47)	489	2.9	57.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	2.7	53.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	393	2.3	46.5
9	Septicemia (A40-A41)	319	1.9	37.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	275	1.6	32.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	265	1.6	31.4
12	Influenza and pneumonia (J09-J18)	212	1.3	25.1
13	Viral hepatitis (B15-B19)	208	1.2	24.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	0.6	12.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	0.4	7.2
...	All other causes (Residual)	2,179	12.9	257.9
Non-Hispanic, black, male, 65-69 years				
...	All causes	15,225	100.0	2,698.9
1	Malignant neoplasms (C00-C97)	4,879	32.0	864.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,043	26.6	716.7
3	Diabetes mellitus (E10-E14)	817	5.4	144.8
4	Cerebrovascular diseases (I60-I69)	745	4.9	132.1
5	Chronic lower respiratory diseases (J40-J47)	597	3.9	105.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	446	2.9	79.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	2.3	62.8
8	Septicemia (A40-A41)	324	2.1	57.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	248	1.6	44.0
10	Influenza and pneumonia (J09-J18)	212	1.4	37.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	207	1.4	36.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	151	1.0	26.8
13	Viral hepatitis (B15-B19)	71	0.5	12.6
14	Pneumonitis due to solids and liquids (J69)	67	0.4	11.9
15	Aortic aneurysm and dissection (I71)	62	0.4	11.0
...	All other causes (Residual)	2,002	13.1	354.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 70-74 years				
...	All causes	14,506	100.0	3,809.7
1	Malignant neoplasms (C00-C97)	4,513	31.1	1,185.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,716	25.6	975.9
3	Cerebrovascular diseases (I60-I69)	853	5.9	224.0
4	Diabetes mellitus (E10-E14)	776	5.3	203.8
5	Chronic lower respiratory diseases (J40-J47)	740	5.1	194.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	447	3.1	117.4
7	Septicemia (A40-A41)	285	2.0	74.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	1.9	73.0
9	Influenza and pneumonia (J09-J18)	251	1.7	65.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	234	1.6	61.5
11	Alzheimer's disease (G30)	126	0.9	33.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	113	0.8	29.7
13	Pneumonitis due to solids and liquids (J69)	85	0.6	22.3
14	Parkinson's disease (G20-G21)	81	0.6	21.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	74	0.5	19.4
...	All other causes (Residual)	1,934	13.3	507.9
Non-Hispanic, black, male, 75-79 years				
...	All causes	14,124	100.0	5,509.6
1	Malignant neoplasms (C00-C97)	3,893	27.6	1,518.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,741	26.5	1,459.3
3	Cerebrovascular diseases (I60-I69)	865	6.1	337.4
4	Chronic lower respiratory diseases (J40-J47)	702	5.0	273.8
5	Diabetes mellitus (E10-E14)	652	4.6	254.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	3.3	183.0
7	Septicemia (A40-A41)	311	2.2	121.3
8	Influenza and pneumonia (J09-J18)	288	2.0	112.3
9	Alzheimer's disease (G30)	252	1.8	98.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.7	95.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	1.7	94.8
12	Parkinson's disease (G20-G21)	116	0.8	45.3
13	Pneumonitis due to solids and liquids (J69)	112	0.8	43.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	28.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.5	26.1
...	All other causes (Residual)	2,095	14.8	817.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,732	100.0	8,286.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,252	25.5	2,116.4
2	Malignant neoplasms (C00-C97)	3,180	25.0	2,069.6
3	Cerebrovascular diseases (I60-I69)	755	5.9	491.4
4	Chronic lower respiratory diseases (J40-J47)	588	4.6	382.7
5	Diabetes mellitus (E10-E14)	577	4.5	375.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	3.4	279.8
7	Alzheimer's disease (G30)	381	3.0	248.0
8	Influenza and pneumonia (J09-J18)	280	2.2	182.2
9	Septicemia (A40-A41)	274	2.2	178.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.9	157.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	1.7	140.6
12	Parkinson's disease (G20-G21)	132	1.0	85.9
13	Pneumonitis due to solids and liquids (J69)	124	1.0	80.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.6	48.2
15	Atherosclerosis (I70)	45	0.4	29.3
...	All other causes (Residual)	2,182	17.1	1,420.1
Non-Hispanic, black, male, 85-89 years				
...	All causes	9,691	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,581	26.6	---
2	Malignant neoplasms (C00-C97)	2,061	21.3	---
3	Cerebrovascular diseases (I60-I69)	571	5.9	---
4	Chronic lower respiratory diseases (J40-J47)	455	4.7	---
5	Alzheimer's disease (G30)	375	3.9	---
6	Diabetes mellitus (E10-E14)	334	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	3.4	---
8	Influenza and pneumonia (J09-J18)	287	3.0	---
9	Septicemia (A40-A41)	225	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	195	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	1.5	---
12	Pneumonitis due to solids and liquids (J69)	131	1.4	---
13	Parkinson's disease (G20-G21)	97	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.6	---
15	Atherosclerosis (I70)	37	0.4	---
...	All other causes (Residual)	1,810	18.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 90-94 years				
...	All causes	5,113	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,450	28.4	---
2	Malignant neoplasms (C00-C97)	823	16.1	---
3	Cerebrovascular diseases (I60-I69)	291	5.7	---
4	Chronic lower respiratory diseases (J40-J47)	236	4.6	---
5	Alzheimer's disease (G30)	217	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	3.3	---
7	Influenza and pneumonia (J09-J18)	157	3.1	---
8	Diabetes mellitus (E10-E14)	152	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	2.4	---
10	Septicemia (A40-A41)	116	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	1.4	---
12	Pneumonitis due to solids and liquids (J69)	54	1.1	---
13	Parkinson's disease (G20-G21)	36	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	18	0.4	---
...	All other causes (Residual)	1,177	23.0	---
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,623	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	490	30.2	---
2	Malignant neoplasms (C00-C97)	193	11.9	---
3	Cerebrovascular diseases (I60-I69)	91	5.6	---
4	Alzheimer's disease (G30)	88	5.4	---
5	Influenza and pneumonia (J09-J18)	71	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	62	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	3.5	---
8	Diabetes mellitus (E10-E14)	39	2.4	---
9	Septicemia (A40-A41)	36	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.0	---
12	Pneumonitis due to solids and liquids (J69)	12	0.7	---
13	Parkinson's disease (G20-G21)	10	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	---
14	Atherosclerosis (I70)	9	0.6	---
...	All other causes (Residual)	405	25.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 100 years and over				
...	All causes	339	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	31.6	---
2	Malignant neoplasms (C00-C97)	24	7.1	---
2	Alzheimer's disease (G30)	24	7.1	---
4	Chronic lower respiratory diseases (J40-J47)	15	4.4	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	4.1	---
5	Influenza and pneumonia (J09-J18)	14	4.1	---
7	Cerebrovascular diseases (I60-I69)	12	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	3.5	---
9	Pneumonitis due to solids and liquids (J69)	8	2.4	---
10	Diabetes mellitus (E10-E14)	7	2.1	---
11	Septicemia (A40-A41)	6	1.8	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	1.2	---
13	Nutritional deficiencies (E40-E64)	2	0.6	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	---
15	Anemias (D50-D64)	1	0.3	---
15	Parkinson's disease (G20-G21)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Diseases of appendix (K35-K38)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	---
...	All other causes (Residual)	79	23.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	142,835	100.0	677.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,561	23.5	159.2
2	Malignant neoplasms (C00-C97)	32,560	22.8	154.4
3	Cerebrovascular diseases (I60-I69)	8,739	6.1	41.4
4	Diabetes mellitus (E10-E14)	6,764	4.7	32.1
5	Chronic lower respiratory diseases (J40-J47)	4,438	3.1	21.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,353	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,049	2.8	19.2
8	Alzheimer's disease (G30)	3,861	2.7	18.3
9	Septicemia (A40-A41)	3,241	2.3	15.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,852	2.0	13.5
11	Influenza and pneumonia (J09-J18)	2,552	1.8	12.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,636	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,357	1.0	6.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	969	0.7	4.6
...	All other causes (Residual)	30,819	21.6	146.1
Non-Hispanic, black, female, 1-4 years				
...	All causes	426	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	25.1	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	12.4	4.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	12.0	4.2
4	Malignant neoplasms (C00-C97)	25	5.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Septicemia (A40-A41)	9	2.1	*
6	Cerebrovascular diseases (I60-I69)	9	2.1	*
8	Influenza and pneumonia (J09-J18)	8	1.9	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	119	27.9	9.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	221	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	31.7	4.7
2	Malignant neoplasms (C00-C97)	25	11.3	1.7
3	Chronic lower respiratory diseases (J40-J47)	23	10.4	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	6.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.0	*
7	Cerebrovascular diseases (I60-I69)	4	1.8	*
7	Influenza and pneumonia (J09-J18)	4	1.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.4	*
10	Septicemia (A40-A41)	2	0.9	*
10	Diabetes mellitus (E10-E14)	2	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
...	All other causes (Residual)	44	19.9	2.9
Non-Hispanic, black, female, 10-14 years				
...	All causes	227	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	21.1	3.1
2	Malignant neoplasms (C00-C97)	35	15.4	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	9.3	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	7.0	*
5	Chronic lower respiratory diseases (J40-J47)	11	4.8	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	4.0	*
7	Diabetes mellitus (E10-E14)	7	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	3.1	*
9	Anemias (D50-D64)	5	2.2	*
10	Cerebrovascular diseases (I60-I69)	4	1.8	*
10	Influenza and pneumonia (J09-J18)	4	1.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
14	Septicemia (A40-A41)	1	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	52	22.9	3.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	533	100.0	32.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	22.3	7.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	19.7	6.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	7.5	2.4
4	Malignant neoplasms (C00-C97)	37	6.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	5.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.6	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	16	3.0	*
8	Diabetes mellitus (E10-E14)	9	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.7	*
10	Anemias (D50-D64)	7	1.3	*
10	Cerebrovascular diseases (I60-I69)	7	1.3	*
12	Septicemia (A40-A41)	5	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Influenza and pneumonia (J09-J18)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	113	21.2	6.9
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,020	100.0	58.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	24.7	14.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	17.5	10.3
3	Malignant neoplasms (C00-C97)	65	6.4	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	6.3	3.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	5.8	3.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	5.0	2.9
7	Anemias (D50-D64)	24	2.4	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	21	2.1	1.2
8	Diabetes mellitus (E10-E14)	21	2.1	1.2
10	Septicemia (A40-A41)	15	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.5	*
12	Cerebrovascular diseases (I60-I69)	12	1.2	*
13	Influenza and pneumonia (J09-J18)	10	1.0	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
...	All other causes (Residual)	216	21.2	12.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,198	100.0	80.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	18.8	15.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	10.9	8.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	10.6	8.6
4	Malignant neoplasms (C00-C97)	115	9.6	7.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	68	5.7	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	50	4.2	3.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	3.4	2.8
8	Diabetes mellitus (E10-E14)	33	2.8	2.2
9	Anemias (D50-D64)	24	2.0	1.6
10	Septicemia (A40-A41)	22	1.8	1.5
11	Cerebrovascular diseases (I60-I69)	21	1.8	1.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.8	1.4
13	Influenza and pneumonia (J09-J18)	15	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (Residual)	282	23.5	19.0
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,595	100.0	110.0
1	Malignant neoplasms (C00-C97)	263	16.5	18.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	13.9	15.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	12.7	13.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	7.6	8.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	103	6.5	7.1
6	Diabetes mellitus (E10-E14)	67	4.2	4.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	65	4.1	4.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.4	2.6
9	Anemias (D50-D64)	30	1.9	2.1
9	Cerebrovascular diseases (I60-I69)	30	1.9	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.8	2.0
12	Chronic lower respiratory diseases (J40-J47)	26	1.6	1.8
13	Influenza and pneumonia (J09-J18)	24	1.5	1.7
14	Septicemia (A40-A41)	19	1.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.1	*
...	All other causes (Residual)	339	21.3	23.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,218	100.0	165.7
1	Malignant neoplasms (C00-C97)	430	19.4	32.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	404	18.2	30.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	10.0	16.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	169	7.6	12.6
5	Diabetes mellitus (E10-E14)	83	3.7	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	3.5	5.8
7	Cerebrovascular diseases (I60-I69)	68	3.1	5.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	56	2.5	4.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	2.0	3.4
10	Anemias (D50-D64)	40	1.8	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.8	3.0
12	Chronic lower respiratory diseases (J40-J47)	34	1.5	2.5
13	Influenza and pneumonia (J09-J18)	33	1.5	2.5
14	Septicemia (A40-A41)	32	1.4	2.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	24	1.1	1.8
...	All other causes (Residual)	460	20.7	34.4
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,379	100.0	238.7
1	Malignant neoplasms (C00-C97)	798	23.6	56.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	585	17.3	41.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	8.1	19.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	190	5.6	13.4
5	Cerebrovascular diseases (I60-I69)	141	4.2	10.0
6	Diabetes mellitus (E10-E14)	122	3.6	8.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	2.8	6.6
8	Septicemia (A40-A41)	83	2.5	5.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.2	5.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	1.9	4.4
11	Chronic lower respiratory diseases (J40-J47)	61	1.8	4.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.5	3.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	50	1.5	3.5
14	Influenza and pneumonia (J09-J18)	45	1.3	3.2
15	Anemias (D50-D64)	39	1.2	2.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.2	2.8
...	All other causes (Residual)	671	19.9	47.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,481	100.0	377.2
1	Malignant neoplasms (C00-C97)	1,520	27.7	104.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,102	20.1	75.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	6.7	25.4
4	Cerebrovascular diseases (I60-I69)	289	5.3	19.9
5	Diabetes mellitus (E10-E14)	240	4.4	16.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	228	4.2	15.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.4	8.9
8	Chronic lower respiratory diseases (J40-J47)	128	2.3	8.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	114	2.1	7.8
10	Septicemia (A40-A41)	104	1.9	7.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.6	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	1.5	5.6
13	Influenza and pneumonia (J09-J18)	63	1.1	4.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	0.8	3.1
15	Anemias (D50-D64)	32	0.6	2.2
...	All other causes (Residual)	946	17.3	65.1
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,464	100.0	579.5
1	Malignant neoplasms (C00-C97)	2,711	32.0	185.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,700	20.1	116.4
3	Cerebrovascular diseases (I60-I69)	468	5.5	32.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	4.5	26.4
5	Diabetes mellitus (E10-E14)	349	4.1	23.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	225	2.7	15.4
7	Chronic lower respiratory diseases (J40-J47)	218	2.6	14.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.6	14.9
9	Septicemia (A40-A41)	177	2.1	12.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.9	11.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	1.6	9.3
12	Influenza and pneumonia (J09-J18)	107	1.3	7.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	0.9	4.9
14	Viral hepatitis (B15-B19)	52	0.6	3.6
15	Anemias (D50-D64)	41	0.5	2.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.5	2.8
...	All other causes (Residual)	1,399	16.5	95.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	10,802	100.0	836.0
1	Malignant neoplasms (C00-C97)	3,663	33.9	283.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,295	21.2	177.6
3	Diabetes mellitus (E10-E14)	543	5.0	42.0
4	Cerebrovascular diseases (I60-I69)	521	4.8	40.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	3.3	27.5
6	Chronic lower respiratory diseases (J40-J47)	320	3.0	24.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	2.8	23.2
8	Septicemia (A40-A41)	256	2.4	19.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	176	1.6	13.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	166	1.5	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.5	12.5
12	Influenza and pneumonia (J09-J18)	131	1.2	10.1
13	Viral hepatitis (B15-B19)	118	1.1	9.1
14	Anemias (D50-D64)	49	0.5	3.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	3.5
...	All other causes (Residual)	1,703	15.8	131.8
Non-Hispanic, black, female, 60-64 years				
...	All causes	11,938	100.0	1,147.6
1	Malignant neoplasms (C00-C97)	4,013	33.6	385.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,752	23.1	264.6
3	Cerebrovascular diseases (I60-I69)	648	5.4	62.3
4	Diabetes mellitus (E10-E14)	642	5.4	61.7
5	Chronic lower respiratory diseases (J40-J47)	419	3.5	40.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	3.0	35.0
7	Septicemia (A40-A41)	294	2.5	28.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	1.9	22.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.7	20.0
10	Influenza and pneumonia (J09-J18)	156	1.3	15.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.2	14.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	115	1.0	11.1
13	Viral hepatitis (B15-B19)	92	0.8	8.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	4.4
15	Pneumonitis due to solids and liquids (J69)	39	0.3	3.7
...	All other causes (Residual)	1,775	14.9	170.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	11,768	100.0	1,613.8
1	Malignant neoplasms (C00-C97)	3,961	33.7	543.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,723	23.1	373.4
3	Diabetes mellitus (E10-E14)	716	6.1	98.2
4	Cerebrovascular diseases (I60-I69)	650	5.5	89.1
5	Chronic lower respiratory diseases (J40-J47)	453	3.8	62.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	401	3.4	55.0
7	Septicemia (A40-A41)	290	2.5	39.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	1.9	30.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.5	23.6
10	Influenza and pneumonia (J09-J18)	166	1.4	22.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.8	13.4
12	Alzheimer's disease (G30)	69	0.6	9.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	8.0
14	Viral hepatitis (B15-B19)	47	0.4	6.4
15	Anemias (D50-D64)	45	0.4	6.2
...	All other causes (Residual)	1,696	14.4	232.6
Non-Hispanic, black, female, 70-74 years				
...	All causes	12,982	100.0	2,442.2
1	Malignant neoplasms (C00-C97)	4,009	30.9	754.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,067	23.6	577.0
3	Cerebrovascular diseases (I60-I69)	786	6.1	147.9
4	Diabetes mellitus (E10-E14)	775	6.0	145.8
5	Chronic lower respiratory diseases (J40-J47)	569	4.4	107.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	473	3.6	89.0
7	Septicemia (A40-A41)	380	2.9	71.5
8	Influenza and pneumonia (J09-J18)	227	1.7	42.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	1.7	42.5
10	Alzheimer's disease (G30)	173	1.3	32.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	1.3	32.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.5	12.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.4	10.7
14	Aortic aneurysm and dissection (I71)	50	0.4	9.4
15	Pneumonitis due to solids and liquids (J69)	38	0.3	7.1
...	All other causes (Residual)	1,915	14.8	360.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,460	100.0	3,619.5
1	Malignant neoplasms (C00-C97)	3,693	25.5	924.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,540	24.5	886.1
3	Cerebrovascular diseases (I60-I69)	1,019	7.0	255.1
4	Diabetes mellitus (E10-E14)	854	5.9	213.8
5	Chronic lower respiratory diseases (J40-J47)	579	4.0	144.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	539	3.7	134.9
7	Alzheimer's disease (G30)	386	2.7	96.6
8	Septicemia (A40-A41)	381	2.6	95.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	311	2.2	77.8
10	Influenza and pneumonia (J09-J18)	261	1.8	65.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.1	40.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	18.5
13	Pneumonitis due to solids and liquids (J69)	69	0.5	17.3
14	Parkinson's disease (G20-G21)	52	0.4	13.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	48	0.3	12.0
...	All other causes (Residual)	2,492	17.2	623.8
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,249	100.0	5,709.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,235	26.1	1,488.2
2	Malignant neoplasms (C00-C97)	3,196	19.7	1,123.1
3	Cerebrovascular diseases (I60-I69)	1,229	7.6	431.9
4	Diabetes mellitus (E10-E14)	862	5.3	302.9
5	Alzheimer's disease (G30)	663	4.1	233.0
6	Chronic lower respiratory diseases (J40-J47)	600	3.7	210.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	589	3.6	207.0
8	Septicemia (A40-A41)	421	2.6	147.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	367	2.3	129.0
10	Influenza and pneumonia (J09-J18)	342	2.1	120.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.1	62.9
12	Pneumonitis due to solids and liquids (J69)	113	0.7	39.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.6	32.3
14	Parkinson's disease (G20-G21)	90	0.6	31.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	51	0.3	17.9
...	All other causes (Residual)	3,220	19.8	1,131.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	16,850	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,721	28.0	---
2	Malignant neoplasms (C00-C97)	2,354	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,330	7.9	---
4	Alzheimer's disease (G30)	1,011	6.0	---
5	Diabetes mellitus (E10-E14)	770	4.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	579	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	521	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	411	2.4	---
9	Influenza and pneumonia (J09-J18)	379	2.2	---
10	Septicemia (A40-A41)	374	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	0.9	---
12	Pneumonitis due to solids and liquids (J69)	123	0.7	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	89	0.5	---
14	Parkinson's disease (G20-G21)	75	0.4	---
15	Atherosclerosis (I70)	59	0.4	---
...	All other causes (Residual)	3,898	23.1	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	12,243	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,558	29.1	---
2	Malignant neoplasms (C00-C97)	1,195	9.8	---
3	Cerebrovascular diseases (I60-I69)	933	7.6	---
4	Alzheimer's disease (G30)	875	7.1	---
5	Diabetes mellitus (E10-E14)	468	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	372	3.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	340	2.8	---
8	Influenza and pneumonia (J09-J18)	316	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	285	2.3	---
10	Septicemia (A40-A41)	227	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	0.9	---
12	Pneumonitis due to solids and liquids (J69)	109	0.9	---
13	Parkinson's disease (G20-G21)	59	0.5	---
14	Atherosclerosis (I70)	48	0.4	---
15	Anemias (D50-D64)	46	0.4	---
...	All other causes (Residual)	3,298	26.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,859	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,708	29.2	---
2	Alzheimer's disease (G30)	488	8.3	---
3	Cerebrovascular diseases (I60-I69)	446	7.6	---
4	Malignant neoplasms (C00-C97)	359	6.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	218	3.7	---
6	Influenza and pneumonia (J09-J18)	170	2.9	---
7	Diabetes mellitus (E10-E14)	159	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	123	2.1	---
10	Septicemia (A40-A41)	102	1.7	---
11	Pneumonitis due to solids and liquids (J69)	57	1.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	0.8	---
13	Anemias (D50-D64)	26	0.4	---
14	Atherosclerosis (I70)	19	0.3	---
15	Parkinson's disease (G20-G21)	18	0.3	---
...	All other causes (Residual)	1,759	30.0	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	1,939	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	664	34.2	---
2	Alzheimer's disease (G30)	158	8.1	---
3	Cerebrovascular diseases (I60-I69)	107	5.5	---
4	Malignant neoplasms (C00-C97)	88	4.5	---
5	Influenza and pneumonia (J09-J18)	66	3.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	3.4	---
7	Diabetes mellitus (E10-E14)	42	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.7	---
9	Chronic lower respiratory diseases (J40-J47)	28	1.4	---
10	Septicemia (A40-A41)	23	1.2	---
11	Pneumonitis due to solids and liquids (J69)	18	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.9	---
13	Atherosclerosis (I70)	9	0.5	---
14	Anemias (D50-D64)	5	0.3	---
15	Aortic aneurysm and dissection (I71)	4	0.2	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.2	---
...	All other causes (Residual)	608	31.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012