

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	163,241	100.0	301.9
1	Malignant neoplasms (C00-C97)	35,147	21.5	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,243	20.4	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,015	7.4	22.2
4	Cerebrovascular diseases (I60-I69)	8,127	5.0	15.0
5	Diabetes mellitus (E10-E14)	7,632	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,141	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	4,827	3.0	8.9
8	Alzheimer's disease (G30)	4,127	2.5	7.6
9	Influenza and pneumonia (J09-J18)	3,592	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,083	1.9	5.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,865	1.8	5.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,571	1.6	4.8
13	Septicemia (A40-A41)	2,435	1.5	4.5
14	Certain conditions originating in the perinatal period (P00-P96)	2,428	1.5	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,092	1.3	3.9
...	All other causes (Residual)	33,916	20.8	62.7
Hispanic, both sexes, 1-4 years				
...	All causes	854	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	32.6	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	14.3	3.0
3	Malignant neoplasms (C00-C97)	90	10.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.7	0.6
6	Influenza and pneumonia (J09-J18)	19	2.2	*
7	Septicemia (A40-A41)	9	1.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.1	*
11	Anemias (D50-D64)	5	0.6	*
11	Meningitis (G00,G03)	5	0.6	*
13	Cerebrovascular diseases (I60-I69)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	213	24.9	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	467	100.0	9.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	28.9	2.6
2	Malignant neoplasms (C00-C97)	96	20.6	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	9.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.9	0.4
5	Chronic lower respiratory diseases (J40-J47)	13	2.8	*
6	Influenza and pneumonia (J09-J18)	12	2.6	*
7	Cerebrovascular diseases (I60-I69)	10	2.1	*
8	Septicemia (A40-A41)	7	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.5	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Anemias (D50-D64)	2	0.4	*
12	Diseases of appendix (K35-K38)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	101	21.6	2.0
Hispanic, both sexes, 10-14 years				
...	All causes	606	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	23.8	3.0
2	Malignant neoplasms (C00-C97)	113	18.6	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	11.7	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.9	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.0	0.5
7	Cerebrovascular diseases (I60-I69)	16	2.6	*
8	Influenza and pneumonia (J09-J18)	10	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.8	*
12	Septicemia (A40-A41)	3	0.5	*
12	Diabetes mellitus (E10-E14)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	126	20.8	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,646	100.0	35.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	636	38.6	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	315	19.1	6.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	233	14.2	5.1
4	Malignant neoplasms (C00-C97)	136	8.3	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	2.7	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.9	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
8	Septicemia (A40-A41)	9	0.5	*
8	Cerebrovascular diseases (I60-I69)	9	0.5	*
8	Influenza and pneumonia (J09-J18)	9	0.5	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.5	*
12	Diabetes mellitus (E10-E14)	5	0.3	*
13	Legal intervention (Y35,Y89.0)	4	0.2	*
14	Anemias (D50-D64)	3	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.2	*
...	All other causes (Residual)	183	11.1	4.0
Hispanic, both sexes, 20-24 years				
...	All causes	3,011	100.0	64.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,308	43.4	28.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	534	17.7	11.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	13.4	8.6
4	Malignant neoplasms (C00-C97)	203	6.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	2.5	1.6
6	Cerebrovascular diseases (I60-I69)	23	0.8	0.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.8	0.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	19	0.6	*
9	Influenza and pneumonia (J09-J18)	18	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.5	*
11	Septicemia (A40-A41)	14	0.5	*
12	Diabetes mellitus (E10-E14)	13	0.4	*
12	Legal intervention (Y35,Y89.0)	13	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	316	10.5	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,995	100.0	68.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,151	38.4	26.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	387	12.9	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	372	12.4	8.5
4	Malignant neoplasms (C00-C97)	267	8.9	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	4.7	3.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	41	1.4	0.9
7	Cerebrovascular diseases (I60-I69)	37	1.2	0.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.0	0.7
9	Diabetes mellitus (E10-E14)	29	1.0	0.7
10	Influenza and pneumonia (J09-J18)	26	0.9	0.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	26	0.9	0.6
12	Chronic lower respiratory diseases (J40-J47)	21	0.7	0.5
13	Septicemia (A40-A41)	20	0.7	0.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.7	0.5
15	Legal intervention (Y35,Y89.0)	13	0.4	*
...	All other causes (Residual)	413	13.8	9.4
Hispanic, both sexes, 30-34 years				
...	All causes	3,346	100.0	76.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,052	31.4	24.1
2	Malignant neoplasms (C00-C97)	457	13.7	10.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	10.1	7.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	9.4	7.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	7.0	5.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	97	2.9	2.2
7	Cerebrovascular diseases (I60-I69)	70	2.1	1.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	62	1.9	1.4
9	Diabetes mellitus (E10-E14)	59	1.8	1.4
10	Influenza and pneumonia (J09-J18)	44	1.3	1.0
11	Septicemia (A40-A41)	30	0.9	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.8	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.7	0.5
15	Chronic lower respiratory diseases (J40-J47)	20	0.6	0.5
...	All other causes (Residual)	492	14.7	11.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,777	100.0	93.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	978	25.9	24.2
2	Malignant neoplasms (C00-C97)	610	16.2	15.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	10.2	9.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	7.4	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	6.3	5.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	169	4.5	4.2
7	Cerebrovascular diseases (I60-I69)	97	2.6	2.4
8	Diabetes mellitus (E10-E14)	92	2.4	2.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	79	2.1	2.0
10	Influenza and pneumonia (J09-J18)	49	1.3	1.2
11	Septicemia (A40-A41)	36	1.0	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	34	0.9	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	23	0.6	0.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.6	0.5
...	All other causes (Residual)	659	17.4	16.3
Hispanic, both sexes, 40-44 years				
...	All causes	4,921	100.0	131.4
1	Malignant neoplasms (C00-C97)	1,009	20.5	26.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	936	19.0	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	607	12.3	16.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	314	6.4	8.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	246	5.0	6.6
6	Cerebrovascular diseases (I60-I69)	210	4.3	5.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	3.6	4.7
8	Diabetes mellitus (E10-E14)	174	3.5	4.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	113	2.3	3.0
10	Influenza and pneumonia (J09-J18)	74	1.5	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.2	1.6
12	Septicemia (A40-A41)	56	1.1	1.5
13	Viral hepatitis (B15-B19)	45	0.9	1.2
14	Chronic lower respiratory diseases (J40-J47)	31	0.6	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.6	0.7
...	All other causes (Residual)	842	17.1	22.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,986	100.0	212.7
1	Malignant neoplasms (C00-C97)	1,614	23.1	49.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	999	14.3	30.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	964	13.8	29.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	613	8.8	18.7
5	Cerebrovascular diseases (I60-I69)	293	4.2	8.9
6	Diabetes mellitus (E10-E14)	281	4.0	8.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	3.1	6.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	155	2.2	4.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	2.1	4.5
10	Viral hepatitis (B15-B19)	132	1.9	4.0
11	Septicemia (A40-A41)	100	1.4	3.0
12	Influenza and pneumonia (J09-J18)	93	1.3	2.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	1.3	2.7
14	Chronic lower respiratory diseases (J40-J47)	53	0.8	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	0.7	1.4
...	All other causes (Residual)	1,187	17.0	36.1
Hispanic, both sexes, 50-54 years				
...	All causes	9,408	100.0	336.4
1	Malignant neoplasms (C00-C97)	2,505	26.6	89.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,621	17.2	58.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	896	9.5	32.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	789	8.4	28.2
5	Diabetes mellitus (E10-E14)	453	4.8	16.2
6	Cerebrovascular diseases (I60-I69)	403	4.3	14.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	221	2.3	7.9
8	Viral hepatitis (B15-B19)	219	2.3	7.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	170	1.8	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	1.7	5.6
11	Septicemia (A40-A41)	131	1.4	4.7
11	Influenza and pneumonia (J09-J18)	131	1.4	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	1.2	3.9
14	Chronic lower respiratory diseases (J40-J47)	105	1.1	3.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.9	2.9
...	All other causes (Residual)	1,419	15.1	50.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	11,089	100.0	504.9
1	Malignant neoplasms (C00-C97)	3,345	30.2	152.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,019	18.2	91.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	805	7.3	36.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	743	6.7	33.8
5	Diabetes mellitus (E10-E14)	646	5.8	29.4
6	Cerebrovascular diseases (I60-I69)	465	4.2	21.2
7	Viral hepatitis (B15-B19)	298	2.7	13.6
8	Chronic lower respiratory diseases (J40-J47)	201	1.8	9.2
9	Influenza and pneumonia (J09-J18)	179	1.6	8.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	1.6	8.1
11	Septicemia (A40-A41)	170	1.5	7.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	1.5	7.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.0	5.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	0.9	4.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.5	2.7
...	All other causes (Residual)	1,603	14.5	73.0
Hispanic, both sexes, 60-64 years				
...	All causes	12,237	100.0	754.0
1	Malignant neoplasms (C00-C97)	3,916	32.0	241.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,506	20.5	154.4
3	Diabetes mellitus (E10-E14)	779	6.4	48.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	662	5.4	40.8
5	Cerebrovascular diseases (I60-I69)	547	4.5	33.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	519	4.2	32.0
7	Chronic lower respiratory diseases (J40-J47)	270	2.2	16.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	2.1	15.8
9	Viral hepatitis (B15-B19)	216	1.8	13.3
10	Septicemia (A40-A41)	199	1.6	12.3
10	Influenza and pneumonia (J09-J18)	199	1.6	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.0	7.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	0.9	6.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	87	0.7	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.5	3.6
...	All other causes (Residual)	1,784	14.6	109.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	12,961	100.0	1,101.2
1	Malignant neoplasms (C00-C97)	4,153	32.0	352.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,822	21.8	239.8
3	Diabetes mellitus (E10-E14)	878	6.8	74.6
4	Cerebrovascular diseases (I60-I69)	606	4.7	51.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	540	4.2	45.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	2.8	31.2
7	Chronic lower respiratory diseases (J40-J47)	357	2.8	30.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	327	2.5	27.8
9	Influenza and pneumonia (J09-J18)	246	1.9	20.9
10	Septicemia (A40-A41)	220	1.7	18.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	139	1.1	11.8
12	Viral hepatitis (B15-B19)	138	1.1	11.7
13	Alzheimer's disease (G30)	94	0.7	8.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	0.5	5.9
15	Parkinson's disease (G20-G21)	62	0.5	5.3
...	All other causes (Residual)	1,942	15.0	165.0
Hispanic, both sexes, 70-74 years				
...	All causes	13,723	100.0	1,693.8
1	Malignant neoplasms (C00-C97)	4,202	30.6	518.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,051	22.2	376.6
3	Diabetes mellitus (E10-E14)	873	6.4	107.8
4	Cerebrovascular diseases (I60-I69)	715	5.2	88.3
5	Chronic lower respiratory diseases (J40-J47)	500	3.6	61.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	383	2.8	47.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	349	2.5	43.1
8	Influenza and pneumonia (J09-J18)	333	2.4	41.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	326	2.4	40.2
10	Septicemia (A40-A41)	224	1.6	27.6
11	Alzheimer's disease (G30)	203	1.5	25.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	178	1.3	22.0
13	Parkinson's disease (G20-G21)	140	1.0	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.6	9.9
15	Viral hepatitis (B15-B19)	73	0.5	9.0
...	All other causes (Residual)	2,093	15.3	258.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	16,210	100.0	2,778.8
1	Malignant neoplasms (C00-C97)	4,222	26.0	723.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,727	23.0	638.9
3	Diabetes mellitus (E10-E14)	988	6.1	169.4
4	Cerebrovascular diseases (I60-I69)	971	6.0	166.5
5	Chronic lower respiratory diseases (J40-J47)	729	4.5	125.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	2.6	71.0
7	Influenza and pneumonia (J09-J18)	410	2.5	70.3
8	Alzheimer's disease (G30)	389	2.4	66.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	2.0	56.7
10	Septicemia (A40-A41)	316	1.9	54.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	313	1.9	53.7
12	Parkinson's disease (G20-G21)	248	1.5	42.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.5	41.5
14	Pneumonitis due to solids and liquids (J69)	112	0.7	19.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.7	18.7
...	All other causes (Residual)	2,689	16.6	461.0
Hispanic, both sexes, 80-84 years				
...	All causes	18,556	100.0	4,604.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,645	25.0	1,152.5
2	Malignant neoplasms (C00-C97)	3,861	20.8	958.0
3	Cerebrovascular diseases (I60-I69)	1,239	6.7	307.4
4	Diabetes mellitus (E10-E14)	960	5.2	238.2
5	Alzheimer's disease (G30)	840	4.5	208.4
6	Chronic lower respiratory diseases (J40-J47)	838	4.5	207.9
7	Influenza and pneumonia (J09-J18)	490	2.6	121.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	2.3	106.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	2.0	90.8
10	Septicemia (A40-A41)	330	1.8	81.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	325	1.8	80.6
12	Parkinson's disease (G20-G21)	313	1.7	77.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	241	1.3	59.8
14	Pneumonitis due to solids and liquids (J69)	130	0.7	32.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	121	0.7	30.0
...	All other causes (Residual)	3,429	18.5	850.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	18,324	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,005	27.3	---
2	Malignant neoplasms (C00-C97)	2,734	14.9	---
3	Cerebrovascular diseases (I60-I69)	1,207	6.6	---
4	Alzheimer's disease (G30)	1,146	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	890	4.9	---
6	Diabetes mellitus (E10-E14)	860	4.7	---
7	Influenza and pneumonia (J09-J18)	600	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	352	1.9	---
11	Septicemia (A40-A41)	298	1.6	---
12	Parkinson's disease (G20-G21)	294	1.6	---
13	Pneumonitis due to solids and liquids (J69)	153	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	134	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	114	0.6	---
...	All other causes (Residual)	3,743	20.4	---
Hispanic, both sexes, 90-94 years				
...	All causes	11,840	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,445	29.1	---
2	Malignant neoplasms (C00-C97)	1,247	10.5	---
3	Alzheimer's disease (G30)	946	8.0	---
4	Cerebrovascular diseases (I60-I69)	841	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	526	4.4	---
6	Diabetes mellitus (E10-E14)	406	3.4	---
7	Influenza and pneumonia (J09-J18)	386	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	1.9	---
11	Parkinson's disease (G20-G21)	164	1.4	---
12	Septicemia (A40-A41)	161	1.4	---
13	Pneumonitis due to solids and liquids (J69)	117	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	48	0.4	---
...	All other causes (Residual)	2,740	23.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	4,423	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,431	32.4	---
2	Alzheimer's disease (G30)	356	8.0	---
3	Malignant neoplasms (C00-C97)	326	7.4	---
4	Cerebrovascular diseases (I60-I69)	298	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	184	4.2	---
6	Influenza and pneumonia (J09-J18)	164	3.7	---
7	Diabetes mellitus (E10-E14)	115	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.6	---
11	Septicemia (A40-A41)	57	1.3	---
12	Pneumonitis due to solids and liquids (J69)	38	0.9	---
13	Atherosclerosis (I70)	27	0.6	---
14	Parkinson's disease (G20-G21)	26	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	16	0.4	---
...	All other causes (Residual)	1,121	25.3	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,103	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	373	33.8	---
2	Alzheimer's disease (G30)	87	7.9	---
3	Influenza and pneumonia (J09-J18)	65	5.9	---
4	Cerebrovascular diseases (I60-I69)	51	4.6	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	3.3	---
6	Malignant neoplasms (C00-C97)	29	2.6	---
6	Chronic lower respiratory diseases (J40-J47)	29	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.6	---
9	Pneumonitis due to solids and liquids (J69)	17	1.5	---
10	Diabetes mellitus (E10-E14)	16	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.1	---
12	Septicemia (A40-A41)	10	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	---
13	Atherosclerosis (I70)	8	0.7	---
15	Nutritional deficiencies (E40-E64)	4	0.4	---
15	Parkinson's disease (G20-G21)	4	0.4	---
...	All other causes (Residual)	336	30.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	88,880	100.0	323.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,377	20.7	66.9
2	Malignant neoplasms (C00-C97)	18,371	20.7	66.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,760	9.9	31.9
4	Diabetes mellitus (E10-E14)	3,934	4.4	14.3
5	Cerebrovascular diseases (I60-I69)	3,841	4.3	14.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,552	4.0	12.9
7	Chronic lower respiratory diseases (J40-J47)	2,536	2.9	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,279	2.6	8.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,132	2.4	7.8
10	Influenza and pneumonia (J09-J18)	1,816	2.0	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,583	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,320	1.5	4.8
13	Alzheimer's disease (G30)	1,298	1.5	4.7
14	Septicemia (A40-A41)	1,226	1.4	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	972	1.1	3.5
...	All other causes (Residual)	16,883	19.0	61.5
Hispanic, male, 1-4 years				
...	All causes	483	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	37.7	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.0	3.0
3	Malignant neoplasms (C00-C97)	48	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	6.4	1.5
5	Influenza and pneumonia (J09-J18)	9	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
8	Septicemia (A40-A41)	4	0.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
10	Meningitis (G00,G03)	3	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Anemias (D50-D64)	2	0.4	*
11	Diseases of appendix (K35-K38)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
...	All other causes (Residual)	113	23.4	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	265	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	30.2	3.1
2	Malignant neoplasms (C00-C97)	51	19.2	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	9.4	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.8	*
5	Chronic lower respiratory diseases (J40-J47)	9	3.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.9	*
7	Septicemia (A40-A41)	4	1.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.5	*
7	Cerebrovascular diseases (I60-I69)	4	1.5	*
7	Influenza and pneumonia (J09-J18)	4	1.5	*
11	Anemias (D50-D64)	2	0.8	*
12	Meningitis (G00,G03)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	63	23.8	2.4
Hispanic, male, 10-14 years				
...	All causes	326	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	28.5	3.8
2	Malignant neoplasms (C00-C97)	55	16.9	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	11.0	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	7.1	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	4.0	*
7	Cerebrovascular diseases (I60-I69)	6	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	6	1.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.2	*
10	Influenza and pneumonia (J09-J18)	3	0.9	*
11	Pneumonitis due to solids and liquids (J69)	2	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
13	Septicemia (A40-A41)	1	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	63	19.3	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,191	100.0	50.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	470	39.5	19.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	278	23.3	11.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	13.5	6.8
4	Malignant neoplasms (C00-C97)	85	7.1	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
8	Septicemia (A40-A41)	7	0.6	*
9	Influenza and pneumonia (J09-J18)	5	0.4	*
10	Cerebrovascular diseases (I60-I69)	4	0.3	*
10	Legal intervention (Y35,Y89.0)	4	0.3	*
12	Diabetes mellitus (E10-E14)	2	0.2	*
12	Aortic aneurysm and dissection (I71)	2	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.2	*
...	All other causes (Residual)	112	9.4	4.7
Hispanic, male, 20-24 years				
...	All causes	2,325	100.0	94.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,043	44.9	42.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	462	19.9	18.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	326	14.0	13.3
4	Malignant neoplasms (C00-C97)	124	5.3	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	2.4	2.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.6	*
8	Legal intervention (Y35,Y89.0)	13	0.6	*
9	Cerebrovascular diseases (I60-I69)	11	0.5	*
10	Influenza and pneumonia (J09-J18)	9	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.3	*
13	Diabetes mellitus (E10-E14)	7	0.3	*
14	Septicemia (A40-A41)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	216	9.3	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,266	100.0	96.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	935	41.3	39.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	334	14.7	14.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	321	14.2	13.7
4	Malignant neoplasms (C00-C97)	144	6.4	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	4.4	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	34	1.5	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.1	1.1
8	Cerebrovascular diseases (I60-I69)	23	1.0	1.0
9	Diabetes mellitus (E10-E14)	19	0.8	*
10	Influenza and pneumonia (J09-J18)	15	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	13	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
14	Septicemia (A40-A41)	10	0.4	*
15	Aortic aneurysm and dissection (I71)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	254	11.2	10.8
Hispanic, male, 30-34 years				
...	All causes	2,366	100.0	103.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	836	35.3	36.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	282	11.9	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	10.6	10.9
4	Malignant neoplasms (C00-C97)	219	9.3	9.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	6.8	7.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	69	2.9	3.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	50	2.1	2.2
8	Cerebrovascular diseases (I60-I69)	46	1.9	2.0
9	Diabetes mellitus (E10-E14)	38	1.6	1.7
10	Influenza and pneumonia (J09-J18)	26	1.1	1.1
11	Septicemia (A40-A41)	20	0.8	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.8	*
13	Legal intervention (Y35,Y89.0)	18	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
...	All other causes (Residual)	306	12.9	13.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,545	100.0	122.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	30.2	36.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	12.1	14.8
3	Malignant neoplasms (C00-C97)	241	9.5	11.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	9.0	10.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	191	7.5	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	124	4.9	5.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	71	2.8	3.4
8	Diabetes mellitus (E10-E14)	56	2.2	2.7
8	Cerebrovascular diseases (I60-I69)	56	2.2	2.7
10	Influenza and pneumonia (J09-J18)	31	1.2	1.5
11	Septicemia (A40-A41)	21	0.8	1.0
12	Legal intervention (Y35,Y89.0)	20	0.8	1.0
13	Aortic aneurysm and dissection (I71)	17	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
...	All other causes (Residual)	375	14.7	18.0
Hispanic, male, 40-44 years				
...	All causes	3,183	100.0	166.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	711	22.3	37.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	450	14.1	23.6
3	Malignant neoplasms (C00-C97)	414	13.0	21.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	250	7.9	13.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	6.1	10.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	4.5	7.5
7	Cerebrovascular diseases (I60-I69)	127	4.0	6.6
8	Diabetes mellitus (E10-E14)	110	3.5	5.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	86	2.7	4.5
10	Influenza and pneumonia (J09-J18)	48	1.5	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.2	2.0
12	Viral hepatitis (B15-B19)	35	1.1	1.8
13	Septicemia (A40-A41)	33	1.0	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.7	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.5	*
...	All other causes (Residual)	506	15.9	26.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,573	100.0	273.5
1	Malignant neoplasms (C00-C97)	751	16.4	44.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	747	16.3	44.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	743	16.2	44.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	477	10.4	28.5
5	Cerebrovascular diseases (I60-I69)	185	4.0	11.1
6	Diabetes mellitus (E10-E14)	184	4.0	11.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	3.7	10.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	124	2.7	7.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	2.7	7.4
10	Viral hepatitis (B15-B19)	106	2.3	6.3
11	Septicemia (A40-A41)	61	1.3	3.6
12	Influenza and pneumonia (J09-J18)	59	1.3	3.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.1	3.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.7	1.9
15	Chronic lower respiratory diseases (J40-J47)	29	0.6	1.7
...	All other causes (Residual)	736	16.1	44.0
Hispanic, male, 50-54 years				
...	All causes	6,110	100.0	437.1
1	Malignant neoplasms (C00-C97)	1,288	21.1	92.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,207	19.8	86.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	11.2	48.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	597	9.8	42.7
5	Diabetes mellitus (E10-E14)	296	4.8	21.2
6	Cerebrovascular diseases (I60-I69)	238	3.9	17.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	2.7	11.9
8	Viral hepatitis (B15-B19)	156	2.6	11.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	140	2.3	10.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.6	7.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	1.5	6.7
12	Influenza and pneumonia (J09-J18)	86	1.4	6.2
13	Chronic lower respiratory diseases (J40-J47)	68	1.1	4.9
14	Septicemia (A40-A41)	66	1.1	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	0.8	3.5
...	All other causes (Residual)	878	14.4	62.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	7,088	100.0	662.2
1	Malignant neoplasms (C00-C97)	1,805	25.5	168.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,492	21.0	139.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	605	8.5	56.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	563	7.9	52.6
5	Diabetes mellitus (E10-E14)	399	5.6	37.3
6	Cerebrovascular diseases (I60-I69)	301	4.2	28.1
7	Viral hepatitis (B15-B19)	211	3.0	19.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	1.8	12.1
9	Influenza and pneumonia (J09-J18)	114	1.6	10.7
10	Chronic lower respiratory diseases (J40-J47)	110	1.6	10.3
11	Septicemia (A40-A41)	98	1.4	9.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.3	8.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	88	1.2	8.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	0.8	5.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	0.7	4.8
...	All other causes (Residual)	969	13.7	90.5
Hispanic, male, 60-64 years				
...	All causes	7,586	100.0	985.3
1	Malignant neoplasms (C00-C97)	2,196	28.9	285.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,784	23.5	231.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	501	6.6	65.1
4	Diabetes mellitus (E10-E14)	424	5.6	55.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	5.0	49.5
6	Cerebrovascular diseases (I60-I69)	322	4.2	41.8
7	Viral hepatitis (B15-B19)	157	2.1	20.4
8	Chronic lower respiratory diseases (J40-J47)	156	2.1	20.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.8	18.1
10	Influenza and pneumonia (J09-J18)	125	1.6	16.2
11	Septicemia (A40-A41)	106	1.4	13.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	1.3	12.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.1	10.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.9	8.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.4	4.4
...	All other causes (Residual)	1,020	13.4	132.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	7,646	100.0	1,411.4
1	Malignant neoplasms (C00-C97)	2,325	30.4	429.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,841	24.1	339.8
3	Diabetes mellitus (E10-E14)	495	6.5	91.4
4	Cerebrovascular diseases (I60-I69)	350	4.6	64.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	342	4.5	63.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	3.3	47.1
7	Chronic lower respiratory diseases (J40-J47)	208	2.7	38.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.1	30.3
9	Influenza and pneumonia (J09-J18)	149	1.9	27.5
10	Septicemia (A40-A41)	124	1.6	22.9
11	Viral hepatitis (B15-B19)	87	1.1	16.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	83	1.1	15.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	0.8	11.4
14	Alzheimer's disease (G30)	47	0.6	8.7
15	Parkinson's disease (G20-G21)	43	0.6	7.9
...	All other causes (Residual)	1,071	14.0	197.7
Hispanic, male, 70-74 years				
...	All causes	7,709	100.0	2,165.3
1	Malignant neoplasms (C00-C97)	2,304	29.9	647.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,899	24.6	533.4
3	Diabetes mellitus (E10-E14)	463	6.0	130.0
4	Cerebrovascular diseases (I60-I69)	387	5.0	108.7
5	Chronic lower respiratory diseases (J40-J47)	305	4.0	85.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	221	2.9	62.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	2.7	58.1
8	Influenza and pneumonia (J09-J18)	188	2.4	52.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	2.3	50.6
10	Septicemia (A40-A41)	117	1.5	32.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.3	28.4
12	Alzheimer's disease (G30)	98	1.3	27.5
13	Parkinson's disease (G20-G21)	94	1.2	26.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	0.6	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	11.5
...	All other causes (Residual)	1,059	13.7	297.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	8,363	100.0	3,400.5
1	Malignant neoplasms (C00-C97)	2,246	26.9	913.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,053	24.5	834.8
3	Diabetes mellitus (E10-E14)	475	5.7	193.1
4	Cerebrovascular diseases (I60-I69)	452	5.4	183.8
5	Chronic lower respiratory diseases (J40-J47)	409	4.9	166.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	221	2.6	89.9
7	Influenza and pneumonia (J09-J18)	215	2.6	87.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	2.2	74.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	164	2.0	66.7
10	Parkinson's disease (G20-G21)	155	1.9	63.0
11	Septicemia (A40-A41)	153	1.8	62.2
12	Alzheimer's disease (G30)	146	1.7	59.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.4	46.4
14	Pneumonitis due to solids and liquids (J69)	68	0.8	27.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.6	22.0
...	All other causes (Residual)	1,256	15.0	510.7
Hispanic, male, 80-84 years				
...	All causes	8,797	100.0	5,467.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,249	25.6	1,397.9
2	Malignant neoplasms (C00-C97)	2,031	23.1	1,262.4
3	Cerebrovascular diseases (I60-I69)	533	6.1	331.3
4	Diabetes mellitus (E10-E14)	428	4.9	266.0
5	Chronic lower respiratory diseases (J40-J47)	424	4.8	263.5
6	Alzheimer's disease (G30)	304	3.5	189.0
7	Influenza and pneumonia (J09-J18)	235	2.7	146.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	2.4	130.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	2.2	121.8
10	Parkinson's disease (G20-G21)	178	2.0	110.6
11	Septicemia (A40-A41)	162	1.8	100.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	154	1.8	95.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.1	59.7
14	Pneumonitis due to solids and liquids (J69)	74	0.8	46.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.8	42.3
...	All other causes (Residual)	1,455	16.5	904.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	7,662	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,200	28.7	---
2	Malignant neoplasms (C00-C97)	1,291	16.8	---
3	Cerebrovascular diseases (I60-I69)	461	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	424	5.5	---
5	Diabetes mellitus (E10-E14)	349	4.6	---
6	Alzheimer's disease (G30)	328	4.3	---
7	Influenza and pneumonia (J09-J18)	260	3.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	2.6	---
10	Parkinson's disease (G20-G21)	152	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.7	---
12	Septicemia (A40-A41)	131	1.7	---
13	Pneumonitis due to solids and liquids (J69)	79	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	---
...	All other causes (Residual)	1,347	17.6	---
Hispanic, male, 90-94 years				
...	All causes	4,330	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,285	29.7	---
2	Malignant neoplasms (C00-C97)	597	13.8	---
3	Alzheimer's disease (G30)	262	6.1	---
4	Cerebrovascular diseases (I60-I69)	242	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	236	5.5	---
6	Diabetes mellitus (E10-E14)	148	3.4	---
7	Influenza and pneumonia (J09-J18)	146	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	2.2	---
11	Parkinson's disease (G20-G21)	82	1.9	---
12	Septicemia (A40-A41)	62	1.4	---
13	Pneumonitis due to solids and liquids (J69)	45	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	15	0.3	---
...	All other causes (Residual)	881	20.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,228	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	382	31.1	---
2	Malignant neoplasms (C00-C97)	146	11.9	---
3	Cerebrovascular diseases (I60-I69)	73	5.9	---
3	Chronic lower respiratory diseases (J40-J47)	73	5.9	---
5	Alzheimer's disease (G30)	66	5.4	---
6	Influenza and pneumonia (J09-J18)	52	4.2	---
7	Diabetes mellitus (E10-E14)	34	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.5	---
11	Septicemia (A40-A41)	17	1.4	---
12	Parkinson's disease (G20-G21)	16	1.3	---
13	Pneumonitis due to solids and liquids (J69)	14	1.1	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	---
15	Anemias (D50-D64)	4	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.3	---
...	All other causes (Residual)	273	22.2	---
Hispanic, male, 100 years and over				
...	All causes	234	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	32.9	---
2	Alzheimer's disease (G30)	17	7.3	---
3	Influenza and pneumonia (J09-J18)	14	6.0	---
4	Cerebrovascular diseases (I60-I69)	12	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	11	4.7	---
6	Pneumonitis due to solids and liquids (J69)	7	3.0	---
7	Malignant neoplasms (C00-C97)	6	2.6	---
8	Diabetes mellitus (E10-E14)	5	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.7	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	---
12	Hyperplasia of prostate (N40)	2	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.9	---
14	Septicemia (A40-A41)	1	0.4	---
14	Parkinson's disease (G20-G21)	1	0.4	---
14	Atherosclerosis (I70)	1	0.4	---
14	Aortic aneurysm and dissection (I71)	1	0.4	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	---
...	All other causes (Residual)	63	26.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	74,361	100.0	279.4
1	Malignant neoplasms (C00-C97)	16,776	22.6	63.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,866	20.0	55.9
3	Cerebrovascular diseases (I60-I69)	4,286	5.8	16.1
4	Diabetes mellitus (E10-E14)	3,698	5.0	13.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,255	4.4	12.2
6	Alzheimer's disease (G30)	2,829	3.8	10.6
7	Chronic lower respiratory diseases (J40-J47)	2,291	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,776	2.4	6.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,589	2.1	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,500	2.0	5.6
11	Septicemia (A40-A41)	1,209	1.6	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,120	1.5	4.2
13	Certain conditions originating in the perinatal period (P00-P96)	1,108	1.5	4.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	813	1.1	3.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	586	0.8	2.2
...	All other causes (Residual)	16,659	22.4	62.6
Hispanic, female, 1-4 years				
...	All causes	371	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	25.9	4.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	15.9	2.9
3	Malignant neoplasms (C00-C97)	42	11.3	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.9	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.0	*
6	Influenza and pneumonia (J09-J18)	10	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.9	*
8	Septicemia (A40-A41)	5	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
10	Anemias (D50-D64)	3	0.8	*
10	Cerebrovascular diseases (I60-I69)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
13	Meningitis (G00,G03)	2	0.5	*
14	Shigellosis and amebiasis (A03,A06)	1	0.3	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	93	25.1	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	202	100.0	8.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	27.2	2.2
2	Malignant neoplasms (C00-C97)	45	22.3	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	9.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	6.4	*
5	Influenza and pneumonia (J09-J18)	8	4.0	*
6	Cerebrovascular diseases (I60-I69)	6	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.0	*
8	Septicemia (A40-A41)	3	1.5	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	1.0	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	35	17.3	1.4
Hispanic, female, 10-14 years				
...	All causes	280	100.0	12.0
1	Malignant neoplasms (C00-C97)	58	20.7	2.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	18.2	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	12.5	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	7.5	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	4.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.6	*
6	Cerebrovascular diseases (I60-I69)	10	3.6	*
8	Influenza and pneumonia (J09-J18)	7	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	1.1	*
11	Septicemia (A40-A41)	2	0.7	*
11	Diabetes mellitus (E10-E14)	2	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
14	Anemias (D50-D64)	1	0.4	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	58	20.7	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	455	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	36.5	7.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	15.8	3.2
3	Malignant neoplasms (C00-C97)	51	11.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	8.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.3	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	5	1.1	*
9	Influenza and pneumonia (J09-J18)	4	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Anemias (D50-D64)	3	0.7	*
10	Diabetes mellitus (E10-E14)	3	0.7	*
13	Septicemia (A40-A41)	2	0.4	*
14	Meningitis (G00,G03)	1	0.2	*
14	Parkinson's disease (G20-G21)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	65	14.3	2.9
Hispanic, female, 20-24 years				
...	All causes	686	100.0	30.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	38.6	12.0
2	Malignant neoplasms (C00-C97)	79	11.5	3.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	11.1	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	10.5	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	2.8	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	19	2.8	*
7	Cerebrovascular diseases (I60-I69)	12	1.7	*
8	Influenza and pneumonia (J09-J18)	9	1.3	*
9	Septicemia (A40-A41)	8	1.2	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.2	*
11	Diabetes mellitus (E10-E14)	6	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	101	14.7	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	729	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	29.6	10.5
2	Malignant neoplasms (C00-C97)	123	16.9	6.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	7.3	2.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	7.0	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	5.6	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	26	3.6	1.3
7	Cerebrovascular diseases (I60-I69)	14	1.9	*
8	Influenza and pneumonia (J09-J18)	11	1.5	*
9	Septicemia (A40-A41)	10	1.4	*
9	Diabetes mellitus (E10-E14)	10	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
...	All other causes (Residual)	140	19.2	6.8
Hispanic, female, 30-34 years				
...	All causes	980	100.0	47.3
1	Malignant neoplasms (C00-C97)	238	24.3	11.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	22.0	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	7.4	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	6.5	3.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	5.6	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.9	1.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	27	2.8	1.3
8	Cerebrovascular diseases (I60-I69)	24	2.4	1.2
9	Diabetes mellitus (E10-E14)	21	2.1	1.0
10	Influenza and pneumonia (J09-J18)	18	1.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.2	*
12	Septicemia (A40-A41)	10	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	10	1.0	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
...	All other causes (Residual)	168	17.1	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,232	100.0	62.9
1	Malignant neoplasms (C00-C97)	369	30.0	18.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	17.0	10.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	6.3	3.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	4.1	2.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	3.7	2.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	45	3.7	2.3
7	Cerebrovascular diseases (I60-I69)	41	3.3	2.1
8	Diabetes mellitus (E10-E14)	36	2.9	1.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	34	2.8	1.7
10	Influenza and pneumonia (J09-J18)	18	1.5	*
11	Septicemia (A40-A41)	15	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
...	All other causes (Residual)	246	20.0	12.6
Hispanic, female, 40-44 years				
...	All causes	1,738	100.0	94.7
1	Malignant neoplasms (C00-C97)	595	34.2	32.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	12.9	12.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	9.0	8.6
4	Cerebrovascular diseases (I60-I69)	83	4.8	4.5
5	Diabetes mellitus (E10-E14)	64	3.7	3.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	64	3.7	3.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.0	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	1.9	1.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	27	1.6	1.5
10	Influenza and pneumonia (J09-J18)	26	1.5	1.4
11	Septicemia (A40-A41)	23	1.3	1.3
12	Pregnancy, childbirth and the puerperium (O00-O99)	22	1.3	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	1.1
14	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
15	Viral hepatitis (B15-B19)	10	0.6	*
...	All other causes (Residual)	319	18.4	17.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,413	100.0	149.6
1	Malignant neoplasms (C00-C97)	863	35.8	53.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	256	10.6	15.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	9.0	13.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	136	5.6	8.4
5	Cerebrovascular diseases (I60-I69)	108	4.5	6.7
6	Diabetes mellitus (E10-E14)	97	4.0	6.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	2.2	3.3
8	Septicemia (A40-A41)	39	1.6	2.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.6	2.4
10	Influenza and pneumonia (J09-J18)	34	1.4	2.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	31	1.3	1.9
12	Viral hepatitis (B15-B19)	26	1.1	1.6
13	Chronic lower respiratory diseases (J40-J47)	24	1.0	1.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.0	1.5
15	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.7	*
...	All other causes (Residual)	449	18.6	27.8
Hispanic, female, 50-54 years				
...	All causes	3,298	100.0	235.8
1	Malignant neoplasms (C00-C97)	1,217	36.9	87.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	414	12.6	29.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	6.5	15.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	192	5.8	13.7
5	Cerebrovascular diseases (I60-I69)	165	5.0	11.8
6	Diabetes mellitus (E10-E14)	157	4.8	11.2
7	Septicemia (A40-A41)	65	2.0	4.6
8	Viral hepatitis (B15-B19)	63	1.9	4.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.8	4.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	1.6	3.9
11	Influenza and pneumonia (J09-J18)	45	1.4	3.2
12	Chronic lower respiratory diseases (J40-J47)	37	1.1	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.9	2.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.9	2.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.5	*
...	All other causes (Residual)	541	16.4	38.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	4,001	100.0	355.4
1	Malignant neoplasms (C00-C97)	1,540	38.5	136.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	527	13.2	46.8
3	Diabetes mellitus (E10-E14)	247	6.2	21.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	200	5.0	17.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	4.5	16.0
6	Cerebrovascular diseases (I60-I69)	164	4.1	14.6
7	Chronic lower respiratory diseases (J40-J47)	91	2.3	8.1
8	Viral hepatitis (B15-B19)	87	2.2	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	2.1	7.4
10	Septicemia (A40-A41)	72	1.8	6.4
11	Influenza and pneumonia (J09-J18)	65	1.6	5.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.1	3.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.9	3.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.5	1.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (Residual)	628	15.7	55.8
Hispanic, female, 60-64 years				
...	All causes	4,651	100.0	545.2
1	Malignant neoplasms (C00-C97)	1,720	37.0	201.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	722	15.5	84.6
3	Diabetes mellitus (E10-E14)	355	7.6	41.6
4	Cerebrovascular diseases (I60-I69)	225	4.8	26.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	161	3.5	18.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	3.0	16.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.5	13.7
8	Chronic lower respiratory diseases (J40-J47)	114	2.5	13.4
9	Septicemia (A40-A41)	93	2.0	10.9
10	Influenza and pneumonia (J09-J18)	74	1.6	8.7
11	Viral hepatitis (B15-B19)	59	1.3	6.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.0	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	2.9
14	Alzheimer's disease (G30)	23	0.5	2.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	2.6
...	All other causes (Residual)	757	16.3	88.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	5,315	100.0	836.7
1	Malignant neoplasms (C00-C97)	1,828	34.4	287.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	981	18.5	154.4
3	Diabetes mellitus (E10-E14)	383	7.2	60.3
4	Cerebrovascular diseases (I60-I69)	256	4.8	40.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	198	3.7	31.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	3.1	25.7
7	Chronic lower respiratory diseases (J40-J47)	149	2.8	23.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.1	17.6
9	Influenza and pneumonia (J09-J18)	97	1.8	15.3
10	Septicemia (A40-A41)	96	1.8	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.1	8.8
12	Viral hepatitis (B15-B19)	51	1.0	8.0
13	Alzheimer's disease (G30)	47	0.9	7.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	3.5
15	Parkinson's disease (G20-G21)	19	0.4	*
...	All other causes (Residual)	857	16.1	134.9
Hispanic, female, 70-74 years				
...	All causes	6,014	100.0	1,324.2
1	Malignant neoplasms (C00-C97)	1,898	31.6	417.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,152	19.2	253.7
3	Diabetes mellitus (E10-E14)	410	6.8	90.3
4	Cerebrovascular diseases (I60-I69)	328	5.5	72.2
5	Chronic lower respiratory diseases (J40-J47)	195	3.2	42.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.8	37.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.7	35.7
8	Influenza and pneumonia (J09-J18)	145	2.4	31.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	2.0	26.2
10	Septicemia (A40-A41)	107	1.8	23.6
11	Alzheimer's disease (G30)	105	1.7	23.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	1.3	17.0
13	Parkinson's disease (G20-G21)	46	0.8	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.6	8.6
15	Viral hepatitis (B15-B19)	38	0.6	8.4
...	All other causes (Residual)	1,024	17.0	225.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	7,847	100.0	2,325.7
1	Malignant neoplasms (C00-C97)	1,976	25.2	585.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,674	21.3	496.1
3	Cerebrovascular diseases (I60-I69)	519	6.6	153.8
4	Diabetes mellitus (E10-E14)	513	6.5	152.0
5	Chronic lower respiratory diseases (J40-J47)	320	4.1	94.8
6	Alzheimer's disease (G30)	243	3.1	72.0
7	Influenza and pneumonia (J09-J18)	195	2.5	57.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	2.5	57.2
9	Septicemia (A40-A41)	163	2.1	48.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.9	44.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.9	44.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	128	1.6	37.9
13	Parkinson's disease (G20-G21)	93	1.2	27.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.7	16.3
15	Pneumonitis due to solids and liquids (J69)	44	0.6	13.0
...	All other causes (Residual)	1,433	18.3	424.7
Hispanic, female, 80-84 years				
...	All causes	9,759	100.0	4,030.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,396	24.6	989.4
2	Malignant neoplasms (C00-C97)	1,830	18.8	755.7
3	Cerebrovascular diseases (I60-I69)	706	7.2	291.5
4	Alzheimer's disease (G30)	536	5.5	221.3
5	Diabetes mellitus (E10-E14)	532	5.5	219.7
6	Chronic lower respiratory diseases (J40-J47)	414	4.2	171.0
7	Influenza and pneumonia (J09-J18)	255	2.6	105.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.2	90.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	1.8	70.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	1.7	70.2
11	Septicemia (A40-A41)	168	1.7	69.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	145	1.5	59.9
13	Parkinson's disease (G20-G21)	135	1.4	55.7
14	Pneumonitis due to solids and liquids (J69)	56	0.6	23.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	54	0.6	22.3
...	All other causes (Residual)	1,973	20.2	814.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	10,662	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,805	26.3	---
2	Malignant neoplasms (C00-C97)	1,443	13.5	---
3	Alzheimer's disease (G30)	818	7.7	---
4	Cerebrovascular diseases (I60-I69)	746	7.0	---
5	Diabetes mellitus (E10-E14)	511	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	466	4.4	---
7	Influenza and pneumonia (J09-J18)	340	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	218	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.7	---
11	Septicemia (A40-A41)	167	1.6	---
12	Parkinson's disease (G20-G21)	142	1.3	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	77	0.7	---
14	Pneumonitis due to solids and liquids (J69)	74	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	---
...	All other causes (Residual)	2,396	22.5	---
Hispanic, female, 90-94 years				
...	All causes	7,510	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,160	28.8	---
2	Alzheimer's disease (G30)	684	9.1	---
3	Malignant neoplasms (C00-C97)	650	8.7	---
4	Cerebrovascular diseases (I60-I69)	599	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	290	3.9	---
6	Diabetes mellitus (E10-E14)	258	3.4	---
7	Influenza and pneumonia (J09-J18)	240	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	1.7	---
11	Septicemia (A40-A41)	99	1.3	---
12	Parkinson's disease (G20-G21)	82	1.1	---
13	Pneumonitis due to solids and liquids (J69)	72	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.5	---
14	Atherosclerosis (I70)	35	0.5	---
...	All other causes (Residual)	1,857	24.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	3,195	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,049	32.8	---
2	Alzheimer's disease (G30)	290	9.1	---
3	Cerebrovascular diseases (I60-I69)	225	7.0	---
4	Malignant neoplasms (C00-C97)	180	5.6	---
5	Influenza and pneumonia (J09-J18)	112	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	111	3.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	3.0	---
8	Diabetes mellitus (E10-E14)	81	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.5	---
11	Septicemia (A40-A41)	40	1.3	---
12	Atherosclerosis (I70)	24	0.8	---
12	Pneumonitis due to solids and liquids (J69)	24	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	---
14	Parkinson's disease (G20-G21)	10	0.3	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	10	0.3	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.3	---
...	All other causes (Residual)	819	25.6	---
Hispanic, female, 100 years and over				
...	All causes	869	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296	34.1	---
2	Alzheimer's disease (G30)	70	8.1	---
3	Influenza and pneumonia (J09-J18)	51	5.9	---
4	Cerebrovascular diseases (I60-I69)	39	4.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	3.7	---
6	Malignant neoplasms (C00-C97)	23	2.6	---
7	Chronic lower respiratory diseases (J40-J47)	18	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.5	---
9	Diabetes mellitus (E10-E14)	11	1.3	---
10	Pneumonitis due to solids and liquids (J69)	10	1.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.2	---
12	Septicemia (A40-A41)	9	1.0	---
13	Atherosclerosis (I70)	7	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	---
15	Nutritional deficiencies (E40-E64)	4	0.5	---
...	All other causes (Residual)	271	31.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,052,660	100.0	1,021.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	488,817	23.8	243.3
2	Malignant neoplasms (C00-C97)	462,850	22.5	230.4
3	Chronic lower respiratory diseases (J40-J47)	131,667	6.4	65.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,773	4.9	50.2
5	Cerebrovascular diseases (I60-I69)	99,759	4.9	49.7
6	Alzheimer's disease (G30)	73,234	3.6	36.4
7	Diabetes mellitus (E10-E14)	51,297	2.5	25.5
8	Influenza and pneumonia (J09-J18)	45,350	2.2	22.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,253	1.7	17.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,173	1.7	17.0
11	Septicemia (A40-A41)	28,431	1.4	14.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,692	1.3	13.3
13	Parkinson's disease (G20-G21)	22,241	1.1	11.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,043	1.1	11.0
15	Pneumonitis due to solids and liquids (J69)	15,630	0.8	7.8
...	All other causes (Residual)	415,450	20.2	206.8
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,977	100.0	23.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	675	34.1	8.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	11.5	2.7
3	Malignant neoplasms (C00-C97)	160	8.1	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	4.1	1.0
6	Influenza and pneumonia (J09-J18)	53	2.7	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	24	1.2	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	15	0.8	*
12	Meningitis (G00,G03)	9	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	6	0.3	*
...	All other causes (Residual)	508	25.7	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,238	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	31.0	3.5
2	Malignant neoplasms (C00-C97)	260	21.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	6.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	4.8	0.5
5	Influenza and pneumonia (J09-J18)	40	3.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	2.8	0.3
7	Cerebrovascular diseases (I60-I69)	22	1.8	0.2
8	Septicemia (A40-A41)	18	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	14	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.7	*
12	Pneumonitis due to solids and liquids (J69)	4	0.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Diabetes mellitus (E10-E14)	3	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
...	All other causes (Residual)	275	22.2	2.5
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,586	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	28.6	4.0
2	Malignant neoplasms (C00-C97)	252	15.9	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	240	15.1	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	5.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.6	0.4
7	Influenza and pneumonia (J09-J18)	31	2.0	0.3
8	Chronic lower respiratory diseases (J40-J47)	24	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	20	1.3	0.2
11	Diabetes mellitus (E10-E14)	15	0.9	*
12	Septicemia (A40-A41)	10	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
15	Aortic aneurysm and dissection (I71)	4	0.3	*
...	All other causes (Residual)	312	19.7	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,253	100.0	43.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,378	45.3	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,237	23.5	10.3
3	Malignant neoplasms (C00-C97)	346	6.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	3.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	2.8	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	1.8	0.8
7	Influenza and pneumonia (J09-J18)	49	0.9	0.4
8	Cerebrovascular diseases (I60-I69)	26	0.5	0.2
9	Diabetes mellitus (E10-E14)	25	0.5	0.2
10	Septicemia (A40-A41)	22	0.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.4	0.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	11	0.2	*
...	All other causes (Residual)	659	12.5	5.5
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	10,839	100.0	83.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,442	50.2	41.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,151	19.8	16.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	502	4.6	3.9
4	Malignant neoplasms (C00-C97)	458	4.2	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	3.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	1.2	1.0
7	Influenza and pneumonia (J09-J18)	83	0.8	0.6
8	Diabetes mellitus (E10-E14)	70	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	55	0.5	0.4
10	Cerebrovascular diseases (I60-I69)	49	0.5	0.4
11	Septicemia (A40-A41)	48	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	32	0.3	0.2
12	Legal intervention (Y35,Y89.0)	32	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.2
...	All other causes (Residual)	1,402	12.9	10.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,634	100.0	100.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,731	45.4	45.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,390	18.9	19.0
3	Malignant neoplasms (C00-C97)	765	6.1	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	618	4.9	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	447	3.5	3.6
6	Diabetes mellitus (E10-E14)	127	1.0	1.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	123	1.0	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.9	1.0
9	Influenza and pneumonia (J09-J18)	106	0.8	0.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	98	0.8	0.8
11	Cerebrovascular diseases (I60-I69)	92	0.7	0.7
12	Septicemia (A40-A41)	71	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	60	0.5	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	46	0.4	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.3	0.3
...	All other causes (Residual)	1,797	14.2	14.3
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	14,949	100.0	120.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,863	39.2	47.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,335	15.6	18.8
3	Malignant neoplasms (C00-C97)	1,306	8.7	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,155	7.7	9.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	477	3.2	3.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	293	2.0	2.4
7	Diabetes mellitus (E10-E14)	215	1.4	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	170	1.1	1.4
9	Influenza and pneumonia (J09-J18)	160	1.1	1.3
10	Cerebrovascular diseases (I60-I69)	145	1.0	1.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	111	0.7	0.9
12	Septicemia (A40-A41)	105	0.7	0.8
13	Chronic lower respiratory diseases (J40-J47)	83	0.6	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.6	0.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	70	0.5	0.6
...	All other causes (Residual)	2,378	15.9	19.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	17,069	100.0	149.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,166	30.3	45.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,392	14.0	20.9
3	Malignant neoplasms (C00-C97)	2,288	13.4	20.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,875	11.0	16.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	531	3.1	4.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	404	2.4	3.5
7	Diabetes mellitus (E10-E14)	365	2.1	3.2
8	Cerebrovascular diseases (I60-I69)	229	1.3	2.0
9	Influenza and pneumonia (J09-J18)	226	1.3	2.0
10	Septicemia (A40-A41)	169	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	156	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	0.8	1.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	114	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	0.6	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.6
...	All other causes (Residual)	2,842	16.7	24.9
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,862	100.0	209.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,954	22.2	46.4
2	Malignant neoplasms (C00-C97)	4,813	17.9	37.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,266	15.9	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,906	10.8	22.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,069	4.0	8.3
6	Diabetes mellitus (E10-E14)	697	2.6	5.4
7	Cerebrovascular diseases (I60-I69)	529	2.0	4.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	1.6	3.3
9	Influenza and pneumonia (J09-J18)	332	1.2	2.6
10	Chronic lower respiratory diseases (J40-J47)	321	1.2	2.5
11	Septicemia (A40-A41)	300	1.1	2.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	231	0.9	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	0.6	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	0.5	0.9
...	All other causes (Residual)	4,556	17.0	35.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	44,615	100.0	322.8
1	Malignant neoplasms (C00-C97)	10,323	23.1	74.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,204	18.4	59.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,114	15.9	51.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,439	7.7	24.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,269	5.1	16.4
6	Diabetes mellitus (E10-E14)	1,266	2.8	9.2
7	Cerebrovascular diseases (I60-I69)	997	2.2	7.2
8	Chronic lower respiratory diseases (J40-J47)	944	2.1	6.8
9	Influenza and pneumonia (J09-J18)	528	1.2	3.8
10	Septicemia (A40-A41)	511	1.1	3.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	442	1.0	3.2
12	Viral hepatitis (B15-B19)	363	0.8	2.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	359	0.8	2.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	0.8	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	237	0.5	1.7
...	All other causes (Residual)	7,276	16.3	52.6
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	77,916	100.0	497.0
1	Malignant neoplasms (C00-C97)	22,111	28.4	141.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,511	19.9	98.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,263	10.6	52.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,088	5.2	26.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,949	5.1	25.2
6	Chronic lower respiratory diseases (J40-J47)	2,764	3.5	17.6
7	Diabetes mellitus (E10-E14)	2,220	2.8	14.2
8	Cerebrovascular diseases (I60-I69)	1,916	2.5	12.2
9	Septicemia (A40-A41)	966	1.2	6.2
10	Influenza and pneumonia (J09-J18)	949	1.2	6.1
11	Viral hepatitis (B15-B19)	892	1.1	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	689	0.9	4.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	559	0.7	3.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	461	0.6	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	453	0.6	2.9
...	All other causes (Residual)	12,125	15.6	77.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	109,650	100.0	715.0
1	Malignant neoplasms (C00-C97)	36,029	32.9	234.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,469	20.5	146.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,451	6.8	48.6
4	Chronic lower respiratory diseases (J40-J47)	5,186	4.7	33.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,681	4.3	30.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,728	3.4	24.3
7	Diabetes mellitus (E10-E14)	3,506	3.2	22.9
8	Cerebrovascular diseases (I60-I69)	2,948	2.7	19.2
9	Septicemia (A40-A41)	1,513	1.4	9.9
10	Influenza and pneumonia (J09-J18)	1,431	1.3	9.3
11	Viral hepatitis (B15-B19)	1,428	1.3	9.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,110	1.0	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	852	0.8	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	466	0.4	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	0.4	2.6
...	All other causes (Residual)	16,451	15.0	107.3
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	135,868	100.0	1,002.1
1	Malignant neoplasms (C00-C97)	48,382	35.6	356.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,879	21.3	213.0
3	Chronic lower respiratory diseases (J40-J47)	8,202	6.0	60.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,450	4.0	40.2
5	Diabetes mellitus (E10-E14)	4,670	3.4	34.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,133	3.0	30.5
7	Cerebrovascular diseases (I60-I69)	4,047	3.0	29.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,683	2.0	19.8
9	Septicemia (A40-A41)	2,152	1.6	15.9
10	Influenza and pneumonia (J09-J18)	1,906	1.4	14.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,683	1.2	12.4
12	Viral hepatitis (B15-B19)	1,201	0.9	8.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,038	0.8	7.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	668	0.5	4.9
15	Aortic aneurysm and dissection (I71)	586	0.4	4.3
...	All other causes (Residual)	20,188	14.9	148.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	165,943	100.0	1,466.2
1	Malignant neoplasms (C00-C97)	60,332	36.4	533.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,144	21.2	310.5
3	Chronic lower respiratory diseases (J40-J47)	13,339	8.0	117.9
4	Diabetes mellitus (E10-E14)	5,722	3.4	50.6
5	Cerebrovascular diseases (I60-I69)	5,620	3.4	49.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,599	2.8	40.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,053	1.8	27.0
8	Septicemia (A40-A41)	2,626	1.6	23.2
9	Influenza and pneumonia (J09-J18)	2,454	1.5	21.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,419	1.5	21.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,960	1.2	17.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,270	0.8	11.2
13	Alzheimer's disease (G30)	1,070	0.6	9.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	999	0.6	8.8
15	Parkinson's disease (G20-G21)	846	0.5	7.5
...	All other causes (Residual)	24,490	14.8	216.4
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	190,006	100.0	2,281.5
1	Malignant neoplasms (C00-C97)	64,304	33.8	772.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,836	21.0	478.3
3	Chronic lower respiratory diseases (J40-J47)	18,322	9.6	220.0
4	Cerebrovascular diseases (I60-I69)	7,733	4.1	92.9
5	Diabetes mellitus (E10-E14)	5,907	3.1	70.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,207	2.2	50.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,236	1.7	38.9
8	Influenza and pneumonia (J09-J18)	3,061	1.6	36.8
9	Septicemia (A40-A41)	3,040	1.6	36.5
10	Alzheimer's disease (G30)	2,689	1.4	32.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,340	1.2	28.1
12	Parkinson's disease (G20-G21)	1,896	1.0	22.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,508	0.8	18.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,494	0.8	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,298	0.7	15.6
...	All other causes (Residual)	29,135	15.3	349.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	222,885	100.0	3,676.5
1	Malignant neoplasms (C00-C97)	63,179	28.3	1,042.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,152	22.1	810.8
3	Chronic lower respiratory diseases (J40-J47)	20,798	9.3	343.1
4	Cerebrovascular diseases (I60-I69)	11,309	5.1	186.5
5	Diabetes mellitus (E10-E14)	6,392	2.9	105.4
6	Alzheimer's disease (G30)	6,391	2.9	105.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,148	2.3	84.9
8	Influenza and pneumonia (J09-J18)	4,295	1.9	70.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,198	1.9	69.2
10	Parkinson's disease (G20-G21)	3,665	1.6	60.5
11	Septicemia (A40-A41)	3,594	1.6	59.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,990	0.9	32.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,772	0.8	29.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,659	0.7	27.4
15	Pneumonitis due to solids and liquids (J69)	1,554	0.7	25.6
...	All other causes (Residual)	37,789	17.0	623.3
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	289,367	100.0	6,185.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,139	24.2	1,499.3
2	Malignant neoplasms (C00-C97)	63,003	21.8	1,346.8
3	Chronic lower respiratory diseases (J40-J47)	23,520	8.1	502.8
4	Cerebrovascular diseases (I60-I69)	17,198	5.9	367.6
5	Alzheimer's disease (G30)	13,245	4.6	283.1
6	Diabetes mellitus (E10-E14)	7,333	2.5	156.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,305	2.5	156.2
8	Influenza and pneumonia (J09-J18)	6,867	2.4	146.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,750	2.0	122.9
10	Parkinson's disease (G20-G21)	5,642	1.9	120.6
11	Septicemia (A40-A41)	4,335	1.5	92.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,115	1.1	66.6
13	Pneumonitis due to solids and liquids (J69)	2,565	0.9	54.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,252	0.8	48.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,343	0.5	28.7
...	All other causes (Residual)	55,755	19.3	1,191.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	333,684	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	91,177	27.3	---
2	Malignant neoplasms (C00-C97)	51,388	15.4	---
3	Cerebrovascular diseases (I60-I69)	21,636	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	21,389	6.4	---
5	Alzheimer's disease (G30)	20,732	6.2	---
6	Influenza and pneumonia (J09-J18)	9,351	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,746	2.6	---
8	Diabetes mellitus (E10-E14)	6,958	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,876	2.1	---
10	Parkinson's disease (G20-G21)	5,878	1.8	---
11	Septicemia (A40-A41)	4,644	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,371	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,623	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,381	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,496	0.4	---
...	All other causes (Residual)	73,038	21.9	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	256,926	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,143	30.8	---
2	Malignant neoplasms (C00-C97)	25,748	10.0	---
3	Alzheimer's disease (G30)	18,622	7.2	---
4	Cerebrovascular diseases (I60-I69)	17,311	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	12,234	4.8	---
6	Influenza and pneumonia (J09-J18)	8,531	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,726	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,200	2.0	---
9	Diabetes mellitus (E10-E14)	4,354	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,263	1.7	---
11	Septicemia (A40-A41)	3,067	1.2	---
12	Parkinson's disease (G20-G21)	3,011	1.2	---
13	Pneumonitis due to solids and liquids (J69)	2,963	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,631	0.6	---
15	Atherosclerosis (I70)	1,257	0.5	---
...	All other causes (Residual)	62,865	24.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	102,027	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,213	32.6	---
2	Alzheimer's disease (G30)	8,131	8.0	---
3	Cerebrovascular diseases (I60-I69)	6,675	6.5	---
4	Malignant neoplasms (C00-C97)	6,589	6.5	---
5	Influenza and pneumonia (J09-J18)	3,873	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	3,684	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,687	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,045	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,881	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,275	1.2	---
11	Diabetes mellitus (E10-E14)	1,245	1.2	---
12	Septicemia (A40-A41)	993	1.0	---
13	Parkinson's disease (G20-G21)	687	0.7	---
14	Atherosclerosis (I70)	667	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	455	0.4	---
...	All other causes (Residual)	27,927	27.4	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	20,753	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,238	34.9	---
2	Alzheimer's disease (G30)	1,546	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,192	5.7	---
4	Influenza and pneumonia (J09-J18)	958	4.6	---
5	Malignant neoplasms (C00-C97)	766	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	594	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	542	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	480	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	324	1.6	---
10	Pneumonitis due to solids and liquids (J69)	257	1.2	---
11	Diabetes mellitus (E10-E14)	199	1.0	---
12	Atherosclerosis (I70)	196	0.9	---
13	Septicemia (A40-A41)	167	0.8	---
14	Parkinson's disease (G20-G21)	81	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.4	---
...	All other causes (Residual)	6,134	29.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,021,135	100.0	1,032.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,294	25.1	259.0
2	Malignant neoplasms (C00-C97)	244,646	24.0	247.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61,391	6.0	62.1
4	Chronic lower respiratory diseases (J40-J47)	61,106	6.0	61.8
5	Cerebrovascular diseases (I60-I69)	40,350	4.0	40.8
6	Diabetes mellitus (E10-E14)	27,807	2.7	28.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,621	2.6	26.9
8	Alzheimer's disease (G30)	22,345	2.2	22.6
9	Influenza and pneumonia (J09-J18)	21,059	2.1	21.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,290	1.7	17.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,213	1.7	17.4
12	Parkinson's disease (G20-G21)	13,370	1.3	13.5
13	Septicemia (A40-A41)	13,364	1.3	13.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,955	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,534	0.8	8.6
...	All other causes (Residual)	180,790	17.7	182.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,149	100.0	26.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	437	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.8	2.4
3	Malignant neoplasms (C00-C97)	96	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.8	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.2	1.1
6	Influenza and pneumonia (J09-J18)	31	2.7	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Septicemia (A40-A41)	13	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Anemias (D50-D64)	4	0.3	*
13	Meningitis (G00,G03)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	285	24.8	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	718	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	33.0	4.2
2	Malignant neoplasms (C00-C97)	140	19.5	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	7.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	5.3	0.7
5	Influenza and pneumonia (J09-J18)	23	3.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.4	*
7	Septicemia (A40-A41)	10	1.4	*
7	Cerebrovascular diseases (I60-I69)	10	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.6	*
13	Anemias (D50-D64)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Aortic aneurysm and dissection (I71)	2	0.3	*
...	All other causes (Residual)	158	22.0	2.8
Non-Hispanic, white, male, 10-14 years				
...	All causes	939	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	30.0	4.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	17.6	2.8
3	Malignant neoplasms (C00-C97)	126	13.4	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	4.8	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	3.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	2.1	0.3
7	Influenza and pneumonia (J09-J18)	18	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	11	1.2	*
11	Diabetes mellitus (E10-E14)	9	1.0	*
12	Septicemia (A40-A41)	6	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Aortic aneurysm and dissection (I71)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	179	19.1	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,611	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,629	45.1	26.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	973	26.9	15.8
3	Malignant neoplasms (C00-C97)	202	5.6	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	3.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	2.7	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.5	0.9
7	Influenza and pneumonia (J09-J18)	27	0.7	0.4
8	Diabetes mellitus (E10-E14)	17	0.5	*
9	Cerebrovascular diseases (I60-I69)	15	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	14	0.4	*
11	Septicemia (A40-A41)	12	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.2	*
14	Aortic aneurysm and dissection (I71)	6	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.1	*
...	All other causes (Residual)	404	11.2	6.6
Non-Hispanic, white, male, 20-24 years				
...	All causes	7,850	100.0	118.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,066	51.8	61.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,759	22.4	26.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	361	4.6	5.4
4	Malignant neoplasms (C00-C97)	254	3.2	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	2.8	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	1.1	1.3
7	Influenza and pneumonia (J09-J18)	52	0.7	0.8
8	Diabetes mellitus (E10-E14)	41	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	31	0.4	0.5
10	Legal intervention (Y35,Y89.0)	29	0.4	0.4
11	Septicemia (A40-A41)	25	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.3	0.3
13	Chronic lower respiratory diseases (J40-J47)	18	0.2	*
14	Pneumonitis due to solids and liquids (J69)	13	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
...	All other causes (Residual)	861	11.0	13.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,880	100.0	139.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,224	47.6	66.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,925	21.7	30.3
3	Malignant neoplasms (C00-C97)	429	4.8	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	411	4.6	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	340	3.8	5.3
6	Diabetes mellitus (E10-E14)	76	0.9	1.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.8	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.7	1.0
9	Influenza and pneumonia (J09-J18)	58	0.7	0.9
10	Cerebrovascular diseases (I60-I69)	52	0.6	0.8
11	Septicemia (A40-A41)	43	0.5	0.7
12	Chronic lower respiratory diseases (J40-J47)	35	0.4	0.6
12	Legal intervention (Y35,Y89.0)	35	0.4	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.3	0.4
...	All other causes (Residual)	1,061	11.9	16.7
Non-Hispanic, white, male, 30-34 years				
...	All causes	10,024	100.0	160.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,267	42.6	68.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,804	18.0	28.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	8.1	12.9
4	Malignant neoplasms (C00-C97)	642	6.4	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	317	3.2	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	192	1.9	3.1
7	Diabetes mellitus (E10-E14)	139	1.4	2.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.0	1.6
9	Influenza and pneumonia (J09-J18)	86	0.9	1.4
10	Cerebrovascular diseases (I60-I69)	79	0.8	1.3
11	Septicemia (A40-A41)	58	0.6	0.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.5	0.9
13	Chronic lower respiratory diseases (J40-J47)	44	0.4	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.4	0.7
15	Legal intervention (Y35,Y89.0)	31	0.3	0.5
...	All other causes (Residual)	1,360	13.6	21.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	10,692	100.0	186.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,517	32.9	61.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,829	17.1	31.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,276	11.9	22.2
4	Malignant neoplasms (C00-C97)	1,026	9.6	17.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	333	3.1	5.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	2.5	4.7
7	Diabetes mellitus (E10-E14)	217	2.0	3.8
8	Cerebrovascular diseases (I60-I69)	127	1.2	2.2
9	Influenza and pneumonia (J09-J18)	117	1.1	2.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.8	1.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	82	0.8	1.4
12	Septicemia (A40-A41)	72	0.7	1.3
12	Chronic lower respiratory diseases (J40-J47)	72	0.7	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.5	0.9
15	Aortic aneurysm and dissection (I71)	43	0.4	0.7
...	All other causes (Residual)	1,573	14.7	27.4
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,505	100.0	256.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,944	23.9	61.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,000	18.2	46.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,184	13.2	33.9
4	Malignant neoplasms (C00-C97)	2,101	12.7	32.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	685	4.2	10.6
6	Diabetes mellitus (E10-E14)	447	2.7	6.9
7	Cerebrovascular diseases (I60-I69)	296	1.8	4.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	285	1.7	4.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	195	1.2	3.0
10	Influenza and pneumonia (J09-J18)	170	1.0	2.6
11	Septicemia (A40-A41)	147	0.9	2.3
12	Chronic lower respiratory diseases (J40-J47)	131	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	0.5	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.5	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	0.5	1.3
...	All other causes (Residual)	2,663	16.1	41.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	27,122	100.0	393.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,905	21.8	85.6
2	Malignant neoplasms (C00-C97)	4,948	18.2	71.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,668	17.2	67.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,514	9.3	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,448	5.3	21.0
6	Diabetes mellitus (E10-E14)	784	2.9	11.4
7	Cerebrovascular diseases (I60-I69)	545	2.0	7.9
8	Chronic lower respiratory diseases (J40-J47)	405	1.5	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	295	1.1	4.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	1.1	4.2
11	Influenza and pneumonia (J09-J18)	284	1.0	4.1
12	Septicemia (A40-A41)	264	1.0	3.8
13	Viral hepatitis (B15-B19)	253	0.9	3.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	0.7	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	164	0.6	2.4
...	All other causes (Residual)	4,167	15.4	60.4
Non-Hispanic, white, male, 50-54 years				
...	All causes	47,836	100.0	616.4
1	Malignant neoplasms (C00-C97)	11,454	23.9	147.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,272	23.6	145.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,384	11.3	69.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,050	6.4	39.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,626	5.5	33.8
6	Diabetes mellitus (E10-E14)	1,425	3.0	18.4
7	Chronic lower respiratory diseases (J40-J47)	1,284	2.7	16.5
8	Cerebrovascular diseases (I60-I69)	1,065	2.2	13.7
9	Viral hepatitis (B15-B19)	621	1.3	8.0
10	Influenza and pneumonia (J09-J18)	534	1.1	6.9
11	Septicemia (A40-A41)	519	1.1	6.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	0.9	5.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	379	0.8	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	372	0.8	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	317	0.7	4.1
...	All other causes (Residual)	7,123	14.9	91.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	67,862	100.0	901.8
1	Malignant neoplasms (C00-C97)	19,817	29.2	263.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,290	24.0	216.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,984	7.3	66.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,307	4.9	43.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,774	4.1	36.9
6	Chronic lower respiratory diseases (J40-J47)	2,635	3.9	35.0
7	Diabetes mellitus (E10-E14)	2,227	3.3	29.6
8	Cerebrovascular diseases (I60-I69)	1,720	2.5	22.9
9	Viral hepatitis (B15-B19)	1,014	1.5	13.5
10	Influenza and pneumonia (J09-J18)	842	1.2	11.2
11	Septicemia (A40-A41)	816	1.2	10.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	635	0.9	8.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	583	0.9	7.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	317	0.5	4.2
15	Aortic aneurysm and dissection (I71)	286	0.4	3.8
...	All other causes (Residual)	9,615	14.2	127.8
Non-Hispanic, white, male, 60-64 years				
...	All causes	82,960	100.0	1,260.1
1	Malignant neoplasms (C00-C97)	27,146	32.7	412.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,421	24.6	310.2
3	Chronic lower respiratory diseases (J40-J47)	4,311	5.2	65.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,675	4.4	55.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,941	3.5	44.7
6	Diabetes mellitus (E10-E14)	2,925	3.5	44.4
7	Cerebrovascular diseases (I60-I69)	2,374	2.9	36.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,046	2.5	31.1
9	Septicemia (A40-A41)	1,182	1.4	18.0
10	Influenza and pneumonia (J09-J18)	1,061	1.3	16.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	961	1.2	14.6
12	Viral hepatitis (B15-B19)	881	1.1	13.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	674	0.8	10.2
14	Aortic aneurysm and dissection (I71)	438	0.5	6.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	391	0.5	5.9
...	All other causes (Residual)	11,533	13.9	175.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	96,581	100.0	1,776.7
1	Malignant neoplasms (C00-C97)	33,553	34.7	617.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,551	24.4	433.2
3	Chronic lower respiratory diseases (J40-J47)	6,608	6.8	121.6
4	Diabetes mellitus (E10-E14)	3,465	3.6	63.7
5	Cerebrovascular diseases (I60-I69)	3,064	3.2	56.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,038	3.1	55.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,006	2.1	36.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,528	1.6	28.1
9	Septicemia (A40-A41)	1,374	1.4	25.3
10	Influenza and pneumonia (J09-J18)	1,372	1.4	25.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,326	1.4	24.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	729	0.8	13.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	600	0.6	11.0
14	Parkinson's disease (G20-G21)	558	0.6	10.3
15	Aortic aneurysm and dissection (I71)	497	0.5	9.1
...	All other causes (Residual)	13,312	13.8	244.9
Non-Hispanic, white, male, 70-74 years				
...	All causes	106,316	100.0	2,729.0
1	Malignant neoplasms (C00-C97)	35,244	33.2	904.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,994	23.5	641.6
3	Chronic lower respiratory diseases (J40-J47)	9,069	8.5	232.8
4	Cerebrovascular diseases (I60-I69)	4,080	3.8	104.7
5	Diabetes mellitus (E10-E14)	3,426	3.2	87.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,564	2.4	65.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,726	1.6	44.3
8	Influenza and pneumonia (J09-J18)	1,676	1.6	43.0
9	Septicemia (A40-A41)	1,581	1.5	40.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,449	1.4	37.2
11	Parkinson's disease (G20-G21)	1,283	1.2	32.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,278	1.2	32.8
13	Alzheimer's disease (G30)	1,140	1.1	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	784	0.7	20.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	768	0.7	19.7
...	All other causes (Residual)	15,254	14.3	391.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	118,536	100.0	4,354.3
1	Malignant neoplasms (C00-C97)	34,257	28.9	1,258.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,847	24.3	1,059.7
3	Chronic lower respiratory diseases (J40-J47)	10,101	8.5	371.0
4	Cerebrovascular diseases (I60-I69)	5,403	4.6	198.5
5	Diabetes mellitus (E10-E14)	3,551	3.0	130.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,925	2.5	107.4
7	Alzheimer's disease (G30)	2,655	2.2	97.5
8	Parkinson's disease (G20-G21)	2,426	2.0	89.1
9	Influenza and pneumonia (J09-J18)	2,257	1.9	82.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,231	1.9	82.0
11	Septicemia (A40-A41)	1,720	1.5	63.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,001	0.8	36.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	972	0.8	35.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	954	0.8	35.0
15	Pneumonitis due to solids and liquids (J69)	929	0.8	34.1
...	All other causes (Residual)	18,307	15.4	672.5
Non-Hispanic, white, male, 80-84 years				
...	All causes	142,541	100.0	7,294.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,001	26.0	1,893.4
2	Malignant neoplasms (C00-C97)	33,311	23.4	1,704.6
3	Chronic lower respiratory diseases (J40-J47)	11,022	7.7	564.0
4	Cerebrovascular diseases (I60-I69)	7,318	5.1	374.5
5	Alzheimer's disease (G30)	4,871	3.4	249.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,800	2.7	194.5
7	Diabetes mellitus (E10-E14)	3,765	2.6	192.7
8	Parkinson's disease (G20-G21)	3,534	2.5	180.8
9	Influenza and pneumonia (J09-J18)	3,474	2.4	177.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,060	2.1	156.6
11	Septicemia (A40-A41)	2,051	1.4	105.0
12	Pneumonitis due to solids and liquids (J69)	1,457	1.0	74.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,279	0.9	65.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,255	0.9	64.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	874	0.6	44.7
...	All other causes (Residual)	24,469	17.2	1,252.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	143,077	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,353	28.9	---
2	Malignant neoplasms (C00-C97)	25,623	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,248	6.5	---
4	Cerebrovascular diseases (I60-I69)	7,516	5.3	---
5	Alzheimer's disease (G30)	6,541	4.6	---
6	Influenza and pneumonia (J09-J18)	4,205	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,978	2.8	---
8	Parkinson's disease (G20-G21)	3,365	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,344	2.3	---
10	Diabetes mellitus (E10-E14)	3,144	2.2	---
11	Septicemia (A40-A41)	2,004	1.4	---
12	Pneumonitis due to solids and liquids (J69)	2,001	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,561	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,213	0.8	---
15	Aortic aneurysm and dissection (I71)	647	0.5	---
...	All other causes (Residual)	27,334	19.1	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	91,199	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,815	32.7	---
2	Malignant neoplasms (C00-C97)	11,612	12.7	---
3	Cerebrovascular diseases (I60-I69)	4,970	5.4	---
4	Chronic lower respiratory diseases (J40-J47)	4,773	5.2	---
5	Alzheimer's disease (G30)	4,723	5.2	---
6	Influenza and pneumonia (J09-J18)	3,289	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,554	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,316	2.5	---
9	Diabetes mellitus (E10-E14)	1,699	1.9	---
10	Parkinson's disease (G20-G21)	1,534	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,478	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,238	1.4	---
13	Septicemia (A40-A41)	1,137	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	695	0.8	---
15	Atherosclerosis (I70)	388	0.4	---
...	All other causes (Residual)	18,978	20.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	27,049	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,491	35.1	---
2	Malignant neoplasms (C00-C97)	2,417	8.9	---
3	Cerebrovascular diseases (I60-I69)	1,448	5.4	---
4	Alzheimer's disease (G30)	1,431	5.3	---
5	Influenza and pneumonia (J09-J18)	1,236	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	1,182	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	787	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	690	2.6	---
9	Pneumonitis due to solids and liquids (J69)	539	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	418	1.5	---
11	Diabetes mellitus (E10-E14)	401	1.5	---
12	Parkinson's disease (G20-G21)	284	1.0	---
13	Septicemia (A40-A41)	266	1.0	---
14	Atherosclerosis (I70)	163	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	162	0.6	---
...	All other causes (Residual)	6,134	22.7	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	3,608	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,361	37.7	---
2	Malignant neoplasms (C00-C97)	219	6.1	---
3	Influenza and pneumonia (J09-J18)	205	5.7	---
4	Cerebrovascular diseases (I60-I69)	190	5.3	---
5	Alzheimer's disease (G30)	185	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	112	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	2.6	---
8	Pneumonitis due to solids and liquids (J69)	90	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.2	---
11	Diabetes mellitus (E10-E14)	42	1.2	---
12	Atherosclerosis (I70)	35	1.0	---
13	Septicemia (A40-A41)	32	0.9	---
14	Parkinson's disease (G20-G21)	26	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	---
15	Aortic aneurysm and dissection (I71)	14	0.4	---
...	All other causes (Residual)	820	22.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,031,525	100.0	1,011.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	232,523	22.5	228.0
2	Malignant neoplasms (C00-C97)	218,204	21.2	214.0
3	Chronic lower respiratory diseases (J40-J47)	70,561	6.8	69.2
4	Cerebrovascular diseases (I60-I69)	59,409	5.8	58.3
5	Alzheimer's disease (G30)	50,889	4.9	49.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,382	3.8	38.6
7	Influenza and pneumonia (J09-J18)	24,291	2.4	23.8
8	Diabetes mellitus (E10-E14)	23,490	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,960	1.6	16.6
10	Septicemia (A40-A41)	15,067	1.5	14.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,088	1.3	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,402	0.9	9.2
13	Parkinson's disease (G20-G21)	8,871	0.9	8.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,632	0.7	7.5
15	Pneumonitis due to solids and liquids (J69)	7,096	0.7	7.0
...	All other causes (Residual)	234,660	22.7	230.1
Non-Hispanic, white, female, 1-4 years				
...	All causes	828	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	28.7	5.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	15.2	3.1
3	Malignant neoplasms (C00-C97)	64	7.7	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.1	0.8
6	Influenza and pneumonia (J09-J18)	22	2.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
9	Septicemia (A40-A41)	9	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	0.8	*
11	Meningitis (G00,G03)	5	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	222	26.8	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	520	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	28.3	2.7
2	Malignant neoplasms (C00-C97)	120	23.1	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	6.7	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.2	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.5	*
6	Influenza and pneumonia (J09-J18)	17	3.3	*
7	Cerebrovascular diseases (I60-I69)	12	2.3	*
8	Septicemia (A40-A41)	8	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
11	Pneumonitis due to solids and liquids (J69)	3	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	118	22.7	2.2
Non-Hispanic, white, female, 10-14 years				
...	All causes	647	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	26.6	3.1
2	Malignant neoplasms (C00-C97)	126	19.5	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	11.6	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.2	0.4
7	Influenza and pneumonia (J09-J18)	13	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
11	Diabetes mellitus (E10-E14)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
13	Septicemia (A40-A41)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
...	All other causes (Residual)	132	20.4	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,642	100.0	28.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	749	45.6	12.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	16.1	4.5
3	Malignant neoplasms (C00-C97)	144	8.8	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.2	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	3.0	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	2.4	0.7
7	Influenza and pneumonia (J09-J18)	22	1.3	0.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	21	1.3	0.4
9	Cerebrovascular diseases (I60-I69)	11	0.7	*
10	Septicemia (A40-A41)	10	0.6	*
11	Diabetes mellitus (E10-E14)	8	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	244	14.9	4.2
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,989	100.0	47.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,376	46.0	21.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	13.1	6.2
3	Malignant neoplasms (C00-C97)	204	6.8	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	4.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	3.9	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	55	1.8	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.5	0.7
8	Influenza and pneumonia (J09-J18)	31	1.0	0.5
9	Diabetes mellitus (E10-E14)	29	1.0	0.5
10	Septicemia (A40-A41)	23	0.8	0.4
11	Cerebrovascular diseases (I60-I69)	18	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.3	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	516	17.3	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,754	100.0	60.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,507	40.1	24.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	465	12.4	7.5
3	Malignant neoplasms (C00-C97)	336	9.0	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	5.5	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	2.9	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	98	2.6	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.6	1.0
8	Diabetes mellitus (E10-E14)	51	1.4	0.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.4	0.8
10	Influenza and pneumonia (J09-J18)	48	1.3	0.8
11	Cerebrovascular diseases (I60-I69)	40	1.1	0.6
12	Septicemia (A40-A41)	28	0.7	0.5
13	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
...	All other causes (Residual)	697	18.6	11.2
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,925	100.0	80.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,596	32.4	26.0
2	Malignant neoplasms (C00-C97)	664	13.5	10.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	531	10.8	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	7.0	5.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	3.2	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	111	2.3	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	101	2.1	1.6
8	Diabetes mellitus (E10-E14)	76	1.5	1.2
9	Influenza and pneumonia (J09-J18)	74	1.5	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.4	1.1
11	Cerebrovascular diseases (I60-I69)	66	1.3	1.1
12	Septicemia (A40-A41)	47	1.0	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.9	0.7
14	Chronic lower respiratory diseases (J40-J47)	39	0.8	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	0.5
...	All other causes (Residual)	975	19.8	15.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,377	100.0	112.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,649	25.9	29.1
2	Malignant neoplasms (C00-C97)	1,262	19.8	22.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	599	9.4	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	8.8	9.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	198	3.1	3.5
6	Diabetes mellitus (E10-E14)	148	2.3	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	2.1	2.3
8	Influenza and pneumonia (J09-J18)	109	1.7	1.9
9	Cerebrovascular diseases (I60-I69)	102	1.6	1.8
10	Septicemia (A40-A41)	97	1.5	1.7
11	Chronic lower respiratory diseases (J40-J47)	84	1.3	1.5
12	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.1	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.7	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.5	0.6
...	All other causes (Residual)	1,226	19.2	21.6
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,357	100.0	161.9
1	Malignant neoplasms (C00-C97)	2,712	26.2	42.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,010	19.4	31.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,266	12.2	19.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	722	7.0	11.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	384	3.7	6.0
6	Diabetes mellitus (E10-E14)	250	2.4	3.9
7	Cerebrovascular diseases (I60-I69)	233	2.2	3.6
8	Chronic lower respiratory diseases (J40-J47)	190	1.8	3.0
9	Influenza and pneumonia (J09-J18)	162	1.6	2.5
10	Septicemia (A40-A41)	153	1.5	2.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	1.4	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.8	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.8	1.3
14	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.6	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	0.7
...	All other causes (Residual)	1,856	17.9	29.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	17,493	100.0	252.5
1	Malignant neoplasms (C00-C97)	5,375	30.7	77.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,446	14.0	35.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,299	13.1	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	925	5.3	13.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	821	4.7	11.9
6	Chronic lower respiratory diseases (J40-J47)	539	3.1	7.8
7	Diabetes mellitus (E10-E14)	482	2.8	7.0
8	Cerebrovascular diseases (I60-I69)	452	2.6	6.5
9	Septicemia (A40-A41)	247	1.4	3.6
10	Influenza and pneumonia (J09-J18)	244	1.4	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	0.9	2.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	0.9	2.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.7	1.7
14	Viral hepatitis (B15-B19)	110	0.6	1.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	1.0
...	All other causes (Residual)	3,058	17.5	44.1
Non-Hispanic, white, female, 50-54 years				
...	All causes	30,080	100.0	379.9
1	Malignant neoplasms (C00-C97)	10,657	35.4	134.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,239	14.1	53.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,879	9.6	36.4
4	Chronic lower respiratory diseases (J40-J47)	1,480	4.9	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,323	4.4	16.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,038	3.5	13.1
7	Cerebrovascular diseases (I60-I69)	851	2.8	10.7
8	Diabetes mellitus (E10-E14)	795	2.6	10.0
9	Septicemia (A40-A41)	447	1.5	5.6
10	Influenza and pneumonia (J09-J18)	415	1.4	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	0.9	3.5
12	Viral hepatitis (B15-B19)	271	0.9	3.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	0.6	2.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	0.5	1.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	0.5	1.7
...	All other causes (Residual)	4,943	16.4	62.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	41,788	100.0	535.0
1	Malignant neoplasms (C00-C97)	16,212	38.8	207.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,179	14.8	79.1
3	Chronic lower respiratory diseases (J40-J47)	2,551	6.1	32.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,467	5.9	31.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,374	3.3	17.6
6	Diabetes mellitus (E10-E14)	1,279	3.1	16.4
7	Cerebrovascular diseases (I60-I69)	1,228	2.9	15.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	954	2.3	12.2
9	Septicemia (A40-A41)	697	1.7	8.9
10	Influenza and pneumonia (J09-J18)	589	1.4	7.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	475	1.1	6.1
12	Viral hepatitis (B15-B19)	414	1.0	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	269	0.6	3.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.5	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	0.4	2.4
...	All other causes (Residual)	6,715	16.1	86.0
Non-Hispanic, white, female, 60-64 years				
...	All causes	52,908	100.0	758.5
1	Malignant neoplasms (C00-C97)	21,236	40.1	304.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,458	16.0	121.3
3	Chronic lower respiratory diseases (J40-J47)	3,891	7.4	55.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,775	3.4	25.4
5	Diabetes mellitus (E10-E14)	1,745	3.3	25.0
6	Cerebrovascular diseases (I60-I69)	1,673	3.2	24.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,192	2.3	17.1
8	Septicemia (A40-A41)	970	1.8	13.9
9	Influenza and pneumonia (J09-J18)	845	1.6	12.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	722	1.4	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	637	1.2	9.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	364	0.7	5.2
13	Alzheimer's disease (G30)	328	0.6	4.7
14	Viral hepatitis (B15-B19)	320	0.6	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	277	0.5	4.0
...	All other causes (Residual)	8,475	16.0	121.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	69,362	100.0	1,179.3
1	Malignant neoplasms (C00-C97)	26,779	38.6	455.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,593	16.7	197.1
3	Chronic lower respiratory diseases (J40-J47)	6,731	9.7	114.4
4	Cerebrovascular diseases (I60-I69)	2,556	3.7	43.5
5	Diabetes mellitus (E10-E14)	2,257	3.3	38.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,561	2.3	26.5
7	Septicemia (A40-A41)	1,252	1.8	21.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,093	1.6	18.6
9	Influenza and pneumonia (J09-J18)	1,082	1.6	18.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,047	1.5	17.8
11	Alzheimer's disease (G30)	596	0.9	10.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	541	0.8	9.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	432	0.6	7.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	399	0.6	6.8
15	Pneumonitis due to solids and liquids (J69)	289	0.4	4.9
...	All other causes (Residual)	11,154	16.1	189.6
Non-Hispanic, white, female, 70-74 years				
...	All causes	83,690	100.0	1,888.1
1	Malignant neoplasms (C00-C97)	29,060	34.7	655.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,842	17.7	334.8
3	Chronic lower respiratory diseases (J40-J47)	9,253	11.1	208.8
4	Cerebrovascular diseases (I60-I69)	3,653	4.4	82.4
5	Diabetes mellitus (E10-E14)	2,481	3.0	56.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,643	2.0	37.1
7	Alzheimer's disease (G30)	1,549	1.9	34.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,510	1.8	34.1
9	Septicemia (A40-A41)	1,459	1.7	32.9
10	Influenza and pneumonia (J09-J18)	1,385	1.7	31.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	891	1.1	20.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	726	0.9	16.4
13	Parkinson's disease (G20-G21)	613	0.7	13.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	514	0.6	11.6
15	Pneumonitis due to solids and liquids (J69)	414	0.5	9.3
...	All other causes (Residual)	13,697	16.4	309.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	104,349	100.0	3,124.2
1	Malignant neoplasms (C00-C97)	28,922	27.7	865.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,305	19.5	607.9
3	Chronic lower respiratory diseases (J40-J47)	10,697	10.3	320.3
4	Cerebrovascular diseases (I60-I69)	5,906	5.7	176.8
5	Alzheimer's disease (G30)	3,736	3.6	111.9
6	Diabetes mellitus (E10-E14)	2,841	2.7	85.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,223	2.1	66.6
8	Influenza and pneumonia (J09-J18)	2,038	2.0	61.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,967	1.9	58.9
10	Septicemia (A40-A41)	1,874	1.8	56.1
11	Parkinson's disease (G20-G21)	1,239	1.2	37.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,065	1.0	31.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	771	0.7	23.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	705	0.7	21.1
15	Pneumonitis due to solids and liquids (J69)	625	0.6	18.7
...	All other causes (Residual)	19,435	18.6	581.9
Non-Hispanic, white, female, 80-84 years				
...	All causes	146,826	100.0	5,390.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,138	22.6	1,216.6
2	Malignant neoplasms (C00-C97)	29,692	20.2	1,090.1
3	Chronic lower respiratory diseases (J40-J47)	12,498	8.5	458.8
4	Cerebrovascular diseases (I60-I69)	9,880	6.7	362.7
5	Alzheimer's disease (G30)	8,374	5.7	307.4
6	Diabetes mellitus (E10-E14)	3,568	2.4	131.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,505	2.4	128.7
8	Influenza and pneumonia (J09-J18)	3,393	2.3	124.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,690	1.8	98.8
10	Septicemia (A40-A41)	2,284	1.6	83.9
11	Parkinson's disease (G20-G21)	2,108	1.4	77.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,836	1.3	67.4
13	Pneumonitis due to solids and liquids (J69)	1,108	0.8	40.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	997	0.7	36.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	717	0.5	26.3
...	All other causes (Residual)	31,038	21.1	1,139.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	190,607	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,824	26.1	---
2	Malignant neoplasms (C00-C97)	25,765	13.5	---
3	Alzheimer's disease (G30)	14,191	7.4	---
4	Cerebrovascular diseases (I60-I69)	14,120	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	12,141	6.4	---
6	Influenza and pneumonia (J09-J18)	5,146	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,768	2.5	---
8	Diabetes mellitus (E10-E14)	3,814	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,532	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,810	1.5	---
11	Septicemia (A40-A41)	2,640	1.4	---
12	Parkinson's disease (G20-G21)	2,513	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,622	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,168	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	969	0.5	---
...	All other causes (Residual)	45,584	23.9	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	165,727	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,328	29.8	---
2	Malignant neoplasms (C00-C97)	14,136	8.5	---
3	Alzheimer's disease (G30)	13,899	8.4	---
4	Cerebrovascular diseases (I60-I69)	12,341	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	7,461	4.5	---
6	Influenza and pneumonia (J09-J18)	5,242	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,172	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,025	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,884	1.7	---
10	Diabetes mellitus (E10-E14)	2,655	1.6	---
11	Septicemia (A40-A41)	1,930	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,485	0.9	---
13	Parkinson's disease (G20-G21)	1,477	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	936	0.6	---
15	Atherosclerosis (I70)	869	0.5	---
...	All other causes (Residual)	43,887	26.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	74,978	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,722	31.6	---
2	Alzheimer's disease (G30)	6,700	8.9	---
3	Cerebrovascular diseases (I60-I69)	5,227	7.0	---
4	Malignant neoplasms (C00-C97)	4,172	5.6	---
5	Influenza and pneumonia (J09-J18)	2,637	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,502	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,900	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,627	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,191	1.6	---
10	Diabetes mellitus (E10-E14)	844	1.1	---
11	Pneumonitis due to solids and liquids (J69)	736	1.0	---
12	Septicemia (A40-A41)	727	1.0	---
13	Atherosclerosis (I70)	504	0.7	---
14	Parkinson's disease (G20-G21)	403	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	293	0.4	---
...	All other causes (Residual)	21,793	29.1	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	17,145	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,877	34.3	---
2	Alzheimer's disease (G30)	1,361	7.9	---
3	Cerebrovascular diseases (I60-I69)	1,002	5.8	---
4	Influenza and pneumonia (J09-J18)	753	4.4	---
5	Malignant neoplasms (C00-C97)	547	3.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	501	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	430	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	1.4	---
10	Pneumonitis due to solids and liquids (J69)	167	1.0	---
11	Atherosclerosis (I70)	161	0.9	---
12	Diabetes mellitus (E10-E14)	157	0.9	---
13	Septicemia (A40-A41)	135	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.4	---
15	Parkinson's disease (G20-G21)	55	0.3	---
...	All other causes (Residual)	5,300	30.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	299,227	100.0	733.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,102	23.8	174.3
2	Malignant neoplasms (C00-C97)	67,272	22.5	164.9
3	Cerebrovascular diseases (I60-I69)	16,111	5.4	39.5
4	Diabetes mellitus (E10-E14)	13,239	4.4	32.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,169	4.4	32.3
6	Chronic lower respiratory diseases (J40-J47)	9,816	3.3	24.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,340	2.8	20.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,950	2.7	19.5
9	Septicemia (A40-A41)	6,193	2.1	15.2
10	Alzheimer's disease (G30)	5,664	1.9	13.9
11	Influenza and pneumonia (J09-J18)	5,494	1.8	13.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,400	1.8	13.2
13	Certain conditions originating in the perinatal period (P00-P96)	3,854	1.3	9.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,673	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,998	1.0	7.3
...	All other causes (Residual)	58,952	19.7	144.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	963	100.0	39.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	29.6	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.8	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	9.1	3.6
4	Malignant neoplasms (C00-C97)	62	6.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	6.0	2.4
6	Chronic lower respiratory diseases (J40-J47)	26	2.7	1.1
7	Influenza and pneumonia (J09-J18)	21	2.2	0.9
8	Septicemia (A40-A41)	18	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Meningitis (G00,G03)	5	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	219	22.7	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	561	100.0	18.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	31.9	5.8
2	Malignant neoplasms (C00-C97)	64	11.4	2.1
3	Chronic lower respiratory diseases (J40-J47)	44	7.8	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	7.3	1.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	5.9	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.9	0.7
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
9	Septicemia (A40-A41)	7	1.2	*
9	Anemias (D50-D64)	7	1.2	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	119	21.2	3.9
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	571	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	25.4	4.7
2	Malignant neoplasms (C00-C97)	55	9.6	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	8.9	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	8.4	1.5
5	Chronic lower respiratory diseases (J40-J47)	42	7.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	6.8	1.3
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.7	0.9
8	Influenza and pneumonia (J09-J18)	15	2.6	*
9	Cerebrovascular diseases (I60-I69)	11	1.9	*
10	Septicemia (A40-A41)	6	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
12	Anemias (D50-D64)	4	0.7	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Meningitis (G00,G03)	2	0.4	*
13	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	117	20.5	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,146	100.0	65.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	874	40.7	26.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	492	22.9	15.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	7.5	4.9
4	Malignant neoplasms (C00-C97)	102	4.8	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	4.1	2.7
6	Chronic lower respiratory diseases (J40-J47)	34	1.6	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.3	0.8
8	Anemias (D50-D64)	22	1.0	0.7
9	Diabetes mellitus (E10-E14)	20	0.9	0.6
10	Cerebrovascular diseases (I60-I69)	18	0.8	*
11	Influenza and pneumonia (J09-J18)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	13	0.6	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
14	Septicemia (A40-A41)	9	0.4	*
15	Diseases of appendix (K35-K38)	6	0.3	*
...	All other causes (Residual)	258	12.0	7.9
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,277	100.0	120.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,819	42.5	51.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	884	20.7	24.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	353	8.3	9.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	4.7	5.6
5	Malignant neoplasms (C00-C97)	137	3.2	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.9	2.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	58	1.4	1.6
8	Diabetes mellitus (E10-E14)	53	1.2	1.5
9	Anemias (D50-D64)	48	1.1	1.3
10	Chronic lower respiratory diseases (J40-J47)	46	1.1	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	0.8
11	Legal intervention (Y35,Y89.0)	30	0.7	0.8
13	Septicemia (A40-A41)	27	0.6	0.8
14	Cerebrovascular diseases (I60-I69)	20	0.5	0.6
15	Influenza and pneumonia (J09-J18)	18	0.4	*
...	All other causes (Residual)	471	11.0	13.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,394	100.0	148.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,354	30.8	45.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	883	20.1	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	7.9	11.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	291	6.6	9.9
5	Malignant neoplasms (C00-C97)	233	5.3	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	176	4.0	6.0
7	Diabetes mellitus (E10-E14)	99	2.3	3.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	86	2.0	2.9
9	Anemias (D50-D64)	52	1.2	1.8
10	Chronic lower respiratory diseases (J40-J47)	46	1.0	1.6
11	Cerebrovascular diseases (I60-I69)	41	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.3
13	Septicemia (A40-A41)	37	0.8	1.3
14	Influenza and pneumonia (J09-J18)	34	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.8	1.2
...	All other causes (Residual)	642	14.6	21.7
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,988	100.0	177.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,091	21.9	38.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	881	17.7	31.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	624	12.5	22.2
4	Malignant neoplasms (C00-C97)	406	8.1	14.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	5.1	9.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	215	4.3	7.6
7	Diabetes mellitus (E10-E14)	122	2.4	4.3
8	Cerebrovascular diseases (I60-I69)	101	2.0	3.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	1.3	2.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.3	2.2
11	Influenza and pneumonia (J09-J18)	62	1.2	2.2
12	Anemias (D50-D64)	61	1.2	2.2
13	Chronic lower respiratory diseases (J40-J47)	53	1.1	1.9
14	Septicemia (A40-A41)	52	1.0	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.8	1.4
...	All other causes (Residual)	900	18.0	32.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	5,647	100.0	222.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,020	18.1	40.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	14.7	32.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	730	12.9	28.7
4	Malignant neoplasms (C00-C97)	634	11.2	24.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	290	5.1	11.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	3.7	8.3
7	Diabetes mellitus (E10-E14)	189	3.3	7.4
8	Cerebrovascular diseases (I60-I69)	183	3.2	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.7	3.8
10	Chronic lower respiratory diseases (J40-J47)	93	1.6	3.7
11	Septicemia (A40-A41)	80	1.4	3.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.1	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.0	2.2
14	Influenza and pneumonia (J09-J18)	55	1.0	2.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	52	0.9	2.0
...	All other causes (Residual)	1,065	18.9	41.9
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	7,932	100.0	297.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,750	22.1	65.6
2	Malignant neoplasms (C00-C97)	1,258	15.9	47.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	916	11.5	34.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	495	6.2	18.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	385	4.9	14.4
6	Diabetes mellitus (E10-E14)	342	4.3	12.8
7	Cerebrovascular diseases (I60-I69)	302	3.8	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	216	2.7	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	1.8	5.5
10	Septicemia (A40-A41)	145	1.8	5.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.6	4.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.5	4.5
13	Chronic lower respiratory diseases (J40-J47)	108	1.4	4.0
14	Influenza and pneumonia (J09-J18)	102	1.3	3.8
15	Aortic aneurysm and dissection (I71)	63	0.8	2.4
...	All other causes (Residual)	1,456	18.4	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	12,316	100.0	456.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,886	23.4	107.0
2	Malignant neoplasms (C00-C97)	2,529	20.5	93.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,069	8.7	39.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	541	4.4	20.1
5	Cerebrovascular diseases (I60-I69)	538	4.4	19.9
6	Diabetes mellitus (E10-E14)	536	4.4	19.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	3.5	15.8
8	Septicemia (A40-A41)	251	2.0	9.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	247	2.0	9.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.0	9.1
11	Chronic lower respiratory diseases (J40-J47)	219	1.8	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.6	7.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	1.6	7.1
14	Influenza and pneumonia (J09-J18)	162	1.3	6.0
15	Aortic aneurysm and dissection (I71)	78	0.6	2.9
...	All other causes (Residual)	2,200	17.9	81.5
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,444	100.0	704.6
1	Malignant neoplasms (C00-C97)	5,060	26.0	183.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,699	24.2	170.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,330	6.8	48.2
4	Cerebrovascular diseases (I60-I69)	917	4.7	33.2
5	Diabetes mellitus (E10-E14)	865	4.4	31.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	621	3.2	22.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	488	2.5	17.7
8	Chronic lower respiratory diseases (J40-J47)	432	2.2	15.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	429	2.2	15.5
10	Septicemia (A40-A41)	385	2.0	14.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	352	1.8	12.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	334	1.7	12.1
13	Influenza and pneumonia (J09-J18)	286	1.5	10.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	0.9	6.1
15	Viral hepatitis (B15-B19)	149	0.8	5.4
...	All other causes (Residual)	2,928	15.1	106.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	26,913	100.0	1,089.7
1	Malignant neoplasms (C00-C97)	7,889	29.3	319.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,648	24.7	269.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,273	4.7	51.5
4	Cerebrovascular diseases (I60-I69)	1,268	4.7	51.3
5	Diabetes mellitus (E10-E14)	1,252	4.7	50.7
6	Chronic lower respiratory diseases (J40-J47)	728	2.7	29.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	686	2.5	27.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	617	2.3	25.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	588	2.2	23.8
10	Septicemia (A40-A41)	519	1.9	21.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	441	1.6	17.9
12	Influenza and pneumonia (J09-J18)	414	1.5	16.8
13	Viral hepatitis (B15-B19)	313	1.2	12.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	229	0.9	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	0.6	6.1
...	All other causes (Residual)	3,898	14.5	157.8
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	30,567	100.0	1,569.2
1	Malignant neoplasms (C00-C97)	9,642	31.5	495.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,506	24.6	385.3
3	Diabetes mellitus (E10-E14)	1,611	5.3	82.7
4	Cerebrovascular diseases (I60-I69)	1,537	5.0	78.9
5	Chronic lower respiratory diseases (J40-J47)	1,057	3.5	54.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	3.4	53.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	846	2.8	43.4
8	Septicemia (A40-A41)	648	2.1	33.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	640	2.1	32.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	464	1.5	23.8
11	Influenza and pneumonia (J09-J18)	440	1.4	22.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	377	1.2	19.4
13	Viral hepatitis (B15-B19)	323	1.1	16.6
14	Pneumonitis due to solids and liquids (J69)	122	0.4	6.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	0.4	6.2
...	All other causes (Residual)	4,200	13.7	215.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	28,682	100.0	2,083.6
1	Malignant neoplasms (C00-C97)	9,171	32.0	666.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,209	25.1	523.7
3	Diabetes mellitus (E10-E14)	1,537	5.4	111.7
4	Cerebrovascular diseases (I60-I69)	1,505	5.2	109.3
5	Chronic lower respiratory diseases (J40-J47)	1,137	4.0	82.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	867	3.0	63.0
7	Septicemia (A40-A41)	688	2.4	50.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	601	2.1	43.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	523	1.8	38.0
10	Influenza and pneumonia (J09-J18)	490	1.7	35.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	330	1.2	24.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	198	0.7	14.4
13	Viral hepatitis (B15-B19)	165	0.6	12.0
14	Pneumonitis due to solids and liquids (J69)	137	0.5	10.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.4	9.3
...	All other causes (Residual)	3,996	13.9	290.3
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	28,826	100.0	3,013.2
1	Malignant neoplasms (C00-C97)	8,607	29.9	899.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,183	24.9	750.9
3	Cerebrovascular diseases (I60-I69)	1,683	5.8	175.9
4	Diabetes mellitus (E10-E14)	1,606	5.6	167.9
5	Chronic lower respiratory diseases (J40-J47)	1,359	4.7	142.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,016	3.5	106.2
7	Septicemia (A40-A41)	649	2.3	67.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	562	1.9	58.7
9	Influenza and pneumonia (J09-J18)	548	1.9	57.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	1.6	49.0
11	Alzheimer's disease (G30)	311	1.1	32.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	181	0.6	18.9
13	Pneumonitis due to solids and liquids (J69)	169	0.6	17.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.4	12.4
15	Parkinson's disease (G20-G21)	116	0.4	12.1
...	All other causes (Residual)	4,248	14.7	444.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	29,473	100.0	4,371.5
1	Malignant neoplasms (C00-C97)	7,691	26.1	1,140.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,410	25.1	1,099.1
3	Cerebrovascular diseases (I60-I69)	1,934	6.6	286.9
4	Diabetes mellitus (E10-E14)	1,606	5.4	238.2
5	Chronic lower respiratory diseases (J40-J47)	1,302	4.4	193.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,027	3.5	152.3
7	Septicemia (A40-A41)	736	2.5	109.2
8	Alzheimer's disease (G30)	663	2.2	98.3
9	Influenza and pneumonia (J09-J18)	581	2.0	86.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	560	1.9	83.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	427	1.4	63.3
12	Parkinson's disease (G20-G21)	208	0.7	30.9
13	Pneumonitis due to solids and liquids (J69)	207	0.7	30.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	22.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	128	0.4	19.0
...	All other causes (Residual)	4,844	16.4	718.5
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,751	100.0	6,620.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,789	26.2	1,733.3
2	Malignant neoplasms (C00-C97)	6,456	21.7	1,436.6
3	Cerebrovascular diseases (I60-I69)	2,139	7.2	476.0
4	Diabetes mellitus (E10-E14)	1,381	4.6	307.3
5	Chronic lower respiratory diseases (J40-J47)	1,241	4.2	276.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,091	3.7	242.8
7	Alzheimer's disease (G30)	1,087	3.7	241.9
8	Septicemia (A40-A41)	694	2.3	154.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	652	2.2	145.1
10	Influenza and pneumonia (J09-J18)	640	2.2	142.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	1.2	81.4
12	Pneumonitis due to solids and liquids (J69)	229	0.8	51.0
13	Parkinson's disease (G20-G21)	227	0.8	50.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.5	34.5
15	Aortic aneurysm and dissection (I71)	100	0.3	22.3
...	All other causes (Residual)	5,504	18.5	1,224.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	27,119	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,369	27.2	---
2	Malignant neoplasms (C00-C97)	4,359	16.1	---
3	Cerebrovascular diseases (I60-I69)	1,884	6.9	---
4	Alzheimer's disease (G30)	1,428	5.3	---
5	Diabetes mellitus (E10-E14)	1,143	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	1,073	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	912	3.4	---
8	Influenza and pneumonia (J09-J18)	739	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	624	2.3	---
10	Septicemia (A40-A41)	617	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	1.3	---
12	Pneumonitis due to solids and liquids (J69)	250	0.9	---
13	Parkinson's disease (G20-G21)	232	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.5	---
15	Aortic aneurysm and dissection (I71)	85	0.3	---
...	All other causes (Residual)	5,927	21.9	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	18,027	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,208	28.9	---
2	Malignant neoplasms (C00-C97)	2,193	12.2	---
3	Cerebrovascular diseases (I60-I69)	1,300	7.2	---
4	Alzheimer's disease (G30)	1,203	6.7	---
5	Diabetes mellitus (E10-E14)	596	3.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	570	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	557	3.1	---
8	Influenza and pneumonia (J09-J18)	497	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	2.6	---
10	Septicemia (A40-A41)	366	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	1.3	---
12	Pneumonitis due to solids and liquids (J69)	164	0.9	---
13	Parkinson's disease (G20-G21)	95	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	---
15	Atherosclerosis (I70)	75	0.4	---
...	All other causes (Residual)	4,423	24.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	7,361	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,222	30.2	---
2	Malignant neoplasms (C00-C97)	593	8.1	---
3	Alzheimer's disease (G30)	579	7.9	---
4	Cerebrovascular diseases (I60-I69)	486	6.6	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	3.1	---
6	Influenza and pneumonia (J09-J18)	219	3.0	---
7	Diabetes mellitus (E10-E14)	212	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.9	---
9	Chronic lower respiratory diseases (J40-J47)	169	2.3	---
10	Septicemia (A40-A41)	142	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	1.2	---
12	Pneumonitis due to solids and liquids (J69)	64	0.9	---
13	Anemias (D50-D64)	28	0.4	---
14	Parkinson's disease (G20-G21)	27	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	---
...	All other causes (Residual)	2,065	28.1	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,468	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	741	30.0	---
2	Alzheimer's disease (G30)	201	8.1	---
3	Cerebrovascular diseases (I60-I69)	182	7.4	---
4	Malignant neoplasms (C00-C97)	122	4.9	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	3.5	---
6	Influenza and pneumonia (J09-J18)	76	3.1	---
7	Septicemia (A40-A41)	63	2.6	---
7	Diabetes mellitus (E10-E14)	63	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.1	---
10	Chronic lower respiratory diseases (J40-J47)	39	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	1.2	---
12	Pneumonitis due to solids and liquids (J69)	19	0.8	---
13	Atherosclerosis (I70)	18	0.7	---
14	Anemias (D50-D64)	11	0.4	---
15	Nutritional deficiencies (E40-E64)	10	0.4	---
...	All other causes (Residual)	755	30.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	152,661	100.0	782.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,562	23.9	187.4
2	Malignant neoplasms (C00-C97)	34,319	22.5	175.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,845	5.8	45.3
4	Cerebrovascular diseases (I60-I69)	7,266	4.8	37.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,851	4.5	35.1
6	Diabetes mellitus (E10-E14)	6,298	4.1	32.3
7	Chronic lower respiratory diseases (J40-J47)	5,019	3.3	25.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,944	2.6	20.2
9	Septicemia (A40-A41)	2,854	1.9	14.6
10	Influenza and pneumonia (J09-J18)	2,657	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,467	1.6	12.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,429	1.6	12.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,102	1.4	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,915	1.3	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.2	9.5
...	All other causes (Residual)	27,283	17.9	139.8
Non-Hispanic, black, male, 1-4 years				
...	All causes	538	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.7	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	12.6	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.7	2.9
5	Malignant neoplasms (C00-C97)	33	6.1	2.7
6	Chronic lower respiratory diseases (J40-J47)	15	2.8	*
7	Septicemia (A40-A41)	12	2.2	*
7	Cerebrovascular diseases (I60-I69)	12	2.2	*
7	Influenza and pneumonia (J09-J18)	12	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	115	21.4	9.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	308	100.0	19.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	33.8	6.6
2	Malignant neoplasms (C00-C97)	32	10.4	2.0
3	Chronic lower respiratory diseases (J40-J47)	28	9.1	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	6.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.2	*
7	Influenza and pneumonia (J09-J18)	8	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.3	*
9	Septicemia (A40-A41)	4	1.3	*
9	Anemias (D50-D64)	4	1.3	*
11	Cerebrovascular diseases (I60-I69)	3	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	1.0	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	60	19.5	3.8
Non-Hispanic, black, male, 10-14 years				
...	All causes	345	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	30.7	6.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	9.9	2.2
3	Malignant neoplasms (C00-C97)	33	9.6	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	9.0	2.0
5	Chronic lower respiratory diseases (J40-J47)	26	7.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.2	*
8	Influenza and pneumonia (J09-J18)	10	2.9	*
9	Cerebrovascular diseases (I60-I69)	7	2.0	*
10	Septicemia (A40-A41)	4	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Anemias (D50-D64)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	50	14.5	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,660	100.0	100.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	798	48.1	48.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	21.7	21.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	7.4	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.4	3.4
5	Malignant neoplasms (C00-C97)	56	3.4	3.4
6	Chronic lower respiratory diseases (J40-J47)	22	1.3	1.3
7	Diabetes mellitus (E10-E14)	16	1.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.0	*
9	Anemias (D50-D64)	13	0.8	*
9	Legal intervention (Y35,Y89.0)	13	0.8	*
11	Cerebrovascular diseases (I60-I69)	10	0.6	*
12	Influenza and pneumonia (J09-J18)	7	0.4	*
13	Septicemia (A40-A41)	4	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
14	Meningitis (G00,G03)	3	0.2	*
14	Diseases of appendix (K35-K38)	3	0.2	*
...	All other causes (Residual)	156	9.4	9.4
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,297	100.0	183.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,652	50.1	92.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	635	19.3	35.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	298	9.0	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	4.1	7.5
5	Malignant neoplasms (C00-C97)	72	2.2	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	59	1.8	3.3
7	Chronic lower respiratory diseases (J40-J47)	33	1.0	1.8
8	Diabetes mellitus (E10-E14)	32	1.0	1.8
9	Anemias (D50-D64)	31	0.9	1.7
10	Legal intervention (Y35,Y89.0)	30	0.9	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
12	Septicemia (A40-A41)	15	0.5	*
13	Cerebrovascular diseases (I60-I69)	11	0.3	*
14	Influenza and pneumonia (J09-J18)	9	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	260	7.9	14.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,156	100.0	218.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,210	38.3	83.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	652	20.7	45.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	7.7	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	7.1	15.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	131	4.2	9.1
5	Malignant neoplasms (C00-C97)	131	4.2	9.1
7	Diabetes mellitus (E10-E14)	52	1.6	3.6
8	Anemias (D50-D64)	32	1.0	2.2
9	Chronic lower respiratory diseases (J40-J47)	28	0.9	1.9
10	Cerebrovascular diseases (I60-I69)	25	0.8	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
12	Septicemia (A40-A41)	17	0.5	*
13	Influenza and pneumonia (J09-J18)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
13	Legal intervention (Y35,Y89.0)	15	0.5	*
...	All other causes (Residual)	348	11.0	24.1
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,333	100.0	248.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	964	28.9	71.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	655	19.7	48.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	404	12.1	30.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	6.4	15.8
5	Malignant neoplasms (C00-C97)	176	5.3	13.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	139	4.2	10.4
7	Diabetes mellitus (E10-E14)	65	2.0	4.8
8	Cerebrovascular diseases (I60-I69)	48	1.4	3.6
9	Influenza and pneumonia (J09-J18)	39	1.2	2.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	2.5
11	Anemias (D50-D64)	26	0.8	1.9
12	Chronic lower respiratory diseases (J40-J47)	25	0.8	1.9
12	Legal intervention (Y35,Y89.0)	25	0.8	1.9
14	Septicemia (A40-A41)	24	0.7	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.7	1.6
...	All other causes (Residual)	475	14.3	35.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,492	100.0	291.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	664	19.0	55.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	646	18.5	53.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	16.9	49.3
4	Malignant neoplasms (C00-C97)	253	7.2	21.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	5.3	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	161	4.6	13.4
7	Diabetes mellitus (E10-E14)	110	3.2	9.2
8	Cerebrovascular diseases (I60-I69)	99	2.8	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.5	4.5
10	Chronic lower respiratory diseases (J40-J47)	50	1.4	4.2
11	Septicemia (A40-A41)	40	1.1	3.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.0	2.9
13	Anemias (D50-D64)	29	0.8	2.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.8	2.3
15	Aortic aneurysm and dissection (I71)	27	0.8	2.3
...	All other causes (Residual)	521	14.9	43.4
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,613	100.0	367.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,129	24.5	90.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	633	13.7	50.4
3	Malignant neoplasms (C00-C97)	455	9.9	36.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	410	8.9	32.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	223	4.8	17.8
6	Diabetes mellitus (E10-E14)	209	4.5	16.7
7	Cerebrovascular diseases (I60-I69)	165	3.6	13.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	3.5	12.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	1.8	6.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	1.6	5.9
11	Septicemia (A40-A41)	70	1.5	5.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.5	5.5
13	Chronic lower respiratory diseases (J40-J47)	67	1.5	5.3
14	Influenza and pneumonia (J09-J18)	56	1.2	4.5
15	Aortic aneurysm and dissection (I71)	49	1.1	3.9
...	All other causes (Residual)	757	16.4	60.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,058	100.0	556.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,822	25.8	143.5
2	Malignant neoplasms (C00-C97)	1,110	15.7	87.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	704	10.0	55.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	354	5.0	27.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	336	4.8	26.5
6	Cerebrovascular diseases (I60-I69)	310	4.4	24.4
7	Diabetes mellitus (E10-E14)	305	4.3	24.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	148	2.1	11.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	2.0	11.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	2.0	10.9
11	Septicemia (A40-A41)	138	2.0	10.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.9	10.5
13	Chronic lower respiratory diseases (J40-J47)	105	1.5	8.3
14	Influenza and pneumonia (J09-J18)	99	1.4	7.8
15	Aortic aneurysm and dissection (I71)	57	0.8	4.5
...	All other causes (Residual)	1,156	16.4	91.1
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,014	100.0	852.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,932	26.6	227.0
2	Malignant neoplasms (C00-C97)	2,416	21.9	187.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	921	8.4	71.3
4	Cerebrovascular diseases (I60-I69)	508	4.6	39.3
5	Diabetes mellitus (E10-E14)	461	4.2	35.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	413	3.7	32.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	300	2.7	23.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	2.4	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	2.4	20.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	207	1.9	16.0
11	Chronic lower respiratory diseases (J40-J47)	201	1.8	15.6
12	Septicemia (A40-A41)	197	1.8	15.3
13	Influenza and pneumonia (J09-J18)	167	1.5	12.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	1.3	10.7
15	Viral hepatitis (B15-B19)	98	0.9	7.6
...	All other causes (Residual)	1,530	13.9	118.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	15,790	100.0	1,386.3
1	Malignant neoplasms (C00-C97)	4,310	27.3	378.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,183	26.5	367.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	5.9	81.6
4	Diabetes mellitus (E10-E14)	722	4.6	63.4
5	Cerebrovascular diseases (I60-I69)	710	4.5	62.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	418	2.6	36.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	398	2.5	34.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	366	2.3	32.1
9	Chronic lower respiratory diseases (J40-J47)	360	2.3	31.6
10	Septicemia (A40-A41)	266	1.7	23.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	261	1.7	22.9
12	Influenza and pneumonia (J09-J18)	226	1.4	19.8
13	Viral hepatitis (B15-B19)	203	1.3	17.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	1.2	16.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	0.7	9.6
...	All other causes (Residual)	2,141	13.6	188.0
Non-Hispanic, black, male, 60-64 years				
...	All causes	17,854	100.0	2,045.3
1	Malignant neoplasms (C00-C97)	5,301	29.7	607.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,687	26.3	536.9
3	Cerebrovascular diseases (I60-I69)	908	5.1	104.0
4	Diabetes mellitus (E10-E14)	877	4.9	100.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	4.2	85.2
6	Chronic lower respiratory diseases (J40-J47)	546	3.1	62.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	503	2.8	57.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	448	2.5	51.3
9	Septicemia (A40-A41)	332	1.9	38.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	276	1.5	31.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	258	1.4	29.6
12	Influenza and pneumonia (J09-J18)	251	1.4	28.8
13	Viral hepatitis (B15-B19)	228	1.3	26.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	0.5	10.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	0.4	8.8
...	All other causes (Residual)	2,330	13.1	266.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	16,079	100.0	2,671.5
1	Malignant neoplasms (C00-C97)	5,038	31.3	837.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,279	26.6	711.0
3	Cerebrovascular diseases (I60-I69)	802	5.0	133.3
4	Diabetes mellitus (E10-E14)	795	4.9	132.1
5	Chronic lower respiratory diseases (J40-J47)	618	3.8	102.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	438	2.7	72.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	2.6	68.6
8	Septicemia (A40-A41)	347	2.2	57.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	1.9	49.8
10	Influenza and pneumonia (J09-J18)	269	1.7	44.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	236	1.5	39.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	138	0.9	22.9
13	Viral hepatitis (B15-B19)	115	0.7	19.1
14	Pneumonitis due to solids and liquids (J69)	87	0.5	14.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.5	13.1
...	All other causes (Residual)	2,125	13.2	353.1
Non-Hispanic, black, male, 70-74 years				
...	All causes	15,354	100.0	3,837.6
1	Malignant neoplasms (C00-C97)	4,638	30.2	1,159.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,975	25.9	993.5
3	Cerebrovascular diseases (I60-I69)	895	5.8	223.7
4	Diabetes mellitus (E10-E14)	777	5.1	194.2
5	Chronic lower respiratory diseases (J40-J47)	776	5.1	194.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	3.0	116.7
7	Septicemia (A40-A41)	323	2.1	80.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	2.0	76.0
9	Influenza and pneumonia (J09-J18)	301	2.0	75.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	277	1.8	69.2
11	Alzheimer's disease (G30)	131	0.9	32.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	120	0.8	30.0
13	Pneumonitis due to solids and liquids (J69)	101	0.7	25.2
14	Parkinson's disease (G20-G21)	80	0.5	20.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	67	0.4	16.7
...	All other causes (Residual)	2,122	13.8	530.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	14,652	100.0	5,530.0
1	Malignant neoplasms (C00-C97)	3,966	27.1	1,496.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,796	25.9	1,432.7
3	Cerebrovascular diseases (I60-I69)	885	6.0	334.0
4	Diabetes mellitus (E10-E14)	729	5.0	275.1
5	Chronic lower respiratory diseases (J40-J47)	700	4.8	264.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	481	3.3	181.5
7	Septicemia (A40-A41)	326	2.2	123.0
8	Influenza and pneumonia (J09-J18)	292	2.0	110.2
9	Alzheimer's disease (G30)	269	1.8	101.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	1.8	97.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	1.8	97.4
12	Parkinson's disease (G20-G21)	138	0.9	52.1
13	Pneumonitis due to solids and liquids (J69)	128	0.9	48.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	29.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.4	24.2
...	All other causes (Residual)	2,284	15.6	862.0
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,930	100.0	8,117.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,369	26.1	2,115.0
2	Malignant neoplasms (C00-C97)	3,132	24.2	1,966.2
3	Cerebrovascular diseases (I60-I69)	801	6.2	502.8
4	Chronic lower respiratory diseases (J40-J47)	614	4.7	385.5
5	Diabetes mellitus (E10-E14)	541	4.2	339.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	3.5	287.5
7	Alzheimer's disease (G30)	379	2.9	237.9
8	Septicemia (A40-A41)	293	2.3	183.9
9	Influenza and pneumonia (J09-J18)	282	2.2	177.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	267	2.1	167.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	1.5	120.5
12	Parkinson's disease (G20-G21)	143	1.1	89.8
13	Pneumonitis due to solids and liquids (J69)	111	0.9	69.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	43.9
15	Aortic aneurysm and dissection (I71)	41	0.3	25.7
...	All other causes (Residual)	2,237	17.3	1,404.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,055	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,693	26.8	---
2	Malignant neoplasms (C00-C97)	1,972	19.6	---
3	Cerebrovascular diseases (I60-I69)	601	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	503	5.0	---
5	Alzheimer's disease (G30)	410	4.1	---
6	Diabetes mellitus (E10-E14)	382	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	3.3	---
8	Influenza and pneumonia (J09-J18)	310	3.1	---
9	Septicemia (A40-A41)	240	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	1.6	---
12	Pneumonitis due to solids and liquids (J69)	125	1.2	---
13	Parkinson's disease (G20-G21)	118	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.5	---
15	Atherosclerosis (I70)	29	0.3	---
...	All other causes (Residual)	1,927	19.2	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	5,538	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,554	28.1	---
2	Malignant neoplasms (C00-C97)	936	16.9	---
3	Cerebrovascular diseases (I60-I69)	323	5.8	---
4	Alzheimer's disease (G30)	258	4.7	---
5	Chronic lower respiratory diseases (J40-J47)	234	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	3.7	---
7	Diabetes mellitus (E10-E14)	169	3.1	---
8	Influenza and pneumonia (J09-J18)	166	3.0	---
9	Septicemia (A40-A41)	128	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	119	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.4	---
12	Pneumonitis due to solids and liquids (J69)	66	1.2	---
13	Parkinson's disease (G20-G21)	38	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	---
15	Anemias (D50-D64)	19	0.3	---
15	Nutritional deficiencies (E40-E64)	19	0.3	---
...	All other causes (Residual)	1,191	21.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,563	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	458	29.3	---
2	Malignant neoplasms (C00-C97)	218	13.9	---
3	Cerebrovascular diseases (I60-I69)	98	6.3	---
4	Alzheimer's disease (G30)	80	5.1	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	4.0	---
6	Influenza and pneumonia (J09-J18)	56	3.6	---
7	Chronic lower respiratory diseases (J40-J47)	49	3.1	---
8	Diabetes mellitus (E10-E14)	47	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	2.2	---
10	Septicemia (A40-A41)	33	2.1	---
11	Pneumonitis due to solids and liquids (J69)	22	1.4	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.2	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.6	---
14	Parkinson's disease (G20-G21)	9	0.6	---
15	Anemias (D50-D64)	4	0.3	---
15	Atherosclerosis (I70)	4	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.3	---
...	All other causes (Residual)	352	22.5	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	321	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	25.9	---
2	Malignant neoplasms (C00-C97)	39	12.1	---
3	Cerebrovascular diseases (I60-I69)	22	6.9	---
4	Influenza and pneumonia (J09-J18)	19	5.9	---
5	Alzheimer's disease (G30)	15	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	13	4.0	---
7	Septicemia (A40-A41)	11	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	3.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.5	---
10	Diabetes mellitus (E10-E14)	6	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.9	---
12	Atherosclerosis (I70)	3	0.9	---
13	Anemias (D50-D64)	2	0.6	---
13	Pneumonitis due to solids and liquids (J69)	2	0.6	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	---
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.6	---
...	All other causes (Residual)	78	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	146,566	100.0	688.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,540	23.6	162.2
2	Malignant neoplasms (C00-C97)	32,953	22.5	154.8
3	Cerebrovascular diseases (I60-I69)	8,845	6.0	41.5
4	Diabetes mellitus (E10-E14)	6,941	4.7	32.6
5	Chronic lower respiratory diseases (J40-J47)	4,797	3.3	22.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,396	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,324	3.0	20.3
8	Alzheimer's disease (G30)	4,018	2.7	18.9
9	Septicemia (A40-A41)	3,339	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,933	2.0	13.8
11	Influenza and pneumonia (J09-J18)	2,837	1.9	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,752	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,244	0.8	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.7	5.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,083	0.7	5.1
...	All other causes (Residual)	31,465	21.5	147.8
Non-Hispanic, black, female, 1-4 years				
...	All causes	425	100.0	35.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	28.2	10.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	15.3	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	8.9	3.2
4	Malignant neoplasms (C00-C97)	29	6.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	11	2.6	*
7	Influenza and pneumonia (J09-J18)	9	2.1	*
8	Septicemia (A40-A41)	6	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	98	23.1	8.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	253	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	29.6	4.9
2	Malignant neoplasms (C00-C97)	32	12.6	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	8.7	1.4
4	Chronic lower respiratory diseases (J40-J47)	16	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.7	*
7	Cerebrovascular diseases (I60-I69)	4	1.6	*
7	Influenza and pneumonia (J09-J18)	4	1.6	*
9	Septicemia (A40-A41)	3	1.2	*
9	Anemias (D50-D64)	3	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
15	Meningococcal infection (A39)	1	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	58	22.9	3.8
Non-Hispanic, black, female, 10-14 years				
...	All causes	226	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	17.3	2.6
2	Malignant neoplasms (C00-C97)	22	9.7	1.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	9.3	1.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	8.8	1.3
5	Chronic lower respiratory diseases (J40-J47)	16	7.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.5	*
8	Influenza and pneumonia (J09-J18)	5	2.2	*
9	Cerebrovascular diseases (I60-I69)	4	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
10	Anemias (D50-D64)	3	1.3	*
12	Septicemia (A40-A41)	2	0.9	*
12	Meningitis (G00,G03)	2	0.9	*
12	Pneumonitis due to solids and liquids (J69)	2	0.9	*
15	Viral hepatitis (B15-B19)	1	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
15	Peptic ulcer (K25-K28)	1	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	59	26.1	3.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	486	100.0	30.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	27.2	8.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	15.6	4.7
3	Malignant neoplasms (C00-C97)	46	9.5	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	7.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	6.2	1.9
6	Chronic lower respiratory diseases (J40-J47)	12	2.5	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	2.3	*
9	Anemias (D50-D64)	9	1.9	*
10	Cerebrovascular diseases (I60-I69)	8	1.6	*
11	Influenza and pneumonia (J09-J18)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Diabetes mellitus (E10-E14)	4	0.8	*
14	Diseases of appendix (K35-K38)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	92	18.9	5.7
Non-Hispanic, black, female, 20-24 years				
...	All causes	980	100.0	55.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	25.4	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	167	17.0	9.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	6.7	3.7
4	Malignant neoplasms (C00-C97)	65	6.6	3.7
5	Pregnancy, childbirth and the puerperium (O00-O99)	58	5.9	3.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	5.6	3.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	24	2.4	1.4
8	Diabetes mellitus (E10-E14)	21	2.1	1.2
9	Anemias (D50-D64)	17	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
11	Septicemia (A40-A41)	12	1.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.1	*
13	Cerebrovascular diseases (I60-I69)	9	0.9	*
13	Influenza and pneumonia (J09-J18)	9	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	198	20.2	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,238	100.0	82.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	18.7	15.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	11.6	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	9.9	8.1
4	Malignant neoplasms (C00-C97)	102	8.2	6.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	86	6.9	5.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	4.0	3.2
7	Diabetes mellitus (E10-E14)	47	3.8	3.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	45	3.6	3.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.7	1.4
10	Septicemia (A40-A41)	20	1.6	1.3
10	Anemias (D50-D64)	20	1.6	1.3
12	Influenza and pneumonia (J09-J18)	19	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
14	Chronic lower respiratory diseases (J40-J47)	18	1.5	*
15	Cerebrovascular diseases (I60-I69)	16	1.3	*
...	All other causes (Residual)	279	22.5	18.5
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,655	100.0	112.5
1	Malignant neoplasms (C00-C97)	230	13.9	15.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	13.7	15.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	13.3	15.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	127	7.7	8.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	76	4.6	5.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	63	3.8	4.3
7	Diabetes mellitus (E10-E14)	57	3.4	3.9
8	Cerebrovascular diseases (I60-I69)	53	3.2	3.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.5	2.8
10	Anemias (D50-D64)	35	2.1	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.9	2.2
12	Septicemia (A40-A41)	28	1.7	1.9
12	Chronic lower respiratory diseases (J40-J47)	28	1.7	1.9
14	Influenza and pneumonia (J09-J18)	23	1.4	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
...	All other causes (Residual)	400	24.2	27.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,155	100.0	160.3
1	Malignant neoplasms (C00-C97)	381	17.7	28.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	16.5	26.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	11.1	17.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	129	6.0	9.6
5	Cerebrovascular diseases (I60-I69)	84	3.9	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	3.9	6.2
7	Diabetes mellitus (E10-E14)	79	3.7	5.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	52	2.4	3.9
9	Chronic lower respiratory diseases (J40-J47)	43	2.0	3.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.9	3.1
11	Septicemia (A40-A41)	40	1.9	3.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.6	2.6
13	Influenza and pneumonia (J09-J18)	32	1.5	2.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	1.2	1.9
15	Anemias (D50-D64)	22	1.0	1.6
...	All other causes (Residual)	510	23.7	37.9
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,319	100.0	235.0
1	Malignant neoplasms (C00-C97)	803	24.2	56.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	18.7	44.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	8.5	20.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	162	4.9	11.5
5	Cerebrovascular diseases (I60-I69)	137	4.1	9.7
6	Diabetes mellitus (E10-E14)	133	4.0	9.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	2.6	6.0
8	Septicemia (A40-A41)	75	2.3	5.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.8	4.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.8	4.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	1.6	3.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.4	3.3
12	Influenza and pneumonia (J09-J18)	46	1.4	3.3
14	Chronic lower respiratory diseases (J40-J47)	41	1.2	2.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	34	1.0	2.4
...	All other causes (Residual)	679	20.5	48.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,258	100.0	368.1
1	Malignant neoplasms (C00-C97)	1,419	27.0	99.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,064	20.2	74.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	6.9	25.6
4	Diabetes mellitus (E10-E14)	231	4.4	16.2
5	Cerebrovascular diseases (I60-I69)	228	4.3	16.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	187	3.6	13.1
7	Chronic lower respiratory diseases (J40-J47)	114	2.2	8.0
8	Septicemia (A40-A41)	113	2.1	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.0	7.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.9	6.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	1.7	6.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	1.2	4.4
12	Influenza and pneumonia (J09-J18)	63	1.2	4.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	1.0	3.6
15	Anemias (D50-D64)	30	0.6	2.1
...	All other causes (Residual)	1,035	19.7	72.5
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,430	100.0	574.3
1	Malignant neoplasms (C00-C97)	2,644	31.4	180.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,767	21.0	120.4
3	Cerebrovascular diseases (I60-I69)	409	4.9	27.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	4.9	27.9
5	Diabetes mellitus (E10-E14)	404	4.8	27.5
6	Chronic lower respiratory diseases (J40-J47)	231	2.7	15.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	208	2.5	14.2
8	Septicemia (A40-A41)	188	2.2	12.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	188	2.2	12.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	2.0	11.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.7	9.9
12	Influenza and pneumonia (J09-J18)	119	1.4	8.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.8	4.8
14	Viral hepatitis (B15-B19)	51	0.6	3.5
15	Anemias (D50-D64)	36	0.4	2.5
...	All other causes (Residual)	1,393	16.5	94.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	11,123	100.0	835.9
1	Malignant neoplasms (C00-C97)	3,579	32.2	268.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,465	22.2	185.2
3	Cerebrovascular diseases (I60-I69)	558	5.0	41.9
4	Diabetes mellitus (E10-E14)	530	4.8	39.8
5	Chronic lower respiratory diseases (J40-J47)	368	3.3	27.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	3.1	25.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	2.9	24.0
8	Septicemia (A40-A41)	253	2.3	19.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	199	1.8	15.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	190	1.7	14.3
11	Influenza and pneumonia (J09-J18)	188	1.7	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	180	1.6	13.5
13	Viral hepatitis (B15-B19)	110	1.0	8.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	3.8
15	Aortic aneurysm and dissection (I71)	41	0.4	3.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.4	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.4	3.1
...	All other causes (Residual)	1,665	15.0	125.1
Non-Hispanic, black, female, 60-64 years				
...	All causes	12,713	100.0	1,182.6
1	Malignant neoplasms (C00-C97)	4,341	34.1	403.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,819	22.2	262.2
3	Diabetes mellitus (E10-E14)	734	5.8	68.3
4	Cerebrovascular diseases (I60-I69)	629	4.9	58.5
5	Chronic lower respiratory diseases (J40-J47)	511	4.0	47.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	2.7	31.9
7	Septicemia (A40-A41)	316	2.5	29.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	2.3	26.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.6	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	192	1.5	17.9
11	Influenza and pneumonia (J09-J18)	189	1.5	17.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	101	0.8	9.4
13	Viral hepatitis (B15-B19)	95	0.7	8.8
14	Pneumonitis due to solids and liquids (J69)	48	0.4	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.3	4.0
...	All other causes (Residual)	1,857	14.6	172.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	12,603	100.0	1,626.8
1	Malignant neoplasms (C00-C97)	4,133	32.8	533.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,930	23.2	378.2
3	Diabetes mellitus (E10-E14)	742	5.9	95.8
4	Cerebrovascular diseases (I60-I69)	703	5.6	90.7
5	Chronic lower respiratory diseases (J40-J47)	519	4.1	67.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	429	3.4	55.4
7	Septicemia (A40-A41)	341	2.7	44.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	1.8	28.8
9	Influenza and pneumonia (J09-J18)	221	1.8	28.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	1.5	24.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	94	0.7	12.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	60	0.5	7.7
13	Alzheimer's disease (G30)	52	0.4	6.7
14	Viral hepatitis (B15-B19)	50	0.4	6.5
14	Pneumonitis due to solids and liquids (J69)	50	0.4	6.5
...	All other causes (Residual)	1,868	14.8	241.1
Non-Hispanic, black, female, 70-74 years				
...	All causes	13,472	100.0	2,420.6
1	Malignant neoplasms (C00-C97)	3,969	29.5	713.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,208	23.8	576.4
3	Diabetes mellitus (E10-E14)	829	6.2	149.0
4	Cerebrovascular diseases (I60-I69)	788	5.8	141.6
5	Chronic lower respiratory diseases (J40-J47)	583	4.3	104.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	549	4.1	98.6
7	Septicemia (A40-A41)	326	2.4	58.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	2.1	51.2
9	Influenza and pneumonia (J09-J18)	247	1.8	44.4
10	Alzheimer's disease (G30)	180	1.3	32.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	1.2	29.6
12	Pneumonitis due to solids and liquids (J69)	68	0.5	12.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	61	0.5	11.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.4	9.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.3	6.8
...	All other causes (Residual)	2,123	15.8	381.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,821	100.0	3,621.4
1	Malignant neoplasms (C00-C97)	3,725	25.1	910.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,614	24.4	883.1
3	Cerebrovascular diseases (I60-I69)	1,049	7.1	256.3
4	Diabetes mellitus (E10-E14)	877	5.9	214.3
5	Chronic lower respiratory diseases (J40-J47)	602	4.1	147.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	3.7	133.4
7	Septicemia (A40-A41)	410	2.8	100.2
8	Alzheimer's disease (G30)	394	2.7	96.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	301	2.0	73.5
10	Influenza and pneumonia (J09-J18)	289	1.9	70.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.1	41.3
12	Pneumonitis due to solids and liquids (J69)	79	0.5	19.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	17.6
14	Parkinson's disease (G20-G21)	70	0.5	17.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.4	15.6
...	All other causes (Residual)	2,560	17.3	625.5
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,821	100.0	5,798.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,420	26.3	1,523.7
2	Malignant neoplasms (C00-C97)	3,324	19.8	1,145.9
3	Cerebrovascular diseases (I60-I69)	1,338	8.0	461.2
4	Diabetes mellitus (E10-E14)	840	5.0	289.6
5	Alzheimer's disease (G30)	708	4.2	244.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	633	3.8	218.2
7	Chronic lower respiratory diseases (J40-J47)	627	3.7	216.1
8	Septicemia (A40-A41)	401	2.4	138.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	385	2.3	132.7
10	Influenza and pneumonia (J09-J18)	358	2.1	123.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.0	60.0
12	Pneumonitis due to solids and liquids (J69)	118	0.7	40.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.5	29.3
14	Parkinson's disease (G20-G21)	84	0.5	29.0
15	Aortic aneurysm and dissection (I71)	59	0.4	20.3
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	59	0.4	20.3
...	All other causes (Residual)	3,208	19.1	1,105.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	17,064	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,676	27.4	---
2	Malignant neoplasms (C00-C97)	2,387	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,283	7.5	---
4	Alzheimer's disease (G30)	1,018	6.0	---
5	Diabetes mellitus (E10-E14)	761	4.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	578	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	570	3.3	---
8	Influenza and pneumonia (J09-J18)	429	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	2.5	---
10	Septicemia (A40-A41)	377	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	1.1	---
12	Pneumonitis due to solids and liquids (J69)	125	0.7	---
13	Parkinson's disease (G20-G21)	114	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	---
15	Aortic aneurysm and dissection (I71)	58	0.3	---
...	All other causes (Residual)	3,998	23.4	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	12,489	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,654	29.3	---
2	Malignant neoplasms (C00-C97)	1,257	10.1	---
3	Cerebrovascular diseases (I60-I69)	977	7.8	---
4	Alzheimer's disease (G30)	945	7.6	---
5	Diabetes mellitus (E10-E14)	427	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	363	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	352	2.8	---
8	Influenza and pneumonia (J09-J18)	331	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	323	2.6	---
10	Septicemia (A40-A41)	238	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	1.2	---
12	Pneumonitis due to solids and liquids (J69)	98	0.8	---
13	Atherosclerosis (I70)	58	0.5	---
14	Parkinson's disease (G20-G21)	57	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.4	---
...	All other causes (Residual)	3,211	25.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,798	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,764	30.4	---
2	Alzheimer's disease (G30)	499	8.6	---
3	Cerebrovascular diseases (I60-I69)	388	6.7	---
4	Malignant neoplasms (C00-C97)	375	6.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	3.4	---
6	Diabetes mellitus (E10-E14)	165	2.8	---
7	Influenza and pneumonia (J09-J18)	163	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	120	2.1	---
10	Septicemia (A40-A41)	109	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	1.2	---
12	Pneumonitis due to solids and liquids (J69)	42	0.7	---
13	Anemias (D50-D64)	24	0.4	---
14	Nutritional deficiencies (E40-E64)	19	0.3	---
15	Parkinson's disease (G20-G21)	18	0.3	---
...	All other causes (Residual)	1,698	29.3	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,147	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	30.6	---
2	Alzheimer's disease (G30)	186	8.7	---
3	Cerebrovascular diseases (I60-I69)	160	7.5	---
4	Malignant neoplasms (C00-C97)	83	3.9	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	3.6	---
6	Diabetes mellitus (E10-E14)	57	2.7	---
6	Influenza and pneumonia (J09-J18)	57	2.7	---
8	Septicemia (A40-A41)	52	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.0	---
10	Chronic lower respiratory diseases (J40-J47)	26	1.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.1	---
12	Pneumonitis due to solids and liquids (J69)	17	0.8	---
13	Atherosclerosis (I70)	15	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (Residual)	673	31.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013