

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	179,457	100.0	317.1
1	Malignant neoplasms (C00-C97)	37,804	21.1	66.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,401	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,806	7.7	24.4
4	Cerebrovascular diseases (I60-I69)	9,795	5.5	17.3
5	Diabetes mellitus (E10-E14)	8,278	4.6	14.6
6	Alzheimer's disease (G30)	6,444	3.6	11.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,018	3.4	10.6
8	Chronic lower respiratory diseases (J40-J47)	5,159	2.9	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,581	2.0	6.3
10	Influenza and pneumonia (J09-J18)	3,497	1.9	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,303	1.8	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,886	1.6	5.1
13	Septicemia (A40-A41)	2,692	1.5	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,454	1.4	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,384	1.3	4.2
...	All other causes (residual)	34,955	19.5	61.8
Hispanic, both sexes, 1-4 years				
...	All causes	815	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	31.7	6.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	14.1	2.8
3	Malignant neoplasms (C00-C97)	75	9.2	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	8.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.6	0.7
6	Influenza and pneumonia (J09-J18)	18	2.2	*
7	Septicemia (A40-A41)	17	2.1	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
11	Anemias (D50-D64)	4	0.5	*
11	Meningitis (G00,G03)	4	0.5	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	192	23.6	4.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	495	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	32.1	3.0
2	Malignant neoplasms (C00-C97)	102	20.6	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.5	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.4	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	Chronic lower respiratory diseases (J40-J47)	12	2.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.8	*
7	Cerebrovascular diseases (I60-I69)	9	1.8	*
9	Septicemia (A40-A41)	7	1.4	*
10	Influenza and pneumonia (J09-J18)	6	1.2	*
11	Meningitis (G00,G03)	4	0.8	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
13	Anemias (D50-D64)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	100	20.2	1.9
Hispanic, both sexes, 10-14 years				
...	All causes	568	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	26.2	3.0
2	Malignant neoplasms (C00-C97)	109	19.2	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	8.1	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	5.6	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	5.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	Septicemia (A40-A41)	9	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
9	Influenza and pneumonia (J09-J18)	8	1.4	*
11	Cerebrovascular diseases (I60-I69)	7	1.2	*
12	Anemias (D50-D64)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Aortic aneurysm and dissection (I71)	2	0.4	*
14	Peptic ulcer (K25-K28)	2	0.4	*
...	All other causes (residual)	130	22.9	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,868	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	742	39.7	15.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	316	16.9	6.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	299	16.0	6.3
4	Malignant neoplasms (C00-C97)	137	7.3	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.7	0.7
7	Cerebrovascular diseases (I60-I69)	18	1.0	*
8	Influenza and pneumonia (J09-J18)	14	0.7	*
9	Septicemia (A40-A41)	9	0.5	*
9	Diabetes mellitus (E10-E14)	9	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.4	*
12	Legal intervention (Y35,Y89.0)	7	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (residual)	201	10.8	4.3
Hispanic, both sexes, 20-24 years				
...	All causes	3,485	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,519	43.6	31.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	605	17.4	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	508	14.6	10.6
4	Malignant neoplasms (C00-C97)	195	5.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.5	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.1	0.8
7	Influenza and pneumonia (J09-J18)	22	0.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	22	0.6	0.5
9	Septicemia (A40-A41)	21	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.6	0.4
11	Cerebrovascular diseases (I60-I69)	14	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
13	Diabetes mellitus (E10-E14)	11	0.3	*
14	Legal intervention (Y35,Y89.0)	10	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (residual)	387	11.1	8.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,396	100.0	74.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,362	40.1	30.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	429	12.6	9.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	394	11.6	8.7
4	Malignant neoplasms (C00-C97)	287	8.5	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	4.7	3.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.4	1.0
7	Cerebrovascular diseases (I60-I69)	39	1.1	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.0	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.0	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.8	0.6
11	Diabetes mellitus (E10-E14)	27	0.8	0.6
12	Influenza and pneumonia (J09-J18)	26	0.8	0.6
13	Legal intervention (Y35,Y89.0)	21	0.6	0.5
14	Septicemia (A40-A41)	20	0.6	0.4
14	Chronic lower respiratory diseases (J40-J47)	20	0.6	0.4
...	All other causes (residual)	470	13.8	10.4
Hispanic, both sexes, 30-34 years				
...	All causes	3,786	100.0	85.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,291	34.1	29.0
2	Malignant neoplasms (C00-C97)	427	11.3	9.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	370	9.8	8.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	362	9.6	8.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	7.4	6.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	123	3.2	2.8
7	Cerebrovascular diseases (I60-I69)	82	2.2	1.8
8	Diabetes mellitus (E10-E14)	79	2.1	1.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.6	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.8	0.7
11	Influenza and pneumonia (J09-J18)	30	0.8	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.7	0.6
13	Septicemia (A40-A41)	26	0.7	0.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.7	0.6
15	Chronic lower respiratory diseases (J40-J47)	19	0.5	*
...	All other causes (residual)	551	14.6	12.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	4,202	100.0	99.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,133	27.0	26.7
2	Malignant neoplasms (C00-C97)	671	16.0	15.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	414	9.9	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	7.5	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	6.8	6.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	218	5.2	5.1
7	Cerebrovascular diseases (I60-I69)	118	2.8	2.8
8	Diabetes mellitus (E10-E14)	92	2.2	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	86	2.0	2.0
10	Influenza and pneumonia (J09-J18)	42	1.0	1.0
11	Septicemia (A40-A41)	40	1.0	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.9	0.9
13	Pregnancy, childbirth and the puerperium (O00-O99)	34	0.8	0.8
14	Aortic aneurysm and dissection (I71)	20	0.5	0.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.4	*
...	All other causes (residual)	675	16.1	15.9
Hispanic, both sexes, 40-44 years				
...	All causes	5,288	100.0	134.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,057	20.0	26.9
2	Malignant neoplasms (C00-C97)	1,047	19.8	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	668	12.6	17.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	376	7.1	9.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	5.1	6.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	4.1	5.6
7	Cerebrovascular diseases (I60-I69)	203	3.8	5.2
8	Diabetes mellitus (E10-E14)	182	3.4	4.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	105	2.0	2.7
10	Septicemia (A40-A41)	65	1.2	1.7
11	Influenza and pneumonia (J09-J18)	60	1.1	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.0	1.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.7	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	32	0.6	0.8
...	All other causes (residual)	876	16.6	22.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	7,159	100.0	204.8
1	Malignant neoplasms (C00-C97)	1,682	23.5	48.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,085	15.2	31.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,040	14.5	29.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	596	8.3	17.1
5	Diabetes mellitus (E10-E14)	282	3.9	8.1
6	Cerebrovascular diseases (I60-I69)	262	3.7	7.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	3.6	7.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	2.4	4.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	140	2.0	4.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	1.7	3.5
11	Viral hepatitis (B15-B19)	113	1.6	3.2
12	Septicemia (A40-A41)	107	1.5	3.1
13	Influenza and pneumonia (J09-J18)	79	1.1	2.3
14	Chronic lower respiratory diseases (J40-J47)	55	0.8	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	0.7	1.5
...	All other causes (residual)	1,120	15.6	32.0
Hispanic, both sexes, 50-54 years				
...	All causes	10,030	100.0	329.5
1	Malignant neoplasms (C00-C97)	2,604	26.0	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,759	17.5	57.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,050	10.5	34.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	881	8.8	28.9
5	Diabetes mellitus (E10-E14)	490	4.9	16.1
6	Cerebrovascular diseases (I60-I69)	403	4.0	13.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	2.5	8.2
8	Viral hepatitis (B15-B19)	167	1.7	5.5
9	Septicemia (A40-A41)	162	1.6	5.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	156	1.6	5.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	1.5	4.9
12	Influenza and pneumonia (J09-J18)	130	1.3	4.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	1.3	4.2
14	Chronic lower respiratory diseases (J40-J47)	112	1.1	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.8	2.6
...	All other causes (residual)	1,510	15.1	49.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	12,128	100.0	494.7
1	Malignant neoplasms (C00-C97)	3,602	29.7	146.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,287	18.9	93.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	913	7.5	37.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	845	7.0	34.5
5	Diabetes mellitus (E10-E14)	659	5.4	26.9
6	Cerebrovascular diseases (I60-I69)	513	4.2	20.9
7	Viral hepatitis (B15-B19)	262	2.2	10.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	1.8	8.9
9	Chronic lower respiratory diseases (J40-J47)	203	1.7	8.3
10	Septicemia (A40-A41)	188	1.6	7.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	1.5	7.3
12	Influenza and pneumonia (J09-J18)	166	1.4	6.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	1.0	5.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	1.0	5.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	0.7	3.4
...	All other causes (residual)	1,759	14.5	71.7
Hispanic, both sexes, 60-64 years				
...	All causes	13,859	100.0	753.9
1	Malignant neoplasms (C00-C97)	4,306	31.1	234.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,809	20.3	152.8
3	Diabetes mellitus (E10-E14)	957	6.9	52.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	854	6.2	46.5
5	Cerebrovascular diseases (I60-I69)	594	4.3	32.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	592	4.3	32.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	2.1	15.9
8	Chronic lower respiratory diseases (J40-J47)	289	2.1	15.7
9	Septicemia (A40-A41)	233	1.7	12.7
10	Viral hepatitis (B15-B19)	231	1.7	12.6
11	Influenza and pneumonia (J09-J18)	186	1.3	10.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.3	9.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	1.0	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	84	0.6	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	3.1
...	All other causes (residual)	2,059	14.9	112.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	14,920	100.0	1,099.0
1	Malignant neoplasms (C00-C97)	4,611	30.9	339.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,132	21.0	230.7
3	Diabetes mellitus (E10-E14)	1,054	7.1	77.6
4	Cerebrovascular diseases (I60-I69)	750	5.0	55.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	661	4.4	48.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	441	3.0	32.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.8	30.3
8	Chronic lower respiratory diseases (J40-J47)	381	2.6	28.1
9	Influenza and pneumonia (J09-J18)	272	1.8	20.0
10	Septicemia (A40-A41)	268	1.8	19.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	216	1.4	15.9
12	Viral hepatitis (B15-B19)	144	1.0	10.6
13	Alzheimer's disease (G30)	117	0.8	8.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	0.7	7.3
15	Parkinson's disease (G20-G21)	80	0.5	5.9
...	All other causes (residual)	2,282	15.3	168.1
Hispanic, both sexes, 70-74 years				
...	All causes	15,355	100.0	1,678.8
1	Malignant neoplasms (C00-C97)	4,522	29.4	494.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,385	22.0	370.1
3	Diabetes mellitus (E10-E14)	1,002	6.5	109.6
4	Cerebrovascular diseases (I60-I69)	811	5.3	88.7
5	Chronic lower respiratory diseases (J40-J47)	576	3.8	63.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	492	3.2	53.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	421	2.7	46.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	2.3	39.3
9	Septicemia (A40-A41)	301	2.0	32.9
10	Alzheimer's disease (G30)	284	1.8	31.1
10	Influenza and pneumonia (J09-J18)	284	1.8	31.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	1.4	23.4
13	Parkinson's disease (G20-G21)	162	1.1	17.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.7	11.0
15	Viral hepatitis (B15-B19)	81	0.5	8.9
...	All other causes (residual)	2,360	15.4	258.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	17,242	100.0	2,670.2
1	Malignant neoplasms (C00-C97)	4,459	25.9	690.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,950	22.9	611.7
3	Cerebrovascular diseases (I60-I69)	1,126	6.5	174.4
4	Diabetes mellitus (E10-E14)	1,032	6.0	159.8
5	Chronic lower respiratory diseases (J40-J47)	694	4.0	107.5
6	Alzheimer's disease (G30)	608	3.5	94.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	447	2.6	69.2
8	Influenza and pneumonia (J09-J18)	415	2.4	64.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	406	2.4	62.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	2.0	53.3
11	Septicemia (A40-A41)	318	1.8	49.2
12	Parkinson's disease (G20-G21)	265	1.5	41.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	263	1.5	40.7
14	Pneumonitis due to solids and liquids (J69)	113	0.7	17.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	16.4
...	All other causes (residual)	2,696	15.6	417.5
Hispanic, both sexes, 80-84 years				
...	All causes	19,847	100.0	4,556.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,885	24.6	1,121.5
2	Malignant neoplasms (C00-C97)	4,031	20.3	925.5
3	Cerebrovascular diseases (I60-I69)	1,522	7.7	349.4
4	Alzheimer's disease (G30)	1,247	6.3	286.3
5	Diabetes mellitus (E10-E14)	1,021	5.1	234.4
6	Chronic lower respiratory diseases (J40-J47)	932	4.7	214.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	530	2.7	121.7
8	Influenza and pneumonia (J09-J18)	524	2.6	120.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	2.1	94.6
10	Parkinson's disease (G20-G21)	368	1.9	84.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	1.7	78.5
12	Septicemia (A40-A41)	300	1.5	68.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	248	1.2	56.9
14	Pneumonitis due to solids and liquids (J69)	167	0.8	38.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.6	29.6
...	All other causes (residual)	3,189	16.1	732.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	20,040	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,465	27.3	---
2	Malignant neoplasms (C00-C97)	3,050	15.2	---
3	Alzheimer's disease (G30)	1,728	8.6	---
4	Cerebrovascular diseases (I60-I69)	1,571	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	913	4.6	---
6	Diabetes mellitus (E10-E14)	819	4.1	---
7	Influenza and pneumonia (J09-J18)	542	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.9	---
11	Parkinson's disease (G20-G21)	352	1.8	---
12	Septicemia (A40-A41)	315	1.6	---
13	Pneumonitis due to solids and liquids (J69)	188	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	142	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	112	0.6	---
...	All other causes (residual)	3,587	17.9	---
Hispanic, both sexes, 90-94 years				
...	All causes	13,807	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,899	28.2	---
2	Alzheimer's disease (G30)	1,590	11.5	---
3	Malignant neoplasms (C00-C97)	1,491	10.8	---
4	Cerebrovascular diseases (I60-I69)	1,179	8.5	---
5	Chronic lower respiratory diseases (J40-J47)	583	4.2	---
6	Diabetes mellitus (E10-E14)	415	3.0	---
7	Influenza and pneumonia (J09-J18)	413	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	277	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	1.9	---
11	Parkinson's disease (G20-G21)	208	1.5	---
12	Septicemia (A40-A41)	181	1.3	---
13	Pneumonitis due to solids and liquids (J69)	150	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	---
15	Nutritional deficiencies (E40-E64)	53	0.4	---
...	All other causes (residual)	2,747	19.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	5,074	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,523	30.0	---
2	Alzheimer's disease (G30)	622	12.3	---
3	Cerebrovascular diseases (I60-I69)	462	9.1	---
4	Malignant neoplasms (C00-C97)	334	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	236	4.7	---
6	Influenza and pneumonia (J09-J18)	173	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	2.7	---
8	Diabetes mellitus (E10-E14)	116	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	1.8	---
11	Pneumonitis due to solids and liquids (J69)	57	1.1	---
12	Septicemia (A40-A41)	52	1.0	---
13	Parkinson's disease (G20-G21)	42	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	---
15	Nutritional deficiencies (E40-E64)	24	0.5	---
...	All other causes (residual)	1,068	21.0	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,282	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	447	34.9	---
2	Alzheimer's disease (G30)	161	12.6	---
3	Cerebrovascular diseases (I60-I69)	85	6.6	---
4	Influenza and pneumonia (J09-J18)	57	4.4	---
5	Malignant neoplasms (C00-C97)	49	3.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	38	3.0	---
8	Diabetes mellitus (E10-E14)	25	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.3	---
11	Pneumonitis due to solids and liquids (J69)	15	1.2	---
12	Septicemia (A40-A41)	12	0.9	---
13	Nutritional deficiencies (E40-E64)	10	0.8	---
14	Atherosclerosis (I70)	8	0.6	---
15	Parkinson's disease (G20-G21)	7	0.5	---
...	All other causes (residual)	292	22.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	98,170	100.0	343.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,225	20.6	70.7
2	Malignant neoplasms (C00-C97)	19,847	20.2	69.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,067	10.3	35.2
4	Cerebrovascular diseases (I60-I69)	4,544	4.6	15.9
5	Diabetes mellitus (E10-E14)	4,426	4.5	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,109	4.2	14.4
7	Chronic lower respiratory diseases (J40-J47)	2,606	2.7	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,587	2.6	9.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,391	2.4	8.4
10	Alzheimer's disease (G30)	2,035	2.1	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,837	1.9	6.4
12	Influenza and pneumonia (J09-J18)	1,735	1.8	6.1
13	Septicemia (A40-A41)	1,370	1.4	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,368	1.4	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,139	1.2	4.0
...	All other causes (residual)	17,884	18.2	62.5
Hispanic, male, 1-4 years				
...	All causes	461	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	33.4	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	9.8	2.1
4	Malignant neoplasms (C00-C97)	40	8.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.6	1.0
6	Influenza and pneumonia (J09-J18)	12	2.6	*
7	Septicemia (A40-A41)	8	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Meningitis (G00,G03)	2	0.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Salmonella infections (A01-A02)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	99	21.5	4.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	301	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	33.9	3.8
2	Malignant neoplasms (C00-C97)	60	19.9	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	8.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.0	*
5	Chronic lower respiratory diseases (J40-J47)	7	2.3	*
6	Septicemia (A40-A41)	5	1.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.7	*
8	Cerebrovascular diseases (I60-I69)	4	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.0	*
9	Influenza and pneumonia (J09-J18)	3	1.0	*
11	Meningitis (G00,G03)	2	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
13	Viral hepatitis (B15-B19)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
...	All other causes (residual)	63	20.9	2.4
Hispanic, male, 10-14 years				
...	All causes	332	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	31.3	4.2
2	Malignant neoplasms (C00-C97)	61	18.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	7.2	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	6.3	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	5.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.7	*
7	Septicemia (A40-A41)	7	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.8	*
9	Cerebrovascular diseases (I60-I69)	4	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
11	Aortic aneurysm and dissection (I71)	2	0.6	*
11	Influenza and pneumonia (J09-J18)	2	0.6	*
11	Peptic ulcer (K25-K28)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
...	All other causes (residual)	64	19.3	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,341	100.0	55.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	549	40.9	22.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	20.2	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	216	16.1	8.9
4	Malignant neoplasms (C00-C97)	81	6.0	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.7	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.3	*
7	Cerebrovascular diseases (I60-I69)	12	0.9	*
8	Influenza and pneumonia (J09-J18)	9	0.7	*
9	Legal intervention (Y35,Y89.0)	7	0.5	*
10	Diabetes mellitus (E10-E14)	6	0.4	*
11	Septicemia (A40-A41)	5	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
12	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.1	*
...	All other causes (residual)	119	8.9	4.9
Hispanic, male, 20-24 years				
...	All causes	2,698	100.0	108.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,199	44.4	48.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	538	19.9	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	15.3	16.5
4	Malignant neoplasms (C00-C97)	127	4.7	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	2.3	2.4
6	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
8	Septicemia (A40-A41)	14	0.5	*
9	Influenza and pneumonia (J09-J18)	12	0.4	*
10	Legal intervention (Y35,Y89.0)	10	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.3	*
11	Diabetes mellitus (E10-E14)	9	0.3	*
13	Cerebrovascular diseases (I60-I69)	7	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.1	*
14	Pneumonitis due to solids and liquids (J69)	4	0.1	*
...	All other causes (residual)	254	9.4	10.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,592	100.0	108.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,117	43.1	46.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	374	14.4	15.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	325	12.5	13.6
4	Malignant neoplasms (C00-C97)	167	6.4	7.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	4.7	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	31	1.2	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.2	1.3
8	Cerebrovascular diseases (I60-I69)	26	1.0	1.1
9	Diabetes mellitus (E10-E14)	21	0.8	0.9
10	Legal intervention (Y35,Y89.0)	20	0.8	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
12	Influenza and pneumonia (J09-J18)	15	0.6	*
13	Septicemia (A40-A41)	13	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	277	10.7	11.6
Hispanic, male, 30-34 years				
...	All causes	2,676	100.0	115.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	37.8	43.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	300	11.2	12.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	10.7	12.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	7.3	8.4
5	Malignant neoplasms (C00-C97)	194	7.2	8.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	96	3.6	4.1
7	Cerebrovascular diseases (I60-I69)	58	2.2	2.5
8	Diabetes mellitus (E10-E14)	48	1.8	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.8	2.0
10	Influenza and pneumonia (J09-J18)	23	0.9	1.0
11	Legal intervention (Y35,Y89.0)	18	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.6	*
13	Septicemia (A40-A41)	14	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
...	All other causes (residual)	345	12.9	14.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,890	100.0	132.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	31.1	41.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	11.2	14.9
3	Malignant neoplasms (C00-C97)	288	10.0	13.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	8.6	11.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	8.2	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	158	5.5	7.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	73	2.5	3.3
8	Cerebrovascular diseases (I60-I69)	70	2.4	3.2
9	Diabetes mellitus (E10-E14)	56	1.9	2.6
10	Influenza and pneumonia (J09-J18)	28	1.0	1.3
11	Septicemia (A40-A41)	25	0.9	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.6	*
14	Aortic aneurysm and dissection (I71)	14	0.5	*
15	Legal intervention (Y35,Y89.0)	11	0.4	*
...	All other causes (residual)	426	14.7	19.5
Hispanic, male, 40-44 years				
...	All causes	3,495	100.0	175.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	23.8	41.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	13.9	24.3
3	Malignant neoplasms (C00-C97)	424	12.1	21.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	276	7.9	13.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	6.4	11.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	5.1	8.9
7	Diabetes mellitus (E10-E14)	133	3.8	6.7
7	Cerebrovascular diseases (I60-I69)	133	3.8	6.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	91	2.6	4.6
10	Septicemia (A40-A41)	41	1.2	2.1
11	Influenza and pneumonia (J09-J18)	35	1.0	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	1.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.7	1.3
14	Aortic aneurysm and dissection (I71)	23	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	1.2
...	All other causes (residual)	538	15.4	27.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,672	100.0	262.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	828	17.7	46.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	17.3	45.4
3	Malignant neoplasms (C00-C97)	778	16.7	43.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	459	9.8	25.8
5	Diabetes mellitus (E10-E14)	191	4.1	10.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	3.9	10.4
7	Cerebrovascular diseases (I60-I69)	147	3.1	8.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	2.6	6.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	118	2.5	6.6
10	Viral hepatitis (B15-B19)	90	1.9	5.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.6	4.1
12	Septicemia (A40-A41)	61	1.3	3.4
13	Influenza and pneumonia (J09-J18)	46	1.0	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	0.7	1.9
15	Chronic lower respiratory diseases (J40-J47)	28	0.6	1.6
...	All other causes (residual)	706	15.1	39.7
Hispanic, male, 50-54 years				
...	All causes	6,558	100.0	430.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,311	20.0	86.0
2	Malignant neoplasms (C00-C97)	1,310	20.0	85.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	12.1	52.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	689	10.5	45.2
5	Diabetes mellitus (E10-E14)	317	4.8	20.8
6	Cerebrovascular diseases (I60-I69)	232	3.5	15.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	2.9	12.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	128	2.0	8.4
9	Viral hepatitis (B15-B19)	126	1.9	8.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	1.6	7.0
11	Septicemia (A40-A41)	106	1.6	7.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.4	6.1
13	Influenza and pneumonia (J09-J18)	74	1.1	4.9
14	Chronic lower respiratory diseases (J40-J47)	57	0.9	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.7	3.1
...	All other causes (residual)	971	14.8	63.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	7,826	100.0	651.1
1	Malignant neoplasms (C00-C97)	1,964	25.1	163.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	21.5	139.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	665	8.5	55.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	640	8.2	53.2
5	Diabetes mellitus (E10-E14)	424	5.4	35.3
6	Cerebrovascular diseases (I60-I69)	330	4.2	27.5
7	Viral hepatitis (B15-B19)	184	2.4	15.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	1.9	12.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.7	11.2
10	Septicemia (A40-A41)	107	1.4	8.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	102	1.3	8.5
12	Chronic lower respiratory diseases (J40-J47)	97	1.2	8.1
13	Influenza and pneumonia (J09-J18)	92	1.2	7.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.1	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	0.9	6.0
...	All other causes (residual)	1,093	14.0	90.9
Hispanic, male, 60-64 years				
...	All causes	8,540	100.0	976.4
1	Malignant neoplasms (C00-C97)	2,372	27.8	271.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,983	23.2	226.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	607	7.1	69.4
4	Diabetes mellitus (E10-E14)	570	6.7	65.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	5.2	50.5
6	Cerebrovascular diseases (I60-I69)	350	4.1	40.0
7	Viral hepatitis (B15-B19)	168	2.0	19.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	1.9	18.4
9	Chronic lower respiratory diseases (J40-J47)	153	1.8	17.5
10	Septicemia (A40-A41)	120	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	1.3	12.7
12	Influenza and pneumonia (J09-J18)	104	1.2	11.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	1.2	11.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.8	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	0.5	4.5
...	All other causes (residual)	1,192	14.0	136.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	8,776	100.0	1,401.2
1	Malignant neoplasms (C00-C97)	2,599	29.6	415.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,010	22.9	320.9
3	Diabetes mellitus (E10-E14)	593	6.8	94.7
4	Cerebrovascular diseases (I60-I69)	449	5.1	71.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	420	4.8	67.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	3.5	48.5
7	Chronic lower respiratory diseases (J40-J47)	231	2.6	36.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	2.6	36.6
9	Influenza and pneumonia (J09-J18)	153	1.7	24.4
10	Septicemia (A40-A41)	151	1.7	24.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.4	20.3
12	Viral hepatitis (B15-B19)	95	1.1	15.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	0.9	13.1
14	Alzheimer's disease (G30)	54	0.6	8.6
15	Parkinson's disease (G20-G21)	50	0.6	8.0
...	All other causes (residual)	1,229	14.0	196.2
Hispanic, male, 70-74 years				
...	All causes	8,477	100.0	2,097.7
1	Malignant neoplasms (C00-C97)	2,429	28.7	601.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,066	24.4	511.2
3	Diabetes mellitus (E10-E14)	527	6.2	130.4
4	Cerebrovascular diseases (I60-I69)	441	5.2	109.1
5	Chronic lower respiratory diseases (J40-J47)	323	3.8	79.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	285	3.4	70.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.6	53.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	2.5	52.7
9	Septicemia (A40-A41)	174	2.1	43.1
10	Influenza and pneumonia (J09-J18)	153	1.8	37.9
11	Alzheimer's disease (G30)	125	1.5	30.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	1.4	29.0
13	Parkinson's disease (G20-G21)	95	1.1	23.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	12.4
15	Viral hepatitis (B15-B19)	48	0.6	11.9
...	All other causes (residual)	1,213	14.3	300.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	8,926	100.0	3,264.0
1	Malignant neoplasms (C00-C97)	2,438	27.3	891.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,189	24.5	800.5
3	Cerebrovascular diseases (I60-I69)	542	6.1	198.2
4	Diabetes mellitus (E10-E14)	494	5.5	180.6
5	Chronic lower respiratory diseases (J40-J47)	357	4.0	130.5
6	Alzheimer's disease (G30)	228	2.6	83.4
7	Influenza and pneumonia (J09-J18)	222	2.5	81.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	208	2.3	76.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	2.3	73.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	2.2	70.2
11	Parkinson's disease (G20-G21)	164	1.8	60.0
12	Septicemia (A40-A41)	150	1.7	54.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.6	53.0
14	Pneumonitis due to solids and liquids (J69)	57	0.6	20.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.6	19.0
...	All other causes (residual)	1,287	14.4	470.6
Hispanic, male, 80-84 years				
...	All causes	9,557	100.0	5,480.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,498	26.1	1,432.5
2	Malignant neoplasms (C00-C97)	2,114	22.1	1,212.3
3	Cerebrovascular diseases (I60-I69)	649	6.8	372.2
4	Chronic lower respiratory diseases (J40-J47)	473	4.9	271.3
5	Diabetes mellitus (E10-E14)	469	4.9	269.0
6	Alzheimer's disease (G30)	460	4.8	263.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	263	2.8	150.8
8	Influenza and pneumonia (J09-J18)	261	2.7	149.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	2.2	119.3
10	Parkinson's disease (G20-G21)	206	2.2	118.1
11	Septicemia (A40-A41)	139	1.5	79.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	138	1.4	79.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	113	1.2	64.8
14	Pneumonitis due to solids and liquids (J69)	90	0.9	51.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.7	39.6
...	All other causes (residual)	1,407	14.7	806.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	8,574	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,367	27.6	---
2	Malignant neoplasms (C00-C97)	1,539	17.9	---
3	Cerebrovascular diseases (I60-I69)	585	6.8	---
4	Alzheimer's disease (G30)	559	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	426	5.0	---
6	Diabetes mellitus (E10-E14)	344	4.0	---
7	Influenza and pneumonia (J09-J18)	248	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	2.7	---
9	Parkinson's disease (G20-G21)	202	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.3	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	1.5	---
12	Septicemia (A40-A41)	123	1.4	---
13	Pneumonitis due to solids and liquids (J69)	103	1.2	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	---
...	All other causes (residual)	1,405	16.4	---
Hispanic, male, 90-94 years				
...	All causes	5,078	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,443	28.4	---
2	Malignant neoplasms (C00-C97)	704	13.9	---
3	Alzheimer's disease (G30)	429	8.4	---
4	Cerebrovascular diseases (I60-I69)	352	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	272	5.4	---
6	Diabetes mellitus (E10-E14)	178	3.5	---
7	Influenza and pneumonia (J09-J18)	151	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.9	---
11	Parkinson's disease (G20-G21)	89	1.8	---
12	Septicemia (A40-A41)	69	1.4	---
13	Pneumonitis due to solids and liquids (J69)	65	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	---
15	Nutritional deficiencies (E40-E64)	19	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.4	---
...	All other causes (residual)	934	18.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,474	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	461	31.3	---
2	Malignant neoplasms (C00-C97)	137	9.3	---
3	Alzheimer's disease (G30)	118	8.0	---
4	Cerebrovascular diseases (I60-I69)	116	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	93	6.3	---
6	Influenza and pneumonia (J09-J18)	58	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	2.7	---
8	Diabetes mellitus (E10-E14)	35	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	2.4	---
11	Pneumonitis due to solids and liquids (J69)	27	1.8	---
12	Septicemia (A40-A41)	13	0.9	---
12	Parkinson's disease (G20-G21)	13	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	---
15	Atherosclerosis (I70)	7	0.5	---
...	All other causes (residual)	275	18.7	---
Hispanic, male, 100 years and over				
...	All causes	302	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	37.7	---
2	Alzheimer's disease (G30)	26	8.6	---
3	Cerebrovascular diseases (I60-I69)	22	7.3	---
4	Influenza and pneumonia (J09-J18)	14	4.6	---
5	Malignant neoplasms (C00-C97)	13	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	13	4.3	---
7	Diabetes mellitus (E10-E14)	9	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.3	---
11	Septicemia (A40-A41)	5	1.7	---
12	Pneumonitis due to solids and liquids (J69)	4	1.3	---
13	Atherosclerosis (I70)	3	1.0	---
14	Parkinson's disease (G20-G21)	2	0.7	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	---
15	Hyperplasia of prostate (N40)	1	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	---
...	All other causes (residual)	49	16.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	81,287	100.0	290.4
1	Malignant neoplasms (C00-C97)	17,957	22.1	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,176	19.9	57.8
3	Cerebrovascular diseases (I60-I69)	5,251	6.5	18.8
4	Alzheimer's disease (G30)	4,409	5.4	15.8
5	Diabetes mellitus (E10-E14)	3,852	4.7	13.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,739	4.6	13.4
7	Chronic lower respiratory diseases (J40-J47)	2,553	3.1	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,909	2.3	6.8
9	Influenza and pneumonia (J09-J18)	1,762	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,744	2.1	6.2
11	Septicemia (A40-A41)	1,322	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,245	1.5	4.4
13	Certain conditions originating in the perinatal period (P00-P96)	1,086	1.3	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	837	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	0.9	2.6
...	All other causes (residual)	16,729	20.6	59.8
Hispanic, female, 1-4 years				
...	All causes	354	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	29.4	5.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.1	2.8
3	Malignant neoplasms (C00-C97)	35	9.9	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.8	1.2
5	Septicemia (A40-A41)	9	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.3	*
7	Influenza and pneumonia (J09-J18)	6	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.4	*
9	Cerebrovascular diseases (I60-I69)	4	1.1	*
10	Anemias (D50-D64)	3	0.8	*
11	Meningitis (G00,G03)	2	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	86	24.3	4.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	194	100.0	7.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	29.4	2.2
2	Malignant neoplasms (C00-C97)	42	21.6	1.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	6.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	6.2	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	3.1	*
6	Cerebrovascular diseases (I60-I69)	5	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	5	2.6	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.1	*
9	Influenza and pneumonia (J09-J18)	3	1.5	*
9	Pneumonitis due to solids and liquids (J69)	3	1.5	*
11	Septicemia (A40-A41)	2	1.0	*
11	Diabetes mellitus (E10-E14)	2	1.0	*
11	Meningitis (G00,G03)	2	1.0	*
14	Meningococcal infection (A39)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (residual)	34	17.5	1.3
Hispanic, female, 10-14 years				
...	All causes	236	100.0	9.7
1	Malignant neoplasms (C00-C97)	48	20.3	2.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	19.1	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	10.6	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.0	*
7	Influenza and pneumonia (J09-J18)	6	2.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.1	*
10	Cerebrovascular diseases (I60-I69)	3	1.3	*
11	Septicemia (A40-A41)	2	0.8	*
11	Anemias (D50-D64)	2	0.8	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	60	25.4	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	527	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	36.6	8.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	15.7	3.6
3	Malignant neoplasms (C00-C97)	56	10.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.4	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.7	*
7	Cerebrovascular diseases (I60-I69)	6	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
9	Influenza and pneumonia (J09-J18)	5	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
12	Septicemia (A40-A41)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
14	Meningitis (G00,G03)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	75	14.2	3.3
Hispanic, female, 20-24 years				
...	All causes	787	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	40.7	13.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	12.1	4.1
3	Malignant neoplasms (C00-C97)	68	8.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	8.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.2	1.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	21	2.7	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.4	*
8	Influenza and pneumonia (J09-J18)	10	1.3	*
9	Septicemia (A40-A41)	7	0.9	*
9	Cerebrovascular diseases (I60-I69)	7	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	127	16.1	5.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	804	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	30.5	11.4
2	Malignant neoplasms (C00-C97)	120	14.9	5.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	8.6	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.8	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.5	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	28	3.5	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	2.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.9	*
9	Cerebrovascular diseases (I60-I69)	13	1.6	*
10	Influenza and pneumonia (J09-J18)	11	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.0	*
13	Septicemia (A40-A41)	7	0.9	*
14	Diabetes mellitus (E10-E14)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
...	All other causes (residual)	162	20.1	7.5
Hispanic, female, 30-34 years				
...	All causes	1,110	100.0	52.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	25.1	13.1
2	Malignant neoplasms (C00-C97)	233	21.0	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	7.7	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	7.5	3.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	2.9
6	Diabetes mellitus (E10-E14)	31	2.8	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.8	1.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.4	1.3
9	Cerebrovascular diseases (I60-I69)	24	2.2	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.3	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.3	*
12	Septicemia (A40-A41)	12	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Influenza and pneumonia (J09-J18)	7	0.6	*
...	All other causes (residual)	188	16.9	8.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,312	100.0	63.8
1	Malignant neoplasms (C00-C97)	383	29.2	18.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	17.9	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	6.8	4.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	5.1	3.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	60	4.6	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	3.9	2.5
7	Cerebrovascular diseases (I60-I69)	48	3.7	2.3
8	Diabetes mellitus (E10-E14)	36	2.7	1.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	34	2.6	1.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.6	1.0
11	Septicemia (A40-A41)	15	1.1	*
12	Influenza and pneumonia (J09-J18)	14	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.0	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (residual)	231	17.6	11.2
Hispanic, female, 40-44 years				
...	All causes	1,793	100.0	92.4
1	Malignant neoplasms (C00-C97)	623	34.7	32.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	12.6	11.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	10.2	9.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	100	5.6	5.2
5	Cerebrovascular diseases (I60-I69)	70	3.9	3.6
6	Diabetes mellitus (E10-E14)	49	2.7	2.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	2.6	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	2.3	2.2
9	Influenza and pneumonia (J09-J18)	25	1.4	1.3
10	Septicemia (A40-A41)	24	1.3	1.2
11	Chronic lower respiratory diseases (J40-J47)	21	1.2	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.8	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.7	*
...	All other causes (residual)	322	18.0	16.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,487	100.0	144.7
1	Malignant neoplasms (C00-C97)	904	36.3	52.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	257	10.3	15.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	9.4	13.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	137	5.5	8.0
5	Cerebrovascular diseases (I60-I69)	115	4.6	6.7
6	Diabetes mellitus (E10-E14)	91	3.7	5.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	2.9	4.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.0	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.9	2.7
10	Septicemia (A40-A41)	46	1.8	2.7
11	Influenza and pneumonia (J09-J18)	33	1.3	1.9
12	Chronic lower respiratory diseases (J40-J47)	27	1.1	1.6
13	Viral hepatitis (B15-B19)	23	0.9	1.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.9	1.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.8	*
...	All other causes (residual)	412	16.6	24.0
Hispanic, female, 50-54 years				
...	All causes	3,472	100.0	228.4
1	Malignant neoplasms (C00-C97)	1,294	37.3	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	448	12.9	29.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	7.3	16.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	192	5.5	12.6
5	Diabetes mellitus (E10-E14)	173	5.0	11.4
6	Cerebrovascular diseases (I60-I69)	171	4.9	11.2
7	Septicemia (A40-A41)	56	1.6	3.7
7	Influenza and pneumonia (J09-J18)	56	1.6	3.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	3.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	1.6	3.7
11	Chronic lower respiratory diseases (J40-J47)	55	1.6	3.6
12	Viral hepatitis (B15-B19)	41	1.2	2.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.9	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.8	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.6	1.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.6	1.4
...	All other causes (residual)	518	14.9	34.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	4,302	100.0	344.2
1	Malignant neoplasms (C00-C97)	1,638	38.1	131.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	607	14.1	48.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	248	5.8	19.8
4	Diabetes mellitus (E10-E14)	235	5.5	18.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	4.8	16.4
6	Cerebrovascular diseases (I60-I69)	183	4.3	14.6
7	Chronic lower respiratory diseases (J40-J47)	106	2.5	8.5
8	Septicemia (A40-A41)	81	1.9	6.5
9	Viral hepatitis (B15-B19)	78	1.8	6.2
10	Influenza and pneumonia (J09-J18)	74	1.7	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	1.5	5.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	1.0	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	0.9	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	*
...	All other causes (residual)	658	15.3	52.6
Hispanic, female, 60-64 years				
...	All causes	5,319	100.0	551.9
1	Malignant neoplasms (C00-C97)	1,934	36.4	200.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	826	15.5	85.7
3	Diabetes mellitus (E10-E14)	387	7.3	40.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	247	4.6	25.6
5	Cerebrovascular diseases (I60-I69)	244	4.6	25.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	2.8	15.6
7	Chronic lower respiratory diseases (J40-J47)	136	2.6	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.5	13.7
9	Septicemia (A40-A41)	113	2.1	11.7
10	Influenza and pneumonia (J09-J18)	82	1.5	8.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.2	6.8
12	Viral hepatitis (B15-B19)	63	1.2	6.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.7	3.6
14	Alzheimer's disease (G30)	31	0.6	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	3.0
...	All other causes (residual)	844	15.9	87.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	6,144	100.0	840.2
1	Malignant neoplasms (C00-C97)	2,012	32.7	275.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,122	18.3	153.4
3	Diabetes mellitus (E10-E14)	461	7.5	63.0
4	Cerebrovascular diseases (I60-I69)	301	4.9	41.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	241	3.9	33.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	3.0	25.0
7	Chronic lower respiratory diseases (J40-J47)	150	2.4	20.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	2.2	18.7
9	Influenza and pneumonia (J09-J18)	119	1.9	16.3
10	Septicemia (A40-A41)	117	1.9	16.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.4	12.2
12	Alzheimer's disease (G30)	63	1.0	8.6
13	Viral hepatitis (B15-B19)	49	0.8	6.7
14	Parkinson's disease (G20-G21)	30	0.5	4.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	26	0.4	3.6
...	All other causes (residual)	1,044	17.0	142.8
Hispanic, female, 70-74 years				
...	All causes	6,878	100.0	1,347.2
1	Malignant neoplasms (C00-C97)	2,093	30.4	410.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,319	19.2	258.4
3	Diabetes mellitus (E10-E14)	475	6.9	93.0
4	Cerebrovascular diseases (I60-I69)	370	5.4	72.5
5	Chronic lower respiratory diseases (J40-J47)	253	3.7	49.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	207	3.0	40.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	3.0	39.8
8	Alzheimer's disease (G30)	159	2.3	31.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	2.1	28.6
10	Influenza and pneumonia (J09-J18)	131	1.9	25.7
11	Septicemia (A40-A41)	127	1.8	24.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.4	19.0
13	Parkinson's disease (G20-G21)	67	1.0	13.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.7	10.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	34	0.5	6.7
...	All other causes (residual)	1,146	16.7	224.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	8,316	100.0	2,234.0
1	Malignant neoplasms (C00-C97)	2,021	24.3	542.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,761	21.2	473.1
3	Cerebrovascular diseases (I60-I69)	584	7.0	156.9
4	Diabetes mellitus (E10-E14)	538	6.5	144.5
5	Alzheimer's disease (G30)	380	4.6	102.1
6	Chronic lower respiratory diseases (J40-J47)	337	4.1	90.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	3.0	66.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	198	2.4	53.2
9	Influenza and pneumonia (J09-J18)	193	2.3	51.8
10	Septicemia (A40-A41)	168	2.0	45.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	1.8	40.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	1.4	31.7
13	Parkinson's disease (G20-G21)	101	1.2	27.1
14	Pneumonitis due to solids and liquids (J69)	56	0.7	15.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.6	14.5
...	All other causes (residual)	1,409	16.9	378.5
Hispanic, female, 80-84 years				
...	All causes	10,290	100.0	3,939.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,387	23.2	913.9
2	Malignant neoplasms (C00-C97)	1,917	18.6	734.0
3	Cerebrovascular diseases (I60-I69)	873	8.5	334.2
4	Alzheimer's disease (G30)	787	7.6	301.3
5	Diabetes mellitus (E10-E14)	552	5.4	211.3
6	Chronic lower respiratory diseases (J40-J47)	459	4.5	175.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	2.6	102.2
8	Influenza and pneumonia (J09-J18)	263	2.6	100.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	204	2.0	78.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	2.0	78.1
11	Parkinson's disease (G20-G21)	162	1.6	62.0
12	Septicemia (A40-A41)	161	1.6	61.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.3	51.7
14	Pneumonitis due to solids and liquids (J69)	77	0.7	29.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.6	23.0
...	All other causes (residual)	1,782	17.3	682.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	11,466	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,098	27.0	---
2	Malignant neoplasms (C00-C97)	1,511	13.2	---
3	Alzheimer's disease (G30)	1,169	10.2	---
4	Cerebrovascular diseases (I60-I69)	986	8.6	---
5	Chronic lower respiratory diseases (J40-J47)	487	4.2	---
6	Diabetes mellitus (E10-E14)	475	4.1	---
7	Influenza and pneumonia (J09-J18)	294	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.8	---
11	Septicemia (A40-A41)	192	1.7	---
12	Parkinson's disease (G20-G21)	150	1.3	---
13	Pneumonitis due to solids and liquids (J69)	85	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.6	---
...	All other causes (residual)	2,182	19.0	---
Hispanic, female, 90-94 years				
...	All causes	8,729	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,456	28.1	---
2	Alzheimer's disease (G30)	1,161	13.3	---
3	Cerebrovascular diseases (I60-I69)	827	9.5	---
4	Malignant neoplasms (C00-C97)	787	9.0	---
5	Chronic lower respiratory diseases (J40-J47)	311	3.6	---
6	Influenza and pneumonia (J09-J18)	262	3.0	---
7	Diabetes mellitus (E10-E14)	237	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	147	1.7	---
11	Parkinson's disease (G20-G21)	119	1.4	---
12	Septicemia (A40-A41)	112	1.3	---
13	Pneumonitis due to solids and liquids (J69)	85	1.0	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	35	0.4	---
15	Nutritional deficiencies (E40-E64)	34	0.4	---
...	All other causes (residual)	1,790	20.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	3,600	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,062	29.5	---
2	Alzheimer's disease (G30)	504	14.0	---
3	Cerebrovascular diseases (I60-I69)	346	9.6	---
4	Malignant neoplasms (C00-C97)	197	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	143	4.0	---
6	Influenza and pneumonia (J09-J18)	115	3.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	2.7	---
8	Diabetes mellitus (E10-E14)	81	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	---
11	Septicemia (A40-A41)	39	1.1	---
12	Pneumonitis due to solids and liquids (J69)	30	0.8	---
13	Parkinson's disease (G20-G21)	29	0.8	---
14	Nutritional deficiencies (E40-E64)	18	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	---
...	All other causes (residual)	792	22.0	---
Hispanic, female, 100 years and over				
...	All causes	980	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	34.0	---
2	Alzheimer's disease (G30)	135	13.8	---
3	Cerebrovascular diseases (I60-I69)	63	6.4	---
4	Influenza and pneumonia (J09-J18)	43	4.4	---
5	Malignant neoplasms (C00-C97)	36	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	25	2.6	---
8	Diabetes mellitus (E10-E14)	16	1.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.3	---
10	Pneumonitis due to solids and liquids (J69)	11	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.0	---
12	Nutritional deficiencies (E40-E64)	9	0.9	---
13	Septicemia (A40-A41)	7	0.7	---
14	Anemias (D50-D64)	6	0.6	---
15	Parkinson's disease (G20-G21)	5	0.5	---
15	Atherosclerosis (I70)	5	0.5	---
...	All other causes (residual)	232	23.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,123,631	100.0	1,055.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,172	23.7	250.0
2	Malignant neoplasms (C00-C97)	467,208	22.0	232.2
3	Chronic lower respiratory diseases (J40-J47)	136,228	6.4	67.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111,827	5.3	55.6
5	Cerebrovascular diseases (I60-I69)	106,830	5.0	53.1
6	Alzheimer's disease (G30)	93,325	4.4	46.4
7	Diabetes mellitus (E10-E14)	53,590	2.5	26.6
8	Influenza and pneumonia (J09-J18)	45,242	2.1	22.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,465	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,481	1.7	17.6
11	Septicemia (A40-A41)	30,240	1.4	15.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,343	1.4	14.6
13	Parkinson's disease (G20-G21)	24,339	1.1	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,628	1.1	11.2
15	Pneumonitis due to solids and liquids (J69)	16,430	0.8	8.2
...	All other causes (residual)	411,283	19.4	204.4
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,895	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	631	33.3	7.6
2	Malignant neoplasms (C00-C97)	206	10.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	10.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	7.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.9	0.7
6	Influenza and pneumonia (J09-J18)	37	2.0	0.4
7	Septicemia (A40-A41)	20	1.1	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Cerebrovascular diseases (I60-I69)	17	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	10	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Anemias (D50-D64)	7	0.4	*
...	All other causes (residual)	511	27.0	6.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,192	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	31.4	3.5
2	Malignant neoplasms (C00-C97)	241	20.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	5.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.6	0.3
6	Influenza and pneumonia (J09-J18)	25	2.1	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.8	0.2
8	Cerebrovascular diseases (I60-I69)	20	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	17	1.4	*
10	Septicemia (A40-A41)	12	1.0	*
11	Anemias (D50-D64)	6	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Meningitis (G00,G03)	2	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
14	Peptic ulcer (K25-K28)	2	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	259	21.7	2.4
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,643	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	424	25.8	3.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	289	17.6	2.6
3	Malignant neoplasms (C00-C97)	235	14.3	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	5.6	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	4.3	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	2.4	0.3
7	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.2
8	Influenza and pneumonia (J09-J18)	22	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	16	1.0	*
12	Diabetes mellitus (E10-E14)	10	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.2	*
...	All other causes (residual)	365	22.2	3.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,456	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,449	44.9	20.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,411	25.9	12.0
3	Malignant neoplasms (C00-C97)	307	5.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	4.3	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	2.0	0.9
7	Influenza and pneumonia (J09-J18)	44	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	29	0.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	0.2
10	Diabetes mellitus (E10-E14)	22	0.4	0.2
11	Septicemia (A40-A41)	18	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	18	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	Legal intervention (Y35,Y89.0)	9	0.2	*
...	All other causes (residual)	602	11.0	5.1
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,080	100.0	87.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,630	50.8	44.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,266	20.5	17.8
3	Malignant neoplasms (C00-C97)	474	4.3	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	446	4.0	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	3.1	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	1.0	0.9
7	Diabetes mellitus (E10-E14)	82	0.7	0.6
8	Influenza and pneumonia (J09-J18)	63	0.6	0.5
9	Cerebrovascular diseases (I60-I69)	54	0.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	51	0.5	0.4
12	Septicemia (A40-A41)	43	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	25	0.2	0.2
15	Legal intervention (Y35,Y89.0)	23	0.2	0.2
...	All other causes (residual)	1,382	12.5	10.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	14,349	100.0	111.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,014	48.9	54.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,495	17.4	19.4
3	Malignant neoplasms (C00-C97)	757	5.3	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	689	4.8	5.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	519	3.6	4.0
6	Diabetes mellitus (E10-E14)	152	1.1	1.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	131	0.9	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	0.9	1.0
9	Influenza and pneumonia (J09-J18)	103	0.7	0.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	103	0.7	0.8
11	Cerebrovascular diseases (I60-I69)	84	0.6	0.7
12	Septicemia (A40-A41)	76	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	47	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.2	0.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.2	0.3
...	All other causes (residual)	1,988	13.9	15.5
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	17,143	100.0	136.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,129	41.6	56.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,682	15.6	21.4
3	Malignant neoplasms (C00-C97)	1,374	8.0	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,239	7.2	9.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	516	3.0	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	392	2.3	3.1
7	Diabetes mellitus (E10-E14)	256	1.5	2.0
8	Cerebrovascular diseases (I60-I69)	172	1.0	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	0.9	1.2
10	Influenza and pneumonia (J09-J18)	123	0.7	1.0
11	Septicemia (A40-A41)	122	0.7	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	99	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	80	0.5	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	0.5
...	All other causes (residual)	2,680	15.6	21.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	19,392	100.0	165.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,632	34.2	56.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,654	13.7	22.6
3	Malignant neoplasms (C00-C97)	2,196	11.3	18.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	10.5	17.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	638	3.3	5.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	429	2.2	3.6
7	Diabetes mellitus (E10-E14)	378	1.9	3.2
8	Cerebrovascular diseases (I60-I69)	266	1.4	2.3
9	Influenza and pneumonia (J09-J18)	173	0.9	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.8	1.4
11	Septicemia (A40-A41)	159	0.8	1.4
12	Chronic lower respiratory diseases (J40-J47)	129	0.7	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	0.6	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	81	0.4	0.7
...	All other causes (residual)	3,265	16.8	27.8
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,245	100.0	218.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,259	23.8	52.2
2	Malignant neoplasms (C00-C97)	4,340	16.5	36.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,950	15.1	32.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,969	11.3	24.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,194	4.5	10.0
6	Diabetes mellitus (E10-E14)	690	2.6	5.8
7	Cerebrovascular diseases (I60-I69)	530	2.0	4.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	1.7	3.6
9	Chronic lower respiratory diseases (J40-J47)	315	1.2	2.6
10	Septicemia (A40-A41)	304	1.2	2.5
11	Influenza and pneumonia (J09-J18)	251	1.0	2.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	177	0.7	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	0.7	1.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	140	0.5	1.2
15	Viral hepatitis (B15-B19)	117	0.4	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	0.4	1.0
...	All other causes (residual)	4,282	16.3	35.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	42,315	100.0	320.2
1	Malignant neoplasms (C00-C97)	9,062	21.4	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,678	18.1	58.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,184	17.0	54.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,388	8.0	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,181	5.2	16.5
6	Diabetes mellitus (E10-E14)	1,301	3.1	9.8
7	Cerebrovascular diseases (I60-I69)	960	2.3	7.3
8	Chronic lower respiratory diseases (J40-J47)	827	2.0	6.3
9	Septicemia (A40-A41)	547	1.3	4.1
10	Influenza and pneumonia (J09-J18)	494	1.2	3.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	444	1.0	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	0.8	2.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	280	0.7	2.1
14	Viral hepatitis (B15-B19)	270	0.6	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.6	1.8
...	All other causes (residual)	7,110	16.8	53.8
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	76,263	100.0	504.2
1	Malignant neoplasms (C00-C97)	20,504	26.9	135.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,969	19.6	99.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,693	11.4	57.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,158	5.5	27.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,033	5.3	26.7
6	Chronic lower respiratory diseases (J40-J47)	2,636	3.5	17.4
7	Diabetes mellitus (E10-E14)	2,334	3.1	15.4
8	Cerebrovascular diseases (I60-I69)	1,915	2.5	12.7
9	Septicemia (A40-A41)	1,003	1.3	6.6
10	Influenza and pneumonia (J09-J18)	835	1.1	5.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	724	0.9	4.8
12	Viral hepatitis (B15-B19)	712	0.9	4.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	0.7	3.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	480	0.6	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	423	0.6	2.8
...	All other causes (residual)	12,294	16.1	81.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	113,442	100.0	733.0
1	Malignant neoplasms (C00-C97)	36,430	32.1	235.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,105	20.4	149.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,239	7.3	53.2
4	Chronic lower respiratory diseases (J40-J47)	5,597	4.9	36.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,961	4.4	32.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,003	3.5	25.9
7	Diabetes mellitus (E10-E14)	3,755	3.3	24.3
8	Cerebrovascular diseases (I60-I69)	3,040	2.7	19.6
9	Septicemia (A40-A41)	1,610	1.4	10.4
10	Influenza and pneumonia (J09-J18)	1,323	1.2	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,174	1.0	7.6
12	Viral hepatitis (B15-B19)	1,120	1.0	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	865	0.8	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	504	0.4	3.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	449	0.4	2.9
...	All other causes (residual)	17,267	15.2	111.6
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	144,070	100.0	1,025.1
1	Malignant neoplasms (C00-C97)	49,264	34.2	350.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,801	21.4	219.2
3	Chronic lower respiratory diseases (J40-J47)	9,060	6.3	64.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,336	4.4	45.1
5	Diabetes mellitus (E10-E14)	5,072	3.5	36.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,767	3.3	33.9
7	Cerebrovascular diseases (I60-I69)	4,308	3.0	30.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,926	2.0	20.8
9	Septicemia (A40-A41)	2,262	1.6	16.1
10	Influenza and pneumonia (J09-J18)	1,879	1.3	13.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,871	1.3	13.3
12	Viral hepatitis (B15-B19)	1,237	0.9	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,210	0.8	8.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	733	0.5	5.2
15	Pneumonitis due to solids and liquids (J69)	647	0.4	4.6
...	All other causes (residual)	21,697	15.1	154.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	179,608	100.0	1,464.5
1	Malignant neoplasms (C00-C97)	63,108	35.1	514.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,997	21.2	309.8
3	Chronic lower respiratory diseases (J40-J47)	14,033	7.8	114.4
4	Diabetes mellitus (E10-E14)	6,470	3.6	52.8
5	Cerebrovascular diseases (I60-I69)	6,040	3.4	49.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,425	3.0	44.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,686	2.1	30.1
8	Septicemia (A40-A41)	2,959	1.6	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,790	1.6	22.7
10	Influenza and pneumonia (J09-J18)	2,631	1.5	21.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,209	1.2	18.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,533	0.9	12.5
13	Alzheimer's disease (G30)	1,384	0.8	11.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,061	0.6	8.7
15	Parkinson's disease (G20-G21)	1,004	0.6	8.2
...	All other causes (residual)	27,278	15.2	222.4
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	204,486	100.0	2,286.1
1	Malignant neoplasms (C00-C97)	66,699	32.6	745.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,294	21.2	484.0
3	Chronic lower respiratory diseases (J40-J47)	19,219	9.4	214.9
4	Cerebrovascular diseases (I60-I69)	8,526	4.2	95.3
5	Diabetes mellitus (E10-E14)	6,643	3.2	74.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,886	2.4	54.6
7	Alzheimer's disease (G30)	3,621	1.8	40.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,538	1.7	39.6
9	Influenza and pneumonia (J09-J18)	3,481	1.7	38.9
10	Septicemia (A40-A41)	3,352	1.6	37.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,641	1.3	29.5
12	Parkinson's disease (G20-G21)	2,199	1.1	24.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,648	0.8	18.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,618	0.8	18.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,499	0.7	16.8
...	All other causes (residual)	31,622	15.5	353.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	232,080	100.0	3,655.3
1	Malignant neoplasms (C00-C97)	64,236	27.7	1,011.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,751	21.9	799.3
3	Chronic lower respiratory diseases (J40-J47)	21,692	9.3	341.7
4	Cerebrovascular diseases (I60-I69)	11,827	5.1	186.3
5	Alzheimer's disease (G30)	8,134	3.5	128.1
6	Diabetes mellitus (E10-E14)	6,590	2.8	103.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,799	2.5	91.3
8	Influenza and pneumonia (J09-J18)	4,527	2.0	71.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,382	1.9	69.0
10	Parkinson's disease (G20-G21)	4,018	1.7	63.3
11	Septicemia (A40-A41)	3,812	1.6	60.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,164	0.9	34.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,979	0.9	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,817	0.8	28.6
15	Pneumonitis due to solids and liquids (J69)	1,722	0.7	27.1
...	All other causes (residual)	38,630	16.6	608.4
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	285,606	100.0	6,176.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,159	24.2	1,495.7
2	Malignant neoplasms (C00-C97)	61,031	21.4	1,319.9
3	Chronic lower respiratory diseases (J40-J47)	22,781	8.0	492.7
4	Cerebrovascular diseases (I60-I69)	17,335	6.1	374.9
5	Alzheimer's disease (G30)	16,388	5.7	354.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,434	2.6	160.8
7	Diabetes mellitus (E10-E14)	7,174	2.5	155.2
8	Influenza and pneumonia (J09-J18)	6,638	2.3	143.6
9	Parkinson's disease (G20-G21)	5,966	2.1	129.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,787	2.0	125.2
11	Septicemia (A40-A41)	4,264	1.5	92.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,091	1.1	66.8
13	Pneumonitis due to solids and liquids (J69)	2,519	0.9	54.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,270	0.8	49.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,461	0.5	31.6
...	All other causes (residual)	52,308	18.3	1,131.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	331,643	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,934	27.1	---
2	Malignant neoplasms (C00-C97)	50,636	15.3	---
3	Alzheimer's disease (G30)	25,114	7.6	---
4	Cerebrovascular diseases (I60-I69)	22,643	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	21,486	6.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,371	2.8	---
7	Influenza and pneumonia (J09-J18)	9,040	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,768	2.0	---
9	Diabetes mellitus (E10-E14)	6,767	2.0	---
10	Parkinson's disease (G20-G21)	6,181	1.9	---
11	Septicemia (A40-A41)	4,860	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,239	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,579	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,396	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,256	0.4	---
...	All other causes (residual)	67,373	20.3	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	271,961	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,632	30.4	---
2	Malignant neoplasms (C00-C97)	27,914	10.3	---
3	Alzheimer's disease (G30)	24,731	9.1	---
4	Cerebrovascular diseases (I60-I69)	19,330	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	13,345	4.9	---
6	Influenza and pneumonia (J09-J18)	8,529	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,631	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,343	2.0	---
9	Diabetes mellitus (E10-E14)	4,380	1.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,267	1.6	---
11	Parkinson's disease (G20-G21)	3,519	1.3	---
12	Septicemia (A40-A41)	3,386	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,259	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,688	0.6	---
15	Atherosclerosis (I70)	1,140	0.4	---
...	All other causes (residual)	60,867	22.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	110,707	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,193	32.7	---
2	Alzheimer's disease (G30)	10,996	9.9	---
3	Cerebrovascular diseases (I60-I69)	8,109	7.3	---
4	Malignant neoplasms (C00-C97)	7,253	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	4,183	3.8	---
6	Influenza and pneumonia (J09-J18)	3,989	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,092	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,072	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,925	1.7	---
10	Pneumonitis due to solids and liquids (J69)	1,344	1.2	---
11	Diabetes mellitus (E10-E14)	1,310	1.2	---
12	Septicemia (A40-A41)	1,153	1.0	---
13	Parkinson's disease (G20-G21)	818	0.7	---
14	Atherosclerosis (I70)	561	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	510	0.5	---
...	All other causes (residual)	27,199	24.6	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	22,721	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,945	35.0	---
2	Alzheimer's disease (G30)	2,125	9.4	---
3	Cerebrovascular diseases (I60-I69)	1,566	6.9	---
4	Influenza and pneumonia (J09-J18)	957	4.2	---
5	Malignant neoplasms (C00-C97)	906	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	658	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	554	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	519	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	1.6	---
10	Pneumonitis due to solids and liquids (J69)	266	1.2	---
11	Diabetes mellitus (E10-E14)	200	0.9	---
12	Septicemia (A40-A41)	190	0.8	---
13	Atherosclerosis (I70)	181	0.8	---
14	Nutritional deficiencies (E40-E64)	126	0.6	---
15	Parkinson's disease (G20-G21)	73	0.3	---
...	All other causes (residual)	6,091	26.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,063,705	100.0	1,072.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,784	24.9	267.0
2	Malignant neoplasms (C00-C97)	247,645	23.3	249.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,715	6.5	69.3
4	Chronic lower respiratory diseases (J40-J47)	62,857	5.9	63.4
5	Cerebrovascular diseases (I60-I69)	43,100	4.1	43.5
6	Diabetes mellitus (E10-E14)	29,813	2.8	30.1
7	Alzheimer's disease (G30)	28,631	2.7	28.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,045	2.6	28.3
9	Influenza and pneumonia (J09-J18)	20,995	2.0	21.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,546	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,351	1.7	18.5
12	Parkinson's disease (G20-G21)	14,778	1.4	14.9
13	Septicemia (A40-A41)	14,309	1.3	14.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,526	0.9	9.6
15	Pneumonitis due to solids and liquids (J69)	9,085	0.9	9.2
...	All other causes (residual)	184,525	17.3	186.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,083	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	37.0	9.4
2	Malignant neoplasms (C00-C97)	116	10.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
6	Influenza and pneumonia (J09-J18)	17	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
8	Cerebrovascular diseases (I60-I69)	10	0.9	*
9	Septicemia (A40-A41)	9	0.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
13	Anemias (D50-D64)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	267	24.7	6.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	669	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	31.7	3.8
2	Malignant neoplasms (C00-C97)	129	19.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	8.1	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.0	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.7	*
6	Influenza and pneumonia (J09-J18)	13	1.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
8	Cerebrovascular diseases (I60-I69)	9	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
10	Septicemia (A40-A41)	5	0.7	*
11	Anemias (D50-D64)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	154	23.0	2.8
Non-Hispanic, white, male, 10-14 years				
...	All causes	965	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	29.1	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	19.2	3.2
3	Malignant neoplasms (C00-C97)	118	12.2	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	5.4	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	12	1.2	*
8	Influenza and pneumonia (J09-J18)	12	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Septicemia (A40-A41)	7	0.7	*
12	Diabetes mellitus (E10-E14)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (residual)	195	20.2	3.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,692	100.0	61.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,664	45.1	27.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,063	28.8	17.6
3	Malignant neoplasms (C00-C97)	179	4.8	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	4.6	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.7	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.6	1.0
7	Influenza and pneumonia (J09-J18)	25	0.7	0.4
8	Septicemia (A40-A41)	12	0.3	*
8	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
11	Diabetes mellitus (E10-E14)	10	0.3	*
11	Cerebrovascular diseases (I60-I69)	10	0.3	*
13	Legal intervention (Y35,Y89.0)	8	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
...	All other causes (residual)	359	9.7	5.9
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,061	100.0	123.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,141	51.4	63.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,885	23.4	28.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	3.9	4.8
4	Malignant neoplasms (C00-C97)	289	3.6	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	2.9	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.8	1.0
7	Diabetes mellitus (E10-E14)	48	0.6	0.7
8	Influenza and pneumonia (J09-J18)	37	0.5	0.6
9	Chronic lower respiratory diseases (J40-J47)	34	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	33	0.4	0.5
11	Legal intervention (Y35,Y89.0)	23	0.3	0.4
12	Septicemia (A40-A41)	20	0.2	0.3
13	Pneumonitis due to solids and liquids (J69)	17	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.2	*
...	All other causes (residual)	891	11.1	13.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	10,097	100.0	154.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,223	51.7	80.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,016	20.0	30.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	435	4.3	6.7
4	Malignant neoplasms (C00-C97)	401	4.0	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	3.5	5.5
6	Diabetes mellitus (E10-E14)	87	0.9	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	0.8	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.7	1.1
9	Influenza and pneumonia (J09-J18)	53	0.5	0.8
10	Cerebrovascular diseases (I60-I69)	38	0.4	0.6
11	Septicemia (A40-A41)	36	0.4	0.6
12	Legal intervention (Y35,Y89.0)	32	0.3	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.3	0.4
14	Aortic aneurysm and dissection (I71)	23	0.2	0.4
14	Chronic lower respiratory diseases (J40-J47)	23	0.2	0.4
...	All other causes (residual)	1,197	11.9	18.3
Non-Hispanic, white, male, 30-34 years				
...	All causes	11,499	100.0	181.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,083	44.2	80.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,091	18.2	33.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	821	7.1	13.0
4	Malignant neoplasms (C00-C97)	676	5.9	10.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	354	3.1	5.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	240	2.1	3.8
7	Diabetes mellitus (E10-E14)	158	1.4	2.5
8	Cerebrovascular diseases (I60-I69)	102	0.9	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.8	1.5
10	Influenza and pneumonia (J09-J18)	65	0.6	1.0
11	Septicemia (A40-A41)	62	0.5	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.4	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.6
14	Legal intervention (Y35,Y89.0)	40	0.3	0.6
...	All other causes (residual)	1,591	13.8	25.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	12,398	100.0	209.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,682	37.8	79.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,994	16.1	33.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,364	11.0	23.0
4	Malignant neoplasms (C00-C97)	1,006	8.1	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	405	3.3	6.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	306	2.5	5.2
7	Diabetes mellitus (E10-E14)	225	1.8	3.8
8	Cerebrovascular diseases (I60-I69)	124	1.0	2.1
9	Influenza and pneumonia (J09-J18)	95	0.8	1.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	0.7	1.5
11	Septicemia (A40-A41)	77	0.6	1.3
12	Chronic lower respiratory diseases (J40-J47)	71	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	0.5	1.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.4	0.9
15	Aortic aneurysm and dissection (I71)	41	0.3	0.7
...	All other causes (residual)	1,804	14.6	30.5
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,039	100.0	266.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,191	26.1	69.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,747	17.1	45.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,194	13.7	36.4
4	Malignant neoplasms (C00-C97)	1,919	12.0	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	741	4.6	12.3
6	Diabetes mellitus (E10-E14)	442	2.8	7.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	1.8	4.8
8	Cerebrovascular diseases (I60-I69)	288	1.8	4.8
9	Septicemia (A40-A41)	138	0.9	2.3
10	Influenza and pneumonia (J09-J18)	134	0.8	2.2
11	Chronic lower respiratory diseases (J40-J47)	131	0.8	2.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	107	0.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.6	1.6
14	Aortic aneurysm and dissection (I71)	90	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	0.5	1.5
...	All other causes (residual)	2,446	15.3	40.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	25,542	100.0	386.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,370	21.0	81.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,722	18.5	71.4
3	Malignant neoplasms (C00-C97)	4,276	16.7	64.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,494	9.8	37.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,320	5.2	20.0
6	Diabetes mellitus (E10-E14)	798	3.1	12.1
7	Cerebrovascular diseases (I60-I69)	506	2.0	7.7
8	Chronic lower respiratory diseases (J40-J47)	358	1.4	5.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	1.2	4.7
10	Septicemia (A40-A41)	277	1.1	4.2
11	Influenza and pneumonia (J09-J18)	260	1.0	3.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	247	1.0	3.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	0.8	3.1
14	Viral hepatitis (B15-B19)	191	0.7	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	154	0.6	2.3
...	All other causes (residual)	4,058	15.9	61.4
Non-Hispanic, white, male, 50-54 years				
...	All causes	46,369	100.0	619.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,682	23.0	142.6
2	Malignant neoplasms (C00-C97)	10,528	22.7	140.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,591	12.1	74.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,034	6.5	40.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,616	5.6	34.9
6	Diabetes mellitus (E10-E14)	1,527	3.3	20.4
7	Chronic lower respiratory diseases (J40-J47)	1,220	2.6	16.3
8	Cerebrovascular diseases (I60-I69)	1,032	2.2	13.8
9	Septicemia (A40-A41)	518	1.1	6.9
10	Influenza and pneumonia (J09-J18)	475	1.0	6.3
11	Viral hepatitis (B15-B19)	464	1.0	6.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	0.9	5.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	369	0.8	4.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	353	0.8	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	0.8	4.7
...	All other causes (residual)	7,175	15.5	95.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	69,459	100.0	913.5
1	Malignant neoplasms (C00-C97)	19,920	28.7	262.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,461	23.7	216.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,463	7.9	71.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,346	4.8	44.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,921	4.2	38.4
6	Chronic lower respiratory diseases (J40-J47)	2,770	4.0	36.4
7	Diabetes mellitus (E10-E14)	2,400	3.5	31.6
8	Cerebrovascular diseases (I60-I69)	1,709	2.5	22.5
9	Septicemia (A40-A41)	870	1.3	11.4
10	Viral hepatitis (B15-B19)	785	1.1	10.3
11	Influenza and pneumonia (J09-J18)	725	1.0	9.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	705	1.0	9.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	578	0.8	7.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	322	0.5	4.2
15	Aortic aneurysm and dissection (I71)	309	0.4	4.1
...	All other causes (residual)	10,175	14.6	133.8
Non-Hispanic, white, male, 60-64 years				
...	All causes	88,164	100.0	1,293.8
1	Malignant neoplasms (C00-C97)	27,909	31.7	409.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,507	24.4	315.6
3	Chronic lower respiratory diseases (J40-J47)	4,816	5.5	70.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,347	4.9	63.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,261	3.7	47.9
6	Diabetes mellitus (E10-E14)	3,159	3.6	46.4
7	Cerebrovascular diseases (I60-I69)	2,461	2.8	36.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,173	2.5	31.9
9	Septicemia (A40-A41)	1,251	1.4	18.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,098	1.2	16.1
11	Influenza and pneumonia (J09-J18)	1,043	1.2	15.3
12	Viral hepatitis (B15-B19)	880	1.0	12.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	790	0.9	11.6
14	Aortic aneurysm and dissection (I71)	441	0.5	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	435	0.5	6.4
...	All other causes (residual)	12,593	14.3	184.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	105,474	100.0	1,790.6
1	Malignant neoplasms (C00-C97)	35,234	33.4	598.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,392	24.1	431.1
3	Chronic lower respiratory diseases (J40-J47)	7,194	6.8	122.1
4	Diabetes mellitus (E10-E14)	3,956	3.8	67.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,625	3.4	61.5
6	Cerebrovascular diseases (I60-I69)	3,313	3.1	56.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,441	2.3	41.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,729	1.6	29.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,557	1.5	26.4
10	Septicemia (A40-A41)	1,519	1.4	25.8
11	Influenza and pneumonia (J09-J18)	1,445	1.4	24.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	916	0.9	15.6
13	Parkinson's disease (G20-G21)	684	0.6	11.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	637	0.6	10.8
15	Alzheimer's disease (G30)	591	0.6	10.0
...	All other causes (residual)	15,241	14.5	258.7
Non-Hispanic, white, male, 70-74 years				
...	All causes	114,786	100.0	2,736.6
1	Malignant neoplasms (C00-C97)	36,680	32.0	874.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,273	23.8	650.2
3	Chronic lower respiratory diseases (J40-J47)	9,439	8.2	225.0
4	Cerebrovascular diseases (I60-I69)	4,407	3.8	105.1
5	Diabetes mellitus (E10-E14)	3,981	3.5	94.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,038	2.6	72.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,990	1.7	47.4
8	Influenza and pneumonia (J09-J18)	1,878	1.6	44.8
9	Septicemia (A40-A41)	1,703	1.5	40.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,609	1.4	38.4
11	Parkinson's disease (G20-G21)	1,503	1.3	35.8
12	Alzheimer's disease (G30)	1,498	1.3	35.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,342	1.2	32.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	892	0.8	21.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	873	0.8	20.8
...	All other causes (residual)	16,680	14.5	397.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	123,883	100.0	4,316.7
1	Malignant neoplasms (C00-C97)	34,739	28.0	1,210.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,866	24.1	1,040.7
3	Chronic lower respiratory diseases (J40-J47)	10,416	8.4	362.9
4	Cerebrovascular diseases (I60-I69)	5,646	4.6	196.7
5	Diabetes mellitus (E10-E14)	3,669	3.0	127.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,385	2.7	118.0
7	Alzheimer's disease (G30)	3,247	2.6	113.1
8	Parkinson's disease (G20-G21)	2,656	2.1	92.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,353	1.9	82.0
10	Influenza and pneumonia (J09-J18)	2,343	1.9	81.6
11	Septicemia (A40-A41)	1,883	1.5	65.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,126	0.9	39.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,084	0.9	37.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,068	0.9	37.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,044	0.8	36.4
...	All other causes (residual)	19,358	15.6	674.5
Non-Hispanic, white, male, 80-84 years				
...	All causes	141,645	100.0	7,236.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,724	25.9	1,876.1
2	Malignant neoplasms (C00-C97)	32,416	22.9	1,656.0
3	Chronic lower respiratory diseases (J40-J47)	10,474	7.4	535.1
4	Cerebrovascular diseases (I60-I69)	7,484	5.3	382.3
5	Alzheimer's disease (G30)	6,049	4.3	309.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,905	2.8	199.5
7	Diabetes mellitus (E10-E14)	3,842	2.7	196.3
8	Parkinson's disease (G20-G21)	3,791	2.7	193.7
9	Influenza and pneumonia (J09-J18)	3,359	2.4	171.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,106	2.2	158.7
11	Septicemia (A40-A41)	2,067	1.5	105.6
12	Pneumonitis due to solids and liquids (J69)	1,465	1.0	74.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,293	0.9	66.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,253	0.9	64.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	874	0.6	44.6
...	All other causes (residual)	23,543	16.6	1,202.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	145,196	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,725	28.7	---
2	Malignant neoplasms (C00-C97)	25,381	17.5	---
3	Chronic lower respiratory diseases (J40-J47)	9,363	6.4	---
4	Cerebrovascular diseases (I60-I69)	8,234	5.7	---
5	Alzheimer's disease (G30)	8,183	5.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,327	3.0	---
7	Influenza and pneumonia (J09-J18)	4,069	2.8	---
8	Parkinson's disease (G20-G21)	3,628	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,441	2.4	---
10	Diabetes mellitus (E10-E14)	3,207	2.2	---
11	Septicemia (A40-A41)	2,115	1.5	---
12	Pneumonitis due to solids and liquids (J69)	1,948	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,492	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,244	0.9	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	649	0.4	---
...	All other causes (residual)	26,190	18.0	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	98,276	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,584	32.1	---
2	Malignant neoplasms (C00-C97)	12,659	12.9	---
3	Alzheimer's disease (G30)	6,426	6.5	---
4	Cerebrovascular diseases (I60-I69)	5,683	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	4,947	5.0	---
6	Influenza and pneumonia (J09-J18)	3,370	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,968	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,359	2.4	---
9	Diabetes mellitus (E10-E14)	1,843	1.9	---
10	Parkinson's disease (G20-G21)	1,763	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,687	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,304	1.3	---
13	Septicemia (A40-A41)	1,288	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	737	0.7	---
15	Atherosclerosis (I70)	364	0.4	---
...	All other causes (residual)	19,294	19.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	30,605	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,813	35.3	---
2	Malignant neoplasms (C00-C97)	2,805	9.2	---
3	Alzheimer's disease (G30)	2,033	6.6	---
4	Cerebrovascular diseases (I60-I69)	1,718	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,362	4.5	---
6	Influenza and pneumonia (J09-J18)	1,319	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	955	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	747	2.4	---
9	Pneumonitis due to solids and liquids (J69)	600	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	488	1.6	---
11	Diabetes mellitus (E10-E14)	429	1.4	---
12	Septicemia (A40-A41)	366	1.2	---
13	Parkinson's disease (G20-G21)	350	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	176	0.6	---
15	Atherosclerosis (I70)	148	0.5	---
...	All other causes (residual)	6,296	20.6	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	4,008	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,513	37.7	---
2	Cerebrovascular diseases (I60-I69)	257	6.4	---
3	Malignant neoplasms (C00-C97)	245	6.1	---
4	Alzheimer's disease (G30)	242	6.0	---
5	Influenza and pneumonia (J09-J18)	210	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	144	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.7	---
9	Pneumonitis due to solids and liquids (J69)	89	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.8	---
11	Septicemia (A40-A41)	47	1.2	---
12	Atherosclerosis (I70)	37	0.9	---
13	Diabetes mellitus (E10-E14)	26	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	---
15	Nutritional deficiencies (E40-E64)	22	0.5	---
...	All other causes (residual)	864	21.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,059,926	100.0	1,038.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,388	22.5	233.6
2	Malignant neoplasms (C00-C97)	219,563	20.7	215.1
3	Chronic lower respiratory diseases (J40-J47)	73,371	6.9	71.9
4	Alzheimer's disease (G30)	64,694	6.1	63.4
5	Cerebrovascular diseases (I60-I69)	63,730	6.0	62.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,112	4.1	42.2
7	Influenza and pneumonia (J09-J18)	24,247	2.3	23.8
8	Diabetes mellitus (E10-E14)	23,777	2.2	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,130	1.6	16.8
10	Septicemia (A40-A41)	15,931	1.5	15.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,102	1.2	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,797	1.0	10.6
13	Parkinson's disease (G20-G21)	9,561	0.9	9.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,420	0.8	8.3
15	Pneumonitis due to solids and liquids (J69)	7,345	0.7	7.2
...	All other causes (residual)	226,758	21.4	222.2
Non-Hispanic, white, female, 1-4 years				
...	All causes	812	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	28.3	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	11.3	2.3
3	Malignant neoplasms (C00-C97)	90	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	7.0	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.9	0.8
6	Influenza and pneumonia (J09-J18)	20	2.5	0.5
7	Septicemia (A40-A41)	11	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
10	Meningitis (G00,G03)	6	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	237	29.2	5.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	523	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	31.0	3.1
2	Malignant neoplasms (C00-C97)	112	21.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.3	*
6	Influenza and pneumonia (J09-J18)	12	2.3	*
8	Cerebrovascular diseases (I60-I69)	11	2.1	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.9	*
10	Septicemia (A40-A41)	7	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Anemias (D50-D64)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
...	All other causes (residual)	104	19.9	2.0
Non-Hispanic, white, female, 10-14 years				
...	All causes	678	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	21.1	2.6
2	Malignant neoplasms (C00-C97)	117	17.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	15.3	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	4.9	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.7	*
7	Influenza and pneumonia (J09-J18)	10	1.5	*
8	Septicemia (A40-A41)	9	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
8	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	8	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Anemias (D50-D64)	3	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	166	24.5	3.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,764	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	785	44.5	13.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	19.7	6.1
3	Malignant neoplasms (C00-C97)	128	7.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	3.9	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	2.8	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	2.8	0.9
7	Cerebrovascular diseases (I60-I69)	19	1.1	*
7	Influenza and pneumonia (J09-J18)	19	1.1	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
10	Diabetes mellitus (E10-E14)	12	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
15	Meningitis (G00,G03)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (residual)	233	13.2	4.1
Non-Hispanic, white, female, 20-24 years				
...	All causes	3,019	100.0	48.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,489	49.3	24.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	381	12.6	6.2
3	Malignant neoplasms (C00-C97)	185	6.1	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	4.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	3.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.7	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.7	0.8
8	Diabetes mellitus (E10-E14)	34	1.1	0.5
9	Influenza and pneumonia (J09-J18)	26	0.9	0.4
10	Septicemia (A40-A41)	23	0.8	0.4
11	Cerebrovascular diseases (I60-I69)	21	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
...	All other causes (residual)	464	15.4	7.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	4,252	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,791	42.1	28.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	11.3	7.6
3	Malignant neoplasms (C00-C97)	356	8.4	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	254	6.0	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	3.8	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	103	2.4	1.6
7	Diabetes mellitus (E10-E14)	65	1.5	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.4	0.9
9	Influenza and pneumonia (J09-J18)	50	1.2	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.1	0.7
11	Cerebrovascular diseases (I60-I69)	46	1.1	0.7
12	Septicemia (A40-A41)	40	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	24	0.6	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (residual)	747	17.6	11.8
Non-Hispanic, white, female, 30-34 years				
...	All causes	5,644	100.0	90.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,046	36.3	32.9
2	Malignant neoplasms (C00-C97)	698	12.4	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	591	10.5	9.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	418	7.4	6.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	2.9	2.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	152	2.7	2.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	99	1.8	1.6
8	Diabetes mellitus (E10-E14)	98	1.7	1.6
9	Cerebrovascular diseases (I60-I69)	70	1.2	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.1	1.0
11	Septicemia (A40-A41)	60	1.1	1.0
12	Influenza and pneumonia (J09-J18)	58	1.0	0.9
13	Chronic lower respiratory diseases (J40-J47)	40	0.7	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.4	0.4
...	All other causes (residual)	1,038	18.4	16.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,994	100.0	119.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,950	27.9	33.4
2	Malignant neoplasms (C00-C97)	1,190	17.0	20.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	9.6	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	660	9.4	11.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	233	3.3	4.0
6	Diabetes mellitus (E10-E14)	153	2.2	2.6
7	Cerebrovascular diseases (I60-I69)	142	2.0	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	1.8	2.1
9	Septicemia (A40-A41)	82	1.2	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	81	1.2	1.4
11	Influenza and pneumonia (J09-J18)	78	1.1	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.0	1.2
13	Chronic lower respiratory diseases (J40-J47)	58	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.7	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.5	0.6
...	All other causes (residual)	1,420	20.3	24.3
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,206	100.0	170.9
1	Malignant neoplasms (C00-C97)	2,421	23.7	40.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,068	20.3	34.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,203	11.8	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	775	7.6	13.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	453	4.4	7.6
6	Diabetes mellitus (E10-E14)	248	2.4	4.2
7	Cerebrovascular diseases (I60-I69)	242	2.4	4.1
8	Chronic lower respiratory diseases (J40-J47)	184	1.8	3.1
9	Septicemia (A40-A41)	166	1.6	2.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	146	1.4	2.4
11	Influenza and pneumonia (J09-J18)	117	1.1	2.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	0.9	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	0.8	1.4
14	Viral hepatitis (B15-B19)	51	0.5	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	49	0.5	0.8
...	All other causes (residual)	1,913	18.7	32.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	16,773	100.0	253.9
1	Malignant neoplasms (C00-C97)	4,786	28.5	72.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,462	14.7	37.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,308	13.8	34.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	894	5.3	13.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	861	5.1	13.0
6	Diabetes mellitus (E10-E14)	503	3.0	7.6
7	Chronic lower respiratory diseases (J40-J47)	469	2.8	7.1
8	Cerebrovascular diseases (I60-I69)	454	2.7	6.9
9	Septicemia (A40-A41)	270	1.6	4.1
10	Influenza and pneumonia (J09-J18)	234	1.4	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	0.9	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	0.8	2.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.6	1.6
14	Viral hepatitis (B15-B19)	79	0.5	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	0.5	1.2
...	All other causes (residual)	2,983	17.8	45.1
Non-Hispanic, white, female, 50-54 years				
...	All causes	29,894	100.0	391.5
1	Malignant neoplasms (C00-C97)	9,976	33.4	130.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,287	14.3	56.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,102	10.4	40.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,417	4.7	18.6
5	Chronic lower respiratory diseases (J40-J47)	1,416	4.7	18.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,124	3.8	14.7
7	Cerebrovascular diseases (I60-I69)	883	3.0	11.6
8	Diabetes mellitus (E10-E14)	807	2.7	10.6
9	Septicemia (A40-A41)	485	1.6	6.4
10	Influenza and pneumonia (J09-J18)	360	1.2	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	1.0	3.8
12	Viral hepatitis (B15-B19)	248	0.8	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	0.6	2.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	0.6	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	0.4	1.7
...	All other causes (residual)	5,011	16.8	65.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	43,983	100.0	558.7
1	Malignant neoplasms (C00-C97)	16,510	37.5	209.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,644	15.1	84.4
3	Chronic lower respiratory diseases (J40-J47)	2,827	6.4	35.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,776	6.3	35.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,615	3.7	20.5
6	Diabetes mellitus (E10-E14)	1,355	3.1	17.2
7	Cerebrovascular diseases (I60-I69)	1,331	3.0	16.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,082	2.5	13.7
9	Septicemia (A40-A41)	740	1.7	9.4
10	Influenza and pneumonia (J09-J18)	598	1.4	7.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	1.1	6.0
12	Viral hepatitis (B15-B19)	335	0.8	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	0.7	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	0.5	3.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	215	0.5	2.7
...	All other causes (residual)	6,963	15.8	88.4
Non-Hispanic, white, female, 60-64 years				
...	All causes	55,906	100.0	772.2
1	Malignant neoplasms (C00-C97)	21,355	38.2	295.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,294	16.6	128.4
3	Chronic lower respiratory diseases (J40-J47)	4,244	7.6	58.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,989	3.6	27.5
5	Diabetes mellitus (E10-E14)	1,913	3.4	26.4
6	Cerebrovascular diseases (I60-I69)	1,847	3.3	25.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,506	2.7	20.8
8	Septicemia (A40-A41)	1,011	1.8	14.0
9	Influenza and pneumonia (J09-J18)	836	1.5	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	773	1.4	10.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	753	1.3	10.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	420	0.8	5.8
13	Viral hepatitis (B15-B19)	357	0.6	4.9
14	Alzheimer's disease (G30)	314	0.6	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	298	0.5	4.1
...	All other causes (residual)	8,996	16.1	124.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	74,134	100.0	1,163.1
1	Malignant neoplasms (C00-C97)	27,874	37.6	437.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,605	17.0	197.8
3	Chronic lower respiratory diseases (J40-J47)	6,839	9.2	107.3
4	Cerebrovascular diseases (I60-I69)	2,727	3.7	42.8
5	Diabetes mellitus (E10-E14)	2,514	3.4	39.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,800	2.4	28.2
7	Septicemia (A40-A41)	1,440	1.9	22.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,245	1.7	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,233	1.7	19.3
10	Influenza and pneumonia (J09-J18)	1,186	1.6	18.6
11	Alzheimer's disease (G30)	793	1.1	12.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	617	0.8	9.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	480	0.6	7.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	424	0.6	6.7
15	Pneumonitis due to solids and liquids (J69)	332	0.4	5.2
...	All other causes (residual)	12,025	16.2	188.7
Non-Hispanic, white, female, 70-74 years				
...	All causes	89,700	100.0	1,888.3
1	Malignant neoplasms (C00-C97)	30,019	33.5	631.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,021	17.9	337.3
3	Chronic lower respiratory diseases (J40-J47)	9,780	10.9	205.9
4	Cerebrovascular diseases (I60-I69)	4,119	4.6	86.7
5	Diabetes mellitus (E10-E14)	2,662	3.0	56.0
6	Alzheimer's disease (G30)	2,123	2.4	44.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,848	2.1	38.9
8	Septicemia (A40-A41)	1,649	1.8	34.7
9	Influenza and pneumonia (J09-J18)	1,603	1.8	33.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,548	1.7	32.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,032	1.2	21.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	775	0.9	16.3
13	Parkinson's disease (G20-G21)	696	0.8	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	607	0.7	12.8
15	Pneumonitis due to solids and liquids (J69)	460	0.5	9.7
...	All other causes (residual)	14,758	16.5	310.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	108,197	100.0	3,109.8
1	Malignant neoplasms (C00-C97)	29,497	27.3	847.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,885	19.3	600.3
3	Chronic lower respiratory diseases (J40-J47)	11,276	10.4	324.1
4	Cerebrovascular diseases (I60-I69)	6,181	5.7	177.7
5	Alzheimer's disease (G30)	4,887	4.5	140.5
6	Diabetes mellitus (E10-E14)	2,921	2.7	84.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,414	2.2	69.4
8	Influenza and pneumonia (J09-J18)	2,184	2.0	62.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,029	1.9	58.3
10	Septicemia (A40-A41)	1,929	1.8	55.4
11	Parkinson's disease (G20-G21)	1,362	1.3	39.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,120	1.0	32.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	853	0.8	24.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	733	0.7	21.1
15	Pneumonitis due to solids and liquids (J69)	707	0.7	20.3
...	All other causes (residual)	19,219	17.8	552.4
Non-Hispanic, white, female, 80-84 years				
...	All causes	143,961	100.0	5,399.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,435	22.5	1,216.5
2	Malignant neoplasms (C00-C97)	28,615	19.9	1,073.2
3	Chronic lower respiratory diseases (J40-J47)	12,307	8.5	461.6
4	Alzheimer's disease (G30)	10,339	7.2	387.8
5	Cerebrovascular diseases (I60-I69)	9,851	6.8	369.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,529	2.5	132.4
7	Diabetes mellitus (E10-E14)	3,332	2.3	125.0
8	Influenza and pneumonia (J09-J18)	3,279	2.3	123.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,681	1.9	100.6
10	Septicemia (A40-A41)	2,197	1.5	82.4
11	Parkinson's disease (G20-G21)	2,175	1.5	81.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,798	1.2	67.4
13	Pneumonitis due to solids and liquids (J69)	1,054	0.7	39.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,017	0.7	38.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	699	0.5	26.2
...	All other causes (residual)	28,653	19.9	1,074.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	186,447	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,209	25.9	---
2	Malignant neoplasms (C00-C97)	25,255	13.5	---
3	Alzheimer's disease (G30)	16,931	9.1	---
4	Cerebrovascular diseases (I60-I69)	14,409	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	12,123	6.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,044	2.7	---
7	Influenza and pneumonia (J09-J18)	4,971	2.7	---
8	Diabetes mellitus (E10-E14)	3,560	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,327	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,747	1.5	---
11	Septicemia (A40-A41)	2,745	1.5	---
12	Parkinson's disease (G20-G21)	2,553	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,631	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,152	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	773	0.4	---
...	All other causes (residual)	41,017	22.0	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	173,685	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,048	29.4	---
2	Alzheimer's disease (G30)	18,305	10.5	---
3	Malignant neoplasms (C00-C97)	15,255	8.8	---
4	Cerebrovascular diseases (I60-I69)	13,647	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	8,398	4.8	---
6	Influenza and pneumonia (J09-J18)	5,159	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,663	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,984	1.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,963	1.7	---
10	Diabetes mellitus (E10-E14)	2,537	1.5	---
11	Septicemia (A40-A41)	2,098	1.2	---
12	Parkinson's disease (G20-G21)	1,756	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,572	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	951	0.5	---
15	Atherosclerosis (I70)	776	0.4	---
...	All other causes (residual)	41,573	23.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	80,102	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,380	31.7	---
2	Alzheimer's disease (G30)	8,963	11.2	---
3	Cerebrovascular diseases (I60-I69)	6,391	8.0	---
4	Malignant neoplasms (C00-C97)	4,448	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	2,821	3.5	---
6	Influenza and pneumonia (J09-J18)	2,670	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,137	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,584	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,178	1.5	---
10	Diabetes mellitus (E10-E14)	881	1.1	---
11	Septicemia (A40-A41)	787	1.0	---
12	Pneumonitis due to solids and liquids (J69)	744	0.9	---
13	Parkinson's disease (G20-G21)	468	0.6	---
14	Atherosclerosis (I70)	413	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	334	0.4	---
...	All other causes (residual)	20,903	26.1	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	18,713	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,432	34.4	---
2	Alzheimer's disease (G30)	1,883	10.1	---
3	Cerebrovascular diseases (I60-I69)	1,309	7.0	---
4	Influenza and pneumonia (J09-J18)	747	4.0	---
5	Malignant neoplasms (C00-C97)	661	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	514	2.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	446	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	445	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	1.4	---
10	Pneumonitis due to solids and liquids (J69)	177	0.9	---
11	Diabetes mellitus (E10-E14)	174	0.9	---
12	Atherosclerosis (I70)	144	0.8	---
13	Septicemia (A40-A41)	143	0.8	---
14	Nutritional deficiencies (E40-E64)	104	0.6	---
15	Parkinson's disease (G20-G21)	55	0.3	---
...	All other causes (residual)	5,222	27.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	315,254	100.0	754.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,093	23.5	177.4
2	Malignant neoplasms (C00-C97)	68,523	21.7	164.0
3	Cerebrovascular diseases (I60-I69)	17,760	5.6	42.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,366	4.9	36.8
5	Diabetes mellitus (E10-E14)	13,693	4.3	32.8
6	Chronic lower respiratory diseases (J40-J47)	10,327	3.3	24.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,083	2.9	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,038	2.9	21.6
9	Alzheimer's disease (G30)	8,072	2.6	19.3
10	Septicemia (A40-A41)	6,576	2.1	15.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,740	1.8	13.7
12	Influenza and pneumonia (J09-J18)	5,554	1.8	13.3
13	Certain conditions originating in the perinatal period (P00-P96)	3,827	1.2	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,379	1.1	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,185	1.0	7.6
...	All other causes (residual)	61,038	19.4	146.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.1	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	9.2	3.8
4	Malignant neoplasms (C00-C97)	53	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	27	2.7	1.1
6	Certain conditions originating in the perinatal period (P00-P96)	27	2.7	1.1
8	Chronic lower respiratory diseases (J40-J47)	23	2.3	0.9
9	Cerebrovascular diseases (I60-I69)	17	1.7	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (residual)	250	24.6	10.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	553	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	32.5	5.8
2	Malignant neoplasms (C00-C97)	61	11.0	2.0
3	Chronic lower respiratory diseases (J40-J47)	49	8.9	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.4	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.9	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.1	0.9
7	Anemias (D50-D64)	12	2.2	*
8	Influenza and pneumonia (J09-J18)	11	2.0	*
9	Cerebrovascular diseases (I60-I69)	10	1.8	*
10	Septicemia (A40-A41)	9	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
13	Diseases of appendix (K35-K38)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	90	16.3	2.9
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	634	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	23.2	4.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	12.6	2.6
3	Malignant neoplasms (C00-C97)	65	10.3	2.1
4	Chronic lower respiratory diseases (J40-J47)	53	8.4	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	7.1	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.7	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.4	0.9
8	Cerebrovascular diseases (I60-I69)	12	1.9	*
9	Influenza and pneumonia (J09-J18)	8	1.3	*
10	Septicemia (A40-A41)	7	1.1	*
10	Diabetes mellitus (E10-E14)	7	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Anemias (D50-D64)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (residual)	135	21.3	4.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,336	100.0	72.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	999	42.8	31.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	23.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	8.5	6.2
4	Malignant neoplasms (C00-C97)	91	3.9	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.6	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.7	1.2
7	Chronic lower respiratory diseases (J40-J47)	30	1.3	0.9
8	Anemias (D50-D64)	24	1.0	0.7
9	Diabetes mellitus (E10-E14)	19	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.7	*
11	Septicemia (A40-A41)	11	0.5	*
11	Influenza and pneumonia (J09-J18)	11	0.5	*
13	Cerebrovascular diseases (I60-I69)	10	0.4	*
14	Legal intervention (Y35,Y89.0)	9	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
...	All other causes (residual)	228	9.8	7.1
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,760	100.0	133.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,014	42.3	56.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,092	22.9	30.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	7.9	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	218	4.6	6.1
5	Malignant neoplasms (C00-C97)	161	3.4	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.5	2.0
7	Chronic lower respiratory diseases (J40-J47)	56	1.2	1.6
8	Anemias (D50-D64)	46	1.0	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	46	1.0	1.3
10	Diabetes mellitus (E10-E14)	45	0.9	1.3
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.6	0.8
13	Legal intervention (Y35,Y89.0)	25	0.5	0.7
14	Influenza and pneumonia (J09-J18)	24	0.5	0.7
15	Septicemia (A40-A41)	23	0.5	0.6
...	All other causes (residual)	509	10.7	14.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	5,006	100.0	154.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,670	33.4	51.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,122	22.4	34.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	7.2	11.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	6.7	10.4
5	Malignant neoplasms (C00-C97)	212	4.2	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	2.9	4.4
7	Diabetes mellitus (E10-E14)	90	1.8	2.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.3	2.0
9	Cerebrovascular diseases (I60-I69)	54	1.1	1.7
10	Anemias (D50-D64)	50	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	48	1.0	1.5
12	Septicemia (A40-A41)	46	0.9	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.9	1.4
14	Influenza and pneumonia (J09-J18)	42	0.8	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.6	1.0
...	All other causes (residual)	689	13.8	21.3
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	5,310	100.0	185.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	22.8	42.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	19.4	36.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	616	11.6	21.5
4	Malignant neoplasms (C00-C97)	419	7.9	14.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	4.8	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	3.4	6.4
7	Diabetes mellitus (E10-E14)	162	3.1	5.7
8	Cerebrovascular diseases (I60-I69)	97	1.8	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.4	2.6
10	Septicemia (A40-A41)	70	1.3	2.4
11	Anemias (D50-D64)	69	1.3	2.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.3	2.4
13	Influenza and pneumonia (J09-J18)	56	1.1	2.0
14	Chronic lower respiratory diseases (J40-J47)	54	1.0	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.8	1.5
...	All other causes (residual)	902	17.0	31.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,273	100.0	233.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,080	17.2	40.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,042	16.6	38.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	859	13.7	32.0
4	Malignant neoplasms (C00-C97)	727	11.6	27.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	260	4.1	9.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	3.5	8.3
7	Diabetes mellitus (E10-E14)	217	3.5	8.1
8	Cerebrovascular diseases (I60-I69)	195	3.1	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.6	3.7
10	Chronic lower respiratory diseases (J40-J47)	83	1.3	3.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	1.3	2.9
12	Septicemia (A40-A41)	74	1.2	2.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	66	1.1	2.5
14	Anemias (D50-D64)	57	0.9	2.1
15	Influenza and pneumonia (J09-J18)	54	0.9	2.0
...	All other causes (residual)	1,160	18.5	43.2
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,029	100.0	307.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,779	22.2	68.1
2	Malignant neoplasms (C00-C97)	1,181	14.7	45.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,003	12.5	38.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	6.8	21.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	333	4.1	12.7
6	Cerebrovascular diseases (I60-I69)	329	4.1	12.6
7	Diabetes mellitus (E10-E14)	323	4.0	12.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	206	2.6	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	2.3	7.1
10	Septicemia (A40-A41)	143	1.8	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.7	5.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	133	1.7	5.1
13	Chronic lower respiratory diseases (J40-J47)	103	1.3	3.9
14	Influenza and pneumonia (J09-J18)	88	1.1	3.4
15	Aortic aneurysm and dissection (I71)	60	0.7	2.3
...	All other causes (residual)	1,481	18.4	56.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	11,774	100.0	446.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,754	23.4	104.5
2	Malignant neoplasms (C00-C97)	2,219	18.8	84.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,196	10.2	45.4
4	Diabetes mellitus (E10-E14)	563	4.8	21.4
5	Cerebrovascular diseases (I60-I69)	547	4.6	20.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	3.7	16.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	425	3.6	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	2.3	10.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	243	2.1	9.2
10	Septicemia (A40-A41)	240	2.0	9.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	188	1.6	7.1
12	Chronic lower respiratory diseases (J40-J47)	187	1.6	7.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	184	1.6	7.0
14	Influenza and pneumonia (J09-J18)	137	1.2	5.2
15	Aortic aneurysm and dissection (I71)	70	0.6	2.7
...	All other causes (residual)	2,125	18.0	80.6
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,398	100.0	703.1
1	Malignant neoplasms (C00-C97)	4,818	24.8	174.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,663	24.0	169.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,522	7.8	55.2
4	Diabetes mellitus (E10-E14)	923	4.8	33.5
5	Cerebrovascular diseases (I60-I69)	887	4.6	32.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	529	2.7	19.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	486	2.5	17.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	2.2	15.2
9	Chronic lower respiratory diseases (J40-J47)	418	2.2	15.2
10	Septicemia (A40-A41)	394	2.0	14.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	1.9	13.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	320	1.6	11.6
13	Influenza and pneumonia (J09-J18)	284	1.5	10.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	0.9	6.5
15	Viral hepatitis (B15-B19)	119	0.6	4.3
...	All other causes (residual)	3,078	15.9	111.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	28,015	100.0	1,078.9
1	Malignant neoplasms (C00-C97)	8,047	28.7	309.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,897	24.6	265.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,452	5.2	55.9
4	Diabetes mellitus (E10-E14)	1,376	4.9	53.0
5	Cerebrovascular diseases (I60-I69)	1,316	4.7	50.7
6	Chronic lower respiratory diseases (J40-J47)	836	3.0	32.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	734	2.6	28.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	637	2.3	24.5
9	Septicemia (A40-A41)	603	2.2	23.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	560	2.0	21.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	472	1.7	18.2
12	Influenza and pneumonia (J09-J18)	366	1.3	14.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	250	0.9	9.6
14	Viral hepatitis (B15-B19)	246	0.9	9.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	0.5	5.5
...	All other causes (residual)	4,079	14.6	157.1
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	32,871	100.0	1,571.1
1	Malignant neoplasms (C00-C97)	9,923	30.2	474.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,171	24.9	390.5
3	Diabetes mellitus (E10-E14)	1,682	5.1	80.4
3	Cerebrovascular diseases (I60-I69)	1,682	5.1	80.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,255	3.8	60.0
6	Chronic lower respiratory diseases (J40-J47)	1,085	3.3	51.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	2.8	44.6
8	Septicemia (A40-A41)	688	2.1	32.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	636	1.9	30.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	594	1.8	28.4
11	Influenza and pneumonia (J09-J18)	494	1.5	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	418	1.3	20.0
13	Viral hepatitis (B15-B19)	333	1.0	15.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	0.5	7.8
15	Pneumonitis due to solids and liquids (J69)	128	0.4	6.1
...	All other causes (residual)	4,686	14.3	224.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	32,541	100.0	2,069.7
1	Malignant neoplasms (C00-C97)	10,023	30.8	637.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,978	24.5	507.4
3	Cerebrovascular diseases (I60-I69)	1,758	5.4	111.8
4	Diabetes mellitus (E10-E14)	1,702	5.2	108.3
5	Chronic lower respiratory diseases (J40-J47)	1,281	3.9	81.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,108	3.4	70.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	783	2.4	49.8
8	Septicemia (A40-A41)	762	2.3	48.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	567	1.7	36.1
10	Influenza and pneumonia (J09-J18)	534	1.6	34.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	439	1.3	27.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	240	0.7	15.3
13	Viral hepatitis (B15-B19)	206	0.6	13.1
14	Alzheimer's disease (G30)	168	0.5	10.7
15	Pneumonitis due to solids and liquids (J69)	165	0.5	10.5
...	All other causes (residual)	4,827	14.8	307.0
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	30,777	100.0	2,964.9
1	Malignant neoplasms (C00-C97)	8,905	28.9	857.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,793	25.3	750.8
3	Cerebrovascular diseases (I60-I69)	1,874	6.1	180.5
4	Diabetes mellitus (E10-E14)	1,596	5.2	153.8
5	Chronic lower respiratory diseases (J40-J47)	1,421	4.6	136.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,040	3.4	100.2
7	Septicemia (A40-A41)	732	2.4	70.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	584	1.9	56.3
9	Influenza and pneumonia (J09-J18)	535	1.7	51.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	1.7	49.9
11	Alzheimer's disease (G30)	467	1.5	45.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	236	0.8	22.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	169	0.5	16.3
14	Pneumonitis due to solids and liquids (J69)	159	0.5	15.3
15	Parkinson's disease (G20-G21)	142	0.5	13.7
...	All other causes (residual)	4,606	15.0	443.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	30,976	100.0	4,311.4
1	Malignant neoplasms (C00-C97)	7,875	25.4	1,096.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,799	25.2	1,085.5
3	Cerebrovascular diseases (I60-I69)	2,165	7.0	301.3
4	Diabetes mellitus (E10-E14)	1,615	5.2	224.8
5	Chronic lower respiratory diseases (J40-J47)	1,390	4.5	193.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,145	3.7	159.4
7	Alzheimer's disease (G30)	909	2.9	126.5
8	Septicemia (A40-A41)	766	2.5	106.6
9	Influenza and pneumonia (J09-J18)	605	2.0	84.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	602	1.9	83.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	498	1.6	69.3
12	Parkinson's disease (G20-G21)	223	0.7	31.0
13	Pneumonitis due to solids and liquids (J69)	222	0.7	30.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.5	23.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	132	0.4	18.4
...	All other causes (residual)	4,862	15.7	676.7
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	30,485	100.0	6,427.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,763	25.5	1,636.7
2	Malignant neoplasms (C00-C97)	6,238	20.5	1,315.2
3	Cerebrovascular diseases (I60-I69)	2,354	7.7	496.3
4	Alzheimer's disease (G30)	1,526	5.0	321.7
5	Diabetes mellitus (E10-E14)	1,409	4.6	297.1
6	Chronic lower respiratory diseases (J40-J47)	1,339	4.4	282.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,076	3.5	226.9
8	Septicemia (A40-A41)	719	2.4	151.6
9	Influenza and pneumonia (J09-J18)	712	2.3	150.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	700	2.3	147.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	452	1.5	95.3
12	Parkinson's disease (G20-G21)	272	0.9	57.3
13	Pneumonitis due to solids and liquids (J69)	235	0.8	49.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	204	0.7	43.0
15	Aortic aneurysm and dissection (I71)	98	0.3	20.7
...	All other causes (residual)	5,388	17.7	1,136.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	27,456	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,395	26.9	---
2	Malignant neoplasms (C00-C97)	4,467	16.3	---
3	Cerebrovascular diseases (I60-I69)	2,124	7.7	---
4	Alzheimer's disease (G30)	1,914	7.0	---
5	Diabetes mellitus (E10-E14)	1,122	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,051	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	942	3.4	---
8	Septicemia (A40-A41)	635	2.3	---
9	Influenza and pneumonia (J09-J18)	629	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	601	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	1.4	---
12	Pneumonitis due to solids and liquids (J69)	245	0.9	---
13	Parkinson's disease (G20-G21)	240	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	171	0.6	---
15	Atherosclerosis (I70)	98	0.4	---
...	All other causes (residual)	5,424	19.8	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	19,189	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,395	28.1	---
2	Malignant neoplasms (C00-C97)	2,219	11.6	---
3	Alzheimer's disease (G30)	1,767	9.2	---
4	Cerebrovascular diseases (I60-I69)	1,473	7.7	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	661	3.4	---
6	Diabetes mellitus (E10-E14)	586	3.1	---
7	Influenza and pneumonia (J09-J18)	566	2.9	---
8	Chronic lower respiratory diseases (J40-J47)	552	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	530	2.8	---
10	Septicemia (A40-A41)	416	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	1.2	---
12	Pneumonitis due to solids and liquids (J69)	192	1.0	---
13	Parkinson's disease (G20-G21)	111	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	---
15	Nutritional deficiencies (E40-E64)	84	0.4	---
...	All other causes (residual)	4,299	22.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	8,333	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,419	29.0	---
2	Alzheimer's disease (G30)	890	10.7	---
3	Malignant neoplasms (C00-C97)	682	8.2	---
4	Cerebrovascular diseases (I60-I69)	624	7.5	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	2.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	2.7	---
7	Influenza and pneumonia (J09-J18)	219	2.6	---
8	Chronic lower respiratory diseases (J40-J47)	208	2.5	---
9	Diabetes mellitus (E10-E14)	201	2.4	---
10	Septicemia (A40-A41)	121	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	1.4	---
12	Pneumonitis due to solids and liquids (J69)	82	1.0	---
13	Nutritional deficiencies (E40-E64)	54	0.6	---
14	Parkinson's disease (G20-G21)	45	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	---
...	All other causes (residual)	2,157	25.9	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,598	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	762	29.3	---
2	Alzheimer's disease (G30)	314	12.1	---
3	Cerebrovascular diseases (I60-I69)	176	6.8	---
4	Malignant neoplasms (C00-C97)	128	4.9	---
5	Influenza and pneumonia (J09-J18)	95	3.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.9	---
8	Diabetes mellitus (E10-E14)	48	1.8	---
8	Chronic lower respiratory diseases (J40-J47)	48	1.8	---
10	Septicemia (A40-A41)	44	1.7	---
11	Pneumonitis due to solids and liquids (J69)	28	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.1	---
13	Nutritional deficiencies (E40-E64)	26	1.0	---
14	Atherosclerosis (I70)	14	0.5	---
15	Anemias (D50-D64)	8	0.3	---
...	All other causes (residual)	711	27.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	161,850	100.0	809.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,638	23.9	193.2
2	Malignant neoplasms (C00-C97)	34,787	21.5	174.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,516	6.5	52.6
4	Cerebrovascular diseases (I60-I69)	7,962	4.9	39.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,909	4.9	39.6
6	Diabetes mellitus (E10-E14)	6,806	4.2	34.0
7	Chronic lower respiratory diseases (J40-J47)	5,236	3.2	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,351	2.7	21.8
9	Septicemia (A40-A41)	3,061	1.9	15.3
10	Influenza and pneumonia (J09-J18)	2,729	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,599	1.6	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,230	1.4	11.2
13	Alzheimer's disease (G30)	2,154	1.3	10.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,130	1.3	10.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,951	1.2	9.8
...	All other causes (residual)	28,791	17.8	144.0
Non-Hispanic, black, male, 1-4 years				
...	All causes	608	100.0	48.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	29.6	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.8	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	8.6	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.3
5	Malignant neoplasms (C00-C97)	26	4.3	2.1
6	Certain conditions originating in the perinatal period (P00-P96)	19	3.1	*
7	Influenza and pneumonia (J09-J18)	17	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
9	Cerebrovascular diseases (I60-I69)	12	2.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.3	*
11	Septicemia (A40-A41)	7	1.2	*
11	Anemias (D50-D64)	7	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (residual)	137	22.5	11.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	317	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	31.9	6.4
2	Malignant neoplasms (C00-C97)	34	10.7	2.1
3	Chronic lower respiratory diseases (J40-J47)	30	9.5	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	7.3	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	7.3	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.7	*
7	Anemias (D50-D64)	5	1.6	*
7	Cerebrovascular diseases (I60-I69)	5	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
9	Influenza and pneumonia (J09-J18)	4	1.3	*
11	Septicemia (A40-A41)	3	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Meningitis (G00,G03)	2	0.6	*
13	Diseases of appendix (K35-K38)	2	0.6	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
...	All other causes (residual)	61	19.2	3.9
Non-Hispanic, black, male, 10-14 years				
...	All causes	386	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	22.3	5.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	15.5	3.9
3	Malignant neoplasms (C00-C97)	34	8.8	2.2
4	Chronic lower respiratory diseases (J40-J47)	33	8.5	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	8.5	2.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.9	*
8	Cerebrovascular diseases (I60-I69)	8	2.1	*
9	Diabetes mellitus (E10-E14)	6	1.6	*
10	Influenza and pneumonia (J09-J18)	5	1.3	*
11	Septicemia (A40-A41)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Anemias (D50-D64)	2	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Meningococcal infection (A39)	1	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	76	19.7	4.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,795	100.0	110.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	901	50.2	55.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	22.6	24.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	8.4	9.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.1	3.4
5	Malignant neoplasms (C00-C97)	50	2.8	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.3	1.4
7	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
8	Diabetes mellitus (E10-E14)	13	0.7	*
9	Legal intervention (Y35,Y89.0)	9	0.5	*
10	Anemias (D50-D64)	8	0.4	*
11	Cerebrovascular diseases (I60-I69)	6	0.3	*
11	Influenza and pneumonia (J09-J18)	6	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.1	*
...	All other causes (residual)	137	7.6	8.4
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,691	100.0	203.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,843	49.9	101.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	820	22.2	45.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	8.2	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	3.8	7.7
5	Malignant neoplasms (C00-C97)	88	2.4	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	43	1.2	2.4
7	Chronic lower respiratory diseases (J40-J47)	37	1.0	2.0
8	Anemias (D50-D64)	27	0.7	1.5
9	Diabetes mellitus (E10-E14)	26	0.7	1.4
10	Legal intervention (Y35,Y89.0)	23	0.6	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.6	1.2
12	Influenza and pneumonia (J09-J18)	15	0.4	*
13	Cerebrovascular diseases (I60-I69)	14	0.4	*
14	Septicemia (A40-A41)	11	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (residual)	268	7.3	14.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,680	100.0	230.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,506	40.9	94.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	22.6	52.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	280	7.6	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	6.1	14.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	109	3.0	6.8
6	Malignant neoplasms (C00-C97)	99	2.7	6.2
7	Diabetes mellitus (E10-E14)	44	1.2	2.8
8	Influenza and pneumonia (J09-J18)	32	0.9	2.0
8	Chronic lower respiratory diseases (J40-J47)	32	0.9	2.0
10	Cerebrovascular diseases (I60-I69)	30	0.8	1.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	1.8
12	Anemias (D50-D64)	27	0.7	1.7
13	Septicemia (A40-A41)	25	0.7	1.6
14	Legal intervention (Y35,Y89.0)	24	0.7	1.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
...	All other causes (residual)	372	10.1	23.3
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,549	100.0	258.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,086	30.6	79.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	769	21.7	55.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	10.5	27.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	5.9	15.3
5	Malignant neoplasms (C00-C97)	169	4.8	12.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	111	3.1	8.1
7	Diabetes mellitus (E10-E14)	87	2.5	6.3
8	Cerebrovascular diseases (I60-I69)	46	1.3	3.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.0	2.7
10	Septicemia (A40-A41)	36	1.0	2.6
10	Chronic lower respiratory diseases (J40-J47)	36	1.0	2.6
12	Influenza and pneumonia (J09-J18)	35	1.0	2.5
13	Anemias (D50-D64)	32	0.9	2.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.6	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.6	1.5
...	All other causes (residual)	458	12.9	33.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,942	100.0	310.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	20.2	62.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	771	19.6	60.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	672	17.0	52.9
4	Malignant neoplasms (C00-C97)	276	7.0	21.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	4.7	14.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	3.7	11.3
7	Diabetes mellitus (E10-E14)	129	3.3	10.1
8	Cerebrovascular diseases (I60-I69)	107	2.7	8.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.3	4.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.2	3.8
11	Chronic lower respiratory diseases (J40-J47)	43	1.1	3.4
12	Septicemia (A40-A41)	36	0.9	2.8
13	Influenza and pneumonia (J09-J18)	34	0.9	2.7
14	Anemias (D50-D64)	30	0.8	2.4
15	Aortic aneurysm and dissection (I71)	27	0.7	2.1
...	All other causes (residual)	592	15.0	46.6
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,668	100.0	380.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,130	24.2	92.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	15.0	57.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	466	10.0	38.0
4	Malignant neoplasms (C00-C97)	421	9.0	34.3
5	Cerebrovascular diseases (I60-I69)	192	4.1	15.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	189	4.0	15.4
7	Diabetes mellitus (E10-E14)	188	4.0	15.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	172	3.7	14.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	2.2	8.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.7	6.6
11	Septicemia (A40-A41)	58	1.2	4.7
12	Chronic lower respiratory diseases (J40-J47)	57	1.2	4.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.2	4.6
14	Aortic aneurysm and dissection (I71)	48	1.0	3.9
15	Influenza and pneumonia (J09-J18)	46	1.0	3.7
...	All other causes (residual)	759	16.3	61.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	6,686	100.0	539.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,700	25.4	137.0
2	Malignant neoplasms (C00-C97)	969	14.5	78.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	832	12.4	67.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	5.4	28.9
5	Diabetes mellitus (E10-E14)	315	4.7	25.4
6	Cerebrovascular diseases (I60-I69)	299	4.5	24.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	255	3.8	20.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.4	12.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	2.1	11.4
10	Septicemia (A40-A41)	123	1.8	9.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	116	1.7	9.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.7	9.0
13	Chronic lower respiratory diseases (J40-J47)	91	1.4	7.3
14	Influenza and pneumonia (J09-J18)	87	1.3	7.0
15	Aortic aneurysm and dissection (I71)	58	0.9	4.7
...	All other causes (residual)	1,069	16.0	86.2
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,126	100.0	861.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,960	26.6	229.3
2	Malignant neoplasms (C00-C97)	2,297	20.6	177.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,068	9.6	82.7
4	Diabetes mellitus (E10-E14)	543	4.9	42.1
5	Cerebrovascular diseases (I60-I69)	496	4.5	38.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	353	3.2	27.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	2.7	23.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	287	2.6	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.2	19.1
10	Chronic lower respiratory diseases (J40-J47)	195	1.8	15.1
11	Septicemia (A40-A41)	194	1.7	15.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.6	14.0
13	Influenza and pneumonia (J09-J18)	169	1.5	13.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.2	10.5
15	Viral hepatitis (B15-B19)	77	0.7	6.0
...	All other causes (residual)	1,623	14.6	125.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	16,180	100.0	1,347.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,335	26.8	361.0
2	Malignant neoplasms (C00-C97)	4,157	25.7	346.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,061	6.6	88.4
4	Diabetes mellitus (E10-E14)	772	4.8	64.3
5	Cerebrovascular diseases (I60-I69)	770	4.8	64.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.5	34.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	402	2.5	33.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	397	2.5	33.1
9	Chronic lower respiratory diseases (J40-J47)	377	2.3	31.4
10	Septicemia (A40-A41)	294	1.8	24.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	1.8	23.9
12	Influenza and pneumonia (J09-J18)	215	1.3	17.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	1.2	16.7
14	Viral hepatitis (B15-B19)	163	1.0	13.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	0.7	9.7
...	All other causes (residual)	2,223	13.7	185.1
Non-Hispanic, black, male, 60-64 years				
...	All causes	19,263	100.0	2,052.3
1	Malignant neoplasms (C00-C97)	5,489	28.5	584.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,116	26.6	545.1
3	Cerebrovascular diseases (I60-I69)	960	5.0	102.3
4	Diabetes mellitus (E10-E14)	952	4.9	101.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	926	4.8	98.7
6	Chronic lower respiratory diseases (J40-J47)	579	3.0	61.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	521	2.7	55.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	440	2.3	46.9
9	Septicemia (A40-A41)	365	1.9	38.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	321	1.7	34.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	310	1.6	33.0
12	Influenza and pneumonia (J09-J18)	289	1.5	30.8
13	Viral hepatitis (B15-B19)	242	1.3	25.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	0.7	14.1
15	Pneumonitis due to solids and liquids (J69)	75	0.4	8.0
...	All other causes (residual)	2,546	13.2	271.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	18,438	100.0	2,684.5
1	Malignant neoplasms (C00-C97)	5,487	29.8	798.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,833	26.2	703.7
3	Cerebrovascular diseases (I60-I69)	981	5.3	142.8
4	Diabetes mellitus (E10-E14)	958	5.2	139.5
5	Chronic lower respiratory diseases (J40-J47)	732	4.0	106.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	558	3.0	81.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	522	2.8	76.0
8	Septicemia (A40-A41)	406	2.2	59.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	313	1.7	45.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	292	1.6	42.5
11	Influenza and pneumonia (J09-J18)	276	1.5	40.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	166	0.9	24.2
13	Viral hepatitis (B15-B19)	145	0.8	21.1
14	Pneumonitis due to solids and liquids (J69)	88	0.5	12.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.4	11.8
...	All other causes (residual)	2,600	14.1	378.6
Non-Hispanic, black, male, 70-74 years				
...	All causes	16,429	100.0	3,780.6
1	Malignant neoplasms (C00-C97)	4,789	29.1	1,102.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,327	26.3	995.7
3	Cerebrovascular diseases (I60-I69)	976	5.9	224.6
4	Diabetes mellitus (E10-E14)	793	4.8	182.5
5	Chronic lower respiratory diseases (J40-J47)	776	4.7	178.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	541	3.3	124.5
7	Septicemia (A40-A41)	378	2.3	87.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	2.0	75.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	293	1.8	67.4
10	Influenza and pneumonia (J09-J18)	277	1.7	63.7
11	Alzheimer's disease (G30)	188	1.1	43.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.9	33.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.6	21.9
14	Parkinson's disease (G20-G21)	90	0.5	20.7
15	Pneumonitis due to solids and liquids (J69)	88	0.5	20.3
...	All other causes (residual)	2,342	14.3	538.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	15,455	100.0	5,424.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,107	26.6	1,441.4
2	Malignant neoplasms (C00-C97)	4,076	26.4	1,430.5
3	Cerebrovascular diseases (I60-I69)	1,013	6.6	355.5
4	Diabetes mellitus (E10-E14)	750	4.9	263.2
5	Chronic lower respiratory diseases (J40-J47)	742	4.8	260.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	543	3.5	190.6
7	Septicemia (A40-A41)	362	2.3	127.0
8	Alzheimer's disease (G30)	339	2.2	119.0
9	Influenza and pneumonia (J09-J18)	318	2.1	111.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	2.0	107.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	260	1.7	91.2
12	Parkinson's disease (G20-G21)	132	0.9	46.3
13	Pneumonitis due to solids and liquids (J69)	127	0.8	44.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	74	0.5	26.0
...	All other causes (residual)	2,226	14.4	781.2
Non-Hispanic, black, male, 80-84 years				
...	All causes	13,485	100.0	7,879.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,540	26.3	2,068.6
2	Malignant neoplasms (C00-C97)	3,060	22.7	1,788.1
3	Cerebrovascular diseases (I60-I69)	902	6.7	527.1
4	Chronic lower respiratory diseases (J40-J47)	649	4.8	379.2
5	Diabetes mellitus (E10-E14)	600	4.4	350.6
6	Alzheimer's disease (G30)	492	3.6	287.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	3.4	265.9
8	Influenza and pneumonia (J09-J18)	352	2.6	205.7
9	Septicemia (A40-A41)	302	2.2	176.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	276	2.0	161.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	1.7	133.2
12	Parkinson's disease (G20-G21)	169	1.3	98.8
13	Pneumonitis due to solids and liquids (J69)	123	0.9	71.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.7	54.9
15	Aortic aneurysm and dissection (I71)	39	0.3	22.8
...	All other causes (residual)	2,204	16.3	1,287.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,140	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,711	26.7	---
2	Malignant neoplasms (C00-C97)	2,039	20.1	---
3	Cerebrovascular diseases (I60-I69)	625	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	480	4.7	---
5	Alzheimer's disease (G30)	475	4.7	---
6	Diabetes mellitus (E10-E14)	407	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	366	3.6	---
8	Influenza and pneumonia (J09-J18)	251	2.5	---
9	Septicemia (A40-A41)	243	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	2.0	---
12	Parkinson's disease (G20-G21)	127	1.3	---
12	Pneumonitis due to solids and liquids (J69)	127	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.6	---
15	Nutritional deficiencies (E40-E64)	44	0.4	---
...	All other causes (residual)	1,774	17.5	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	5,884	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,634	27.8	---
2	Malignant neoplasms (C00-C97)	957	16.3	---
3	Cerebrovascular diseases (I60-I69)	373	6.3	---
4	Alzheimer's disease (G30)	353	6.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	218	3.7	---
7	Influenza and pneumonia (J09-J18)	193	3.3	---
8	Diabetes mellitus (E10-E14)	172	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	2.5	---
10	Septicemia (A40-A41)	134	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	1.4	---
12	Pneumonitis due to solids and liquids (J69)	78	1.3	---
13	Parkinson's disease (G20-G21)	46	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	---
15	Nutritional deficiencies (E40-E64)	25	0.4	---
...	All other causes (residual)	1,190	20.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,918	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	562	29.3	---
2	Malignant neoplasms (C00-C97)	232	12.1	---
3	Alzheimer's disease (G30)	150	7.8	---
4	Cerebrovascular diseases (I60-I69)	112	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	77	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	3.6	---
7	Influenza and pneumonia (J09-J18)	55	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	2.1	---
10	Diabetes mellitus (E10-E14)	39	2.0	---
11	Septicemia (A40-A41)	33	1.7	---
12	Pneumonitis due to solids and liquids (J69)	24	1.3	---
13	Parkinson's disease (G20-G21)	20	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	---
14	Nutritional deficiencies (E40-E64)	14	0.7	---
...	All other causes (residual)	434	22.6	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	383	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	29.5	---
2	Malignant neoplasms (C00-C97)	33	8.6	---
3	Alzheimer's disease (G30)	30	7.8	---
4	Cerebrovascular diseases (I60-I69)	21	5.5	---
5	Influenza and pneumonia (J09-J18)	16	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	14	3.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	3.1	---
9	Septicemia (A40-A41)	9	2.3	---
9	Diabetes mellitus (E10-E14)	9	2.3	---
11	Nutritional deficiencies (E40-E64)	5	1.3	---
11	Pneumonitis due to solids and liquids (J69)	5	1.3	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.0	---
14	Atherosclerosis (I70)	2	0.5	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.5	---
...	All other causes (residual)	93	24.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	153,404	100.0	704.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,455	23.1	162.8
2	Malignant neoplasms (C00-C97)	33,736	22.0	154.9
3	Cerebrovascular diseases (I60-I69)	9,798	6.4	45.0
4	Diabetes mellitus (E10-E14)	6,887	4.5	31.6
5	Alzheimer's disease (G30)	5,918	3.9	27.2
6	Chronic lower respiratory diseases (J40-J47)	5,091	3.3	23.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,850	3.2	22.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,732	3.1	21.7
9	Septicemia (A40-A41)	3,515	2.3	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,141	2.0	14.4
11	Influenza and pneumonia (J09-J18)	2,825	1.8	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,697	1.1	7.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,252	0.8	5.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,149	0.7	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,129	0.7	5.2
...	All other causes (residual)	32,229	21.0	148.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	408	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	12.0	4.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.0	3.4
4	Malignant neoplasms (C00-C97)	27	6.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.9	2.0
6	Influenza and pneumonia (J09-J18)	10	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	2.0	*
9	Septicemia (A40-A41)	7	1.7	*
10	Anemias (D50-D64)	6	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Cerebrovascular diseases (I60-I69)	5	1.2	*
13	Meningitis (G00,G03)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	101	24.8	8.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	236	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	33.5	5.1
2	Malignant neoplasms (C00-C97)	27	11.4	1.8
3	Chronic lower respiratory diseases (J40-J47)	19	8.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	7.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	6.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	5.5	*
7	Anemias (D50-D64)	7	3.0	*
7	Influenza and pneumonia (J09-J18)	7	3.0	*
9	Septicemia (A40-A41)	6	2.5	*
10	Cerebrovascular diseases (I60-I69)	5	2.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
14	Pneumonitis due to solids and liquids (J69)	1	0.4	*
14	Diseases of appendix (K35-K38)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	30	12.7	2.0
Non-Hispanic, black, female, 10-14 years				
...	All causes	248	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	24.6	4.0
2	Malignant neoplasms (C00-C97)	31	12.5	2.1
3	Chronic lower respiratory diseases (J40-J47)	20	8.1	1.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	8.1	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	5.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.2	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.8	*
8	Cerebrovascular diseases (I60-I69)	4	1.6	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.6	*
10	Septicemia (A40-A41)	3	1.2	*
10	Influenza and pneumonia (J09-J18)	3	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Anemias (D50-D64)	2	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
15	Meningococcal infection (A39)	1	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Hernia (K40-K46)	1	0.4	*
...	All other causes (residual)	53	21.4	3.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	541	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	27.9	9.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	18.1	6.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	8.9	3.0
4	Malignant neoplasms (C00-C97)	41	7.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.4	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	17	3.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	3.1	*
8	Anemias (D50-D64)	16	3.0	*
9	Chronic lower respiratory diseases (J40-J47)	12	2.2	*
10	Septicemia (A40-A41)	7	1.3	*
11	Diabetes mellitus (E10-E14)	6	1.1	*
12	Influenza and pneumonia (J09-J18)	5	0.9	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	83	15.3	5.3
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,069	100.0	60.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	25.4	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	16.0	9.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	7.4	4.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	6.9	4.2
5	Malignant neoplasms (C00-C97)	73	6.8	4.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	46	4.3	2.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	27	2.5	1.5
8	Anemias (D50-D64)	19	1.8	*
8	Diabetes mellitus (E10-E14)	19	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	19	1.8	*
11	Cerebrovascular diseases (I60-I69)	15	1.4	*
12	Septicemia (A40-A41)	12	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	Influenza and pneumonia (J09-J18)	9	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
...	All other causes (residual)	217	20.3	12.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,326	100.0	80.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	21.8	17.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	12.4	10.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	10.4	8.4
4	Malignant neoplasms (C00-C97)	113	8.5	6.9
5	Pregnancy, childbirth and the puerperium (O00-O99)	64	4.8	3.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	4.2	3.4
7	Diabetes mellitus (E10-E14)	46	3.5	2.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	2.6	2.1
9	Cerebrovascular diseases (I60-I69)	24	1.8	1.5
10	Anemias (D50-D64)	23	1.7	1.4
11	Septicemia (A40-A41)	21	1.6	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.1	*
...	All other causes (residual)	289	21.8	17.6
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,761	100.0	118.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	14.9	17.7
2	Malignant neoplasms (C00-C97)	250	14.2	16.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	13.9	16.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	126	7.2	8.5
5	Diabetes mellitus (E10-E14)	75	4.3	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	71	4.0	4.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	69	3.9	4.6
8	Cerebrovascular diseases (I60-I69)	51	2.9	3.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	2.4	2.8
10	Anemias (D50-D64)	37	2.1	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.1	2.5
12	Septicemia (A40-A41)	34	1.9	2.3
13	Influenza and pneumonia (J09-J18)	21	1.2	1.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.1	*
...	All other causes (residual)	403	22.9	27.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,331	100.0	164.9
1	Malignant neoplasms (C00-C97)	451	19.3	31.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	15.9	26.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	12.2	20.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	116	5.0	8.2
5	Diabetes mellitus (E10-E14)	88	3.8	6.2
5	Cerebrovascular diseases (I60-I69)	88	3.8	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	3.8	6.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	66	2.8	4.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.1	3.5
10	Chronic lower respiratory diseases (J40-J47)	40	1.7	2.8
11	Septicemia (A40-A41)	38	1.6	2.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	1.6	2.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.2	2.1
14	Anemias (D50-D64)	27	1.2	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.2	1.9
...	All other causes (residual)	532	22.8	37.6
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,361	100.0	242.4
1	Malignant neoplasms (C00-C97)	760	22.6	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	649	19.3	46.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	9.0	21.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	144	4.3	10.4
5	Cerebrovascular diseases (I60-I69)	137	4.1	9.9
6	Diabetes mellitus (E10-E14)	135	4.0	9.7
7	Septicemia (A40-A41)	85	2.5	6.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	2.5	6.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	2.5	6.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	76	2.3	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.5	3.8
12	Chronic lower respiratory diseases (J40-J47)	46	1.4	3.3
13	Influenza and pneumonia (J09-J18)	42	1.2	3.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.2	2.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.0	2.5
...	All other causes (residual)	695	20.7	50.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,088	100.0	364.5
1	Malignant neoplasms (C00-C97)	1,250	24.6	89.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,054	20.7	75.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	7.2	26.1
4	Diabetes mellitus (E10-E14)	248	4.9	17.8
4	Cerebrovascular diseases (I60-I69)	248	4.9	17.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	170	3.3	12.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	127	2.5	9.1
8	Septicemia (A40-A41)	117	2.3	8.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.1	7.7
10	Chronic lower respiratory diseases (J40-J47)	96	1.9	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.4	5.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	1.4	5.1
13	Influenza and pneumonia (J09-J18)	50	1.0	3.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	0.9	3.3
15	Anemias (D50-D64)	39	0.8	2.8
...	All other causes (residual)	1,029	20.2	73.7
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,272	100.0	563.6
1	Malignant neoplasms (C00-C97)	2,521	30.5	171.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,703	20.6	116.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	5.5	30.9
4	Cerebrovascular diseases (I60-I69)	391	4.7	26.6
5	Diabetes mellitus (E10-E14)	380	4.6	25.9
6	Chronic lower respiratory diseases (J40-J47)	223	2.7	15.2
7	Septicemia (A40-A41)	200	2.4	13.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	199	2.4	13.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	176	2.1	12.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.1	11.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	139	1.7	9.5
12	Influenza and pneumonia (J09-J18)	115	1.4	7.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.7	4.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	0.5	2.9
15	Viral hepatitis (B15-B19)	42	0.5	2.9
...	All other causes (residual)	1,455	17.6	99.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	11,835	100.0	847.8
1	Malignant neoplasms (C00-C97)	3,890	32.9	278.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,562	21.6	183.5
3	Diabetes mellitus (E10-E14)	604	5.1	43.3
4	Cerebrovascular diseases (I60-I69)	546	4.6	39.1
5	Chronic lower respiratory diseases (J40-J47)	459	3.9	32.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	3.3	28.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	2.7	23.1
8	Septicemia (A40-A41)	309	2.6	22.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	235	2.0	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.6	13.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	163	1.4	11.7
12	Influenza and pneumonia (J09-J18)	151	1.3	10.8
13	Viral hepatitis (B15-B19)	83	0.7	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	0.4	3.6
15	Pneumonitis due to solids and liquids (J69)	47	0.4	3.4
...	All other causes (residual)	1,837	15.5	131.6
Non-Hispanic, black, female, 60-64 years				
...	All causes	13,608	100.0	1,179.6
1	Malignant neoplasms (C00-C97)	4,434	32.6	384.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,055	22.5	264.8
3	Diabetes mellitus (E10-E14)	730	5.4	63.3
4	Cerebrovascular diseases (I60-I69)	722	5.3	62.6
5	Chronic lower respiratory diseases (J40-J47)	506	3.7	43.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	3.0	35.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	2.4	28.5
8	Septicemia (A40-A41)	323	2.4	28.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	2.0	23.7
10	Influenza and pneumonia (J09-J18)	205	1.5	17.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	196	1.4	17.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	108	0.8	9.4
13	Viral hepatitis (B15-B19)	91	0.7	7.9
14	Pneumonitis due to solids and liquids (J69)	53	0.4	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.4	4.5
...	All other causes (residual)	2,119	15.6	183.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	14,103	100.0	1,592.8
1	Malignant neoplasms (C00-C97)	4,536	32.2	512.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,145	22.3	355.2
3	Cerebrovascular diseases (I60-I69)	777	5.5	87.8
4	Diabetes mellitus (E10-E14)	744	5.3	84.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	550	3.9	62.1
6	Chronic lower respiratory diseases (J40-J47)	549	3.9	62.0
7	Septicemia (A40-A41)	356	2.5	40.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	275	1.9	31.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	1.9	29.5
10	Influenza and pneumonia (J09-J18)	258	1.8	29.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	126	0.9	14.2
12	Alzheimer's disease (G30)	91	0.6	10.3
13	Pneumonitis due to solids and liquids (J69)	77	0.5	8.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	74	0.5	8.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	7.8
...	All other causes (residual)	2,215	15.7	250.2
Non-Hispanic, black, female, 70-74 years				
...	All causes	14,348	100.0	2,377.6
1	Malignant neoplasms (C00-C97)	4,116	28.7	682.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,466	24.2	574.3
3	Cerebrovascular diseases (I60-I69)	898	6.3	148.8
4	Diabetes mellitus (E10-E14)	803	5.6	133.1
5	Chronic lower respiratory diseases (J40-J47)	645	4.5	106.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	499	3.5	82.7
7	Septicemia (A40-A41)	354	2.5	58.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	291	2.0	48.2
9	Alzheimer's disease (G30)	279	1.9	46.2
10	Influenza and pneumonia (J09-J18)	258	1.8	42.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	1.3	31.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.6	14.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	12.3
14	Pneumonitis due to solids and liquids (J69)	71	0.5	11.8
15	Parkinson's disease (G20-G21)	52	0.4	8.6
...	All other causes (residual)	2,264	15.8	375.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	15,521	100.0	3,580.2
1	Malignant neoplasms (C00-C97)	3,799	24.5	876.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,692	23.8	851.6
3	Cerebrovascular diseases (I60-I69)	1,152	7.4	265.7
4	Diabetes mellitus (E10-E14)	865	5.6	199.5
5	Chronic lower respiratory diseases (J40-J47)	648	4.2	149.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	602	3.9	138.9
7	Alzheimer's disease (G30)	570	3.7	131.5
8	Septicemia (A40-A41)	404	2.6	93.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	2.2	78.9
10	Influenza and pneumonia (J09-J18)	287	1.8	66.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	1.2	44.5
12	Pneumonitis due to solids and liquids (J69)	95	0.6	21.9
13	Parkinson's disease (G20-G21)	91	0.6	21.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.6	20.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.4	13.4
...	All other causes (residual)	2,636	17.0	608.0
Non-Hispanic, black, female, 80-84 years				
...	All causes	17,000	100.0	5,607.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,223	24.8	1,393.0
2	Malignant neoplasms (C00-C97)	3,178	18.7	1,048.3
3	Cerebrovascular diseases (I60-I69)	1,452	8.5	478.9
4	Alzheimer's disease (G30)	1,034	6.1	341.1
5	Diabetes mellitus (E10-E14)	809	4.8	266.9
6	Chronic lower respiratory diseases (J40-J47)	690	4.1	227.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	621	3.7	204.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	2.5	139.9
9	Septicemia (A40-A41)	417	2.5	137.5
10	Influenza and pneumonia (J09-J18)	360	2.1	118.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	1.3	73.9
12	Pneumonitis due to solids and liquids (J69)	112	0.7	36.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.6	36.3
14	Parkinson's disease (G20-G21)	103	0.6	34.0
15	Aortic aneurysm and dissection (I71)	59	0.3	19.5
...	All other causes (residual)	3,184	18.7	1,050.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	17,316	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,684	27.1	---
2	Malignant neoplasms (C00-C97)	2,428	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,499	8.7	---
4	Alzheimer's disease (G30)	1,439	8.3	---
5	Diabetes mellitus (E10-E14)	715	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	576	3.3	---
7	Chronic lower respiratory diseases (J40-J47)	571	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	393	2.3	---
9	Septicemia (A40-A41)	392	2.3	---
10	Influenza and pneumonia (J09-J18)	378	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	1.2	---
12	Pneumonitis due to solids and liquids (J69)	118	0.7	---
13	Parkinson's disease (G20-G21)	113	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	---
15	Atherosclerosis (I70)	60	0.3	---
...	All other causes (residual)	3,644	21.0	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	13,305	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,761	28.3	---
2	Alzheimer's disease (G30)	1,414	10.6	---
3	Malignant neoplasms (C00-C97)	1,262	9.5	---
4	Cerebrovascular diseases (I60-I69)	1,100	8.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	3.1	---
6	Diabetes mellitus (E10-E14)	414	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	380	2.9	---
8	Influenza and pneumonia (J09-J18)	373	2.8	---
9	Chronic lower respiratory diseases (J40-J47)	334	2.5	---
10	Septicemia (A40-A41)	282	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.1	---
12	Pneumonitis due to solids and liquids (J69)	114	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	---
14	Parkinson's disease (G20-G21)	65	0.5	---
15	Nutritional deficiencies (E40-E64)	59	0.4	---
...	All other causes (residual)	3,109	23.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	6,415	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,857	28.9	---
2	Alzheimer's disease (G30)	740	11.5	---
3	Cerebrovascular diseases (I60-I69)	512	8.0	---
4	Malignant neoplasms (C00-C97)	450	7.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	2.9	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	2.7	---
7	Influenza and pneumonia (J09-J18)	164	2.6	---
8	Diabetes mellitus (E10-E14)	162	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	131	2.0	---
10	Septicemia (A40-A41)	88	1.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.2	---
12	Pneumonitis due to solids and liquids (J69)	58	0.9	---
13	Nutritional deficiencies (E40-E64)	40	0.6	---
14	Atherosclerosis (I70)	32	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	---
...	All other causes (residual)	1,716	26.7	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,215	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	649	29.3	---
2	Alzheimer's disease (G30)	284	12.8	---
3	Cerebrovascular diseases (I60-I69)	155	7.0	---
4	Malignant neoplasms (C00-C97)	95	4.3	---
5	Influenza and pneumonia (J09-J18)	79	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.8	---
8	Diabetes mellitus (E10-E14)	39	1.8	---
9	Septicemia (A40-A41)	35	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.1	---
12	Pneumonitis due to solids and liquids (J69)	23	1.0	---
13	Nutritional deficiencies (E40-E64)	21	0.9	---
14	Atherosclerosis (I70)	12	0.5	---
15	Anemias (D50-D64)	8	0.4	---
...	All other causes (residual)	616	27.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015