

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	141,576	100.0	292.4
1	Malignant neoplasms (C00-C97)	29,935	21.1	61.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,611	20.9	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,654	7.5	22.0
4	Cerebrovascular diseases (I60-I69)	7,065	5.0	14.6
5	Diabetes mellitus (E10-E14)	6,311	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,303	3.0	8.9
7	Chronic lower respiratory diseases (J40-J47)	4,026	2.8	8.3
8	Influenza and pneumonia (J09-J18)	3,679	2.6	7.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,179	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,107	2.2	6.4
11	Alzheimer's disease (G30)	2,986	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,618	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,573	1.8	5.3
14	Septicemia (A40-A41)	2,077	1.5	4.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,851	1.3	3.8
...	All other causes (Residual)	27,601	19.5	57.0
Hispanic, both sexes, 1-4 years				
...	All causes	1,004	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	32.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.4	2.8
3	Malignant neoplasms (C00-C97)	99	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	7.5	1.7
5	Influenza and pneumonia (J09-J18)	34	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.6	0.6
7	Septicemia (A40-A41)	17	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
9	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	1.0	*
12	Meningitis (G00,G03)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
...	All other causes (Residual)	247	24.6	5.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,172	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	28.3	3.8
2	Malignant neoplasms (C00-C97)	216	18.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	6.1	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	5.9	0.8
5	Influenza and pneumonia (J09-J18)	63	5.4	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	2.0	0.3
9	Chronic lower respiratory diseases (J40-J47)	19	1.6	*
10	Septicemia (A40-A41)	15	1.3	*
11	Cerebrovascular diseases (I60-I69)	10	0.9	*
12	Diabetes mellitus (E10-E14)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Diseases of appendix (K35-K38)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (Residual)	233	19.9	2.6
Hispanic, both sexes, 15-24 years				
...	All causes	5,187	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,999	38.5	25.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,168	22.5	14.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	583	11.2	7.4
4	Malignant neoplasms (C00-C97)	320	6.2	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	2.9	1.9
6	Influenza and pneumonia (J09-J18)	88	1.7	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.4	0.9
8	Cerebrovascular diseases (I60-I69)	42	0.8	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	41	0.8	0.5
10	Septicemia (A40-A41)	36	0.7	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.6	0.4
12	Diabetes mellitus (E10-E14)	24	0.5	0.3
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	584	11.3	7.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,369	100.0	77.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,054	32.2	25.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	845	13.3	10.3
3	Malignant neoplasms (C00-C97)	628	9.9	7.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	598	9.4	7.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	6.0	4.7
6	Influenza and pneumonia (J09-J18)	178	2.8	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	124	1.9	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.9	1.5
9	Cerebrovascular diseases (I60-I69)	117	1.8	1.4
10	Diabetes mellitus (E10-E14)	90	1.4	1.1
11	Pregnancy, childbirth and the puerperium (O00-O99)	84	1.3	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	1.0	0.8
13	Septicemia (A40-A41)	59	0.9	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.7	0.5
15	Chronic lower respiratory diseases (J40-J47)	40	0.6	0.5
...	All other causes (Residual)	936	14.7	11.4
Hispanic, both sexes, 35-44 years				
...	All causes	8,640	100.0	122.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,756	20.3	24.9
2	Malignant neoplasms (C00-C97)	1,513	17.5	21.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	924	10.7	13.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	516	6.0	7.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	506	5.9	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	482	5.6	6.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	360	4.2	5.1
8	Cerebrovascular diseases (I60-I69)	285	3.3	4.0
9	Influenza and pneumonia (J09-J18)	263	3.0	3.7
10	Diabetes mellitus (E10-E14)	215	2.5	3.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	1.4	1.7
12	Septicemia (A40-A41)	95	1.1	1.3
13	Viral hepatitis (B15-B19)	87	1.0	1.2
14	Chronic lower respiratory diseases (J40-J47)	67	0.8	0.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.5	0.7
...	All other causes (Residual)	1,404	16.3	19.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	15,226	100.0	294.6
1	Malignant neoplasms (C00-C97)	3,732	24.5	72.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,500	16.4	48.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,560	10.2	30.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,191	7.8	23.0
5	Cerebrovascular diseases (I60-I69)	628	4.1	12.2
6	Diabetes mellitus (E10-E14)	612	4.0	11.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	451	3.0	8.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	403	2.6	7.8
9	Viral hepatitis (B15-B19)	379	2.5	7.3
10	Influenza and pneumonia (J09-J18)	368	2.4	7.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	265	1.7	5.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	253	1.7	4.9
13	Septicemia (A40-A41)	228	1.5	4.4
14	Chronic lower respiratory diseases (J40-J47)	165	1.1	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	0.7	2.1
...	All other causes (Residual)	2,383	15.7	46.1
Hispanic, both sexes, 55-64 years				
...	All causes	19,251	100.0	642.9
1	Malignant neoplasms (C00-C97)	5,972	31.0	199.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,036	21.0	134.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,169	6.1	39.0
4	Diabetes mellitus (E10-E14)	1,168	6.1	39.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	884	4.6	29.5
6	Cerebrovascular diseases (I60-I69)	858	4.5	28.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	2.4	15.2
8	Viral hepatitis (B15-B19)	391	2.0	13.1
9	Septicemia (A40-A41)	344	1.8	11.5
10	Chronic lower respiratory diseases (J40-J47)	324	1.7	10.8
11	Influenza and pneumonia (J09-J18)	315	1.6	10.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	217	1.1	7.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	208	1.1	6.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.0	6.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	0.5	3.3
...	All other causes (Residual)	2,619	13.6	87.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	22,379	100.0	1,428.6
1	Malignant neoplasms (C00-C97)	6,951	31.1	443.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,168	23.1	329.9
3	Diabetes mellitus (E10-E14)	1,464	6.5	93.5
4	Cerebrovascular diseases (I60-I69)	1,168	5.2	74.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	736	3.3	47.0
6	Chronic lower respiratory diseases (J40-J47)	693	3.1	44.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	666	3.0	42.5
8	Influenza and pneumonia (J09-J18)	513	2.3	32.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	2.3	32.2
10	Septicemia (A40-A41)	384	1.7	24.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	284	1.3	18.1
12	Alzheimer's disease (G30)	200	0.9	12.8
13	Viral hepatitis (B15-B19)	154	0.7	9.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.6	8.0
15	Parkinson's disease (G20-G21)	115	0.5	7.3
...	All other causes (Residual)	3,253	14.5	207.7
Hispanic, both sexes, 75-84 years				
...	All causes	30,498	100.0	3,504.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,025	26.3	922.0
2	Malignant neoplasms (C00-C97)	7,110	23.3	816.9
3	Cerebrovascular diseases (I60-I69)	2,012	6.6	231.2
4	Diabetes mellitus (E10-E14)	1,746	5.7	200.6
5	Chronic lower respiratory diseases (J40-J47)	1,423	4.7	163.5
6	Alzheimer's disease (G30)	975	3.2	112.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	881	2.9	101.2
8	Influenza and pneumonia (J09-J18)	847	2.8	97.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	570	1.9	65.5
10	Septicemia (A40-A41)	498	1.6	57.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	455	1.5	52.3
12	Parkinson's disease (G20-G21)	428	1.4	49.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	391	1.3	44.9
14	Pneumonitis due to solids and liquids (J69)	192	0.6	22.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	177	0.6	20.3
...	All other causes (Residual)	4,768	15.6	547.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	26,397	100.0	8,145.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,262	31.3	2,549.4
2	Malignant neoplasms (C00-C97)	3,377	12.8	1,042.0
3	Cerebrovascular diseases (I60-I69)	1,901	7.2	586.6
4	Alzheimer's disease (G30)	1,764	6.7	544.3
5	Chronic lower respiratory diseases (J40-J47)	1,254	4.8	386.9
6	Diabetes mellitus (E10-E14)	985	3.7	303.9
7	Influenza and pneumonia (J09-J18)	956	3.6	295.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	643	2.4	198.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	488	1.8	150.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	1.8	146.3
11	Septicemia (A40-A41)	351	1.3	108.3
12	Parkinson's disease (G20-G21)	280	1.1	86.4
13	Pneumonitis due to solids and liquids (J69)	262	1.0	80.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.6	51.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.5	42.3
...	All other causes (Residual)	5,095	19.3	1,572.2
Hispanic, male, all ages				
...	All causes	78,157	100.0	311.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,301	20.9	65.1
2	Malignant neoplasms (C00-C97)	15,772	20.2	62.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,841	10.0	31.3
4	Cerebrovascular diseases (I60-I69)	3,288	4.2	13.1
5	Diabetes mellitus (E10-E14)	3,212	4.1	12.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,969	3.8	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	3.4	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,135	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,024	2.6	8.1
10	Influenza and pneumonia (J09-J18)	1,882	2.4	7.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,593	2.0	6.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,462	1.9	5.8
13	Septicemia (A40-A41)	1,041	1.3	4.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	985	1.3	3.9
15	Alzheimer's disease (G30)	973	1.2	3.9
...	All other causes (Residual)	14,044	18.0	56.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	539	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	38.6	9.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.8	2.6
3	Malignant neoplasms (C00-C97)	44	8.2	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.9	1.4
5	Influenza and pneumonia (J09-J18)	16	3.0	*
6	Septicemia (A40-A41)	12	2.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
12	Meningococcal infection (A39)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
12	Hernia (K40-K46)	1	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	131	24.3	5.9
Hispanic, male, 5-14 years				
...	All causes	650	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	29.2	4.2
2	Malignant neoplasms (C00-C97)	118	18.2	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	6.0	0.9
5	Influenza and pneumonia (J09-J18)	33	5.1	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.9	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.1	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
10	Cerebrovascular diseases (I60-I69)	6	0.9	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
12	Septicemia (A40-A41)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Anemias (D50-D64)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	122	18.8	2.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,992	100.0	96.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,590	39.8	38.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,044	26.2	25.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	488	12.2	11.7
4	Malignant neoplasms (C00-C97)	204	5.1	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.5	2.4
6	Influenza and pneumonia (J09-J18)	40	1.0	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	25	0.6	0.6
9	Septicemia (A40-A41)	21	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.4	*
11	Legal intervention (Y35,Y89.0)	15	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.2	*
13	Diabetes mellitus (E10-E14)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
...	All other causes (Residual)	371	9.3	8.9
Hispanic, male, 25-34 years				
...	All causes	4,688	100.0	104.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,674	35.7	37.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	721	15.4	16.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	501	10.7	11.1
4	Malignant neoplasms (C00-C97)	336	7.2	7.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	5.8	6.0
6	Influenza and pneumonia (J09-J18)	112	2.4	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	107	2.3	2.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	99	2.1	2.2
9	Cerebrovascular diseases (I60-I69)	67	1.4	1.5
10	Diabetes mellitus (E10-E14)	55	1.2	1.2
11	Septicemia (A40-A41)	40	0.9	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	0.9
13	Legal intervention (Y35,Y89.0)	30	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	22	0.5	0.5
...	All other causes (Residual)	583	12.4	12.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,811	100.0	153.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,396	24.0	36.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	711	12.2	18.8
3	Malignant neoplasms (C00-C97)	660	11.4	17.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	7.3	11.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	416	7.2	11.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	391	6.7	10.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	278	4.8	7.4
8	Cerebrovascular diseases (I60-I69)	170	2.9	4.5
8	Influenza and pneumonia (J09-J18)	170	2.9	4.5
10	Diabetes mellitus (E10-E14)	131	2.3	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.3	2.0
12	Viral hepatitis (B15-B19)	57	1.0	1.5
13	Septicemia (A40-A41)	55	0.9	1.5
14	Chronic lower respiratory diseases (J40-J47)	31	0.5	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	0.5	0.7
...	All other causes (Residual)	821	14.1	21.7
Hispanic, male, 45-54 years				
...	All causes	9,955	100.0	375.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,874	18.8	70.7
2	Malignant neoplasms (C00-C97)	1,867	18.8	70.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,166	11.7	44.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	931	9.4	35.1
5	Diabetes mellitus (E10-E14)	392	3.9	14.8
6	Cerebrovascular diseases (I60-I69)	387	3.9	14.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	362	3.6	13.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	335	3.4	12.6
9	Viral hepatitis (B15-B19)	275	2.8	10.4
10	Influenza and pneumonia (J09-J18)	220	2.2	8.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	2.1	7.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	1.6	6.0
13	Septicemia (A40-A41)	147	1.5	5.5
14	Chronic lower respiratory diseases (J40-J47)	83	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	0.7	2.5
...	All other causes (Residual)	1,485	14.9	56.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	11,941	100.0	822.5
1	Malignant neoplasms (C00-C97)	3,251	27.2	223.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,851	23.9	196.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	846	7.1	58.3
4	Diabetes mellitus (E10-E14)	698	5.8	48.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	681	5.7	46.9
6	Cerebrovascular diseases (I60-I69)	492	4.1	33.9
7	Viral hepatitis (B15-B19)	267	2.2	18.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	2.2	18.3
9	Influenza and pneumonia (J09-J18)	185	1.5	12.7
10	Septicemia (A40-A41)	182	1.5	12.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	1.5	12.5
12	Chronic lower respiratory diseases (J40-J47)	177	1.5	12.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	154	1.3	10.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	0.9	7.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	0.6	5.1
...	All other causes (Residual)	1,526	12.8	105.1
Hispanic, male, 65-74 years				
...	All causes	12,623	100.0	1,783.0
1	Malignant neoplasms (C00-C97)	3,851	30.5	543.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,189	25.3	450.4
3	Diabetes mellitus (E10-E14)	799	6.3	112.9
4	Cerebrovascular diseases (I60-I69)	653	5.2	92.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	456	3.6	64.4
6	Chronic lower respiratory diseases (J40-J47)	392	3.1	55.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	2.7	48.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	310	2.5	43.8
9	Influenza and pneumonia (J09-J18)	270	2.1	38.1
10	Septicemia (A40-A41)	192	1.5	27.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	164	1.3	23.2
12	Viral hepatitis (B15-B19)	86	0.7	12.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.7	12.1
14	Alzheimer's disease (G30)	84	0.7	11.9
15	Parkinson's disease (G20-G21)	77	0.6	10.9
...	All other causes (Residual)	1,671	13.2	236.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	14,911	100.0	4,109.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,110	27.6	1,132.8
2	Malignant neoplasms (C00-C97)	3,828	25.7	1,055.1
3	Cerebrovascular diseases (I60-I69)	858	5.8	236.5
4	Diabetes mellitus (E10-E14)	758	5.1	208.9
5	Chronic lower respiratory diseases (J40-J47)	730	4.9	201.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	421	2.8	116.0
7	Influenza and pneumonia (J09-J18)	404	2.7	111.4
8	Alzheimer's disease (G30)	382	2.6	105.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	2.0	82.1
10	Parkinson's disease (G20-G21)	250	1.7	68.9
11	Septicemia (A40-A41)	223	1.5	61.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	188	1.3	51.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	165	1.1	45.5
14	Pneumonitis due to solids and liquids (J69)	99	0.7	27.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	90	0.6	24.8
...	All other causes (Residual)	2,107	14.1	580.7
Hispanic, male, 85 years and over				
...	All causes	10,039	100.0	8,768.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,118	31.1	2,723.5
2	Malignant neoplasms (C00-C97)	1,605	16.0	1,401.9
3	Cerebrovascular diseases (I60-I69)	607	6.0	530.2
4	Chronic lower respiratory diseases (J40-J47)	555	5.5	484.8
5	Alzheimer's disease (G30)	481	4.8	420.1
6	Influenza and pneumonia (J09-J18)	395	3.9	345.0
7	Diabetes mellitus (E10-E14)	366	3.6	319.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	280	2.8	244.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	2.1	182.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	1.8	159.0
11	Parkinson's disease (G20-G21)	142	1.4	124.0
12	Septicemia (A40-A41)	138	1.4	120.5
13	Pneumonitis due to solids and liquids (J69)	121	1.2	105.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.8	67.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.5	46.3
...	All other causes (Residual)	1,710	17.0	1,493.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	63,419	100.0	271.5
1	Malignant neoplasms (C00-C97)	14,163	22.3	60.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,310	21.0	57.0
3	Cerebrovascular diseases (I60-I69)	3,777	6.0	16.2
4	Diabetes mellitus (E10-E14)	3,099	4.9	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,813	4.4	12.0
6	Alzheimer's disease (G30)	2,013	3.2	8.6
7	Chronic lower respiratory diseases (J40-J47)	2,002	3.2	8.6
8	Influenza and pneumonia (J09-J18)	1,797	2.8	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,514	2.4	6.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,334	2.1	5.7
11	Certain conditions originating in the perinatal period (P00-P96)	1,156	1.8	4.9
12	Septicemia (A40-A41)	1,036	1.6	4.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	882	1.4	3.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	791	1.2	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	0.9	2.3
...	All other causes (Residual)	13,188	20.8	56.4
Hispanic, female, 1-4 years				
...	All causes	465	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	24.9	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	14.2	3.1
3	Malignant neoplasms (C00-C97)	55	11.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	9.2	2.0
5	Influenza and pneumonia (J09-J18)	18	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.3	*
11	Septicemia (A40-A41)	5	1.1	*
12	Meningitis (G00,G03)	3	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	111	23.9	5.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	522	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	27.2	3.3
2	Malignant neoplasms (C00-C97)	98	18.8	2.3
3	Influenza and pneumonia (J09-J18)	30	5.7	0.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.2	0.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.1	*
8	Septicemia (A40-A41)	12	2.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.9	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Anemias (D50-D64)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Diseases of appendix (K35-K38)	3	0.6	*
...	All other causes (Residual)	115	22.0	2.7
Hispanic, female, 15-24 years				
...	All causes	1,195	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	34.2	10.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	10.4	3.3
3	Malignant neoplasms (C00-C97)	116	9.7	3.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	7.9	2.5
5	Influenza and pneumonia (J09-J18)	48	4.0	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	1.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	41	3.4	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.6	0.8
9	Cerebrovascular diseases (I60-I69)	17	1.4	*
10	Septicemia (A40-A41)	15	1.3	*
10	Diabetes mellitus (E10-E14)	15	1.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
...	All other causes (Residual)	198	16.6	5.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,681	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	22.6	10.3
2	Malignant neoplasms (C00-C97)	292	17.4	7.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	7.4	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	6.6	3.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	5.8	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	84	5.0	2.3
7	Influenza and pneumonia (J09-J18)	66	3.9	1.8
8	Cerebrovascular diseases (I60-I69)	50	3.0	1.4
9	Diabetes mellitus (E10-E14)	35	2.1	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	1.5	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.5	0.7
12	Septicemia (A40-A41)	19	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
...	All other causes (Residual)	323	19.2	8.8
Hispanic, female, 35-44 years				
...	All causes	2,829	100.0	86.2
1	Malignant neoplasms (C00-C97)	853	30.2	26.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	12.7	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	213	7.5	6.5
4	Cerebrovascular diseases (I60-I69)	115	4.1	3.5
5	Influenza and pneumonia (J09-J18)	93	3.3	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	3.3	2.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	91	3.2	2.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	3.2	2.7
9	Diabetes mellitus (E10-E14)	84	3.0	2.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	82	2.9	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.6	1.4
12	Pregnancy, childbirth and the puerperium (O00-O99)	44	1.6	1.3
13	Septicemia (A40-A41)	40	1.4	1.2
14	Chronic lower respiratory diseases (J40-J47)	36	1.3	1.1
15	Viral hepatitis (B15-B19)	30	1.1	0.9
...	All other causes (Residual)	559	19.8	17.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,271	100.0	209.3
1	Malignant neoplasms (C00-C97)	1,865	35.4	74.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	626	11.9	24.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	7.5	15.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	260	4.9	10.3
5	Cerebrovascular diseases (I60-I69)	241	4.6	9.6
6	Diabetes mellitus (E10-E14)	220	4.2	8.7
7	Influenza and pneumonia (J09-J18)	148	2.8	5.9
8	Viral hepatitis (B15-B19)	104	2.0	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	1.8	3.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	89	1.7	3.5
11	Chronic lower respiratory diseases (J40-J47)	82	1.6	3.3
12	Septicemia (A40-A41)	81	1.5	3.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	1.3	2.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	1.1	2.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	0.8	1.7
...	All other causes (Residual)	898	17.0	35.7
Hispanic, female, 55-64 years				
...	All causes	7,310	100.0	473.9
1	Malignant neoplasms (C00-C97)	2,721	37.2	176.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	16.2	76.8
3	Diabetes mellitus (E10-E14)	470	6.4	30.5
4	Cerebrovascular diseases (I60-I69)	366	5.0	23.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	323	4.4	20.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	2.8	13.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	2.6	12.3
8	Septicemia (A40-A41)	162	2.2	10.5
9	Chronic lower respiratory diseases (J40-J47)	147	2.0	9.5
10	Influenza and pneumonia (J09-J18)	130	1.8	8.4
11	Viral hepatitis (B15-B19)	124	1.7	8.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	1.1	5.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.7	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.6	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.5	2.3
...	All other causes (Residual)	1,077	14.7	69.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,756	100.0	1,136.3
1	Malignant neoplasms (C00-C97)	3,100	31.8	361.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,979	20.3	230.5
3	Diabetes mellitus (E10-E14)	665	6.8	77.5
4	Cerebrovascular diseases (I60-I69)	515	5.3	60.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	3.3	37.6
6	Chronic lower respiratory diseases (J40-J47)	301	3.1	35.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	280	2.9	32.6
8	Influenza and pneumonia (J09-J18)	243	2.5	28.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	2.0	22.7
10	Septicemia (A40-A41)	192	2.0	22.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.2	14.0
12	Alzheimer's disease (G30)	116	1.2	13.5
13	Viral hepatitis (B15-B19)	68	0.7	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.6	6.9
15	Parkinson's disease (G20-G21)	38	0.4	4.4
...	All other causes (Residual)	1,562	16.0	181.9
Hispanic, female, 75-84 years				
...	All causes	15,587	100.0	3,071.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,915	25.1	771.3
2	Malignant neoplasms (C00-C97)	3,282	21.1	646.6
3	Cerebrovascular diseases (I60-I69)	1,154	7.4	227.4
4	Diabetes mellitus (E10-E14)	988	6.3	194.7
5	Chronic lower respiratory diseases (J40-J47)	693	4.4	136.5
6	Alzheimer's disease (G30)	593	3.8	116.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	3.0	90.6
8	Influenza and pneumonia (J09-J18)	443	2.8	87.3
9	Septicemia (A40-A41)	275	1.8	54.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	1.7	53.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	267	1.7	52.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	1.4	44.5
13	Parkinson's disease (G20-G21)	178	1.1	35.1
14	Pneumonitis due to solids and liquids (J69)	93	0.6	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.6	17.1
...	All other causes (Residual)	2,661	17.1	524.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	16,358	100.0	7,804.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,144	31.4	2,454.3
2	Malignant neoplasms (C00-C97)	1,772	10.8	845.4
3	Cerebrovascular diseases (I60-I69)	1,294	7.9	617.4
4	Alzheimer's disease (G30)	1,283	7.8	612.1
5	Chronic lower respiratory diseases (J40-J47)	699	4.3	333.5
6	Diabetes mellitus (E10-E14)	619	3.8	295.3
7	Influenza and pneumonia (J09-J18)	561	3.4	267.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	363	2.2	173.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	306	1.9	146.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	1.6	126.4
11	Septicemia (A40-A41)	213	1.3	101.6
12	Pneumonitis due to solids and liquids (J69)	141	0.9	67.3
13	Parkinson's disease (G20-G21)	138	0.8	65.8
14	Atherosclerosis (I70)	93	0.6	44.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.6	43.4
...	All other causes (Residual)	3,376	20.6	1,610.7
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,944,606	100.0	961.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	485,779	25.0	240.3
2	Malignant neoplasms (C00-C97)	457,189	23.5	226.2
3	Chronic lower respiratory diseases (J40-J47)	122,605	6.3	60.6
4	Cerebrovascular diseases (I60-I69)	101,703	5.2	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,416	4.7	45.2
6	Alzheimer's disease (G30)	69,695	3.6	34.5
7	Diabetes mellitus (E10-E14)	47,851	2.5	23.7
8	Influenza and pneumonia (J09-J18)	42,752	2.2	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,670	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,813	1.6	15.2
11	Septicemia (A40-A41)	26,533	1.4	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,520	1.2	11.1
13	Parkinson's disease (G20-G21)	18,554	1.0	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,552	1.0	9.2
15	Pneumonitis due to solids and liquids (J69)	13,489	0.7	6.7
...	All other causes (Residual)	359,485	18.5	177.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,181	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	773	35.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	10.5	2.5
3	Malignant neoplasms (C00-C97)	180	8.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	6.5	1.5
5	Influenza and pneumonia (J09-J18)	75	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.2	0.8
7	Septicemia (A40-A41)	30	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	27	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
12	Meningitis (G00,G03)	11	0.5	*
13	Anemias (D50-D64)	10	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
...	All other causes (Residual)	545	25.0	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,867	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	895	31.2	3.8
2	Malignant neoplasms (C00-C97)	493	17.2	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	6.7	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	5.5	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	117	4.1	0.5
6	Influenza and pneumonia (J09-J18)	104	3.6	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.2	0.4
8	Cerebrovascular diseases (I60-I69)	43	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.4	0.2
10	Septicemia (A40-A41)	28	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	26	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	14	0.5	*
14	Anemias (D50-D64)	10	0.3	*
14	Meningitis (G00,G03)	10	0.3	*
...	All other causes (Residual)	632	22.0	2.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,150	100.0	65.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,564	49.9	32.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,046	17.8	11.6
3	Malignant neoplasms (C00-C97)	945	5.5	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	755	4.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	499	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	1.6	1.0
7	Influenza and pneumonia (J09-J18)	223	1.3	0.9
8	Cerebrovascular diseases (I60-I69)	97	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	93	0.5	0.4
10	Diabetes mellitus (E10-E14)	89	0.5	0.3
11	Septicemia (A40-A41)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	71	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.2	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	33	0.2	0.1
...	All other causes (Residual)	2,321	13.5	8.9
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	25,203	100.0	101.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,887	39.2	39.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,983	15.8	16.0
3	Malignant neoplasms (C00-C97)	2,206	8.8	8.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,688	6.7	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	930	3.7	3.7
6	Influenza and pneumonia (J09-J18)	434	1.7	1.7
7	Diabetes mellitus (E10-E14)	303	1.2	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	250	1.0	1.0
9	Cerebrovascular diseases (I60-I69)	245	1.0	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	242	1.0	1.0
11	Septicemia (A40-A41)	192	0.8	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	165	0.7	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	140	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	136	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.4	0.5
...	All other causes (Residual)	4,289	17.0	17.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	47,736	100.0	181.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,002	23.0	41.7
2	Malignant neoplasms (C00-C97)	8,213	17.2	31.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,847	14.3	26.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,508	11.5	20.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,618	3.4	6.1
6	Diabetes mellitus (E10-E14)	1,045	2.2	4.0
7	Cerebrovascular diseases (I60-I69)	962	2.0	3.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	943	2.0	3.6
9	Influenza and pneumonia (J09-J18)	717	1.5	2.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	639	1.3	2.4
11	Septicemia (A40-A41)	519	1.1	2.0
12	Chronic lower respiratory diseases (J40-J47)	469	1.0	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	0.8	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	0.7	1.3
15	Viral hepatitis (B15-B19)	247	0.5	0.9
...	All other causes (Residual)	8,299	17.4	31.5
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	130,740	100.0	412.1
1	Malignant neoplasms (C00-C97)	36,485	27.9	115.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,169	19.3	79.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,361	11.7	48.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,564	5.8	23.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,022	4.6	19.0
6	Chronic lower respiratory diseases (J40-J47)	3,699	2.8	11.7
7	Diabetes mellitus (E10-E14)	3,554	2.7	11.2
8	Cerebrovascular diseases (I60-I69)	3,424	2.6	10.8
9	Influenza and pneumonia (J09-J18)	1,899	1.5	6.0
10	Viral hepatitis (B15-B19)	1,771	1.4	5.6
11	Septicemia (A40-A41)	1,504	1.2	4.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,151	0.9	3.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,023	0.8	3.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	979	0.7	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	683	0.5	2.2
...	All other causes (Residual)	20,452	15.6	64.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	224,375	100.0	846.8
1	Malignant neoplasms (C00-C97)	81,811	36.5	308.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,396	21.6	182.7
3	Chronic lower respiratory diseases (J40-J47)	12,258	5.5	46.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,011	4.5	37.8
5	Diabetes mellitus (E10-E14)	7,289	3.2	27.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6,770	3.0	25.6
7	Cerebrovascular diseases (I60-I69)	6,570	2.9	24.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,221	2.3	19.7
9	Septicemia (A40-A41)	3,091	1.4	11.7
10	Influenza and pneumonia (J09-J18)	3,021	1.3	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,771	1.2	10.5
12	Viral hepatitis (B15-B19)	2,057	0.9	7.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,411	0.6	5.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,118	0.5	4.2
15	Aortic aneurysm and dissection (I71)	911	0.4	3.4
...	All other causes (Residual)	31,669	14.1	119.5
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	315,532	100.0	1,925.7
1	Malignant neoplasms (C00-C97)	114,690	36.3	699.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,466	21.7	417.8
3	Chronic lower respiratory diseases (J40-J47)	28,100	8.9	171.5
4	Cerebrovascular diseases (I60-I69)	12,705	4.0	77.5
5	Diabetes mellitus (E10-E14)	9,951	3.2	60.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,311	2.3	44.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,338	1.7	32.6
8	Septicemia (A40-A41)	4,817	1.5	29.4
9	Influenza and pneumonia (J09-J18)	4,693	1.5	28.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,104	1.3	25.0
11	Alzheimer's disease (G30)	3,469	1.1	21.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,639	0.8	16.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,181	0.7	13.3
14	Parkinson's disease (G20-G21)	2,079	0.7	12.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,034	0.6	12.4
...	All other causes (Residual)	42,955	13.6	262.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	524,664	100.0	4,904.4
1	Malignant neoplasms (C00-C97)	132,982	25.3	1,243.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	130,386	24.9	1,218.8
3	Chronic lower respiratory diseases (J40-J47)	44,529	8.5	416.2
4	Cerebrovascular diseases (I60-I69)	31,136	5.9	291.0
5	Alzheimer's disease (G30)	20,305	3.9	189.8
6	Diabetes mellitus (E10-E14)	13,759	2.6	128.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,803	2.2	110.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,395	2.2	106.5
9	Influenza and pneumonia (J09-J18)	11,357	2.2	106.2
10	Parkinson's disease (G20-G21)	8,279	1.6	77.4
11	Septicemia (A40-A41)	8,129	1.5	76.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,942	0.9	46.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,980	0.8	37.2
14	Pneumonitis due to solids and liquids (J69)	3,936	0.8	36.8
15	Aortic aneurysm and dissection (I71)	3,010	0.6	28.1
...	All other causes (Residual)	84,736	16.2	792.1
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	642,473	100.0	13,477.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	203,988	31.8	4,279.3
2	Malignant neoplasms (C00-C97)	79,137	12.3	1,660.2
3	Cerebrovascular diseases (I60-I69)	46,426	7.2	973.9
4	Alzheimer's disease (G30)	45,219	7.0	948.6
5	Chronic lower respiratory diseases (J40-J47)	33,286	5.2	698.3
6	Influenza and pneumonia (J09-J18)	20,132	3.1	422.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,236	2.4	319.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,441	2.2	302.9
9	Diabetes mellitus (E10-E14)	11,845	1.8	248.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,111	1.4	191.1
11	Septicemia (A40-A41)	8,072	1.3	169.3
12	Parkinson's disease (G20-G21)	7,746	1.2	162.5
13	Pneumonitis due to solids and liquids (J69)	6,931	1.1	145.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,148	0.6	87.0
15	Atherosclerosis (I70)	3,762	0.6	78.9
...	All other causes (Residual)	132,993	20.7	2,790.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	959,014	100.0	965.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,489	25.8	249.2
2	Malignant neoplasms (C00-C97)	239,301	25.0	241.0
3	Chronic lower respiratory diseases (J40-J47)	57,370	6.0	57.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,607	5.9	57.0
5	Cerebrovascular diseases (I60-I69)	39,912	4.2	40.2
6	Diabetes mellitus (E10-E14)	24,993	2.6	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,248	2.5	24.4
8	Alzheimer's disease (G30)	21,087	2.2	21.2
9	Influenza and pneumonia (J09-J18)	19,736	2.1	19.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,735	1.8	17.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,494	1.5	14.6
12	Septicemia (A40-A41)	12,167	1.3	12.3
13	Parkinson's disease (G20-G21)	10,919	1.1	11.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,316	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	7,068	0.7	7.1
...	All other causes (Residual)	158,572	16.5	159.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,262	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	500	39.6	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	8.8	2.4
3	Malignant neoplasms (C00-C97)	106	8.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.2	0.9
6	Influenza and pneumonia (J09-J18)	32	2.5	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.4	*
8	Septicemia (A40-A41)	17	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Anemias (D50-D64)	8	0.6	*
12	Meningitis (G00,G03)	8	0.6	*
14	Meningococcal infection (A39)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	298	23.6	6.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,684	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	574	34.1	4.8
2	Malignant neoplasms (C00-C97)	277	16.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	6.5	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	4.0	0.6
6	Influenza and pneumonia (J09-J18)	51	3.0	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.8	0.4
8	Cerebrovascular diseases (I60-I69)	26	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
11	Septicemia (A40-A41)	11	0.7	*
12	Meningitis (G00,G03)	7	0.4	*
13	Diabetes mellitus (E10-E14)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
14	Diseases of appendix (K35-K38)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (Residual)	351	20.8	2.9
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,318	100.0	91.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,301	51.2	46.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,533	20.6	18.9
3	Malignant neoplasms (C00-C97)	578	4.7	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	536	4.4	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	1.3	1.2
7	Influenza and pneumonia (J09-J18)	129	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	54	0.4	0.4
9	Diabetes mellitus (E10-E14)	53	0.4	0.4
10	Septicemia (A40-A41)	45	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	41	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.2	0.2
12	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	21	0.2	0.2
...	All other causes (Residual)	1,449	11.8	10.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	17,165	100.0	136.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,286	42.4	58.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,218	18.7	25.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,194	7.0	9.5
4	Malignant neoplasms (C00-C97)	1,095	6.4	8.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	626	3.6	5.0
6	Influenza and pneumonia (J09-J18)	225	1.3	1.8
7	Diabetes mellitus (E10-E14)	173	1.0	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	151	0.9	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.9	1.2
10	Cerebrovascular diseases (I60-I69)	136	0.8	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	98	0.6	0.8
12	Septicemia (A40-A41)	92	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	63	0.4	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.3	0.4
...	All other causes (Residual)	2,562	14.9	20.4
Non-Hispanic, white, male, 35-44 years				
...	All causes	29,834	100.0	225.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,589	25.4	57.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,846	16.2	36.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,209	14.1	31.8
4	Malignant neoplasms (C00-C97)	3,615	12.1	27.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,024	3.4	7.7
6	Diabetes mellitus (E10-E14)	687	2.3	5.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	616	2.1	4.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	525	1.8	4.0
9	Cerebrovascular diseases (I60-I69)	516	1.7	3.9
10	Influenza and pneumonia (J09-J18)	392	1.3	3.0
11	Septicemia (A40-A41)	251	0.8	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	211	0.7	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	209	0.7	1.6
14	Chronic lower respiratory diseases (J40-J47)	203	0.7	1.5
15	Viral hepatitis (B15-B19)	162	0.5	1.2
...	All other causes (Residual)	4,779	16.0	36.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	80,827	100.0	513.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,565	23.0	118.0
2	Malignant neoplasms (C00-C97)	18,487	22.9	117.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,215	12.6	64.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,680	7.0	36.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,077	5.0	25.9
6	Diabetes mellitus (E10-E14)	2,284	2.8	14.5
7	Cerebrovascular diseases (I60-I69)	1,903	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,806	2.2	11.5
9	Viral hepatitis (B15-B19)	1,242	1.5	7.9
10	Influenza and pneumonia (J09-J18)	1,002	1.2	6.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	853	1.1	5.4
12	Septicemia (A40-A41)	813	1.0	5.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	681	0.8	4.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	668	0.8	4.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	468	0.6	3.0
...	All other causes (Residual)	12,083	14.9	76.8
Non-Hispanic, white, male, 55-64 years				
...	All causes	136,875	100.0	1,057.5
1	Malignant neoplasms (C00-C97)	45,236	33.0	349.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,455	25.2	266.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,841	5.0	52.9
4	Chronic lower respiratory diseases (J40-J47)	6,306	4.6	48.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,797	3.5	37.1
6	Diabetes mellitus (E10-E14)	4,538	3.3	35.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,986	2.9	30.8
8	Cerebrovascular diseases (I60-I69)	3,740	2.7	28.9
9	Influenza and pneumonia (J09-J18)	1,689	1.2	13.0
10	Septicemia (A40-A41)	1,633	1.2	12.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,602	1.2	12.4
12	Viral hepatitis (B15-B19)	1,523	1.1	11.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	906	0.7	7.0
14	Aortic aneurysm and dissection (I71)	685	0.5	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	658	0.5	5.1
...	All other causes (Residual)	18,280	13.4	141.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	178,722	100.0	2,332.6
1	Malignant neoplasms (C00-C97)	63,024	35.3	822.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,727	24.5	570.7
3	Chronic lower respiratory diseases (J40-J47)	13,926	7.8	181.8
4	Cerebrovascular diseases (I60-I69)	6,697	3.7	87.4
5	Diabetes mellitus (E10-E14)	5,802	3.2	75.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,588	2.6	59.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,942	1.6	38.4
8	Influenza and pneumonia (J09-J18)	2,595	1.5	33.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,553	1.4	33.3
10	Septicemia (A40-A41)	2,459	1.4	32.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,174	1.2	28.4
12	Alzheimer's disease (G30)	1,539	0.9	20.1
13	Parkinson's disease (G20-G21)	1,398	0.8	18.2
14	Aortic aneurysm and dissection (I71)	1,198	0.7	15.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,194	0.7	15.6
...	All other causes (Residual)	22,906	12.8	299.0
Non-Hispanic, white, male, 75-84 years				
...	All causes	262,553	100.0	5,857.5
1	Malignant neoplasms (C00-C97)	70,578	26.9	1,574.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,219	26.4	1,544.3
3	Chronic lower respiratory diseases (J40-J47)	21,310	8.1	475.4
4	Cerebrovascular diseases (I60-I69)	13,316	5.1	297.1
5	Alzheimer's disease (G30)	7,549	2.9	168.4
6	Diabetes mellitus (E10-E14)	6,911	2.6	154.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,376	2.4	142.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,910	2.3	131.9
9	Influenza and pneumonia (J09-J18)	5,763	2.2	128.6
10	Parkinson's disease (G20-G21)	5,105	1.9	113.9
11	Septicemia (A40-A41)	3,846	1.5	85.8
12	Pneumonitis due to solids and liquids (J69)	2,230	0.8	49.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,198	0.8	49.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,095	0.8	46.7
15	Aortic aneurysm and dissection (I71)	1,699	0.6	37.9
...	All other causes (Residual)	38,448	14.6	857.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	231,198	100.0	15,361.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,962	32.4	4,980.5
2	Malignant neoplasms (C00-C97)	36,278	15.7	2,410.3
3	Chronic lower respiratory diseases (J40-J47)	13,677	5.9	908.7
4	Cerebrovascular diseases (I60-I69)	13,471	5.8	895.0
5	Alzheimer's disease (G30)	11,690	5.1	776.7
6	Influenza and pneumonia (J09-J18)	7,799	3.4	518.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,275	2.7	416.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,997	2.6	398.4
9	Diabetes mellitus (E10-E14)	4,537	2.0	301.4
10	Parkinson's disease (G20-G21)	4,125	1.8	274.1
11	Pneumonitis due to solids and liquids (J69)	3,279	1.4	217.9
12	Septicemia (A40-A41)	2,957	1.3	196.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,527	1.1	167.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,825	0.8	121.3
15	Atherosclerosis (I70)	1,191	0.5	79.1
...	All other causes (Residual)	40,608	17.6	2,698.0
Non-Hispanic, white, female, all ages				
...	All causes	985,592	100.0	958.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,290	24.2	231.7
2	Malignant neoplasms (C00-C97)	217,888	22.1	211.9
3	Chronic lower respiratory diseases (J40-J47)	65,235	6.6	63.4
4	Cerebrovascular diseases (I60-I69)	61,791	6.3	60.1
5	Alzheimer's disease (G30)	48,608	4.9	47.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,809	3.5	33.8
7	Influenza and pneumonia (J09-J18)	23,016	2.3	22.4
8	Diabetes mellitus (E10-E14)	22,858	2.3	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,935	1.8	17.4
10	Septicemia (A40-A41)	14,366	1.5	14.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,236	1.1	10.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,026	0.8	7.8
13	Parkinson's disease (G20-G21)	7,635	0.8	7.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,565	0.7	6.4
15	Pneumonitis due to solids and liquids (J69)	6,421	0.7	6.2
...	All other causes (Residual)	200,913	20.4	195.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	919	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	29.7	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	12.9	2.7
3	Malignant neoplasms (C00-C97)	74	8.1	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	7.9	1.6
5	Influenza and pneumonia (J09-J18)	43	4.7	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
7	Septicemia (A40-A41)	13	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
8	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Meningitis (G00,G03)	3	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
...	All other causes (Residual)	240	26.1	5.4
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,183	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.1	2.8
2	Malignant neoplasms (C00-C97)	216	18.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	6.9	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	4.7	0.5
5	Influenza and pneumonia (J09-J18)	53	4.5	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.1	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.7	0.4
8	Septicemia (A40-A41)	17	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.4	*
8	Cerebrovascular diseases (I60-I69)	17	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
14	Anemias (D50-D64)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	274	23.2	2.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,832	100.0	37.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,263	46.8	17.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	513	10.6	4.0
3	Malignant neoplasms (C00-C97)	367	7.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	4.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	2.4	0.9
7	Influenza and pneumonia (J09-J18)	94	1.9	0.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	93	1.9	0.7
9	Cerebrovascular diseases (I60-I69)	43	0.9	0.3
10	Diabetes mellitus (E10-E14)	36	0.7	0.3
11	Chronic lower respiratory diseases (J40-J47)	30	0.6	0.2
12	Septicemia (A40-A41)	28	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
14	Pneumonitis due to solids and liquids (J69)	12	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Meningitis (G00,G03)	11	0.2	*
...	All other causes (Residual)	815	16.9	6.4
Non-Hispanic, white, female, 25-34 years				
...	All causes	8,038	100.0	65.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,601	32.4	21.1
2	Malignant neoplasms (C00-C97)	1,111	13.8	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	765	9.5	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	494	6.1	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	3.8	2.5
6	Influenza and pneumonia (J09-J18)	209	2.6	1.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	165	2.1	1.3
8	Diabetes mellitus (E10-E14)	130	1.6	1.1
9	Cerebrovascular diseases (I60-I69)	109	1.4	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.3	0.8
11	Septicemia (A40-A41)	100	1.2	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	73	0.9	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.7	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.5	0.3
...	All other causes (Residual)	1,682	20.9	13.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	17,902	100.0	136.2
1	Malignant neoplasms (C00-C97)	4,598	25.7	35.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,413	19.1	26.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,001	11.2	15.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,299	7.3	9.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	594	3.3	4.5
6	Cerebrovascular diseases (I60-I69)	446	2.5	3.4
7	Diabetes mellitus (E10-E14)	358	2.0	2.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	1.8	2.5
9	Influenza and pneumonia (J09-J18)	325	1.8	2.5
10	Septicemia (A40-A41)	268	1.5	2.0
11	Chronic lower respiratory diseases (J40-J47)	266	1.5	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	0.8	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	114	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	105	0.6	0.8
...	All other causes (Residual)	3,500	19.6	26.6
Non-Hispanic, white, female, 45-54 years				
...	All causes	49,913	100.0	312.1
1	Malignant neoplasms (C00-C97)	17,998	36.1	112.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,604	13.2	41.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,146	10.3	32.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,945	3.9	12.2
5	Chronic lower respiratory diseases (J40-J47)	1,893	3.8	11.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,884	3.8	11.8
7	Cerebrovascular diseases (I60-I69)	1,521	3.0	9.5
8	Diabetes mellitus (E10-E14)	1,270	2.5	7.9
9	Influenza and pneumonia (J09-J18)	897	1.8	5.6
10	Septicemia (A40-A41)	691	1.4	4.3
11	Viral hepatitis (B15-B19)	529	1.1	3.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	470	0.9	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	311	0.6	1.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	241	0.5	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	215	0.4	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	215	0.4	1.3
...	All other causes (Residual)	8,083	16.2	50.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	87,500	100.0	645.6
1	Malignant neoplasms (C00-C97)	36,575	41.8	269.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,941	15.9	102.9
3	Chronic lower respiratory diseases (J40-J47)	5,952	6.8	43.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,170	3.6	23.4
5	Cerebrovascular diseases (I60-I69)	2,830	3.2	20.9
6	Diabetes mellitus (E10-E14)	2,751	3.1	20.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,973	2.3	14.6
8	Septicemia (A40-A41)	1,458	1.7	10.8
9	Influenza and pneumonia (J09-J18)	1,332	1.5	9.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,235	1.4	9.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,169	1.3	8.6
12	Viral hepatitis (B15-B19)	534	0.6	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	505	0.6	3.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	460	0.5	3.4
15	Alzheimer's disease (G30)	336	0.4	2.5
...	All other causes (Residual)	13,279	15.2	98.0
Non-Hispanic, white, female, 65-74 years				
...	All causes	136,810	100.0	1,568.3
1	Malignant neoplasms (C00-C97)	51,666	37.8	592.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,739	18.1	283.6
3	Chronic lower respiratory diseases (J40-J47)	14,174	10.4	162.5
4	Cerebrovascular diseases (I60-I69)	6,008	4.4	68.9
5	Diabetes mellitus (E10-E14)	4,149	3.0	47.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,723	2.0	31.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,396	1.8	27.5
8	Septicemia (A40-A41)	2,358	1.7	27.0
9	Influenza and pneumonia (J09-J18)	2,098	1.5	24.0
10	Alzheimer's disease (G30)	1,930	1.4	22.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,551	1.1	17.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,014	0.7	11.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	840	0.6	9.6
14	Parkinson's disease (G20-G21)	681	0.5	7.8
15	Aortic aneurysm and dissection (I71)	621	0.5	7.1
...	All other causes (Residual)	19,862	14.5	227.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	262,111	100.0	4,217.0
1	Malignant neoplasms (C00-C97)	62,404	23.8	1,004.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,167	23.3	984.1
3	Chronic lower respiratory diseases (J40-J47)	23,219	8.9	373.6
4	Cerebrovascular diseases (I60-I69)	17,820	6.8	286.7
5	Alzheimer's disease (G30)	12,756	4.9	205.2
6	Diabetes mellitus (E10-E14)	6,848	2.6	110.2
7	Influenza and pneumonia (J09-J18)	5,594	2.1	90.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,485	2.1	88.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,427	2.1	87.3
10	Septicemia (A40-A41)	4,283	1.6	68.9
11	Parkinson's disease (G20-G21)	3,174	1.2	51.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,847	1.1	45.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,782	0.7	28.7
14	Pneumonitis due to solids and liquids (J69)	1,706	0.7	27.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,330	0.5	21.4
...	All other causes (Residual)	46,269	17.7	744.4
Non-Hispanic, white, female, 85 years and over				
...	All causes	411,275	100.0	12,609.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	129,026	31.4	3,955.7
2	Malignant neoplasms (C00-C97)	42,859	10.4	1,314.0
3	Alzheimer's disease (G30)	33,529	8.2	1,027.9
4	Cerebrovascular diseases (I60-I69)	32,955	8.0	1,010.3
5	Chronic lower respiratory diseases (J40-J47)	19,609	4.8	601.2
6	Influenza and pneumonia (J09-J18)	12,333	3.0	378.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,239	2.2	283.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,166	2.0	250.4
9	Diabetes mellitus (E10-E14)	7,308	1.8	224.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,584	1.6	201.9
11	Septicemia (A40-A41)	5,115	1.2	156.8
12	Pneumonitis due to solids and liquids (J69)	3,652	0.9	112.0
13	Parkinson's disease (G20-G21)	3,621	0.9	111.0
14	Atherosclerosis (I70)	2,571	0.6	78.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,323	0.6	71.2
...	All other causes (Residual)	92,385	22.5	2,832.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	282,982	100.0	728.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,811	24.3	177.1
2	Malignant neoplasms (C00-C97)	63,967	22.6	164.6
3	Cerebrovascular diseases (I60-I69)	15,718	5.6	40.4
4	Diabetes mellitus (E10-E14)	11,833	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,810	4.2	30.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,727	3.1	22.5
7	Chronic lower respiratory diseases (J40-J47)	8,444	3.0	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,733	2.7	19.9
9	Septicemia (A40-A41)	6,131	2.2	15.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,307	1.9	13.7
11	Influenza and pneumonia (J09-J18)	5,214	1.8	13.4
12	Alzheimer's disease (G30)	5,042	1.8	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,819	1.7	12.4
14	Certain conditions originating in the perinatal period (P00-P96)	4,456	1.6	11.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,518	0.9	6.5
...	All other causes (Residual)	52,452	18.5	135.0
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,039	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	27.7	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	13.8	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	8.4	3.5
4	Malignant neoplasms (C00-C97)	56	5.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	33	3.2	1.3
7	Influenza and pneumonia (J09-J18)	29	2.8	1.2
8	Septicemia (A40-A41)	19	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	19	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	12	1.2	*
12	Cerebrovascular diseases (I60-I69)	9	0.9	*
13	Anemias (D50-D64)	8	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	260	25.0	10.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,266	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	29.8	6.3
2	Malignant neoplasms (C00-C97)	128	10.1	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	8.3	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	70	5.5	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.1	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	3.7	0.8
8	Influenza and pneumonia (J09-J18)	45	3.6	0.7
9	Septicemia (A40-A41)	22	1.7	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.6	0.3
11	Anemias (D50-D64)	19	1.5	*
12	Cerebrovascular diseases (I60-I69)	18	1.4	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	248	19.6	4.1
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,602	100.0	99.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,760	41.8	41.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	19.8	19.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	433	6.6	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	5.1	5.1
5	Malignant neoplasms (C00-C97)	284	4.3	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	106	1.6	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.4	1.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	88	1.3	1.3
9	Influenza and pneumonia (J09-J18)	85	1.3	1.3
9	Chronic lower respiratory diseases (J40-J47)	85	1.3	1.3
11	Anemias (D50-D64)	66	1.0	1.0
12	Diabetes mellitus (E10-E14)	50	0.8	0.8
13	Cerebrovascular diseases (I60-I69)	39	0.6	0.6
14	Septicemia (A40-A41)	31	0.5	0.5
15	Legal intervention (Y35,Y89.0)	30	0.5	0.5
...	All other causes (Residual)	805	12.2	12.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	9,031	100.0	160.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,291	25.4	40.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,566	17.3	27.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	949	10.5	16.8
4	Malignant neoplasms (C00-C97)	626	6.9	11.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	596	6.6	10.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	463	5.1	8.2
7	Diabetes mellitus (E10-E14)	189	2.1	3.3
8	Cerebrovascular diseases (I60-I69)	143	1.6	2.5
8	Influenza and pneumonia (J09-J18)	143	1.6	2.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	129	1.4	2.3
11	Anemias (D50-D64)	111	1.2	2.0
12	Septicemia (A40-A41)	107	1.2	1.9
13	Chronic lower respiratory diseases (J40-J47)	100	1.1	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	1.1	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.9	1.4
...	All other causes (Residual)	1,442	16.0	25.5
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	14,951	100.0	284.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,863	19.1	54.4
2	Malignant neoplasms (C00-C97)	2,133	14.3	40.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,736	11.6	33.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,368	9.1	26.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,175	7.9	22.3
6	Cerebrovascular diseases (I60-I69)	543	3.6	10.3
7	Diabetes mellitus (E10-E14)	524	3.5	10.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	398	2.7	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	2.1	5.9
10	Septicemia (A40-A41)	271	1.8	5.2
11	Influenza and pneumonia (J09-J18)	252	1.7	4.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	201	1.3	3.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	184	1.2	3.5
14	Chronic lower respiratory diseases (J40-J47)	179	1.2	3.4
15	Anemias (D50-D64)	131	0.9	2.5
...	All other causes (Residual)	2,682	17.9	51.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	35,062	100.0	664.3
1	Malignant neoplasms (C00-C97)	8,528	24.3	161.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,088	23.1	153.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,398	6.8	45.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,845	5.3	35.0
5	Cerebrovascular diseases (I60-I69)	1,780	5.1	33.7
6	Diabetes mellitus (E10-E14)	1,328	3.8	25.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	856	2.4	16.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	799	2.3	15.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	714	2.0	13.5
10	Chronic lower respiratory diseases (J40-J47)	702	2.0	13.3
11	Septicemia (A40-A41)	630	1.8	11.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	554	1.6	10.5
13	Influenza and pneumonia (J09-J18)	519	1.5	9.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	366	1.0	6.9
15	Viral hepatitis (B15-B19)	334	1.0	6.3
...	All other causes (Residual)	5,621	16.0	106.5
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	50,089	100.0	1,404.5
1	Malignant neoplasms (C00-C97)	15,544	31.0	435.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,776	25.5	358.2
3	Cerebrovascular diseases (I60-I69)	2,579	5.1	72.3
4	Diabetes mellitus (E10-E14)	2,428	4.8	68.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,608	3.2	45.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,415	2.8	39.7
7	Chronic lower respiratory diseases (J40-J47)	1,363	2.7	38.2
8	Septicemia (A40-A41)	1,076	2.1	30.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,033	2.1	29.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	928	1.9	26.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	824	1.6	23.1
12	Influenza and pneumonia (J09-J18)	640	1.3	17.9
13	Viral hepatitis (B15-B19)	514	1.0	14.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	0.5	7.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	225	0.4	6.3
...	All other causes (Residual)	6,867	13.7	192.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	51,335	100.0	2,699.0
1	Malignant neoplasms (C00-C97)	16,230	31.6	853.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,392	26.1	704.1
3	Cerebrovascular diseases (I60-I69)	2,979	5.8	156.6
4	Diabetes mellitus (E10-E14)	2,748	5.4	144.5
5	Chronic lower respiratory diseases (J40-J47)	2,057	4.0	108.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,919	3.7	100.9
7	Septicemia (A40-A41)	1,261	2.5	66.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	865	1.7	45.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	810	1.6	42.6
10	Influenza and pneumonia (J09-J18)	808	1.6	42.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	367	0.7	19.3
12	Alzheimer's disease (G30)	361	0.7	19.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	274	0.5	14.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	220	0.4	11.6
15	Pneumonitis due to solids and liquids (J69)	219	0.4	11.5
...	All other causes (Residual)	6,825	13.3	358.8
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	56,785	100.0	5,261.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,338	27.0	1,421.0
2	Malignant neoplasms (C00-C97)	13,839	24.4	1,282.1
3	Cerebrovascular diseases (I60-I69)	3,965	7.0	367.3
4	Diabetes mellitus (E10-E14)	2,745	4.8	254.3
5	Chronic lower respiratory diseases (J40-J47)	2,317	4.1	214.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,290	4.0	212.2
7	Alzheimer's disease (G30)	1,656	2.9	153.4
8	Septicemia (A40-A41)	1,449	2.6	134.2
9	Influenza and pneumonia (J09-J18)	1,234	2.2	114.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,071	1.9	99.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	713	1.3	66.1
12	Pneumonitis due to solids and liquids (J69)	416	0.7	38.5
13	Parkinson's disease (G20-G21)	319	0.6	29.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	291	0.5	27.0
15	Aortic aneurysm and dissection (I71)	198	0.3	18.3
...	All other causes (Residual)	8,944	15.8	828.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	48,832	100.0	13,734.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,833	30.4	4,171.8
2	Malignant neoplasms (C00-C97)	6,584	13.5	1,851.8
3	Cerebrovascular diseases (I60-I69)	3,619	7.4	1,017.9
4	Alzheimer's disease (G30)	2,945	6.0	828.3
5	Diabetes mellitus (E10-E14)	1,812	3.7	509.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,766	3.6	496.7
7	Chronic lower respiratory diseases (J40-J47)	1,529	3.1	430.0
8	Influenza and pneumonia (J09-J18)	1,378	2.8	387.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,233	2.5	346.8
10	Septicemia (A40-A41)	1,177	2.4	331.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	614	1.3	172.7
12	Pneumonitis due to solids and liquids (J69)	475	1.0	133.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	248	0.5	69.8
14	Parkinson's disease (G20-G21)	244	0.5	68.6
15	Atherosclerosis (I70)	212	0.4	59.6
...	All other causes (Residual)	10,163	20.8	2,858.4
Non-Hispanic, black, male, all ages				
...	All causes	144,197	100.0	777.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,935	24.2	188.3
2	Malignant neoplasms (C00-C97)	33,065	22.9	178.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,894	5.5	42.5
4	Cerebrovascular diseases (I60-I69)	6,892	4.8	37.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,605	4.6	35.6
6	Diabetes mellitus (E10-E14)	5,430	3.8	29.3
7	Chronic lower respiratory diseases (J40-J47)	4,457	3.1	24.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,930	2.7	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,476	2.4	18.7
10	Septicemia (A40-A41)	2,721	1.9	14.7
11	Certain conditions originating in the perinatal period (P00-P96)	2,533	1.8	13.7
12	Influenza and pneumonia (J09-J18)	2,463	1.7	13.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,045	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,649	1.1	8.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,636	1.1	8.8
...	All other causes (Residual)	24,466	17.0	131.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	561	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	30.1	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	14.1	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	8.6	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.9	1.7
5	Malignant neoplasms (C00-C97)	20	3.6	1.6
6	Chronic lower respiratory diseases (J40-J47)	18	3.2	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	6	1.1	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	141	25.1	11.1
Non-Hispanic, black, male, 5-14 years				
...	All causes	735	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	32.4	7.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	9.9	2.4
3	Malignant neoplasms (C00-C97)	60	8.2	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	6.4	1.5
5	Chronic lower respiratory diseases (J40-J47)	43	5.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.8	1.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	4.8	1.1
8	Influenza and pneumonia (J09-J18)	22	3.0	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.8	*
10	Cerebrovascular diseases (I60-I69)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
12	Anemias (D50-D64)	7	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	128	17.4	4.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,985	100.0	148.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,483	49.8	74.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	937	18.8	27.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	360	7.2	10.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	4.5	6.7
5	Malignant neoplasms (C00-C97)	158	3.2	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	1.3	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.1	1.7
8	Chronic lower respiratory diseases (J40-J47)	53	1.1	1.6
9	Influenza and pneumonia (J09-J18)	40	0.8	1.2
10	Anemias (D50-D64)	36	0.7	1.1
11	Legal intervention (Y35,Y89.0)	29	0.6	0.9
12	Diabetes mellitus (E10-E14)	25	0.5	0.7
13	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
14	Septicemia (A40-A41)	12	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (Residual)	474	9.5	14.1
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,112	100.0	221.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,014	33.0	72.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,187	19.4	43.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	573	9.4	20.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	376	6.2	13.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	342	5.6	12.4
6	Malignant neoplasms (C00-C97)	280	4.6	10.1
7	Diabetes mellitus (E10-E14)	109	1.8	3.9
8	Cerebrovascular diseases (I60-I69)	79	1.3	2.9
9	Influenza and pneumonia (J09-J18)	62	1.0	2.2
10	Anemias (D50-D64)	60	1.0	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.9	2.0
12	Chronic lower respiratory diseases (J40-J47)	52	0.9	1.9
13	Septicemia (A40-A41)	50	0.8	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.7	1.4
15	Legal intervention (Y35,Y89.0)	37	0.6	1.3
...	All other causes (Residual)	795	13.0	28.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	8,826	100.0	356.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,839	20.8	74.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,238	14.0	50.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	994	11.3	40.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	844	9.6	34.1
5	Malignant neoplasms (C00-C97)	820	9.3	33.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	3.7	13.3
7	Diabetes mellitus (E10-E14)	327	3.7	13.2
8	Cerebrovascular diseases (I60-I69)	288	3.3	11.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	1.8	6.4
10	Septicemia (A40-A41)	130	1.5	5.3
11	Influenza and pneumonia (J09-J18)	117	1.3	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	113	1.3	4.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.3	4.5
14	Chronic lower respiratory diseases (J40-J47)	106	1.2	4.3
15	Anemias (D50-D64)	62	0.7	2.5
...	All other causes (Residual)	1,349	15.3	54.5
Non-Hispanic, black, male, 45-54 years				
...	All causes	20,066	100.0	823.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,110	25.5	209.8
2	Malignant neoplasms (C00-C97)	4,156	20.7	170.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,610	8.0	66.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,213	6.0	49.8
5	Cerebrovascular diseases (I60-I69)	916	4.6	37.6
6	Diabetes mellitus (E10-E14)	772	3.8	31.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	560	2.8	23.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	529	2.6	21.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	2.3	18.8
10	Chronic lower respiratory diseases (J40-J47)	333	1.7	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	326	1.6	13.4
12	Septicemia (A40-A41)	320	1.6	13.1
13	Influenza and pneumonia (J09-J18)	275	1.4	11.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	275	1.4	11.3
15	Viral hepatitis (B15-B19)	212	1.1	8.7
...	All other causes (Residual)	3,001	15.0	123.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	29,527	100.0	1,859.5
1	Malignant neoplasms (C00-C97)	8,699	29.5	547.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,005	27.1	504.1
3	Cerebrovascular diseases (I60-I69)	1,462	5.0	92.1
4	Diabetes mellitus (E10-E14)	1,293	4.4	81.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,136	3.8	71.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	757	2.6	47.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	756	2.6	47.6
8	Chronic lower respiratory diseases (J40-J47)	749	2.5	47.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	638	2.2	40.2
10	Septicemia (A40-A41)	565	1.9	35.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	463	1.6	29.2
12	Viral hepatitis (B15-B19)	372	1.3	23.4
13	Influenza and pneumonia (J09-J18)	353	1.2	22.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	223	0.8	14.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	0.5	8.9
...	All other causes (Residual)	3,914	13.3	246.5
Non-Hispanic, black, male, 65-74 years				
...	All causes	27,768	100.0	3,498.9
1	Malignant neoplasms (C00-C97)	8,877	32.0	1,118.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,628	27.5	961.2
3	Cerebrovascular diseases (I60-I69)	1,565	5.6	197.2
4	Diabetes mellitus (E10-E14)	1,331	4.8	167.7
5	Chronic lower respiratory diseases (J40-J47)	1,136	4.1	143.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	924	3.3	116.4
7	Septicemia (A40-A41)	597	2.1	75.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	1.9	67.7
9	Influenza and pneumonia (J09-J18)	465	1.7	58.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	404	1.5	50.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	234	0.8	29.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	197	0.7	24.8
13	Alzheimer's disease (G30)	163	0.6	20.5
14	Pneumonitis due to solids and liquids (J69)	114	0.4	14.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.4	13.9
...	All other causes (Residual)	3,486	12.6	439.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	25,790	100.0	6,494.0
1	Malignant neoplasms (C00-C97)	7,050	27.3	1,775.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,951	27.0	1,750.3
3	Cerebrovascular diseases (I60-I69)	1,586	6.1	399.4
4	Chronic lower respiratory diseases (J40-J47)	1,273	4.9	320.5
5	Diabetes mellitus (E10-E14)	1,082	4.2	272.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	979	3.8	246.5
7	Septicemia (A40-A41)	599	2.3	150.8
8	Influenza and pneumonia (J09-J18)	591	2.3	148.8
9	Alzheimer's disease (G30)	589	2.3	148.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	400	1.6	100.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	1.6	100.7
12	Pneumonitis due to solids and liquids (J69)	208	0.8	52.4
13	Parkinson's disease (G20-G21)	183	0.7	46.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.5	33.2
15	Aortic aneurysm and dissection (I71)	88	0.3	22.2
...	All other causes (Residual)	3,679	14.3	926.4
Non-Hispanic, black, male, 85 years and over				
...	All causes	15,289	100.0	15,823.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,473	29.3	4,629.5
2	Malignant neoplasms (C00-C97)	2,936	19.2	3,038.7
3	Cerebrovascular diseases (I60-I69)	932	6.1	964.6
4	Chronic lower respiratory diseases (J40-J47)	689	4.5	713.1
5	Alzheimer's disease (G30)	633	4.1	655.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	560	3.7	579.6
7	Diabetes mellitus (E10-E14)	488	3.2	505.1
8	Influenza and pneumonia (J09-J18)	479	3.1	495.8
9	Septicemia (A40-A41)	384	2.5	397.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	307	2.0	317.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	1.4	222.5
12	Pneumonitis due to solids and liquids (J69)	185	1.2	191.5
13	Parkinson's disease (G20-G21)	126	0.8	130.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	76.6
15	Atherosclerosis (I70)	57	0.4	59.0
...	All other causes (Residual)	2,751	18.0	2,847.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	138,785	100.0	683.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,876	24.4	166.8
2	Malignant neoplasms (C00-C97)	30,902	22.3	152.2
3	Cerebrovascular diseases (I60-I69)	8,826	6.4	43.5
4	Diabetes mellitus (E10-E14)	6,403	4.6	31.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,797	3.5	23.6
6	Chronic lower respiratory diseases (J40-J47)	3,987	2.9	19.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,916	2.8	19.3
8	Alzheimer's disease (G30)	3,612	2.6	17.8
9	Septicemia (A40-A41)	3,410	2.5	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,774	2.0	13.7
11	Influenza and pneumonia (J09-J18)	2,751	2.0	13.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,923	1.4	9.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,831	1.3	9.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,128	0.8	5.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	882	0.6	4.3
...	All other causes (Residual)	27,767	20.0	136.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	478	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.9	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	13.4	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	8.2	3.2
4	Malignant neoplasms (C00-C97)	36	7.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.9	2.3
6	Influenza and pneumonia (J09-J18)	16	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	15	3.1	*
8	Septicemia (A40-A41)	12	2.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.9	*
10	Cerebrovascular diseases (I60-I69)	5	1.0	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	119	24.9	9.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	531	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	26.2	4.7
2	Malignant neoplasms (C00-C97)	68	12.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	6.0	1.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.0	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.5	1.0
6	Chronic lower respiratory diseases (J40-J47)	27	5.1	0.9
7	Influenza and pneumonia (J09-J18)	23	4.3	0.8
8	Septicemia (A40-A41)	12	2.3	*
8	Anemias (D50-D64)	12	2.3	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
11	Cerebrovascular diseases (I60-I69)	7	1.3	*
13	Diabetes mellitus (E10-E14)	6	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
...	All other causes (Residual)	119	22.4	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,617	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	23.0	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	17.1	8.5
3	Malignant neoplasms (C00-C97)	126	7.8	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	7.2	3.6
5	Pregnancy, childbirth and the puerperium (O00-O99)	88	5.4	2.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	4.5	2.2
7	Influenza and pneumonia (J09-J18)	45	2.8	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.5	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.2	1.1
10	Chronic lower respiratory diseases (J40-J47)	32	2.0	1.0
11	Anemias (D50-D64)	30	1.9	0.9
12	Diabetes mellitus (E10-E14)	25	1.5	0.8
13	Septicemia (A40-A41)	19	1.2	*
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.1	*
...	All other causes (Residual)	303	18.7	9.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,919	100.0	101.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	13.0	13.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	376	12.9	13.0
3	Malignant neoplasms (C00-C97)	346	11.9	12.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	9.5	9.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	254	8.7	8.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	129	4.4	4.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	3.0	3.0
8	Influenza and pneumonia (J09-J18)	81	2.8	2.8
9	Diabetes mellitus (E10-E14)	80	2.7	2.8
10	Cerebrovascular diseases (I60-I69)	64	2.2	2.2
11	Septicemia (A40-A41)	57	2.0	2.0
12	Anemias (D50-D64)	51	1.7	1.8
13	Chronic lower respiratory diseases (J40-J47)	48	1.6	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.5	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.3	1.3
...	All other causes (Residual)	610	20.9	21.2
Non-Hispanic, black, female, 35-44 years				
...	All causes	6,125	100.0	220.0
1	Malignant neoplasms (C00-C97)	1,313	21.4	47.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,024	16.7	36.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	524	8.6	18.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	498	8.1	17.9
5	Cerebrovascular diseases (I60-I69)	255	4.2	9.2
6	Diabetes mellitus (E10-E14)	197	3.2	7.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	3.0	6.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.5	5.5
9	Septicemia (A40-A41)	141	2.3	5.1
10	Influenza and pneumonia (J09-J18)	135	2.2	4.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	1.4	3.2
12	Chronic lower respiratory diseases (J40-J47)	73	1.2	2.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.2	2.6
14	Anemias (D50-D64)	69	1.1	2.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	1.1	2.5
...	All other causes (Residual)	1,333	21.8	47.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	14,996	100.0	527.6
1	Malignant neoplasms (C00-C97)	4,372	29.2	153.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,978	19.9	104.8
3	Cerebrovascular diseases (I60-I69)	864	5.8	30.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	788	5.3	27.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	632	4.2	22.2
6	Diabetes mellitus (E10-E14)	556	3.7	19.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	398	2.7	14.0
8	Chronic lower respiratory diseases (J40-J47)	369	2.5	13.0
9	Septicemia (A40-A41)	310	2.1	10.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	270	1.8	9.5
11	Influenza and pneumonia (J09-J18)	244	1.6	8.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	228	1.5	8.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	1.0	5.4
14	Viral hepatitis (B15-B19)	122	0.8	4.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	0.6	3.2
...	All other causes (Residual)	2,620	17.5	92.2
Non-Hispanic, black, female, 55-64 years				
...	All causes	20,562	100.0	1,039.4
1	Malignant neoplasms (C00-C97)	6,845	33.3	346.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,771	23.2	241.2
3	Diabetes mellitus (E10-E14)	1,135	5.5	57.4
4	Cerebrovascular diseases (I60-I69)	1,117	5.4	56.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	658	3.2	33.3
6	Chronic lower respiratory diseases (J40-J47)	614	3.0	31.0
7	Septicemia (A40-A41)	511	2.5	25.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	2.3	23.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	361	1.8	18.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	290	1.4	14.7
11	Influenza and pneumonia (J09-J18)	287	1.4	14.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	277	1.3	14.0
13	Viral hepatitis (B15-B19)	142	0.7	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	5.3
15	Pneumonitis due to solids and liquids (J69)	76	0.4	3.8
...	All other causes (Residual)	2,902	14.1	146.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	23,567	100.0	2,126.3
1	Malignant neoplasms (C00-C97)	7,353	31.2	663.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,764	24.5	520.0
3	Diabetes mellitus (E10-E14)	1,417	6.0	127.8
4	Cerebrovascular diseases (I60-I69)	1,414	6.0	127.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	995	4.2	89.8
6	Chronic lower respiratory diseases (J40-J47)	921	3.9	83.1
7	Septicemia (A40-A41)	664	2.8	59.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	461	2.0	41.6
9	Influenza and pneumonia (J09-J18)	343	1.5	30.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	1.2	24.6
11	Alzheimer's disease (G30)	198	0.8	17.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	133	0.6	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.5	9.9
14	Pneumonitis due to solids and liquids (J69)	105	0.4	9.5
15	Aortic aneurysm and dissection (I71)	78	0.3	7.0
...	All other causes (Residual)	3,338	14.2	301.2
Non-Hispanic, black, female, 75-84 years				
...	All causes	30,995	100.0	4,543.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,387	27.1	1,229.3
2	Malignant neoplasms (C00-C97)	6,789	21.9	995.1
3	Cerebrovascular diseases (I60-I69)	2,379	7.7	348.7
4	Diabetes mellitus (E10-E14)	1,663	5.4	243.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,311	4.2	192.2
6	Alzheimer's disease (G30)	1,067	3.4	156.4
7	Chronic lower respiratory diseases (J40-J47)	1,044	3.4	153.0
8	Septicemia (A40-A41)	850	2.7	124.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	671	2.2	98.4
10	Influenza and pneumonia (J09-J18)	643	2.1	94.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	1.0	45.9
12	Pneumonitis due to solids and liquids (J69)	208	0.7	30.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	159	0.5	23.3
14	Parkinson's disease (G20-G21)	136	0.4	19.9
15	Atherosclerosis (I70)	111	0.4	16.3
...	All other causes (Residual)	5,264	17.0	771.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	33,543	100.0	12,954.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,360	30.9	4,001.1
2	Malignant neoplasms (C00-C97)	3,648	10.9	1,408.9
3	Cerebrovascular diseases (I60-I69)	2,687	8.0	1,037.7
4	Alzheimer's disease (G30)	2,312	6.9	892.9
5	Diabetes mellitus (E10-E14)	1,324	3.9	511.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,206	3.6	465.8
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	926	2.8	357.6
8	Influenza and pneumonia (J09-J18)	899	2.7	347.2
9	Chronic lower respiratory diseases (J40-J47)	840	2.5	324.4
10	Septicemia (A40-A41)	793	2.4	306.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	1.2	154.1
12	Pneumonitis due to solids and liquids (J69)	290	0.9	112.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	174	0.5	67.2
14	Atherosclerosis (I70)	155	0.5	59.9
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	127	0.4	49.0
...	All other causes (Residual)	7,403	22.1	2,859.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable