

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	139,241	100.0	296.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,951	20.8	61.7
2	Malignant neoplasms (C00-C97)	28,851	20.7	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,080	8.0	23.6
4	Cerebrovascular diseases (I60-I69)	7,121	5.1	15.2
5	Diabetes mellitus (E10-E14)	6,544	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,091	2.9	8.7
7	Chronic lower respiratory diseases (J40-J47)	3,949	2.8	8.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,331	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,176	2.3	6.8
10	Alzheimer's disease (G30)	3,005	2.2	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,903	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,873	2.1	6.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,345	1.7	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,011	1.4	4.3
15	Septicemia (A40-A41)	2,004	1.4	4.3
...	All other causes (Residual)	27,006	19.4	57.5
Hispanic, both sexes, 1-4 years				
...	All causes	1,029	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	32.0	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.6	3.1
3	Malignant neoplasms (C00-C97)	105	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	7.7	1.9
5	Influenza and pneumonia (J09-J18)	31	3.0	0.7
6	Septicemia (A40-A41)	28	2.7	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	2.6	0.6
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Anemias (D50-D64)	4	0.4	*
13	Meningitis (G00,G03)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	240	23.3	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,020	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	29.8	3.6
2	Malignant neoplasms (C00-C97)	186	18.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	6.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	6.2	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.4
6	Cerebrovascular diseases (I60-I69)	26	2.5	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	2.2	0.3
8	Influenza and pneumonia (J09-J18)	16	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
11	Septicemia (A40-A41)	11	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Diabetes mellitus (E10-E14)	6	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
...	All other causes (Residual)	232	22.7	2.7
Hispanic, both sexes, 15-24 years				
...	All causes	5,203	100.0	69.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,152	41.4	28.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,217	23.4	16.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	511	9.8	6.8
4	Malignant neoplasms (C00-C97)	350	6.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.4	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.2	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	39	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	33	0.6	0.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.4
10	Influenza and pneumonia (J09-J18)	29	0.6	0.4
11	Diabetes mellitus (E10-E14)	21	0.4	0.3
11	Legal intervention (Y35,Y89.0)	21	0.4	0.3
13	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
14	Septicemia (A40-A41)	19	0.4	*
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	551	10.6	7.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,017	100.0	98.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,093	27.3	26.9
2	Malignant neoplasms (C00-C97)	2,143	14.3	14.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,476	9.8	9.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,247	8.3	8.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,050	7.0	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	591	3.9	3.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	580	3.9	3.8
8	Cerebrovascular diseases (I60-I69)	418	2.8	2.8
9	Diabetes mellitus (E10-E14)	287	1.9	1.9
10	Influenza and pneumonia (J09-J18)	157	1.0	1.0
11	Septicemia (A40-A41)	130	0.9	0.9
12	Viral hepatitis (B15-B19)	115	0.8	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.8	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	0.7	0.7
15	Pregnancy, childbirth and the puerperium (O00-O99)	106	0.7	0.7
...	All other causes (Residual)	2,398	16.0	15.8
Hispanic, both sexes, 45-64 years				
...	All causes	32,872	100.0	419.5
1	Malignant neoplasms (C00-C97)	9,324	28.4	119.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,178	18.8	78.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,404	7.3	30.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,208	6.7	28.2
5	Diabetes mellitus (E10-E14)	1,795	5.5	22.9
6	Cerebrovascular diseases (I60-I69)	1,433	4.4	18.3
7	Viral hepatitis (B15-B19)	820	2.5	10.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	683	2.1	8.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	667	2.0	8.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	559	1.7	7.1
11	Chronic lower respiratory diseases (J40-J47)	491	1.5	6.3
12	Influenza and pneumonia (J09-J18)	475	1.4	6.1
13	Septicemia (A40-A41)	443	1.3	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	371	1.1	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	307	0.9	3.9
...	All other causes (Residual)	4,714	14.3	60.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	78,201	100.0	2,938.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,242	27.2	798.2
2	Malignant neoplasms (C00-C97)	16,725	21.4	628.4
3	Cerebrovascular diseases (I60-I69)	5,169	6.6	194.2
4	Diabetes mellitus (E10-E14)	4,434	5.7	166.6
5	Chronic lower respiratory diseases (J40-J47)	3,324	4.3	124.9
6	Alzheimer's disease (G30)	2,958	3.8	111.1
7	Influenza and pneumonia (J09-J18)	2,422	3.1	91.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,061	2.6	77.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,611	2.1	60.5
10	Septicemia (A40-A41)	1,304	1.7	49.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,285	1.6	48.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,145	1.5	43.0
13	Parkinson's disease (G20-G21)	810	1.0	30.4
14	Pneumonitis due to solids and liquids (J69)	535	0.7	20.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	434	0.6	16.3
...	All other causes (Residual)	12,742	16.3	478.8
Hispanic, male, all ages				
...	All causes	76,861	100.0	316.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,498	20.2	63.9
2	Malignant neoplasms (C00-C97)	15,283	19.9	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,363	10.9	34.5
4	Cerebrovascular diseases (I60-I69)	3,370	4.4	13.9
5	Diabetes mellitus (E10-E14)	3,314	4.3	13.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,850	3.7	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,777	3.6	11.4
8	Chronic lower respiratory diseases (J40-J47)	2,019	2.6	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,955	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,645	2.1	6.8
11	Influenza and pneumonia (J09-J18)	1,544	2.0	6.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,456	1.9	6.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,061	1.4	4.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,037	1.3	4.3
15	Septicemia (A40-A41)	973	1.3	4.0
...	All other causes (Residual)	13,716	17.8	56.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	587	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	36.1	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	11.1	3.0
3	Malignant neoplasms (C00-C97)	52	8.9	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.1	*
6	Septicemia (A40-A41)	14	2.4	*
6	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.7	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Anemias (D50-D64)	2	0.3	*
12	Meningitis (G00,G03)	2	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	145	24.7	6.8
Hispanic, male, 5-14 years				
...	All causes	548	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	34.7	4.4
2	Malignant neoplasms (C00-C97)	95	17.3	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.5	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	5.3	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	2.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.0	*
8	Influenza and pneumonia (J09-J18)	7	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
10	Septicemia (A40-A41)	4	0.7	*
10	Anemias (D50-D64)	4	0.7	*
12	Diseases of appendix (K35-K38)	3	0.5	*
13	Meningococcal infection (A39)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	120	21.9	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,079	100.0	103.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,740	42.7	44.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,087	26.6	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	10.4	10.8
4	Malignant neoplasms (C00-C97)	217	5.3	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	2.3	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.0
7	Legal intervention (Y35,Y89.0)	21	0.5	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.4	*
8	Cerebrovascular diseases (I60-I69)	17	0.4	*
10	Influenza and pneumonia (J09-J18)	16	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.3	*
12	Diabetes mellitus (E10-E14)	13	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
14	Septicemia (A40-A41)	8	0.2	*
14	Anemias (D50-D64)	8	0.2	*
...	All other causes (Residual)	351	8.6	8.9
Hispanic, male, 25-44 years				
...	All causes	10,763	100.0	130.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,397	31.6	41.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,250	11.6	15.2
3	Malignant neoplasms (C00-C97)	994	9.2	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	936	8.7	11.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	887	8.2	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	460	4.3	5.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	436	4.1	5.3
8	Cerebrovascular diseases (I60-I69)	269	2.5	3.3
9	Diabetes mellitus (E10-E14)	189	1.8	2.3
10	Influenza and pneumonia (J09-J18)	93	0.9	1.1
11	Viral hepatitis (B15-B19)	81	0.8	1.0
12	Septicemia (A40-A41)	75	0.7	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	44	0.4	0.5
15	Legal intervention (Y35,Y89.0)	44	0.4	0.5
...	All other causes (Residual)	1,471	13.7	17.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	20,930	100.0	533.9
1	Malignant neoplasms (C00-C97)	4,932	23.6	125.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,409	21.1	112.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,867	8.9	47.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,712	8.2	43.7
5	Diabetes mellitus (E10-E14)	1,059	5.1	27.0
6	Cerebrovascular diseases (I60-I69)	869	4.2	22.2
7	Viral hepatitis (B15-B19)	594	2.8	15.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	518	2.5	13.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	451	2.2	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	1.8	9.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	1.4	7.4
12	Influenza and pneumonia (J09-J18)	288	1.4	7.3
12	Chronic lower respiratory diseases (J40-J47)	288	1.4	7.3
14	Septicemia (A40-A41)	250	1.2	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	0.8	4.4
...	All other causes (Residual)	2,858	13.7	72.9
Hispanic, male, 65 years and over				
...	All causes	36,666	100.0	3,215.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,972	27.2	874.5
2	Malignant neoplasms (C00-C97)	8,982	24.5	787.7
3	Cerebrovascular diseases (I60-I69)	2,180	5.9	191.2
4	Diabetes mellitus (E10-E14)	2,051	5.6	179.9
5	Chronic lower respiratory diseases (J40-J47)	1,659	4.5	145.5
6	Influenza and pneumonia (J09-J18)	1,103	3.0	96.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	990	2.7	86.8
8	Alzheimer's disease (G30)	944	2.6	82.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	843	2.3	73.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	673	1.8	59.0
11	Septicemia (A40-A41)	585	1.6	51.3
12	Parkinson's disease (G20-G21)	453	1.2	39.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	453	1.2	39.7
14	Pneumonitis due to solids and liquids (J69)	262	0.7	23.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.5	17.5
...	All other causes (Residual)	5,316	14.5	466.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	62,380	100.0	274.9
1	Malignant neoplasms (C00-C97)	13,568	21.8	59.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,453	21.6	59.3
3	Cerebrovascular diseases (I60-I69)	3,751	6.0	16.5
4	Diabetes mellitus (E10-E14)	3,230	5.2	14.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,717	4.4	12.0
6	Alzheimer's disease (G30)	2,039	3.3	9.0
7	Chronic lower respiratory diseases (J40-J47)	1,930	3.1	8.5
8	Influenza and pneumonia (J09-J18)	1,632	2.6	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,447	2.3	6.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,241	2.0	5.5
11	Certain conditions originating in the perinatal period (P00-P96)	1,228	2.0	5.4
12	Septicemia (A40-A41)	1,031	1.7	4.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	950	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	856	1.4	3.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	554	0.9	2.4
...	All other causes (Residual)	12,753	20.4	56.2
Hispanic, female, 1-4 years				
...	All causes	442	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	26.5	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	14.7	3.2
3	Malignant neoplasms (C00-C97)	53	12.0	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	10.0	2.2
5	Influenza and pneumonia (J09-J18)	17	3.8	*
6	Septicemia (A40-A41)	14	3.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.0	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	89	20.1	4.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	472	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	24.2	2.8
2	Malignant neoplasms (C00-C97)	91	19.3	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	8.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	0.5
6	Cerebrovascular diseases (I60-I69)	15	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	10	2.1	*
9	Influenza and pneumonia (J09-J18)	9	1.9	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.9	*
11	Septicemia (A40-A41)	7	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.1	*
14	Diabetes mellitus (E10-E14)	4	0.8	*
14	Meningitis (G00,G03)	4	0.8	*
...	All other causes (Residual)	105	22.2	2.5
Hispanic, female, 15-24 years				
...	All causes	1,124	100.0	31.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	36.7	11.5
2	Malignant neoplasms (C00-C97)	133	11.8	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	11.6	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	7.7	2.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	39	3.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.8	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	2.2	0.7
8	Cerebrovascular diseases (I60-I69)	16	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.3	*
10	Influenza and pneumonia (J09-J18)	13	1.2	*
11	Septicemia (A40-A41)	11	1.0	*
12	Anemias (D50-D64)	9	0.8	*
13	Diabetes mellitus (E10-E14)	8	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
...	All other causes (Residual)	182	16.2	5.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,254	100.0	61.2
1	Malignant neoplasms (C00-C97)	1,149	27.0	16.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	696	16.4	10.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	311	7.3	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	226	5.3	3.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	3.8	2.3
6	Cerebrovascular diseases (I60-I69)	149	3.5	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	144	3.4	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	131	3.1	1.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	106	2.5	1.5
10	Diabetes mellitus (E10-E14)	98	2.3	1.4
11	Influenza and pneumonia (J09-J18)	64	1.5	0.9
12	Septicemia (A40-A41)	55	1.3	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.1	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.0	0.6
15	Chronic lower respiratory diseases (J40-J47)	41	1.0	0.6
...	All other causes (Residual)	832	19.6	12.0
Hispanic, female, 45-64 years				
...	All causes	11,942	100.0	305.0
1	Malignant neoplasms (C00-C97)	4,392	36.8	112.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,769	14.8	45.2
3	Diabetes mellitus (E10-E14)	736	6.2	18.8
4	Cerebrovascular diseases (I60-I69)	564	4.7	14.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	4.5	13.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	496	4.2	12.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	309	2.6	7.9
8	Viral hepatitis (B15-B19)	226	1.9	5.8
9	Chronic lower respiratory diseases (J40-J47)	203	1.7	5.2
10	Septicemia (A40-A41)	193	1.6	4.9
11	Influenza and pneumonia (J09-J18)	187	1.6	4.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	149	1.2	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	1.1	3.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	0.9	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	0.7	2.1
...	All other causes (Residual)	1,856	15.5	47.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	41,535	100.0	2,730.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,270	27.1	741.0
2	Malignant neoplasms (C00-C97)	7,743	18.6	509.1
3	Cerebrovascular diseases (I60-I69)	2,989	7.2	196.5
4	Diabetes mellitus (E10-E14)	2,383	5.7	156.7
5	Alzheimer's disease (G30)	2,014	4.8	132.4
6	Chronic lower respiratory diseases (J40-J47)	1,665	4.0	109.5
7	Influenza and pneumonia (J09-J18)	1,319	3.2	86.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,071	2.6	70.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	1.8	50.5
10	Septicemia (A40-A41)	719	1.7	47.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	692	1.7	45.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	612	1.5	40.2
13	Parkinson's disease (G20-G21)	357	0.9	23.5
14	Pneumonitis due to solids and liquids (J69)	273	0.7	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	234	0.6	15.4
...	All other causes (Residual)	7,426	17.9	488.2
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,981,034	100.0	982.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,096	25.4	249.4
2	Malignant neoplasms (C00-C97)	457,084	23.1	226.6
3	Chronic lower respiratory diseases (J40-J47)	126,146	6.4	62.5
4	Cerebrovascular diseases (I60-I69)	106,134	5.4	52.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,722	4.8	47.0
6	Alzheimer's disease (G30)	73,219	3.7	36.3
7	Diabetes mellitus (E10-E14)	49,388	2.5	24.5
8	Influenza and pneumonia (J09-J18)	45,780	2.3	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,439	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,281	1.5	15.0
11	Septicemia (A40-A41)	26,692	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,210	1.1	11.0
13	Parkinson's disease (G20-G21)	18,474	0.9	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,351	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	14,135	0.7	7.0
...	All other causes (Residual)	359,883	18.2	178.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,373	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	788	33.2	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	256	10.8	2.8
3	Malignant neoplasms (C00-C97)	225	9.5	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.1	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.4	0.9
6	Influenza and pneumonia (J09-J18)	60	2.5	0.7
7	Septicemia (A40-A41)	41	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	30	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	12	0.5	*
13	Anemias (D50-D64)	10	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	612	25.8	6.7
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	3,070	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,091	35.5	4.7
2	Malignant neoplasms (C00-C97)	509	16.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	5.5	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	3.8	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	3.4	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.6	0.2
7	Influenza and pneumonia (J09-J18)	49	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	47	1.5	0.2
10	Septicemia (A40-A41)	32	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	29	0.9	0.1
12	Diabetes mellitus (E10-E14)	18	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	18	0.6	*
14	Anemias (D50-D64)	13	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (Residual)	666	21.7	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	18,560	100.0	70.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,911	53.4	37.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,025	16.3	11.5
3	Malignant neoplasms (C00-C97)	949	5.1	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	4.9	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	511	2.8	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	1.4	1.0
7	Influenza and pneumonia (J09-J18)	125	0.7	0.5
8	Diabetes mellitus (E10-E14)	110	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	107	0.6	0.4
10	Septicemia (A40-A41)	73	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	63	0.3	0.2
12	Chronic lower respiratory diseases (J40-J47)	62	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	28	0.2	0.1
...	All other causes (Residual)	2,327	12.5	8.9
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	73,943	100.0	142.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,071	29.8	42.6
2	Malignant neoplasms (C00-C97)	10,330	14.0	20.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,548	12.9	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,913	12.1	17.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,142	2.9	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,920	2.6	3.7
7	Diabetes mellitus (E10-E14)	1,349	1.8	2.6
8	Cerebrovascular diseases (I60-I69)	1,253	1.7	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	990	1.3	1.9
10	Influenza and pneumonia (J09-J18)	756	1.0	1.5
11	Septicemia (A40-A41)	706	1.0	1.4
12	Chronic lower respiratory diseases (J40-J47)	641	0.9	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	569	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	0.7	0.9
15	Viral hepatitis (B15-B19)	284	0.4	0.5
...	All other causes (Residual)	11,982	16.2	23.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	350,678	100.0	609.4
1	Malignant neoplasms (C00-C97)	116,532	33.2	202.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,778	21.3	129.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,633	7.3	44.5
4	Chronic lower respiratory diseases (J40-J47)	15,586	4.4	27.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,335	3.5	21.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	12,244	3.5	21.3
7	Diabetes mellitus (E10-E14)	10,900	3.1	18.9
8	Cerebrovascular diseases (I60-I69)	9,855	2.8	17.1
9	Septicemia (A40-A41)	4,628	1.3	8.0
10	Influenza and pneumonia (J09-J18)	4,136	1.2	7.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,769	1.1	6.5
12	Viral hepatitis (B15-B19)	3,631	1.0	6.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,948	0.6	3.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,773	0.5	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,608	0.5	2.8
...	All other causes (Residual)	51,322	14.6	89.2
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,519,812	100.0	4,843.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	418,513	27.5	1,333.7
2	Malignant neoplasms (C00-C97)	328,499	21.6	1,046.9
3	Chronic lower respiratory diseases (J40-J47)	109,795	7.2	349.9
4	Cerebrovascular diseases (I60-I69)	94,771	6.2	302.0
5	Alzheimer's disease (G30)	72,486	4.8	231.0
6	Influenza and pneumonia (J09-J18)	40,547	2.7	129.2
7	Diabetes mellitus (E10-E14)	37,006	2.4	117.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,543	2.3	110.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,068	2.0	99.0
10	Septicemia (A40-A41)	21,110	1.4	67.3
11	Parkinson's disease (G20-G21)	18,072	1.2	57.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,176	1.1	51.6
13	Pneumonitis due to solids and liquids (J69)	12,877	0.8	41.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,217	0.7	32.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8,025	0.5	25.6
...	All other causes (Residual)	266,107	17.5	848.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	969,288	100.0	978.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252,572	26.1	254.9
2	Malignant neoplasms (C00-C97)	238,846	24.6	241.1
3	Chronic lower respiratory diseases (J40-J47)	59,284	6.1	59.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,173	6.1	59.7
5	Cerebrovascular diseases (I60-I69)	41,092	4.2	41.5
6	Diabetes mellitus (E10-E14)	25,303	2.6	25.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,835	2.5	24.1
8	Alzheimer's disease (G30)	21,780	2.2	22.0
9	Influenza and pneumonia (J09-J18)	20,516	2.1	20.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,535	1.8	17.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,452	1.5	14.6
12	Septicemia (A40-A41)	12,092	1.2	12.2
13	Parkinson's disease (G20-G21)	10,845	1.1	10.9
14	Pneumonitis due to solids and liquids (J69)	7,369	0.8	7.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,088	0.7	7.2
...	All other causes (Residual)	157,506	16.2	159.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,365	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	479	35.1	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.0	2.9
3	Malignant neoplasms (C00-C97)	127	9.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	7.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	2.9	0.8
6	Influenza and pneumonia (J09-J18)	27	2.0	0.6
7	Septicemia (A40-A41)	17	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
9	Cerebrovascular diseases (I60-I69)	13	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Anemias (D50-D64)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	367	26.9	7.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,806	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	705	39.0	5.9
2	Malignant neoplasms (C00-C97)	264	14.6	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	6.0	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	4.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.4	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.2
8	Influenza and pneumonia (J09-J18)	27	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	21	1.2	0.2
10	Septicemia (A40-A41)	17	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
12	Diabetes mellitus (E10-E14)	11	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
14	Anemias (D50-D64)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	370	20.5	3.1
Non-Hispanic, white, male, 15-24 years				
...	All causes	13,575	100.0	100.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,382	54.4	54.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,530	18.6	18.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	689	5.1	5.1
4	Malignant neoplasms (C00-C97)	584	4.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	351	2.6	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.2	1.2
7	Influenza and pneumonia (J09-J18)	84	0.6	0.6
8	Diabetes mellitus (E10-E14)	69	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	65	0.5	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	34	0.3	0.3
12	Septicemia (A40-A41)	29	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
...	All other causes (Residual)	1,492	11.0	11.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	48,140	100.0	185.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,890	33.0	61.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,417	15.4	28.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,296	13.1	24.2
4	Malignant neoplasms (C00-C97)	4,726	9.8	18.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,505	3.1	5.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,239	2.6	4.8
7	Diabetes mellitus (E10-E14)	827	1.7	3.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	778	1.6	3.0
9	Cerebrovascular diseases (I60-I69)	659	1.4	2.5
10	Influenza and pneumonia (J09-J18)	405	0.8	1.6
11	Septicemia (A40-A41)	357	0.7	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	294	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	286	0.6	1.1
15	Aortic aneurysm and dissection (I71)	182	0.4	0.7
...	All other causes (Residual)	6,959	14.5	26.8
Non-Hispanic, white, male, 45-64 years				
...	All causes	215,621	100.0	760.5
1	Malignant neoplasms (C00-C97)	63,134	29.3	222.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,832	25.0	189.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,304	8.0	61.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,371	4.3	33.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,613	4.0	30.4
6	Chronic lower respiratory diseases (J40-J47)	8,020	3.7	28.3
7	Diabetes mellitus (E10-E14)	6,751	3.1	23.8
8	Cerebrovascular diseases (I60-I69)	5,468	2.5	19.3
9	Viral hepatitis (B15-B19)	2,593	1.2	9.1
10	Septicemia (A40-A41)	2,459	1.1	8.7
11	Influenza and pneumonia (J09-J18)	2,347	1.1	8.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,213	1.0	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,493	0.7	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,226	0.6	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,112	0.5	3.9
...	All other causes (Residual)	29,685	13.8	104.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	681,707	100.0	5,090.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	191,887	28.1	1,432.9
2	Malignant neoplasms (C00-C97)	169,984	24.9	1,269.3
3	Chronic lower respiratory diseases (J40-J47)	50,901	7.5	380.1
4	Cerebrovascular diseases (I60-I69)	34,836	5.1	260.1
5	Alzheimer's disease (G30)	21,452	3.1	160.2
6	Diabetes mellitus (E10-E14)	17,643	2.6	131.7
7	Influenza and pneumonia (J09-J18)	17,565	2.6	131.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,026	2.5	127.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,972	2.2	111.8
10	Parkinson's disease (G20-G21)	10,589	1.6	79.1
11	Septicemia (A40-A41)	9,165	1.3	68.4
12	Pneumonitis due to solids and liquids (J69)	6,605	1.0	49.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,701	0.8	42.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,232	0.8	39.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,590	0.7	34.3
...	All other causes (Residual)	103,559	15.2	773.3
Non-Hispanic, white, female, all ages				
...	All causes	1,011,746	100.0	985.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,524	24.8	244.0
2	Malignant neoplasms (C00-C97)	218,238	21.6	212.6
3	Chronic lower respiratory diseases (J40-J47)	66,862	6.6	65.1
4	Cerebrovascular diseases (I60-I69)	65,042	6.4	63.4
5	Alzheimer's disease (G30)	51,439	5.1	50.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,549	3.5	34.6
7	Influenza and pneumonia (J09-J18)	25,264	2.5	24.6
8	Diabetes mellitus (E10-E14)	24,085	2.4	23.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,904	1.8	17.4
10	Septicemia (A40-A41)	14,600	1.4	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,263	1.1	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,758	0.8	7.6
13	Parkinson's disease (G20-G21)	7,629	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,766	0.7	6.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,446	0.6	6.3
...	All other causes (Residual)	202,377	20.0	197.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,008	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	30.7	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	11.9	2.7
3	Malignant neoplasms (C00-C97)	98	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.1	0.9
6	Influenza and pneumonia (J09-J18)	33	3.3	0.7
7	Septicemia (A40-A41)	24	2.4	0.5
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Meningococcal infection (A39)	5	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (Residual)	250	24.8	5.6
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,264	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	30.5	3.4
2	Malignant neoplasms (C00-C97)	245	19.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	6.6	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	3.8	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	3.4	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.2	0.4
7	Cerebrovascular diseases (I60-I69)	26	2.1	0.2
8	Influenza and pneumonia (J09-J18)	22	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.2
10	Septicemia (A40-A41)	15	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
13	Diabetes mellitus (E10-E14)	7	0.6	*
14	Anemias (D50-D64)	6	0.5	*
15	Meningitis (G00,G03)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	290	22.9	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,985	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,529	50.7	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	495	9.9	3.9
3	Malignant neoplasms (C00-C97)	365	7.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	225	4.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	3.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	2.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.3	0.5
8	Septicemia (A40-A41)	44	0.9	0.3
9	Cerebrovascular diseases (I60-I69)	42	0.8	0.3
10	Diabetes mellitus (E10-E14)	41	0.8	0.3
10	Influenza and pneumonia (J09-J18)	41	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	28	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.2	*
...	All other causes (Residual)	797	16.0	6.2
Non-Hispanic, white, female, 25-44 years				
...	All causes	25,803	100.0	100.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,181	24.0	24.0
2	Malignant neoplasms (C00-C97)	5,604	21.7	21.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,617	10.1	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,131	8.3	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	681	2.6	2.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	637	2.5	2.5
7	Cerebrovascular diseases (I60-I69)	594	2.3	2.3
8	Diabetes mellitus (E10-E14)	522	2.0	2.0
9	Influenza and pneumonia (J09-J18)	351	1.4	1.4
10	Septicemia (A40-A41)	349	1.4	1.4
11	Chronic lower respiratory diseases (J40-J47)	347	1.3	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	249	1.0	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	240	0.9	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	212	0.8	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	0.8	0.8
...	All other causes (Residual)	4,885	18.9	19.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	135,057	100.0	462.6
1	Malignant neoplasms (C00-C97)	53,398	39.5	182.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,946	15.5	71.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,329	6.2	28.5
4	Chronic lower respiratory diseases (J40-J47)	7,566	5.6	25.9
5	Cerebrovascular diseases (I60-I69)	4,387	3.2	15.0
6	Diabetes mellitus (E10-E14)	4,149	3.1	14.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,631	2.7	12.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,964	2.2	10.2
9	Septicemia (A40-A41)	2,169	1.6	7.4
10	Influenza and pneumonia (J09-J18)	1,789	1.3	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,556	1.2	5.3
12	Viral hepatitis (B15-B19)	1,038	0.8	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	722	0.5	2.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	674	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	575	0.4	2.0
...	All other causes (Residual)	21,164	15.7	72.5
Non-Hispanic, white, female, 65 years and over				
...	All causes	838,105	100.0	4,659.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	226,626	27.0	1,259.9
2	Malignant neoplasms (C00-C97)	158,515	18.9	881.2
3	Cerebrovascular diseases (I60-I69)	59,935	7.2	333.2
4	Chronic lower respiratory diseases (J40-J47)	58,894	7.0	327.4
5	Alzheimer's disease (G30)	51,034	6.1	283.7
6	Influenza and pneumonia (J09-J18)	22,982	2.7	127.8
7	Diabetes mellitus (E10-E14)	19,363	2.3	107.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,517	2.1	97.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,096	1.9	89.5
10	Septicemia (A40-A41)	11,945	1.4	66.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,475	1.2	58.2
12	Parkinson's disease (G20-G21)	7,483	0.9	41.6
13	Pneumonitis due to solids and liquids (J69)	6,272	0.7	34.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,985	0.6	27.7
15	Atherosclerosis (I70)	4,066	0.5	22.6
...	All other causes (Residual)	161,917	19.3	900.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	285,522	100.0	745.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,918	24.5	182.5
2	Malignant neoplasms (C00-C97)	63,279	22.2	165.2
3	Cerebrovascular diseases (I60-I69)	16,527	5.8	43.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,215	4.3	31.9
5	Diabetes mellitus (E10-E14)	11,934	4.2	31.1
6	Chronic lower respiratory diseases (J40-J47)	8,679	3.0	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,564	3.0	22.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,206	2.9	21.4
9	Septicemia (A40-A41)	6,359	2.2	16.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,686	2.0	14.8
11	Influenza and pneumonia (J09-J18)	5,396	1.9	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,028	1.8	13.1
13	Alzheimer's disease (G30)	5,024	1.8	13.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,627	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,464	0.9	6.4
...	All other causes (Residual)	51,616	18.1	134.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,076	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	26.2	11.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	14.0	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	10.4	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.9	2.6
5	Malignant neoplasms (C00-C97)	51	4.7	2.1
6	Influenza and pneumonia (J09-J18)	44	4.1	1.8
7	Chronic lower respiratory diseases (J40-J47)	22	2.0	0.9
8	Septicemia (A40-A41)	21	2.0	0.9
9	Anemias (D50-D64)	18	1.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.4	*
11	Cerebrovascular diseases (I60-I69)	13	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	258	24.0	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,258	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	30.2	6.3
2	Malignant neoplasms (C00-C97)	148	11.8	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	10.6	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	5.8	1.2
5	Chronic lower respiratory diseases (J40-J47)	72	5.7	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.1	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	2.4	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	19	1.5	*
10	Influenza and pneumonia (J09-J18)	16	1.3	*
11	Anemias (D50-D64)	14	1.1	*
12	Septicemia (A40-A41)	13	1.0	*
13	Diabetes mellitus (E10-E14)	12	1.0	*
14	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	252	20.0	4.2
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,092	100.0	109.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,992	42.2	46.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,480	20.9	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	501	7.1	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	5.2	5.7
5	Malignant neoplasms (C00-C97)	280	3.9	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	113	1.6	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	1.5	1.6
8	Chronic lower respiratory diseases (J40-J47)	75	1.1	1.2
9	Diabetes mellitus (E10-E14)	67	0.9	1.0
10	Anemias (D50-D64)	64	0.9	1.0
11	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.8	0.9
12	Septicemia (A40-A41)	45	0.6	0.7
13	Cerebrovascular diseases (I60-I69)	41	0.6	0.6
13	Influenza and pneumonia (J09-J18)	41	0.6	0.6
15	Legal intervention (Y35,Y89.0)	37	0.5	0.6
...	All other causes (Residual)	825	11.6	12.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	24,695	100.0	227.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,915	15.9	36.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,642	14.7	33.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,381	13.7	31.2
4	Malignant neoplasms (C00-C97)	2,880	11.7	26.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,145	8.7	19.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	883	3.6	8.1
7	Cerebrovascular diseases (I60-I69)	755	3.1	7.0
8	Diabetes mellitus (E10-E14)	695	2.8	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	1.6	3.7
10	Septicemia (A40-A41)	360	1.5	3.3
11	Chronic lower respiratory diseases (J40-J47)	304	1.2	2.8
12	Anemias (D50-D64)	276	1.1	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	262	1.1	2.4
14	Influenza and pneumonia (J09-J18)	259	1.0	2.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	225	0.9	2.1
...	All other causes (Residual)	4,316	17.5	39.8
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	83,755	100.0	972.4
1	Malignant neoplasms (C00-C97)	23,495	28.1	272.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,495	24.5	237.9
3	Cerebrovascular diseases (I60-I69)	4,440	5.3	51.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,095	4.9	47.5
5	Diabetes mellitus (E10-E14)	3,624	4.3	42.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,084	3.7	35.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,302	2.7	26.7
8	Chronic lower respiratory diseases (J40-J47)	2,005	2.4	23.3
9	Septicemia (A40-A41)	1,773	2.1	20.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,669	2.0	19.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,425	1.7	16.5
12	Influenza and pneumonia (J09-J18)	1,159	1.4	13.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,021	1.2	11.9
14	Viral hepatitis (B15-B19)	829	1.0	9.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	486	0.6	5.6
...	All other causes (Residual)	11,853	14.2	137.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	159,439	100.0	4,888.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,893	28.2	1,376.3
2	Malignant neoplasms (C00-C97)	36,409	22.8	1,116.2
3	Cerebrovascular diseases (I60-I69)	11,219	7.0	343.9
4	Diabetes mellitus (E10-E14)	7,532	4.7	230.9
5	Chronic lower respiratory diseases (J40-J47)	6,189	3.9	189.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,780	3.6	177.2
7	Alzheimer's disease (G30)	4,956	3.1	151.9
8	Septicemia (A40-A41)	4,043	2.5	123.9
9	Influenza and pneumonia (J09-J18)	3,816	2.4	117.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,322	2.1	101.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,195	1.4	67.3
12	Pneumonitis due to solids and liquids (J69)	1,176	0.7	36.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	711	0.4	21.8
14	Parkinson's disease (G20-G21)	703	0.4	21.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	566	0.4	17.4
...	All other causes (Residual)	25,929	16.3	794.9
Non-Hispanic, black, male, all ages				
...	All causes	145,168	100.0	794.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,965	24.1	191.3
2	Malignant neoplasms (C00-C97)	32,654	22.5	178.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,282	5.7	45.3
4	Cerebrovascular diseases (I60-I69)	7,135	4.9	39.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,047	4.9	38.6
6	Diabetes mellitus (E10-E14)	5,378	3.7	29.4
7	Chronic lower respiratory diseases (J40-J47)	4,497	3.1	24.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,893	2.7	21.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,724	2.6	20.4
10	Septicemia (A40-A41)	2,841	2.0	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,608	1.8	14.3
12	Influenza and pneumonia (J09-J18)	2,542	1.8	13.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,194	1.5	12.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,721	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,653	1.1	9.0
...	All other causes (Residual)	24,034	16.6	131.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	619	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	27.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.5	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	10.3	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.7	2.8
5	Malignant neoplasms (C00-C97)	32	5.2	2.6
6	Influenza and pneumonia (J09-J18)	26	4.2	2.1
7	Septicemia (A40-A41)	11	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	144	23.3	11.6
Non-Hispanic, black, male, 5-14 years				
...	All causes	750	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	32.3	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	11.1	2.7
3	Malignant neoplasms (C00-C97)	68	9.1	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.1	1.5
5	Chronic lower respiratory diseases (J40-J47)	44	5.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.7	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	2.9	0.7
8	Cerebrovascular diseases (I60-I69)	14	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
10	Influenza and pneumonia (J09-J18)	10	1.3	*
11	Diabetes mellitus (E10-E14)	7	0.9	*
12	Septicemia (A40-A41)	6	0.8	*
12	Anemias (D50-D64)	6	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	147	19.6	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,423	100.0	164.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,672	49.3	81.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,101	20.3	33.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	417	7.7	12.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	255	4.7	7.7
5	Malignant neoplasms (C00-C97)	163	3.0	4.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.2	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	59	1.1	1.8
8	Anemias (D50-D64)	43	0.8	1.3
9	Diabetes mellitus (E10-E14)	41	0.8	1.2
9	Chronic lower respiratory diseases (J40-J47)	41	0.8	1.2
11	Legal intervention (Y35,Y89.0)	36	0.7	1.1
12	Cerebrovascular diseases (I60-I69)	23	0.4	0.7
13	Septicemia (A40-A41)	22	0.4	0.7
14	Influenza and pneumonia (J09-J18)	18	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
...	All other causes (Residual)	454	8.4	13.8
Non-Hispanic, black, male, 25-44 years				
...	All causes	15,318	100.0	295.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,192	20.8	61.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,468	16.1	47.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,414	15.8	46.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,268	8.3	24.5
5	Malignant neoplasms (C00-C97)	1,151	7.5	22.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	736	4.8	14.2
7	Diabetes mellitus (E10-E14)	377	2.5	7.3
8	Cerebrovascular diseases (I60-I69)	368	2.4	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	1.4	4.1
10	Septicemia (A40-A41)	191	1.2	3.7
11	Chronic lower respiratory diseases (J40-J47)	165	1.1	3.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	1.0	3.0
13	Influenza and pneumonia (J09-J18)	149	1.0	2.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.9	2.6
15	Anemias (D50-D64)	126	0.8	2.4
...	All other causes (Residual)	2,206	14.4	42.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	48,748	100.0	1,243.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,618	25.9	321.8
2	Malignant neoplasms (C00-C97)	12,541	25.7	319.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,901	6.0	74.0
4	Cerebrovascular diseases (I60-I69)	2,415	5.0	61.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,162	4.4	55.1
6	Diabetes mellitus (E10-E14)	1,953	4.0	49.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,249	2.6	31.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,162	2.4	29.6
9	Chronic lower respiratory diseases (J40-J47)	1,063	2.2	27.1
10	Septicemia (A40-A41)	909	1.9	23.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	841	1.7	21.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	806	1.7	20.6
13	Influenza and pneumonia (J09-J18)	667	1.4	17.0
14	Viral hepatitis (B15-B19)	586	1.2	14.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	407	0.8	10.4
...	All other causes (Residual)	6,468	13.3	164.9
Non-Hispanic, black, male, 65 years and over				
...	All causes	69,750	100.0	5,552.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,543	28.0	1,555.9
2	Malignant neoplasms (C00-C97)	18,687	26.8	1,487.7
3	Cerebrovascular diseases (I60-I69)	4,286	6.1	341.2
4	Chronic lower respiratory diseases (J40-J47)	3,162	4.5	251.7
5	Diabetes mellitus (E10-E14)	2,997	4.3	238.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,386	3.4	190.0
7	Septicemia (A40-A41)	1,648	2.4	131.2
8	Influenza and pneumonia (J09-J18)	1,639	2.3	130.5
9	Alzheimer's disease (G30)	1,363	2.0	108.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,221	1.8	97.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,187	1.7	94.5
12	Pneumonitis due to solids and liquids (J69)	574	0.8	45.7
13	Parkinson's disease (G20-G21)	370	0.5	29.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	352	0.5	28.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	310	0.4	24.7
...	All other causes (Residual)	10,025	14.4	798.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	140,354	100.0	700.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,953	24.9	174.4
2	Malignant neoplasms (C00-C97)	30,625	21.8	152.8
3	Cerebrovascular diseases (I60-I69)	9,392	6.7	46.9
4	Diabetes mellitus (E10-E14)	6,556	4.7	32.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,671	3.3	23.3
6	Chronic lower respiratory diseases (J40-J47)	4,182	3.0	20.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.6
8	Alzheimer's disease (G30)	3,630	2.6	18.1
9	Septicemia (A40-A41)	3,518	2.5	17.6
10	Influenza and pneumonia (J09-J18)	2,854	2.0	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,834	2.0	14.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,019	1.4	10.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,962	1.4	9.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.8	5.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	811	0.6	4.0
...	All other causes (Residual)	27,255	19.4	136.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	457	100.0	38.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	24.5	9.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	13.3	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	10.5	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.3	2.4
5	Malignant neoplasms (C00-C97)	19	4.2	*
6	Influenza and pneumonia (J09-J18)	18	3.9	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.4	*
8	Septicemia (A40-A41)	10	2.2	*
9	Anemias (D50-D64)	9	2.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	1.8	*
11	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Salmonella infections (A01-A02)	1	0.2	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	111	24.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	508	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	27.2	4.7
2	Malignant neoplasms (C00-C97)	80	15.7	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	9.8	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.7	1.0
5	Chronic lower respiratory diseases (J40-J47)	28	5.5	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	5.3	0.9
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.6	*
7	Anemias (D50-D64)	8	1.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	1.6	*
10	Septicemia (A40-A41)	7	1.4	*
11	Influenza and pneumonia (J09-J18)	6	1.2	*
12	Diabetes mellitus (E10-E14)	5	1.0	*
12	Cerebrovascular diseases (I60-I69)	5	1.0	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
15	Meningococcal infection (A39)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	97	19.1	3.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,669	100.0	52.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	22.7	11.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	19.2	10.0
3	Malignant neoplasms (C00-C97)	117	7.0	3.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	7.0	3.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	5.0	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.4	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	54	3.2	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.4	1.2
9	Chronic lower respiratory diseases (J40-J47)	34	2.0	1.1
10	Diabetes mellitus (E10-E14)	26	1.6	0.8
11	Septicemia (A40-A41)	23	1.4	0.7
11	Influenza and pneumonia (J09-J18)	23	1.4	0.7
13	Anemias (D50-D64)	21	1.3	0.7
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
...	All other causes (Residual)	331	19.8	10.3
Non-Hispanic, black, female, 25-44 years				
...	All causes	9,377	100.0	165.8
1	Malignant neoplasms (C00-C97)	1,729	18.4	30.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,501	16.0	26.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	913	9.7	16.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	877	9.4	15.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	450	4.8	8.0
6	Cerebrovascular diseases (I60-I69)	387	4.1	6.8
7	Diabetes mellitus (E10-E14)	318	3.4	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	2.0	3.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	174	1.9	3.1
10	Septicemia (A40-A41)	169	1.8	3.0
11	Anemias (D50-D64)	150	1.6	2.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	1.6	2.6
13	Chronic lower respiratory diseases (J40-J47)	139	1.5	2.5
14	Influenza and pneumonia (J09-J18)	110	1.2	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	106	1.1	1.9
...	All other causes (Residual)	2,024	21.6	35.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-64 years				
...	All causes	35,007	100.0	746.1
1	Malignant neoplasms (C00-C97)	10,954	31.3	233.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,877	22.5	167.9
3	Cerebrovascular diseases (I60-I69)	2,025	5.8	43.2
4	Diabetes mellitus (E10-E14)	1,671	4.8	35.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,194	3.4	25.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,053	3.0	22.4
7	Chronic lower respiratory diseases (J40-J47)	942	2.7	20.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	922	2.6	19.6
9	Septicemia (A40-A41)	864	2.5	18.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	619	1.8	13.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	507	1.4	10.8
12	Influenza and pneumonia (J09-J18)	492	1.4	10.5
13	Viral hepatitis (B15-B19)	243	0.7	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	180	0.5	3.8
15	Anemias (D50-D64)	129	0.4	2.7
...	All other causes (Residual)	5,335	15.2	113.7
Non-Hispanic, black, female, 65 years and over				
...	All causes	89,689	100.0	4,471.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,350	28.3	1,263.9
2	Malignant neoplasms (C00-C97)	17,722	19.8	883.6
3	Cerebrovascular diseases (I60-I69)	6,933	7.7	345.7
4	Diabetes mellitus (E10-E14)	4,535	5.1	226.1
5	Alzheimer's disease (G30)	3,593	4.0	179.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,394	3.8	169.2
7	Chronic lower respiratory diseases (J40-J47)	3,027	3.4	150.9
8	Septicemia (A40-A41)	2,395	2.7	119.4
9	Influenza and pneumonia (J09-J18)	2,177	2.4	108.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,101	2.3	104.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,008	1.1	50.3
12	Pneumonitis due to solids and liquids (J69)	602	0.7	30.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	401	0.4	20.0
14	Parkinson's disease (G20-G21)	333	0.4	16.6
15	Atherosclerosis (I70)	327	0.4	16.3
...	All other causes (Residual)	15,791	17.6	787.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable