

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (\*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	169,387	100.0	305.8
1	Malignant neoplasms (C00-C97)	36,447	21.5	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,021	20.1	61.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,413	7.3	22.4
4	Cerebrovascular diseases (I60-I69)	8,713	5.1	15.7
5	Diabetes mellitus (E10-E14)	7,795	4.6	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,658	3.3	10.2
7	Alzheimer's disease (G30)	4,934	2.9	8.9
8	Chronic lower respiratory diseases (J40-J47)	4,795	2.8	8.7
9	Influenza and pneumonia (J09-J18)	3,875	2.3	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,273	1.9	5.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,246	1.9	5.9
12	Septicemia (A40-A41)	2,613	1.5	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,610	1.5	4.7
14	Certain conditions originating in the perinatal period (P00-P96)	2,422	1.4	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,139	1.3	3.9
...	All other causes (residual)	34,433	20.3	62.2
Hispanic, both sexes, 1-4 years				
...	All causes	769	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	28.7	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	12.6	2.4
3	Malignant neoplasms (C00-C97)	81	10.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.8	0.7
6	Influenza and pneumonia (J09-J18)	20	2.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
10	Cerebrovascular diseases (I60-I69)	7	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Nutritional deficiencies (E40-E64)	4	0.5	*
12	Meningitis (G00,G03)	4	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (residual)	196	25.5	4.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,116	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	28.0	3.1
2	Malignant neoplasms (C00-C97)	219	19.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.2	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	5.9	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	4.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	2.4	0.3
6	Cerebrovascular diseases (I60-I69)	27	2.4	0.3
8	Chronic lower respiratory diseases (J40-J47)	24	2.2	0.2
9	Influenza and pneumonia (J09-J18)	23	2.1	0.2
10	Septicemia (A40-A41)	16	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
13	Anemias (D50-D64)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.4	*
14	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	233	20.9	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,905	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,044	41.7	21.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	812	16.6	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	14.7	7.6
4	Malignant neoplasms (C00-C97)	364	7.4	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	2.6	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.3	0.7
7	Cerebrovascular diseases (I60-I69)	34	0.7	0.4
8	Influenza and pneumonia (J09-J18)	31	0.6	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.6	0.3
10	Septicemia (A40-A41)	24	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.2
12	Legal intervention (Y35,Y89.0)	20	0.4	0.2
13	Diabetes mellitus (E10-E14)	18	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (residual)	567	11.6	6.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,743	100.0	93.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,323	27.5	25.6
2	Malignant neoplasms (C00-C97)	2,329	14.8	13.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,429	9.1	8.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,348	8.6	8.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,201	7.6	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	684	4.3	4.1
7	Cerebrovascular diseases (I60-I69)	430	2.7	2.5
8	Diabetes mellitus (E10-E14)	360	2.3	2.1
9	Influenza and pneumonia (J09-J18)	295	1.9	1.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	272	1.7	1.6
11	Septicemia (A40-A41)	176	1.1	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	117	0.7	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.7	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	0.7	0.7
15	Chronic lower respiratory diseases (J40-J47)	83	0.5	0.5
...	All other causes (residual)	2,473	15.7	14.7
Hispanic, both sexes, 45-64 years				
...	All causes	41,406	100.0	398.7
1	Malignant neoplasms (C00-C97)	11,721	28.3	112.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,502	18.1	72.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	3,166	7.6	30.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,122	7.5	30.1
5	Diabetes mellitus (E10-E14)	2,276	5.5	21.9
6	Cerebrovascular diseases (I60-I69)	1,676	4.0	16.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	826	2.0	8.0
8	Influenza and pneumonia (J09-J18)	816	2.0	7.9
9	Viral hepatitis (B15-B19)	779	1.9	7.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	721	1.7	6.9
11	Septicemia (A40-A41)	709	1.7	6.8
12	Chronic lower respiratory diseases (J40-J47)	596	1.4	5.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	530	1.3	5.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	400	1.0	3.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	361	0.9	3.5
...	All other causes (residual)	6,205	15.0	59.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	100,670	100.0	2,834.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,842	24.7	699.5
2	Malignant neoplasms (C00-C97)	21,722	21.6	611.7
3	Cerebrovascular diseases (I60-I69)	6,524	6.5	183.7
4	Diabetes mellitus (E10-E14)	5,135	5.1	144.6
5	Alzheimer's disease (G30)	4,864	4.8	137.0
6	Chronic lower respiratory diseases (J40-J47)	4,059	4.0	114.3
7	Influenza and pneumonia (J09-J18)	2,655	2.6	74.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,406	2.4	67.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,237	2.2	63.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,800	1.8	50.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,670	1.7	47.0
12	Septicemia (A40-A41)	1,644	1.6	46.3
13	Parkinson's disease (G20-G21)	1,312	1.3	36.9
14	Pneumonitis due to solids and liquids (J69)	683	0.7	19.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	590	0.6	16.6
...	All other causes (residual)	18,527	18.4	521.7
Hispanic, male, all ages				
...	All causes	92,474	100.0	330.1
1	Malignant neoplasms (C00-C97)	19,040	20.6	68.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,884	20.4	67.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,969	9.7	32.0
4	Diabetes mellitus (E10-E14)	4,149	4.5	14.8
5	Cerebrovascular diseases (I60-I69)	4,092	4.4	14.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,911	4.2	14.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,584	2.8	9.2
8	Chronic lower respiratory diseases (J40-J47)	2,433	2.6	8.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,122	2.3	7.6
10	Influenza and pneumonia (J09-J18)	1,975	2.1	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,700	1.8	6.1
12	Alzheimer's disease (G30)	1,600	1.7	5.7
13	Certain conditions originating in the perinatal period (P00-P96)	1,369	1.5	4.9
14	Septicemia (A40-A41)	1,319	1.4	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,016	1.1	3.6
...	All other causes (residual)	17,311	18.7	61.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	421	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	32.8	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	11.2	2.2
3	Malignant neoplasms (C00-C97)	37	8.8	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.3	*
6	Influenza and pneumonia (J09-J18)	9	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
10	Nutritional deficiencies (E40-E64)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	111	26.4	5.3
Hispanic, male, 5-14 years				
...	All causes	638	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	30.3	3.8
2	Malignant neoplasms (C00-C97)	132	20.7	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.7	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	5.5	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
8	Cerebrovascular diseases (I60-I69)	12	1.9	*
9	Septicemia (A40-A41)	11	1.7	*
9	Influenza and pneumonia (J09-J18)	11	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Anemias (D50-D64)	4	0.6	*
13	Nutritional deficiencies (E40-E64)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
15	Diseases of appendix (K35-K38)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (residual)	129	20.2	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,669	100.0	75.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,572	42.8	32.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	691	18.8	14.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	15.3	11.6
4	Malignant neoplasms (C00-C97)	237	6.5	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	2.5	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.0	0.8
7	Cerebrovascular diseases (I60-I69)	21	0.6	0.4
8	Influenza and pneumonia (J09-J18)	19	0.5	*
8	Legal intervention (Y35,Y89.0)	19	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.4	*
11	Septicemia (A40-A41)	13	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
13	Diabetes mellitus (E10-E14)	10	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
15	Anemias (D50-D64)	5	0.1	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
...	All other causes (residual)	346	9.4	7.1
Hispanic, male, 25-44 years				
...	All causes	10,935	100.0	125.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,422	31.3	39.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,101	10.1	12.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,069	9.8	12.2
4	Malignant neoplasms (C00-C97)	1,033	9.4	11.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,016	9.3	11.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	523	4.8	6.0
7	Cerebrovascular diseases (I60-I69)	263	2.4	3.0
8	Diabetes mellitus (E10-E14)	222	2.0	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	208	1.9	2.4
10	Influenza and pneumonia (J09-J18)	186	1.7	2.1
11	Septicemia (A40-A41)	107	1.0	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	0.6	0.8
13	Legal intervention (Y35,Y89.0)	65	0.6	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.5	0.7
15	Aortic aneurysm and dissection (I71)	50	0.5	0.6
...	All other causes (residual)	1,544	14.1	17.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	26,292	100.0	510.6
1	Malignant neoplasms (C00-C97)	6,142	23.4	119.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,359	20.4	104.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	2,416	9.2	46.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,327	8.9	45.2
5	Diabetes mellitus (E10-E14)	1,425	5.4	27.7
6	Cerebrovascular diseases (I60-I69)	1,074	4.1	20.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	641	2.4	12.4
8	Viral hepatitis (B15-B19)	560	2.1	10.9
9	Influenza and pneumonia (J09-J18)	477	1.8	9.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	437	1.7	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	1.6	8.4
12	Septicemia (A40-A41)	410	1.6	8.0
13	Chronic lower respiratory diseases (J40-J47)	336	1.3	6.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	1.1	5.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	264	1.0	5.1
...	All other causes (residual)	3,707	14.1	72.0
Hispanic, male, 65 years and over				
...	All causes	47,887	100.0	3,128.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,301	25.7	803.6
2	Malignant neoplasms (C00-C97)	11,454	23.9	748.3
3	Cerebrovascular diseases (I60-I69)	2,709	5.7	177.0
4	Diabetes mellitus (E10-E14)	2,491	5.2	162.7
5	Chronic lower respiratory diseases (J40-J47)	2,016	4.2	131.7
6	Alzheimer's disease (G30)	1,565	3.3	102.2
7	Influenza and pneumonia (J09-J18)	1,253	2.6	81.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,227	2.6	80.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,182	2.5	77.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	968	2.0	63.2
11	Septicemia (A40-A41)	753	1.6	49.2
12	Parkinson's disease (G20-G21)	720	1.5	47.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	702	1.5	45.9
14	Pneumonitis due to solids and liquids (J69)	391	0.8	25.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	304	0.6	19.9
...	All other causes (residual)	7,851	16.4	512.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	76,913	100.0	281.0
1	Malignant neoplasms (C00-C97)	17,407	22.6	63.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,137	19.7	55.3
3	Cerebrovascular diseases (I60-I69)	4,621	6.0	16.9
4	Diabetes mellitus (E10-E14)	3,646	4.7	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,444	4.5	12.6
6	Alzheimer's disease (G30)	3,334	4.3	12.2
7	Chronic lower respiratory diseases (J40-J47)	2,362	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,900	2.5	6.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,747	2.3	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,573	2.0	5.7
11	Septicemia (A40-A41)	1,294	1.7	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,123	1.5	4.1
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.4	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.1	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	662	0.9	2.4
...	All other causes (residual)	16,737	21.8	61.2
Hispanic, female, 1-4 years				
...	All causes	348	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	23.9	4.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	14.4	2.5
3	Malignant neoplasms (C00-C97)	44	12.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	11.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.3	*
6	Influenza and pneumonia (J09-J18)	11	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.0	*
8	Meningitis (G00,G03)	4	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
10	Septicemia (A40-A41)	3	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Diabetes mellitus (E10-E14)	1	0.3	*
15	Nutritional deficiencies (E40-E64)	1	0.3	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	71	20.4	3.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	478	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	25.1	2.4
2	Malignant neoplasms (C00-C97)	87	18.2	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.7	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	6.5	0.6
6	Cerebrovascular diseases (I60-I69)	15	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
8	Influenza and pneumonia (J09-J18)	12	2.5	*
9	Chronic lower respiratory diseases (J40-J47)	10	2.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
12	Septicemia (A40-A41)	5	1.0	*
13	Diabetes mellitus (E10-E14)	4	0.8	*
14	Hernia (K40-K46)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	95	19.9	1.9
Hispanic, female, 15-24 years				
...	All causes	1,236	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	38.2	10.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	12.7	3.4
3	Malignant neoplasms (C00-C97)	127	10.3	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	9.8	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.9	0.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	29	2.3	0.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	2.3	0.6
8	Cerebrovascular diseases (I60-I69)	13	1.1	*
9	Influenza and pneumonia (J09-J18)	12	1.0	*
10	Septicemia (A40-A41)	11	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
14	Anemias (D50-D64)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (residual)	197	15.9	4.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,808	100.0	59.2
1	Malignant neoplasms (C00-C97)	1,296	27.0	16.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	901	18.7	11.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	7.5	4.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	247	5.1	3.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	3.8	2.3
6	Cerebrovascular diseases (I60-I69)	167	3.5	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	161	3.3	2.0
8	Diabetes mellitus (E10-E14)	138	2.9	1.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	117	2.4	1.4
10	Influenza and pneumonia (J09-J18)	109	2.3	1.3
11	Septicemia (A40-A41)	69	1.4	0.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.3	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	1.1	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.0	0.6
15	Chronic lower respiratory diseases (J40-J47)	36	0.7	0.4
...	All other causes (residual)	861	17.9	10.6
Hispanic, female, 45-64 years				
...	All causes	15,114	100.0	288.7
1	Malignant neoplasms (C00-C97)	5,579	36.9	106.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,143	14.2	40.9
3	Diabetes mellitus (E10-E14)	851	5.6	16.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	795	5.3	15.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	750	5.0	14.3
6	Cerebrovascular diseases (I60-I69)	602	4.0	11.5
7	Influenza and pneumonia (J09-J18)	339	2.2	6.5
8	Septicemia (A40-A41)	299	2.0	5.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	291	1.9	5.6
10	Chronic lower respiratory diseases (J40-J47)	260	1.7	5.0
11	Viral hepatitis (B15-B19)	219	1.4	4.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	1.2	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	136	0.9	2.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	93	0.6	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.5	1.5
...	All other causes (residual)	2,496	16.5	47.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	52,783	100.0	2,612.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,541	23.8	620.6
2	Malignant neoplasms (C00-C97)	10,268	19.5	508.1
3	Cerebrovascular diseases (I60-I69)	3,815	7.2	188.8
4	Alzheimer's disease (G30)	3,299	6.3	163.3
5	Diabetes mellitus (E10-E14)	2,644	5.0	130.8
6	Chronic lower respiratory diseases (J40-J47)	2,043	3.9	101.1
7	Influenza and pneumonia (J09-J18)	1,402	2.7	69.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,224	2.3	60.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,010	1.9	50.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	968	1.8	47.9
11	Septicemia (A40-A41)	891	1.7	44.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	832	1.6	41.2
13	Parkinson's disease (G20-G21)	592	1.1	29.3
14	Pneumonitis due to solids and liquids (J69)	292	0.6	14.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	286	0.5	14.2
...	All other causes (residual)	10,676	20.2	528.3
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,066,949	100.0	1,028.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	489,926	23.7	243.7
2	Malignant neoplasms (C00-C97)	466,271	22.6	231.9
3	Chronic lower respiratory diseases (J40-J47)	129,519	6.3	64.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,682	5.1	52.1
5	Cerebrovascular diseases (I60-I69)	102,326	5.0	50.9
6	Alzheimer's disease (G30)	80,014	3.9	39.8
7	Diabetes mellitus (E10-E14)	51,944	2.5	25.8
8	Influenza and pneumonia (J09-J18)	43,377	2.1	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,444	1.7	17.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,691	1.7	17.3
11	Septicemia (A40-A41)	28,868	1.4	14.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,830	1.3	13.8
13	Parkinson's disease (G20-G21)	22,917	1.1	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,511	1.0	10.7
15	Pneumonitis due to solids and liquids (J69)	15,644	0.8	7.8
...	All other causes (residual)	411,985	19.9	204.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,876	100.0	22.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	655	34.9	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	9.6	2.2
3	Malignant neoplasms (C00-C97)	176	9.4	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	7.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.0	0.7
6	Influenza and pneumonia (J09-J18)	53	2.8	0.6
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
14	Anemias (D50-D64)	6	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (residual)	493	26.3	5.9
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,684	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	769	28.7	3.5
2	Malignant neoplasms (C00-C97)	451	16.8	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	285	10.6	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	6.6	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	3.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.4	0.4
7	Influenza and pneumonia (J09-J18)	52	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	42	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	37	1.4	0.2
11	Septicemia (A40-A41)	26	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
14	Anemias (D50-D64)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (residual)	562	20.9	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,040	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,743	48.3	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,503	21.8	14.1
3	Malignant neoplasms (C00-C97)	859	5.4	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	607	3.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	493	3.1	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	1.4	0.9
7	Influenza and pneumonia (J09-J18)	123	0.8	0.5
8	Diabetes mellitus (E10-E14)	101	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	88	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	75	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	71	0.4	0.3
12	Septicemia (A40-A41)	65	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.2	0.1
15	Legal intervention (Y35,Y89.0)	27	0.2	0.1
...	All other causes (residual)	1,986	12.4	8.0
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	73,534	100.0	149.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,939	32.6	48.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,280	14.0	20.9
3	Malignant neoplasms (C00-C97)	9,078	12.3	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,903	10.7	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,085	2.8	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,779	2.4	3.6
7	Diabetes mellitus (E10-E14)	1,430	1.9	2.9
8	Cerebrovascular diseases (I60-I69)	1,099	1.5	2.2
9	Influenza and pneumonia (J09-J18)	996	1.4	2.0
10	Septicemia (A40-A41)	687	0.9	1.4
11	Chronic lower respiratory diseases (J40-J47)	682	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	564	0.8	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	435	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	401	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	347	0.5	0.7
...	All other causes (residual)	11,829	16.1	24.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	372,977	100.0	642.3
1	Malignant neoplasms (C00-C97)	116,708	31.3	201.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,873	20.3	130.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,192	7.8	50.3
4	Chronic lower respiratory diseases (J40-J47)	17,418	4.7	30.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,415	4.1	26.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,349	3.8	24.7
7	Diabetes mellitus (E10-E14)	11,926	3.2	20.5
8	Cerebrovascular diseases (I60-I69)	10,079	2.7	17.4
9	Influenza and pneumonia (J09-J18)	5,547	1.5	9.6
10	Septicemia (A40-A41)	5,374	1.4	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,877	1.0	6.7
12	Viral hepatitis (B15-B19)	3,851	1.0	6.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,717	0.7	4.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,674	0.4	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,492	0.4	2.6
...	All other causes (residual)	57,485	15.4	99.0
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,589,400	100.0	4,367.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	405,361	25.5	1,113.9
2	Malignant neoplasms (C00-C97)	338,956	21.3	931.5
3	Chronic lower respiratory diseases (J40-J47)	111,278	7.0	305.8
4	Cerebrovascular diseases (I60-I69)	90,952	5.7	249.9
5	Alzheimer's disease (G30)	79,270	5.0	217.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,810	2.6	114.9
7	Diabetes mellitus (E10-E14)	38,472	2.4	105.7
8	Influenza and pneumonia (J09-J18)	36,535	2.3	100.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,323	1.9	83.3
10	Septicemia (A40-A41)	22,626	1.4	62.2
11	Parkinson's disease (G20-G21)	22,383	1.4	61.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,567	1.2	51.0
13	Pneumonitis due to solids and liquids (J69)	14,050	0.9	38.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,296	0.7	31.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10,312	0.6	28.3
...	All other causes (residual)	317,209	20.0	871.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,035,345	100.0	1,045.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	258,473	25.0	261.0
2	Malignant neoplasms (C00-C97)	246,872	23.8	249.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63,807	6.2	64.4
4	Chronic lower respiratory diseases (J40-J47)	60,428	5.8	61.0
5	Cerebrovascular diseases (I60-I69)	41,410	4.0	41.8
6	Diabetes mellitus (E10-E14)	28,743	2.8	29.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,410	2.6	27.7
8	Alzheimer's disease (G30)	24,321	2.3	24.6
9	Influenza and pneumonia (J09-J18)	20,633	2.0	20.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,878	1.7	18.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,844	1.7	18.0
12	Parkinson's disease (G20-G21)	13,830	1.3	14.0
13	Septicemia (A40-A41)	13,514	1.3	13.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,984	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,623	0.8	8.7
...	All other causes (residual)	182,575	17.6	184.3
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,060	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	36.9	9.2
2	Malignant neoplasms (C00-C97)	110	10.4	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.7	0.7
5	Influenza and pneumonia (J09-J18)	29	2.7	0.7
7	Septicemia (A40-A41)	13	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
10	Cerebrovascular diseases (I60-I69)	7	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
...	All other causes (residual)	259	24.4	6.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,620	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	31.5	4.5
2	Malignant neoplasms (C00-C97)	256	15.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	11.8	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	4.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.8	0.4
7	Influenza and pneumonia (J09-J18)	26	1.6	0.2
8	Chronic lower respiratory diseases (J40-J47)	25	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	24	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	10	0.6	*
11	Diabetes mellitus (E10-E14)	10	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	9	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
15	Anemias (D50-D64)	4	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (residual)	322	19.9	2.8
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,508	100.0	90.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,707	49.6	44.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,850	24.8	22.4
3	Malignant neoplasms (C00-C97)	509	4.4	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	447	3.9	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	318	2.8	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	1.1	1.0
7	Diabetes mellitus (E10-E14)	69	0.6	0.5
8	Influenza and pneumonia (J09-J18)	66	0.6	0.5
9	Chronic lower respiratory diseases (J40-J47)	49	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	48	0.4	0.4
11	Septicemia (A40-A41)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.2	0.2
12	Legal intervention (Y35,Y89.0)	24	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.1	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.1	*
...	All other causes (residual)	1,203	10.5	9.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	47,258	100.0	190.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,783	35.5	67.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,878	16.7	31.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,420	11.5	21.9
4	Malignant neoplasms (C00-C97)	4,184	8.9	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,248	2.6	5.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,221	2.6	4.9
7	Diabetes mellitus (E10-E14)	903	1.9	3.6
8	Cerebrovascular diseases (I60-I69)	594	1.3	2.4
9	Influenza and pneumonia (J09-J18)	520	1.1	2.1
10	Septicemia (A40-A41)	334	0.7	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	332	0.7	1.3
12	Chronic lower respiratory diseases (J40-J47)	331	0.7	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	319	0.7	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	0.5	1.0
15	Aortic aneurysm and dissection (I71)	168	0.4	0.7
...	All other causes (residual)	6,779	14.3	27.3
Non-Hispanic, white, male, 45-64 years				
...	All causes	228,091	100.0	797.2
1	Malignant neoplasms (C00-C97)	63,322	27.8	221.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,075	23.7	189.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,314	8.5	67.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,632	4.7	37.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,404	4.6	36.4
6	Chronic lower respiratory diseases (J40-J47)	8,773	3.8	30.7
7	Diabetes mellitus (E10-E14)	7,598	3.3	26.6
8	Cerebrovascular diseases (I60-I69)	5,713	2.5	20.0
9	Influenza and pneumonia (J09-J18)	3,095	1.4	10.8
10	Septicemia (A40-A41)	2,838	1.2	9.9
11	Viral hepatitis (B15-B19)	2,691	1.2	9.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,310	1.0	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,746	0.8	6.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,137	0.5	4.0
15	Aortic aneurysm and dissection (I71)	1,086	0.5	3.8
...	All other causes (residual)	33,357	14.6	116.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	739,947	100.0	4,562.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	198,501	26.8	1,223.8
2	Malignant neoplasms (C00-C97)	178,466	24.1	1,100.3
3	Chronic lower respiratory diseases (J40-J47)	51,230	6.9	315.9
4	Cerebrovascular diseases (I60-I69)	34,993	4.7	215.7
5	Alzheimer's disease (G30)	24,007	3.2	148.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,766	2.8	128.0
7	Diabetes mellitus (E10-E14)	20,161	2.7	124.3
8	Influenza and pneumonia (J09-J18)	16,859	2.3	103.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,280	2.1	94.2
10	Parkinson's disease (G20-G21)	13,481	1.8	83.1
11	Septicemia (A40-A41)	10,248	1.4	63.2
12	Pneumonitis due to solids and liquids (J69)	7,671	1.0	47.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,079	1.0	43.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,180	0.8	38.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,035	0.8	37.2
...	All other causes (residual)	128,990	17.4	795.3
Non-Hispanic, white, female, all ages				
...	All causes	1,031,604	100.0	1,011.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,453	22.4	226.9
2	Malignant neoplasms (C00-C97)	219,399	21.3	215.1
3	Chronic lower respiratory diseases (J40-J47)	69,091	6.7	67.7
4	Cerebrovascular diseases (I60-I69)	60,916	5.9	59.7
5	Alzheimer's disease (G30)	55,693	5.4	54.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40,875	4.0	40.1
7	Diabetes mellitus (E10-E14)	23,201	2.2	22.7
8	Influenza and pneumonia (J09-J18)	22,744	2.2	22.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,813	1.6	16.5
10	Septicemia (A40-A41)	15,354	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,527	1.2	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,986	1.0	9.8
13	Parkinson's disease (G20-G21)	9,087	0.9	8.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,034	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,021	0.7	6.9
...	All other causes (residual)	229,410	22.2	224.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	816	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	32.4	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	9.7	2.0
3	Malignant neoplasms (C00-C97)	66	8.1	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.3	0.7
6	Influenza and pneumonia (J09-J18)	24	2.9	0.6
7	Septicemia (A40-A41)	19	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
13	Meningitis (G00,G03)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.4	*
...	All other causes (residual)	221	27.1	5.5
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,064	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	24.2	2.4
2	Malignant neoplasms (C00-C97)	195	18.3	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	8.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	7.9	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.1	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.8	0.4
7	Influenza and pneumonia (J09-J18)	26	2.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.8	*
9	Cerebrovascular diseases (I60-I69)	18	1.7	*
10	Septicemia (A40-A41)	16	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
13	Anemias (D50-D64)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Tuberculosis (A16-A19)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
...	All other causes (residual)	241	22.7	2.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,532	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,036	44.9	16.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	653	14.4	5.4
3	Malignant neoplasms (C00-C97)	350	7.7	2.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	3.9	1.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	2.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	71	1.6	0.6
8	Influenza and pneumonia (J09-J18)	57	1.3	0.5
9	Cerebrovascular diseases (I60-I69)	40	0.9	0.3
10	Diabetes mellitus (E10-E14)	32	0.7	0.3
11	Septicemia (A40-A41)	27	0.6	0.2
12	Chronic lower respiratory diseases (J40-J47)	26	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Pneumonitis due to solids and liquids (J69)	11	0.2	*
...	All other causes (residual)	760	16.8	6.3
Non-Hispanic, white, female, 25-44 years				
...	All causes	26,276	100.0	107.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,156	27.2	29.3
2	Malignant neoplasms (C00-C97)	4,894	18.6	20.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,483	9.4	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,402	9.1	9.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	837	3.2	3.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	558	2.1	2.3
7	Diabetes mellitus (E10-E14)	527	2.0	2.2
8	Cerebrovascular diseases (I60-I69)	505	1.9	2.1
9	Influenza and pneumonia (J09-J18)	476	1.8	2.0
10	Septicemia (A40-A41)	353	1.3	1.4
11	Chronic lower respiratory diseases (J40-J47)	351	1.3	1.4
12	Pregnancy, childbirth and the puerperium (O00-O99)	347	1.3	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	0.6	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	0.4
...	All other causes (residual)	4,878	18.6	20.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	144,886	100.0	491.8
1	Malignant neoplasms (C00-C97)	53,386	36.8	181.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,798	15.0	74.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,878	6.8	33.5
4	Chronic lower respiratory diseases (J40-J47)	8,645	6.0	29.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,011	3.5	17.0
6	Cerebrovascular diseases (I60-I69)	4,366	3.0	14.8
7	Diabetes mellitus (E10-E14)	4,328	3.0	14.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,717	2.6	12.6
9	Septicemia (A40-A41)	2,536	1.8	8.6
10	Influenza and pneumonia (J09-J18)	2,452	1.7	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,567	1.1	5.3
12	Viral hepatitis (B15-B19)	1,160	0.8	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	971	0.7	3.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	681	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	671	0.5	2.3
...	All other causes (residual)	23,719	16.4	80.5
Non-Hispanic, white, female, 65 years and over				
...	All causes	849,453	100.0	4,211.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	206,860	24.4	1,025.6
2	Malignant neoplasms (C00-C97)	160,490	18.9	795.7
3	Chronic lower respiratory diseases (J40-J47)	60,048	7.1	297.7
4	Cerebrovascular diseases (I60-I69)	55,959	6.6	277.4
5	Alzheimer's disease (G30)	55,263	6.5	274.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,044	2.5	104.3
7	Influenza and pneumonia (J09-J18)	19,676	2.3	97.5
8	Diabetes mellitus (E10-E14)	18,311	2.2	90.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,043	1.8	74.6
10	Septicemia (A40-A41)	12,378	1.5	61.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,488	1.4	57.0
12	Parkinson's disease (G20-G21)	8,902	1.0	44.1
13	Pneumonitis due to solids and liquids (J69)	6,379	0.8	31.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,261	0.6	26.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,132	0.5	20.5
...	All other causes (residual)	188,219	22.2	933.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	303,844	100.0	735.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,894	23.7	174.0
2	Malignant neoplasms (C00-C97)	68,104	22.4	164.8
3	Cerebrovascular diseases (I60-I69)	16,883	5.6	40.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,820	4.5	33.4
5	Diabetes mellitus (E10-E14)	13,264	4.4	32.1
6	Chronic lower respiratory diseases (J40-J47)	9,762	3.2	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,496	2.8	20.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,767	2.6	18.8
9	Alzheimer's disease (G30)	6,493	2.1	15.7
10	Septicemia (A40-A41)	6,321	2.1	15.3
11	Influenza and pneumonia (J09-J18)	5,462	1.8	13.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,322	1.8	12.9
13	Certain conditions originating in the perinatal period (P00-P96)	3,810	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,527	1.2	8.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,011	1.0	7.3
...	All other causes (residual)	59,908	19.7	145.0
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	967	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	28.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	14.8	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	10.0	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.6	2.2
5	Malignant neoplasms (C00-C97)	50	5.2	2.0
6	Influenza and pneumonia (J09-J18)	28	2.9	1.1
7	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.8
8	Septicemia (A40-A41)	12	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Anemias (D50-D64)	6	0.6	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	232	24.0	9.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,162	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	28.1	5.3
2	Malignant neoplasms (C00-C97)	142	12.2	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	8.9	1.7
4	Chronic lower respiratory diseases (J40-J47)	71	6.1	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.0	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.2	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	4.7	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.7	0.3
9	Anemias (D50-D64)	17	1.5	*
10	Influenza and pneumonia (J09-J18)	16	1.4	*
11	Septicemia (A40-A41)	15	1.3	*
11	Cerebrovascular diseases (I60-I69)	15	1.3	*
13	Diabetes mellitus (E10-E14)	10	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	221	19.0	3.6
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,425	100.0	94.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,632	41.0	38.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,466	22.8	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	519	8.1	7.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	4.4	4.1
5	Malignant neoplasms (C00-C97)	261	4.1	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	71	1.1	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.0	1.0
9	Anemias (D50-D64)	64	1.0	0.9
10	Diabetes mellitus (E10-E14)	59	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.9	0.8
12	Cerebrovascular diseases (I60-I69)	48	0.7	0.7
13	Influenza and pneumonia (J09-J18)	36	0.6	0.5
14	Septicemia (A40-A41)	32	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.5	0.4
15	Legal intervention (Y35,Y89.0)	29	0.5	0.4
...	All other causes (residual)	695	10.8	10.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	23,146	100.0	206.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,744	16.2	33.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,717	16.1	33.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,549	15.3	31.7
4	Malignant neoplasms (C00-C97)	2,566	11.1	22.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,002	4.3	9.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	982	4.2	8.8
7	Diabetes mellitus (E10-E14)	777	3.4	6.9
8	Cerebrovascular diseases (I60-I69)	658	2.8	5.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	1.5	3.1
10	Chronic lower respiratory diseases (J40-J47)	306	1.3	2.7
11	Influenza and pneumonia (J09-J18)	293	1.3	2.6
12	Septicemia (A40-A41)	268	1.2	2.4
13	Anemias (D50-D64)	266	1.1	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	1.0	2.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	239	1.0	2.1
...	All other causes (residual)	4,191	18.1	37.4
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	90,376	100.0	906.0
1	Malignant neoplasms (C00-C97)	25,183	27.9	252.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,964	24.3	220.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,894	5.4	49.1
4	Cerebrovascular diseases (I60-I69)	4,389	4.9	44.0
5	Diabetes mellitus (E10-E14)	4,291	4.7	43.0
6	Chronic lower respiratory diseases (J40-J47)	2,448	2.7	24.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,211	2.4	22.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,060	2.3	20.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,979	2.2	19.8
10	Septicemia (A40-A41)	1,895	2.1	19.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,482	1.6	14.9
12	Influenza and pneumonia (J09-J18)	1,400	1.5	14.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,066	1.2	10.7
14	Viral hepatitis (B15-B19)	819	0.9	8.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	598	0.7	6.0
...	All other causes (residual)	13,697	15.2	137.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	175,050	100.0	4,295.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,674	26.1	1,120.6
2	Malignant neoplasms (C00-C97)	39,892	22.8	978.8
3	Cerebrovascular diseases (I60-I69)	11,741	6.7	288.1
4	Diabetes mellitus (E10-E14)	8,125	4.6	199.4
5	Chronic lower respiratory diseases (J40-J47)	6,838	3.9	167.8
6	Alzheimer's disease (G30)	6,388	3.6	156.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,881	3.4	144.3
8	Septicemia (A40-A41)	4,047	2.3	99.3
9	Influenza and pneumonia (J09-J18)	3,618	2.1	88.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,592	2.1	88.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,732	1.6	67.0
12	Pneumonitis due to solids and liquids (J69)	1,354	0.8	33.2
13	Parkinson's disease (G20-G21)	1,080	0.6	26.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	894	0.5	21.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	807	0.5	19.8
...	All other causes (residual)	32,387	18.5	794.6
Non-Hispanic, black, male, all ages				
...	All causes	154,836	100.0	783.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,265	24.1	188.5
2	Malignant neoplasms (C00-C97)	34,539	22.3	174.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,316	6.0	47.1
4	Cerebrovascular diseases (I60-I69)	7,650	4.9	38.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,718	4.3	34.0
6	Diabetes mellitus (E10-E14)	6,369	4.1	32.2
7	Chronic lower respiratory diseases (J40-J47)	4,940	3.2	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,983	2.6	20.2
9	Septicemia (A40-A41)	2,938	1.9	14.9
10	Influenza and pneumonia (J09-J18)	2,656	1.7	13.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,440	1.6	12.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,321	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.4	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,876	1.2	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,873	1.2	9.5
...	All other causes (residual)	27,818	18.0	140.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	560	100.0	45.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	30.9	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.6	2.1
5	Malignant neoplasms (C00-C97)	25	4.5	2.0
6	Influenza and pneumonia (J09-J18)	18	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Septicemia (A40-A41)	7	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Anemias (D50-D64)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	133	23.8	10.7
Non-Hispanic, black, male, 5-14 years				
...	All causes	701	100.0	22.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	31.2	7.0
2	Malignant neoplasms (C00-C97)	75	10.7	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	9.8	2.2
4	Chronic lower respiratory diseases (J40-J47)	47	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.7	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.7	1.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.7	1.3
8	Influenza and pneumonia (J09-J18)	9	1.3	*
9	Septicemia (A40-A41)	8	1.1	*
9	Anemias (D50-D64)	8	1.1	*
11	Cerebrovascular diseases (I60-I69)	7	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
13	Diabetes mellitus (E10-E14)	5	0.7	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
15	Diseases of appendix (K35-K38)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	114	16.3	3.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,935	100.0	142.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,388	48.4	69.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,092	22.1	31.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	8.7	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.4	4.9
5	Malignant neoplasms (C00-C97)	153	3.1	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.0	1.4
7	Chronic lower respiratory diseases (J40-J47)	47	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.7	1.1
9	Diabetes mellitus (E10-E14)	33	0.7	1.0
10	Anemias (D50-D64)	32	0.6	0.9
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.8
11	Legal intervention (Y35,Y89.0)	29	0.6	0.8
13	Influenza and pneumonia (J09-J18)	26	0.5	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Septicemia (A40-A41)	12	0.2	*
...	All other causes (residual)	393	8.0	11.4
Non-Hispanic, black, male, 25-44 years				
...	All causes	14,564	100.0	271.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,136	21.5	58.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,678	18.4	50.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,321	15.9	43.3
4	Malignant neoplasms (C00-C97)	1,011	6.9	18.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	808	5.5	15.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	607	4.2	11.3
7	Diabetes mellitus (E10-E14)	426	2.9	8.0
8	Cerebrovascular diseases (I60-I69)	389	2.7	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.3	3.6
10	Chronic lower respiratory diseases (J40-J47)	170	1.2	3.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	157	1.1	2.9
12	Anemias (D50-D64)	151	1.0	2.8
13	Influenza and pneumonia (J09-J18)	147	1.0	2.7
14	Septicemia (A40-A41)	134	0.9	2.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	106	0.7	2.0
...	All other causes (residual)	2,131	14.6	39.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	51,936	100.0	1,124.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,748	26.5	297.6
2	Malignant neoplasms (C00-C97)	13,009	25.0	281.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,383	6.5	73.2
4	Cerebrovascular diseases (I60-I69)	2,482	4.8	53.7
5	Diabetes mellitus (E10-E14)	2,396	4.6	51.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,396	2.7	30.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,251	2.4	27.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,234	2.4	26.7
9	Chronic lower respiratory diseases (J40-J47)	1,197	2.3	25.9
10	Septicemia (A40-A41)	975	1.9	21.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	868	1.7	18.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	863	1.7	18.7
13	Influenza and pneumonia (J09-J18)	769	1.5	16.6
14	Viral hepatitis (B15-B19)	549	1.1	11.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	455	0.9	9.8
...	All other causes (residual)	7,361	14.2	159.3
Non-Hispanic, black, male, 65 years and over				
...	All causes	78,444	100.0	4,808.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,899	26.6	1,281.1
2	Malignant neoplasms (C00-C97)	20,264	25.8	1,242.1
3	Cerebrovascular diseases (I60-I69)	4,725	6.0	289.6
4	Diabetes mellitus (E10-E14)	3,507	4.5	215.0
5	Chronic lower respiratory diseases (J40-J47)	3,464	4.4	212.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,524	3.2	154.7
7	Septicemia (A40-A41)	1,775	2.3	108.8
8	Alzheimer's disease (G30)	1,774	2.3	108.7
9	Influenza and pneumonia (J09-J18)	1,660	2.1	101.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,545	2.0	94.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,409	1.8	86.4
12	Pneumonitis due to solids and liquids (J69)	686	0.9	42.1
13	Parkinson's disease (G20-G21)	620	0.8	38.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	516	0.7	31.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	369	0.5	22.6
...	All other causes (residual)	12,707	16.2	778.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014



LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	149,008	100.0	691.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,629	23.2	160.7
2	Malignant neoplasms (C00-C97)	33,565	22.5	155.8
3	Cerebrovascular diseases (I60-I69)	9,233	6.2	42.8
4	Diabetes mellitus (E10-E14)	6,895	4.6	32.0
5	Chronic lower respiratory diseases (J40-J47)	4,822	3.2	22.4
6	Alzheimer's disease (G30)	4,675	3.1	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,513	3.0	20.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,504	3.0	20.9
9	Septicemia (A40-A41)	3,383	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,882	1.9	13.4
11	Influenza and pneumonia (J09-J18)	2,806	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,676	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,206	0.8	5.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,138	0.8	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,049	0.7	4.9
...	All other causes (residual)	32,032	21.5	148.6
Non-Hispanic, black, female, 1-4 years				
...	All causes	407	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	16.0	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.1	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.9	2.3
5	Malignant neoplasms (C00-C97)	25	6.1	2.1
6	Chronic lower respiratory diseases (J40-J47)	11	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.5	*
8	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
13	Salmonella infections (A01-A02)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Malaria (B50-B54)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (residual)	94	23.1	7.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	461	100.0	15.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	23.4	3.5
2	Malignant neoplasms (C00-C97)	67	14.5	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	7.4	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	6.5	1.0
5	Chronic lower respiratory diseases (J40-J47)	24	5.2	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.6	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	3.3	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.3	*
9	Anemias (D50-D64)	9	2.0	*
10	Cerebrovascular diseases (I60-I69)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
11	Influenza and pneumonia (J09-J18)	7	1.5	*
13	Diabetes mellitus (E10-E14)	5	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (residual)	105	22.8	3.4
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,490	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	25.1	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	16.4	7.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	7.4	3.3
4	Malignant neoplasms (C00-C97)	108	7.2	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	6.1	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.8	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	34	2.3	1.0
8	Anemias (D50-D64)	32	2.1	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.9	0.8
10	Diabetes mellitus (E10-E14)	26	1.7	0.8
11	Chronic lower respiratory diseases (J40-J47)	24	1.6	0.7
12	Septicemia (A40-A41)	20	1.3	0.6
13	Cerebrovascular diseases (I60-I69)	19	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.8	*
15	Influenza and pneumonia (J09-J18)	10	0.7	*
...	All other causes (residual)	302	20.3	9.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	8,582	100.0	147.0
1	Malignant neoplasms (C00-C97)	1,555	18.1	26.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,423	16.6	24.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,039	12.1	17.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	413	4.8	7.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	395	4.6	6.8
6	Diabetes mellitus (E10-E14)	351	4.1	6.0
7	Cerebrovascular diseases (I60-I69)	269	3.1	4.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	239	2.8	4.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	2.0	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	1.8	2.7
11	Influenza and pneumonia (J09-J18)	146	1.7	2.5
12	Chronic lower respiratory diseases (J40-J47)	136	1.6	2.3
13	Septicemia (A40-A41)	134	1.6	2.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	118	1.4	2.0
15	Anemias (D50-D64)	115	1.3	2.0
...	All other causes (residual)	1,919	22.4	32.9
Non-Hispanic, black, female, 45-64 years				
...	All causes	38,440	100.0	717.8
1	Malignant neoplasms (C00-C97)	12,174	31.7	227.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,216	21.4	153.4
3	Cerebrovascular diseases (I60-I69)	1,907	5.0	35.6
4	Diabetes mellitus (E10-E14)	1,895	4.9	35.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,511	3.9	28.2
6	Chronic lower respiratory diseases (J40-J47)	1,251	3.3	23.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	977	2.5	18.2
8	Septicemia (A40-A41)	920	2.4	17.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	728	1.9	13.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	664	1.7	12.4
11	Influenza and pneumonia (J09-J18)	631	1.6	11.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	614	1.6	11.5
13	Viral hepatitis (B15-B19)	270	0.7	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	203	0.5	3.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.4	3.0
...	All other causes (residual)	6,321	16.4	118.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	96,606	100.0	3,952.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,775	25.6	1,013.6
2	Malignant neoplasms (C00-C97)	19,628	20.3	803.0
3	Cerebrovascular diseases (I60-I69)	7,016	7.3	287.0
4	Diabetes mellitus (E10-E14)	4,618	4.8	188.9
5	Alzheimer's disease (G30)	4,614	4.8	188.8
6	Chronic lower respiratory diseases (J40-J47)	3,374	3.5	138.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,357	3.5	137.3
8	Septicemia (A40-A41)	2,272	2.4	93.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,183	2.3	89.3
10	Influenza and pneumonia (J09-J18)	1,958	2.0	80.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,187	1.2	48.6
12	Pneumonitis due to solids and liquids (J69)	668	0.7	27.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	525	0.5	21.5
14	Parkinson's disease (G20-G21)	460	0.5	18.8
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	293	0.3	12.0
...	All other causes (residual)	19,678	20.4	805.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2014