

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All causes											
All origins 3/, both sexes											
2007.....	803.6	684.5	28.6	13.7	16.9	61.9	98.3	99.4	110.8	145.8	221.6
2006.....	810.4	690.7	28.4	13.9	16.6	64.4	100.2	100.9	111.9	148.7	229.3
2005.....	825.9	692.5	29.4	14.5	18.1	65.1	97.6	97.5	111.4	149.6	233.4
2004.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7	231.7
2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1	241.3
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3	242.6
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9	240.5
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2	239.1
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8	236.3
All origins 3/, male											
2007.....	809.9	747.8	31.3	15.0	19.9	86.8	145.2	140.2	148.2	185.4	276.8
2006.....	814.8	756.3	30.5	15.4	19.6	90.7	148.0	143.4	150.4	189.0	285.9
2005.....	827.2	762.3	33.4	15.6	21.5	91.6	143.9	138.7	148.1	189.4	292.7
2004.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8	289.5
2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7	304.1
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6	306.4
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7	304.8
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4	305.9
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6	304.0
All origins 3/, female											
2007.....	797.4	618.1	25.7	12.3	13.8	35.7	48.4	56.6	72.3	105.7	166.7
2006.....	806.1	622.0	26.3	12.3	13.3	36.8	49.0	56.5	72.5	107.9	173.1
2005.....	824.6	619.4	25.1	13.4	14.4	37.2	48.2	54.4	73.7	109.3	174.7
2004.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3	174.5
2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3	179.2
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1	179.7
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2	177.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2	173.4
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4	169.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Hispanic, both sexes

2007.....	297.8	587.4	26.0	13.4	14.9	57.9	95.3	79.4	80.9	103.4	154.0
2006.....	300.1	590.6	26.4	11.8	16.6	65.3	95.0	79.4	83.7	108.2	164.0
2005.....	307.3	614.1	28.9	12.0	15.4	67.1	94.2	80.8	83.0	109.8	171.7
2004.....	296.2	586.9	27.3	12.6	16.3	64.2	87.3	77.1	79.4	112.7	168.3
2003.....	305.8	610.1	30.2	14.0	17.3	67.2	85.7	81.3	86.6	116.0	178.5
2002.....	302.2	592.7	29.8	14.4	16.6	65.2	87.9	74.7	84.2	115.5	181.8
2001.....	306.8	572.7	30.6	13.0	16.6	63.1	86.9	80.6	88.7	129.5	185.3
2000.....	303.8	596.3	29.6	14.1	17.5	61.6	83.3	76.7	93.2	129.4	192.2
1999.....	305.7	583.8	30.9	13.3	18.3	61.0	81.1	77.5	94.1	132.9	189.4

Hispanic, male

2007.....	321.8	632.7	28.0	14.6	17.2	86.8	143.0	110.7	109.4	134.8	202.8
2006.....	323.9	640.7	28.8	13.5	19.3	98.0	141.4	111.7	113.9	142.4	215.6
2005.....	334.4	670.2	33.2	12.8	17.9	98.6	139.6	115.8	115.1	145.1	224.7
2004.....	321.1	636.5	30.2	13.8	19.8	96.5	130.9	108.8	109.7	151.5	222.3
2003.....	330.7	665.5	33.8	16.0	20.9	98.5	124.6	115.7	120.7	153.8	232.5
2002.....	328.7	644.0	34.2	15.9	19.0	97.0	128.9	108.7	116.5	151.4	242.6
2001.....	332.9	624.4	33.8	14.9	18.6	92.0	128.2	115.2	121.1	175.5	249.0
2000.....	331.3	637.1	31.5	15.0	21.3	90.5	123.1	110.3	131.1	172.5	259.1
1999.....	332.6	623.4	32.9	14.9	21.2	87.8	119.1	111.1	130.8	183.0	255.7

Hispanic, female

2007.....	272.1	539.9	23.8	12.2	12.5	27.2	40.1	40.0	46.8	67.5	99.8
2006.....	274.6	538.3	24.0	10.1	13.7	30.5	40.0	38.7	47.7	69.0	107.0
2005.....	278.2	555.4	24.5	11.2	12.7	33.6	39.5	36.9	45.3	69.5	113.8
2004.....	269.7	535.1	24.3	11.3	12.7	29.7	33.9	37.7	44.2	68.6	109.8
2003.....	279.3	552.4	26.3	12.0	13.6	33.5	37.4	39.1	47.1	73.4	120.5
2002.....	274.0	539.1	25.3	12.9	14.0	30.8	37.2	33.3	46.9	75.6	117.3
2001.....	279.0	518.9	27.2	11.0	14.5	31.1	36.1	39.4	51.3	79.0	118.3
2000.....	274.6	553.6	27.5	13.2	13.6	28.7	34.6	37.5	49.8	82.4	122.2
1999.....	277.2	542.3	28.7	11.6	15.2	30.5	35.3	38.0	52.1	78.3	120.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/
Under 1 year 2/
1-4 years
5-9 years
10-14 years
15-19 years
20-24 years
25-29 years
30-34 years
35-39 years
40-44 years

All causes

Non-Hispanic 4/, both sexes

2007.....	892.0	709.8	29.2	13.7	17.4	62.7	98.7	104.1	118.1	154.2	232.5
2006.....	897.1	714.6	28.9	14.4	16.5	64.1	100.9	106.0	118.5	156.3	238.9
2005.....	911.2	708.4	29.4	15.1	18.6	64.6	98.0	101.3	117.5	156.8	241.8
2004.....	899.4	705.6	30.5	15.2	19.1	66.2	95.2	100.3	113.8	158.3	239.8
2003.....	924.4	716.2	31.7	14.7	19.5	65.9	98.3	98.2	116.0	165.7	248.7
2002.....	928.8	713.0	31.4	15.3	20.1	67.9	96.3	99.1	116.6	168.0	249.2
2001.....	926.2	703.5	33.8	15.8	19.6	67.3	96.2	99.2	117.5	170.6	245.8
2000.....	929.6	760.4	32.9	16.1	20.7	67.7	95.3	93.8	113.5	162.4	242.8
1999.....	929.9	760.7	34.8	17.5	20.7	69.7	92.4	92.5	116.1	163.7	239.9

Non-Hispanic 4/, male

2007.....	899.8	777.9	32.2	15.1	20.4	86.7	145.3	147.9	158.6	196.2	289.3
2006.....	902.8	784.5	30.8	15.9	19.6	88.9	149.0	151.8	159.7	198.4	296.7
2005.....	912.6	781.7	33.3	16.2	22.2	90.0	144.3	144.5	155.9	197.9	302.1
2004.....	900.9	778.7	32.9	17.0	22.1	89.6	139.3	142.5	150.4	201.3	298.4
2003.....	922.9	798.1	35.2	16.2	23.5	90.6	143.5	137.6	154.5	210.6	312.7
2002.....	928.0	782.6	35.4	16.8	23.7	93.0	142.8	140.6	155.7	215.0	313.3
2001.....	923.6	773.2	37.6	17.0	23.6	93.6	143.3	139.8	156.2	218.5	309.6
2000.....	928.1	836.8	36.8	18.1	24.7	93.4	140.9	132.2	150.3	208.8	309.0
1999.....	932.2	834.9	38.8	19.3	24.9	95.8	135.7	129.6	154.9	209.7	307.1

Non-Hispanic 4/, female

2007.....	884.5	638.3	26.1	12.3	14.1	37.5	50.0	60.2	78.0	112.8	177.0
2006.....	891.7	641.4	26.9	12.8	13.2	38.0	50.6	60.3	77.8	114.9	182.4
2005.....	910.0	631.6	25.3	13.9	14.8	37.9	49.8	58.1	79.5	116.1	182.9
2004.....	898.0	628.9	28.1	13.4	15.9	41.7	49.7	58.1	77.5	116.0	182.7
2003.....	925.9	630.7	28.0	13.2	15.2	39.9	52.0	58.8	77.8	121.6	186.1
2002.....	929.5	640.1	27.2	13.7	16.2	41.6	49.1	57.8	77.8	121.9	186.7
2001.....	928.6	630.4	29.9	14.4	15.4	39.7	48.7	58.8	79.1	123.4	183.5
2000.....	931.0	680.1	28.8	14.0	16.4	41.0	49.3	55.6	77.1	116.8	178.2
1999.....	927.8	682.9	30.6	15.7	16.3	42.5	48.6	55.7	77.7	118.5	174.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All causes											
Non-Hispanic White, both sexes											
2007 5/.....	964.1	559.7	25.5	12.7	15.7	58.0	91.1	96.0	109.4	142.6	216.8
2006 5/.....	968.5	564.2	25.0	12.9	15.0	59.1	93.2	96.5	107.9	143.5	221.5
2005 5/.....	981.8	562.6	26.2	13.5	16.8	60.4	89.3	90.7	106.0	143.4	224.0
2004 5/.....	967.8	564.4	26.8	13.5	17.4	63.4	87.2	89.1	102.2	145.2	220.9
2003 5/.....	993.6	576.5	27.6	13.3	17.6	62.9	88.9	86.2	104.4	149.5	227.3
2002.....	997.5	575.9	27.1	13.7	18.1	65.5	86.6	86.5	103.9	151.1	228.1
2001.....	991.1	555.4	30.1	14.5	18.0	63.4	85.4	87.2	104.4	153.1	220.9
2000.....	993.2	596.5	28.5	14.7	19.4	64.0	84.1	80.6	98.7	144.5	216.4
1999.....	990.7	593.1	30.1	15.6	19.1	65.3	80.2	80.3	101.3	144.5	211.0
Non-Hispanic White, male											
2007 5/.....	960.4	616.8	28.1	13.7	18.3	77.2	132.2	135.7	146.3	181.9	271.1
2006 5/.....	962.0	621.9	26.7	14.2	18.0	79.0	136.5	137.2	145.2	183.0	278.0
2005 5/.....	970.6	625.7	29.9	14.5	20.1	82.0	130.1	128.2	140.0	182.0	283.2
2004 5/.....	957.4	625.2	29.0	15.2	20.4	83.4	126.6	126.2	135.2	185.6	279.0
2003 5/.....	979.1	647.2	30.2	14.7	21.2	84.3	128.3	119.2	139.4	192.2	289.4
2002.....	983.9	643.5	30.3	14.9	21.3	87.6	126.7	121.2	139.3	194.0	290.9
2001.....	975.6	611.6	33.7	15.2	21.7	86.1	125.6	122.4	139.3	197.9	282.4
2000.....	978.5	658.7	32.4	16.6	23.3	86.1	122.9	112.7	132.2	188.0	278.5
1999.....	979.6	658.1	33.4	17.0	22.7	87.1	116.1	112.1	135.7	187.1	272.8
Non-Hispanic White, female											
2007 5/.....	967.6	499.6	22.7	11.6	13.0	37.8	47.8	55.6	72.0	103.0	162.7
2006 5/.....	974.7	503.7	23.2	11.7	11.9	38.1	47.6	55.3	70.2	103.7	165.1
2005 5/.....	992.6	496.5	22.2	12.4	13.3	37.6	46.8	52.4	71.6	104.4	165.0
2004 5/.....	977.7	500.7	24.4	11.6	14.3	42.4	46.3	51.3	68.7	104.4	163.0
2003 5/.....	1,007.6	502.5	24.9	11.9	13.8	40.4	48.3	52.4	68.9	106.6	165.4
2002.....	1,010.6	504.8	23.8	12.3	14.8	42.0	45.6	51.3	68.1	108.0	165.4
2001.....	1,006.1	496.4	26.3	13.7	14.1	39.6	44.3	51.4	69.0	108.2	159.7
2000.....	1,007.3	530.9	24.4	12.6	15.2	41.0	44.5	48.0	64.7	100.8	154.5
1999.....	1,001.3	524.6	26.7	14.1	15.2	42.4	43.3	48.2	66.5	101.7	149.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All causes											
Non-Hispanic Black, both sexes											
2007 5/.....	750.7	1,309.1	43.7	18.6	24.6	85.7	142.2	161.9	192.6	254.0	366.0
2006 5/.....	759.1	1,339.2	44.3	20.6	23.1	87.1	146.7	171.3	201.6	262.6	384.5
2005 5/.....	774.4	1,323.9	42.5	21.8	25.9	86.2	147.6	168.3	204.6	263.2	389.7
2004 5/.....	768.8	1,297.2	45.5	22.3	26.2	83.7	143.3	174.2	198.6	263.3	397.2
2003 5/.....	788.8	1,290.0	48.0	20.6	26.7	82.3	152.2	175.4	201.8	285.1	421.1
2002.....	792.8	1,280.3	48.4	22.0	28.3	83.8	151.7	179.6	207.4	292.2	422.8
2001.....	798.1	1,354.4	49.5	21.4	26.6	88.7	156.7	174.9	207.1	294.6	438.1
2000.....	805.5	1,477.8	51.7	22.6	27.5	89.9	160.5	175.2	213.6	290.5	443.1
1999.....	812.1	1,510.2	54.6	26.2	29.0	94.8	160.6	169.8	212.0	301.4	457.4
Non-Hispanic Black, male											
2007 5/.....	805.1	1,426.3	47.5	21.1	29.8	131.4	219.1	236.6	267.0	326.5	458.5
2006 5/.....	815.3	1,453.3	48.2	23.7	27.6	134.7	223.0	253.7	279.9	336.8	471.1
2005 5/.....	825.7	1,451.5	47.6	23.6	31.6	131.3	227.8	248.7	280.3	334.8	476.2
2004 5/.....	818.7	1,427.6	49.5	23.8	29.7	124.6	216.9	255.3	269.0	334.7	480.0
2003 5/.....	840.6	1,427.7	55.1	22.8	32.6	124.3	232.0	259.9	273.1	357.4	520.3
2002.....	842.3	1,369.3	55.8	24.8	34.4	124.1	235.3	266.6	281.5	376.4	514.5
2001.....	849.7	1,484.2	55.1	24.6	32.5	133.9	244.7	255.7	279.6	375.1	538.9
2000.....	859.5	1,623.2	56.1	25.4	33.0	134.0	249.2	256.9	283.2	371.1	558.6
1999.....	872.8	1,650.8	61.5	30.1	36.3	142.3	249.7	247.5	288.0	382.5	583.2
Non-Hispanic Black, female											
2007 5/.....	701.0	1,187.1	39.8	15.9	19.1	38.8	63.4	90.3	125.1	189.3	284.7
2006 5/.....	708.0	1,220.1	40.3	17.3	18.4	38.2	68.6	93.5	130.9	196.5	308.4
2005 5/.....	727.6	1,190.8	37.2	20.0	19.9	39.9	66.2	92.5	136.3	199.2	313.6
2004 5/.....	723.4	1,161.6	41.3	20.7	22.7	41.7	69.4	98.6	135.2	199.5	324.4
2003 5/.....	741.6	1,146.9	40.7	18.4	20.6	39.1	73.3	97.5	137.5	220.8	333.8
2002.....	748.0	1,187.5	40.8	19.2	22.0	42.1	70.5	100.1	141.0	217.4	342.0
2001.....	751.2	1,217.8	43.6	18.1	20.5	42.0	72.2	101.4	142.2	223.2	349.0
2000.....	756.7	1,327.7	47.1	19.7	21.7	44.9	76.5	101.3	151.1	219.2	341.0
1999.....	757.3	1,365.4	47.4	22.1	21.5	46.3	76.1	99.5	144.0	229.5	346.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Septicemia (A40-A41)

All origins 3/, both sexes

2007.....	11.5	6.6	0.5	0.2	0.2	0.2	0.5	0.6	0.9	1.5	2.6
2006.....	11.4	6.5	0.5	0.2	0.2	0.2	0.4	0.6	0.8	1.5	2.4
2005.....	11.5	7.4	0.5	0.2	0.2	0.3	0.4	0.6	0.9	1.5	2.3
2004.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5	2.3
2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5	2.6
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5	2.3
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4	2.3
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4	2.5
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4	2.2

All origins 3/, male

2007.....	10.7	7.0	0.5	0.2	0.2	0.2	0.6	0.6	0.9	1.6	2.7
2006.....	10.5	6.8	0.5	0.2	0.3	0.2	0.4	0.6	0.9	1.5	2.6
2005.....	10.5	8.1	0.6	0.2	0.2	0.3	0.5	0.7	0.8	1.4	2.4
2004.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5	2.4
2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6	2.5
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5	2.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5	2.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4	2.6
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5	2.6

All origins 3/, female

2007.....	12.4	6.3	0.4	*	*	0.3	0.5	0.6	0.9	1.5	2.6
2006.....	12.3	6.2	0.6	*	*	0.3	0.4	0.6	0.8	1.6	2.2
2005.....	12.5	6.5	0.5	*	0.2	0.3	0.4	0.6	1.0	1.6	2.3
2004.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4	2.3
2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4	2.7
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5	2.2
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2	2.1
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4	2.3
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Septicemia (A40-A41)

Non-Hispanic 4/, both sexes

2007.....	12.8	6.7	0.5	0.2	0.2	0.3	0.6	0.7	0.9	1.7	2.9
2006.....	12.7	6.3	0.6	0.2	0.2	0.2	0.4	0.6	0.9	1.6	2.6
2005.....	12.8	7.4	0.6	0.2	0.2	0.3	0.4	0.7	1.0	1.6	2.4
2004.....	12.5	6.7	0.5	0.2	0.2	0.3	0.5	0.6	1.2	1.5	2.4
2003.....	12.9	7.2	0.6	0.2	0.2	0.3	0.5	0.7	0.9	1.6	2.7
2002.....	12.9	7.1	0.5	0.2	0.3	0.2	0.4	0.7	1.0	1.6	2.4
2001.....	12.4	8.1	0.7	0.1	0.2	0.3	0.4	0.6	1.0	1.5	2.3
2000.....	12.1	6.9	0.7	0.2	0.1	0.2	0.3	0.5	0.9	1.5	2.5
1999.....	12.0	7.9	0.6	0.2	0.2	0.2	0.6	0.6	0.9	1.4	2.3

Non-Hispanic 4/, male

2007.....	11.9	7.0	0.6	*	*	0.2	0.6	0.7	0.9	1.7	2.9
2006.....	11.7	6.5	0.6	*	0.3	0.2	0.4	0.7	0.9	1.6	2.8
2005.....	11.7	8.2	0.6	0.3	0.3	0.3	0.5	0.8	0.8	1.5	2.5
2004.....	11.5	7.9	0.4	*	*	0.3	0.5	0.6	1.2	1.6	2.4
2003.....	11.6	7.6	0.8	*	*	0.4	0.5	0.7	1.0	1.6	2.6
2002.....	11.6	7.3	0.5	0.2	0.3	*	0.4	0.7	1.0	1.5	2.5
2001.....	11.2	8.8	0.9	*	*	0.3	0.5	0.6	0.9	1.6	2.6
2000.....	10.7	6.7	0.7	0.2	*	0.2	0.4	0.7	1.0	1.5	2.6
1999.....	10.7	9.4	0.6	*	*	*	0.6	0.6	1.0	1.5	2.6

Non-Hispanic 4/, female

2007.....	13.7	6.4	0.4	*	*	0.3	0.5	0.6	0.9	1.6	2.8
2006.....	13.6	6.2	0.5	*	*	0.3	0.4	0.6	0.9	1.7	2.3
2005.....	13.8	6.6	0.5	*	*	0.3	0.4	0.7	1.1	1.7	2.3
2004.....	13.5	5.4	0.5	*	*	0.2	0.4	0.7	1.1	1.4	2.5
2003.....	14.2	6.8	0.4	0.3	*	0.3	0.5	0.8	0.9	1.5	2.8
2002.....	14.1	6.8	0.5	*	0.3	0.3	0.3	0.7	0.9	1.6	2.4
2001.....	13.5	7.4	0.6	*	*	0.3	0.4	0.5	1.1	1.3	2.1
2000.....	13.5	7.0	0.6	*	*	*	*	0.4	0.9	1.5	2.4
1999.....	13.2	6.2	0.5	0.3	*	*	0.5	0.6	0.8	1.3	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Septicemia (A40-A41)

Non-Hispanic White, both sexes

2007 5/.....	12.9	4.2	0.5	*	0.2	0.2	0.4	0.5	0.8	1.3	2.4
2006 5/.....	12.7	4.0	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.3	2.1
2005 5/.....	12.7	4.3	0.5	*	0.2	0.2	0.4	0.5	0.7	1.3	2.0
2004 5/.....	12.5	4.8	0.4	0.2	*	0.2	0.3	0.5	0.9	1.3	1.8
2003 5/.....	12.9	4.2	0.5	0.2	0.2	0.2	0.4	0.5	0.7	1.2	2.1
2002.....	12.9	5.5	0.5	0.2	0.2	0.2	0.3	0.6	0.8	1.3	2.0
2001.....	12.3	5.1	0.7	*	*	0.2	0.3	0.4	0.7	1.1	1.8
2000.....	11.9	4.8	0.5	0.2	*	0.2	0.2	0.4	0.7	1.1	2.0
1999.....	11.7	5.3	0.5	0.2	*	0.2	0.4	0.5	0.7	1.0	1.7

Non-Hispanic White, male

2007 5/.....	11.8	4.4	0.6	*	*	*	0.5	0.6	0.8	1.4	2.4
2006 5/.....	11.7	4.1	0.5	*	*	*	*	0.5	0.7	1.3	2.3
2005 5/.....	11.5	5.3	0.6	*	*	*	0.4	0.6	0.6	1.2	2.1
2004 5/.....	11.4	5.7	*	*	*	*	0.3	0.5	0.9	1.4	1.8
2003 5/.....	11.4	3.9	0.6	*	*	0.3	0.4	0.5	0.7	1.3	2.0
2002.....	11.5	5.6	0.5	*	*	*	0.4	0.6	0.7	1.2	2.0
2001.....	11.0	5.8	0.9	*	*	*	*	0.5	0.6	1.2	2.1
2000.....	10.4	4.7	0.5	*	*	*	0.3	0.4	0.7	1.2	2.1
1999.....	10.2	6.7	0.5	*	*	*	0.6	0.5	0.8	1.1	1.9

Non-Hispanic White, female

2007 5/.....	13.9	4.1	*	*	*	*	0.3	0.4	0.7	1.3	2.3
2006 5/.....	13.7	3.9	0.4	*	*	*	*	0.4	0.7	1.2	1.9
2005 5/.....	13.9	3.3	*	*	*	*	*	0.5	0.9	1.4	1.9
2004 5/.....	13.6	3.9	0.5	*	*	*	*	0.5	0.9	1.2	1.9
2003 5/.....	14.3	4.5	*	*	*	*	0.5	0.6	0.7	1.1	2.2
2002.....	14.2	5.4	*	*	*	0.3	*	0.5	0.8	1.3	2.0
2001.....	13.6	4.4	0.5	*	*	*	*	0.4	0.8	0.9	1.5
2000.....	13.4	4.8	0.6	*	*	*	*	*	0.6	1.0	1.9
1999.....	13.1	3.8	0.5	*	*	*	*	0.4	0.6	0.9	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Septicemia (A40-A41)

Non-Hispanic Black, both sexes

2007 5/.....	16.4	16.7	*	*	*	*	1.1	1.6	1.7	4.0	6.0
2006 5/.....	16.0	15.6	*	*	*	*	1.0	1.5	1.9	4.0	5.9
2005 5/.....	16.5	19.9	0.8	*	*	0.6	0.8	1.6	2.4	3.8	5.6
2004 5/.....	16.1	14.2	*	*	*	*	1.1	1.3	2.6	2.7	6.4
2003 5/.....	16.8	18.8	1.0	*	*	*	0.8	1.7	2.3	3.7	6.5
2002.....	16.8	13.1	*	*	*	*	*	1.6	2.3	3.5	5.6
2001.....	16.4	20.3	0.9	*	*	*	0.8	1.1	2.7	3.9	5.7
2000.....	16.7	16.6	1.1	*	*	*	0.8	1.5	2.6	3.8	5.9
1999.....	16.6	18.6	*	*	*	*	1.0	1.4	1.8	3.8	6.4

Non-Hispanic Black, male

2007 5/.....	15.4	17.9	*	*	*	*	*	1.5	*	3.9	6.0
2006 5/.....	14.9	16.3	*	*	*	*	*	*	2.2	3.7	6.6
2005 5/.....	15.5	20.2	*	*	*	*	*	1.8	2.1	3.7	5.7
2004 5/.....	15.1	15.8	*	*	*	*	*	*	3.0	2.6	7.0
2003 5/.....	15.8	21.7	*	*	*	*	*	1.6	2.9	3.8	6.5
2002.....	15.5	13.9	*	*	*	*	*	*	2.5	3.8	6.1
2001.....	15.4	21.4	*	*	*	*	*	*	2.6	3.9	5.9
2000.....	15.3	16.4	*	*	*	*	*	2.2	2.6	3.6	6.0
1999.....	15.7	20.8	*	*	*	*	*	*	2.2	4.1	7.9

Non-Hispanic Black, female

2007 5/.....	17.2	15.5	*	*	*	*	*	1.6	1.9	4.1	6.1
2006 5/.....	17.0	14.8	*	*	*	*	*	1.6	1.6	4.2	5.3
2005 5/.....	17.4	19.5	*	*	*	*	*	1.5	2.6	3.8	5.6
2004 5/.....	17.0	12.6	*	*	*	*	*	*	2.2	2.9	5.9
2003 5/.....	17.7	15.8	*	*	*	*	*	1.7	1.7	3.7	6.4
2002.....	18.0	12.3	*	*	*	*	*	2.0	2.1	3.1	5.2
2001.....	17.2	19.1	*	*	*	*	*	*	2.7	3.9	5.6
2000.....	17.9	16.9	*	*	*	*	*	*	2.6	4.1	5.8
1999.....	17.4	16.4	*	*	*	*	*	1.7	1.5	3.6	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Viral hepatitis (B15-B19)

All origins 3/, both sexes

2007.....	2.5	*	*	*	*	*	*	0.1	0.2	0.5	1.7
2006.....	2.4	*	*	*	*	*	*	*	0.2	0.6	1.8
2005.....	1.9	*	*	*	*	*	*	0.1	0.2	0.6	1.7
2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7	2.0
2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8	2.1
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0	2.3
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9	2.6
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0	2.5
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9	2.3

All origins 3/, male

2007.....	3.3	*	*	*	*	*	*	*	0.3	0.8	2.2
2006.....	3.2	*	*	*	*	*	*	*	0.3	0.8	2.5
2005.....	2.5	*	*	*	*	*	*	*	*	0.8	2.3
2004.....	2.4	*	*	*	*	*	*	*	0.2	0.8	2.8
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2	2.9
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4	3.2
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2	3.7
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3	3.5
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2	3.5

All origins 3/, female

2007.....	1.6	*	*	*	*	*	*	*	*	0.3	1.2
2006.....	1.6	*	*	*	*	*	*	*	*	0.4	1.1
2005.....	1.3	*	*	*	*	*	*	*	*	0.4	1.1
2004.....	1.3	*	*	*	*	*	*	*	*	0.5	1.3
2003.....	1.3	*	*	*	*	*	*	*	*	0.4	1.2
2002.....	1.4	*	*	*	*	*	*	*	*	0.6	1.4
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Viral hepatitis (B15-B19)

Non-Hispanic 4/, both sexes

2007.....	2.4	*	*	*	*	*	*	*	0.2	0.5	1.6
2006.....	2.4	*	*	*	*	*	*	*	0.2	0.6	1.7
2005.....	1.9	*	*	*	*	*	*	*	0.2	0.5	1.5
2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7	1.9
2003.....	1.9	*	*	*	*	*	*	*	0.2	0.8	2.0
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0	2.2
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.9	2.5
2000.....	1.9	*	*	*	*	*	*	0.1	0.2	0.9	2.3
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.8	2.2

Non-Hispanic 4/, male

2007.....	3.3	*	*	*	*	*	*	*	0.3	0.8	1.9
2006.....	3.2	*	*	*	*	*	*	*	0.3	0.8	2.2
2005.....	2.5	*	*	*	*	*	*	*	*	0.7	2.0
2004.....	2.5	*	*	*	*	*	*	*	*	0.8	2.4
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2	2.8
2002.....	2.7	*	*	*	*	*	*	*	0.4	1.4	3.1
2001.....	2.5	*	*	*	*	*	*	*	0.4	1.1	3.5
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.2	3.2
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.1	3.3

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	*	*	*	*	0.3	1.2
2006.....	1.6	*	*	*	*	*	*	*	*	0.4	1.1
2005.....	1.3	*	*	*	*	*	*	*	*	0.3	1.0
2004.....	1.3	*	*	*	*	*	*	*	*	0.5	1.3
2003.....	1.3	*	*	*	*	*	*	*	*	0.4	1.2
2002.....	1.4	*	*	*	*	*	*	*	*	0.6	1.4
2001.....	1.4	*	*	*	*	*	*	*	*	0.6	1.5
2000.....	1.4	*	*	*	*	*	*	*	*	0.6	1.5
1999.....	1.2	*	*	*	*	*	*	*	0.3	0.6	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Viral hepatitis (B15-B19)

Non-Hispanic White, both sexes

2007 5/.....	2.4	*	*	*	*	*	*	*	0.2	0.5	1.5
2006 5/.....	2.3	*	*	*	*	*	*	*	0.2	0.5	1.7
2005 5/.....	1.8	*	*	*	*	*	*	*	*	0.5	1.5
2004 5/.....	1.8	*	*	*	*	*	*	*	0.2	0.6	1.9
2003 5/.....	1.8	*	*	*	*	*	*	*	0.2	0.8	1.9
2002.....	1.9	*	*	*	*	*	*	*	0.2	1.0	2.1
2001.....	1.8	*	*	*	*	*	*	*	0.2	0.9	2.4
2000.....	1.8	*	*	*	*	*	*	*	0.2	0.9	2.2
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.8	2.0

Non-Hispanic White, male

2007 5/.....	3.3	*	*	*	*	*	*	*	*	0.8	1.8
2006 5/.....	3.1	*	*	*	*	*	*	*	*	0.6	2.2
2005 5/.....	2.5	*	*	*	*	*	*	*	*	0.7	2.1
2004 5/.....	2.4	*	*	*	*	*	*	*	*	0.7	2.5
2003 5/.....	2.4	*	*	*	*	*	*	*	*	1.2	2.7
2002.....	2.6	*	*	*	*	*	*	*	0.4	1.5	3.0
2001.....	2.5	*	*	*	*	*	*	*	0.4	1.2	3.4
2000.....	2.3	*	*	*	*	*	*	*	*	1.2	3.0
1999.....	2.1	*	*	*	*	*	*	*	0.4	1.1	3.0

Non-Hispanic White, female

2007 5/.....	1.6	*	*	*	*	*	*	*	*	0.3	1.2
2006 5/.....	1.5	*	*	*	*	*	*	*	*	0.4	1.1
2005 5/.....	1.2	*	*	*	*	*	*	*	*	0.3	1.0
2004 5/.....	1.2	*	*	*	*	*	*	*	*	0.5	1.3
2003 5/.....	1.2	*	*	*	*	*	*	*	*	0.5	1.1
2002.....	1.3	*	*	*	*	*	*	*	*	0.5	1.3
2001.....	1.2	*	*	*	*	*	*	*	*	0.6	1.4
2000.....	1.3	*	*	*	*	*	*	*	*	0.6	1.4
1999.....	1.1	*	*	*	*	*	*	*	*	0.5	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes

2007.....	3.7	*	*	*	*	0.2	0.6	2.0	3.5	6.5	10.0
2006.....	4.0	*	*	*	*	0.2	0.8	2.0	3.9	7.5	10.8
2005.....	4.2	*	*	*	*	0.1	0.6	2.1	4.5	8.1	11.7
2004.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5	12.3
2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2	12.8
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9	13.4
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3	13.7
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8	13.5
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0	14.6

All origins 3/, male

2007.....	5.4	*	*	*	*	*	0.7	2.3	4.1	8.7	14.4
2006.....	5.9	*	*	*	*	0.2	0.9	2.4	4.7	10.1	15.6
2005.....	6.3	*	*	*	*	*	0.7	2.4	5.5	11.2	17.2
2004.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3	17.9
2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9	19.1
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7	19.8
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4	20.5
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2	20.4
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5	22.6

All origins 3/, female

2007.....	2.1	*	*	*	*	*	0.5	1.6	2.8	4.4	5.5
2006.....	2.2	*	*	*	*	*	0.7	1.6	3.0	4.9	5.9
2005.....	2.2	*	*	*	*	*	0.5	1.7	3.5	4.9	6.2
2004.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6	6.7
2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5	6.6
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2	7.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3	7.0
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4	6.7
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6	6.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, both sexes

2007.....	3.3	*	*	*	*	*	0.8	1.5	3.0	5.0	10.1
2006.....	3.6	*	*	*	*	*	0.6	1.5	3.4	7.2	10.6
2005.....	3.9	*	*	*	*	*	0.8	1.5	3.4	7.2	12.4
2004.....	4.3	*	*	*	*	*	*	1.7	3.9	9.7	12.4
2003.....	4.7	*	*	*	*	*	0.6	2.0	4.7	9.8	15.2
2002.....	4.8	*	*	*	*	*	0.6	2.1	5.1	11.2	16.1
2001.....	5.0	*	*	*	*	*	0.8	2.3	6.6	11.9	16.3
2000.....	5.4	*	*	*	*	*	*	2.9	8.5	13.5	16.5
1999.....	5.8	*	*	*	*	*	0.9	3.5	9.6	15.5	19.7

Hispanic, male

2007.....	5.0	*	*	*	*	*	1.1	2.1	3.9	7.7	14.7
2006.....	5.6	*	*	*	*	*	*	2.0	4.9	10.5	15.7
2005.....	5.9	*	*	*	*	*	1.1	2.1	4.7	10.9	19.0
2004.....	6.4	*	*	*	*	*	*	2.4	5.9	14.4	18.1
2003.....	7.1	*	*	*	*	*	*	2.6	6.7	14.2	21.9
2002.....	7.2	*	*	*	*	*	*	2.9	7.4	16.9	23.9
2001.....	7.6	*	*	*	*	*	*	2.9	9.4	17.5	24.9
2000.....	8.2	*	*	*	*	*	*	3.5	12.6	21.0	24.9
1999.....	8.7	*	*	*	*	*	*	4.9	13.9	23.8	29.2

Hispanic, female

2007.....	1.5	*	*	*	*	*	*	*	1.8	1.9	5.1
2006.....	1.6	*	*	*	*	*	*	*	1.6	3.3	4.9
2005.....	1.6	*	*	*	*	*	*	*	1.9	2.9	5.2
2004.....	1.9	*	*	*	*	*	*	*	1.7	4.5	6.1
2003.....	2.2	*	*	*	*	*	*	1.2	2.4	4.8	7.9
2002.....	2.2	*	*	*	*	*	*	1.2	2.4	4.8	7.8
2001.....	2.3	*	*	*	*	*	*	1.5	3.5	5.8	7.3
2000.....	2.4	*	*	*	*	*	*	2.2	3.9	5.4	7.6
1999.....	2.6	*	*	*	*	*	*	1.9	4.6	6.4	9.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 4/, both sexes

2007.....	3.8	*	*	*	*	0.2	0.6	2.1	3.6	6.8	9.9
2006.....	4.1	*	*	*	*	0.2	0.8	2.1	4.0	7.5	10.7
2005.....	4.3	*	*	*	*	0.1	0.6	2.2	4.7	8.2	11.4
2004.....	4.4	*	*	*	*	0.1	0.8	2.3	5.3	9.4	12.2
2003.....	4.6	*	*	*	0.1	0.1	0.7	2.4	5.7	11.3	12.3
2002.....	4.8	*	*	*	*	0.2	0.7	2.7	6.6	11.9	12.9
2001.....	4.9	*	*	*	*	0.2	1.0	3.2	7.4	12.3	13.2
2000.....	5.0	*	*	*	0.1	0.2	0.8	3.4	8.7	12.5	12.9
1999.....	5.2	*	*	0.2	0.1	0.2	0.9	3.9	9.5	12.6	13.9

Non-Hispanic 4/, male

2007.....	5.5	*	*	*	*	*	0.6	2.3	4.1	8.9	14.3
2006.....	6.0	*	*	*	*	*	0.9	2.5	4.7	10.0	15.5
2005.....	6.3	*	*	*	*	*	0.6	2.5	5.7	11.2	16.7
2004.....	6.5	*	*	*	*	*	0.9	2.6	6.4	13.1	17.7
2003.....	7.0	*	*	*	*	*	0.7	2.8	7.4	16.1	18.5
2002.....	7.3	*	*	*	*	*	0.7	3.0	8.7	17.6	19.0
2001.....	7.5	*	*	*	*	*	0.8	3.8	10.3	18.4	19.6
2000.....	7.7	*	*	*	*	*	0.8	3.8	11.7	18.6	19.5
1999.....	8.0	*	*	*	*	*	0.8	5.1	13.4	18.7	21.4

Non-Hispanic 4/, female

2007.....	2.2	*	*	*	*	*	0.6	1.8	3.0	4.8	5.5
2006.....	2.3	*	*	*	*	*	0.7	1.8	3.3	5.1	6.1
2005.....	2.3	*	*	*	*	*	0.6	1.8	3.8	5.2	6.3
2004.....	2.4	*	*	*	*	*	0.8	2.0	4.1	5.7	6.7
2003.....	2.4	*	*	*	*	*	0.7	2.0	3.9	6.6	6.4
2002.....	2.5	*	*	*	*	*	0.7	2.5	4.5	6.3	7.0
2001.....	2.5	*	*	*	*	*	1.1	2.6	4.4	6.3	6.9
2000.....	2.5	*	*	*	*	*	0.8	3.0	5.7	6.4	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.8	5.5	6.6	6.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic White, both sexes

2007 5/.....	1.6	*	*	*	*	*	*	0.5	1.1	2.7	4.4
2006 5/.....	1.8	*	*	*	*	*	0.2	0.5	1.3	3.2	5.0
2005 5/.....	1.9	*	*	*	*	*	*	0.5	1.7	3.4	5.6
2004 5/.....	1.9	*	*	*	*	*	0.2	0.5	1.8	4.4	5.8
2003 5/.....	2.1	*	*	*	*	*	0.2	0.5	2.4	5.1	6.0
2002.....	2.1	*	*	*	*	*	*	0.9	2.4	5.7	5.8
2001.....	2.2	*	*	*	*	*	0.2	0.9	3.0	6.0	6.1
2000.....	2.3	*	*	*	*	*	0.2	1.1	3.5	6.3	6.1
1999.....	2.4	*	*	*	*	*	0.2	1.5	4.4	6.5	6.4

Non-Hispanic White, male

2007 5/.....	2.7	*	*	*	*	*	*	0.6	1.6	4.3	7.6
2006 5/.....	3.0	*	*	*	*	*	*	0.7	1.8	4.9	8.5
2005 5/.....	3.1	*	*	*	*	*	*	0.7	2.5	5.6	9.4
2004 5/.....	3.3	*	*	*	*	*	*	0.7	2.9	6.9	9.8
2003 5/.....	3.6	*	*	*	*	*	*	0.9	3.7	8.4	10.2
2002.....	3.7	*	*	*	*	*	*	1.3	3.8	9.6	9.9
2001.....	3.8	*	*	*	*	*	*	1.3	4.9	10.2	10.5
2000.....	4.0	*	*	*	*	*	*	1.5	5.6	10.6	10.2
1999.....	4.2	*	*	*	*	*	*	2.2	7.3	10.9	10.9

Non-Hispanic White, female

2007 5/.....	0.5	*	*	*	*	*	*	*	0.7	1.1	1.3
2006 5/.....	0.6	*	*	*	*	*	*	*	0.7	1.4	1.5
2005 5/.....	0.6	*	*	*	*	*	*	0.4	1.0	1.2	1.9
2004 5/.....	0.6	*	*	*	*	*	*	*	0.8	1.9	1.8
2003 5/.....	0.6	*	*	*	*	*	*	*	1.0	1.8	1.8
2002.....	0.6	*	*	*	*	*	*	0.5	1.0	1.8	1.7
2001.....	0.6	*	*	*	*	*	*	0.5	1.2	1.7	1.7
2000.....	0.7	*	*	*	*	*	*	0.7	1.5	2.1	1.9
1999.....	0.7	*	*	*	*	*	*	0.8	1.5	2.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic Black, both sexes

2007 5/.....	16.8	*	*	*	*	0.7	2.7	9.8	16.6	30.4	43.0
2006 5/.....	17.9	*	*	*	*	0.9	3.7	10.4	18.6	33.1	45.9
2005 5/.....	18.5	*	*	*	*	0.7	2.9	10.1	21.3	35.7	48.1
2004 5/.....	19.4	*	*	*	*	0.8	3.9	11.4	24.3	38.7	52.8
2003 5/.....	20.1	*	*	*	*	0.7	3.3	11.6	24.3	47.5	53.3
2002.....	21.3	*	*	*	*	0.8	3.4	12.2	30.2	49.3	58.6
2001.....	21.6	*	*	*	*	1.1	4.6	15.2	32.3	50.4	59.7
2000.....	22.0	*	*	*	*	0.7	3.9	15.8	38.4	51.0	58.2
1999.....	22.3	*	*	0.8	*	0.7	4.2	17.2	39.1	51.3	63.5

Non-Hispanic Black, male

2007 5/.....	22.7	*	*	*	*	*	2.9	10.7	18.5	37.0	58.6
2006 5/.....	24.4	*	*	*	*	*	4.3	11.9	21.0	41.8	62.3
2005 5/.....	25.8	*	*	*	*	*	3.1	11.6	24.7	46.2	66.6
2004 5/.....	26.7	*	*	*	*	*	4.0	12.6	27.5	51.9	71.8
2003 5/.....	28.5	*	*	*	*	*	3.3	13.2	30.0	63.9	75.9
2002.....	30.3	*	*	*	*	*	3.5	12.2	38.0	69.9	81.3
2001.....	30.8	*	*	*	*	*	3.7	17.6	43.5	71.6	84.0
2000.....	31.7	*	*	*	*	*	3.8	17.1	49.4	73.0	85.3
1999.....	32.6	*	*	*	*	*	3.5	21.8	52.2	71.7	96.1

Non-Hispanic Black, female

2007 5/.....	11.4	*	*	*	*	*	2.6	9.0	14.9	24.6	29.3
2006 5/.....	12.1	*	*	*	*	*	3.1	9.0	16.5	25.4	31.4
2005 5/.....	11.8	*	*	*	*	*	2.6	8.7	18.2	26.3	31.8
2004 5/.....	12.8	*	*	*	*	*	3.8	10.3	21.4	27.0	36.0
2003 5/.....	12.6	*	*	*	*	*	3.3	10.2	19.1	33.0	33.4
2002.....	13.2	*	*	*	*	*	3.4	12.3	23.2	31.1	38.6
2001.....	13.3	*	*	*	*	*	5.4	13.0	22.3	31.5	38.2
2000.....	13.1	*	*	*	*	*	4.0	14.6	28.5	31.5	34.3
1999.....	13.0	*	*	*	*	*	4.9	13.1	27.4	33.2	34.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/, both sexes

2007.....	1.9	3.6	0.4	0.1	0.1	0.2	0.3	0.3	0.4	0.5	0.6
2006.....	2.0	4.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.5	0.8
2005.....	2.6	4.1	0.4	0.2	0.1	0.2	0.2	0.2	0.4	0.7	1.1
2004.....	2.6	4.1	0.4	0.1	0.1	0.2	0.3	0.3	0.4	0.7	1.2
2003.....	2.6	4.3	0.5	0.1	0.1	0.2	0.2	0.3	0.4	0.7	1.3
2002.....	2.3	5.6	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7	0.9
2000.....	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7	0.8
1999.....	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4

All origins 3/, male

2007.....	2.0	3.9	0.4	*	0.2	0.2	0.3	0.4	0.4	0.5	0.7
2006.....	2.1	4.5	0.5	*	*	0.2	0.2	0.3	0.3	0.6	1.0
2005.....	2.9	4.8	0.5	*	*	0.2	0.2	0.3	0.5	0.9	1.5
2004.....	2.8	4.8	0.4	*	*	0.2	0.4	0.2	0.6	0.8	1.5
2003.....	2.8	4.2	0.5	*	*	0.3	0.2	0.3	0.5	0.9	1.6
2002.....	2.6	6.1	0.5	*	0.2	0.2	0.3	0.4	0.5	0.9	1.7
2001.....	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8	1.2
2000.....	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9	1.0
1999.....	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3	1.7

All origins 3/, female

2007.....	1.9	3.4	0.4	*	*	*	0.2	0.2	0.3	0.5	0.5
2006.....	1.9	3.6	0.3	*	0.2	0.2	0.2	0.2	0.4	0.3	0.6
2005.....	2.3	3.4	0.3	0.2	*	0.2	0.2	0.2	0.3	0.5	0.7
2004.....	2.3	3.4	0.4	*	*	0.2	*	0.4	0.3	0.6	0.9
2003.....	2.3	4.5	0.5	*	*	*	*	0.3	0.3	0.6	0.9
2002.....	2.1	5.0	0.3	0.3	*	0.2	0.2	0.3	0.4	0.6	0.8
2001.....	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6	0.6
2000.....	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5	0.6
1999.....	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 4/, both sexes

2007.....	2.1	3.5	0.4	0.1	0.1	0.2	0.3	0.2	0.3	0.5	0.6
2006.....	2.1	4.4	0.4	0.1	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2005.....	2.7	4.3	0.4	0.2	*	0.2	0.2	0.2	0.4	0.6	1.0
2004.....	2.7	4.3	0.4	*	0.1	0.2	0.2	0.3	0.4	0.7	1.1
2003.....	2.7	4.8	0.5	0.1	*	0.2	0.2	0.3	0.4	0.7	1.2
2002.....	2.4	5.7	0.4	0.2	0.2	0.2	0.3	0.3	0.4	0.8	1.2
2001.....	2.1	4.6	0.5	0.2	0.1	0.2	0.2	0.3	0.4	0.7	0.9
2000.....	2.1	4.7	0.4	0.1	0.2	0.2	0.2	0.3	0.4	0.7	0.8
1999.....	2.8	5.5	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4

Non-Hispanic 4/, male

2007.....	2.2	3.8	0.3	*	*	*	0.3	0.3	0.4	0.5	0.7
2006.....	2.2	4.7	0.5	*	*	*	*	0.3	0.3	0.6	0.9
2005.....	3.0	5.1	0.5	*	*	*	*	*	0.5	0.8	1.4
2004.....	3.0	5.2	0.4	*	*	*	0.3	*	0.6	0.8	1.3
2003.....	2.9	4.8	0.5	*	*	0.3	0.2	0.3	0.5	0.8	1.5
2002.....	2.7	5.9	0.5	*	*	*	0.3	0.4	0.4	0.9	1.6
2001.....	2.2	4.8	0.5	*	*	*	0.3	0.3	0.5	0.8	1.2
2000.....	2.2	5.1	0.5	*	*	0.3	*	0.3	0.4	0.9	1.0
1999.....	2.8	6.2	0.6	*	0.2	0.2	0.3	0.4	0.8	1.2	1.6

Non-Hispanic 4/, female

2007.....	2.0	3.2	0.4	*	*	*	*	*	0.3	0.5	0.5
2006.....	2.1	4.1	*	*	*	*	0.3	0.3	0.3	0.3	0.5
2005.....	2.5	3.4	*	*	*	*	*	*	0.3	0.5	0.7
2004.....	2.4	3.3	0.3	*	*	0.2	*	0.4	0.3	0.6	0.9
2003.....	2.5	4.8	0.5	*	*	*	*	0.3	0.3	0.6	0.8
2002.....	2.2	5.4	*	0.3	*	*	*	*	0.4	0.6	0.8
2001.....	2.1	4.5	0.4	*	*	*	*	0.3	0.4	0.6	0.6
2000.....	2.0	4.3	0.4	*	*	*	*	0.3	0.4	0.5	0.6
1999.....	2.7	4.7	0.4	*	*	*	*	0.5	0.6	0.8	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic White, both sexes

2007 5/.....	2.2	2.9	0.4	*	*	*	0.2	0.2	0.3	0.4	0.4
2006 5/.....	2.2	2.8	0.4	*	0.2	0.2	0.2	0.3	0.2	0.3	0.6
2005 5/.....	2.8	3.7	0.4	0.2	*	*	0.2	*	0.3	0.6	0.9
2004 5/.....	2.8	3.7	0.3	*	*	0.2	0.2	0.2	0.3	0.6	1.0
2003 5/.....	2.8	3.8	0.4	*	*	0.2	0.2	0.3	0.4	0.6	1.1
2002.....	2.5	4.8	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.6	1.0
2001.....	2.1	3.3	0.4	0.2	*	0.2	0.2	0.2	0.3	0.6	0.8
2000.....	2.1	3.5	0.4	*	0.2	0.2	0.2	0.2	0.3	0.5	0.7
1999.....	2.7	4.4	0.5	0.2	*	0.2	0.2	0.3	0.5	0.7	1.0

Non-Hispanic White, male

2007 5/.....	2.2	3.1	*	*	*	*	*	*	*	0.4	0.5
2006 5/.....	2.3	3.1	0.4	*	*	*	*	*	*	0.5	0.7
2005 5/.....	3.1	4.4	0.4	*	*	*	*	*	0.3	0.8	1.2
2004 5/.....	3.0	4.6	*	*	*	*	*	*	0.4	0.7	1.1
2003 5/.....	2.9	3.6	*	*	*	*	*	*	0.4	0.7	1.4
2002.....	2.7	4.8	0.4	*	*	*	*	*	*	0.7	1.3
2001.....	2.2	3.3	0.4	*	*	*	*	*	0.4	0.6	1.0
2000.....	2.2	3.9	*	*	*	*	*	*	*	0.6	0.8
1999.....	2.7	5.5	0.6	*	*	*	*	*	0.5	1.0	1.3

Non-Hispanic White, female

2007 5/.....	2.1	2.7	*	*	*	*	*	*	*	0.3	0.3
2006 5/.....	2.2	2.4	*	*	*	*	*	*	*	*	0.5
2005 5/.....	2.5	2.9	*	*	*	*	*	*	*	0.3	0.6
2004 5/.....	2.5	2.7	*	*	*	*	*	*	*	0.6	0.8
2003 5/.....	2.6	4.0	*	*	*	*	*	*	*	0.5	0.8
2002.....	2.3	4.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	2.1	3.4	*	*	*	*	*	*	*	0.5	0.5
2000.....	2.0	3.1	*	*	*	*	*	*	0.3	0.4	0.5
1999.....	2.6	3.3	*	*	*	*	*	0.4	0.4	0.5	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic Black, both sexes

2007 5/.....	2.0	5.9	*	*	*	*	*	*	1.0	1.1	1.4
2006 5/.....	2.2	10.2	*	*	*	*	*	*	1.0	1.1	1.5
2005 5/.....	2.6	6.8	*	*	*	*	*	*	1.1	1.2	1.8
2004 5/.....	2.6	6.8	*	*	*	*	*	*	1.0	1.2	2.1
2003 5/.....	2.6	8.4	0.9	*	*	*	*	*	0.8	1.5	2.1
2002.....	2.6	9.0	*	*	*	*	*	*	1.1	1.9	2.5
2001.....	2.3	10.0	*	*	*	*	*	*	1.1	1.5	1.8
2000.....	2.3	10.4	*	*	*	*	*	*	1.3	1.7	1.7
1999.....	3.6	10.0	1.0	*	*	*	*	1.2	2.2	2.6	3.6

Non-Hispanic Black, male

2007 5/.....	2.2	6.5	*	*	*	*	*	*	*	*	*
2006 5/.....	2.3	10.5	*	*	*	*	*	*	*	*	2.2
2005 5/.....	2.9	8.4	*	*	*	*	*	*	*	*	2.7
2004 5/.....	3.0	7.4	*	*	*	*	*	*	*	*	2.5
2003 5/.....	3.0	9.3	*	*	*	*	*	*	*	1.6	2.7
2002.....	3.0	10.0	*	*	*	*	*	*	*	2.1	3.1
2001.....	2.5	11.5	*	*	*	*	*	*	*	1.7	2.6
2000.....	2.6	10.6	*	*	*	*	*	*	*	2.3	2.2
1999.....	3.7	10.0	*	*	*	*	*	*	2.7	2.7	3.6

Non-Hispanic Black, female

2007 5/.....	1.8	*	*	*	*	*	*	*	*	*	1.4
2006 5/.....	2.1	10.0	*	*	*	*	*	*	*	*	*
2005 5/.....	2.3	*	*	*	*	*	*	*	*	*	*
2004 5/.....	2.3	*	*	*	*	*	*	*	*	*	1.8
2003 5/.....	2.2	7.4	*	*	*	*	*	*	*	*	1.5
2002.....	2.2	7.9	*	*	*	*	*	*	*	1.7	1.9
2001.....	2.1	8.4	*	*	*	*	*	*	*	*	*
2000.....	2.1	10.2	*	*	*	*	*	*	*	*	*
1999.....	3.5	10.0	*	*	*	*	*	*	1.7	2.5	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms (C00-C97)

All origins 3/, both sexes

2007.....	186.6	1.7	2.2	2.4	2.4	3.1	4.7	6.3	11.0	20.0	41.2
2006.....	187.0	1.8	2.3	2.3	2.2	3.2	4.6	6.7	11.5	20.8	42.3
2005.....	188.7	1.8	2.3	2.5	2.5	3.5	4.7	6.7	11.3	20.8	44.6
2004.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5	44.3
2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0	46.1
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5	47.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6	48.7
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5	48.8
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2	49.4

All origins 3/, male

2007.....	197.0	1.8	2.3	2.5	2.3	3.6	5.5	6.6	9.9	16.5	36.0
2006.....	196.6	1.8	2.5	2.6	2.4	3.8	5.4	7.0	10.3	17.2	37.1
2005.....	198.9	2.1	2.6	2.6	2.8	4.1	5.5	7.3	10.4	17.4	39.5
2004.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1	39.2
2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3	41.7
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8	42.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0	44.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2	44.4
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6	45.5

All origins 3/, female

2007.....	176.5	1.6	2.2	2.3	2.4	2.7	3.8	5.9	12.1	23.6	46.3
2006.....	177.6	1.8	2.1	2.1	2.0	2.5	3.7	6.4	12.7	24.4	47.6
2005.....	178.8	1.5	2.0	2.3	2.2	2.8	3.9	6.0	12.2	24.3	49.6
2004.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9	49.2
2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7	50.5
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2	52.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1	53.5
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8	53.2
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7	53.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Malignant neoplasms (C00-C97)											
Hispanic, both sexes											
2007.....	60.8	*	1.8	2.8	2.8	3.7	5.6	5.5	9.5	16.0	28.2
2006.....	60.1	*	2.2	2.2	2.4	3.4	5.3	5.8	8.6	15.5	28.6
2005.....	61.3	*	2.2	2.4	2.6	4.0	4.9	5.5	8.2	14.7	30.5
2004.....	59.3	*	2.0	2.6	3.0	4.1	4.5	5.0	8.9	15.9	28.6
2003.....	60.3	*	3.0	2.9	2.9	3.8	4.4	5.5	8.8	16.2	30.7
2002.....	59.7	*	2.3	2.8	3.0	3.8	5.1	4.7	9.1	15.8	31.9
2001.....	60.5	*	2.8	2.4	2.7	4.5	4.1	5.7	8.9	18.3	34.4
2000.....	59.9	*	2.5	2.5	3.4	3.8	4.5	5.8	9.0	18.2	33.9
1999.....	59.6	*	2.5	2.5	3.0	3.1	5.1	5.8	9.1	17.3	33.3
Hispanic, male											
2007.....	61.6	*	1.9	2.8	2.6	4.5	6.2	5.2	7.3	12.1	23.6
2006.....	60.4	*	2.6	2.4	2.2	3.8	6.4	5.6	6.5	11.0	21.8
2005.....	63.0	*	2.8	2.3	2.6	5.0	5.5	5.6	7.5	11.3	25.3
2004.....	60.9	*	1.6	2.9	3.1	5.5	5.5	4.8	8.2	13.0	25.1
2003.....	61.5	*	3.6	3.6	2.8	5.4	4.6	5.5	8.2	12.7	24.8
2002.....	61.2	*	2.2	2.7	3.1	4.9	5.5	5.0	7.8	11.9	26.3
2001.....	62.2	*	2.9	2.6	2.4	5.7	5.4	5.2	7.5	15.1	29.9
2000.....	61.3	*	2.9	3.0	4.1	4.3	4.8	5.7	8.2	13.8	28.0
1999.....	61.2	*	2.3	2.5	3.1	3.4	5.8	5.6	7.6	13.9	27.2
Hispanic, female											
2007.....	59.9	*	1.6	2.7	3.0	2.9	4.9	5.8	12.1	20.5	33.3
2006.....	59.7	*	1.8	2.0	2.5	2.9	4.0	6.1	11.0	20.6	36.1
2005.....	59.5	*	1.6	2.4	2.5	3.0	4.1	5.3	8.9	18.6	36.2
2004.....	57.7	*	2.4	2.2	2.8	2.7	3.3	5.3	9.7	19.1	32.3
2003.....	59.1	*	2.4	2.2	3.0	2.1	4.1	5.4	9.4	20.0	37.1
2002.....	58.1	*	2.4	2.9	2.9	2.6	4.5	4.4	10.6	20.2	37.9
2001.....	58.7	*	2.7	2.1	3.0	3.2	2.7	6.4	10.4	21.8	39.2
2000.....	58.5	*	2.2	2.0	2.6	3.4	4.1	5.8	9.9	23.0	40.0
1999.....	58.0	*	2.7	2.5	2.9	2.7	4.2	6.1	10.7	20.9	39.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms (C00-C97)

Non-Hispanic 4/, both sexes

2007.....	208.7	1.6	2.4	2.3	2.3	3.0	4.5	6.5	11.4	20.8	43.3
2006.....	208.7	1.8	2.3	2.4	2.1	3.1	4.4	6.9	12.2	21.8	44.5
2005.....	209.8	1.8	2.3	2.5	2.4	3.4	4.6	6.9	12.0	22.0	46.6
2004.....	209.4	1.7	2.6	2.7	2.2	3.4	4.7	6.9	12.0	22.4	46.4
2003.....	211.9	2.0	2.3	2.5	2.6	3.3	4.7	7.1	12.3	24.1	48.1
2002.....	213.4	1.7	2.6	2.7	2.4	3.5	4.9	7.1	13.1	24.7	49.4
2001.....	213.9	1.6	2.7	2.4	2.4	3.4	5.1	7.7	13.3	25.4	50.3
2000.....	215.6	2.5	2.7	2.3	2.4	3.6	5.2	7.2	13.1	25.3	50.3
1999.....	215.5	1.7	2.7	2.5	2.4	3.8	5.3	7.6	13.0	26.2	51.0

Non-Hispanic 4/, male

2007.....	222.2	1.9	2.4	2.4	2.3	3.4	5.3	7.0	10.6	17.5	38.1
2006.....	221.3	1.7	2.5	2.6	2.4	3.8	5.2	7.4	11.3	18.5	39.6
2005.....	222.8	2.4	2.6	2.7	2.8	3.9	5.4	7.7	11.1	18.7	41.7
2004.....	221.8	1.8	2.8	3.0	2.3	3.8	5.5	7.3	10.8	19.0	41.3
2003.....	224.3	1.9	2.6	2.7	2.7	3.9	5.2	7.1	11.4	20.5	44.0
2002.....	226.7	1.9	2.8	3.0	2.7	3.9	5.7	7.6	12.0	21.1	44.9
2001.....	227.2	1.6	2.9	2.4	2.7	4.0	5.8	7.7	11.9	21.8	45.5
2000.....	228.7	2.6	2.9	2.5	2.6	4.3	6.2	7.3	11.8	22.3	46.2
1999.....	229.9	1.6	3.0	2.7	2.6	4.5	6.1	8.1	12.3	22.7	47.4

Non-Hispanic 4/, female

2007.....	195.9	1.3	2.3	2.2	2.2	2.6	3.6	5.9	12.1	24.1	48.3
2006.....	196.7	1.9	2.2	2.1	1.8	2.4	3.6	6.5	13.1	25.1	49.2
2005.....	197.5	*	2.1	2.3	2.1	2.8	3.8	6.1	12.8	25.3	51.4
2004.....	197.5	1.6	2.4	2.4	2.0	3.0	3.9	6.6	13.2	25.8	51.4
2003.....	200.0	2.1	2.1	2.3	2.4	2.6	4.2	7.2	13.2	27.7	52.1
2002.....	200.8	1.5	2.4	2.4	2.1	3.1	4.1	6.6	14.2	28.2	53.8
2001.....	201.2	1.6	2.5	2.5	2.2	2.8	4.4	7.6	14.6	29.0	55.0
2000.....	203.1	2.4	2.6	2.2	2.2	3.0	4.2	7.2	14.3	28.3	54.4
1999.....	201.9	1.8	2.5	2.3	2.2	3.1	4.5	7.1	13.7	29.7	54.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms (C00-C97)

Non-Hispanic White, both sexes

2007 5/.....	226.9	1.8	2.5	2.5	2.3	3.0	4.3	6.3	11.0	20.4	41.5
2006 5/.....	227.1	2.1	2.3	2.4	2.1	3.1	4.3	6.7	11.9	21.1	42.5
2005 5/.....	227.7	2.0	2.3	2.5	2.5	3.4	4.6	6.5	11.5	21.2	45.0
2004 5/.....	227.0	1.9	2.7	2.8	2.1	3.4	4.4	6.8	11.6	21.7	44.0
2003 5/.....	229.5	2.2	2.4	2.7	2.5	3.4	4.7	7.0	11.9	23.2	45.8
2002.....	230.9	1.6	2.6	2.8	2.4	3.5	5.1	6.9	12.7	23.9	47.2
2001.....	230.6	1.8	2.9	2.5	2.3	3.4	5.1	7.4	12.8	24.5	47.7
2000.....	232.2	2.6	2.8	2.4	2.4	3.5	5.1	7.3	12.7	24.0	46.9
1999.....	231.5	1.6	3.0	2.4	2.3	3.7	5.1	7.6	12.6	24.9	48.1

Non-Hispanic White, male

2007 5/.....	240.6	2.0	2.4	2.5	2.3	3.4	5.1	7.0	10.5	17.6	37.0
2006 5/.....	239.9	1.9	2.5	2.7	2.3	3.8	5.2	7.4	11.3	18.4	38.3
2005 5/.....	240.7	2.7	2.5	2.6	2.8	4.0	5.4	7.4	10.6	18.4	41.0
2004 5/.....	239.2	2.0	2.8	3.1	2.3	3.8	5.1	7.5	11.0	19.0	39.2
2003 5/.....	241.8	2.1	2.6	2.9	2.5	4.0	5.2	7.3	11.2	20.6	42.0
2002.....	243.8	2.0	2.8	2.9	2.6	3.9	5.9	7.4	11.8	20.5	43.2
2001.....	243.4	1.7	3.1	2.3	2.4	4.0	5.8	7.7	12.0	21.2	43.2
2000.....	244.4	2.9	3.1	2.5	2.6	4.1	6.1	7.4	11.8	21.5	42.9
1999.....	245.1	1.7	3.3	2.7	2.5	4.2	5.8	8.1	12.0	22.1	44.8

Non-Hispanic White, female

2007 5/.....	213.7	*	2.5	2.4	2.2	2.5	3.5	5.5	11.5	23.3	46.0
2006 5/.....	214.7	2.2	2.1	2.2	1.8	2.4	3.4	5.9	12.6	23.8	46.6
2005 5/.....	215.1	*	2.2	2.4	2.1	2.8	3.7	5.5	12.3	24.2	49.0
2004 5/.....	215.3	1.8	2.5	2.4	1.8	2.9	3.7	6.1	12.3	24.4	48.7
2003 5/.....	217.6	2.3	2.2	2.4	2.4	2.7	4.1	6.7	12.6	26.0	49.6
2002.....	218.5	*	2.3	2.5	2.2	3.0	4.3	6.5	13.5	27.2	51.2
2001.....	218.4	1.8	2.8	2.6	2.2	2.8	4.4	7.2	13.7	27.7	52.2
2000.....	220.6	2.4	2.4	2.4	2.2	2.9	4.1	7.1	13.5	26.5	50.9
1999.....	218.4	*	2.7	2.1	2.1	3.0	4.3	7.1	13.1	27.7	51.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms (C00-C97)

Non-Hispanic Black, both sexes

2007 5/.....	166.3	*	2.0	1.9	2.5	3.1	5.0	8.0	15.1	26.5	59.4
2006 5/.....	165.5	*	2.5	2.3	2.2	3.1	5.1	8.4	15.8	29.3	61.5
2005 5/.....	167.4	*	2.5	2.7	2.6	3.1	5.2	9.4	16.2	29.0	62.3
2004 5/.....	167.5	*	2.5	2.5	2.7	3.9	6.1	9.2	15.6	29.4	66.0
2003 5/.....	170.0	*	2.3	2.4	3.0	3.0	5.2	8.9	16.2	32.1	68.2
2002.....	171.5	*	3.0	2.6	2.6	3.6	4.6	8.4	17.0	32.7	70.0
2001.....	172.8	*	2.0	2.4	2.9	3.7	5.3	9.6	17.0	33.7	72.2
2000.....	174.9	*	2.8	2.2	2.4	4.5	6.1	8.3	16.3	35.4	76.1
1999.....	176.9	*	2.0	2.5	3.1	4.6	6.9	8.8	16.4	36.9	74.9

Non-Hispanic Black, male

2007 5/.....	180.1	*	2.4	1.9	2.6	3.2	5.7	7.7	12.6	19.6	50.6
2006 5/.....	179.2	*	2.7	2.6	2.5	3.9	5.5	8.0	13.5	23.0	52.5
2005 5/.....	181.8	*	2.8	3.1	3.1	3.4	6.3	9.4	15.4	22.5	51.6
2004 5/.....	182.4	*	2.8	2.5	2.6	4.3	7.1	8.1	12.2	20.6	58.0
2003 5/.....	184.8	*	2.4	2.6	3.3	3.6	5.8	7.3	13.9	22.5	62.7
2002.....	187.8	*	2.7	3.3	3.0	4.2	5.2	8.6	14.7	26.5	61.9
2001.....	190.8	*	2.4	2.6	3.4	4.3	6.1	8.9	12.9	27.6	64.3
2000.....	194.9	*	2.6	2.5	2.7	5.5	7.4	7.6	13.3	28.3	71.8
1999.....	197.4	*	2.0	2.4	3.4	5.6	7.7	9.2	14.5	28.6	69.9

Non-Hispanic Black, female

2007 5/.....	153.7	*	1.6	1.9	2.3	3.0	4.4	8.3	17.3	32.6	67.0
2006 5/.....	153.0	*	2.4	2.1	1.8	2.3	4.7	8.9	17.8	34.9	69.5
2005 5/.....	154.3	*	2.1	2.3	2.1	2.8	4.2	9.5	16.8	34.8	71.7
2004 5/.....	153.9	*	2.2	2.5	2.8	3.5	5.2	10.3	18.6	37.2	73.2
2003 5/.....	156.5	*	2.1	2.2	2.6	2.4	4.5	10.4	18.4	40.7	73.0
2002.....	156.7	*	3.2	1.9	2.2	3.1	4.0	8.3	19.0	38.2	77.1
2001.....	156.5	*	*	2.2	2.3	3.0	4.5	10.3	20.8	39.2	79.2
2000.....	156.9	*	3.0	1.8	2.0	3.6	4.8	8.9	18.9	41.8	79.8
1999.....	158.3	*	2.0	2.7	2.7	3.5	6.1	8.4	18.1	44.3	79.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/, both sexes

2007.....	2.7	*	*	*	*	*	*	*	0.2	0.3	0.8
2006.....	2.6	*	*	*	*	*	*	*	0.1	0.4	0.9
2005.....	2.6	*	*	*	*	*	*	0.1	0.2	0.3	0.8
2004.....	2.7	*	*	*	*	*	*	*	0.1	0.3	0.9
2003.....	2.7	*	*	*	*	*	*	*	0.2	0.3	0.9
2002.....	2.7	*	*	*	*	*	*	0.1	0.2	0.4	0.8
2001.....	2.7	*	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	2.7	*	*	*	*	*	*	*	0.2	0.4	0.9
1999.....	2.7	*	*	*	*	*	*	*	0.1	0.4	0.9

All origins 3/, male

2007.....	3.7	*	*	*	*	*	*	*	*	0.5	1.1
2006.....	3.6	*	*	*	*	*	*	*	*	0.5	1.2
2005.....	3.6	*	*	*	*	*	*	*	*	0.5	1.2
2004.....	3.7	*	*	*	*	*	*	*	0.2	0.3	1.2
2003.....	3.7	*	*	*	*	*	*	*	0.3	0.4	1.4
2002.....	3.7	*	*	*	*	*	*	*	0.2	0.4	1.2
2001.....	3.7	*	*	*	*	*	*	*	0.2	0.5	1.2
2000.....	3.6	*	*	*	*	*	*	*	*	0.5	1.4
1999.....	3.6	*	*	*	*	*	*	*	*	0.5	1.3

All origins 3/, female

2007.....	1.7	*	*	*	*	*	*	*	*	0.2	0.6
2006.....	1.6	*	*	*	*	*	*	*	*	0.2	0.5
2005.....	1.7	*	*	*	*	*	*	*	*	0.2	0.4
2004.....	1.7	*	*	*	*	*	*	*	*	0.2	0.5
2003.....	1.7	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	1.7	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.8	*	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.8	*	*	*	*	*	*	*	*	0.2	0.4
1999.....	1.8	*	*	*	*	*	*	*	*	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 4/, both sexes

2007.....	3.0	*	*	*	*	*	*	*	0.2	0.4	0.9
2006.....	2.9	*	*	*	*	*	*	*	0.2	0.4	0.9
2005.....	2.9	*	*	*	*	*	*	*	0.2	0.4	0.9
2004.....	3.0	*	*	*	*	*	*	*	0.2	0.3	0.9
2003.....	3.0	*	*	*	*	*	*	*	0.2	0.3	1.0
2002.....	3.0	*	*	*	*	*	*	0.1	0.2	0.4	0.8
2001.....	3.0	*	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	2.9	*	*	*	*	*	*	*	0.2	0.4	1.0
1999.....	2.9	*	*	*	*	*	*	*	0.2	0.4	0.9

Non-Hispanic 4/, male

2007.....	4.2	*	*	*	*	*	*	*	*	0.5	1.2
2006.....	4.0	*	*	*	*	*	*	*	*	0.5	1.4
2005.....	4.1	*	*	*	*	*	*	*	*	0.6	1.3
2004.....	4.1	*	*	*	*	*	*	*	0.2	0.4	1.4
2003.....	4.1	*	*	*	*	*	*	*	0.3	0.4	1.5
2002.....	4.1	*	*	*	*	*	*	*	0.3	0.5	1.2
2001.....	4.1	*	*	*	*	*	*	*	0.3	0.6	1.3
2000.....	3.9	*	*	*	*	*	*	*	*	0.6	1.5
1999.....	4.0	*	*	*	*	*	*	*	*	0.5	1.4

Non-Hispanic 4/, female

2007.....	1.9	*	*	*	*	*	*	*	*	0.2	0.6
2006.....	1.8	*	*	*	*	*	*	*	*	0.3	0.5
2005.....	1.9	*	*	*	*	*	*	*	*	*	0.5
2004.....	1.9	*	*	*	*	*	*	*	*	0.2	0.5
2003.....	1.9	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	1.9	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.9	*	*	*	*	*	*	*	*	0.3	0.5
2000.....	2.0	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	2.0	*	*	*	*	*	*	*	*	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic White, both sexes

2007 5/.....	3.2	*	*	*	*	*	*	*	*	0.3	0.8
2006 5/.....	3.0	*	*	*	*	*	*	*	*	0.3	0.8
2005 5/.....	3.0	*	*	*	*	*	*	*	*	0.3	0.9
2004 5/.....	3.1	*	*	*	*	*	*	*	*	0.2	0.7
2003 5/.....	3.1	*	*	*	*	*	*	*	0.2	0.3	0.9
2002.....	3.1	*	*	*	*	*	*	*	0.2	0.3	0.7
2001.....	3.1	*	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	3.0	*	*	*	*	*	*	*	*	0.3	0.8
1999.....	3.0	*	*	*	*	*	*	*	*	0.3	0.8

Non-Hispanic White, male

2007 5/.....	4.3	*	*	*	*	*	*	*	*	0.4	1.0
2006 5/.....	4.1	*	*	*	*	*	*	*	*	0.4	1.2
2005 5/.....	4.1	*	*	*	*	*	*	*	*	0.4	1.2
2004 5/.....	4.1	*	*	*	*	*	*	*	*	*	1.1
2003 5/.....	4.1	*	*	*	*	*	*	*	*	0.4	1.4
2002.....	4.1	*	*	*	*	*	*	*	*	0.4	1.0
2001.....	4.1	*	*	*	*	*	*	*	*	0.4	1.1
2000.....	3.9	*	*	*	*	*	*	*	*	0.4	1.1
1999.....	3.9	*	*	*	*	*	*	*	*	0.4	1.2

Non-Hispanic White, female

2007 5/.....	2.0	*	*	*	*	*	*	*	*	*	0.5
2006 5/.....	2.0	*	*	*	*	*	*	*	*	*	0.5
2005 5/.....	2.0	*	*	*	*	*	*	*	*	*	0.5
2004 5/.....	2.0	*	*	*	*	*	*	*	*	*	0.4
2003 5/.....	2.0	*	*	*	*	*	*	*	*	*	0.3
2002.....	2.1	*	*	*	*	*	*	*	*	*	0.4
2001.....	2.1	*	*	*	*	*	*	*	*	*	0.4
2000.....	2.1	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	2.1	*	*	*	*	*	*	*	*	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of stomach (C16)

All origins 3/, both sexes

2007.....	3.8	*	*	*	*	*	*	0.2	0.4	0.8	1.3
2006.....	3.8	*	*	*	*	*	*	0.2	0.4	0.8	1.3
2005.....	3.9	*	*	*	*	*	*	0.1	0.4	0.7	1.3
2004.....	4.0	*	*	*	*	*	*	0.1	0.5	0.7	1.4
2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8	1.3
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7	1.2
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8	1.3
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7	1.4
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8	1.4

All origins 3/, male

2007.....	4.5	*	*	*	*	*	*	0.3	0.3	0.8	1.5
2006.....	4.5	*	*	*	*	*	*	*	0.4	0.7	1.4
2005.....	4.6	*	*	*	*	*	*	*	0.4	0.8	1.6
2004.....	4.8	*	*	*	*	*	*	*	0.4	0.7	1.7
2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8	1.5
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7	1.4
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9	1.4
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9	1.7
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9	1.7

All origins 3/, female

2007.....	3.0	*	*	*	*	*	*	*	0.5	0.9	1.1
2006.....	3.1	*	*	*	*	*	*	0.2	0.4	0.9	1.2
2005.....	3.2	*	*	*	*	*	*	*	0.4	0.5	1.1
2004.....	3.3	*	*	*	*	*	*	*	0.5	0.6	1.2
2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7	1.0
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7	1.1
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6	1.2
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of stomach (C16)

Non-Hispanic 4/, both sexes

2007.....	3.9	*	*	*	*	*	*	0.2	0.3	0.7	1.2
2006.....	3.9	*	*	*	*	*	*	0.1	0.4	0.7	1.2
2005.....	4.0	*	*	*	*	*	*	*	0.4	0.6	1.2
2004.....	4.2	*	*	*	*	*	*	0.1	0.4	0.5	1.3
2003.....	4.3	*	*	*	*	*	*	0.1	0.4	0.7	1.3
2002.....	4.4	*	*	*	*	*	*	0.2	0.4	0.6	1.1
2001.....	4.5	*	*	*	*	*	*	*	0.4	0.7	1.1
2000.....	4.7	*	*	*	*	*	*	0.2	0.4	0.6	1.3
1999.....	4.7	*	*	*	*	*	*	0.2	0.4	0.8	1.3

Non-Hispanic 4/, male

2007.....	4.8	*	*	*	*	*	*	*	*	0.7	1.4
2006.....	4.8	*	*	*	*	*	*	*	0.4	0.7	1.3
2005.....	4.8	*	*	*	*	*	*	*	0.3	0.8	1.4
2004.....	5.0	*	*	*	*	*	*	*	0.3	0.6	1.5
2003.....	5.2	*	*	*	*	*	*	*	0.4	0.8	1.5
2002.....	5.3	*	*	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	5.4	*	*	*	*	*	*	*	0.5	0.8	1.2
2000.....	5.5	*	*	*	*	*	*	*	0.4	0.8	1.6
1999.....	5.7	*	*	*	*	*	*	*	0.4	0.9	1.6

Non-Hispanic 4/, female

2007.....	3.1	*	*	*	*	*	*	*	0.4	0.7	0.9
2006.....	3.1	*	*	*	*	*	*	*	0.4	0.7	1.1
2005.....	3.3	*	*	*	*	*	*	*	0.4	0.5	1.0
2004.....	3.3	*	*	*	*	*	*	*	0.4	0.5	1.0
2003.....	3.5	*	*	*	*	*	*	*	0.4	0.7	1.0
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.5	0.9
2001.....	3.6	*	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	3.8	*	*	*	*	*	*	*	0.4	0.4	1.0
1999.....	3.8	*	*	*	*	*	*	*	0.3	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of stomach (C16)

Non-Hispanic White, both sexes

2007 5/.....	3.6	*	*	*	*	*	*	*	0.3	0.6	1.0
2006 5/.....	3.6	*	*	*	*	*	*	*	0.3	0.5	0.8
2005 5/.....	3.7	*	*	*	*	*	*	*	0.2	0.4	0.9
2004 5/.....	3.9	*	*	*	*	*	*	*	0.3	0.5	0.9
2003 5/.....	4.0	*	*	*	*	*	*	*	0.3	0.6	1.0
2002.....	4.1	*	*	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	4.2	*	*	*	*	*	*	*	0.3	0.5	0.8
2000.....	4.4	*	*	*	*	*	*	*	0.3	0.4	0.9
1999.....	4.4	*	*	*	*	*	*	*	0.3	0.6	0.9

Non-Hispanic White, male

2007 5/.....	4.4	*	*	*	*	*	*	*	*	0.6	1.2
2006 5/.....	4.5	*	*	*	*	*	*	*	*	0.5	0.9
2005 5/.....	4.4	*	*	*	*	*	*	*	*	0.5	1.1
2004 5/.....	4.7	*	*	*	*	*	*	*	0.3	0.5	1.1
2003 5/.....	4.8	*	*	*	*	*	*	*	0.3	0.7	1.1
2002.....	5.0	*	*	*	*	*	*	*	*	0.6	1.0
2001.....	5.0	*	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	5.2	*	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	5.4	*	*	*	*	*	*	*	0.3	0.7	1.0

Non-Hispanic White, female

2007 5/.....	2.8	*	*	*	*	*	*	*	0.3	0.5	0.7
2006 5/.....	2.8	*	*	*	*	*	*	*	*	0.4	0.8
2005 5/.....	3.0	*	*	*	*	*	*	*	*	0.4	0.7
2004 5/.....	3.1	*	*	*	*	*	*	*	*	0.5	0.6
2003 5/.....	3.3	*	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.2	*	*	*	*	*	*	*	*	0.4	0.6
2001.....	3.4	*	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.5	*	*	*	*	*	*	*	0.3	0.3	0.7
1999.....	3.5	*	*	*	*	*	*	*	*	0.5	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes

2007.....	17.8	*	*	*	*	*	0.1	0.5	0.9	1.9	4.1
2006.....	17.9	*	*	*	*	*	0.2	0.5	1.1	2.0	4.0
2005.....	18.0	*	*	*	*	*	0.2	0.4	1.1	1.9	4.0
2004.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0	3.9
2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1	3.8
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0	4.3
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3	4.1
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1	4.0
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9	4.0

All origins 3/, male

2007.....	18.2	*	*	*	*	*	*	0.5	1.0	2.1	4.4
2006.....	18.3	*	*	*	*	*	*	0.5	1.1	2.1	4.2
2005.....	18.4	*	*	*	*	*	0.2	0.5	1.1	2.1	4.2
2004.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1	4.2
2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2	4.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1	4.9
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6	4.3
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4	4.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0	4.1

All origins 3/, female

2007.....	17.3	*	*	*	*	*	*	0.4	0.9	1.7	3.8
2006.....	17.5	*	*	*	*	*	0.2	0.5	1.1	1.9	3.8
2005.....	17.6	*	*	*	*	*	*	0.4	1.0	1.7	3.7
2004.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0	3.6
2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0	3.4
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9	3.6
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9	3.8
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8	3.5
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 4/, both sexes

2007.....	19.8	*	*	*	*	*	0.1	0.5	1.0	2.0	4.3
2006.....	19.9	*	*	*	*	*	0.1	0.5	1.2	2.1	4.2
2005.....	20.0	*	*	*	*	*	0.2	0.5	1.2	2.0	4.2
2004.....	20.3	*	*	*	*	*	0.2	0.5	1.0	2.1	4.1
2003.....	21.3	*	*	*	*	*	0.1	0.5	1.0	2.1	4.0
2002.....	21.7	*	*	*	*	*	0.2	0.5	1.1	2.1	4.4
2001.....	22.0	*	*	*	*	*	0.2	0.5	1.2	2.4	4.2
2000.....	22.4	*	*	*	*	*	0.1	0.4	1.0	2.2	4.0
1999.....	22.4	*	*	*	*	*	0.2	0.5	1.0	2.0	4.2

Non-Hispanic 4/, male

2007.....	20.4	*	*	*	*	*	*	0.6	1.0	2.3	4.5
2006.....	20.4	*	*	*	*	*	*	0.5	1.2	2.3	4.6
2005.....	20.5	*	*	*	*	*	*	0.6	1.3	2.2	4.4
2004.....	20.7	*	*	*	*	*	0.3	0.5	1.0	2.1	4.4
2003.....	21.7	*	*	*	*	*	*	0.6	1.1	2.3	4.4
2002.....	22.3	*	*	*	*	*	0.3	0.5	1.2	2.2	5.2
2001.....	22.3	*	*	*	*	*	*	0.4	1.3	2.7	4.4
2000.....	22.8	*	*	*	*	*	*	0.5	1.1	2.5	4.6
1999.....	22.7	*	*	*	*	*	*	0.7	1.1	2.1	4.3

Non-Hispanic 4/, female

2007.....	19.2	*	*	*	*	*	*	0.4	0.9	1.8	4.0
2006.....	19.4	*	*	*	*	*	*	0.5	1.2	1.9	3.9
2005.....	19.4	*	*	*	*	*	*	0.4	1.0	1.7	3.9
2004.....	19.9	*	*	*	*	*	*	0.4	0.9	2.1	3.9
2003.....	20.9	*	*	*	*	*	*	0.4	1.0	2.0	3.6
2002.....	21.2	*	*	*	*	*	*	0.4	0.9	2.0	3.7
2001.....	21.7	*	*	*	*	*	*	0.5	1.0	2.0	3.9
2000.....	22.1	*	*	*	*	*	*	0.4	0.8	1.8	3.5
1999.....	22.2	*	*	*	*	*	*	0.3	0.9	1.9	4.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic White, both sexes

2007 5/.....	21.1	*	*	*	*	*	*	0.5	1.0	2.0	4.0
2006 5/.....	21.2	*	*	*	*	*	*	0.5	1.1	2.0	3.8
2005 5/.....	21.2	*	*	*	*	*	0.2	0.4	1.1	1.9	3.9
2004 5/.....	21.7	*	*	*	*	*	*	0.4	0.9	2.0	3.8
2003 5/.....	22.7	*	*	*	*	*	*	0.5	1.0	1.9	3.5
2002.....	23.2	*	*	*	*	*	0.2	0.5	1.1	2.0	4.0
2001.....	23.5	*	*	*	*	*	0.2	0.5	1.1	2.2	3.8
2000.....	24.0	*	*	*	*	*	*	0.4	0.9	1.9	3.6
1999.....	24.0	*	*	*	*	*	0.2	0.4	0.9	1.9	3.9

Non-Hispanic White, male

2007 5/.....	21.7	*	*	*	*	*	*	0.5	1.0	2.2	4.2
2006 5/.....	21.6	*	*	*	*	*	*	0.4	1.2	2.1	4.2
2005 5/.....	21.8	*	*	*	*	*	*	0.5	1.2	2.2	4.3
2004 5/.....	22.2	*	*	*	*	*	*	0.5	1.0	2.1	4.0
2003 5/.....	23.2	*	*	*	*	*	*	0.6	1.0	2.1	3.9
2002.....	23.8	*	*	*	*	*	0.3	0.6	1.2	2.0	4.7
2001.....	23.9	*	*	*	*	*	*	0.4	1.3	2.6	4.0
2000.....	24.2	*	*	*	*	*	*	0.4	1.1	2.3	4.1
1999.....	24.3	*	*	*	*	*	*	0.6	1.0	2.1	4.1

Non-Hispanic White, female

2007 5/.....	20.5	*	*	*	*	*	*	0.4	0.9	1.8	3.8
2006 5/.....	20.7	*	*	*	*	*	*	0.5	1.0	1.9	3.5
2005 5/.....	20.6	*	*	*	*	*	*	0.4	1.1	1.6	3.5
2004 5/.....	21.3	*	*	*	*	*	*	0.4	0.8	1.8	3.6
2003 5/.....	22.2	*	*	*	*	*	*	0.4	1.0	1.8	3.1
2002.....	22.7	*	*	*	*	*	*	0.4	1.0	1.9	3.3
2001.....	23.1	*	*	*	*	*	*	0.5	0.9	1.9	3.7
2000.....	23.8	*	*	*	*	*	*	0.4	0.6	1.5	3.2
1999.....	23.7	*	*	*	*	*	*	*	0.9	1.7	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic Black, both sexes

2007 5/.....	18.0	*	*	*	*	*	*	0.8	1.2	2.5	6.1
2006 5/.....	18.1	*	*	*	*	*	*	0.7	2.0	3.2	7.1
2005 5/.....	18.2	*	*	*	*	*	*	*	1.7	2.8	5.9
2004 5/.....	17.8	*	*	*	*	*	*	*	1.4	2.9	6.1
2003 5/.....	18.8	*	*	*	*	*	*	*	1.2	3.5	6.8
2002.....	18.9	*	*	*	*	*	*	*	1.0	2.7	7.4
2001.....	19.2	*	*	*	*	*	*	*	1.6	3.4	6.6
2000.....	19.4	*	*	*	*	*	*	*	1.6	3.8	6.7
1999.....	19.3	*	*	*	*	*	*	0.8	1.3	3.2	6.7

Non-Hispanic Black, male

2007 5/.....	18.8	*	*	*	*	*	*	*	*	3.0	6.9
2006 5/.....	18.9	*	*	*	*	*	*	*	1.8	3.7	7.6
2005 5/.....	18.5	*	*	*	*	*	*	*	2.3	2.9	6.0
2004 5/.....	18.0	*	*	*	*	*	*	*	*	2.5	6.8
2003 5/.....	19.0	*	*	*	*	*	*	*	*	3.4	7.4
2002.....	19.3	*	*	*	*	*	*	*	*	3.3	8.6
2001.....	19.0	*	*	*	*	*	*	*	1.9	4.0	7.6
2000.....	19.8	*	*	*	*	*	*	*	1.8	4.2	8.2
1999.....	19.2	*	*	*	*	*	*	*	*	3.1	6.5

Non-Hispanic Black, female

2007 5/.....	17.2	*	*	*	*	*	*	*	*	2.1	5.3
2006 5/.....	17.3	*	*	*	*	*	*	*	2.1	2.7	6.7
2005 5/.....	17.9	*	*	*	*	*	*	*	*	2.7	5.8
2004 5/.....	17.6	*	*	*	*	*	*	*	1.4	3.3	5.5
2003 5/.....	18.7	*	*	*	*	*	*	*	*	3.6	6.3
2002.....	18.4	*	*	*	*	*	*	*	*	2.2	6.4
2001.....	19.4	*	*	*	*	*	*	*	*	3.0	5.7
2000.....	19.1	*	*	*	*	*	*	*	1.4	3.5	5.3
1999.....	19.5	*	*	*	*	*	*	*	*	3.2	6.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes

2007.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.5	1.2
2006.....	5.5	*	*	*	*	*	*	0.1	0.3	0.5	1.1
2005.....	5.4	*	0.1	*	*	*	0.1	0.2	0.3	0.5	1.2
2004.....	5.2	*	0.1	*	*	*	0.1	0.1	0.3	0.5	1.2
2003.....	5.1	*	0.1	*	*	*	0.1	0.1	0.3	0.5	1.2
2002.....	4.9	*	0.2	*	*	*	*	0.2	0.3	0.6	1.3
2001.....	4.7	*	*	*	*	*	*	0.2	0.2	0.6	1.2
2000.....	4.6	*	0.1	*	*	*	*	0.1	0.3	0.5	1.4
1999.....	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6	1.4

All origins 3/, male

2007.....	7.6	*	*	*	*	*	*	0.2	0.4	0.7	1.7
2006.....	7.3	*	*	*	*	*	*	0.2	0.3	0.6	1.6
2005.....	7.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6
2004.....	6.8	*	*	*	*	*	*	0.2	0.3	0.8	1.8
2003.....	6.6	*	*	*	*	*	*	*	0.4	0.8	1.8
2002.....	6.4	*	*	*	*	*	*	0.2	0.4	0.8	2.0
2001.....	6.1	*	*	*	*	*	*	*	0.3	0.9	1.8
2000.....	5.9	*	*	*	*	*	*	*	0.4	0.7	2.0
1999.....	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8	2.1

All origins 3/, female

2007.....	3.8	*	*	*	*	*	*	*	*	0.4	0.7
2006.....	3.8	*	*	*	*	*	*	*	0.3	0.3	0.7
2005.....	3.7	*	*	*	*	*	*	*	*	0.3	0.8
2004.....	3.7	*	*	*	*	*	*	*	0.2	0.2	0.6
2003.....	3.5	*	*	*	*	*	*	*	0.2	0.3	0.7
2002.....	3.4	*	*	*	*	*	*	*	0.3	0.3	0.6
2001.....	3.4	*	*	*	*	*	*	*	*	0.3	0.5
2000.....	3.3	*	*	*	*	*	*	*	*	0.3	0.8
1999.....	3.3	*	*	*	*	*	*	*	*	0.3	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic White, both sexes

2007 5/.....	5.8	*	*	*	*	*	*	*	0.2	0.4	0.9
2006 5/.....	5.6	*	*	*	*	*	*	*	0.2	0.3	0.9
2005 5/.....	5.5	*	*	*	*	*	*	*	0.2	0.3	1.0
2004 5/.....	5.3	*	*	*	*	*	*	*	0.2	0.4	0.9
2003 5/.....	5.1	*	*	*	*	*	*	*	0.2	0.4	1.0
2002.....	5.0	*	*	*	*	*	*	*	0.2	0.4	1.0
2001.....	4.8	*	*	*	*	*	*	*	0.1	0.4	0.8
2000.....	4.6	*	*	*	*	*	*	*	0.2	0.3	0.9
1999.....	4.5	*	0.2	*	*	*	*	*	*	0.4	1.1

Non-Hispanic White, male

2007 5/.....	7.7	*	*	*	*	*	*	*	*	0.6	1.1
2006 5/.....	7.3	*	*	*	*	*	*	*	*	0.4	1.1
2005 5/.....	7.2	*	*	*	*	*	*	*	*	0.3	1.3
2004 5/.....	6.8	*	*	*	*	*	*	*	*	0.5	1.2
2003 5/.....	6.7	*	*	*	*	*	*	*	*	0.5	1.3
2002.....	6.4	*	*	*	*	*	*	*	*	0.6	1.6
2001.....	6.1	*	*	*	*	*	*	*	*	0.6	1.1
2000.....	5.9	*	*	*	*	*	*	*	*	0.3	1.3
1999.....	5.7	*	*	*	*	*	*	*	*	0.5	1.7

Non-Hispanic White, female

2007 5/.....	4.0	*	*	*	*	*	*	*	*	0.3	0.6
2006 5/.....	4.0	*	*	*	*	*	*	*	*	*	0.7
2005 5/.....	3.8	*	*	*	*	*	*	*	*	*	0.6
2004 5/.....	3.8	*	*	*	*	*	*	*	*	*	0.5
2003 5/.....	3.6	*	*	*	*	*	*	*	*	0.3	0.6
2002.....	3.6	*	*	*	*	*	*	*	*	0.3	0.5
2001.....	3.4	*	*	*	*	*	*	*	*	*	0.4
2000.....	3.4	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	3.4	*	*	*	*	*	*	*	*	0.3	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of pancreas (C25)

All origins 3/, both sexes

2007.....	11.3	*	*	*	*	*	*	*	0.2	0.6	1.8
2006.....	11.2	*	*	*	*	*	*	*	0.2	0.6	1.7
2005.....	11.1	*	*	*	*	*	*	0.1	0.2	0.6	1.9
2004.....	10.8	*	*	*	*	*	*	*	0.2	0.6	1.7
2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7	1.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8	1.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6	1.8
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8	1.9

All origins 3/, male

2007.....	11.5	*	*	*	*	*	*	*	0.3	0.8	2.2
2006.....	11.2	*	*	*	*	*	*	*	*	0.8	2.0
2005.....	11.1	*	*	*	*	*	*	*	0.2	0.8	2.2
2004.....	10.9	*	*	*	*	*	*	*	0.3	0.7	2.2
2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9	2.0
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8	2.4
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9	2.0
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8	2.1
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9	2.4

All origins 3/, female

2007.....	11.1	*	*	*	*	*	*	*	*	0.5	1.5
2006.....	11.1	*	*	*	*	*	*	*	*	0.3	1.4
2005.....	11.0	*	*	*	*	*	*	*	*	0.5	1.6
2004.....	10.7	*	*	*	*	*	*	*	*	0.5	1.3
2003.....	10.6	*	*	*	*	*	*	*	*	0.5	1.4
2002.....	10.5	*	*	*	*	*	*	*	*	0.6	1.4
2001.....	10.6	*	*	*	*	*	*	*	*	0.6	1.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5	1.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/, both sexes

2007.....	52.6	*	*	*	*	*	*	0.2	0.5	1.7	6.8
2006.....	53.0	*	*	*	*	*	*	0.2	0.6	2.0	7.1
2005.....	53.7	*	*	*	*	*	*	0.2	0.5	2.1	8.2
2004.....	53.8	*	*	*	*	*	*	0.2	0.5	2.2	8.5
2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4	8.5
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7	9.2
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9	9.5
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9	9.3
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9	9.3

All origins 3/, male

2007.....	59.4	*	*	*	*	*	*	0.2	0.5	1.7	6.7
2006.....	60.5	*	*	*	*	*	*	0.2	0.6	2.0	7.3
2005.....	61.8	*	*	*	*	*	*	*	0.6	2.2	8.5
2004.....	62.0	*	*	*	*	*	*	0.3	0.6	2.2	9.1
2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4	9.5
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6	9.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8	10.5
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3	10.5
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8	10.5

All origins 3/, female

2007.....	46.0	*	*	*	*	*	*	*	0.4	1.7	7.0
2006.....	45.7	*	*	*	*	*	*	*	0.5	2.0	6.9
2005.....	45.9	*	*	*	*	*	*	0.2	0.4	2.0	8.0
2004.....	45.9	*	*	*	*	*	*	*	0.5	2.2	7.9
2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5	7.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7	8.8
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9	8.4
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6	8.1
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0	8.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 4/, both sexes

2007.....	60.1	*	*	*	*	*	*	0.2	0.5	1.9	7.6
2006.....	60.4	*	*	*	*	*	*	0.2	0.7	2.3	7.8
2005.....	60.9	*	*	*	*	*	*	0.2	0.5	2.3	9.1
2004.....	60.9	*	*	*	*	*	*	0.2	0.5	2.5	9.4
2003.....	61.2	*	*	*	*	*	*	0.1	0.6	2.7	9.3
2002.....	61.4	*	*	*	*	*	*	0.1	0.7	3.0	10.0
2001.....	61.3	*	*	*	*	*	*	0.2	0.6	3.2	10.2
2000.....	61.5	*	*	*	*	*	*	0.2	0.8	3.2	10.0
1999.....	60.5	*	*	*	*	*	*	0.2	0.7	3.2	9.9

Non-Hispanic 4/, male

2007.....	68.3	*	*	*	*	*	*	0.3	0.6	1.9	7.5
2006.....	69.3	*	*	*	*	*	*	0.3	0.7	2.3	8.1
2005.....	70.3	*	*	*	*	*	*	*	0.7	2.4	9.4
2004.....	70.4	*	*	*	*	*	*	0.3	0.6	2.5	10.1
2003.....	71.1	*	*	*	*	*	*	*	0.7	2.7	10.4
2002.....	71.8	*	*	*	*	*	*	*	0.7	3.0	10.5
2001.....	72.6	*	*	*	*	*	*	*	0.6	3.1	11.3
2000.....	73.2	*	*	*	*	*	*	0.3	0.8	3.6	11.3
1999.....	72.7	*	*	*	*	*	*	0.3	0.8	3.1	11.2

Non-Hispanic 4/, female

2007.....	52.3	*	*	*	*	*	*	*	0.4	1.8	7.7
2006.....	51.9	*	*	*	*	*	*	*	0.6	2.2	7.6
2005.....	51.9	*	*	*	*	*	*	*	0.4	2.2	8.8
2004.....	51.8	*	*	*	*	*	*	*	0.5	2.5	8.7
2003.....	51.8	*	*	*	*	*	*	*	0.6	2.7	8.2
2002.....	51.6	*	*	*	*	*	*	*	0.6	2.9	9.6
2001.....	50.5	*	*	*	*	*	*	*	0.7	3.3	9.0
2000.....	50.4	*	*	*	*	*	*	*	0.9	2.8	8.7
1999.....	48.8	*	*	*	*	*	*	*	0.6	3.2	8.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic White, both sexes

2007 5/.....	66.7	*	*	*	*	*	*	0.2	0.5	1.9	7.7
2006 5/.....	66.9	*	*	*	*	*	*	0.2	0.6	2.3	7.8
2005 5/.....	67.3	*	*	*	*	*	*	0.2	0.5	2.4	9.0
2004 5/.....	67.1	*	*	*	*	*	*	0.2	0.4	2.5	9.1
2003 5/.....	67.5	*	*	*	*	*	*	*	0.7	2.7	9.1
2002.....	67.7	*	*	*	*	*	*	*	0.6	3.0	9.7
2001.....	67.3	*	*	*	*	*	*	0.2	0.7	3.2	9.8
2000.....	67.4	*	*	*	*	*	*	0.2	0.9	3.0	9.4
1999.....	65.9	*	*	*	*	*	*	0.2	0.7	3.0	9.5

Non-Hispanic White, male

2007 5/.....	74.6	*	*	*	*	*	*	*	0.6	2.0	7.5
2006 5/.....	75.8	*	*	*	*	*	*	*	0.7	2.3	8.0
2005 5/.....	76.8	*	*	*	*	*	*	*	0.7	2.4	9.1
2004 5/.....	76.5	*	*	*	*	*	*	*	0.5	2.4	9.6
2003 5/.....	77.3	*	*	*	*	*	*	*	0.7	2.6	9.9
2002.....	77.8	*	*	*	*	*	*	*	0.7	2.9	9.8
2001.....	78.4	*	*	*	*	*	*	*	0.6	3.0	10.5
2000.....	78.9	*	*	*	*	*	*	*	0.8	3.2	10.0
1999.....	78.1	*	*	*	*	*	*	*	0.8	2.9	10.4

Non-Hispanic White, female

2007 5/.....	59.0	*	*	*	*	*	*	*	0.5	1.8	8.0
2006 5/.....	58.4	*	*	*	*	*	*	*	0.5	2.2	7.6
2005 5/.....	58.2	*	*	*	*	*	*	*	0.4	2.4	8.9
2004 5/.....	58.1	*	*	*	*	*	*	*	0.4	2.7	8.7
2003 5/.....	58.1	*	*	*	*	*	*	*	0.6	2.7	8.3
2002.....	57.9	*	*	*	*	*	*	*	0.5	3.1	9.5
2001.....	56.6	*	*	*	*	*	*	*	0.7	3.4	9.1
2000.....	56.4	*	*	*	*	*	*	*	0.9	2.7	8.7
1999.....	54.3	*	*	*	*	*	*	*	0.6	3.1	8.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic Black, both sexes

2007 5/.....	42.9	*	*	*	*	*	*	*	*	1.9	8.6
2006 5/.....	43.3	*	*	*	*	*	*	*	0.9	2.7	9.7
2005 5/.....	44.0	*	*	*	*	*	*	*	*	2.5	11.3
2004 5/.....	44.2	*	*	*	*	*	*	*	*	2.5	13.1
2003 5/.....	44.5	*	*	*	*	*	*	*	*	3.0	12.8
2002.....	44.7	*	*	*	*	*	*	*	*	3.5	14.5
2001.....	44.6	*	*	*	*	*	*	*	*	3.6	14.9
2000.....	45.4	*	*	*	*	*	*	*	0.9	5.0	16.1
1999.....	45.6	*	*	*	*	*	*	*	*	4.7	14.9

Non-Hispanic Black, male

2007 5/.....	53.6	*	*	*	*	*	*	*	*	1.7	9.7
2006 5/.....	54.1	*	*	*	*	*	*	*	*	3.0	10.6
2005 5/.....	55.0	*	*	*	*	*	*	*	*	3.0	12.4
2004 5/.....	56.3	*	*	*	*	*	*	*	*	2.8	15.0
2003 5/.....	57.0	*	*	*	*	*	*	*	*	3.0	16.1
2002.....	58.1	*	*	*	*	*	*	*	*	4.4	17.5
2001.....	59.2	*	*	*	*	*	*	*	*	4.0	18.8
2000.....	60.3	*	*	*	*	*	*	*	*	6.0	21.9
1999.....	60.9	*	*	*	*	*	*	*	*	4.7	19.1

Non-Hispanic Black, female

2007 5/.....	33.2	*	*	*	*	*	*	*	*	2.1	7.7
2006 5/.....	33.5	*	*	*	*	*	*	*	*	2.5	9.0
2005 5/.....	34.0	*	*	*	*	*	*	*	*	2.0	10.3
2004 5/.....	33.2	*	*	*	*	*	*	*	*	2.3	11.5
2003 5/.....	33.1	*	*	*	*	*	*	*	*	2.9	9.9
2002.....	32.6	*	*	*	*	*	*	*	*	2.7	11.9
2001.....	31.3	*	*	*	*	*	*	*	*	3.3	11.4
2000.....	31.9	*	*	*	*	*	*	*	*	4.1	10.9
1999.....	31.8	*	*	*	*	*	*	*	*	4.7	11.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant melanoma of skin (C43)

All origins 3/, both sexes

2007.....	2.8	*	*	*	*	*	0.1	0.4	0.5	0.8	1.4
2006.....	2.8	*	*	*	*	*	0.2	0.3	0.7	0.9	1.4
2005.....	2.8	*	*	*	*	*	0.2	0.5	0.6	0.9	1.5
2004.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9	1.4
2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9	1.6
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0	1.7
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0	1.8
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0	1.7
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0	1.7

All origins 3/, male

2007.....	3.7	*	*	*	*	*	*	0.4	0.6	0.8	1.6
2006.....	3.7	*	*	*	*	*	0.2	0.3	0.7	1.2	1.6
2005.....	3.6	*	*	*	*	*	*	0.6	0.8	0.9	1.9
2004.....	3.5	*	*	*	*	*	*	0.4	0.7	1.0	1.8
2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1	1.9
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1	2.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2	2.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0	2.1
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3	2.2

All origins 3/, female

2007.....	1.9	*	*	*	*	*	*	0.4	0.4	0.7	1.2
2006.....	2.0	*	*	*	*	*	*	*	0.6	0.6	1.2
2005.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8	1.1
2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.8	1.1
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7	1.3
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9	1.2
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7	1.4
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0	1.4
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant melanoma of skin (C43)

Non-Hispanic 4/, both sexes

2007.....	3.2	*	*	*	*	*	0.1	0.4	0.6	0.8	1.6
2006.....	3.2	*	*	*	*	*	0.2	0.3	0.8	1.0	1.6
2005.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.0	1.7
2004.....	3.1	*	*	*	*	*	0.2	0.4	0.7	1.0	1.6
2003.....	3.0	*	*	*	*	*	0.2	0.4	0.7	1.0	1.8
2002.....	2.9	*	*	*	*	*	0.2	0.4	0.7	1.1	1.8
2001.....	3.0	*	*	*	*	*	0.2	0.4	0.6	1.1	1.9
2000.....	3.0	*	*	*	*	*	0.2	0.4	0.7	1.1	1.9
1999.....	2.9	*	*	*	*	*	0.2	0.4	0.7	1.2	1.9

Non-Hispanic 4/, male

2007.....	4.3	*	*	*	*	*	*	0.4	0.7	0.9	1.8
2006.....	4.3	*	*	*	*	*	0.2	0.4	0.9	1.3	1.9
2005.....	4.2	*	*	*	*	*	*	0.7	0.9	1.0	2.2
2004.....	4.1	*	*	*	*	*	*	0.5	0.9	1.2	2.0
2003.....	4.0	*	*	*	*	*	*	0.5	0.9	1.2	2.1
2002.....	3.8	*	*	*	*	*	*	0.5	0.9	1.2	2.3
2001.....	3.9	*	*	*	*	*	*	0.5	0.7	1.3	2.4
2000.....	3.7	*	*	*	*	*	*	0.4	0.9	1.1	2.3
1999.....	3.7	*	*	*	*	*	*	0.5	0.8	1.5	2.4

Non-Hispanic 4/, female

2007.....	2.2	*	*	*	*	*	*	0.4	0.5	0.7	1.3
2006.....	2.2	*	*	*	*	*	*	*	0.7	0.7	1.3
2005.....	2.3	*	*	*	*	*	*	0.3	0.5	1.0	1.2
2004.....	2.1	*	*	*	*	*	*	0.3	0.6	0.9	1.2
2003.....	2.2	*	*	*	*	*	*	0.3	0.6	0.8	1.4
2002.....	2.1	*	*	*	*	*	*	0.4	0.5	1.0	1.3
2001.....	2.1	*	*	*	*	*	*	0.4	0.6	0.8	1.5
2000.....	2.2	*	*	*	*	*	*	0.3	0.6	1.1	1.5
1999.....	2.1	*	*	*	*	*	*	0.3	0.7	0.8	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant melanoma of skin (C43)

Non-Hispanic White, both sexes

2007 5/.....	4.0	*	*	*	*	*	0.2	0.5	0.8	1.0	1.9
2006 5/.....	4.0	*	*	*	*	*	0.2	0.4	1.0	1.3	1.9
2005 5/.....	4.0	*	*	*	*	*	0.2	0.7	0.9	1.3	2.0
2004 5/.....	3.8	*	*	*	*	*	0.2	0.6	0.9	1.3	2.0
2003 5/.....	3.8	*	*	*	*	*	0.2	0.5	0.9	1.3	2.2
2002.....	3.6	*	*	*	*	*	0.2	0.6	0.9	1.4	2.2
2001.....	3.7	*	*	*	*	*	0.2	0.6	0.8	1.3	2.4
2000.....	3.6	*	*	*	*	*	0.3	0.4	0.9	1.3	2.2
1999.....	3.5	*	*	*	*	*	0.2	0.5	0.9	1.4	2.3

Non-Hispanic White, male

2007 5/.....	5.4	*	*	*	*	*	*	0.5	0.9	1.2	2.3
2006 5/.....	5.3	*	*	*	*	*	*	0.5	1.2	1.7	2.3
2005 5/.....	5.2	*	*	*	*	*	*	1.0	1.1	1.3	2.6
2004 5/.....	5.1	*	*	*	*	*	*	0.7	1.1	1.5	2.4
2003 5/.....	4.9	*	*	*	*	*	*	0.7	1.1	1.6	2.6
2002.....	4.7	*	*	*	*	*	*	0.6	1.1	1.5	2.8
2001.....	4.8	*	*	*	*	*	*	0.6	0.9	1.6	2.9
2000.....	4.6	*	*	*	*	*	*	0.5	1.1	1.3	2.7
1999.....	4.5	*	*	*	*	*	*	0.6	1.0	1.8	3.0

Non-Hispanic White, female

2007 5/.....	2.7	*	*	*	*	*	*	0.5	0.7	0.9	1.6
2006 5/.....	2.7	*	*	*	*	*	*	*	0.9	0.9	1.5
2005 5/.....	2.8	*	*	*	*	*	*	0.5	0.7	1.2	1.5
2004 5/.....	2.6	*	*	*	*	*	*	0.4	0.8	1.1	1.6
2003 5/.....	2.6	*	*	*	*	*	*	0.4	0.7	1.0	1.7
2002.....	2.6	*	*	*	*	*	*	0.5	0.6	1.2	1.7
2001.....	2.6	*	*	*	*	*	*	0.5	0.7	1.1	1.8
2000.....	2.7	*	*	*	*	*	*	0.4	0.7	1.4	1.8
1999.....	2.5	*	*	*	*	*	*	0.4	0.8	1.1	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of breast (C50)

All origins 3/, both sexes

2007.....	13.6	*	*	*	*	*	*	0.3	1.4	3.5	6.6
2006.....	13.8	*	*	*	*	*	*	0.4	1.5	3.8	6.9
2005.....	14.0	*	*	*	*	*	*	0.4	1.4	3.8	7.4
2004.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9	7.3
2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3	7.9
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1	7.9
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3	8.2
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3	8.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3	7.9

All origins 3/, male

2007.....	0.2	*	*	*	*	*	*	*	*	*	*
2006.....	0.3	*	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*

All origins 3/, female

2007.....	26.5	*	*	*	*	*	*	0.6	2.9	7.0	13.0
2006.....	26.9	*	*	*	*	*	*	0.7	3.0	7.5	13.8
2005.....	27.3	*	*	*	*	*	*	0.8	2.8	7.7	14.7
2004.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8	14.5
2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5	15.6
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2	15.6
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6	16.2
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5	16.4
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5	15.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of breast (C50)

Hispanic, both sexes

2007.....	4.6	*	*	*	*	*	*	*	1.1	2.7	4.4
2006.....	4.7	*	*	*	*	*	*	*	1.3	2.6	5.1
2005.....	4.5	*	*	*	*	*	*	*	0.8	2.6	5.1
2004.....	4.7	*	*	*	*	*	*	*	1.0	2.8	4.7
2003.....	4.7	*	*	*	*	*	*	*	1.2	2.7	6.2
2002.....	4.5	*	*	*	*	*	*	*	1.3	2.4	5.4
2001.....	4.8	*	*	*	*	*	*	*	1.1	3.5	5.7
2000.....	4.8	*	*	*	*	*	*	*	1.2	3.2	5.6
1999.....	4.6	*	*	*	*	*	*	*	1.1	2.7	5.6

Hispanic, male

2007.....	*	*	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Hispanic, female

2007.....	9.4	*	*	*	*	*	*	*	2.3	5.8	9.2
2006.....	9.6	*	*	*	*	*	*	*	2.8	5.5	10.6
2005.....	9.4	*	*	*	*	*	*	*	1.7	5.5	10.7
2004.....	9.6	*	*	*	*	*	*	*	2.1	5.9	9.7
2003.....	9.8	*	*	*	*	*	*	*	2.7	5.8	12.9
2002.....	9.2	*	*	*	*	*	*	*	2.7	5.1	11.0
2001.....	9.7	*	*	*	*	*	*	*	2.5	7.3	11.5
2000.....	9.7	*	*	*	*	*	*	*	2.6	6.6	11.3
1999.....	9.3	*	*	*	*	*	*	*	2.4	5.7	11.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of breast (C50)

Non-Hispanic 4/, both sexes

2007.....	15.2	*	*	*	*	*	*	0.3	1.5	3.6	6.9
2006.....	15.3	*	*	*	*	*	*	0.4	1.5	4.0	7.2
2005.....	15.6	*	*	*	*	*	*	0.4	1.6	4.1	7.7
2004.....	15.6	*	*	*	*	*	*	0.4	1.7	4.1	7.7
2003.....	15.9	*	*	*	*	*	*	0.3	1.8	4.5	8.1
2002.....	16.0	*	*	*	*	*	*	0.4	1.8	4.4	8.2
2001.....	16.1	*	*	*	*	*	*	0.5	1.9	4.4	8.5
2000.....	16.5	*	*	*	*	*	*	0.4	2.0	4.5	8.6
1999.....	16.3	*	*	*	*	*	*	0.4	1.8	4.5	8.1

Non-Hispanic 4/, male

2007.....	0.3	*	*	*	*	*	*	*	*	*	*
2006.....	0.3	*	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2007.....	29.4	*	*	*	*	*	*	0.7	3.0	7.2	13.6
2006.....	29.7	*	*	*	*	*	*	0.7	3.0	7.9	14.2
2005.....	30.1	*	*	*	*	*	*	0.8	3.1	8.1	15.2
2004.....	30.2	*	*	*	*	*	*	0.8	3.4	8.1	15.1
2003.....	30.8	*	*	*	*	*	*	0.6	3.6	8.9	15.9
2002.....	31.0	*	*	*	*	*	*	0.7	3.6	8.7	16.1
2001.....	31.1	*	*	*	*	*	*	0.9	3.8	8.8	16.7
2000.....	31.8	*	*	*	*	*	*	0.8	4.0	8.8	16.9
1999.....	31.4	*	*	*	*	*	*	0.9	3.6	8.9	16.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of breast (C50)

Non-Hispanic White, both sexes

2007 5/.....	16.0	*	*	*	*	*	*	0.2	1.2	3.1	6.1
2006 5/.....	16.2	*	*	*	*	*	*	0.3	1.3	3.4	6.5
2005 5/.....	16.3	*	*	*	*	*	*	0.3	1.3	3.4	6.9
2004 5/.....	16.4	*	*	*	*	*	*	0.3	1.4	3.5	6.7
2003 5/.....	16.8	*	*	*	*	*	*	*	1.5	4.0	7.2
2002.....	16.9	*	*	*	*	*	*	0.3	1.5	3.8	7.2
2001.....	17.0	*	*	*	*	*	*	0.4	1.6	3.9	7.5
2000.....	17.4	*	*	*	*	*	*	0.4	1.7	4.0	7.7
1999.....	17.1	*	*	*	*	*	*	0.4	1.5	3.9	7.2

Non-Hispanic White, male

2007 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2005 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2004 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2003 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, female

2007 5/.....	31.1	*	*	*	*	*	*	0.5	2.4	6.3	12.1
2006 5/.....	31.4	*	*	*	*	*	*	0.5	2.5	6.8	12.8
2005 5/.....	31.8	*	*	*	*	*	*	0.7	2.6	6.9	13.7
2004 5/.....	32.0	*	*	*	*	*	*	0.6	2.9	7.0	13.3
2003 5/.....	32.6	*	*	*	*	*	*	*	3.1	8.0	14.3
2002.....	32.9	*	*	*	*	*	*	0.5	3.0	7.6	14.3
2001.....	32.9	*	*	*	*	*	*	0.7	3.2	7.8	15.0
2000.....	33.8	*	*	*	*	*	*	0.7	3.4	7.8	15.3
1999.....	33.3	*	*	*	*	*	*	0.7	3.1	7.8	14.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of breast (C50)

Non-Hispanic Black, both sexes

2007 5/.....	15.2	*	*	*	*	*	*	0.9	3.3	7.1	12.8
2006 5/.....	15.1	*	*	*	*	*	*	0.8	3.4	8.0	13.0
2005 5/.....	15.5	*	*	*	*	*	*	0.8	3.1	8.1	14.0
2004 5/.....	15.0	*	*	*	*	*	*	1.0	3.4	8.3	14.0
2003 5/.....	15.5	*	*	*	*	*	*	0.9	3.5	8.3	14.5
2002.....	15.4	*	*	*	*	*	*	0.9	3.8	8.4	15.6
2001.....	15.5	*	*	*	*	*	*	1.2	4.2	7.8	15.2
2000.....	15.3	*	*	*	*	*	*	0.8	3.8	8.0	15.2
1999.....	15.5	*	*	*	*	*	*	0.9	3.8	9.1	15.1

Non-Hispanic Black, male

2007 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2005 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2004 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2003 5/.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	28.8	*	*	*	*	*	*	1.7	6.2	13.5	24.1
2006 5/.....	28.5	*	*	*	*	*	*	1.6	6.4	15.0	24.3
2005 5/.....	29.4	*	*	*	*	*	*	1.6	5.9	15.4	26.2
2004 5/.....	28.4	*	*	*	*	*	*	1.9	6.4	15.7	26.3
2003 5/.....	29.4	*	*	*	*	*	*	1.7	6.7	15.6	26.9
2002.....	29.1	*	*	*	*	*	*	1.7	7.1	15.6	29.3
2001.....	29.2	*	*	*	*	*	*	2.3	7.8	14.7	28.3
2000.....	28.9	*	*	*	*	*	*	*	7.2	14.8	28.4
1999.....	29.2	*	*	*	*	*	*	1.8	7.1	16.8	28.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes

2007.....	1.3	*	*	*	*	*	*	0.2	0.7	1.3	1.7
2006.....	1.3	*	*	*	*	*	*	0.2	0.7	1.2	1.7
2005.....	1.3	*	*	*	*	*	*	0.2	0.6	1.1	1.7
2004.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1	1.7
2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2	1.7
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3	1.7
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3	1.8
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3	1.8
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	2.6	*	*	*	*	*	*	0.5	1.4	2.6	3.3
2006.....	2.6	*	*	*	*	*	*	0.5	1.4	2.4	3.4
2005.....	2.6	*	*	*	*	*	*	0.4	1.3	2.3	3.4
2004.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2	3.4
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3	3.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5	3.4
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6	3.7
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6	3.5
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

Hispanic, both sexes

2007.....	1.0	*	*	*	*	*	*	*	0.8	0.9	1.8
2006.....	1.0	*	*	*	*	*	*	*	0.6	1.3	1.5
2005.....	1.0	*	*	*	*	*	*	*	*	1.4	1.9
2004.....	1.0	*	*	*	*	*	*	*	*	1.2	2.0
2003.....	1.0	*	*	*	*	*	*	*	*	1.2	1.7
2002.....	1.0	*	*	*	*	*	*	*	0.6	1.3	1.7
2001.....	1.1	*	*	*	*	*	*	*	0.8	1.4	2.4
2000.....	1.2	*	*	*	*	*	*	*	0.6	1.9	2.3
1999.....	1.1	*	*	*	*	*	*	*	0.8	1.2	2.4

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	2.2	*	*	*	*	*	*	*	1.8	1.9	3.8
2006.....	2.2	*	*	*	*	*	*	*	1.2	2.7	3.1
2005.....	2.2	*	*	*	*	*	*	*	*	3.0	3.9
2004.....	2.2	*	*	*	*	*	*	*	*	2.5	4.2
2003.....	2.0	*	*	*	*	*	*	*	*	2.5	3.6
2002.....	2.1	*	*	*	*	*	*	*	1.3	2.8	3.4
2001.....	2.3	*	*	*	*	*	*	*	1.7	3.0	5.0
2000.....	2.5	*	*	*	*	*	*	*	1.4	4.0	4.8
1999.....	2.3	*	*	*	*	*	*	*	1.7	2.6	4.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 4/, both sexes

2007.....	1.4	*	*	*	*	*	*	0.3	0.6	1.4	1.7
2006.....	1.4	*	*	*	*	*	*	0.2	0.7	1.1	1.7
2005.....	1.4	*	*	*	*	*	*	0.2	0.7	1.1	1.7
2004.....	1.4	*	*	*	*	*	*	0.4	0.7	1.1	1.6
2003.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2	1.7
2002.....	1.4	*	*	*	*	*	*	0.3	0.8	1.2	1.7
2001.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8
2000.....	1.5	*	*	*	*	*	*	0.4	0.8	1.2	1.7
1999.....	1.6	*	*	*	*	*	*	0.3	0.7	1.3	1.8

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	2.7	*	*	*	*	*	*	0.5	1.3	2.8	3.3
2006.....	2.7	*	*	*	*	*	*	0.4	1.4	2.3	3.4
2005.....	2.7	*	*	*	*	*	*	0.4	1.4	2.1	3.3
2004.....	2.6	*	*	*	*	*	*	0.7	1.5	2.1	3.2
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3	3.3
2002.....	2.8	*	*	*	*	*	*	0.5	1.6	2.4	3.4
2001.....	2.9	*	*	*	*	*	*	0.5	1.5	2.5	3.5
2000.....	3.0	*	*	*	*	*	*	0.8	1.5	2.4	3.4
1999.....	3.0	*	*	*	*	*	*	0.7	1.5	2.5	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic White, both sexes

2007 5/.....	1.3	*	*	*	*	*	*	0.2	0.5	1.4	1.4
2006 5/.....	1.3	*	*	*	*	*	*	0.2	0.6	1.1	1.5
2005 5/.....	1.3	*	*	*	*	*	*	0.2	0.7	1.0	1.5
2004 5/.....	1.2	*	*	*	*	*	*	0.3	0.7	1.0	1.4
2003 5/.....	1.3	*	*	*	*	*	*	0.2	0.6	1.0	1.5
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.1	1.5
2001.....	1.4	*	*	*	*	*	*	0.2	0.7	1.2	1.6
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.1	1.4
1999.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2	1.6

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	2.5	*	*	*	*	*	*	0.4	1.1	2.7	2.9
2006 5/.....	2.5	*	*	*	*	*	*	0.4	1.2	2.3	3.0
2005 5/.....	2.5	*	*	*	*	*	*	0.4	1.5	2.0	3.0
2004 5/.....	2.4	*	*	*	*	*	*	0.6	1.4	2.0	2.9
2003 5/.....	2.5	*	*	*	*	*	*	0.5	1.3	2.0	3.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.6	2.3	3.0
2001.....	2.7	*	*	*	*	*	*	0.5	1.4	2.4	3.2
2000.....	2.7	*	*	*	*	*	*	0.8	1.4	2.1	2.9
1999.....	2.8	*	*	*	*	*	*	0.8	1.5	2.3	3.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic Black, both sexes

2007 5/.....	2.1	*	*	*	*	*	*	*	1.4	2.0	3.0
2006 5/.....	2.1	*	*	*	*	*	*	*	1.2	1.6	3.3
2005 5/.....	2.1	*	*	*	*	*	*	*	0.8	1.7	3.1
2004 5/.....	2.1	*	*	*	*	*	*	*	1.0	1.7	2.9
2003 5/.....	2.2	*	*	*	*	*	*	*	1.2	2.1	2.5
2002.....	2.3	*	*	*	*	*	*	*	1.1	2.1	3.0
2001.....	2.2	*	*	*	*	*	*	*	1.2	1.9	3.0
2000.....	2.5	*	*	*	*	*	*	*	1.5	2.1	3.2
1999.....	2.4	*	*	*	*	*	*	*	1.0	2.1	3.0

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	4.0	*	*	*	*	*	*	*	2.7	3.8	5.6
2006 5/.....	3.9	*	*	*	*	*	*	*	2.3	3.1	6.2
2005 5/.....	4.0	*	*	*	*	*	*	*	1.5	3.2	5.9
2004 5/.....	4.0	*	*	*	*	*	*	*	1.8	3.1	5.4
2003 5/.....	4.1	*	*	*	*	*	*	*	2.3	4.0	4.6
2002.....	4.3	*	*	*	*	*	*	*	2.1	4.0	5.7
2001.....	4.2	*	*	*	*	*	*	*	2.2	3.6	5.6
2000.....	4.7	*	*	*	*	*	*	*	2.8	4.0	6.0
1999.....	4.5	*	*	*	*	*	*	*	1.9	4.0	5.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins 3/, both sexes

2007.....	2.5	*	*	*	*	*	*	*	0.1	0.3	0.5
2006.....	2.5	*	*	*	*	*	*	*	*	0.2	0.4
2005.....	2.4	*	*	*	*	*	*	*	*	0.3	0.4
2004.....	2.4	*	*	*	*	*	*	*	*	0.2	0.5
2003.....	2.4	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	2.4	*	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	2.4	*	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	2.3	*	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	2.3	*	*	*	*	*	*	*	*	0.2	0.5

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	4.9	*	*	*	*	*	*	*	0.2	0.6	1.0
2006.....	4.9	*	*	*	*	*	*	*	*	0.4	0.9
2005.....	4.7	*	*	*	*	*	*	*	*	0.5	0.8
2004.....	4.7	*	*	*	*	*	*	*	*	0.5	1.0
2003.....	4.7	*	*	*	*	*	*	*	*	0.3	1.0
2002.....	4.7	*	*	*	*	*	*	*	0.2	0.4	0.9
2001.....	4.7	*	*	*	*	*	*	*	0.3	0.4	0.8
2000.....	4.6	*	*	*	*	*	*	*	0.2	0.4	0.9
1999.....	4.5	*	*	*	*	*	*	*	*	0.3	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 4/, both sexes

2007.....	2.8	*	*	*	*	*	*	*	*	0.3	0.5
2006.....	2.7	*	*	*	*	*	*	*	*	0.2	0.5
2005.....	2.6	*	*	*	*	*	*	*	*	0.3	0.4
2004.....	2.6	*	*	*	*	*	*	*	*	0.3	0.5
2003.....	2.6	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	2.6	*	*	*	*	*	*	*	*	0.2	0.4
2001.....	2.6	*	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	2.6	*	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	2.5	*	*	*	*	*	*	*	*	0.2	0.5

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	5.4	*	*	*	*	*	*	*	*	0.5	1.0
2006.....	5.4	*	*	*	*	*	*	*	*	0.4	0.9
2005.....	5.2	*	*	*	*	*	*	*	*	0.5	0.7
2004.....	5.2	*	*	*	*	*	*	*	*	0.5	1.0
2003.....	5.1	*	*	*	*	*	*	*	*	0.3	1.0
2002.....	5.1	*	*	*	*	*	*	*	*	0.4	0.8
2001.....	5.1	*	*	*	*	*	*	*	0.2	0.5	0.8
2000.....	5.0	*	*	*	*	*	*	*	0.3	0.4	0.8
1999.....	4.9	*	*	*	*	*	*	*	*	0.3	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic White, both sexes

2007 5/.....	2.8	*	*	*	*	*	*	*	*	0.2	0.4
2006 5/.....	2.8	*	*	*	*	*	*	*	*	0.2	0.5
2005 5/.....	2.7	*	*	*	*	*	*	*	*	0.2	0.3
2004 5/.....	2.7	*	*	*	*	*	*	*	*	0.3	0.5
2003 5/.....	2.7	*	*	*	*	*	*	*	*	0.2	0.4
2002.....	2.6	*	*	*	*	*	*	*	*	0.2	0.4
2001.....	2.7	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	2.6	*	*	*	*	*	*	*	*	0.2	0.4
1999.....	2.6	*	*	*	*	*	*	*	*	0.2	0.4

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	5.4	*	*	*	*	*	*	*	*	0.5	0.8
2006 5/.....	5.5	*	*	*	*	*	*	*	*	0.4	0.9
2005 5/.....	5.3	*	*	*	*	*	*	*	*	0.5	0.7
2004 5/.....	5.3	*	*	*	*	*	*	*	*	0.5	1.0
2003 5/.....	5.2	*	*	*	*	*	*	*	*	0.3	0.8
2002.....	5.1	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	5.2	*	*	*	*	*	*	*	*	0.4	0.8
2000.....	5.1	*	*	*	*	*	*	*	*	0.4	0.7
1999.....	5.1	*	*	*	*	*	*	*	*	0.3	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of ovary (C56)

All origins 3/, both sexes

2007.....	4.8	*	*	*	*	*	*	0.1	0.2	0.5	1.1
2006.....	5.0	*	*	*	*	*	*	0.1	0.3	0.5	1.3
2005.....	5.0	*	*	*	*	*	*	0.2	0.3	0.5	1.3
2004.....	5.0	*	*	*	*	*	0.1	0.2	0.3	0.6	1.4
2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6	1.4
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7	1.4
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7	1.4
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6	1.5
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7	1.5

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	9.6	*	*	*	*	*	*	0.3	0.5	1.0	2.3
2006.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1	2.5
2005.....	9.8	*	*	*	*	*	*	0.3	0.5	1.0	2.7
2004.....	9.9	*	*	*	*	*	0.2	0.3	0.6	1.2	2.8
2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3	2.8
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4	2.8
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4	2.8
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3	2.9
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of ovary (C56)

Non-Hispanic 4/, both sexes

2007.....	5.4	*	*	*	*	*	*	0.1	0.3	0.5	1.2
2006.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6	1.3
2005.....	5.5	*	*	*	*	*	*	0.1	0.3	0.5	1.4
2004.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6	1.5
2003.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6	1.4
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7	1.4
2001.....	5.6	*	*	*	*	*	*	0.2	0.4	0.7	1.4
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.7	1.5
1999.....	5.3	*	*	*	*	*	*	0.1	0.3	0.7	1.6

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	10.6	*	*	*	*	*	*	0.3	0.5	0.9	2.3
2006.....	10.8	*	*	*	*	*	*	0.3	0.6	1.1	2.6
2005.....	10.8	*	*	*	*	*	*	0.3	0.6	1.1	2.8
2004.....	10.8	*	*	*	*	*	*	0.4	0.7	1.2	2.9
2003.....	10.9	*	*	*	*	*	*	0.4	0.6	1.3	2.8
2002.....	11.0	*	*	*	*	*	0.3	0.4	0.6	1.5	2.7
2001.....	10.9	*	*	*	*	*	*	0.4	0.7	1.4	2.8
2000.....	10.7	*	*	*	*	*	*	0.4	0.7	1.3	3.0
1999.....	10.4	*	*	*	*	*	*	0.3	0.6	1.5	3.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of ovary (C56)

Non-Hispanic White, both sexes

2007 5/.....	6.1	*	*	*	*	*	*	*	0.3	0.5	1.3
2006 5/.....	6.2	*	*	*	*	*	*	0.2	0.3	0.6	1.4
2005 5/.....	6.2	*	*	*	*	*	*	*	0.3	0.6	1.4
2004 5/.....	6.2	*	*	*	*	*	*	*	0.3	0.6	1.5
2003 5/.....	6.3	*	*	*	*	*	*	0.2	0.3	0.6	1.5
2002.....	6.3	*	*	*	*	*	0.2	0.2	0.3	0.8	1.4
2001.....	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.5
2000.....	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.6
1999.....	6.0	*	*	*	*	*	*	*	0.3	0.8	1.7

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	12.0	*	*	*	*	*	*	*	0.5	1.0	2.6
2006 5/.....	12.3	*	*	*	*	*	*	0.3	0.7	1.2	2.7
2005 5/.....	12.2	*	*	*	*	*	*	*	0.6	1.1	2.8
2004 5/.....	12.2	*	*	*	*	*	*	*	0.7	1.2	3.0
2003 5/.....	12.3	*	*	*	*	*	*	0.4	0.7	1.2	3.0
2002.....	12.4	*	*	*	*	*	0.3	0.5	0.6	1.5	2.8
2001.....	12.2	*	*	*	*	*	*	0.4	0.7	1.4	2.9
2000.....	12.0	*	*	*	*	*	*	0.5	0.7	1.4	3.2
1999.....	11.7	*	*	*	*	*	*	*	0.5	1.5	3.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins 3/, both sexes

2007.....	4.2	*	*	*	*	*	0.1	*	0.2	0.3	0.8
2006.....	4.1	*	*	0.1	*	*	*	*	0.2	0.4	0.8
2005.....	4.2	*	*	0.1	*	*	*	*	0.2	0.4	0.9
2004.....	4.2	*	*	0.1	*	*	*	0.1	0.1	0.4	0.8
2003.....	4.2	*	*	0.1	*	*	*	*	0.2	0.4	1.0
2002.....	4.2	*	*	0.1	*	*	*	*	0.2	0.4	0.9
2001.....	4.2	*	0.1	0.1	*	*	*	*	0.2	0.3	1.2
2000.....	4.2	*	*	0.1	*	*	*	*	0.1	0.4	1.1
1999.....	4.0	*	*	0.1	*	*	*	0.1	0.1	0.5	1.1

All origins 3/, male

2007.....	5.4	*	*	*	*	*	*	*	0.3	0.4	1.1
2006.....	5.2	*	*	*	*	*	*	*	0.3	0.5	1.2
2005.....	5.3	*	*	*	*	*	*	*	0.2	0.4	1.2
2004.....	5.3	*	*	*	*	*	*	*	*	0.5	1.2
2003.....	5.4	*	*	*	*	*	*	*	0.3	0.5	1.5
2002.....	5.4	*	*	*	*	*	*	*	0.2	0.6	1.3
2001.....	5.4	*	*	*	*	*	*	*	*	0.5	1.6
2000.....	5.3	*	*	*	*	*	*	*	*	0.5	1.6
1999.....	5.0	*	*	*	*	*	*	*	0.2	0.6	1.5

All origins 3/, female

2007.....	3.1	*	*	*	*	*	*	*	*	0.2	0.5
2006.....	3.1	*	*	*	*	*	*	*	*	0.3	0.4
2005.....	3.1	*	*	*	*	*	*	*	*	0.4	0.7
2004.....	3.1	*	*	*	*	*	*	*	*	0.2	0.5
2003.....	3.1	*	*	*	*	*	*	*	*	0.2	0.6
2002.....	3.1	*	*	*	*	*	*	*	*	0.3	0.5
2001.....	3.1	*	*	0.2	*	*	*	*	*	*	0.9
2000.....	3.1	*	*	*	*	*	*	*	*	0.3	0.7
1999.....	3.0	*	*	*	*	*	*	*	*	0.3	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic White, both sexes

2007 5/.....	5.1	*	*	*	*	*	*	*	0.2	0.3	0.8
2006 5/.....	5.1	*	*	*	*	*	*	*	*	0.3	0.8
2005 5/.....	5.1	*	*	*	*	*	*	*	*	0.4	0.9
2004 5/.....	5.1	*	*	*	*	*	*	*	*	0.3	0.8
2003 5/.....	5.1	*	*	0.2	*	*	*	*	0.2	0.3	1.1
2002.....	5.1	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	5.1	*	*	*	*	*	*	*	*	0.3	1.2
2000.....	5.0	*	*	*	*	*	*	*	*	0.3	1.1
1999.....	4.7	*	*	*	*	*	*	*	*	0.4	1.1

Non-Hispanic White, male

2007 5/.....	6.5	*	*	*	*	*	*	*	*	0.4	1.1
2006 5/.....	6.4	*	*	*	*	*	*	*	*	0.4	1.3
2005 5/.....	6.5	*	*	*	*	*	*	*	*	0.4	1.1
2004 5/.....	6.5	*	*	*	*	*	*	*	*	0.4	1.1
2003 5/.....	6.6	*	*	*	*	*	*	*	*	0.5	1.5
2002.....	6.5	*	*	*	*	*	*	*	*	0.5	1.3
2001.....	6.5	*	*	*	*	*	*	*	*	0.4	1.5
2000.....	6.3	*	*	*	*	*	*	*	*	0.5	1.6
1999.....	6.0	*	*	*	*	*	*	*	*	0.5	1.5

Non-Hispanic White, female

2007 5/.....	3.8	*	*	*	*	*	*	*	*	*	0.6
2006 5/.....	3.8	*	*	*	*	*	*	*	*	0.3	0.4
2005 5/.....	3.8	*	*	*	*	*	*	*	*	0.3	0.7
2004 5/.....	3.8	*	*	*	*	*	*	*	*	*	0.5
2003 5/.....	3.8	*	*	*	*	*	*	*	*	*	0.6
2002.....	3.8	*	*	*	*	*	*	*	*	*	0.5
2001.....	3.8	*	*	*	*	*	*	*	*	*	0.8
2000.....	3.8	*	*	*	*	*	*	*	*	*	0.7
1999.....	3.5	*	*	*	*	*	*	*	*	0.3	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes

2007.....	4.4	*	0.7	0.9	0.6	0.4	0.5	0.7	1.2	1.5	2.4
2006.....	4.3	*	0.6	0.9	0.6	0.5	0.5	0.8	1.1	1.6	2.3
2005.....	4.4	*	0.5	1.0	0.7	0.6	0.6	0.8	1.1	1.6	2.7
2004.....	4.4	*	0.6	0.9	0.7	0.5	0.5	0.7	1.1	1.7	2.4
2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9	2.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7	2.8
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8	2.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0	2.9
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9	2.8

All origins 3/, male

2007.....	4.9	*	0.7	0.8	0.6	0.4	0.6	0.9	1.4	2.0	2.9
2006.....	4.9	*	0.7	0.9	0.7	0.5	0.5	0.9	1.2	1.9	2.9
2005.....	5.0	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	2.0	3.5
2004.....	4.9	*	0.7	1.0	0.7	0.6	0.7	0.8	1.3	2.1	2.9
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3	3.4
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0	3.5
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2	3.5
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6	3.7
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3	3.4

All origins 3/, female

2007.....	3.9	*	0.6	0.9	0.7	0.4	0.4	0.5	1.0	1.1	1.8
2006.....	3.7	*	0.6	0.8	0.6	0.4	0.5	0.7	1.0	1.2	1.8
2005.....	3.9	*	0.6	1.0	0.6	0.4	0.5	0.5	0.8	1.2	1.9
2004.....	3.9	*	0.6	0.9	0.7	0.5	0.4	0.6	0.8	1.2	1.9
2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5	2.4
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4	2.1
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4	2.1
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4	2.0
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 4/, both sexes

2007.....	4.8	*	0.7	0.9	0.7	0.5	0.5	0.8	1.3	1.6	2.5
2006.....	4.8	*	0.6	0.9	0.6	0.5	0.5	0.8	1.3	1.8	2.5
2005.....	4.9	*	0.5	1.0	0.7	0.6	0.6	0.9	1.2	1.7	2.8
2004.....	4.8	*	0.7	1.0	0.7	0.5	0.6	0.8	1.2	1.8	2.5
2003.....	4.9	*	0.7	0.9	0.8	0.6	0.6	0.9	1.3	2.1	3.1
2002.....	4.9	*	0.7	1.1	0.6	0.5	0.6	0.8	1.4	1.9	2.9
2001.....	4.8	*	0.6	0.9	0.7	0.5	0.7	0.8	1.5	1.9	2.9
2000.....	4.9	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.1	3.0
1999.....	5.0	*	0.7	0.9	0.6	0.7	0.6	1.0	1.5	2.0	2.9

Non-Hispanic 4/, male

2007.....	5.5	*	0.7	0.8	0.6	0.4	0.5	1.0	1.5	2.1	3.2
2006.....	5.4	*	0.6	0.9	0.7	0.6	0.6	0.9	1.5	2.1	3.1
2005.....	5.5	*	0.5	1.0	0.8	0.7	0.6	1.2	1.5	2.2	3.7
2004.....	5.4	*	0.7	1.1	0.7	0.5	0.7	0.9	1.5	2.3	3.1
2003.....	5.5	*	0.7	0.8	0.8	0.7	0.7	1.0	1.5	2.5	3.7
2002.....	5.5	*	0.8	1.2	0.7	0.6	0.8	0.8	1.7	2.3	3.7
2001.....	5.5	*	0.6	0.8	0.7	0.6	0.7	1.0	1.9	2.3	3.7
2000.....	5.5	*	0.7	0.7	0.8	0.6	0.8	1.0	1.8	2.8	3.9
1999.....	5.6	*	0.7	0.9	0.6	0.8	0.7	1.1	1.9	2.5	3.5

Non-Hispanic 4/, female

2007.....	4.2	*	0.7	0.9	0.7	0.5	0.4	0.5	1.0	1.2	1.9
2006.....	4.1	*	0.6	0.9	0.6	0.4	0.5	0.7	1.1	1.4	1.9
2005.....	4.2	*	0.5	1.0	0.5	0.4	0.5	0.5	1.0	1.3	2.0
2004.....	4.2	*	0.6	0.8	0.6	0.4	0.4	0.7	0.9	1.3	2.0
2003.....	4.3	*	0.6	0.9	0.8	0.5	0.6	0.9	1.0	1.7	2.5
2002.....	4.2	*	0.7	0.9	0.5	0.4	0.3	0.7	1.1	1.5	2.2
2001.....	4.2	*	0.6	1.0	0.6	0.4	0.6	0.5	1.1	1.4	2.1
2000.....	4.2	*	0.7	0.8	0.6	0.4	0.6	0.7	0.8	1.5	2.0
1999.....	4.4	*	0.7	0.9	0.7	0.5	0.4	0.8	1.0	1.6	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic White, both sexes

2007 5/.....	5.7	*	0.8	1.0	0.7	0.5	0.5	0.9	1.5	1.9	2.8
2006 5/.....	5.5	*	0.6	0.9	0.7	0.5	0.6	0.9	1.5	2.0	2.8
2005 5/.....	5.6	*	0.5	1.0	0.7	0.6	0.6	1.0	1.5	1.9	3.2
2004 5/.....	5.6	*	0.7	1.0	0.6	0.5	0.6	0.9	1.3	2.1	2.8
2003 5/.....	5.6	*	0.7	0.9	0.8	0.6	0.7	1.1	1.4	2.4	3.5
2002.....	5.6	*	0.6	1.1	0.6	0.6	0.6	0.8	1.6	2.2	3.3
2001.....	5.6	*	0.7	1.0	0.6	0.5	0.7	0.9	1.7	2.2	3.2
2000.....	5.6	*	0.7	0.8	0.7	0.5	0.7	1.0	1.4	2.4	3.2
1999.....	5.7	*	0.7	0.9	0.6	0.6	0.6	1.2	1.7	2.3	3.2

Non-Hispanic White, male

2007 5/.....	6.4	*	0.8	1.0	0.6	0.5	0.6	1.2	1.6	2.5	3.5
2006 5/.....	6.3	*	0.7	1.0	0.8	0.6	0.6	1.1	1.7	2.4	3.5
2005 5/.....	6.4	*	0.4	0.9	0.9	0.7	0.7	1.3	1.8	2.4	4.2
2004 5/.....	6.3	*	0.8	1.1	0.6	0.6	0.7	1.0	1.7	2.6	3.3
2003 5/.....	6.3	*	0.8	0.8	0.8	0.7	0.8	1.1	1.8	2.8	4.1
2002.....	6.4	*	0.7	1.1	0.7	0.7	0.9	0.9	1.9	2.6	4.2
2001.....	6.3	*	0.7	0.8	0.6	0.6	0.8	1.2	2.2	2.8	4.1
2000.....	6.3	*	0.7	0.7	0.8	0.6	0.8	1.2	1.9	3.1	4.2
1999.....	6.4	*	0.8	1.0	0.6	0.7	0.7	1.4	2.1	2.8	3.7

Non-Hispanic White, female

2007 5/.....	4.9	*	0.8	1.1	0.7	0.5	0.4	0.5	1.3	1.4	2.2
2006 5/.....	4.8	*	0.5	0.9	0.5	0.4	0.5	0.6	1.3	1.6	2.1
2005 5/.....	4.9	*	0.6	1.0	0.6	0.4	0.6	0.6	1.1	1.4	2.2
2004 5/.....	4.9	*	0.7	0.8	0.6	0.4	0.5	0.8	0.9	1.6	2.3
2003 5/.....	4.9	*	0.6	0.9	0.8	0.6	0.6	1.1	1.1	2.0	2.9
2002.....	4.9	*	0.6	1.0	0.6	0.5	0.3	0.8	1.3	1.7	2.4
2001.....	4.8	*	0.6	1.1	0.6	0.5	0.6	0.6	1.3	1.6	2.4
2000.....	4.9	*	0.7	0.8	0.7	0.4	0.7	0.8	1.0	1.6	2.2
1999.....	5.0	*	0.7	0.8	0.7	0.6	0.5	1.0	1.2	1.8	2.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes

2007.....	18.2	0.6	0.7	0.8	1.0	1.2	1.7	1.7	2.1	2.6	4.2
2006.....	18.4	0.8	0.8	0.6	0.9	1.3	1.9	1.9	2.2	2.8	4.2
2005.....	18.6	0.6	0.8	0.8	0.9	1.3	1.8	1.8	2.2	2.7	4.1
2004.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7	4.1
2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1	4.7
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2	4.6
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5	4.8
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4	4.8
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7	4.8

All origins 3/, male

2007.....	20.3	*	0.7	0.9	1.1	1.4	2.0	2.1	2.4	3.0	5.2
2006.....	20.3	*	0.9	0.7	1.0	1.6	2.3	2.3	2.6	3.1	5.1
2005.....	20.6	*	1.0	0.9	1.0	1.6	2.2	2.1	2.4	3.0	4.9
2004.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2	5.0
2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5	5.8
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7	5.5
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2	5.9
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9	5.7
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4	5.9

All origins 3/, female

2007.....	16.2	*	0.7	0.6	0.9	1.1	1.4	1.4	1.7	2.1	3.2
2006.....	16.5	*	0.8	0.5	0.7	0.9	1.3	1.5	1.8	2.4	3.3
2005.....	16.6	*	0.6	0.6	0.7	1.0	1.3	1.5	2.0	2.3	3.3
2004.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2	3.2
2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6	3.5
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6	3.7
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7	3.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0	3.9
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0	3.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Hispanic, both sexes

2007	7.2	*	0.5	1.1	1.5	1.7	2.4	2.0	2.1	2.3	4.4
2006	7.0	*	0.7	0.6	1.0	1.8	2.2	1.9	2.0	2.3	4.1
2005	7.3	*	0.8	1.0	1.2	1.8	2.5	1.8	2.1	2.3	3.4
2004	6.9	*	0.8	1.2	1.6	2.0	2.0	1.7	2.2	2.5	4.0
2003	7.1	*	1.3	1.1	1.5	1.5	2.2	1.6	2.4	2.6	4.3
2002	7.3	*	0.9	1.2	1.3	1.9	2.3	1.7	2.2	3.3	4.5
2001	7.4	*	1.3	1.2	1.2	2.4	1.9	2.1	2.4	3.0	4.9
2000	7.5	*	1.0	0.8	1.4	1.7	2.2	1.9	1.9	3.0	5.0
1999	7.5	*	1.0	0.8	1.6	1.5	2.6	1.9	2.4	3.2	4.6

Hispanic, male

2007	7.5	*	*	1.3	1.4	1.9	2.5	2.0	2.0	2.4	5.1
2006	7.4	*	*	*	*	2.1	2.9	2.2	2.0	2.3	4.8
2005	7.8	*	1.1	1.1	1.4	2.4	3.0	2.0	2.4	2.7	4.0
2004	7.6	*	*	1.7	1.9	2.6	2.7	1.8	2.7	2.6	5.8
2003	7.4	*	1.5	1.6	1.3	2.3	2.4	1.9	2.9	3.3	5.3
2002	7.6	*	*	1.1	1.4	2.6	2.7	2.1	2.6	3.7	5.1
2001	8.3	*	1.4	1.5	1.2	3.1	2.4	2.3	3.1	3.8	6.5
2000	8.0	*	*	*	2.0	2.0	2.6	2.4	2.3	2.9	6.1
1999	7.6	*	*	*	1.5	1.4	2.8	1.9	2.5	4.0	4.5

Hispanic, female

2007	7.0	*	*	1.0	1.6	1.4	2.3	2.0	2.1	2.1	3.8
2006	6.6	*	*	*	*	1.4	1.4	1.6	2.0	2.3	3.3
2005	6.7	*	*	*	*	1.2	1.8	1.5	1.7	2.0	2.8
2004	6.2	*	*	*	1.4	1.3	1.2	1.5	1.7	2.4	2.0
2003	6.8	*	*	*	1.7	*	2.0	1.2	1.9	1.9	3.3
2002	7.0	*	*	1.4	1.2	*	1.9	*	1.7	2.9	3.9
2001	6.5	*	*	*	1.3	1.7	1.3	1.8	1.6	2.2	3.2
2000	6.9	*	*	*	*	1.4	1.6	1.3	1.4	3.1	3.8
1999	7.4	*	*	*	1.6	1.5	2.2	1.9	2.2	2.4	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 4/, both sexes

2007.....	20.2	0.7	0.7	0.7	0.8	1.2	1.6	1.7	2.1	2.6	4.1
2006.....	20.3	0.8	0.8	0.6	0.9	1.2	1.8	1.9	2.2	2.8	4.3
2005.....	20.4	*	0.8	0.7	0.8	1.2	1.6	1.8	2.2	2.8	4.2
2004.....	20.5	*	0.8	0.8	0.7	1.3	1.6	1.8	2.3	2.7	4.1
2003.....	21.0	0.8	0.8	0.8	0.9	1.2	1.6	2.0	2.4	3.1	4.7
2002.....	21.3	0.8	0.7	0.7	0.9	1.4	1.8	2.0	2.7	3.1	4.6
2001.....	21.6	0.6	0.9	0.7	0.9	1.4	2.0	2.3	2.7	3.5	4.8
2000.....	21.8	1.0	0.9	0.8	0.9	1.6	2.1	2.2	2.6	3.5	4.8
1999.....	21.9	0.7	0.9	0.7	0.9	1.5	2.2	2.2	2.6	3.7	4.9

Non-Hispanic 4/, male

2007.....	22.7	*	0.7	0.9	1.0	1.3	1.9	2.1	2.5	3.1	5.2
2006.....	22.7	*	0.9	0.7	1.0	1.5	2.2	2.3	2.7	3.3	5.2
2005.....	22.8	*	1.0	0.9	1.0	1.5	2.0	2.2	2.4	3.1	5.0
2004.....	22.6	*	0.9	1.0	0.7	1.4	2.0	2.2	2.7	3.3	4.9
2003.....	23.3	*	0.8	0.9	1.1	1.4	1.9	2.3	3.0	3.6	5.9
2002.....	23.6	*	0.8	0.8	1.2	1.6	2.1	2.4	3.1	3.7	5.5
2001.....	23.5	*	1.0	0.8	1.1	1.6	2.3	2.6	3.2	4.2	5.8
2000.....	23.7	*	1.0	1.0	1.0	1.9	2.6	2.3	3.2	4.0	5.7
1999.....	23.7	*	0.9	0.7	1.1	1.8	2.7	2.5	3.1	4.4	6.0

Non-Hispanic 4/, female

2007.....	17.7	*	0.7	0.5	0.7	1.0	1.2	1.3	1.6	2.2	3.1
2006.....	18.1	*	0.8	0.5	0.7	0.8	1.3	1.5	1.7	2.4	3.3
2005.....	18.2	*	0.6	0.5	0.7	1.0	1.3	1.5	2.1	2.4	3.3
2004.....	18.5	*	0.6	0.7	0.6	1.1	1.3	1.5	1.9	2.2	3.3
2003.....	18.9	*	0.7	0.8	0.8	0.9	1.3	1.7	1.7	2.7	3.6
2002.....	19.2	*	0.7	0.7	0.7	1.2	1.6	1.6	2.3	2.6	3.6
2001.....	19.7	*	0.9	0.6	0.7	1.1	1.7	2.0	2.3	2.8	3.7
2000.....	20.1	*	0.9	0.6	0.8	1.2	1.5	2.1	2.0	2.9	3.9
1999.....	20.1	*	0.8	0.6	0.8	1.2	1.7	1.9	2.1	3.0	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic White, both sexes

2007 5/.....	22.4	*	0.7	0.7	0.9	1.1	1.5	1.6	2.0	2.6	3.7
2006 5/.....	22.6	*	0.9	0.6	0.8	1.2	1.7	2.0	2.1	2.7	4.1
2005 5/.....	22.8	*	0.7	0.7	0.8	1.2	1.6	1.6	2.0	2.7	3.9
2004 5/.....	22.7	*	0.8	0.9	0.6	1.3	1.6	1.7	2.2	2.6	4.0
2003 5/.....	23.4	1.0	0.8	0.9	0.9	1.2	1.6	2.0	2.2	3.0	4.6
2002.....	23.7	*	0.8	0.8	0.9	1.4	2.0	2.0	2.6	3.0	4.5
2001.....	23.8	*	1.1	0.7	0.9	1.4	2.0	2.1	2.6	3.4	4.5
2000.....	24.2	1.3	1.0	0.8	1.0	1.5	1.9	2.1	2.5	3.3	4.6
1999.....	24.1	*	1.0	0.7	0.9	1.4	2.1	2.2	2.5	3.7	4.7

Non-Hispanic White, male

2007 5/.....	25.3	*	0.6	0.9	1.0	1.2	1.9	2.0	2.5	3.0	4.8
2006 5/.....	25.2	*	0.9	0.6	1.0	1.5	2.2	2.4	2.6	3.1	4.9
2005 5/.....	25.4	*	0.8	0.8	1.0	1.5	1.9	1.9	2.1	3.1	4.8
2004 5/.....	25.1	*	0.9	1.0	0.7	1.4	1.8	2.2	2.6	3.2	4.8
2003 5/.....	26.0	*	0.9	1.0	1.0	1.4	1.8	2.3	2.9	3.4	5.7
2002.....	26.2	*	0.9	0.8	1.1	1.7	2.2	2.4	2.9	3.5	5.5
2001.....	25.9	*	1.0	0.7	1.0	1.6	2.3	2.3	3.2	3.9	5.5
2000.....	26.2	*	1.1	1.0	1.1	1.8	2.5	2.2	3.0	3.9	5.7
1999.....	26.1	*	1.0	0.8	1.0	1.7	2.8	2.4	2.9	4.3	5.7

Non-Hispanic White, female

2007 5/.....	19.6	*	0.8	0.5	0.7	0.9	1.2	1.2	1.5	2.2	2.5
2006 5/.....	20.2	*	0.9	0.5	0.6	0.7	1.2	1.5	1.6	2.3	3.3
2005 5/.....	20.2	*	0.6	0.6	0.6	1.0	1.2	1.2	2.0	2.4	3.1
2004 5/.....	20.4	*	0.6	0.7	0.5	1.1	1.3	1.2	1.8	2.0	3.2
2003 5/.....	20.8	*	0.7	0.7	0.8	0.9	1.3	1.7	1.6	2.6	3.4
2002.....	21.2	*	0.7	0.8	0.7	1.1	1.7	1.6	2.3	2.5	3.6
2001.....	21.8	*	1.1	0.6	0.8	1.1	1.7	1.8	2.1	2.9	3.5
2000.....	22.3	*	0.9	0.7	0.9	1.1	1.4	1.9	2.0	2.8	3.5
1999.....	22.2	*	0.9	0.6	0.7	1.2	1.5	1.9	2.1	3.0	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic Black, both sexes

2007 5/.....	13.6	*	0.8	*	1.0	1.3	1.8	2.3	2.6	3.3	7.3
2006 5/.....	13.8	*	*	*	0.9	1.3	2.0	2.0	3.0	4.3	6.2
2005 5/.....	13.4	*	1.0	*	0.8	1.4	1.9	2.9	3.9	3.3	5.9
2004 5/.....	14.0	*	*	*	0.8	1.4	2.0	3.0	3.3	3.8	5.4
2003 5/.....	13.9	*	*	0.7	1.1	1.1	1.8	2.6	3.4	4.3	6.5
2002.....	14.1	*	*	*	1.0	1.4	1.5	2.5	3.5	3.8	5.8
2001.....	14.5	*	*	0.8	1.1	1.5	2.2	3.3	3.7	4.7	6.8
2000.....	14.3	*	*	0.6	0.7	2.0	2.7	2.9	3.2	4.5	6.4
1999.....	14.5	*	*	0.6	1.3	1.8	3.1	2.7	3.3	4.7	6.5

Non-Hispanic Black, male

2007 5/.....	14.5	*	*	*	1.3	1.3	2.1	2.8	2.8	4.2	8.8
2006 5/.....	15.5	*	*	*	*	1.6	2.3	2.2	3.7	5.1	8.3
2005 5/.....	14.6	*	*	*	*	1.4	2.5	3.0	4.9	3.9	7.0
2004 5/.....	14.8	*	*	*	*	1.4	2.5	3.1	3.8	4.5	6.2
2003 5/.....	14.5	*	*	*	1.3	*	2.2	2.6	4.4	4.9	8.1
2002.....	15.1	*	*	*	1.2	1.6	1.4	2.9	4.6	5.1	7.3
2001.....	15.6	*	*	*	1.7	1.8	2.2	3.9	4.1	6.8	8.4
2000.....	15.2	*	*	*	*	2.4	3.3	2.7	4.6	4.8	6.7
1999.....	15.6	*	*	*	1.7	2.1	3.2	3.5	4.3	5.5	8.5

Non-Hispanic Black, female

2007 5/.....	12.8	*	*	*	*	1.2	1.5	1.8	2.4	2.6	5.9
2006 5/.....	12.3	*	*	*	*	*	1.7	1.9	2.4	3.5	4.4
2005 5/.....	12.2	*	*	*	*	1.3	*	2.8	3.1	2.9	5.0
2004 5/.....	13.2	*	*	*	*	1.4	1.4	2.8	2.8	3.1	4.7
2003 5/.....	13.3	*	*	*	*	*	1.4	2.5	2.5	3.8	5.0
2002.....	13.1	*	*	*	*	*	1.6	2.1	2.6	2.6	4.4
2001.....	13.5	*	*	*	*	*	2.2	2.8	3.3	2.9	5.4
2000.....	13.5	*	*	*	*	1.5	2.1	3.0	1.9	4.3	6.2
1999.....	13.6	*	*	*	*	1.4	2.9	2.1	2.3	4.0	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hodgkin's disease (C81)

All origins 3/, both sexes

2007.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2006.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3	0.4
2005.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2004.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4	0.3
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4	0.3
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4	0.3

All origins 3/, male

2007.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.4	0.4
2006.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.6
2005.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4	0.5
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5	0.4

All origins 3/, female

2007.....	0.4	*	*	*	*	*	*	0.2	0.2	0.2	0.2
2006.....	0.4	*	*	*	*	*	*	0.3	0.2	0.3	0.2
2005.....	0.4	*	*	*	*	*	0.3	0.2	0.3	0.2	0.2
2004.....	0.4	*	*	*	*	*	*	0.3	0.3	0.2	0.3
2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3	0.2
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hodgkin's disease (C81)

Non-Hispanic 4/, both sexes

2007.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3	0.3
2006.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.4
2005.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.2	0.4	0.4	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.5	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.4	0.3

Non-Hispanic 4/, male

2007.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2006.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.6
2005.....	0.5	*	*	*	*	*	*	0.4	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.6	0.4	0.5
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.5	0.5	0.5	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.5	0.5	0.5	0.4

Non-Hispanic 4/, female

2007.....	0.4	*	*	*	*	*	*	*	0.3	0.2	0.2
2006.....	0.4	*	*	*	*	*	*	0.3	0.3	0.3	0.2
2005.....	0.4	*	*	*	*	*	0.3	0.2	0.3	0.2	0.2
2004.....	0.4	*	*	*	*	*	*	0.4	0.4	0.2	0.3
2003.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.2
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2000.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3	0.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hodgkin's disease (C81)

Non-Hispanic White, both sexes

2007 5/.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2006 5/.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.4
2005 5/.....	0.5	*	*	*	*	*	0.3	0.2	0.3	0.3	0.3
2004 5/.....	0.5	*	*	*	*	*	0.2	0.3	0.4	0.3	0.3
2003 5/.....	0.6	*	*	*	*	*	0.2	0.4	0.5	0.4	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.3
1999.....	0.6	*	*	*	*	0.2	0.4	0.5	0.4	0.5	0.4

Non-Hispanic White, male

2007 5/.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.4	0.4
2006 5/.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5	0.6
2005 5/.....	0.5	*	*	*	*	*	*	*	*	0.3	0.3
2004 5/.....	0.6	*	*	*	*	*	*	0.4	0.4	0.3	0.4
2003 5/.....	0.6	*	*	*	*	*	*	0.4	0.6	0.5	0.5
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2001.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5	0.5
2000.....	0.6	*	*	*	*	*	0.3	0.4	0.6	0.4	0.4
1999.....	0.7	*	*	*	*	*	0.4	0.5	0.5	0.6	0.5

Non-Hispanic White, female

2007 5/.....	0.4	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.4	*	*	*	*	*	*	0.3	*	0.4	*
2005 5/.....	0.5	*	*	*	*	*	0.3	*	*	*	*
2004 5/.....	0.4	*	*	*	*	*	*	*	*	*	0.3
2003 5/.....	0.5	*	*	*	*	*	*	0.4	0.4	0.4	0.3
2002.....	0.5	*	*	*	*	*	0.4	0.4	0.5	0.4	0.3
2001.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3	0.3
2000.....	0.5	*	*	*	*	*	*	0.3	0.5	0.4	0.3
1999.....	0.5	*	*	*	*	*	*	0.4	0.3	0.4	0.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Non-Hodgkin's lymphoma (C82-C85)

All origins 3/, both sexes

2007.....	6.8	*	*	*	0.1	0.2	0.4	0.5	0.5	0.9	1.5
2006.....	6.9	*	*	*	0.1	0.2	0.4	0.5	0.6	0.8	1.5
2005.....	7.0	*	*	*	0.1	0.2	0.4	0.4	0.7	0.9	1.6
2004.....	7.1	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9	1.5
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1	1.7
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2	1.9
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4	2.0
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3	2.1
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4	2.1

All origins 3/, male

2007.....	7.4	*	*	*	*	0.3	0.5	0.6	0.7	1.1	2.1
2006.....	7.4	*	*	*	*	0.3	0.5	0.6	0.7	1.0	1.9
2005.....	7.6	*	*	*	0.2	0.3	0.5	0.5	0.8	1.1	2.2
2004.....	7.6	*	*	*	*	0.3	0.5	0.8	0.8	1.2	2.1
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4	2.2
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5	2.4
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8	2.7
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7	2.6
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8	2.8

All origins 3/, female

2007.....	6.2	*	*	*	*	*	0.3	0.4	0.4	0.7	0.9
2006.....	6.4	*	*	*	*	*	0.3	0.4	0.5	0.7	1.1
2005.....	6.5	*	*	*	*	*	0.3	0.4	0.5	0.7	1.1
2004.....	6.6	*	*	*	*	*	0.3	0.4	0.6	0.6	0.9
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7	1.2
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8	1.3
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0	1.3
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9	1.5
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic 4/, both sexes

2007.....	7.5	*	*	*	*	0.2	0.4	0.5	0.6	0.9	1.5
2006.....	7.6	*	*	*	0.1	0.2	0.4	0.5	0.6	0.9	1.5
2005.....	7.8	*	*	*	0.2	0.2	0.4	0.5	0.7	1.0	1.7
2004.....	7.9	*	*	*	*	0.3	0.4	0.6	0.7	1.0	1.5
2003.....	8.1	*	*	*	0.1	0.3	0.4	0.5	0.7	1.1	1.7
2002.....	8.3	*	*	*	0.2	0.2	0.5	0.6	0.9	1.2	1.9
2001.....	8.6	*	*	*	0.1	0.3	0.5	0.7	0.9	1.4	2.0
2000.....	8.8	*	*	0.1	0.1	0.3	0.7	0.7	0.9	1.3	2.0
1999.....	8.9	*	*	*	0.1	0.3	0.5	0.6	0.9	1.5	2.1

Non-Hispanic 4/, male

2007.....	8.3	*	*	*	*	0.3	0.5	0.6	0.8	1.1	2.1
2006.....	8.3	*	*	*	*	0.3	0.5	0.6	0.8	1.0	2.0
2005.....	8.5	*	*	*	0.3	0.3	0.5	0.5	0.9	1.2	2.3
2004.....	8.5	*	*	*	*	0.3	0.5	0.9	0.8	1.3	2.0
2003.....	8.8	*	*	*	*	0.3	0.5	0.6	1.0	1.4	2.2
2002.....	9.0	*	*	*	0.3	0.3	0.6	0.8	1.1	1.5	2.5
2001.....	9.2	*	*	*	*	0.4	0.5	0.8	1.1	1.8	2.7
2000.....	9.4	*	*	*	*	0.5	0.9	0.7	1.1	1.8	2.6
1999.....	9.4	*	*	*	0.2	0.3	0.7	0.8	1.1	1.9	2.8

Non-Hispanic 4/, female

2007.....	6.8	*	*	*	*	*	0.3	0.3	0.4	0.7	0.9
2006.....	7.0	*	*	*	*	*	0.3	0.4	0.5	0.8	1.1
2005.....	7.1	*	*	*	*	*	0.3	0.4	0.5	0.7	1.1
2004.....	7.3	*	*	*	*	*	0.3	0.3	0.6	0.6	1.0
2003.....	7.4	*	*	*	*	*	0.3	0.5	0.5	0.8	1.3
2002.....	7.7	*	*	*	*	*	0.4	0.5	0.6	0.8	1.3
2001.....	8.0	*	*	*	*	*	0.4	0.6	0.7	1.0	1.4
2000.....	8.3	*	*	*	*	*	0.5	0.6	0.6	0.9	1.5
1999.....	8.4	*	*	*	*	0.2	0.3	0.4	0.6	1.1	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic White, both sexes

2007 5/.....	8.6	*	*	*	*	0.2	0.4	0.4	0.5	1.0	1.3
2006 5/.....	8.7	*	*	*	*	0.2	0.3	0.5	0.6	0.8	1.3
2005 5/.....	8.9	*	*	*	0.2	0.2	0.3	0.4	0.6	0.9	1.6
2004 5/.....	9.0	*	*	*	*	0.3	0.4	0.5	0.6	1.0	1.4
2003 5/.....	9.3	*	*	*	*	0.2	0.4	0.4	0.6	1.0	1.7
2002.....	9.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1	1.8
2001.....	9.7	*	*	*	*	0.3	0.4	0.6	0.8	1.3	2.0
2000.....	10.1	*	*	*	0.2	0.3	0.6	0.6	0.8	1.3	2.0
1999.....	10.1	*	*	*	*	0.2	0.5	0.6	0.8	1.4	2.0

Non-Hispanic White, male

2007 5/.....	9.4	*	*	*	*	*	0.4	0.5	0.8	1.2	1.8
2006 5/.....	9.4	*	*	*	*	*	0.4	0.6	0.7	0.9	1.8
2005 5/.....	9.7	*	*	*	0.3	0.4	0.4	0.4	0.7	1.1	2.2
2004 5/.....	9.6	*	*	*	*	0.3	0.5	0.8	0.7	1.3	1.9
2003 5/.....	10.1	*	*	*	*	0.3	0.5	0.5	0.9	1.4	2.1
2002.....	10.2	*	*	*	0.3	*	0.6	0.8	0.9	1.4	2.4
2001.....	10.4	*	*	*	*	0.4	0.5	0.7	1.1	1.7	2.6
2000.....	10.6	*	*	*	*	0.5	0.8	0.6	1.0	1.7	2.7
1999.....	10.6	*	*	*	*	*	0.7	0.7	1.0	1.8	2.7

Non-Hispanic White, female

2007 5/.....	7.9	*	*	*	*	*	*	*	*	0.8	0.7
2006 5/.....	8.1	*	*	*	*	*	*	0.4	0.4	0.7	0.9
2005 5/.....	8.2	*	*	*	*	*	*	0.4	0.5	0.6	0.9
2004 5/.....	8.4	*	*	*	*	*	0.3	*	0.5	0.6	0.9
2003 5/.....	8.5	*	*	*	*	*	0.3	0.4	0.4	0.7	1.2
2002.....	8.8	*	*	*	*	*	0.3	0.4	0.5	0.8	1.3
2001.....	9.2	*	*	*	*	*	0.4	0.5	0.6	1.0	1.3
2000.....	9.5	*	*	*	*	*	0.4	0.5	0.6	0.8	1.4
1999.....	9.6	*	*	*	*	*	*	0.4	0.6	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic Black, both sexes

2007 5/.....	3.6	*	*	*	*	*	*	0.9	0.9	1.0	2.8
2006 5/.....	3.9	*	*	*	*	*	*	0.8	0.9	1.9	2.5
2005 5/.....	3.7	*	*	*	*	*	0.7	*	1.5	1.5	2.5
2004 5/.....	3.8	*	*	*	*	*	0.7	1.2	1.1	1.3	1.8
2003 5/.....	3.7	*	*	*	*	*	*	0.9	1.3	1.5	2.4
2002.....	4.0	*	*	*	*	*	*	0.9	1.5	1.3	2.7
2001.....	4.1	*	*	*	*	*	*	1.2	1.2	2.1	2.5
2000.....	3.9	*	*	*	*	*	1.0	1.2	1.0	1.9	2.4
1999.....	4.1	*	*	*	*	*	*	0.9	1.3	2.2	2.7

Non-Hispanic Black, male

2007 5/.....	4.1	*	*	*	*	*	*	*	*	*	3.9
2006 5/.....	4.4	*	*	*	*	*	*	*	*	2.2	3.5
2005 5/.....	4.2	*	*	*	*	*	*	*	2.0	1.7	3.0
2004 5/.....	4.1	*	*	*	*	*	*	*	*	*	2.3
2003 5/.....	3.9	*	*	*	*	*	*	*	1.9	1.7	3.1
2002.....	4.5	*	*	*	*	*	*	*	2.3	2.0	3.8
2001.....	4.7	*	*	*	*	*	*	*	*	3.2	3.5
2000.....	4.3	*	*	*	*	*	*	*	*	2.6	2.6
1999.....	4.7	*	*	*	*	*	*	*	1.8	2.7	4.0

Non-Hispanic Black, female

2007 5/.....	3.2	*	*	*	*	*	*	*	*	*	1.8
2006 5/.....	3.4	*	*	*	*	*	*	*	*	1.6	1.6
2005 5/.....	3.3	*	*	*	*	*	*	*	*	*	2.1
2004 5/.....	3.4	*	*	*	*	*	*	*	*	*	*
2003 5/.....	3.5	*	*	*	*	*	*	*	*	1.4	1.7
2002.....	3.6	*	*	*	*	*	*	*	*	*	1.7
2001.....	3.7	*	*	*	*	*	*	*	*	*	1.6
2000.....	3.5	*	*	*	*	*	*	*	*	*	2.1
1999.....	3.6	*	*	*	*	*	*	*	*	1.7	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Leukemia (C91-C95)

All origins 3/, both sexes

2007.....	7.2	0.5	0.6	0.7	0.9	0.9	1.1	1.0	1.2	1.2	1.9
2006.....	7.3	0.8	0.8	0.5	0.7	1.0	1.2	1.1	1.2	1.4	1.8
2005.....	7.3	0.5	0.7	0.6	0.7	1.1	1.2	1.1	1.2	1.4	1.7
2004.....	7.3	0.6	0.8	0.8	0.7	1.0	1.1	0.9	1.2	1.4	1.8
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5	2.1
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5	1.9
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6	1.9
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6	1.9
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6	1.9

All origins 3/, male

2007.....	8.3	*	0.6	0.8	0.9	1.1	1.2	1.1	1.3	1.3	2.1
2006.....	8.4	*	0.8	0.6	0.8	1.2	1.5	1.3	1.4	1.6	2.0
2005.....	8.4	*	1.0	0.8	0.8	1.2	1.5	1.3	1.2	1.4	1.8
2004.....	8.3	*	0.9	1.0	0.8	1.2	1.3	1.0	1.4	1.5	2.1
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6	2.4
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6	2.2
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8	2.1
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6	2.1
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8	2.1

All origins 3/, female

2007.....	6.2	*	0.7	0.5	0.8	0.8	0.9	0.8	1.1	1.1	1.6
2006.....	6.3	*	0.8	0.5	0.6	0.7	0.9	0.8	1.0	1.3	1.7
2005.....	6.2	*	0.5	0.5	0.7	0.9	0.8	0.9	1.2	1.3	1.6
2004.....	6.3	*	0.6	0.6	0.7	0.8	0.8	0.8	0.9	1.2	1.6
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4	1.7
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3	1.6
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3	1.7
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6	1.7
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Leukemia (C91-C95)											
Hispanic, both sexes											
2007.....	2.8	*	*	1.0	1.4	1.4	1.6	1.2	1.5	1.1	2.2
2006.....	2.8	*	0.7	0.6	0.9	1.4	1.7	1.2	1.3	1.5	1.9
2005.....	3.0	*	0.7	0.9	1.2	1.6	1.9	1.2	1.4	1.4	1.7
2004.....	2.8	*	0.8	1.1	1.5	1.5	1.5	0.9	1.3	1.4	1.7
2003.....	2.9	*	1.2	1.0	1.3	1.2	1.5	1.0	1.3	1.3	2.2
2002.....	2.8	*	0.8	1.1	1.2	1.7	1.5	1.0	1.2	2.0	2.2
2001.....	3.0	*	1.3	1.2	1.1	2.0	1.5	1.3	1.6	1.5	2.3
2000.....	2.9	*	0.9	0.8	1.4	1.5	1.5	1.1	1.1	1.8	2.0
1999.....	2.9	*	0.9	0.8	1.4	1.2	1.9	1.5	1.2	1.9	2.1
Hispanic, male											
2007.....	3.0	*	*	1.2	1.2	1.6	1.8	1.3	1.5	1.2	2.2
2006.....	3.0	*	*	*	*	1.6	2.0	1.5	1.3	1.3	2.0
2005.....	3.4	*	*	*	1.3	2.1	2.4	1.4	1.6	1.7	1.8
2004.....	3.1	*	*	1.5	1.7	2.0	2.0	1.0	1.7	1.4	2.2
2003.....	3.2	*	1.4	1.5	1.1	1.8	1.6	1.1	1.7	1.4	2.5
2002.....	3.0	*	*	*	1.3	2.2	1.8	1.2	1.7	2.1	2.5
2001.....	3.4	*	1.4	1.3	*	2.5	2.0	1.6	2.0	2.0	2.7
2000.....	3.3	*	*	*	2.0	1.7	1.6	1.6	1.3	1.6	2.4
1999.....	3.1	*	*	*	1.3	*	2.3	1.5	1.4	2.0	1.8
Hispanic, female											
2007.....	2.7	*	*	*	1.5	1.2	1.5	1.1	1.5	*	2.1
2006.....	2.6	*	*	*	*	1.1	1.3	*	1.4	1.7	1.7
2005.....	2.5	*	*	*	*	*	1.4	*	1.1	*	1.5
2004.....	2.4	*	*	*	1.3	*	*	*	*	1.4	*
2003.....	2.6	*	*	*	1.5	*	1.4	*	*	*	1.9
2002.....	2.6	*	*	1.2	*	*	*	*	*	1.9	1.9
2001.....	2.6	*	*	*	1.3	1.3	*	*	*	*	1.9
2000.....	2.6	*	*	*	*	*	1.4	*	*	2.1	*
1999.....	2.8	*	*	*	1.5	1.5	1.6	1.5	*	1.8	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cause of death	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Leukemia (C91-C95)											
Non-Hispanic 4/, both sexes											
2007.....	8.0	*	0.7	0.6	0.7	0.9	1.0	0.9	1.1	1.2	1.8
2006.....	8.1	0.7	0.8	0.5	0.7	0.9	1.1	1.0	1.2	1.4	1.8
2005.....	8.0	*	0.8	0.6	0.6	1.0	1.0	1.0	1.2	1.4	1.7
2004.....	8.0	*	0.8	0.7	0.6	0.9	1.0	0.9	1.2	1.4	1.9
2003.....	8.1	0.8	0.7	0.7	0.8	0.8	1.0	1.1	1.1	1.5	2.1
2002.....	8.2	*	0.6	0.7	0.7	1.1	1.1	1.0	1.4	1.4	1.9
2001.....	8.2	*	0.9	0.7	0.8	1.0	1.2	1.2	1.3	1.5	1.8
2000.....	8.2	1.0	0.9	0.7	0.7	1.1	1.1	1.2	1.2	1.6	1.9
1999.....	8.2	*	0.7	0.6	0.8	1.1	1.3	1.1	1.3	1.6	1.9
Non-Hispanic 4/, male											
2007.....	9.3	*	0.7	0.8	0.8	1.0	1.1	1.1	1.3	1.3	2.0
2006.....	9.4	*	0.8	0.6	0.8	1.1	1.4	1.2	1.4	1.6	2.0
2005.....	9.3	*	1.0	0.7	0.7	1.0	1.3	1.2	1.1	1.4	1.8
2004.....	9.2	*	0.9	0.8	0.6	1.0	1.1	0.9	1.4	1.5	2.1
2003.....	9.3	*	0.8	0.7	0.9	1.1	1.1	1.3	1.4	1.6	2.4
2002.....	9.4	*	0.7	0.7	0.8	1.2	1.2	1.2	1.5	1.5	2.2
2001.....	9.3	*	0.9	0.7	0.9	1.1	1.4	1.4	1.5	1.7	2.0
2000.....	9.3	*	0.9	0.8	0.9	1.4	1.4	1.3	1.5	1.6	2.1
1999.....	9.2	*	0.8	0.6	0.9	1.3	1.6	1.3	1.4	1.8	2.1
Non-Hispanic 4/, female											
2007.....	6.8	*	0.7	0.5	0.6	0.7	0.8	0.7	1.0	1.1	1.5
2006.....	6.9	*	0.8	0.5	0.6	0.6	0.8	0.8	0.9	1.2	1.6
2005.....	6.8	*	0.5	0.4	0.6	0.9	0.7	0.8	1.2	1.3	1.6
2004.....	6.9	*	0.6	0.6	0.5	0.8	0.8	0.8	0.9	1.2	1.7
2003.....	7.0	*	0.7	0.7	0.6	0.6	0.8	0.8	0.8	1.5	1.7
2002.....	7.0	*	0.5	0.6	0.6	0.9	0.9	0.8	1.2	1.3	1.6
2001.....	7.2	*	0.9	0.6	0.6	0.9	1.0	1.0	1.1	1.4	1.7
2000.....	7.2	*	0.9	0.5	0.6	0.9	0.7	1.1	0.9	1.5	1.7
1999.....	7.2	*	0.7	0.6	0.7	0.8	1.0	1.0	1.2	1.4	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Leukemia (C91-C95)

Non-Hispanic White, both sexes

2007 5/.....	9.1	*	0.7	0.6	0.8	0.8	0.9	0.9	1.1	1.2	1.6
2006 5/.....	9.2	*	0.9	0.5	0.7	0.9	1.1	1.1	1.1	1.4	1.8
2005 5/.....	9.1	*	0.7	0.6	0.6	1.0	1.0	0.9	1.1	1.4	1.7
2004 5/.....	9.0	*	0.7	0.8	0.5	0.9	1.0	0.8	1.2	1.3	1.8
2003 5/.....	9.1	1.0	0.8	0.8	0.7	0.8	0.9	1.1	1.0	1.4	2.0
2002.....	9.1	*	0.7	0.7	0.7	1.1	1.2	1.0	1.3	1.4	2.0
2001.....	9.1	*	1.0	0.6	0.7	1.0	1.2	1.0	1.3	1.5	1.8
2000.....	9.2	1.3	0.9	0.7	0.8	1.0	1.1	1.1	1.1	1.5	1.8
1999.....	9.1	*	0.8	0.6	0.7	1.0	1.2	1.2	1.2	1.6	1.9

Non-Hispanic White, male

2007 5/.....	10.6	*	0.6	0.7	0.9	0.9	1.1	1.0	1.2	1.3	1.9
2006 5/.....	10.6	*	0.8	0.5	0.9	1.2	1.5	1.4	1.3	1.6	1.9
2005 5/.....	10.5	*	0.8	0.7	0.6	1.0	1.3	1.2	1.1	1.4	1.7
2004 5/.....	10.3	*	0.8	0.9	0.6	1.0	1.1	1.0	1.4	1.4	2.0
2003 5/.....	10.5	*	0.8	0.9	0.9	1.1	1.1	1.3	1.3	1.4	2.4
2002.....	10.5	*	0.8	0.7	0.8	1.2	1.3	1.2	1.5	1.5	2.3
2001.....	10.3	*	1.0	0.6	0.8	1.1	1.4	1.2	1.5	1.6	1.9
2000.....	10.4	*	1.0	0.8	0.9	1.2	1.4	1.2	1.3	1.6	2.1
1999.....	10.2	*	0.8	0.7	0.8	1.2	1.6	1.3	1.3	1.8	2.1

Non-Hispanic White, female

2007 5/.....	7.6	*	0.8	0.4	0.6	0.7	0.8	0.7	1.0	1.1	1.4
2006 5/.....	7.8	*	0.9	0.5	0.5	0.7	0.8	0.8	0.9	1.2	1.7
2005 5/.....	7.7	*	0.5	0.5	0.5	0.9	0.8	0.7	1.2	1.5	1.6
2004 5/.....	7.7	*	0.6	0.7	0.5	0.8	0.8	0.7	1.0	1.1	1.7
2003 5/.....	7.8	*	0.7	0.7	0.6	0.6	0.7	0.9	0.7	1.5	1.7
2002.....	7.8	*	0.5	0.7	0.6	0.9	1.0	0.7	1.2	1.3	1.7
2001.....	8.0	*	1.1	0.6	0.7	0.8	1.0	0.9	1.1	1.4	1.7
2000.....	8.1	*	0.9	0.6	0.7	0.8	0.7	1.0	0.8	1.5	1.6
1999.....	8.0	*	0.9	0.5	0.6	0.8	0.9	1.0	1.2	1.4	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Leukemia (C91-C95)

Non-Hispanic Black, both sexes

2007 5/.....	4.7	*	*	*	0.8	0.9	1.1	1.1	1.3	1.4	2.7
2006 5/.....	4.7	*	*	*	0.7	0.9	1.2	0.9	1.4	1.6	2.2
2005 5/.....	4.5	*	1.0	*	0.7	1.1	1.0	1.6	1.6	1.2	2.0
2004 5/.....	5.0	*	*	*	0.7	1.1	0.9	1.2	1.5	1.8	2.3
2003 5/.....	4.8	*	*	*	0.9	0.7	1.1	1.0	1.4	2.1	2.6
2002.....	4.8	*	*	*	0.7	1.0	0.8	1.1	1.5	1.5	1.7
2001.....	4.9	*	*	0.8	0.9	0.9	1.2	1.5	1.4	1.9	2.3
2000.....	4.9	*	*	*	0.6	1.5	1.3	1.5	1.9	1.7	2.5
1999.....	4.9	*	*	*	1.1	1.2	1.8	1.1	1.4	1.8	1.9

Non-Hispanic Black, male

2007 5/.....	5.2	*	*	*	*	*	1.3	*	*	1.7	3.1
2006 5/.....	5.5	*	*	*	*	*	1.5	*	1.7	1.8	3.1
2005 5/.....	5.1	*	*	*	*	*	1.6	1.6	*	*	2.3
2004 5/.....	5.5	*	*	*	*	*	*	*	1.8	2.2	2.4
2003 5/.....	5.2	*	*	*	*	*	1.4	*	1.8	2.6	3.0
2002.....	5.3	*	*	*	*	*	*	*	1.7	1.9	1.8
2001.....	5.4	*	*	*	1.5	*	*	1.8	1.8	2.4	2.7
2000.....	5.5	*	*	*	*	2.1	1.8	*	2.6	1.5	2.5
1999.....	5.4	*	*	*	1.5	1.5	1.7	*	1.8	1.9	2.3

Non-Hispanic Black, female

2007 5/.....	4.3	*	*	*	*	*	*	*	*	*	2.3
2006 5/.....	4.0	*	*	*	*	*	*	*	*	*	1.5
2005 5/.....	4.1	*	*	*	*	*	*	1.5	1.6	*	1.7
2004 5/.....	4.5	*	*	*	*	*	*	*	*	1.5	2.1
2003 5/.....	4.4	*	*	*	*	*	*	*	*	1.7	2.1
2002.....	4.4	*	*	*	*	*	*	*	*	*	1.6
2001.....	4.4	*	*	*	*	*	*	*	*	1.4	2.0
2000.....	4.4	*	*	*	*	*	*	1.6	*	1.9	2.4
1999.....	4.4	*	*	*	*	*	1.9	*	*	1.8	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes

2007.....	21.0	0.5	0.7	0.6	0.6	1.1	1.5	1.5	2.0	2.9	5.1
2006.....	21.1	0.5	0.7	0.7	0.5	1.1	1.3	1.6	1.9	2.7	5.4
2005.....	21.2	*	0.8	0.6	0.8	1.3	1.5	1.5	2.0	2.9	5.3
2004.....	21.1	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.1	5.3
2003.....	21.5	*	0.7	0.6	0.7	1.2	1.3	1.6	1.9	3.1	5.3
2002.....	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3	5.8
2001.....	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2	5.9
2000.....	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4	5.9
1999.....	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6	6.7

All origins 3/, male

2007.....	22.1	*	0.7	0.6	0.5	1.4	2.0	1.8	2.4	3.1	5.2
2006.....	21.9	*	0.7	0.8	0.5	1.3	1.8	1.9	2.4	2.9	5.6
2005.....	22.1	*	0.8	0.6	0.8	1.4	1.9	1.9	2.4	3.1	5.9
2004.....	21.9	*	0.7	0.8	0.8	1.5	1.8	1.9	2.3	3.6	5.5
2003.....	22.3	*	0.8	0.8	0.7	1.5	1.6	1.9	2.4	3.5	5.9
2002.....	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7	6.2
2001.....	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5	6.4
2000.....	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7	6.4
1999.....	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1	7.6

All origins 3/, female

2007.....	19.8	*	0.7	0.6	0.7	0.8	1.0	1.1	1.5	2.8	4.9
2006.....	20.2	*	0.6	0.6	0.5	0.9	0.8	1.3	1.4	2.5	5.2
2005.....	20.4	*	0.7	0.6	0.7	1.2	1.0	1.0	1.6	2.7	4.7
2004.....	20.3	*	0.9	0.5	0.6	1.0	0.9	1.1	1.7	2.5	5.1
2003.....	20.8	*	0.5	0.5	0.7	0.8	1.1	1.2	1.4	2.7	4.8
2002.....	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8	5.4
2001.....	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0	5.3
2000.....	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0	5.3
1999.....	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Hispanic, both sexes

2007	7.5	*	*	0.6	0.6	1.2	1.7	1.3	1.9	2.6	3.7
2006	7.7	*	0.6	0.5	0.7	1.2	1.7	1.1	1.5	2.1	4.2
2005	7.6	*	0.6	*	0.6	1.3	1.4	1.5	2.0	2.0	4.2
2004	7.2	*	0.6	*	0.6	1.1	1.4	1.3	1.4	1.8	3.6
2003	7.7	*	0.6	0.6	0.7	1.5	1.2	1.3	1.6	2.7	3.9
2002	7.8	*	*	0.6	0.9	1.2	1.9	1.1	1.8	2.2	4.0
2001	7.7	*	*	*	0.9	1.3	1.2	1.0	1.4	2.8	4.3
2000	7.8	*	0.8	*	0.9	1.4	1.1	1.4	1.7	2.1	3.6
1999	8.1	*	*	0.7	0.7	1.3	1.4	1.5	1.7	2.5	4.9

Hispanic, male

2007	7.6	*	*	*	*	1.6	2.6	1.6	2.0	2.8	3.7
2006	7.6	*	*	*	*	1.3	2.5	1.4	2.0	1.9	3.5
2005	7.8	*	*	*	*	1.6	1.7	1.7	2.3	2.2	5.0
2004	7.1	*	*	*	*	1.8	1.9	1.5	1.6	2.0	3.9
2003	7.9	*	*	*	*	2.1	1.2	1.9	2.3	2.7	4.0
2002	7.8	*	*	*	*	1.5	2.2	1.5	2.0	2.3	4.0
2001	7.7	*	*	*	*	1.5	2.0	1.1	2.0	2.7	4.2
2000	7.7	*	*	*	*	1.5	1.3	1.6	2.3	2.2	3.8
1999	8.0	*	*	*	*	1.7	2.0	1.9	1.8	2.5	4.8

Hispanic, female

2007	7.5	*	*	*	*	*	*	*	1.7	2.3	3.6
2006	7.8	*	*	*	*	*	*	*	*	2.4	4.8
2005	7.4	*	*	*	*	*	*	1.2	1.6	1.9	3.4
2004	7.3	*	*	*	*	*	*	*	1.2	1.6	3.2
2003	7.6	*	*	*	*	*	1.2	*	*	2.7	3.8
2002	7.7	*	*	*	*	*	1.5	*	1.7	2.1	3.9
2001	7.7	*	*	*	*	*	*	*	*	2.8	4.4
2000	7.9	*	*	*	*	*	*	*	*	1.9	3.4
1999	8.2	*	*	*	*	*	*	*	1.5	2.5	4.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 4/, both sexes

2007.....	23.3	*	0.8	0.6	0.6	1.1	1.5	1.5	2.0	3.0	5.3
2006.....	23.4	*	0.7	0.7	0.5	1.1	1.3	1.7	2.0	2.8	5.6
2005.....	23.5	*	0.8	0.6	0.8	1.3	1.5	1.5	2.0	3.1	5.5
2004.....	23.3	0.6	0.8	0.7	0.7	1.3	1.4	1.6	2.1	3.3	5.6
2003.....	23.7	*	0.7	0.6	0.7	1.1	1.3	1.6	1.9	3.2	5.5
2002.....	24.0	*	0.9	0.7	0.7	1.1	1.4	1.6	2.2	3.4	6.0
2001.....	24.1	*	0.8	0.6	0.8	1.2	1.4	1.8	2.1	3.3	6.0
2000.....	24.4	0.7	0.8	0.6	0.6	1.2	1.5	1.5	2.0	3.6	6.1
1999.....	25.8	*	0.8	0.7	0.7	1.2	1.5	1.6	2.3	3.8	6.8

Non-Hispanic 4/, male

2007.....	24.8	*	0.7	0.6	0.5	1.3	1.9	1.9	2.5	3.2	5.5
2006.....	24.5	*	0.8	0.8	0.5	1.3	1.7	2.1	2.5	3.1	6.0
2005.....	24.5	*	0.8	0.7	0.9	1.4	2.0	2.0	2.4	3.3	6.0
2004.....	24.4	*	0.8	0.8	0.8	1.5	1.8	2.0	2.4	3.9	5.8
2003.....	24.6	*	0.8	0.8	0.7	1.4	1.7	1.8	2.4	3.6	6.1
2002.....	25.0	*	0.9	0.8	0.7	1.3	1.9	2.2	2.6	4.0	6.4
2001.....	24.9	*	1.0	0.7	0.8	1.4	1.9	2.2	2.5	3.6	6.6
2000.....	25.2	*	0.9	0.6	0.7	1.3	2.0	2.0	2.4	4.0	6.7
1999.....	26.8	*	1.0	0.8	0.8	1.5	1.7	2.1	2.9	4.3	7.9

Non-Hispanic 4/, female

2007.....	21.9	*	0.8	0.6	0.7	0.8	1.1	1.2	1.5	2.8	5.1
2006.....	22.2	*	0.6	0.6	0.4	0.8	0.8	1.4	1.6	2.5	5.2
2005.....	22.4	*	0.8	0.6	0.8	1.2	1.0	1.0	1.6	2.8	4.9
2004.....	22.3	*	0.8	0.6	0.6	1.0	1.0	1.2	1.7	2.7	5.4
2003.....	22.8	*	0.6	0.5	0.7	0.8	1.0	1.4	1.5	2.7	4.9
2002.....	22.9	*	0.8	0.6	0.7	1.0	0.8	1.1	1.7	2.9	5.6
2001.....	23.3	*	0.6	0.6	0.7	0.9	0.9	1.4	1.7	3.0	5.4
2000.....	23.7	*	0.8	0.6	0.6	1.0	1.0	1.1	1.7	3.2	5.5
1999.....	25.0	*	0.7	0.7	0.6	0.9	1.2	1.1	1.6	3.3	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic White, both sexes

2007 5/.....	25.5	*	0.8	0.6	0.6	1.1	1.4	1.5	1.9	2.9	5.1
2006 5/.....	25.6	*	0.6	0.8	0.5	1.1	1.3	1.6	2.1	2.8	5.5
2005 5/.....	25.6	*	0.8	0.6	0.8	1.3	1.5	1.4	1.9	3.1	5.3
2004 5/.....	25.4	*	0.8	0.7	0.7	1.2	1.3	1.7	2.1	3.2	5.4
2003 5/.....	25.9	*	0.6	0.7	0.6	1.2	1.4	1.6	1.8	3.2	5.2
2002.....	26.0	*	0.8	0.7	0.7	1.2	1.3	1.6	2.1	3.4	5.7
2001.....	26.0	*	0.9	0.6	0.7	1.2	1.5	1.9	2.0	3.2	5.8
2000.....	26.5	*	0.8	0.7	0.6	1.1	1.6	1.7	2.0	3.4	5.6
1999.....	27.9	*	0.9	0.7	0.6	1.2	1.4	1.7	2.2	3.6	6.4

Non-Hispanic White, male

2007 5/.....	27.2	*	0.7	0.6	0.5	1.4	1.9	1.9	2.4	3.0	5.2
2006 5/.....	27.0	*	0.7	0.9	0.5	1.3	1.8	2.1	2.6	3.1	5.9
2005 5/.....	26.8	*	0.8	0.7	0.7	1.4	2.0	1.9	2.3	3.4	5.9
2004 5/.....	26.5	*	0.9	0.8	0.8	1.4	1.7	2.1	2.6	3.9	5.5
2003 5/.....	26.8	*	0.7	0.9	0.6	1.5	1.7	2.0	2.2	3.8	5.7
2002.....	27.2	*	0.8	0.7	0.6	1.3	1.8	2.1	2.6	3.9	5.9
2001.....	26.8	*	1.1	0.7	0.8	1.4	2.0	2.3	2.4	3.5	6.4
2000.....	27.2	*	1.0	0.7	0.7	1.3	2.0	2.2	2.5	3.8	6.0
1999.....	28.8	*	1.1	0.8	0.7	1.4	1.6	2.3	2.8	4.2	7.4

Non-Hispanic White, female

2007 5/.....	23.9	*	0.8	0.6	0.7	0.8	1.0	1.2	1.5	2.8	5.0
2006 5/.....	24.3	*	0.5	0.6	0.5	0.8	0.8	1.2	1.6	2.4	5.0
2005 5/.....	24.5	*	0.7	0.5	0.8	1.2	1.0	0.8	1.5	2.8	4.7
2004 5/.....	24.4	*	0.8	0.6	0.5	1.1	0.9	1.2	1.6	2.6	5.3
2003 5/.....	25.0	*	0.6	0.5	0.7	0.8	1.0	1.2	1.4	2.5	4.7
2002.....	24.9	*	0.7	0.6	0.8	1.0	0.9	1.1	1.6	2.9	5.5
2001.....	25.3	*	0.7	0.6	0.7	0.9	1.0	1.5	1.6	2.8	5.1
2000.....	25.8	*	0.6	0.7	0.6	0.9	1.1	1.1	1.6	2.9	5.2
1999.....	27.0	*	0.8	0.6	0.5	1.0	1.2	1.0	1.6	3.1	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic Black, both sexes

2007 5/.....	17.3	*	*	*	0.6	1.0	1.6	1.5	2.9	3.9	7.3
2006 5/.....	17.5	*	1.0	0.7	*	1.1	1.4	2.2	2.2	3.3	7.4
2005 5/.....	17.7	*	*	0.8	1.1	1.0	1.6	2.1	2.7	3.7	7.3
2004 5/.....	17.8	*	*	0.8	1.1	1.4	1.9	1.5	2.1	4.5	7.7
2003 5/.....	17.9	*	0.9	0.7	0.9	0.9	1.4	1.9	2.8	3.8	7.9
2002.....	18.7	*	1.1	0.8	0.8	1.1	1.6	2.0	2.8	4.7	8.2
2001.....	18.9	*	*	*	0.8	1.2	1.4	2.1	2.7	4.3	7.8
2000.....	18.9	*	0.9	*	0.7	1.6	1.4	1.3	2.0	5.4	10.1
1999.....	20.2	*	*	0.7	0.9	1.3	1.7	1.6	2.5	5.1	10.4

Non-Hispanic Black, male

2007 5/.....	18.2	*	*	*	*	1.2	1.7	1.7	3.9	4.5	7.8
2006 5/.....	18.1	*	*	*	*	1.4	1.4	2.2	2.8	3.4	7.6
2005 5/.....	18.6	*	*	*	1.4	*	2.0	2.5	3.4	3.9	7.8
2004 5/.....	19.0	*	*	*	*	1.8	2.2	1.8	2.0	4.9	8.5
2003 5/.....	18.8	*	*	*	*	*	1.8	1.7	3.1	3.7	9.4
2002.....	19.4	*	*	*	*	1.4	2.3	2.6	2.9	5.4	9.9
2001.....	20.0	*	*	*	*	1.5	2.1	2.6	3.0	4.4	8.3
2000.....	20.0	*	*	*	*	1.8	2.3	1.6	2.1	5.6	12.0
1999.....	21.3	*	*	*	*	1.9	2.1	1.6	3.2	5.2	11.9

Non-Hispanic Black, female

2007 5/.....	16.5	*	*	*	*	*	1.5	*	2.0	3.3	6.8
2006 5/.....	16.9	*	*	*	*	*	1.3	2.3	1.7	3.1	7.3
2005 5/.....	16.8	*	*	*	*	*	*	1.8	2.0	3.6	6.8
2004 5/.....	16.7	*	*	*	1.2	*	1.5	*	2.2	4.2	7.0
2003 5/.....	17.1	*	*	*	*	*	*	2.1	2.4	4.0	6.6
2002.....	18.0	*	*	*	*	*	*	1.5	2.7	4.2	6.8
2001.....	17.8	*	*	*	*	*	*	1.6	2.4	4.3	7.4
2000.....	17.8	*	*	*	*	1.4	*	*	1.9	5.2	8.5
1999.....	19.2	*	*	*	*	*	*	1.5	1.8	4.9	9.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/
Under 1 year 2/
1-4 years
5-9 years
10-14 years
15-19 years
20-24 years
25-29 years
30-34 years
35-39 years
40-44 years

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes

2007.....	4.7	1.4	0.4	0.2	0.2	0.1	0.2	0.3	0.3	0.7	0.9
2006.....	4.7	1.6	0.4	0.2	0.2	0.3	0.2	0.3	0.4	0.6	0.8
2005.....	4.6	1.4	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.0
2004.....	4.6	1.7	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.6	0.8
2003.....	4.7	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.7	0.9
2002.....	4.6	1.7	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6	1.0
2001.....	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7	0.9
2000.....	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7	0.9
1999.....	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6	0.9

All origins 3/, male

2007.....	4.9	1.1	0.3	0.2	0.3	*	0.2	0.4	0.4	0.7	1.0
2006.....	4.7	1.6	0.3	0.2	*	0.2	0.2	0.3	0.4	0.7	0.9
2005.....	4.7	1.0	0.4	*	*	0.3	0.2	0.4	0.5	0.7	1.0
2004.....	4.7	1.7	0.3	0.2	0.2	0.2	0.4	0.4	0.6	0.6	0.9
2003.....	4.7	1.6	0.4	0.2	0.2	0.2	0.3	0.4	0.6	0.7	1.0
2002.....	4.6	1.5	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7	1.0
2001.....	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7	1.0
2000.....	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8	0.9
1999.....	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7	1.1

All origins 3/, female

2007.....	4.5	1.7	0.4	0.2	*	*	0.2	0.2	0.3	0.7	0.8
2006.....	4.7	1.6	0.5	*	0.2	0.3	0.2	0.4	0.4	0.5	0.8
2005.....	4.6	1.8	0.3	0.2	*	0.2	*	0.3	0.4	0.7	1.0
2004.....	4.6	1.8	0.3	0.2	*	0.2	0.2	0.3	0.5	0.6	0.7
2003.....	4.7	1.5	0.3	*	*	*	0.2	0.3	0.5	0.6	0.7
2002.....	4.6	2.0	0.4	0.2	*	0.2	*	0.3	0.5	0.5	0.9
2001.....	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7	0.8
2000.....	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6	1.0
1999.....	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 4/, both sexes

2007.....	5.3	1.2	0.3	0.2	0.2	0.1	0.2	0.4	0.4	0.7	1.0
2006.....	5.2	1.5	0.4	0.2	0.2	0.2	0.2	0.4	0.4	0.6	0.9
2005.....	5.2	1.3	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.7	1.1
2004.....	5.1	1.7	0.4	0.2	0.2	0.2	0.3	0.3	0.6	0.7	0.8
2003.....	5.2	1.5	0.4	0.2	0.2	0.2	0.2	0.4	0.5	0.7	0.9
2002.....	5.1	1.6	0.4	0.2	0.2	0.2	0.2	0.4	0.5	0.7	1.0
2001.....	5.3	1.4	0.4	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.9
2000.....	5.3	1.7	0.4	0.3	0.2	0.2	0.3	0.4	0.5	0.7	0.9
1999.....	5.2	1.6	0.4	0.3	0.2	0.3	0.3	0.3	0.6	0.7	0.9

Non-Hispanic 4/, male

2007.....	5.5	*	0.3	*	0.3	*	*	0.4	0.5	0.7	1.1
2006.....	5.3	1.6	0.3	*	*	0.3	*	0.3	0.4	0.7	0.9
2005.....	5.2	*	0.4	*	*	0.3	0.2	0.4	0.5	0.7	1.1
2004.....	5.2	1.5	0.4	*	0.3	*	0.4	0.4	0.6	0.7	0.9
2003.....	5.2	1.6	0.4	*	*	*	0.3	0.4	0.6	0.8	1.1
2002.....	5.1	1.3	0.4	*	*	*	0.3	0.4	0.5	0.8	1.1
2001.....	5.3	1.5	0.4	*	0.3	*	0.4	0.4	0.5	0.8	1.0
2000.....	5.3	1.8	0.5	0.3	*	0.3	0.4	0.4	0.5	0.8	0.9
1999.....	5.3	1.6	0.5	0.3	*	0.2	0.3	0.3	0.7	0.8	1.0

Non-Hispanic 4/, female

2007.....	5.0	1.5	0.3	*	*	*	*	0.3	0.3	0.7	0.9
2006.....	5.2	1.4	0.5	*	*	0.2	*	0.4	0.4	0.5	0.9
2005.....	5.1	1.8	*	*	*	*	*	0.3	0.5	0.7	1.0
2004.....	5.0	1.9	0.4	*	*	0.3	0.2	0.3	0.5	0.7	0.7
2003.....	5.1	1.3	*	*	*	*	*	0.3	0.5	0.6	0.7
2002.....	5.1	2.0	0.4	*	*	*	*	0.4	0.5	0.6	0.9
2001.....	5.2	1.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.7	0.9
2000.....	5.3	1.7	*	0.4	*	*	*	0.4	0.5	0.6	0.9
1999.....	5.1	1.7	0.4	0.3	*	0.3	*	0.3	0.5	0.6	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic White, both sexes

2007 5/.....	5.9	1.2	0.4	0.2	0.2	*	0.2	0.3	0.4	0.7	1.0
2006 5/.....	5.9	1.5	0.5	0.2	0.2	0.2	0.2	0.3	0.3	0.5	0.8
2005 5/.....	5.8	1.2	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.7	1.0
2004 5/.....	5.7	1.8	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7	0.7
2003 5/.....	5.8	1.4	0.3	*	0.2	0.2	0.2	0.4	0.5	0.6	0.9
2002.....	5.7	1.8	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.7	0.9
2001.....	5.9	1.3	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7	0.9
2000.....	5.9	1.8	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7	0.9
1999.....	5.7	1.5	0.4	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.8

Non-Hispanic White, male

2007 5/.....	6.2	*	*	*	*	*	*	0.5	0.4	0.7	1.1
2006 5/.....	6.0	*	*	*	*	*	*	*	0.4	0.7	0.9
2005 5/.....	5.9	*	0.4	*	*	0.3	*	0.4	0.5	0.8	1.1
2004 5/.....	5.9	*	0.4	*	*	*	0.4	0.4	0.6	0.6	0.9
2003 5/.....	5.9	1.7	*	*	*	*	*	0.4	0.5	0.8	1.1
2002.....	5.8	*	*	*	*	*	*	0.4	0.6	0.8	1.1
2001.....	6.0	*	*	*	*	*	0.3	0.4	0.5	0.8	0.9
2000.....	5.9	1.8	0.5	0.3	*	*	0.4	0.4	0.5	0.8	0.9
1999.....	5.9	*	*	*	*	*	*	*	0.6	0.7	1.0

Non-Hispanic White, female

2007 5/.....	5.7	*	*	*	*	*	*	*	*	0.7	0.8
2006 5/.....	5.8	*	0.6	*	*	*	*	0.4	*	0.4	0.7
2005 5/.....	5.7	*	*	*	*	*	*	*	0.5	0.6	1.0
2004 5/.....	5.6	2.0	*	*	*	*	*	*	0.4	0.7	0.6
2003 5/.....	5.8	*	*	*	*	*	*	0.4	0.6	0.5	0.7
2002.....	5.6	2.1	*	*	*	*	*	*	0.4	0.5	0.8
2001.....	5.8	*	*	*	*	*	*	0.3	0.4	0.7	0.8
2000.....	5.9	1.8	*	0.3	*	*	*	*	0.5	0.6	0.9
1999.....	5.6	*	*	0.3	*	*	*	*	0.4	0.5	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Anemias (D50-D64)

All origins 3/, both sexes

2007.....	1.6	*	*	*	*	0.2	0.3	0.3	0.3	0.4	0.5
2006.....	1.3	*	0.2	*	*	0.2	0.3	0.4	0.3	0.4	0.5
2005.....	1.6	*	0.1	*	*	0.2	0.3	0.4	0.4	0.4	0.5
2004.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.4	0.4	0.4	0.5
2003.....	1.6	0.5	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4	0.4
2002.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5	0.4
2001.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4	0.6
2000.....	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4	0.5
1999.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4	0.5

All origins 3/, male

2007.....	1.3	*	*	*	*	*	0.3	0.4	0.3	0.4	0.5
2006.....	1.1	*	*	*	*	0.2	0.3	0.4	0.3	0.3	0.5
2005.....	1.2	*	*	*	*	0.2	0.3	0.4	0.4	0.4	0.5
2004.....	1.3	*	*	*	*	0.3	0.4	0.4	0.5	0.5	0.5
2003.....	1.3	*	*	*	*	*	0.3	0.4	0.4	0.3	0.5
2002.....	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5	0.4
2001.....	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4	0.4
2000.....	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3	0.4
1999.....	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4	0.5

All origins 3/, female

2007.....	1.9	*	*	*	*	0.2	0.2	0.3	0.3	0.3	0.5
2006.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.5	0.5
2005.....	1.9	*	*	*	*	*	0.3	0.4	0.3	0.4	0.5
2004.....	1.8	*	*	*	*	*	0.4	0.5	0.3	0.3	0.5
2003.....	1.9	*	*	*	*	*	0.3	0.3	0.4	0.4	0.4
2002.....	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4	0.5
2001.....	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4	0.7
2000.....	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4	0.5
1999.....	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Anemias (D50-D64)

Non-Hispanic 4/, both sexes

2007.....	1.8	*	*	*	*	0.2	0.3	0.4	0.4	0.4	0.5
2006.....	1.5	*	0.2	*	*	0.2	0.3	0.4	0.4	0.4	0.5
2005.....	1.7	*	*	*	*	0.2	0.3	0.5	0.4	0.4	0.5
2004.....	1.7	*	0.2	0.1	0.1	0.3	0.4	0.5	0.5	0.4	0.5
2003.....	1.8	0.6	*	*	0.2	0.2	0.4	0.4	0.5	0.4	0.5
2002.....	1.8	*	*	0.2	0.1	0.2	0.4	0.5	0.5	0.5	0.4
2001.....	1.8	*	*	0.1	0.2	0.2	0.3	0.4	0.5	0.4	0.6
2000.....	1.8	*	0.3	0.1	0.1	0.2	0.3	0.3	0.5	0.4	0.5
1999.....	1.8	*	0.2	0.1	0.2	0.2	0.5	0.4	0.5	0.4	0.5

Non-Hispanic 4/, male

2007.....	1.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.5
2006.....	1.2	*	*	*	*	*	0.4	0.5	0.3	0.4	0.5
2005.....	1.4	*	*	*	*	*	0.4	0.5	0.5	0.4	0.6
2004.....	1.4	*	*	*	*	0.3	0.4	0.5	0.6	0.6	0.5
2003.....	1.4	*	*	*	*	*	0.4	0.5	0.5	0.4	0.5
2002.....	1.4	*	*	*	*	0.2	0.5	0.7	0.5	0.5	0.4
2001.....	1.4	*	*	*	*	0.3	0.4	0.5	0.6	0.4	0.5
2000.....	1.3	*	*	*	*	*	0.4	0.3	0.5	0.3	0.4
1999.....	1.4	*	*	*	*	0.3	0.5	0.4	0.5	0.4	0.5

Non-Hispanic 4/, female

2007.....	2.1	*	*	*	*	*	0.2	0.3	0.4	0.4	0.5
2006.....	1.7	*	*	*	*	*	0.3	0.4	0.4	0.5	0.5
2005.....	2.1	*	*	*	*	*	0.3	0.5	0.4	0.4	0.5
2004.....	2.0	*	*	*	*	*	0.4	0.5	0.3	0.3	0.6
2003.....	2.1	*	*	*	*	*	0.4	0.4	0.5	0.5	0.4
2002.....	2.1	*	*	*	*	*	0.3	0.4	0.5	0.5	0.5
2001.....	2.2	*	*	*	*	*	0.3	0.3	0.4	0.4	0.8
2000.....	2.2	*	*	*	*	*	0.3	0.3	0.5	0.5	0.6
1999.....	2.1	*	*	*	*	*	0.4	0.4	0.5	0.4	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Anemias (D50-D64)

Non-Hispanic Black, both sexes

2007 5/.....	2.5	*	*	*	*	0.8	1.3	1.8	2.0	2.3	2.8
2006 5/.....	2.4	*	*	*	*	0.7	1.4	2.0	2.1	2.3	3.0
2005 5/.....	2.7	*	*	*	*	0.9	1.6	2.4	2.6	2.3	2.9
2004 5/.....	2.7	*	*	*	*	0.9	2.0	2.3	2.5	2.5	2.5
2003 5/.....	2.8	*	*	*	*	*	2.0	2.1	2.3	2.2	2.6
2002.....	2.9	*	*	*	*	0.9	1.8	2.6	2.4	2.6	2.5
2001.....	2.8	*	*	*	*	0.9	1.6	2.0	2.5	2.2	3.3
2000.....	2.7	*	*	*	*	*	1.7	1.8	2.9	2.3	2.7
1999.....	2.9	*	0.9	*	*	0.8	2.4	2.1	3.2	2.3	2.8

Non-Hispanic Black, male

2007 5/.....	2.4	*	*	*	*	*	1.6	2.1	2.2	2.7	3.2
2006 5/.....	2.1	*	*	*	*	*	1.6	2.2	2.1	2.0	3.0
2005 5/.....	2.5	*	*	*	*	*	1.8	2.6	2.9	2.8	3.2
2004 5/.....	2.6	*	*	*	*	1.3	2.0	2.5	3.1	3.2	2.6
2003 5/.....	2.8	*	*	*	*	*	1.8	2.3	2.4	2.2	3.0
2002.....	2.7	*	*	*	*	*	2.3	3.3	2.6	2.7	2.4
2001.....	2.6	*	*	*	*	*	1.7	2.6	3.4	2.3	2.8
2000.....	2.4	*	*	*	*	*	1.7	1.6	2.8	2.0	2.3
1999.....	2.7	*	*	*	*	*	2.9	2.4	3.4	2.6	3.0

Non-Hispanic Black, female

2007 5/.....	2.6	*	*	*	*	*	*	1.5	1.8	2.0	2.5
2006 5/.....	2.6	*	*	*	*	*	*	1.9	2.2	2.5	3.0
2005 5/.....	2.9	*	*	*	*	*	1.3	2.2	2.3	1.9	2.7
2004 5/.....	2.8	*	*	*	*	*	2.1	2.1	2.0	1.9	2.5
2003 5/.....	2.9	*	*	*	*	*	2.2	1.8	2.3	2.3	2.1
2002.....	3.0	*	*	*	*	*	*	1.9	2.1	2.5	2.6
2001.....	2.9	*	*	*	*	*	1.5	*	1.8	2.1	3.8
2000.....	3.1	*	*	*	*	*	1.7	2.0	3.0	2.6	3.1
1999.....	3.1	*	*	*	*	*	1.8	1.9	2.9	2.1	2.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

All origins 3/, both sexes

2007.....	23.7	*	*	*	*	0.3	0.5	1.0	2.0	3.5	5.6
2006.....	24.2	*	*	*	0.1	0.2	0.6	1.2	2.1	3.3	6.2
2005.....	25.3	*	*	*	0.2	0.3	0.6	1.0	2.0	3.6	5.6
2004.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4	5.7
2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3	5.8
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5	6.0
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3	5.4
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0	5.5
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2	5.5

All origins 3/, male

2007.....	23.9	*	*	*	*	0.3	0.6	1.1	2.4	4.5	6.8
2006.....	24.4	*	*	*	*	0.2	0.7	1.3	2.4	4.0	7.6
2005.....	25.0	*	*	*	*	0.4	0.7	1.1	2.5	4.3	7.2
2004.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2	7.1
2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9	7.4
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1	7.7
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9	6.6
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5	6.8
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8	6.4

All origins 3/, female

2007.....	23.5	*	*	*	*	0.2	0.5	0.9	1.7	2.6	4.4
2006.....	24.0	*	*	*	*	*	0.5	1.1	1.9	2.7	4.7
2005.....	25.7	*	*	*	0.2	0.3	0.6	0.9	1.5	2.9	4.1
2004.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5	4.3
2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8	4.2
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0	4.4
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7	4.2
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5	4.3
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6	4.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Hispanic, both sexes

2007.....	14.1	*	*	*	*	*	*	0.5	1.0	2.7	4.1
2006.....	14.2	*	*	*	*	*	*	0.7	1.1	1.9	3.8
2005.....	15.6	*	*	*	*	*	*	0.7	1.1	2.0	5.0
2004.....	14.5	*	*	*	*	*	*	0.5	1.2	2.0	4.2
2003.....	15.5	*	*	*	*	*	*	0.7	1.0	1.9	5.0
2002.....	15.3	*	*	*	*	*	*	0.9	0.7	2.3	5.0
2001.....	15.3	*	*	*	*	*	*	0.6	0.9	2.3	4.0
2000.....	15.1	*	*	*	*	*	*	0.6	1.4	1.7	4.6
1999.....	15.3	*	*	*	*	*	*	*	1.0	2.1	4.6

Hispanic, male

2007.....	13.6	*	*	*	*	*	*	*	1.2	3.6	4.9
2006.....	13.7	*	*	*	*	*	*	*	1.1	2.2	4.7
2005.....	14.9	*	*	*	*	*	*	*	1.1	2.8	6.6
2004.....	13.4	*	*	*	*	*	*	*	1.3	2.2	4.9
2003.....	14.4	*	*	*	*	*	*	*	*	2.2	5.9
2002.....	13.9	*	*	*	*	*	*	*	*	2.5	6.1
2001.....	13.6	*	*	*	*	*	*	*	*	2.6	5.4
2000.....	13.8	*	*	*	*	*	*	*	1.9	1.9	5.7
1999.....	13.4	*	*	*	*	*	*	*	*	2.2	4.7

Hispanic, female

2007.....	14.6	*	*	*	*	*	*	*	*	1.7	3.2
2006.....	14.7	*	*	*	*	*	*	*	*	1.5	2.8
2005.....	16.3	*	*	*	*	*	*	*	*	*	3.2
2004.....	15.7	*	*	*	*	*	*	*	*	1.8	3.5
2003.....	16.6	*	*	*	*	*	*	*	*	1.5	4.1
2002.....	16.7	*	*	*	*	*	*	*	*	2.1	3.8
2001.....	17.1	*	*	*	*	*	*	*	*	2.0	2.4
2000.....	16.5	*	*	*	*	*	*	*	*	1.6	3.4
1999.....	17.3	*	*	*	*	*	*	*	*	1.9	4.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Non-Hispanic 4/, both sexes

2007.....	25.3	*	*	*	*	0.3	0.6	1.1	2.3	3.7	5.9
2006.....	25.9	*	*	*	*	0.2	0.6	1.3	2.4	3.6	6.5
2005.....	26.9	*	*	*	0.2	0.4	0.7	1.1	2.3	3.9	5.7
2004.....	26.5	*	*	*	0.2	0.3	0.6	1.0	2.2	3.6	5.9
2003.....	27.0	*	*	*	*	0.2	0.6	1.3	2.3	3.6	5.9
2002.....	26.9	*	*	*	0.2	0.3	0.6	1.3	2.2	3.7	6.1
2001.....	26.4	*	*	*	0.2	0.2	0.6	1.3	2.0	3.5	5.5
2000.....	25.9	*	*	*	*	0.3	0.7	1.2	2.1	3.2	5.6
1999.....	25.7	*	*	*	*	0.3	0.6	1.1	2.1	3.3	5.5

Non-Hispanic 4/, male

2007.....	25.8	*	*	*	*	0.3	0.7	1.3	2.7	4.6	7.2
2006.....	26.3	*	*	*	*	0.2	0.7	1.5	2.8	4.4	8.1
2005.....	26.8	*	*	*	*	0.4	0.8	1.3	2.9	4.6	7.2
2004.....	26.2	*	*	*	*	0.3	0.6	1.2	2.5	4.6	7.4
2003.....	26.4	*	*	*	*	0.3	0.7	1.4	2.8	4.2	7.6
2002.....	25.8	*	*	*	*	0.3	0.7	1.4	2.8	4.4	7.9
2001.....	25.0	*	*	*	*	0.3	0.6	1.4	2.4	4.1	6.7
2000.....	24.2	*	*	*	*	0.3	0.8	1.3	2.6	3.7	6.9
1999.....	24.1	*	*	*	*	0.3	0.6	1.0	2.4	4.1	6.6

Non-Hispanic 4/, female

2007.....	24.9	*	*	*	*	0.3	0.5	1.0	1.9	2.7	4.6
2006.....	25.5	*	*	*	*	*	0.6	1.1	2.1	2.9	5.0
2005.....	27.1	*	*	*	0.2	0.3	0.7	1.0	1.7	3.3	4.2
2004.....	26.8	*	*	*	*	*	0.6	0.8	2.0	2.6	4.4
2003.....	27.6	*	*	*	*	*	0.5	1.2	1.8	3.0	4.3
2002.....	27.9	*	*	*	*	0.3	0.5	1.2	1.7	3.1	4.5
2001.....	27.8	*	*	*	*	*	0.5	1.1	1.6	2.8	4.4
2000.....	27.6	*	*	*	*	*	0.6	1.0	1.7	2.7	4.4
1999.....	27.3	*	*	*	*	0.3	0.6	1.1	1.7	2.6	4.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Non-Hispanic White, both sexes

2007 5/.....	24.9	*	*	*	*	0.2	0.5	0.8	1.9	3.0	4.9
2006 5/.....	25.4	*	*	*	*	0.2	0.5	0.9	1.9	3.0	5.6
2005 5/.....	26.5	*	*	*	*	0.3	0.6	0.8	1.7	3.3	4.9
2004 5/.....	26.1	*	*	*	*	0.2	0.5	0.8	1.9	3.0	5.1
2003 5/.....	26.6	*	*	*	*	0.2	0.5	0.9	2.0	3.0	5.0
2002.....	26.4	*	*	*	*	0.2	0.5	1.0	2.0	3.2	5.3
2001.....	26.0	*	*	*	*	0.2	0.4	0.9	1.7	2.9	4.8
2000.....	25.4	*	*	*	*	0.2	0.5	0.9	1.8	2.8	5.0
1999.....	25.0	*	*	*	*	0.3	0.5	1.0	1.8	2.9	4.7

Non-Hispanic White, male

2007 5/.....	25.9	*	*	*	*	*	0.6	1.1	2.2	3.8	6.2
2006 5/.....	26.3	*	*	*	*	*	0.5	1.1	2.1	3.6	7.0
2005 5/.....	26.8	*	*	*	*	*	0.7	0.9	2.2	3.7	6.5
2004 5/.....	26.3	*	*	*	*	*	0.5	0.9	2.1	3.9	6.5
2003 5/.....	26.6	*	*	*	*	*	0.6	0.8	2.5	3.6	6.6
2002.....	26.0	*	*	*	*	*	0.5	1.2	2.4	3.9	6.9
2001.....	25.1	*	*	*	*	*	0.5	0.9	1.9	3.4	6.2
2000.....	24.3	*	*	*	*	*	0.7	1.0	2.3	3.2	6.3
1999.....	24.0	*	*	*	*	0.3	0.5	1.0	2.0	3.6	5.6

Non-Hispanic White, female

2007 5/.....	23.9	*	*	*	*	*	0.3	0.6	1.5	2.2	3.6
2006 5/.....	24.5	*	*	*	*	*	0.5	0.8	1.7	2.5	4.1
2005 5/.....	26.3	*	*	*	*	*	0.4	0.7	1.3	2.9	3.3
2004 5/.....	25.9	*	*	*	*	*	0.5	0.7	1.6	2.2	3.7
2003 5/.....	26.5	*	*	*	*	*	0.4	1.0	1.5	2.4	3.4
2002.....	26.8	*	*	*	*	*	0.5	0.9	1.5	2.6	3.7
2001.....	26.9	*	*	*	*	*	*	0.9	1.5	2.4	3.5
2000.....	26.5	*	*	*	*	*	0.4	0.8	1.3	2.4	3.7
1999.....	26.0	*	*	*	*	*	0.5	0.9	1.5	2.1	3.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Non-Hispanic Black, both sexes

2007 5/.....	32.4	*	*	*	*	0.6	1.1	2.9	5.1	7.8	12.6
2006 5/.....	33.6	*	*	*	*	*	1.3	3.5	5.5	8.2	13.0
2005 5/.....	34.4	*	*	*	*	0.7	1.6	3.0	5.3	8.4	11.1
2004 5/.....	34.4	*	*	*	*	*	1.4	2.1	4.7	7.7	11.8
2003 5/.....	35.0	*	*	*	*	*	1.0	3.4	4.6	7.7	12.1
2002.....	34.8	*	*	*	*	*	1.4	2.9	4.5	7.5	12.3
2001.....	34.1	*	*	*	*	*	1.4	3.3	4.4	7.2	11.0
2000.....	34.0	*	*	*	*	*	1.5	2.9	4.3	6.2	10.4
1999.....	34.1	*	*	*	*	*	1.2	1.9	4.2	6.8	11.6

Non-Hispanic Black, male

2007 5/.....	29.9	*	*	*	*	*	*	3.0	6.2	9.7	14.6
2006 5/.....	31.7	*	*	*	*	*	1.5	3.7	7.1	10.3	16.4
2005 5/.....	31.8	*	*	*	*	*	1.4	3.1	7.0	11.2	12.7
2004 5/.....	31.3	*	*	*	*	*	*	2.6	5.5	9.9	14.4
2003 5/.....	30.9	*	*	*	*	*	*	4.4	5.5	8.8	14.9
2002.....	30.0	*	*	*	*	*	1.8	3.1	6.0	8.7	15.4
2001.....	29.4	*	*	*	*	*	1.5	4.1	5.8	9.0	12.0
2000.....	28.3	*	*	*	*	*	1.6	3.3	4.7	7.4	12.0
1999.....	28.6	*	*	*	*	*	*	1.7	5.1	8.0	13.5

Non-Hispanic Black, female

2007 5/.....	34.6	*	*	*	*	*	1.5	2.8	4.2	6.2	10.8
2006 5/.....	35.3	*	*	*	*	*	*	3.3	4.1	6.3	9.9
2005 5/.....	36.7	*	*	*	*	*	1.8	2.8	3.8	5.8	9.7
2004 5/.....	37.2	*	*	*	*	*	1.7	1.7	4.0	5.8	9.6
2003 5/.....	38.7	*	*	*	*	*	*	2.4	3.9	6.6	9.6
2002.....	39.2	*	*	*	*	*	*	2.8	3.3	6.4	9.7
2001.....	38.4	*	*	*	*	*	*	2.6	3.0	5.6	10.1
2000.....	39.1	*	*	*	*	*	1.5	2.6	3.9	5.1	8.9
1999.....	39.0	*	*	*	*	*	*	2.0	3.4	5.6	9.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes

2007.....	267.3	13.4	1.4	0.8	0.9	2.1	4.4	7.2	12.4	23.4	45.1
2006.....	275.1	12.3	1.3	0.7	1.1	2.2	4.3	7.2	13.2	23.4	46.5
2005.....	288.8	12.2	1.3	0.8	1.0	2.3	4.3	7.3	12.8	24.1	47.2
2004.....	293.3	13.8	1.5	0.6	1.0	2.2	4.2	7.3	12.6	24.3	47.8
2003.....	310.3	13.9	1.5	0.7	1.0	2.5	4.5	7.1	13.2	25.1	50.3
2002.....	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3	49.7
2001.....	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0	48.8
2000.....	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2	49.6
1999.....	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9	50.0

All origins 3/, male

2007.....	261.0	14.8	1.2	0.8	0.9	2.6	5.4	9.1	16.0	31.2	61.8
2006.....	267.7	13.4	1.4	0.8	1.2	2.6	5.4	9.5	17.6	31.5	62.7
2005.....	277.9	13.4	1.5	0.8	1.1	3.0	5.5	9.5	16.6	32.0	64.1
2004.....	281.2	14.7	1.5	0.5	1.1	2.8	5.3	9.2	16.3	32.1	64.6
2003.....	296.3	15.2	1.4	0.7	1.2	3.1	5.3	8.5	17.0	33.4	67.7
2002.....	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9	67.9
2001.....	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1	66.5
2000.....	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3	68.0
1999.....	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2	68.9

All origins 3/, female

2007.....	273.3	11.9	1.6	0.7	0.9	1.5	3.3	5.3	8.7	15.5	28.4
2006.....	282.4	11.2	1.3	0.6	0.9	1.7	3.1	4.8	8.8	15.3	30.5
2005.....	299.3	11.0	1.1	0.9	0.8	1.7	3.0	5.0	9.0	16.1	30.5
2004.....	304.9	12.9	1.5	0.7	1.0	1.6	3.0	5.3	8.9	16.4	31.2
2003.....	323.9	12.5	1.5	0.6	0.8	2.0	3.6	5.6	9.3	16.8	33.2
2002.....	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7	31.6
2001.....	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0	31.5
2000.....	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1	31.5
1999.....	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6	31.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Major cardiovascular diseases (I00-I78)

Hispanic, both sexes

2007.....	85.0	11.9	1.2	0.7	0.7	1.7	3.6	5.2	7.4	14.2	25.9
2006.....	86.5	10.2	1.1	0.6	1.0	2.1	3.4	4.8	7.8	13.0	27.7
2005.....	91.4	10.5	1.1	0.6	0.9	2.0	3.5	5.0	6.9	13.8	29.0
2004.....	89.3	11.5	1.2	0.5	0.7	1.7	3.2	4.1	7.2	14.2	29.1
2003.....	93.4	11.0	1.3	*	0.9	1.9	3.0	4.8	8.0	13.5	28.6
2002.....	94.3	13.1	1.2	*	1.0	1.8	3.3	4.7	7.5	13.8	28.2
2001.....	96.2	12.0	1.5	0.7	0.9	2.1	2.8	4.4	8.6	14.1	26.7
2000.....	96.5	13.7	1.3	0.6	0.8	1.9	3.2	4.8	8.2	14.5	29.7
1999.....	99.2	15.2	1.4	0.6	0.8	2.2	2.9	4.6	8.5	15.6	28.6

Hispanic, male

2007.....	86.0	11.9	1.1	*	*	1.9	4.9	6.6	9.5	18.9	37.7
2006.....	87.1	11.1	1.3	*	1.2	2.7	4.4	6.5	10.7	18.0	38.1
2005.....	92.3	11.3	1.5	*	1.1	2.6	4.0	6.6	9.4	18.8	40.2
2004.....	90.6	11.7	1.6	*	*	2.5	4.3	5.7	10.0	19.6	40.4
2003.....	92.6	12.2	1.5	*	*	2.4	4.0	5.7	10.2	18.3	39.2
2002.....	94.2	12.2	*	*	*	2.2	4.3	6.5	9.8	17.4	39.5
2001.....	95.4	12.7	1.4	*	*	2.4	3.5	5.6	11.2	19.0	36.5
2000.....	96.0	13.7	1.4	*	1.2	2.4	3.7	6.0	10.4	18.2	40.0
1999.....	99.2	12.7	1.4	*	*	2.5	3.5	6.1	11.4	20.8	40.8

Hispanic, female

2007.....	84.0	11.9	1.4	*	*	1.5	2.1	3.4	5.0	8.9	12.8
2006.....	85.9	9.3	*	*	*	1.4	2.2	2.7	4.4	7.3	16.3
2005.....	90.4	9.7	*	*	*	1.3	2.9	3.0	4.1	8.2	16.7
2004.....	87.9	11.3	*	*	*	*	1.9	2.1	3.8	8.0	16.8
2003.....	94.3	9.7	*	*	*	1.3	1.9	3.8	5.5	8.0	17.2
2002.....	94.4	14.0	1.3	*	1.2	1.4	2.0	2.5	4.9	9.8	16.2
2001.....	97.1	11.2	1.6	*	*	1.7	1.8	3.0	5.6	8.7	16.5
2000.....	96.9	13.8	*	*	*	*	2.6	3.4	5.7	10.5	18.9
1999.....	99.1	17.9	1.4	*	*	1.8	2.1	2.8	5.2	10.0	15.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Major cardiovascular diseases (I00-I78)

Non-Hispanic 4/, both sexes

2007.....	299.2	13.8	1.5	0.8	1.0	2.2	4.5	7.7	13.7	25.3	48.2
2006.....	307.3	12.9	1.4	0.7	1.1	2.2	4.5	7.8	14.5	25.4	49.4
2005.....	321.4	12.8	1.4	0.9	1.0	2.4	4.5	7.9	14.2	26.0	49.9
2004.....	325.9	14.4	1.5	0.6	1.1	2.3	4.4	8.0	13.8	26.1	50.3
2003.....	343.8	14.5	1.5	0.7	1.0	2.6	4.8	7.6	14.2	27.0	53.0
2002.....	352.2	16.5	1.5	0.7	1.1	2.5	4.0	7.7	13.6	27.1	52.2
2001.....	356.7	15.5	1.9	0.7	1.1	2.2	4.5	7.7	13.8	26.6	51.3
2000.....	365.6	17.3	1.6	0.7	1.1	2.5	4.5	7.0	12.8	25.4	51.5
1999.....	372.7	17.4	1.5	0.7	1.1	2.8	4.4	6.8	13.3	27.1	52.0

Non-Hispanic 4/, male

2007.....	293.3	15.6	1.3	0.8	1.0	2.8	5.5	9.8	17.8	33.9	66.0
2006.....	300.1	14.1	1.4	0.8	1.2	2.6	5.7	10.4	19.4	34.2	66.7
2005.....	310.3	14.0	1.5	0.8	1.1	3.0	5.8	10.2	18.4	34.6	67.7
2004.....	313.4	15.5	1.4	0.6	1.1	2.8	5.5	10.1	17.8	34.5	68.0
2003.....	329.5	15.9	1.4	0.8	1.2	3.2	5.7	9.2	18.5	36.1	71.3
2002.....	336.4	17.6	1.7	0.6	1.4	3.3	5.1	9.7	17.4	35.5	71.5
2001.....	338.1	15.8	2.0	0.6	1.2	2.7	5.6	9.5	17.4	35.2	69.8
2000.....	347.7	18.3	1.7	0.8	1.2	2.9	5.4	8.6	16.2	33.0	70.8
1999.....	354.2	19.1	1.7	0.8	1.1	3.5	5.1	8.1	17.0	35.9	71.6

Non-Hispanic 4/, female

2007.....	304.8	11.8	1.6	0.7	0.9	1.5	3.6	5.7	9.6	16.7	30.9
2006.....	314.1	11.7	1.4	0.7	1.0	1.7	3.3	5.2	9.7	16.7	32.5
2005.....	332.0	11.4	1.3	0.9	0.8	1.7	3.0	5.5	10.0	17.5	32.4
2004.....	337.9	13.2	1.6	0.7	1.0	1.7	3.2	6.0	9.9	17.8	33.0
2003.....	357.5	13.1	1.6	0.7	0.8	2.1	3.9	5.9	10.1	18.1	35.1
2002.....	367.2	15.3	1.3	0.8	0.9	1.7	2.9	5.7	9.8	18.9	33.4
2001.....	374.3	15.2	1.8	0.8	1.1	1.7	3.4	6.0	10.1	18.2	33.1
2000.....	382.6	16.3	1.5	0.6	1.1	2.1	3.7	5.5	9.5	18.0	32.7
1999.....	390.2	15.5	1.4	0.6	1.0	2.1	3.7	5.6	9.7	18.5	33.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Major cardiovascular diseases (I00-I78)

Non-Hispanic White, both sexes

2007 5/.....	324.4	10.4	1.2	0.7	0.8	1.8	3.6	6.1	11.0	21.2	41.9
2006 5/.....	333.1	9.5	1.2	0.6	1.0	1.9	3.9	6.2	11.7	21.0	43.4
2005 5/.....	348.4	9.8	1.1	0.7	0.8	2.0	3.6	6.1	11.4	21.7	43.2
2004 5/.....	353.1	12.0	1.2	0.5	0.9	1.8	3.7	6.2	11.3	22.4	43.7
2003 5/.....	372.4	10.9	1.2	0.6	0.9	2.2	3.8	5.9	11.7	22.8	45.4
2002.....	381.9	13.4	1.4	0.6	0.9	2.2	3.2	6.0	10.8	22.9	45.6
2001.....	385.8	12.1	1.5	0.6	1.0	1.9	3.5	6.2	11.3	22.5	43.3
2000.....	395.5	14.6	1.4	0.6	0.9	2.1	3.5	5.6	10.4	21.4	43.9
1999.....	402.7	14.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.8	43.8

Non-Hispanic White, male

2007 5/.....	315.2	11.7	0.9	0.7	0.9	2.2	4.1	7.7	14.5	28.8	58.3
2006 5/.....	322.3	10.2	1.1	0.7	1.2	2.3	5.1	8.4	16.0	28.8	60.7
2005 5/.....	333.6	10.7	1.3	0.7	1.0	2.4	4.8	8.2	15.0	29.4	60.8
2004 5/.....	336.9	13.0	1.1	0.5	1.0	2.3	4.7	8.1	14.8	30.0	61.8
2003 5/.....	354.1	11.3	0.9	0.6	1.1	2.7	4.5	7.3	15.5	31.4	63.5
2002.....	362.2	14.8	1.6	0.6	1.1	2.9	4.2	7.7	14.1	30.6	64.9
2001.....	363.0	13.0	1.6	0.6	1.1	2.2	4.5	7.8	14.6	30.4	61.8
2000.....	373.6	15.5	1.6	0.7	1.1	2.4	4.3	7.0	13.7	28.9	62.9
1999.....	379.8	16.1	1.5	0.8	0.9	2.9	4.0	6.5	14.3	30.9	62.8

Non-Hispanic White, female

2007 5/.....	333.4	9.0	1.4	0.7	0.7	1.3	3.1	4.4	7.4	13.5	25.6
2006 5/.....	343.6	8.8	1.3	0.6	0.7	1.4	2.7	4.0	7.3	13.2	26.1
2005 5/.....	362.8	8.9	0.9	0.7	0.7	1.5	2.3	3.9	7.8	13.9	25.7
2004 5/.....	368.7	10.9	1.2	0.5	0.9	1.4	2.6	4.3	7.7	14.8	25.6
2003 5/.....	390.0	10.4	1.4	0.6	0.7	1.8	3.0	4.4	7.8	14.1	27.4
2002.....	400.8	12.0	1.1	0.7	0.7	1.5	2.2	4.2	7.5	15.2	26.4
2001.....	407.6	11.0	1.5	0.7	0.9	1.5	2.5	4.6	8.0	14.6	24.8
2000.....	416.5	13.6	1.1	0.4	0.8	1.7	2.8	4.2	7.0	13.8	25.0
1999.....	424.7	12.1	1.1	0.6	0.9	1.9	3.1	4.4	7.4	14.6	24.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Major cardiovascular diseases (I00-I78)

Non-Hispanic Black, both sexes

2007 5/.....	249.2	25.9	2.3	1.0	1.5	4.1	9.0	15.9	29.8	52.8	92.1
2006 5/.....	255.2	26.3	2.2	1.0	1.7	3.8	7.9	16.3	32.1	55.0	93.2
2005 5/.....	265.1	23.5	2.4	1.6	1.6	4.4	8.7	17.4	30.9	54.8	96.9
2004 5/.....	269.2	22.4	3.0	1.2	1.9	4.3	7.9	17.6	29.9	51.7	100.4
2003 5/.....	282.2	26.6	2.9	1.3	1.7	4.5	9.8	16.9	30.3	54.7	107.0
2002.....	285.1	28.3	2.1	0.8	2.0	4.0	8.1	17.4	30.7	54.7	101.5
2001.....	289.7	30.1	3.7	1.0	1.6	4.1	9.8	17.1	29.5	53.3	108.4
2000.....	294.2	28.8	2.5	1.0	1.9	4.8	9.5	15.7	28.8	52.8	106.7
1999.....	298.5	30.2	2.6	1.1	1.9	5.0	9.3	15.1	28.0	56.8	112.2

Non-Hispanic Black, male

2007 5/.....	253.7	31.5	2.5	*	1.3	5.6	11.9	19.8	37.9	70.3	122.9
2006 5/.....	259.6	29.6	2.7	*	1.6	4.4	9.0	21.5	41.5	72.7	115.5
2005 5/.....	264.0	26.8	2.4	1.5	1.7	5.9	11.1	21.6	39.6	71.8	121.7
2004 5/.....	266.0	24.8	2.9	*	1.9	5.7	10.1	21.2	36.7	68.0	122.0
2003 5/.....	277.5	31.7	3.3	1.5	1.7	5.8	11.7	20.6	37.4	68.5	133.4
2002.....	277.2	29.1	2.0	*	2.4	5.1	9.7	21.6	38.2	68.8	126.0
2001.....	281.2	29.7	3.8	*	1.6	5.2	11.7	21.0	36.4	67.9	132.7
2000.....	284.3	30.0	2.3	*	1.9	5.3	10.8	18.9	33.9	63.5	133.3
1999.....	291.1	31.5	2.7	*	2.2	6.5	11.2	18.6	34.5	71.6	141.1

Non-Hispanic Black, female

2007 5/.....	245.1	20.1	2.1	*	1.7	2.5	6.1	12.1	22.4	37.2	64.9
2006 5/.....	251.3	22.8	1.8	*	1.8	3.2	6.7	11.4	23.7	39.2	73.6
2005 5/.....	266.1	20.1	2.4	1.8	1.5	2.8	6.3	13.4	23.0	39.5	75.1
2004 5/.....	272.2	20.0	3.0	1.4	1.9	2.8	5.8	14.3	23.7	37.1	81.5
2003 5/.....	286.5	21.3	2.4	*	1.6	3.2	8.0	13.5	23.8	42.5	83.8
2002.....	292.3	27.4	2.2	*	1.6	2.9	6.6	13.5	24.0	42.2	80.0
2001.....	297.4	30.6	3.6	1.3	1.5	2.9	7.9	13.6	23.3	40.4	86.9
2000.....	303.1	27.5	2.9	*	2.0	4.4	8.4	12.9	24.2	43.3	83.2
1999.....	305.1	28.9	2.4	*	1.6	3.3	7.4	11.9	22.2	43.8	86.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes

2007.....	204.3	10.0	1.1	0.6	0.6	1.6	3.5	6.0	10.0	18.4	36.1
2006.....	211.0	8.4	1.0	0.5	0.8	1.7	3.4	5.8	10.6	18.6	37.3
2005.....	220.0	8.7	0.9	0.5	0.7	1.8	3.5	5.9	10.3	19.4	37.7
2004.....	222.2	10.3	1.2	0.4	0.8	1.8	3.2	5.7	10.0	19.4	38.3
2003.....	235.6	11.0	1.2	0.5	0.8	1.9	3.6	5.6	10.5	20.1	40.5
2002.....	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3	40.2
2001.....	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9	39.1
2000.....	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1	39.4
1999.....	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6	40.1

All origins 3/, male

2007.....	208.4	10.9	1.0	0.6	0.7	2.1	4.3	7.9	13.4	25.4	51.5
2006.....	214.0	8.8	1.1	0.5	0.9	2.2	4.4	7.9	14.7	25.9	52.4
2005.....	221.1	9.4	1.0	0.5	0.8	2.5	4.6	7.9	13.7	26.8	53.6
2004.....	222.8	10.9	1.1	0.3	0.8	2.3	4.2	7.4	13.6	26.5	54.2
2003.....	235.0	12.1	1.1	0.6	0.9	2.4	4.4	6.9	13.9	27.8	56.9
2002.....	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8	57.7
2001.....	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4	55.8
2000.....	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0	57.0
1999.....	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5	58.1

All origins 3/, female

2007.....	200.2	9.0	1.1	0.5	0.6	1.1	2.6	4.1	6.5	11.3	20.9
2006.....	208.0	7.9	0.9	0.4	0.7	1.2	2.4	3.7	6.5	11.3	22.4
2005.....	218.9	8.0	0.9	0.6	0.6	1.2	2.2	3.7	6.8	12.0	21.8
2004.....	221.6	9.7	1.2	0.5	0.7	1.2	2.2	3.9	6.4	12.3	22.7
2003.....	236.2	9.8	1.3	0.5	0.6	1.4	2.7	4.2	7.1	12.5	24.3
2002.....	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8	22.9
2001.....	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4	22.6
2000.....	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2	22.2
1999.....	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6	22.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

Hispanic, both sexes

2007.....	63.8	8.8	1.0	0.5	*	1.3	2.7	4.0	5.2	10.1	18.5
2006.....	65.3	6.5	0.8	*	0.7	1.3	2.2	3.4	5.9	9.4	19.1
2005.....	69.2	7.6	0.6	*	0.6	1.3	2.4	4.0	5.0	9.7	20.9
2004.....	67.2	8.9	1.1	*	*	1.3	2.3	2.8	5.2	9.9	21.1
2003.....	70.9	8.7	1.1	*	0.6	1.4	2.4	3.5	6.0	10.3	20.1
2002.....	71.9	10.2	0.8	*	*	1.2	2.5	3.4	5.5	9.9	20.3
2001.....	73.3	9.7	1.1	*	0.6	1.4	1.9	3.2	6.0	10.1	18.1
2000.....	73.1	11.2	1.0	*	0.7	1.5	2.2	3.3	5.4	10.1	20.7
1999.....	76.2	12.7	1.2	*	*	1.7	2.0	2.9	6.3	11.7	20.0

Hispanic, male

2007.....	66.6	9.3	*	*	*	1.6	3.7	5.3	7.1	13.8	28.2
2006.....	67.7	7.1	*	*	*	1.9	3.1	4.8	8.4	13.3	28.0
2005.....	72.1	8.0	*	*	*	2.0	2.7	5.5	7.2	14.0	30.6
2004.....	70.2	8.9	1.4	*	*	1.9	3.3	4.1	7.7	14.1	31.3
2003.....	72.2	9.5	*	*	*	1.8	3.4	4.4	7.9	14.5	28.9
2002.....	74.0	9.6	*	*	*	1.6	3.4	4.8	7.5	12.9	30.0
2001.....	74.6	10.3	*	*	*	1.6	2.5	4.4	8.0	14.6	26.6
2000.....	74.7	11.2	1.4	*	*	2.1	2.7	4.3	7.2	13.4	29.9
1999.....	77.7	10.3	1.4	*	*	1.9	2.4	4.4	8.7	16.5	30.0

Hispanic, female

2007.....	60.8	8.3	1.2	*	*	*	1.4	2.5	3.0	5.9	7.8
2006.....	62.6	5.9	*	*	*	*	*	1.7	2.8	5.0	9.4
2005.....	66.2	7.2	*	*	*	*	2.1	2.2	2.5	4.8	10.3
2004.....	64.1	9.0	*	*	*	*	*	1.2	2.2	5.2	10.0
2003.....	69.6	7.8	*	*	*	*	*	2.4	3.7	5.6	10.7
2002.....	69.7	10.8	*	*	*	*	1.3	1.7	3.1	6.6	10.0
2001.....	71.8	9.0	*	*	*	*	*	1.9	3.7	5.2	9.2
2000.....	71.5	11.2	*	*	*	*	1.6	2.1	3.3	6.4	11.0
1999.....	74.6	15.2	*	*	*	1.5	1.5	*	3.6	6.5	9.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic 4/, both sexes

2007.....	228.8	10.2	1.1	0.6	0.7	1.7	3.7	6.5	11.2	20.1	39.0
2006.....	235.8	8.9	1.0	0.5	0.8	1.8	3.7	6.4	11.8	20.4	40.1
2005.....	244.9	9.0	1.0	0.6	0.7	2.0	3.7	6.3	11.5	21.2	40.1
2004.....	247.0	10.7	1.2	0.4	0.8	1.8	3.4	6.4	11.1	21.1	40.6
2003.....	261.0	11.5	1.2	0.6	0.8	2.0	3.8	6.1	11.4	21.8	43.0
2002.....	267.1	12.9	1.1	0.5	0.8	2.1	3.2	6.2	10.8	21.9	42.6
2001.....	270.7	12.3	1.5	0.5	0.9	1.7	3.6	6.1	10.9	21.3	41.4
2000.....	277.3	13.3	1.2	0.5	0.8	2.1	3.5	5.6	10.2	20.3	41.3
1999.....	284.3	14.0	1.2	0.6	0.8	2.4	3.5	5.5	10.7	21.6	42.1

Non-Hispanic 4/, male

2007.....	234.6	11.3	1.0	0.6	0.7	2.2	4.5	8.5	15.2	27.9	55.5
2006.....	240.3	9.4	1.1	0.5	0.9	2.2	4.6	8.7	16.3	28.5	56.4
2005.....	247.1	9.8	1.0	0.6	0.7	2.6	5.1	8.6	15.3	29.3	57.2
2004.....	248.5	11.5	1.1	0.3	0.9	2.3	4.4	8.2	15.0	28.8	57.4
2003.....	261.4	12.8	1.1	0.6	0.9	2.5	4.6	7.6	15.2	30.1	60.5
2002.....	267.0	13.7	1.2	0.4	1.0	2.8	4.1	8.1	14.6	30.3	61.3
2001.....	268.0	12.0	1.7	0.4	0.9	2.2	4.6	7.6	14.4	29.3	59.2
2000.....	275.2	13.7	1.4	0.6	0.9	2.4	4.3	7.0	13.4	27.7	59.7
1999.....	282.0	14.8	1.3	0.6	0.9	3.0	4.2	6.6	14.0	30.0	60.9

Non-Hispanic 4/, female

2007.....	223.3	9.1	1.1	0.5	0.7	1.1	2.9	4.5	7.3	12.4	22.9
2006.....	231.4	8.4	1.0	0.4	0.7	1.3	2.6	4.2	7.3	12.5	24.2
2005.....	242.8	8.3	1.0	0.6	0.7	1.3	2.3	4.1	7.7	13.2	23.4
2004.....	245.5	9.8	1.3	0.5	0.8	1.4	2.4	4.5	7.2	13.4	24.3
2003.....	260.5	10.2	1.3	0.6	0.6	1.5	3.0	4.6	7.7	13.6	25.9
2002.....	267.2	12.0	1.1	0.6	0.6	1.4	2.2	4.3	7.1	13.7	24.4
2001.....	273.3	12.7	1.4	0.6	0.8	1.3	2.6	4.5	7.5	13.4	24.2
2000.....	279.3	12.8	1.1	0.4	0.7	1.7	2.7	4.2	7.0	12.9	23.3
1999.....	286.4	13.3	1.1	0.5	0.8	1.7	2.8	4.3	7.4	13.4	23.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic White, both sexes

2007 5/.....	249.9	7.2	0.8	0.5	0.6	1.4	2.9	5.1	9.2	17.3	35.0
2006 5/.....	257.4	6.5	0.9	0.4	0.7	1.5	3.2	5.2	9.7	17.4	36.8
2005 5/.....	267.1	7.0	0.8	0.5	0.6	1.6	3.0	5.0	9.4	18.2	36.1
2004 5/.....	269.1	9.1	0.9	0.3	0.7	1.5	2.8	4.9	9.2	18.6	36.4
2003 5/.....	284.0	8.7	0.9	0.5	0.7	1.7	3.0	4.8	9.5	19.1	38.1
2002.....	290.8	10.7	1.0	0.4	0.7	1.8	2.4	4.8	8.7	19.2	38.5
2001.....	294.1	9.8	1.2	0.4	0.8	1.4	2.7	4.8	9.0	18.4	36.2
2000.....	301.3	11.4	1.1	0.4	0.7	1.7	2.7	4.3	8.3	17.6	36.5
1999.....	308.2	11.3	1.0	0.5	0.7	2.1	2.7	4.3	8.8	18.6	36.5

Non-Hispanic White, male

2007 5/.....	254.3	8.1	0.6	0.5	0.6	1.7	3.4	6.6	12.8	24.2	50.2
2006 5/.....	260.3	6.8	0.9	0.4	0.8	1.9	4.2	7.1	13.7	24.8	53.2
2005 5/.....	267.8	7.5	0.8	0.5	0.6	2.1	4.2	7.1	12.6	25.6	53.0
2004 5/.....	269.1	9.8	0.8	*	0.8	1.8	3.7	6.5	12.7	25.8	53.5
2003 5/.....	282.9	9.4	0.7	0.5	0.9	2.0	3.5	6.1	13.1	27.2	55.5
2002.....	289.2	11.6	1.1	0.4	0.9	2.4	3.2	6.4	12.0	26.9	57.2
2001.....	289.5	10.3	1.3	0.4	0.8	1.7	3.5	6.3	12.1	26.0	53.9
2000.....	297.5	12.0	1.2	0.5	0.8	1.9	3.3	5.5	11.4	24.9	54.8
1999.....	304.0	12.4	1.1	0.5	0.7	2.5	3.2	5.3	11.9	26.4	54.9

Non-Hispanic White, female

2007 5/.....	245.5	6.4	1.0	0.5	0.5	1.0	2.5	3.5	5.7	10.2	19.8
2006 5/.....	254.7	6.1	0.9	0.4	0.5	1.0	2.1	3.2	5.6	9.9	20.4
2005 5/.....	266.4	6.5	0.7	0.4	0.5	1.2	1.7	2.8	6.1	10.6	19.2
2004 5/.....	269.1	8.3	1.0	0.4	0.6	1.1	1.9	3.3	5.7	11.3	19.3
2003 5/.....	285.1	8.0	1.1	0.5	0.5	1.3	2.5	3.4	5.9	11.0	20.8
2002.....	292.3	9.6	0.8	0.5	0.5	1.2	1.6	3.3	5.4	11.4	19.8
2001.....	298.4	9.2	1.1	0.5	0.7	1.1	1.9	3.4	5.9	10.8	18.5
2000.....	304.9	10.8	0.9	0.3	0.5	1.4	2.0	3.1	5.2	10.2	18.2
1999.....	312.3	10.2	0.8	0.5	0.7	1.6	2.2	3.4	5.7	10.7	18.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic Black, both sexes

2007 5/.....	184.7	20.5	1.7	0.8	1.3	3.3	7.6	13.7	23.4	40.3	68.9
2006 5/.....	189.3	17.9	1.8	*	1.4	3.3	6.4	13.3	25.5	42.4	68.5
2005 5/.....	196.3	16.5	1.8	1.2	1.2	3.6	7.2	13.7	24.7	43.3	71.8
2004 5/.....	198.7	15.6	2.5	0.9	1.5	3.7	6.5	13.9	23.1	39.6	76.0
2003 5/.....	209.4	20.4	2.3	1.1	1.3	3.5	8.1	13.6	23.7	41.2	81.1
2002.....	212.2	21.9	1.7	*	1.4	3.5	6.7	14.1	24.1	42.1	76.5
2001.....	215.5	22.9	3.1	0.8	1.3	3.5	7.8	13.5	23.5	41.3	81.6
2000.....	218.6	20.5	2.0	0.9	1.4	4.2	7.8	13.4	22.3	39.4	78.8
1999.....	223.8	24.8	2.1	1.0	1.4	4.3	8.0	12.3	22.5	43.8	84.7

Non-Hispanic Black, male

2007 5/.....	193.7	23.5	2.2	*	*	4.6	10.0	17.9	30.9	55.1	96.4
2006 5/.....	198.8	18.8	2.1	*	1.5	3.9	7.5	17.7	34.2	57.0	88.2
2005 5/.....	201.5	19.0	*	*	1.3	5.2	9.4	17.3	32.4	58.3	94.3
2004 5/.....	203.2	17.0	2.5	*	1.5	4.8	8.3	17.4	29.6	52.9	96.2
2003 5/.....	212.6	24.5	2.5	*	1.4	4.6	10.4	16.6	29.5	53.3	105.4
2002.....	212.9	23.0	*	*	1.6	4.4	8.3	18.5	31.1	56.0	99.1
2001.....	215.7	20.7	3.4	*	1.4	4.5	10.0	16.7	30.3	54.1	104.2
2000.....	217.6	20.1	2.0	*	1.5	4.9	9.2	16.4	27.6	50.0	102.5
1999.....	224.9	24.6	2.3	*	1.7	6.0	9.7	15.6	28.4	57.9	111.5

Non-Hispanic Black, female

2007 5/.....	176.4	17.3	*	*	1.4	1.9	5.2	9.6	16.7	27.1	44.7
2006 5/.....	180.6	17.0	*	*	*	2.6	5.3	9.2	17.6	29.5	51.1
2005 5/.....	191.6	14.0	2.1	1.4	1.2	1.9	4.9	10.2	17.7	29.9	52.0
2004 5/.....	194.6	14.2	2.4	*	1.6	2.5	4.7	10.7	17.3	27.8	58.2
2003 5/.....	206.5	16.1	2.0	*	*	2.4	5.7	10.8	18.6	30.3	59.6
2002.....	211.6	20.8	1.9	*	*	2.5	5.3	10.1	17.7	29.7	56.6
2001.....	215.4	25.2	2.9	*	*	2.4	5.8	10.7	17.5	30.0	61.6
2000.....	219.5	20.8	2.0	*	1.4	3.4	6.5	10.6	17.5	30.0	58.0
1999.....	222.8	25.0	1.9	*	*	2.6	6.5	9.4	17.2	31.4	61.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins 3/, both sexes

2007.....	1.1	*	*	*	*	*	*	*	*	0.1	0.2
2006.....	1.1	*	*	*	*	*	*	*	0.1	0.1	0.2
2005.....	1.1	*	*	*	*	*	*	*	*	0.1	0.2
2004.....	1.1	*	*	*	*	*	*	*	*	0.2	0.3
2003.....	1.2	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.2	*	*	*	*	*	*	*	0.1	0.2	0.2
2001.....	1.2	*	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	1.3	*	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	1.3	*	*	*	*	*	*	*	0.1	0.2	0.3

All origins 3/, male

2007.....	0.7	*	*	*	*	*	*	*	*	*	*
2006.....	0.7	*	*	*	*	*	*	*	*	*	*
2005.....	0.7	*	*	*	*	*	*	*	*	*	0.2
2004.....	0.7	*	*	*	*	*	*	*	*	*	0.2
2003.....	0.8	*	*	*	*	*	*	*	*	*	0.2
2002.....	0.8	*	*	*	*	*	*	*	*	*	0.2
2001.....	0.7	*	*	*	*	*	*	*	*	*	0.3
2000.....	0.8	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	0.8	*	*	*	*	*	*	*	*	*	0.3

All origins 3/, female

2007.....	1.4	*	*	*	*	*	*	*	*	*	0.3
2006.....	1.5	*	*	*	*	*	*	*	*	*	0.2
2005.....	1.5	*	*	*	*	*	*	*	*	*	0.3
2004.....	1.5	*	*	*	*	*	*	*	*	*	0.3
2003.....	1.7	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.7	*	*	*	*	*	*	*	*	*	0.3
2001.....	1.7	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.8	*	*	*	*	*	*	*	*	0.2	0.5
1999.....	1.9	*	*	*	*	*	*	*	*	0.2	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hypertensive heart disease (I11)

All origins 3/, both sexes

2007.....	10.2	*	*	*	*	*	0.2	0.5	1.1	2.1	4.2
2006.....	9.9	*	*	*	*	*	0.2	0.5	0.9	2.2	4.0
2005.....	9.9	*	*	*	*	*	0.2	0.6	1.1	2.2	4.0
2004.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9	4.0
2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9	3.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8	3.5
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6	3.2
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4	2.9
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5	2.8

All origins 3/, male

2007.....	9.6	*	*	*	*	*	0.3	0.8	1.6	3.0	5.8
2006.....	9.3	*	*	*	*	*	0.3	0.7	1.3	3.2	5.2
2005.....	9.1	*	*	*	*	*	0.2	0.7	1.4	3.2	5.4
2004.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8	5.2
2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7	5.2
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6	4.9
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1	4.5
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9	3.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0	3.9

All origins 3/, female

2007.....	10.8	*	*	*	*	*	*	0.3	0.7	1.2	2.7
2006.....	10.6	*	*	*	*	*	*	0.3	0.5	1.1	2.8
2005.....	10.7	*	*	*	*	*	*	0.4	0.7	1.2	2.5
2004.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0	2.7
2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2	2.6
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0	2.2
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1	2.0
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0	2.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hypertensive heart disease (I11)

Non-Hispanic 4/, both sexes

2007	11.3	*	*	*	*	*	0.2	0.6	1.3	2.3	4.5
2006	11.0	*	*	*	*	*	0.2	0.6	1.0	2.3	4.2
2005	10.9	*	*	*	*	*	0.2	0.6	1.2	2.4	4.2
2004	10.7	*	*	*	*	*	0.1	0.6	1.1	2.0	4.2
2003	10.6	*	*	*	*	*	0.2	0.5	1.2	2.0	4.1
2002	10.0	*	*	*	*	*	0.1	0.5	1.1	1.9	3.7
2001	9.4	*	*	*	*	*	*	0.4	1.0	1.6	3.4
2000	9.2	*	*	*	*	*	0.2	0.4	0.8	1.5	3.0
1999	8.8	*	*	*	*	*	0.2	0.3	0.8	1.6	2.9

Non-Hispanic 4/, male

2007	10.6	*	*	*	*	*	0.3	0.8	1.8	3.3	6.1
2006	10.3	*	*	*	*	*	0.4	0.8	1.5	3.4	5.5
2005	10.0	*	*	*	*	*	0.2	0.8	1.6	3.5	5.7
2004	9.8	*	*	*	*	*	*	0.8	1.7	3.0	5.5
2003	9.5	*	*	*	*	*	0.3	0.7	1.5	2.8	5.3
2002	8.9	*	*	*	*	*	*	0.8	1.5	2.8	5.0
2001	8.1	*	*	*	*	*	*	0.4	1.4	2.1	4.6
2000	7.8	*	*	*	*	*	0.3	0.4	1.0	1.9	4.0
1999	7.4	*	*	*	*	*	*	0.4	1.1	2.1	4.1

Non-Hispanic 4/, female

2007	12.0	*	*	*	*	*	*	0.3	0.8	1.3	3.0
2006	11.7	*	*	*	*	*	*	0.3	0.6	1.2	3.0
2005	11.8	*	*	*	*	*	*	0.4	0.8	1.3	2.7
2004	11.6	*	*	*	*	*	*	0.3	0.6	1.1	2.9
2003	11.7	*	*	*	*	*	*	0.3	0.8	1.3	2.8
2002	11.1	*	*	*	*	*	*	*	0.7	1.1	2.3
2001	10.7	*	*	*	*	*	*	0.3	0.6	1.1	2.1
2000	10.6	*	*	*	*	*	*	0.3	0.5	1.0	2.1
1999	10.2	*	*	*	*	*	*	*	0.6	1.1	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hypertensive heart disease (I11)

Non-Hispanic White, both sexes

2007 5/.....	10.6	*	*	*	*	*	*	0.3	0.7	1.4	3.1
2006 5/.....	10.3	*	*	*	*	*	0.2	0.3	0.6	1.7	3.1
2005 5/.....	10.1	*	*	*	*	*	*	0.4	0.8	1.6	2.8
2004 5/.....	9.8	*	*	*	*	*	*	0.3	0.7	1.3	2.9
2003 5/.....	9.7	*	*	*	*	*	*	0.3	0.7	1.3	2.7
2002.....	9.1	*	*	*	*	*	*	0.3	0.6	1.3	2.5
2001.....	8.6	*	*	*	*	*	*	0.2	0.6	1.0	2.1
2000.....	8.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0
1999.....	7.9	*	*	*	*	*	*	*	0.5	0.9	1.8

Non-Hispanic White, male

2007 5/.....	9.4	*	*	*	*	*	*	0.4	1.0	2.2	4.3
2006 5/.....	9.0	*	*	*	*	*	0.3	0.4	0.9	2.6	4.3
2005 5/.....	8.8	*	*	*	*	*	*	0.6	1.1	2.6	4.1
2004 5/.....	8.6	*	*	*	*	*	*	0.5	1.1	2.0	4.1
2003 5/.....	8.2	*	*	*	*	*	*	0.6	1.0	2.0	3.9
2002.....	7.6	*	*	*	*	*	*	0.4	0.9	1.9	3.7
2001.....	6.9	*	*	*	*	*	*	*	0.9	1.5	3.1
2000.....	6.6	*	*	*	*	*	*	*	0.6	1.3	2.9
1999.....	6.2	*	*	*	*	*	*	*	0.8	1.2	2.7

Non-Hispanic White, female

2007 5/.....	11.7	*	*	*	*	*	*	*	0.4	0.6	2.0
2006 5/.....	11.4	*	*	*	*	*	*	*	*	0.7	1.9
2005 5/.....	11.3	*	*	*	*	*	*	*	0.4	0.7	1.6
2004 5/.....	11.0	*	*	*	*	*	*	*	*	0.7	1.7
2003 5/.....	11.1	*	*	*	*	*	*	*	0.4	0.7	1.4
2002.....	10.5	*	*	*	*	*	*	*	0.3	0.7	1.2
2001.....	10.2	*	*	*	*	*	*	*	*	0.6	1.0
2000.....	9.9	*	*	*	*	*	*	*	*	0.5	1.2
1999.....	9.6	*	*	*	*	*	*	*	*	0.5	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Hypertensive heart disease (I11)

Non-Hispanic Black, both sexes

2007 5/.....	18.4	*	*	*	*	*	0.8	1.6	4.6	7.2	13.4
2006 5/.....	17.9	*	*	*	*	*	*	1.8	3.4	6.6	11.6
2005 5/.....	18.3	*	*	*	*	*	0.8	1.7	3.6	7.0	12.5
2004 5/.....	18.2	*	*	*	*	*	*	2.2	3.4	6.2	12.7
2003 5/.....	18.5	*	*	*	*	*	0.7	1.5	3.6	6.2	13.2
2002.....	17.7	*	*	*	*	*	*	1.6	4.1	6.0	11.6
2001.....	16.5	*	*	*	*	*	*	1.1	3.3	5.2	11.8
2000.....	16.6	*	*	*	*	*	*	1.4	2.9	5.1	9.9
1999.....	16.0	*	*	*	*	*	*	1.0	2.7	5.9	10.4

Non-Hispanic Black, male

2007 5/.....	19.9	*	*	*	*	*	*	2.3	6.3	10.1	18.2
2006 5/.....	19.5	*	*	*	*	*	*	2.8	4.7	9.3	13.2
2005 5/.....	19.1	*	*	*	*	*	*	1.8	4.3	9.5	16.6
2004 5/.....	19.0	*	*	*	*	*	*	2.6	4.7	9.0	15.0
2003 5/.....	18.7	*	*	*	*	*	*	1.8	4.5	7.6	15.4
2002.....	18.0	*	*	*	*	*	*	2.5	5.6	8.7	14.7
2001.....	16.3	*	*	*	*	*	*	*	4.2	6.0	15.1
2000.....	16.1	*	*	*	*	*	*	1.6	3.7	6.0	12.6
1999.....	15.6	*	*	*	*	*	*	*	2.9	7.6	13.3

Non-Hispanic Black, female

2007 5/.....	17.1	*	*	*	*	*	*	*	3.0	4.6	9.2
2006 5/.....	16.5	*	*	*	*	*	*	*	2.3	4.2	10.2
2005 5/.....	17.5	*	*	*	*	*	*	1.6	2.9	4.7	8.9
2004 5/.....	17.5	*	*	*	*	*	*	1.9	2.2	3.8	10.6
2003 5/.....	18.4	*	*	*	*	*	*	*	2.8	4.9	11.3
2002.....	17.4	*	*	*	*	*	*	*	2.8	3.5	8.9
2001.....	16.7	*	*	*	*	*	*	*	2.4	4.6	8.9
2000.....	17.1	*	*	*	*	*	*	*	2.2	4.3	7.6
1999.....	16.4	*	*	*	*	*	*	*	2.6	4.5	7.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Ischemic heart diseases (I20-I25)

All origins 3/, both sexes

2007.....	134.7	0.6	*	*	*	0.2	0.5	1.6	3.7	8.4	20.2
2006.....	142.1	0.5	*	*	*	0.2	0.5	1.4	3.9	8.8	21.5
2005.....	150.4	*	*	*	*	0.1	0.6	1.4	3.6	9.1	21.6
2004.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4	22.4
2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1	23.5
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1	23.8
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9	23.4
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8	23.6
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4	24.2

All origins 3/, male

2007.....	145.3	*	*	*	*	0.2	0.7	2.3	5.7	12.6	30.9
2006.....	152.2	*	*	*	*	0.3	0.7	2.1	6.1	13.2	32.7
2005.....	159.0	*	*	*	*	0.2	0.9	2.2	5.4	13.6	33.0
2004.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9	34.0
2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2	35.5
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2	36.5
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0	35.8
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7	37.0
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8	37.9

All origins 3/, female

2007.....	124.4	*	*	*	*	*	0.3	0.8	1.6	4.2	9.4
2006.....	132.3	*	*	*	*	*	0.2	0.6	1.7	4.4	10.3
2005.....	142.0	*	*	*	*	*	0.2	0.6	1.9	4.6	10.4
2004.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9	10.8
2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9	11.6
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9	11.2
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9	11.1
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0	10.3
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Ischemic heart diseases (I20-I25)

Non-Hispanic 4/, both sexes

2007.....	150.4	*	*	*	*	0.2	0.6	1.7	4.2	9.3	21.9
2006.....	158.2	0.6	*	*	*	0.2	0.5	1.5	4.4	9.7	23.2
2005.....	166.7	*	*	*	*	0.1	0.6	1.6	4.1	10.1	23.0
2004.....	170.3	*	*	*	*	0.1	0.5	1.7	4.3	10.3	23.8
2003.....	182.5	0.6	*	*	*	0.2	0.6	1.6	4.3	10.9	25.1
2002.....	189.0	*	*	*	*	0.2	0.5	1.4	3.9	11.0	25.3
2001.....	193.6	0.7	*	*	*	0.2	0.7	1.6	4.4	10.8	24.9
2000.....	200.5	0.8	*	*	*	0.2	0.5	1.3	3.7	10.6	24.7
1999.....	207.1	0.8	*	*	*	0.3	0.5	1.3	4.0	11.1	25.5

Non-Hispanic 4/, male

2007.....	163.3	*	*	*	*	0.3	0.8	2.5	6.5	13.9	33.7
2006.....	170.6	*	*	*	*	0.3	0.7	2.4	6.9	14.6	35.6
2005.....	177.2	*	*	*	*	0.2	1.0	2.5	6.0	15.1	35.2
2004.....	179.9	*	*	*	*	*	0.7	2.4	6.6	15.3	36.2
2003.....	191.5	*	*	*	*	*	0.8	2.2	6.5	16.7	38.1
2002.....	197.6	*	*	*	*	*	0.7	2.2	5.7	16.8	39.0
2001.....	200.4	*	*	*	*	0.2	1.0	2.4	6.4	16.3	38.2
2000.....	207.6	*	*	*	*	*	0.7	2.1	5.5	15.8	38.9
1999.....	214.1	*	*	*	*	0.4	0.7	1.8	6.0	16.8	40.0

Non-Hispanic 4/, female

2007.....	138.1	*	*	*	*	*	0.4	0.9	1.8	4.7	10.4
2006.....	146.4	*	*	*	*	*	0.2	0.7	1.8	4.9	11.1
2005.....	156.7	*	*	*	*	*	*	0.6	2.1	5.1	11.2
2004.....	161.1	*	*	*	*	*	0.3	1.0	2.1	5.4	11.6
2003.....	173.9	*	*	*	*	*	0.5	0.9	2.0	5.3	12.4
2002.....	180.7	*	*	*	*	*	0.3	0.7	2.1	5.3	12.0
2001.....	187.2	*	*	*	*	*	0.4	0.8	2.3	5.3	11.9
2000.....	193.8	*	*	*	*	*	0.3	0.6	1.9	5.4	10.9
1999.....	200.6	*	*	*	*	*	0.3	0.9	2.1	5.6	11.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Ischemic heart diseases (I20-I25)

Non-Hispanic White, both sexes

2007 5/.....	166.0	*	*	*	*	0.2	0.5	1.4	3.9	8.9	21.4
2006 5/.....	174.5	*	*	*	*	0.2	0.5	1.4	4.1	9.2	22.9
2005 5/.....	183.9	*	*	*	*	*	0.5	1.3	3.9	9.5	22.5
2004 5/.....	187.7	*	*	*	*	*	0.3	1.5	4.0	10.3	23.1
2003 5/.....	200.8	*	*	*	*	0.2	0.6	1.4	4.1	10.8	24.1
2002.....	208.0	*	*	*	*	0.2	0.4	1.3	3.7	10.6	24.5
2001.....	212.6	*	*	*	*	*	0.6	1.4	4.1	10.4	23.6
2000.....	220.2	*	*	*	*	0.2	0.5	1.1	3.5	10.2	23.4
1999.....	227.2	*	*	*	*	0.2	0.4	1.2	3.7	10.7	23.9

Non-Hispanic White, male

2007 5/.....	179.9	*	*	*	*	*	0.7	2.1	6.1	13.5	33.2
2006 5/.....	187.5	*	*	*	*	0.3	0.7	2.3	6.7	14.0	35.9
2005 5/.....	195.1	*	*	*	*	*	0.9	2.2	5.7	14.3	35.1
2004 5/.....	198.0	*	*	*	*	*	0.5	2.2	6.2	15.2	36.1
2003 5/.....	210.2	*	*	*	*	*	0.7	2.0	6.3	16.7	37.1
2002.....	217.2	*	*	*	*	*	0.6	2.1	5.5	16.3	38.5
2001.....	219.6	*	*	*	*	*	0.9	2.2	6.0	15.8	37.2
2000.....	227.6	*	*	*	*	*	0.6	1.8	5.3	15.5	37.8
1999.....	234.2	*	*	*	*	0.3	0.6	1.6	5.5	16.2	38.3

Non-Hispanic White, female

2007 5/.....	152.6	*	*	*	*	*	*	0.8	1.7	4.3	9.7
2006 5/.....	161.9	*	*	*	*	*	*	0.6	1.6	4.3	10.0
2005 5/.....	173.1	*	*	*	*	*	*	0.5	2.1	4.6	9.9
2004 5/.....	177.7	*	*	*	*	*	*	0.8	1.9	5.3	10.1
2003 5/.....	191.8	*	*	*	*	*	0.4	0.8	1.9	4.8	11.1
2002.....	199.1	*	*	*	*	*	*	0.6	1.9	4.8	10.5
2001.....	205.8	*	*	*	*	*	*	0.6	2.1	4.9	10.1
2000.....	213.1	*	*	*	*	*	*	0.5	1.6	4.9	9.0
1999.....	220.5	*	*	*	*	*	*	0.9	1.9	5.1	9.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Ischemic heart diseases (I20-I25)

Non-Hispanic Black, both sexes

2007 5/.....	110.5	*	*	*	*	*	1.1	3.0	6.5	13.3	28.4
2006 5/.....	116.5	*	*	*	*	*	0.8	2.4	6.5	15.3	30.6
2005 5/.....	121.7	*	*	*	*	*	0.9	2.7	5.6	16.1	31.4
2004 5/.....	124.8	*	*	*	*	*	1.3	3.0	6.8	13.1	34.2
2003 5/.....	133.2	*	*	*	*	*	1.2	2.5	6.2	14.6	36.0
2002.....	137.0	*	*	*	*	*	0.9	2.2	5.8	15.5	36.1
2001.....	141.2	*	*	*	*	*	1.5	2.9	6.6	15.4	37.5
2000.....	144.4	*	*	*	*	*	0.8	2.6	6.2	15.0	37.7
1999.....	147.9	*	*	*	*	*	1.2	2.0	6.1	16.4	40.7

Non-Hispanic Black, male

2007 5/.....	118.0	*	*	*	*	*	1.6	4.4	10.4	19.6	41.9
2006 5/.....	124.5	*	*	*	*	*	*	3.4	9.7	21.5	42.4
2005 5/.....	127.0	*	*	*	*	*	1.3	4.4	8.3	23.8	42.8
2004 5/.....	129.2	*	*	*	*	*	1.9	3.5	10.0	19.0	46.1
2003 5/.....	137.6	*	*	*	*	*	1.5	3.3	9.5	20.4	51.5
2002.....	139.5	*	*	*	*	*	*	2.9	7.7	22.5	49.8
2001.....	143.2	*	*	*	*	*	2.0	4.0	9.5	22.5	52.0
2000.....	145.7	*	*	*	*	*	*	3.9	8.2	20.8	52.5
1999.....	150.0	*	*	*	*	*	*	3.1	8.8	23.8	58.1

Non-Hispanic Black, female

2007 5/.....	103.6	*	*	*	*	*	*	1.6	3.0	7.8	16.6
2006 5/.....	109.2	*	*	*	*	*	*	1.5	3.5	9.8	20.2
2005 5/.....	116.8	*	*	*	*	*	*	*	3.1	9.2	21.3
2004 5/.....	120.8	*	*	*	*	*	*	2.5	3.9	7.8	23.7
2003 5/.....	129.1	*	*	*	*	*	*	1.8	3.3	9.4	22.4
2002.....	134.8	*	*	*	*	*	*	1.6	4.1	9.3	23.9
2001.....	139.4	*	*	*	*	*	*	1.8	4.0	9.2	24.7
2000.....	143.2	*	*	*	*	*	*	*	4.4	9.9	24.7
1999.....	146.0	*	*	*	*	*	*	*	3.6	9.9	25.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Acute myocardial infarction (I21-I22)

All origins 3/, both sexes

2007.....	44.1	*	*	*	*	*	0.2	0.6	1.4	3.2	7.8
2006.....	47.2	*	*	*	*	0.1	0.2	0.6	1.5	3.6	8.5
2005.....	50.9	*	*	*	*	*	0.3	0.6	1.3	3.5	8.7
2004.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8	8.9
2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1	9.6
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0	9.9
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2	9.9
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0	9.9
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1	10.4

All origins 3/, male

2007.....	48.2	*	*	*	*	*	0.2	0.8	2.1	4.6	11.6
2006.....	51.6	*	*	*	*	*	0.2	0.9	2.3	5.2	13.0
2005.....	54.8	*	*	*	*	*	0.4	1.0	2.0	5.1	13.1
2004.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7	13.2
2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2	14.4
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0	14.8
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2	15.0
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8	15.5
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2	16.0

All origins 3/, female

2007.....	40.0	*	*	*	*	*	*	0.3	0.7	1.8	4.0
2006.....	43.0	*	*	*	*	*	*	0.3	0.8	1.9	4.1
2005.....	47.2	*	*	*	*	*	*	0.3	0.7	2.0	4.4
2004.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9	4.5
2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0	5.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1	5.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1	4.9
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2	4.4
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0	4.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Acute myocardial infarction (I21-I22)

Non-Hispanic 4/, both sexes

2007.....	49.2	*	*	*	*	*	0.2	0.7	1.6	3.5	8.6
2006.....	52.6	*	*	*	*	*	0.2	0.7	1.7	4.0	9.2
2005.....	56.5	*	*	*	*	*	0.3	0.7	1.5	3.9	9.4
2004.....	59.2	*	*	*	*	*	0.2	0.7	1.8	4.2	9.5
2003.....	64.9	*	*	*	*	*	0.3	0.6	1.6	4.5	10.3
2002.....	68.7	*	*	*	*	*	0.3	0.5	1.6	4.4	10.6
2001.....	71.4	*	*	*	*	*	0.3	0.6	1.9	4.6	10.6
2000.....	75.3	*	*	*	*	*	0.2	0.6	1.4	4.3	10.5
1999.....	78.2	*	*	*	*	0.1	0.2	0.6	1.8	4.4	11.0

Non-Hispanic 4/, male

2007.....	54.2	*	*	*	*	*	0.2	0.9	2.5	5.1	12.9
2006.....	57.9	*	*	*	*	*	0.2	1.0	2.6	5.9	14.2
2005.....	61.2	*	*	*	*	*	0.4	1.1	2.3	5.7	14.3
2004.....	64.0	*	*	*	*	*	0.4	0.9	2.6	6.3	14.2
2003.....	69.7	*	*	*	*	*	0.3	0.9	2.4	6.9	15.5
2002.....	73.5	*	*	*	*	*	0.3	0.8	2.3	6.6	15.9
2001.....	75.9	*	*	*	*	*	0.4	0.8	2.7	6.9	16.0
2000.....	80.3	*	*	*	*	*	*	0.9	2.0	6.3	16.4
1999.....	83.8	*	*	*	*	*	0.3	0.7	2.5	6.6	17.0

Non-Hispanic 4/, female

2007.....	44.4	*	*	*	*	*	*	0.4	0.8	2.0	4.5
2006.....	47.6	*	*	*	*	*	*	0.4	0.8	2.1	4.4
2005.....	52.0	*	*	*	*	*	*	0.3	0.7	2.2	4.6
2004.....	54.7	*	*	*	*	*	*	0.5	0.9	2.1	4.9
2003.....	60.3	*	*	*	*	*	0.2	0.4	0.9	2.2	5.3
2002.....	64.1	*	*	*	*	*	*	0.3	0.9	2.2	5.4
2001.....	67.1	*	*	*	*	*	*	0.4	1.1	2.3	5.3
2000.....	70.6	*	*	*	*	*	*	0.3	0.8	2.3	4.7
1999.....	72.9	*	*	*	*	*	*	0.4	1.0	2.2	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Acute myocardial infarction (I21-I22)

Non-Hispanic White, both sexes

2007 5/.....	54.2	*	*	*	*	*	0.2	0.6	1.6	3.6	8.6
2006 5/.....	58.0	*	*	*	*	*	0.2	0.7	1.6	4.0	9.3
2005 5/.....	62.3	*	*	*	*	*	0.2	0.5	1.5	3.8	9.3
2004 5/.....	65.3	*	*	*	*	*	0.2	0.6	1.8	4.2	9.4
2003 5/.....	71.5	*	*	*	*	*	0.2	0.6	1.6	4.6	10.1
2002.....	75.5	*	*	*	*	*	0.2	0.5	1.6	4.3	10.4
2001.....	78.4	*	*	*	*	*	0.2	0.5	1.9	4.5	10.3
2000.....	82.7	*	*	*	*	*	0.2	0.5	1.4	4.4	10.1
1999.....	86.0	*	*	*	*	*	*	0.5	1.7	4.4	10.7

Non-Hispanic White, male

2007 5/.....	60.0	*	*	*	*	*	*	0.8	2.4	5.3	13.0
2006 5/.....	64.0	*	*	*	*	*	*	0.9	2.6	6.0	14.5
2005 5/.....	67.8	*	*	*	*	*	0.4	0.9	2.2	5.6	14.5
2004 5/.....	70.9	*	*	*	*	*	*	0.9	2.8	6.3	14.5
2003 5/.....	77.0	*	*	*	*	*	*	0.9	2.3	7.1	15.3
2002.....	81.2	*	*	*	*	*	*	0.7	2.3	6.7	15.9
2001.....	83.7	*	*	*	*	*	0.3	0.6	2.7	6.8	16.1
2000.....	88.5	*	*	*	*	*	*	0.7	2.0	6.5	16.2
1999.....	92.4	*	*	*	*	*	*	0.7	2.3	6.7	16.8

Non-Hispanic White, female

2007 5/.....	48.7	*	*	*	*	*	*	0.3	0.7	1.9	4.3
2006 5/.....	52.2	*	*	*	*	*	*	0.4	0.7	1.9	4.2
2005 5/.....	57.0	*	*	*	*	*	*	*	0.7	2.1	4.2
2004 5/.....	59.9	*	*	*	*	*	*	0.4	0.8	2.1	4.2
2003 5/.....	66.2	*	*	*	*	*	*	*	0.9	2.0	4.8
2002.....	70.1	*	*	*	*	*	*	*	0.8	2.0	5.0
2001.....	73.4	*	*	*	*	*	*	0.4	1.1	2.2	4.5
2000.....	77.2	*	*	*	*	*	*	*	0.7	2.2	3.9
1999.....	79.8	*	*	*	*	*	*	0.4	1.0	2.0	4.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Acute myocardial infarction (I21-I22)

Non-Hispanic Black, both sexes

2007 5/.....	36.6	*	*	*	*	*	*	1.0	2.6	4.5	10.6
2006 5/.....	38.7	*	*	*	*	*	*	1.0	2.4	5.4	11.5
2005 5/.....	41.1	*	*	*	*	*	*	1.5	2.1	5.6	12.3
2004 5/.....	43.4	*	*	*	*	*	*	1.2	2.4	5.1	12.8
2003 5/.....	47.3	*	*	*	*	*	*	0.9	2.4	5.9	14.8
2002.....	50.2	*	*	*	*	*	*	1.0	2.4	5.7	14.2
2001.....	51.6	*	*	*	*	*	*	1.1	2.6	6.0	14.6
2000.....	54.0	*	*	*	*	*	*	1.2	2.3	5.1	15.3
1999.....	55.1	*	*	*	*	*	*	0.9	2.5	5.6	15.8

Non-Hispanic Black, male

2007 5/.....	38.1	*	*	*	*	*	*	1.4	4.0	6.3	15.2
2006 5/.....	40.6	*	*	*	*	*	*	1.5	3.5	7.1	16.4
2005 5/.....	41.8	*	*	*	*	*	*	2.2	3.1	7.8	16.5
2004 5/.....	43.9	*	*	*	*	*	*	*	2.9	8.0	15.8
2003 5/.....	48.2	*	*	*	*	*	*	*	3.7	7.9	20.6
2002.....	50.1	*	*	*	*	*	*	*	3.1	7.8	19.6
2001.....	50.9	*	*	*	*	*	*	*	3.5	8.3	18.8
2000.....	53.7	*	*	*	*	*	*	1.7	2.7	6.7	20.9
1999.....	55.2	*	*	*	*	*	*	*	3.6	7.8	21.6

Non-Hispanic Black, female

2007 5/.....	35.3	*	*	*	*	*	*	*	*	2.8	6.6
2006 5/.....	37.0	*	*	*	*	*	*	*	1.5	3.9	7.1
2005 5/.....	40.5	*	*	*	*	*	*	*	*	3.7	8.7
2004 5/.....	43.0	*	*	*	*	*	*	*	1.9	2.6	10.1
2003 5/.....	46.5	*	*	*	*	*	*	*	*	4.0	9.6
2002.....	50.4	*	*	*	*	*	*	*	1.7	3.7	9.5
2001.....	52.3	*	*	*	*	*	*	*	1.8	3.9	10.9
2000.....	54.2	*	*	*	*	*	*	*	1.9	3.7	10.3
1999.....	54.9	*	*	*	*	*	*	*	1.5	3.6	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes

2007.....	89.3	*	*	*	*	0.1	0.3	0.9	2.2	5.1	12.0
2006.....	93.5	*	*	*	*	*	0.3	0.8	2.3	5.1	12.7
2005.....	98.2	*	*	*	*	*	0.3	0.8	2.3	5.5	12.7
2004.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5	13.2
2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8	13.5
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0	13.6
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6	13.2
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7	13.4
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2	13.5

All origins 3/, male

2007.....	95.6	*	*	*	*	*	0.5	1.4	3.5	7.7	18.8
2006.....	99.2	*	*	*	*	*	0.4	1.2	3.7	7.9	19.4
2005.....	102.9	*	*	*	*	*	0.5	1.2	3.4	8.4	19.5
2004.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1	20.4
2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8	20.7
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2	21.4
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6	20.4
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7	21.1
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4	21.5

All origins 3/, female

2007.....	83.1	*	*	*	*	*	*	0.5	0.9	2.3	5.2
2006.....	88.0	*	*	*	*	*	*	0.3	0.9	2.3	6.0
2005.....	93.7	*	*	*	*	*	*	0.3	1.2	2.5	5.9
2004.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0	6.1
2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8	6.5
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8	6.0
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7	6.1
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8	5.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1	5.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 4/, both sexes

2007.....	99.7	*	*	*	*	0.1	0.4	1.0	2.5	5.6	12.9
2006.....	104.1	*	*	*	*	*	0.3	0.8	2.6	5.6	13.7
2005.....	108.9	*	*	*	*	*	0.3	0.8	2.5	6.0	13.3
2004.....	110.1	*	*	*	*	*	0.3	1.0	2.5	6.0	14.0
2003.....	116.3	*	*	*	*	*	0.3	0.9	2.6	6.3	14.4
2002.....	118.9	*	*	*	*	*	0.3	0.9	2.2	6.5	14.5
2001.....	120.9	*	*	*	*	*	0.4	1.0	2.4	6.1	14.0
2000.....	123.9	*	*	*	*	*	0.3	0.7	2.2	6.1	14.0
1999.....	127.6	*	*	*	*	0.1	0.3	0.8	2.2	6.6	14.2

Non-Hispanic 4/, male

2007.....	107.4	*	*	*	*	*	0.6	1.5	3.9	8.6	20.3
2006.....	111.1	*	*	*	*	*	0.5	1.3	4.2	8.6	21.0
2005.....	114.6	*	*	*	*	*	0.5	1.4	3.7	9.3	20.5
2004.....	114.9	*	*	*	*	*	0.3	1.5	3.9	8.8	21.7
2003.....	120.4	*	*	*	*	*	0.4	1.3	4.1	9.6	22.1
2002.....	122.7	*	*	*	*	*	0.4	1.3	3.3	10.0	22.7
2001.....	123.1	*	*	*	*	*	0.6	1.6	3.6	9.3	21.8
2000.....	126.0	*	*	*	*	*	0.4	1.2	3.4	9.3	22.0
1999.....	128.9	*	*	*	*	*	0.4	1.1	3.4	10.0	22.6

Non-Hispanic 4/, female

2007.....	92.3	*	*	*	*	*	*	0.5	1.0	2.6	5.7
2006.....	97.4	*	*	*	*	*	*	0.3	1.0	2.6	6.5
2005.....	103.4	*	*	*	*	*	*	0.3	1.4	2.8	6.4
2004.....	105.6	*	*	*	*	*	*	0.5	1.1	3.3	6.5
2003.....	112.4	*	*	*	*	*	0.2	0.5	1.2	3.0	6.9
2002.....	115.4	*	*	*	*	*	*	0.4	1.1	3.1	6.4
2001.....	118.8	*	*	*	*	*	*	0.4	1.1	2.9	6.5
2000.....	122.0	*	*	*	*	*	*	*	1.1	3.0	6.1
1999.....	126.4	*	*	*	*	*	*	0.5	1.1	3.4	6.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic White, both sexes

2007 5/.....	110.1	*	*	*	*	*	0.3	0.8	2.3	5.2	12.4
2006 5/.....	114.9	*	*	*	*	*	0.3	0.7	2.4	5.1	13.3
2005 5/.....	120.2	*	*	*	*	*	0.3	0.8	2.4	5.5	12.9
2004 5/.....	121.4	*	*	*	*	*	0.2	0.9	2.2	5.9	13.4
2003 5/.....	128.0	*	*	*	*	*	0.3	0.8	2.5	6.1	13.8
2002.....	131.0	*	*	*	*	*	0.2	0.9	2.1	6.1	13.8
2001.....	132.7	*	*	*	*	*	0.3	0.9	2.1	5.7	13.1
2000.....	136.1	*	*	*	*	*	0.2	0.6	2.0	5.7	13.1
1999.....	139.9	*	*	*	*	*	0.2	0.7	2.0	6.2	13.0

Non-Hispanic White, male

2007 5/.....	118.2	*	*	*	*	*	0.5	1.2	3.7	8.0	19.7
2006 5/.....	121.9	*	*	*	*	*	0.4	1.3	4.0	7.9	21.0
2005 5/.....	125.9	*	*	*	*	*	0.4	1.3	3.5	8.5	20.2
2004 5/.....	126.1	*	*	*	*	*	*	1.4	3.4	8.7	21.2
2003 5/.....	131.9	*	*	*	*	*	0.4	1.1	3.9	9.3	21.3
2002.....	134.5	*	*	*	*	*	0.4	1.3	3.1	9.5	22.2
2001.....	134.4	*	*	*	*	*	0.5	1.6	3.2	8.8	20.7
2000.....	137.7	*	*	*	*	*	0.4	1.0	3.2	8.8	21.2
1999.....	140.4	*	*	*	*	*	*	0.9	3.1	9.3	21.1

Non-Hispanic White, female

2007 5/.....	102.4	*	*	*	*	*	*	0.5	1.0	2.4	5.2
2006 5/.....	108.1	*	*	*	*	*	*	*	0.9	2.3	5.7
2005 5/.....	114.7	*	*	*	*	*	*	*	1.4	2.5	5.6
2004 5/.....	116.9	*	*	*	*	*	*	0.4	1.1	3.1	5.7
2003 5/.....	124.3	*	*	*	*	*	*	0.5	1.0	2.8	6.2
2002.....	127.7	*	*	*	*	*	*	0.4	1.0	2.8	5.5
2001.....	131.1	*	*	*	*	*	*	*	1.0	2.6	5.5
2000.....	134.6	*	*	*	*	*	*	*	0.9	2.6	4.9
1999.....	139.3	*	*	*	*	*	*	0.5	0.9	3.1	4.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic Black, both sexes

2007 5/.....	72.5	*	*	*	*	*	0.7	1.9	3.7	8.4	17.3
2006 5/.....	76.4	*	*	*	*	*	*	1.4	3.8	9.4	18.6
2005 5/.....	79.1	*	*	*	*	*	*	1.1	3.4	10.2	18.6
2004 5/.....	80.3	*	*	*	*	*	*	1.7	4.3	7.9	20.9
2003 5/.....	84.7	*	*	*	*	*	*	1.5	3.8	8.5	20.6
2002.....	85.5	*	*	*	*	*	*	1.2	3.3	9.8	21.2
2001.....	88.2	*	*	*	*	*	0.8	1.8	3.9	9.2	22.2
2000.....	89.1	*	*	*	*	*	*	1.3	3.7	9.7	21.9
1999.....	91.6	*	*	*	*	*	*	1.2	3.5	10.4	24.4

Non-Hispanic Black, male

2007 5/.....	78.4	*	*	*	*	*	*	2.8	6.2	12.8	25.9
2006 5/.....	82.3	*	*	*	*	*	*	1.8	5.9	13.8	25.3
2005 5/.....	83.6	*	*	*	*	*	*	2.1	5.1	15.9	25.6
2004 5/.....	83.9	*	*	*	*	*	*	2.1	7.0	10.8	29.5
2003 5/.....	87.9	*	*	*	*	*	*	2.2	5.7	12.2	30.1
2002.....	88.0	*	*	*	*	*	*	*	4.3	14.6	29.9
2001.....	90.8	*	*	*	*	*	*	2.5	6.0	13.8	32.1
2000.....	90.4	*	*	*	*	*	*	2.2	5.1	13.7	30.8
1999.....	93.3	*	*	*	*	*	*	1.8	5.2	15.3	35.6

Non-Hispanic Black, female

2007 5/.....	67.2	*	*	*	*	*	*	*	1.5	4.5	9.7
2006 5/.....	71.0	*	*	*	*	*	*	*	1.9	5.4	12.8
2005 5/.....	75.0	*	*	*	*	*	*	*	1.9	5.2	12.4
2004 5/.....	76.9	*	*	*	*	*	*	*	1.8	5.3	13.2
2003 5/.....	81.7	*	*	*	*	*	*	*	2.0	5.1	12.2
2002.....	83.3	*	*	*	*	*	*	*	2.4	5.5	13.6
2001.....	86.0	*	*	*	*	*	*	*	2.0	5.0	13.4
2000.....	88.0	*	*	*	*	*	*	*	2.5	6.0	14.1
1999.....	90.1	*	*	*	*	*	*	*	2.0	6.0	14.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins 3/, both sexes

2007.....	19.6	*	*	*	*	*	*	0.3	0.7	1.9	4.5
2006.....	20.4	*	*	*	*	*	*	0.2	0.8	1.9	4.6
2005.....	21.2	*	*	*	*	*	*	0.2	0.7	1.9	4.4
2004.....	21.7	*	*	*	*	*	*	0.2	0.8	1.7	4.8
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9	4.7
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9	4.8
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8	4.6
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7	4.4
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0	4.7

All origins 3/, male

2007.....	22.5	*	*	*	*	*	*	0.5	1.2	2.9	7.3
2006.....	23.2	*	*	*	*	*	*	0.4	1.4	3.1	7.0
2005.....	23.6	*	*	*	*	*	*	0.4	1.1	3.0	7.0
2004.....	23.9	*	*	*	*	*	*	0.3	1.2	2.5	7.7
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9	7.4
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0	7.8
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8	7.2
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8	7.1
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1	7.6

All origins 3/, female

2007.....	16.8	*	*	*	*	*	*	*	0.2	0.8	1.8
2006.....	17.7	*	*	*	*	*	*	*	0.3	0.7	2.2
2005.....	18.8	*	*	*	*	*	*	*	0.3	0.8	1.9
2004.....	19.5	*	*	*	*	*	*	*	0.3	0.8	2.0
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9	2.0
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8	1.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8	2.1
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7	1.8
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 4/, both sexes

2007.....	21.7	*	*	*	*	*	*	0.3	0.8	2.1	4.8
2006.....	22.5	*	*	*	*	*	*	0.2	0.9	2.1	4.9
2005.....	23.3	*	*	*	*	*	*	0.2	0.8	2.1	4.7
2004.....	23.8	*	*	*	*	*	*	0.2	0.8	1.9	5.1
2003.....	25.3	*	*	*	*	*	*	0.3	0.8	2.1	5.0
2002.....	25.9	*	*	*	*	*	*	0.2	0.7	2.0	5.0
2001.....	26.0	*	*	*	*	*	*	0.2	0.7	1.9	4.9
2000.....	26.9	*	*	*	*	*	*	0.2	0.6	1.8	4.6
1999.....	27.9	*	*	*	*	*	*	0.2	0.7	2.1	4.9

Non-Hispanic 4/, male

2007.....	24.9	*	*	*	*	*	*	0.5	1.4	3.3	7.8
2006.....	25.5	*	*	*	*	*	*	0.4	1.6	3.4	7.6
2005.....	25.8	*	*	*	*	*	*	0.4	1.2	3.3	7.5
2004.....	26.2	*	*	*	*	*	*	0.4	1.3	2.8	8.1
2003.....	27.5	*	*	*	*	*	*	0.4	1.5	3.2	7.8
2002.....	27.8	*	*	*	*	*	*	0.3	1.1	3.2	8.2
2001.....	27.5	*	*	*	*	*	*	0.3	1.0	3.1	7.6
2000.....	28.2	*	*	*	*	*	*	0.3	1.0	2.9	7.4
1999.....	28.9	*	*	*	*	*	*	*	1.0	3.3	8.0

Non-Hispanic 4/, female

2007.....	18.6	*	*	*	*	*	*	*	0.3	0.9	2.0
2006.....	19.6	*	*	*	*	*	*	*	0.3	0.9	2.3
2005.....	20.8	*	*	*	*	*	*	*	0.3	0.9	2.0
2004.....	21.5	*	*	*	*	*	*	*	0.3	0.9	2.2
2003.....	23.3	*	*	*	*	*	*	*	*	1.0	2.2
2002.....	24.0	*	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	24.6	*	*	*	*	*	*	*	0.4	0.8	2.2
2000.....	25.6	*	*	*	*	*	*	*	0.3	0.7	1.9
1999.....	27.0	*	*	*	*	*	*	*	0.3	1.0	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic Black, both sexes

2007 5/.....	23.1	*	*	*	*	*	*	0.7	1.3	3.4	6.6
2006 5/.....	24.4	*	*	*	*	*	*	*	1.6	3.5	7.9
2005 5/.....	25.3	*	*	*	*	*	*	*	1.2	4.0	6.7
2004 5/.....	25.7	*	*	*	*	*	*	*	1.6	2.0	7.7
2003 5/.....	26.7	*	*	*	*	*	*	*	1.3	2.7	7.4
2002.....	27.7	*	*	*	*	*	*	*	1.3	3.6	8.3
2001.....	28.5	*	*	*	*	*	*	*	1.0	3.2	8.5
2000.....	29.3	*	*	*	*	*	*	*	0.9	3.0	7.6
1999.....	30.7	*	*	*	*	*	*	*	1.2	3.9	9.3

Non-Hispanic Black, male

2007 5/.....	27.9	*	*	*	*	*	*	*	2.4	5.3	10.2
2006 5/.....	29.5	*	*	*	*	*	*	*	2.6	5.8	10.9
2005 5/.....	29.7	*	*	*	*	*	*	*	1.8	6.3	9.3
2004 5/.....	29.7	*	*	*	*	*	*	*	2.7	3.0	10.9
2003 5/.....	31.2	*	*	*	*	*	*	*	2.3	4.0	11.6
2002.....	31.9	*	*	*	*	*	*	*	1.9	5.6	12.2
2001.....	32.3	*	*	*	*	*	*	*	*	4.8	12.8
2000.....	33.2	*	*	*	*	*	*	*	*	4.8	11.2
1999.....	34.5	*	*	*	*	*	*	*	1.8	6.0	13.9

Non-Hispanic Black, female

2007 5/.....	18.7	*	*	*	*	*	*	*	*	1.8	3.4
2006 5/.....	19.8	*	*	*	*	*	*	*	*	1.5	5.3
2005 5/.....	21.3	*	*	*	*	*	*	*	*	1.9	4.5
2004 5/.....	22.1	*	*	*	*	*	*	*	*	*	4.9
2003 5/.....	22.7	*	*	*	*	*	*	*	*	1.5	3.8
2002.....	24.0	*	*	*	*	*	*	*	*	1.8	5.0
2001.....	25.0	*	*	*	*	*	*	*	*	1.7	4.7
2000.....	25.8	*	*	*	*	*	*	*	*	1.5	4.4
1999.....	27.1	*	*	*	*	*	*	*	*	2.0	5.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes

2007.....	69.7	*	*	*	*	0.1	0.3	0.6	1.5	3.2	7.5
2006.....	73.1	*	*	*	*	*	0.2	0.5	1.5	3.2	8.1
2005.....	77.0	*	*	*	*	*	0.2	0.5	1.6	3.6	8.2
2004.....	77.8	*	*	*	*	*	0.2	0.7	1.5	3.9	8.4
2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9	8.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1	8.8
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8	8.6
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0	9.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3	8.8

All origins 3/, male

2007.....	73.2	*	*	*	*	*	0.4	0.9	2.3	4.8	11.5
2006.....	76.1	*	*	*	*	*	0.3	0.9	2.3	4.8	12.3
2005.....	79.3	*	*	*	*	*	0.4	0.9	2.3	5.4	12.5
2004.....	79.4	*	*	*	*	*	0.3	1.0	2.3	5.6	12.7
2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9	13.3
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2	13.6
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8	13.2
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9	14.0
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3	13.9

All origins 3/, female

2007.....	66.3	*	*	*	*	*	*	0.3	0.7	1.5	3.4
2006.....	70.3	*	*	*	*	*	*	*	0.6	1.6	3.8
2005.....	74.8	*	*	*	*	*	*	0.2	0.9	1.7	4.0
2004.....	76.2	*	*	*	*	*	*	0.4	0.7	2.1	4.1
2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9	4.5
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0	4.2
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9	4.0
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1	4.0
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 4/, both sexes

2007.....	78.0	*	*	*	*	*	0.3	0.7	1.7	3.5	8.1
2006.....	81.6	*	*	*	*	*	0.2	0.6	1.6	3.5	8.7
2005.....	85.6	*	*	*	*	*	0.3	0.6	1.8	3.9	8.6
2004.....	86.3	*	*	*	*	*	0.2	0.8	1.7	4.2	8.9
2003.....	91.0	*	*	*	*	*	0.3	0.6	1.8	4.2	9.5
2002.....	93.1	*	*	*	*	*	0.2	0.7	1.5	4.5	9.4
2001.....	94.9	*	*	*	*	*	0.3	0.8	1.6	4.1	9.2
2000.....	97.0	*	*	*	*	*	0.2	0.5	1.6	4.3	9.4
1999.....	99.7	*	*	*	*	0.1	0.2	0.6	1.6	4.5	9.3

Non-Hispanic 4/, male

2007.....	82.5	*	*	*	*	*	0.5	0.9	2.6	5.3	12.5
2006.....	85.6	*	*	*	*	*	0.3	1.0	2.6	5.2	13.4
2005.....	88.8	*	*	*	*	*	0.4	1.0	2.6	5.9	13.0
2004.....	88.7	*	*	*	*	*	0.3	1.1	2.6	6.0	13.6
2003.....	92.9	*	*	*	*	*	0.3	0.9	2.6	6.4	14.3
2002.....	94.9	*	*	*	*	*	0.3	1.1	2.2	6.8	14.5
2001.....	95.6	*	*	*	*	*	0.4	1.3	2.5	6.2	14.1
2000.....	97.8	*	*	*	*	*	0.3	0.9	2.4	6.4	14.7
1999.....	100.0	*	*	*	*	*	*	0.8	2.4	6.6	14.6

Non-Hispanic 4/, female

2007.....	73.6	*	*	*	*	*	*	0.4	0.8	1.7	3.7
2006.....	77.9	*	*	*	*	*	*	*	0.7	1.8	4.2
2005.....	82.6	*	*	*	*	*	*	*	1.0	1.9	4.4
2004.....	84.1	*	*	*	*	*	*	0.4	0.8	2.4	4.4
2003.....	89.2	*	*	*	*	*	*	0.4	0.9	2.0	4.7
2002.....	91.4	*	*	*	*	*	*	0.3	0.8	2.3	4.5
2001.....	94.2	*	*	*	*	*	*	0.3	0.8	2.1	4.3
2000.....	96.3	*	*	*	*	*	*	*	0.8	2.3	4.2
1999.....	99.4	*	*	*	*	*	*	0.4	0.7	2.4	4.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic White, both sexes

2007 5/.....	87.6	*	*	*	*	*	0.3	0.6	1.6	3.3	7.8
2006 5/.....	91.6	*	*	*	*	*	0.2	0.5	1.6	3.2	8.7
2005 5/.....	96.2	*	*	*	*	*	0.2	0.6	1.8	3.7	8.4
2004 5/.....	96.8	*	*	*	*	*	0.2	0.7	1.5	4.1	8.6
2003 5/.....	101.8	*	*	*	*	*	0.2	0.5	1.7	4.0	9.1
2002.....	104.3	*	*	*	*	*	0.2	0.7	1.4	4.3	9.2
2001.....	106.0	*	*	*	*	*	0.2	0.7	1.5	3.9	8.7
2000.....	108.4	*	*	*	*	*	*	0.5	1.5	4.0	8.9
1999.....	111.3	*	*	*	*	*	*	0.6	1.5	4.3	8.6

Non-Hispanic White, male

2007 5/.....	92.7	*	*	*	*	*	0.4	0.8	2.5	5.0	12.2
2006 5/.....	95.9	*	*	*	*	*	*	0.9	2.5	4.8	13.7
2005 5/.....	99.6	*	*	*	*	*	0.3	1.0	2.5	5.6	12.8
2004 5/.....	99.3	*	*	*	*	*	*	1.0	2.3	6.0	13.3
2003 5/.....	103.9	*	*	*	*	*	*	0.7	2.5	6.2	14.0
2002.....	106.2	*	*	*	*	*	*	1.1	2.1	6.6	14.5
2001.....	106.6	*	*	*	*	*	0.4	1.3	2.3	5.9	13.7
2000.....	109.2	*	*	*	*	*	*	0.8	2.3	6.1	14.3
1999.....	111.4	*	*	*	*	*	*	0.7	2.3	6.3	13.8

Non-Hispanic White, female

2007 5/.....	82.7	*	*	*	*	*	*	0.3	0.7	1.6	3.5
2006 5/.....	87.5	*	*	*	*	*	*	*	0.7	1.5	3.8
2005 5/.....	92.9	*	*	*	*	*	*	*	1.1	1.7	4.0
2004 5/.....	94.3	*	*	*	*	*	*	*	0.7	2.2	3.9
2003 5/.....	99.8	*	*	*	*	*	*	*	0.8	1.7	4.2
2002.....	102.4	*	*	*	*	*	*	*	0.7	2.1	4.0
2001.....	105.3	*	*	*	*	*	*	*	0.7	1.9	3.7
2000.....	107.7	*	*	*	*	*	*	*	0.7	2.0	3.5
1999.....	111.1	*	*	*	*	*	*	0.4	0.6	2.2	3.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic Black, both sexes

2007 5/.....	49.4	*	*	*	*	*	*	1.1	2.4	5.0	10.7
2006 5/.....	51.9	*	*	*	*	*	*	0.8	2.2	5.9	10.7
2005 5/.....	53.8	*	*	*	*	*	*	0.7	2.3	6.2	11.9
2004 5/.....	54.6	*	*	*	*	*	*	1.4	2.7	5.9	13.2
2003 5/.....	57.9	*	*	*	*	*	*	1.1	2.5	5.8	13.1
2002.....	57.8	*	*	*	*	*	*	1.0	2.0	6.2	12.9
2001.....	59.7	*	*	*	*	*	*	1.3	2.9	6.0	13.7
2000.....	59.8	*	*	*	*	*	*	0.9	2.8	6.6	14.3
1999.....	61.0	*	*	*	*	*	*	0.9	2.3	6.5	15.1

Non-Hispanic Black, male

2007 5/.....	50.5	*	*	*	*	*	*	1.6	3.8	7.6	15.7
2006 5/.....	52.7	*	*	*	*	*	*	*	3.3	8.0	14.5
2005 5/.....	53.9	*	*	*	*	*	*	*	3.3	9.5	16.3
2004 5/.....	54.2	*	*	*	*	*	*	1.7	4.2	7.8	18.7
2003 5/.....	56.7	*	*	*	*	*	*	*	3.4	8.2	18.5
2002.....	56.1	*	*	*	*	*	*	*	2.4	9.0	17.7
2001.....	58.4	*	*	*	*	*	*	1.9	4.6	9.0	19.4
2000.....	57.3	*	*	*	*	*	*	*	3.9	8.9	19.6
1999.....	58.8	*	*	*	*	*	*	*	3.4	9.3	21.6

Non-Hispanic Black, female

2007 5/.....	48.5	*	*	*	*	*	*	*	*	2.7	6.3
2006 5/.....	51.2	*	*	*	*	*	*	*	*	3.9	7.5
2005 5/.....	53.7	*	*	*	*	*	*	*	*	3.2	7.9
2004 5/.....	54.9	*	*	*	*	*	*	*	*	4.2	8.3
2003 5/.....	59.1	*	*	*	*	*	*	*	1.6	3.7	8.4
2002.....	59.3	*	*	*	*	*	*	*	1.6	3.7	8.7
2001.....	60.9	*	*	*	*	*	*	*	*	3.4	8.7
2000.....	62.1	*	*	*	*	*	*	*	1.9	4.6	9.7
1999.....	63.0	*	*	*	*	*	*	*	*	4.0	9.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

All origins 3/, both sexes

2007.....	57.3	9.3	1.0	0.5	0.6	1.4	2.7	3.8	5.0	7.6	11.3
2006.....	56.9	7.7	0.9	0.4	0.7	1.5	2.7	3.9	5.6	7.4	11.4
2005.....	57.6	8.3	0.9	0.5	0.6	1.7	2.7	3.8	5.4	7.8	11.6
2004.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8	11.5
2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8	12.6
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1	12.4
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0	11.9
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5	12.3
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2	12.5

All origins 3/, male

2007.....	51.9	10.3	0.9	0.6	0.6	1.8	3.3	4.7	6.0	9.5	14.3
2006.....	51.0	8.0	1.0	0.5	0.8	1.8	3.3	4.9	7.0	9.2	14.0
2005.....	51.4	8.9	0.9	0.4	0.7	2.3	3.4	4.9	6.7	9.6	14.8
2004.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4	14.4
2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5	15.6
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6	15.9
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9	14.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1	15.4
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3	15.6

All origins 3/, female

2007.....	62.5	8.3	1.1	0.4	0.6	1.0	2.2	2.9	4.0	5.8	8.3
2006.....	62.6	7.5	0.9	0.4	0.6	1.1	2.0	2.8	4.2	5.6	8.9
2005.....	63.5	7.7	0.8	0.5	0.6	1.1	1.9	2.7	4.1	6.0	8.5
2004.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1	8.6
2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1	9.5
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6	9.0
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0	8.9
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9	9.2
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1	9.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

Hispanic, both sexes

2007	14.3	8.1	1.0	0.5	*	1.1	2.3	2.4	2.9	4.6	6.3
2006	13.7	6.3	0.8	*	0.6	1.1	1.7	2.4	3.3	3.6	6.0
2005	13.9	7.5	0.6	*	0.5	1.2	1.8	2.8	2.7	4.1	6.2
2004	13.2	8.5	1.0	*	*	1.2	1.8	2.0	2.6	4.2	6.5
2003	14.1	8.3	1.0	*	0.5	1.2	1.6	2.3	3.4	4.3	6.3
2002	13.3	9.7	0.7	*	*	1.1	1.9	2.3	2.9	3.9	6.6
2001	13.3	9.2	1.0	*	*	1.2	1.3	2.1	3.2	4.1	5.9
2000	13.6	10.2	1.0	*	0.6	1.5	1.9	2.3	3.3	4.6	6.7
1999	13.8	12.2	1.2	*	*	1.5	1.7	1.9	3.7	5.4	7.4

Hispanic, male

2007	14.0	8.7	*	*	*	1.3	3.1	2.9	3.7	5.9	8.9
2006	13.5	6.9	*	*	*	1.4	2.4	3.2	4.6	4.4	8.0
2005	13.7	8.0	*	*	*	1.8	2.0	3.7	3.8	5.9	8.1
2004	13.0	8.4	1.3	*	*	1.7	2.6	3.1	3.6	5.6	8.9
2003	13.6	9.1	*	*	*	1.4	2.2	2.7	4.1	5.6	8.6
2002	12.9	9.1	*	*	*	1.5	2.6	3.1	3.7	4.6	8.6
2001	12.6	9.6	*	*	*	1.3	1.8	2.6	3.8	5.3	8.0
2000	13.2	9.9	1.3	*	*	2.0	2.2	3.0	4.1	5.4	8.3
1999	13.7	10.3	1.4	*	*	1.5	2.0	2.8	4.9	6.7	10.0

Hispanic, female

2007	14.7	7.5	1.1	*	*	*	1.3	1.7	2.0	3.2	3.4
2006	13.9	5.7	*	*	*	*	*	1.3	1.7	2.6	3.8
2005	14.1	7.0	*	*	*	*	1.7	1.6	1.4	2.1	4.2
2004	13.5	8.6	*	*	*	*	*	*	1.5	2.7	3.9
2003	14.5	7.6	*	*	*	*	*	1.9	2.5	2.8	3.9
2002	13.7	10.3	*	*	*	*	*	1.2	1.9	3.1	4.4
2001	13.9	8.7	*	*	*	*	*	1.4	2.5	2.8	3.6
2000	14.0	10.6	*	*	*	*	1.4	1.5	2.3	3.7	5.0
1999	13.9	14.1	*	*	*	1.4	*	*	2.4	4.0	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

Non-Hispanic 4/, both sexes

2007.....	64.8	9.6	1.0	0.5	0.6	1.4	2.8	4.2	5.5	8.2	12.1
2006.....	64.3	8.2	1.0	0.4	0.7	1.5	2.9	4.2	6.1	8.1	12.2
2005.....	64.8	8.6	1.0	0.6	0.7	1.8	2.8	4.1	6.0	8.5	12.4
2004.....	63.6	10.3	1.1	0.4	0.8	1.6	2.7	3.9	5.5	8.4	12.1
2003.....	65.4	10.9	1.1	0.5	0.7	1.8	2.9	3.9	5.8	8.4	13.3
2002.....	65.7	12.2	1.1	0.5	0.7	1.9	2.4	4.2	5.6	8.7	13.1
2001.....	65.2	11.6	1.5	0.5	0.8	1.5	2.7	4.0	5.4	8.5	12.5
2000.....	65.1	12.4	1.1	0.5	0.8	1.8	2.8	3.8	5.5	7.9	12.8
1999.....	65.6	13.1	1.1	0.5	0.8	2.1	2.8	3.8	5.6	8.5	13.0

Non-Hispanic 4/, male

2007.....	59.0	10.8	0.9	0.5	0.7	1.9	3.3	5.2	6.6	10.2	15.3
2006.....	57.8	8.3	1.0	0.4	0.8	1.9	3.5	5.4	7.6	10.2	14.9
2005.....	58.0	9.2	1.0	0.5	0.7	2.4	3.7	5.1	7.4	10.4	15.8
2004.....	57.0	11.0	1.0	0.3	0.8	2.0	3.4	4.8	6.6	10.1	15.1
2003.....	58.6	12.0	1.0	0.6	0.9	2.2	3.4	4.5	7.0	10.2	16.5
2002.....	58.7	12.8	1.1	0.4	0.9	2.6	3.1	5.0	7.1	10.4	16.7
2001.....	57.8	11.0	1.6	0.4	0.8	1.9	3.3	4.7	6.4	10.6	15.6
2000.....	58.0	12.7	1.2	0.6	0.9	2.1	3.3	4.4	6.6	9.6	16.1
1999.....	58.6	13.8	1.3	0.5	0.8	2.6	3.2	4.3	6.7	10.8	16.2

Non-Hispanic 4/, female

2007.....	70.4	8.4	1.1	0.5	0.6	1.0	2.4	3.1	4.4	6.3	9.1
2006.....	70.4	8.0	0.9	0.4	0.7	1.1	2.3	3.1	4.7	6.1	9.6
2005.....	71.2	7.9	1.0	0.6	0.7	1.2	1.9	3.0	4.6	6.7	9.1
2004.....	69.9	9.6	1.2	0.5	0.7	1.3	2.0	3.0	4.4	6.7	9.2
2003.....	71.8	9.7	1.2	0.5	0.5	1.3	2.4	3.3	4.6	6.7	10.2
2002.....	72.3	11.6	1.0	0.5	0.5	1.2	1.8	3.3	4.1	7.1	9.5
2001.....	72.3	12.2	1.3	0.5	0.7	1.1	2.1	3.3	4.4	6.5	9.5
2000.....	71.8	12.0	1.0	0.4	0.7	1.5	2.3	3.2	4.4	6.2	9.6
1999.....	72.2	12.3	1.0	0.5	0.7	1.5	2.4	3.2	4.5	6.4	9.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

Non-Hispanic White, both sexes

2007 5/.....	71.0	6.6	0.8	0.4	0.5	1.2	2.3	3.3	4.5	6.7	10.2
2006 5/.....	70.4	5.8	0.8	0.4	0.6	1.3	2.5	3.4	4.8	6.4	10.5
2005 5/.....	70.6	6.7	0.7	0.4	0.5	1.5	2.4	3.2	4.6	6.9	10.5
2004 5/.....	69.3	8.7	0.8	0.3	0.6	1.3	2.3	3.1	4.4	6.9	10.1
2003 5/.....	71.0	8.1	0.8	0.4	0.6	1.5	2.3	3.0	4.6	6.8	11.1
2002.....	71.3	10.0	0.9	0.4	0.6	1.7	1.9	3.2	4.3	7.1	11.3
2001.....	70.6	9.1	1.1	0.4	0.7	1.3	2.1	3.1	4.3	6.8	10.1
2000.....	70.3	10.5	0.9	0.4	0.6	1.4	2.1	2.9	4.4	6.2	10.6
1999.....	70.4	10.5	0.9	0.4	0.6	1.8	2.2	2.9	4.4	6.8	10.5

Non-Hispanic White, male

2007 5/.....	63.4	7.4	0.6	0.4	0.6	1.5	2.5	4.1	5.5	8.3	12.5
2006 5/.....	62.1	5.7	0.8	0.4	0.7	1.6	3.1	4.3	5.9	8.0	12.8
2005 5/.....	62.2	6.9	0.8	0.4	0.6	1.9	3.2	4.2	5.6	8.6	13.6
2004 5/.....	60.9	9.3	0.7	*	0.7	1.5	3.0	3.7	5.4	8.4	12.9
2003 5/.....	62.6	8.7	0.7	0.4	0.8	1.9	2.7	3.4	5.7	8.3	14.2
2002.....	62.7	10.7	1.1	0.4	0.8	2.2	2.4	3.8	5.5	8.5	14.8
2001.....	61.4	9.4	1.2	0.3	0.8	1.5	2.6	3.7	5.0	8.5	13.2
2000.....	61.6	10.9	1.1	0.5	0.7	1.6	2.5	3.4	5.4	7.8	13.6
1999.....	61.7	11.6	1.1	0.5	0.6	2.2	2.5	3.3	5.4	8.7	13.6

Non-Hispanic White, female

2007 5/.....	78.3	5.8	1.0	0.4	0.5	0.9	2.1	2.4	3.5	5.1	7.9
2006 5/.....	78.4	5.8	0.8	0.4	0.5	0.9	1.8	2.5	3.6	4.8	8.3
2005 5/.....	78.8	6.4	0.6	0.4	0.5	1.1	1.5	2.2	3.5	5.2	7.5
2004 5/.....	77.3	8.2	0.9	*	0.5	1.0	1.6	2.4	3.4	5.3	7.3
2003 5/.....	79.1	7.5	1.0	0.4	0.4	1.1	2.0	2.5	3.4	5.3	8.0
2002.....	79.6	9.3	0.8	0.5	0.5	1.1	1.3	2.5	3.1	5.7	7.8
2001.....	79.4	8.8	1.0	0.5	0.6	1.0	1.6	2.6	3.5	5.1	7.0
2000.....	78.7	10.1	0.8	*	0.5	1.2	1.6	2.4	3.4	4.7	7.6
1999.....	78.8	9.4	0.8	0.4	0.6	1.4	2.0	2.4	3.5	4.9	7.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

Non-Hispanic Black, both sexes

2007 5/.....	53.0	20.0	1.6	0.7	1.2	2.7	5.5	8.9	11.8	18.7	25.5
2006 5/.....	52.0	16.8	1.7	*	1.2	2.9	5.1	9.0	14.8	19.5	24.8
2005 5/.....	53.4	15.6	1.8	1.1	1.2	3.3	5.4	9.0	14.5	19.3	26.2
2004 5/.....	52.7	15.3	2.4	0.8	1.4	3.1	4.6	8.4	12.4	18.8	27.1
2003 5/.....	54.4	19.8	2.0	1.0	1.1	3.1	6.1	9.3	13.2	18.9	29.7
2002.....	54.3	20.9	1.6	*	1.2	3.1	5.3	10.0	13.4	19.7	27.1
2001.....	54.7	21.3	3.0	0.8	1.2	3.0	5.8	9.1	12.8	19.1	30.0
2000.....	54.6	19.6	2.0	0.8	1.4	3.8	6.3	9.1	12.4	18.5	29.1
1999.....	56.2	23.2	2.0	0.8	1.3	3.7	5.9	9.0	13.0	20.1	30.8

Non-Hispanic Black, male

2007 5/.....	53.1	23.2	2.1	*	*	3.8	7.1	10.8	13.7	23.9	34.3
2006 5/.....	52.1	17.9	2.0	*	1.4	3.6	5.5	11.4	18.7	25.0	30.9
2005 5/.....	52.6	18.4	*	*	1.3	4.8	6.9	11.0	18.4	23.5	33.0
2004 5/.....	52.3	16.4	2.5	*	1.3	4.0	5.7	10.8	14.4	22.6	32.5
2003 5/.....	53.4	23.6	2.2	*	1.3	4.0	7.7	11.2	14.8	23.5	35.6
2002.....	52.6	21.8	*	*	1.4	4.0	6.4	12.5	17.0	23.6	32.8
2001.....	53.5	18.8	3.3	*	1.3	3.9	7.3	11.1	15.5	24.1	34.6
2000.....	53.3	19.1	2.0	*	1.5	4.5	7.1	10.5	14.9	22.4	35.3
1999.....	56.2	22.9	2.3	*	1.5	5.1	7.1	10.8	15.9	25.1	37.0

Non-Hispanic Black, female

2007 5/.....	52.8	16.7	*	*	1.3	1.6	3.8	7.1	10.0	14.1	17.8
2006 5/.....	51.9	15.8	*	*	*	2.1	4.6	6.7	11.4	14.6	19.4
2005 5/.....	54.2	12.7	2.1	*	*	1.8	3.9	7.1	11.0	15.6	20.2
2004 5/.....	53.1	14.2	2.3	*	1.4	2.1	3.5	6.1	10.6	15.4	22.3
2003 5/.....	55.4	15.8	1.8	*	*	2.1	4.5	7.6	11.8	14.9	24.5
2002.....	55.9	19.9	1.9	*	*	2.3	4.2	7.7	10.3	16.1	22.2
2001.....	55.8	23.8	2.7	*	*	2.1	4.3	7.2	10.4	14.7	26.0
2000.....	55.8	20.1	1.9	*	1.3	3.1	5.5	7.8	10.2	15.0	23.6
1999.....	56.3	23.6	1.8	*	*	2.2	4.7	7.4	10.4	15.7	25.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All origins 3/, both sexes

2007.....	0.3	0.6	*	*	*	*	0.1	0.1	0.1	0.2	0.2
2006.....	0.3	*	0.1	*	*	*	0.1	0.1	*	0.2	0.2
2005.....	0.3	0.5	*	*	*	*	*	0.1	0.1	0.2	0.2
2004.....	0.3	0.5	0.2	*	*	*	*	*	0.1	0.1	0.2
2003.....	0.3	0.8	0.1	*	*	0.1	0.2	0.1	0.1	0.1	0.2
2002.....	0.3	0.6	0.1	*	*	*	*	*	0.1	0.2	0.2
2001.....	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2	0.2
2000.....	0.3	0.5	0.2	*	*	*	0.1	0.1	0.1	0.2	0.2
1999.....	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2	0.3

All origins 3/, male

2007.....	0.3	*	*	*	*	*	*	*	*	0.3	0.2
2006.....	0.3	*	*	*	*	*	*	*	*	0.2	0.2
2005.....	0.3	*	*	*	*	*	*	*	*	*	0.3
2004.....	0.3	*	*	*	*	*	*	*	*	0.2	0.2
2003.....	0.3	*	*	*	*	*	0.2	*	*	0.2	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.2	*	0.3
2001.....	0.3	*	*	*	*	*	*	*	*	0.2	*
2000.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.3	0.3

All origins 3/, female

2007.....	0.3	*	*	*	*	*	*	*	*	*	*
2006.....	0.3	*	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	0.2	*
2004.....	0.3	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	0.2
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	0.3
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	0.2	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.7	*	*	*	*	0.1	0.1	0.1	0.2	0.2
2006.....	0.3	*	*	*	*	*	0.1	0.2	*	0.2	0.2
2005.....	0.3	*	*	*	*	*	*	0.1	0.1	0.2	0.2
2004.....	0.3	*	*	*	*	*	*	*	0.1	0.2	0.2
2003.....	0.3	0.9	*	*	*	*	0.2	*	0.1	0.2	0.3
2002.....	0.3	0.7	0.2	*	*	*	*	*	0.2	0.2	0.2
2001.....	0.3	0.9	0.2	*	*	*	0.1	*	0.1	0.2	0.2
2000.....	0.3	*	*	*	*	*	*	0.1	0.1	0.2	0.2
1999.....	0.3	0.8	0.2	*	*	*	*	*	0.1	0.2	0.3

Non-Hispanic 4/, male

2007.....	0.3	*	*	*	*	*	*	*	*	0.3	0.2
2006.....	0.3	*	*	*	*	*	*	*	*	0.3	0.2
2005.....	0.3	*	*	*	*	*	*	*	*	*	0.3
2004.....	0.3	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.2	*	0.3
2001.....	0.3	*	*	*	*	*	*	*	*	0.2	*
2000.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3
1999.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3

Non-Hispanic 4/, female

2007.....	0.3	*	*	*	*	*	*	*	*	*	*
2006.....	0.3	*	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	0.2
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	0.3
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	0.2	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Heart failure (I50)

All origins 3/, both sexes

2007.....	18.8	0.5	*	*	*	*	0.1	0.1	0.3	0.5	1.0
2006.....	20.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0
2005.....	19.9	0.7	*	*	*	*	0.1	0.2	0.3	0.5	1.0
2004.....	19.5	0.7	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2003.....	19.8	0.9	*	*	*	*	0.1	0.1	0.3	0.4	1.0
2002.....	19.6	0.6	*	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	20.0	0.6	*	*	*	*	*	0.2	0.2	0.5	0.8
2000.....	19.8	0.8	*	*	*	0.1	*	0.1	0.2	0.3	0.8
1999.....	19.7	*	*	*	*	*	*	0.1	0.2	0.4	0.8

All origins 3/, male

2007.....	15.4	*	*	*	*	*	*	*	0.4	0.6	1.3
2006.....	16.2	*	*	*	*	*	*	*	0.4	0.6	1.2
2005.....	15.8	*	*	*	*	*	*	0.2	0.4	0.6	1.3
2004.....	15.4	*	*	*	*	*	*	0.3	0.3	0.7	1.0
2003.....	15.7	1.4	*	*	*	*	*	*	0.4	0.5	1.3
2002.....	15.3	*	*	*	*	*	*	0.2	0.4	0.5	1.0
2001.....	15.5	*	*	*	*	*	*	0.3	0.3	0.7	0.9
2000.....	15.3	1.1	*	*	*	*	*	*	0.2	0.4	1.1
1999.....	15.2	*	*	*	*	*	*	*	*	0.5	1.0

All origins 3/, female

2007.....	22.0	*	*	*	*	*	*	*	0.2	0.4	0.6
2006.....	24.0	*	*	*	*	*	*	*	0.2	0.4	0.8
2005.....	23.9	*	*	*	*	*	*	*	0.2	0.3	0.6
2004.....	23.4	*	*	*	*	*	*	*	0.2	0.4	0.7
2003.....	23.7	*	*	*	*	*	*	*	0.2	0.4	0.7
2002.....	23.7	*	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	24.3	*	*	*	*	*	*	*	0.2	0.4	0.7
2000.....	24.1	*	*	*	*	*	*	*	*	0.3	0.6
1999.....	24.0	*	*	*	*	*	*	*	*	0.3	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Heart failure (I50)

Non-Hispanic 4/, both sexes

2007.....	21.3	*	*	*	*	*	0.2	0.1	0.3	0.5	1.0
2006.....	22.9	*	*	*	*	*	0.1	0.2	0.3	0.6	1.1
2005.....	22.5	0.7	*	*	*	*	0.2	0.2	0.3	0.5	1.0
2004.....	22.0	0.8	*	*	*	*	0.1	0.2	0.3	0.6	0.9
2003.....	22.2	1.0	*	*	*	*	*	0.2	0.4	0.5	1.1
2002.....	22.0	*	*	*	*	*	*	0.2	0.4	0.5	0.8
2001.....	22.4	*	*	*	*	*	*	0.2	0.3	0.5	0.9
2000.....	22.1	0.8	*	*	*	0.1	*	0.2	0.2	0.4	0.9
1999.....	21.8	*	*	*	*	*	*	0.2	0.2	0.4	0.9

Non-Hispanic 4/, male

2007.....	17.6	*	*	*	*	*	*	*	0.5	0.7	1.4
2006.....	18.5	*	*	*	*	*	*	*	0.5	0.7	1.3
2005.....	18.0	*	*	*	*	*	*	*	0.5	0.7	1.4
2004.....	17.5	*	*	*	*	*	*	0.3	0.4	0.7	1.1
2003.....	17.7	1.4	*	*	*	*	*	*	0.5	0.5	1.3
2002.....	17.3	*	*	*	*	*	*	0.3	0.4	0.6	1.1
2001.....	17.4	*	*	*	*	*	*	0.3	0.3	0.7	1.0
2000.....	17.2	*	*	*	*	*	*	*	0.2	0.4	1.1
1999.....	16.9	*	*	*	*	*	*	*	*	0.5	1.1

Non-Hispanic 4/, female

2007.....	24.9	*	*	*	*	*	*	*	*	0.4	0.7
2006.....	27.1	*	*	*	*	*	*	*	*	0.5	0.8
2005.....	26.9	*	*	*	*	*	*	*	*	0.4	0.7
2004.....	26.3	*	*	*	*	*	*	*	0.2	0.4	0.8
2003.....	26.5	*	*	*	*	*	*	*	0.3	0.4	0.8
2002.....	26.5	*	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	27.1	*	*	*	*	*	*	*	*	0.4	0.8
2000.....	26.7	*	*	*	*	*	*	*	*	0.3	0.6
1999.....	26.5	*	*	*	*	*	*	*	*	0.3	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Heart failure (I50)

Non-Hispanic White, both sexes

2007 5/.....	24.1	*	*	*	*	*	*	*	0.2	0.4	0.7
2006 5/.....	26.0	*	*	*	*	*	*	*	0.2	0.4	0.8
2005 5/.....	25.4	*	*	*	*	*	*	*	0.2	0.4	0.7
2004 5/.....	24.8	*	*	*	*	*	*	*	0.2	0.4	0.7
2003 5/.....	25.0	*	*	*	*	*	*	*	0.3	0.4	0.7
2002.....	24.7	*	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	25.1	*	*	*	*	*	*	0.2	0.2	0.4	0.6
2000.....	24.7	*	*	*	*	*	*	*	0.2	0.3	0.7
1999.....	24.3	*	*	*	*	*	*	*	*	0.3	0.7

Non-Hispanic White, male

2007 5/.....	19.7	*	*	*	*	*	*	*	*	0.5	0.9
2006 5/.....	20.7	*	*	*	*	*	*	*	*	0.4	0.9
2005 5/.....	20.0	*	*	*	*	*	*	*	*	0.5	0.9
2004 5/.....	19.5	*	*	*	*	*	*	*	*	0.6	0.9
2003 5/.....	19.7	*	*	*	*	*	*	*	0.3	0.4	0.9
2002.....	19.2	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	19.3	*	*	*	*	*	*	*	*	0.5	0.8
2000.....	19.0	*	*	*	*	*	*	*	*	0.3	1.0
1999.....	18.6	*	*	*	*	*	*	*	*	0.4	0.9

Non-Hispanic White, female

2007 5/.....	28.4	*	*	*	*	*	*	*	*	*	0.5
2006 5/.....	31.0	*	*	*	*	*	*	*	*	0.4	0.7
2005 5/.....	30.5	*	*	*	*	*	*	*	*	*	0.6
2004 5/.....	30.0	*	*	*	*	*	*	*	*	*	0.6
2003 5/.....	30.1	*	*	*	*	*	*	*	*	0.3	0.4
2002.....	30.0	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	30.7	*	*	*	*	*	*	*	*	0.3	0.5
2000.....	30.2	*	*	*	*	*	*	*	*	*	0.4
1999.....	29.9	*	*	*	*	*	*	*	*	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes

2007.....	37.8	8.2	0.8	0.4	0.5	1.2	2.4	3.5	4.5	6.7	9.9
2006.....	36.0	6.9	0.7	0.3	0.6	1.3	2.4	3.5	5.1	6.6	10.0
2005.....	37.0	7.1	0.7	0.4	0.5	1.5	2.4	3.5	4.9	7.0	10.3
2004.....	36.5	8.7	0.9	0.3	0.6	1.4	2.3	3.2	4.5	6.9	10.2
2003.....	38.0	8.6	0.9	0.4	0.6	1.5	2.4	3.3	4.9	7.0	11.0
2002.....	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3	11.2
2001.....	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0	10.5
2000.....	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8	10.9
1999.....	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4	11.1

All origins 3/, male

2007.....	35.7	9.0	0.7	0.4	0.5	1.6	2.9	4.3	5.3	8.4	12.3
2006.....	34.0	6.9	0.8	0.4	0.7	1.7	3.0	4.6	6.3	8.2	12.3
2005.....	34.9	7.7	0.8	0.4	0.6	2.1	3.1	4.4	6.0	8.7	12.9
2004.....	34.5	9.2	0.9	0.2	0.7	1.8	3.0	4.1	5.5	8.3	12.9
2003.....	35.8	9.2	0.9	0.4	0.7	1.8	2.9	3.7	5.8	8.6	13.8
2002.....	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8	14.2
2001.....	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8	13.4
2000.....	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3	13.6
1999.....	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3	13.9

All origins 3/, female

2007.....	39.9	7.3	0.9	0.3	0.5	0.9	1.8	2.6	3.6	5.1	7.4
2006.....	38.0	6.8	0.7	0.3	0.5	1.0	1.8	2.5	3.8	4.9	7.8
2005.....	39.0	6.4	0.7	0.4	0.4	1.0	1.7	2.4	3.7	5.3	7.6
2004.....	38.5	8.2	0.8	0.4	0.6	1.0	1.6	2.4	3.5	5.5	7.6
2003.....	40.1	7.9	1.0	0.3	0.4	1.1	1.8	2.8	3.9	5.5	8.4
2002.....	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9	8.2
2001.....	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3	7.6
2000.....	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3	8.2
1999.....	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4	8.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Hispanic, both sexes

2007.....	9.8	7.4	0.9	*	*	0.9	2.0	2.2	2.6	4.1	5.2
2006.....	9.2	5.3	0.6	*	*	0.9	1.5	2.2	2.9	3.3	5.3
2005.....	9.5	6.4	*	*	*	1.1	1.7	2.5	2.4	3.8	5.3
2004.....	9.1	7.2	0.7	*	*	1.0	1.7	1.8	2.5	3.5	5.7
2003.....	9.7	6.8	0.9	*	*	1.0	1.4	2.1	3.1	4.0	5.7
2002.....	9.3	8.6	0.6	*	*	1.0	1.8	2.2	2.6	3.6	6.0
2001.....	9.3	8.0	0.8	*	*	1.2	1.2	1.8	2.7	3.6	5.1
2000.....	9.6	8.7	0.7	*	0.6	1.4	1.8	2.1	3.0	4.1	5.9
1999.....	9.8	11.5	0.9	*	*	1.3	1.4	1.8	3.3	4.7	6.5

Hispanic, male

2007.....	10.0	7.9	*	*	*	1.0	2.8	2.7	3.2	5.3	7.1
2006.....	9.6	5.8	*	*	*	1.2	2.1	3.1	4.2	4.1	7.1
2005.....	10.0	6.9	*	*	*	1.6	1.8	3.4	3.3	5.5	6.8
2004.....	9.4	6.9	*	*	*	1.5	2.5	2.8	3.4	4.6	8.1
2003.....	9.8	7.0	*	*	*	*	2.0	2.4	3.6	5.3	7.6
2002.....	9.6	8.0	*	*	*	1.4	2.5	2.9	3.3	4.4	7.9
2001.....	9.3	7.9	*	*	*	1.3	1.6	2.2	3.3	4.4	7.3
2000.....	9.9	8.1	*	*	*	2.0	2.2	2.8	3.8	4.9	7.3
1999.....	10.3	9.8	*	*	*	1.3	1.7	2.5	4.3	5.8	8.9

Hispanic, female

2007.....	9.7	6.7	1.1	*	*	*	*	1.6	1.8	2.8	3.0
2006.....	8.8	4.6	*	*	*	*	*	1.1	1.4	2.3	3.3
2005.....	9.0	5.9	*	*	*	*	1.5	1.5	1.3	1.8	3.7
2004.....	8.8	7.4	*	*	*	*	*	*	1.3	2.3	3.0
2003.....	9.6	6.6	*	*	*	*	*	1.6	2.4	2.6	3.6
2002.....	9.0	9.3	*	*	*	*	*	1.2	1.8	2.7	4.0
2001.....	9.2	8.0	*	*	*	*	*	1.3	1.9	2.7	2.9
2000.....	9.3	9.3	*	*	*	*	1.4	*	2.1	3.2	4.5
1999.....	9.2	13.3	*	*	*	*	*	*	2.1	3.5	4.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 4/, both sexes

2007.....	42.7	8.3	0.8	0.4	0.6	1.3	2.5	3.8	4.9	7.3	10.7
2006.....	40.6	7.3	0.8	0.3	0.7	1.4	2.6	3.9	5.6	7.2	10.7
2005.....	41.5	7.2	0.8	0.4	0.5	1.6	2.6	3.7	5.4	7.6	11.0
2004.....	40.9	9.1	0.9	0.3	0.7	1.5	2.4	3.6	4.9	7.6	10.8
2003.....	42.4	9.0	0.9	0.4	0.6	1.6	2.6	3.5	5.2	7.5	11.7
2002.....	42.9	10.9	0.8	0.4	0.6	1.7	2.2	3.8	4.9	7.9	11.8
2001.....	42.1	10.1	1.2	0.3	0.7	1.3	2.4	3.6	4.9	7.5	11.1
2000.....	42.3	11.0	0.9	0.4	0.6	1.6	2.5	3.4	5.0	7.2	11.4
1999.....	43.0	11.9	0.9	0.4	0.7	1.9	2.6	3.4	5.2	7.7	11.6

Non-Hispanic 4/, male

2007.....	40.5	9.3	0.7	0.4	0.6	1.7	3.0	4.7	5.8	9.0	13.3
2006.....	38.5	7.2	0.8	0.4	0.8	1.8	3.2	5.0	6.9	9.0	13.1
2005.....	39.2	7.9	0.8	0.4	0.6	2.2	3.4	4.7	6.7	9.3	13.9
2004.....	38.8	9.7	0.9	*	0.7	1.8	3.1	4.4	6.0	9.0	13.6
2003.....	40.0	9.9	0.8	0.5	0.8	1.9	3.1	4.1	6.3	9.2	14.5
2002.....	40.6	11.6	0.8	0.3	0.8	2.3	2.8	4.6	6.3	9.5	14.9
2001.....	39.6	9.3	1.4	0.3	0.7	1.7	3.1	4.2	5.8	9.4	14.1
2000.....	40.0	10.9	1.0	0.5	0.7	1.9	3.0	3.9	6.1	8.8	14.3
1999.....	40.9	12.7	1.0	0.4	0.8	2.4	2.9	3.9	6.2	9.7	14.4

Non-Hispanic 4/, female

2007.....	44.9	7.4	0.8	0.4	0.6	0.9	2.0	2.8	4.0	5.5	8.1
2006.....	42.7	7.4	0.8	0.3	0.6	1.0	2.0	2.8	4.4	5.4	8.4
2005.....	43.7	6.5	0.8	0.5	0.5	1.1	1.7	2.7	4.2	5.9	8.1
2004.....	43.0	8.3	0.9	0.4	0.6	1.1	1.8	2.8	3.9	6.1	8.2
2003.....	44.6	8.1	1.0	0.3	0.4	1.2	2.1	3.0	4.2	5.9	8.9
2002.....	45.1	10.1	0.8	0.4	0.4	1.1	1.6	3.0	3.7	6.4	8.7
2001.....	44.5	11.0	1.0	0.4	0.6	1.0	1.8	3.0	4.0	5.7	8.1
2000.....	44.5	11.0	0.8	0.2	0.5	1.3	2.1	2.9	4.0	5.6	8.6
1999.....	44.9	11.1	0.9	0.4	0.6	1.5	2.2	3.0	4.2	5.7	8.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic White, both sexes

2007 5/.....	46.1	5.9	0.6	0.4	0.4	1.1	2.0	2.9	4.0	6.0	9.1
2006 5/.....	43.7	5.4	0.6	0.3	0.6	1.2	2.2	3.1	4.4	5.7	9.3
2005 5/.....	44.5	5.9	0.7	0.3	0.5	1.4	2.1	2.9	4.2	6.3	9.5
2004 5/.....	43.7	7.9	0.7	0.3	0.6	1.2	2.1	2.8	4.0	6.2	9.1
2003 5/.....	45.2	7.1	0.7	0.3	0.6	1.3	2.0	2.7	4.2	6.2	9.9
2002.....	45.8	9.2	0.7	0.3	0.6	1.5	1.7	2.8	3.8	6.5	10.2
2001.....	44.7	8.1	0.9	0.3	0.6	1.1	1.9	2.8	3.9	6.1	9.1
2000.....	44.9	9.6	0.8	0.3	0.5	1.2	1.9	2.6	4.1	5.7	9.5
1999.....	45.4	9.8	0.8	0.4	0.6	1.7	2.1	2.6	4.1	6.1	9.5

Non-Hispanic White, male

2007 5/.....	42.9	6.5	*	*	0.5	1.3	2.3	3.7	4.9	7.3	11.1
2006 5/.....	40.5	5.2	0.6	*	0.7	1.5	2.9	4.0	5.4	7.2	11.4
2005 5/.....	41.4	6.2	0.7	0.4	0.5	1.7	2.9	3.9	5.1	7.8	12.3
2004 5/.....	40.6	8.4	0.7	*	0.6	1.4	2.7	3.4	4.9	7.5	11.7
2003 5/.....	42.0	7.5	0.6	0.4	0.7	1.7	2.3	3.0	5.3	7.6	12.6
2002.....	42.7	9.8	0.8	0.3	0.7	2.1	2.2	3.4	4.8	7.8	13.2
2001.....	41.3	8.1	1.0	*	0.7	1.3	2.4	3.4	4.5	7.7	12.0
2000.....	41.7	9.6	0.8	0.4	0.6	1.4	2.3	3.1	5.0	7.1	12.0
1999.....	42.4	10.8	0.9	0.4	0.6	2.0	2.3	3.1	5.0	7.9	12.2

Non-Hispanic White, female

2007 5/.....	49.3	5.3	0.7	0.4	0.4	0.8	1.7	2.2	3.2	4.6	7.2
2006 5/.....	46.7	5.6	0.6	*	0.5	0.9	1.5	2.2	3.4	4.2	7.2
2005 5/.....	47.6	5.5	0.6	*	0.4	1.0	1.4	2.0	3.2	4.7	6.6
2004 5/.....	46.7	7.3	0.7	*	0.5	1.0	1.5	2.2	3.0	5.0	6.5
2003 5/.....	48.3	6.7	0.8	*	0.4	1.0	1.7	2.3	3.1	4.8	7.2
2002.....	48.9	8.5	0.6	0.4	0.4	1.0	1.1	2.3	2.7	5.1	7.1
2001.....	47.9	8.1	0.7	*	0.5	0.8	1.4	2.2	3.2	4.4	6.1
2000.....	47.9	9.6	0.7	*	0.3	1.0	1.5	2.1	3.1	4.3	6.9
1999.....	48.2	8.7	0.6	0.4	0.6	1.4	1.9	2.2	3.3	4.3	6.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic Black, both sexes

2007 5/.....	37.7	17.0	1.3	*	1.2	2.5	4.9	8.3	10.3	16.3	21.5
2006 5/.....	36.5	14.3	1.5	*	1.1	2.6	4.6	8.2	13.2	17.0	21.3
2005 5/.....	37.4	12.7	1.2	0.9	0.9	3.1	4.9	8.0	13.0	16.7	22.4
2004 5/.....	37.7	12.8	1.9	*	1.2	2.8	4.2	7.6	11.1	16.4	24.0
2003 5/.....	38.8	15.2	1.6	0.8	0.9	2.6	5.4	8.4	11.6	16.6	25.0
2002.....	39.1	17.8	1.2	*	1.0	2.9	4.9	9.1	11.8	17.7	24.0
2001.....	39.3	18.2	2.4	*	1.1	2.7	5.1	8.3	11.7	17.0	25.8
2000.....	39.5	16.1	1.5	0.7	1.1	3.4	5.7	8.3	11.3	16.5	25.7
1999.....	40.9	20.9	1.5	0.6	1.1	3.4	5.4	8.2	11.9	18.0	26.9

Non-Hispanic Black, male

2007 5/.....	39.1	19.6	1.7	*	*	3.5	6.2	10.3	11.8	21.0	28.4
2006 5/.....	38.2	14.8	1.7	*	1.3	3.2	4.9	10.7	16.5	21.6	26.5
2005 5/.....	38.8	15.2	*	*	*	4.4	6.5	9.7	16.4	20.4	27.2
2004 5/.....	39.0	13.9	1.9	*	*	3.8	5.3	9.8	12.7	20.2	28.6
2003 5/.....	39.9	18.3	1.9	*	*	3.4	6.9	10.2	12.7	20.7	30.3
2002.....	39.6	19.7	*	*	1.2	3.6	5.9	11.5	14.9	21.3	28.8
2001.....	40.3	16.0	2.7	*	1.2	3.5	6.6	9.7	14.1	21.1	30.8
2000.....	40.5	15.0	1.7	*	1.2	4.0	6.4	9.5	13.7	20.0	31.2
1999.....	42.8	20.8	*	*	1.5	4.8	6.5	9.7	14.5	22.5	32.6

Non-Hispanic Black, female

2007 5/.....	36.4	14.2	*	*	1.3	1.4	3.5	6.4	9.0	12.1	15.4
2006 5/.....	34.9	13.8	*	*	*	2.0	4.2	5.8	10.3	12.9	16.8
2005 5/.....	36.2	10.1	*	*	*	1.6	3.3	6.4	10.0	13.5	18.1
2004 5/.....	36.5	11.6	1.8	*	1.3	1.7	3.2	5.6	9.8	13.0	19.9
2003 5/.....	37.8	11.9	*	*	*	1.9	3.8	6.8	10.7	12.9	20.4
2002.....	38.7	15.8	*	*	*	2.1	4.0	7.0	9.1	14.5	19.8
2001.....	38.5	20.5	2.1	*	*	1.8	3.7	6.9	9.6	13.3	21.4
2000.....	38.6	17.3	*	*	*	2.7	5.1	7.1	9.0	13.4	20.9
1999.....	39.2	21.1	*	*	*	2.0	4.4	6.8	9.6	14.1	22.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic 4/, both sexes

2007.....	8.8	*	*	*	*	*	*	0.2	0.3	0.6	1.2
2006.....	8.8	*	*	*	*	*	*	0.1	0.5	0.6	1.3
2005.....	9.3	*	*	*	*	*	*	0.2	0.3	0.6	1.3
2004.....	8.7	*	*	*	*	*	*	0.2	0.4	0.7	1.0
2003.....	8.3	*	*	*	*	*	*	0.1	0.4	0.7	1.1
2002.....	7.7	*	*	*	*	*	*	0.2	0.3	0.6	1.0
2001.....	7.4	*	*	*	*	*	*	0.2	0.4	0.5	1.1
2000.....	7.0	*	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	6.6	*	*	*	*	*	*	0.1	0.3	0.5	1.0

Non-Hispanic 4/, male

2007.....	7.0	*	*	*	*	*	*	*	0.4	0.8	1.7
2006.....	7.1	*	*	*	*	*	*	*	0.6	0.8	1.6
2005.....	7.2	*	*	*	*	*	*	*	0.4	0.7	1.6
2004.....	6.8	*	*	*	*	*	*	0.3	0.4	0.9	1.3
2003.....	6.3	*	*	*	*	*	*	*	0.5	0.8	1.3
2002.....	5.9	*	*	*	*	*	*	*	0.3	0.7	1.3
2001.....	5.7	*	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	5.4	*	*	*	*	*	*	*	0.3	0.5	1.4
1999.....	5.1	*	*	*	*	*	*	*	0.4	0.7	1.2

Non-Hispanic 4/, female

2007.....	10.5	*	*	*	*	*	*	*	0.3	0.5	0.8
2006.....	10.5	*	*	*	*	*	*	*	0.4	0.4	1.0
2005.....	11.3	*	*	*	*	*	*	*	*	0.5	1.1
2004.....	10.5	*	*	*	*	*	*	*	0.3	0.4	0.8
2003.....	10.2	*	*	*	*	*	*	*	*	0.6	1.0
2002.....	9.4	*	*	*	*	*	*	*	0.2	0.5	0.7
2001.....	9.0	*	*	*	*	*	*	*	0.4	0.4	0.8
2000.....	8.5	*	*	*	*	*	*	*	0.2	0.4	0.8
1999.....	8.0	*	*	*	*	*	*	*	*	0.4	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic Black, both sexes

2007 5/.....	12.1	*	*	*	*	*	*	*	1.4	2.4	4.0
2006 5/.....	12.8	*	*	*	*	*	*	*	2.0	2.3	5.3
2005 5/.....	13.1	*	*	*	*	*	*	*	1.4	2.3	4.8
2004 5/.....	12.8	*	*	*	*	*	*	0.8	1.3	2.7	3.7
2003 5/.....	12.2	*	*	*	*	*	*	*	1.4	2.3	4.0
2002.....	11.5	*	*	*	*	*	*	*	1.0	2.8	3.9
2001.....	11.3	*	*	*	*	*	*	*	1.4	1.9	4.5
2000.....	11.0	*	*	*	*	*	*	*	1.1	2.1	4.2
1999.....	10.7	*	*	*	*	*	*	*	1.0	2.3	4.1

Non-Hispanic Black, male

2007 5/.....	10.8	*	*	*	*	*	*	*	*	2.5	5.4
2006 5/.....	11.7	*	*	*	*	*	*	*	2.2	3.1	6.7
2005 5/.....	11.8	*	*	*	*	*	*	*	1.7	2.5	5.5
2004 5/.....	11.1	*	*	*	*	*	*	*	*	3.6	4.4
2003 5/.....	10.8	*	*	*	*	*	*	*	2.2	2.4	4.6
2002.....	9.9	*	*	*	*	*	*	*	*	3.7	4.9
2001.....	10.0	*	*	*	*	*	*	*	*	2.3	5.5
2000.....	9.5	*	*	*	*	*	*	*	*	2.2	5.4
1999.....	9.4	*	*	*	*	*	*	*	*	2.8	4.8

Non-Hispanic Black, female

2007 5/.....	13.2	*	*	*	*	*	*	*	*	2.4	2.7
2006 5/.....	13.9	*	*	*	*	*	*	*	1.7	1.6	4.0
2005 5/.....	14.3	*	*	*	*	*	*	*	*	2.1	4.1
2004 5/.....	14.3	*	*	*	*	*	*	*	*	1.9	3.1
2003 5/.....	13.4	*	*	*	*	*	*	*	*	2.2	3.4
2002.....	13.0	*	*	*	*	*	*	*	*	2.1	3.1
2001.....	12.4	*	*	*	*	*	*	*	1.4	1.5	3.5
2000.....	12.4	*	*	*	*	*	*	*	*	2.1	3.2
1999.....	11.8	*	*	*	*	*	*	*	*	1.9	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes

2007.....	45.1	3.1	0.3	0.2	0.2	0.3	0.6	0.9	1.7	3.5	6.4
2006.....	45.8	3.4	0.3	0.2	0.2	0.3	0.6	0.9	1.7	3.4	6.7
2005.....	48.4	3.1	0.4	0.3	0.2	0.4	0.6	1.0	1.8	3.3	6.9
2004.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4	7.1
2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6	7.4
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6	7.1
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8	7.2
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9	7.7
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0	7.5

All origins 3/, male

2007.....	36.4	3.5	0.2	*	0.2	0.4	0.7	0.8	1.6	3.8	6.8
2006.....	37.0	3.9	0.3	0.3	0.3	0.3	0.7	1.0	1.8	3.7	6.9
2005.....	38.8	3.5	0.5	0.3	0.3	0.3	0.6	1.1	1.9	3.4	7.0
2004.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7	7.4
2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8	7.6
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6	6.9
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9	7.4
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8	7.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9	7.8

All origins 3/, female

2007.....	53.5	2.6	0.4	0.2	0.3	0.3	0.5	0.9	1.7	3.2	6.0
2006.....	54.4	2.9	0.4	*	0.2	0.4	0.6	0.8	1.6	3.1	6.5
2005.....	57.8	2.6	0.3	0.3	*	0.4	0.6	0.8	1.6	3.2	6.7
2004.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1	6.8
2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3	7.2
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7	7.2
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6	7.1
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0	7.5
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1	7.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

Hispanic, both sexes

2007.....	15.6	2.8	*	*	*	*	0.7	0.9	1.6	3.2	5.7
2006.....	15.8	3.4	*	*	*	0.6	0.9	0.9	1.4	2.8	6.7
2005.....	16.0	2.6	*	*	*	*	0.8	0.7	1.4	3.3	6.1
2004.....	16.4	2.2	*	*	*	*	0.7	0.9	1.5	3.2	6.9
2003.....	16.7	*	*	*	*	*	*	0.9	1.5	2.5	6.8
2002.....	16.6	2.4	*	*	*	*	*	1.1	1.5	3.0	6.1
2001.....	17.4	*	*	*	*	*	0.8	1.0	2.0	3.4	7.0
2000.....	17.5	*	*	*	*	*	0.8	1.1	2.3	3.4	7.5
1999.....	17.4	*	*	*	*	*	0.7	1.4	1.8	3.3	6.9

Hispanic, male

2007.....	14.1	*	*	*	*	*	*	1.0	1.7	3.8	7.1
2006.....	14.3	*	*	*	*	*	*	1.2	1.6	3.6	7.6
2005.....	14.4	*	*	*	*	*	*	*	1.6	3.9	6.8
2004.....	15.0	*	*	*	*	*	*	1.2	1.7	4.3	7.7
2003.....	14.9	*	*	*	*	*	*	*	1.7	2.9	8.0
2002.....	15.0	*	*	*	*	*	*	1.4	1.7	3.4	7.1
2001.....	15.7	*	*	*	*	*	*	1.0	2.3	3.7	7.3
2000.....	15.8	*	*	*	*	*	*	1.2	2.6	3.6	7.8
1999.....	16.1	*	*	*	*	*	*	1.4	2.3	3.4	8.6

Hispanic, female

2007.....	17.1	*	*	*	*	*	*	*	1.4	2.6	4.2
2006.....	17.5	*	*	*	*	*	*	*	1.2	1.9	5.6
2005.....	17.7	*	*	*	*	*	*	*	1.3	2.6	5.4
2004.....	17.9	*	*	*	*	*	*	*	1.3	2.0	6.0
2003.....	18.6	*	*	*	*	*	*	*	1.3	2.1	5.5
2002.....	18.4	*	*	*	*	*	*	*	1.3	2.5	5.1
2001.....	19.1	*	*	*	*	*	*	*	1.6	3.1	6.6
2000.....	19.4	*	*	*	*	*	*	*	1.9	3.1	7.1
1999.....	18.8	*	*	*	*	*	*	1.4	*	3.1	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

Non-Hispanic 4/, both sexes

2007.....	50.3	3.2	0.3	0.2	0.2	0.3	0.6	0.8	1.7	3.5	6.5
2006.....	50.9	3.5	0.4	0.2	0.2	0.3	0.6	0.9	1.8	3.5	6.7
2005.....	53.8	3.2	0.4	0.3	0.2	0.3	0.5	1.0	1.8	3.3	6.9
2004.....	56.7	3.3	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4	7.1
2003.....	60.1	2.6	0.3	0.2	0.2	0.5	0.7	1.0	2.0	3.7	7.4
2002.....	62.4	3.0	0.3	0.2	0.2	0.3	0.6	0.9	1.9	3.7	7.2
2001.....	63.2	2.8	0.3	0.2	0.2	0.3	0.6	1.1	1.9	3.8	7.3
2000.....	65.4	3.6	0.3	0.1	0.3	0.4	0.7	1.1	1.8	3.9	7.6
1999.....	65.7	2.9	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.1	7.5

Non-Hispanic 4/, male

2007.....	40.5	3.8	*	*	*	0.4	0.6	0.8	1.6	3.8	6.7
2006.....	41.1	4.1	0.3	0.3	0.3	0.3	0.7	1.0	1.9	3.7	6.7
2005.....	43.0	3.6	0.5	0.3	0.3	0.3	0.5	1.1	2.0	3.2	7.0
2004.....	45.0	3.7	0.3	*	*	0.4	0.7	1.1	1.7	3.6	7.4
2003.....	47.6	2.8	*	*	*	0.5	0.7	1.1	2.1	4.0	7.6
2002.....	48.9	3.5	0.4	*	0.3	0.3	0.7	0.9	1.8	3.6	6.9
2001.....	49.7	3.3	0.3	*	*	0.3	0.6	1.1	1.9	3.9	7.3
2000.....	51.5	4.2	*	*	*	0.4	0.7	1.1	1.8	3.8	7.7
1999.....	51.5	3.7	0.3	*	*	0.3	0.6	0.9	1.8	4.0	7.7

Non-Hispanic 4/, female

2007.....	59.6	2.5	0.5	*	0.3	0.3	0.5	0.9	1.7	3.3	6.3
2006.....	60.3	2.8	0.4	*	*	0.3	0.5	0.8	1.7	3.3	6.6
2005.....	64.1	2.8	*	0.3	*	0.4	0.6	0.9	1.7	3.4	6.9
2004.....	67.8	3.0	*	*	*	0.3	0.7	1.1	2.0	3.3	6.9
2003.....	72.0	2.4	0.3	*	*	0.4	0.7	0.9	1.9	3.4	7.3
2002.....	75.3	2.5	*	*	*	*	0.5	0.9	2.0	3.9	7.5
2001.....	76.1	2.4	0.4	*	*	0.3	0.6	1.0	2.0	3.7	7.2
2000.....	78.7	2.8	0.4	*	0.3	0.3	0.7	1.0	1.9	4.1	7.6
1999.....	79.2	2.0	*	*	0.2	0.3	0.6	1.0	1.8	4.2	7.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

Non-Hispanic White, both sexes

2007 5/.....	53.5	2.8	0.3	0.2	0.2	0.3	0.5	0.7	1.2	2.7	4.8
2006 5/.....	54.2	2.4	0.4	0.2	0.2	0.3	0.5	0.7	1.4	2.5	4.9
2005 5/.....	57.4	2.4	0.3	0.2	0.2	0.3	0.4	0.8	1.5	2.6	5.1
2004 5/.....	60.6	2.6	0.3	0.2	0.2	0.3	0.6	0.9	1.4	2.8	5.4
2003 5/.....	64.2	2.0	0.2	*	0.2	0.4	0.5	0.8	1.6	2.6	5.3
2002.....	67.0	2.2	0.3	0.2	0.2	0.2	0.6	0.7	1.5	2.8	5.4
2001.....	67.6	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.6	3.0	5.4
2000.....	70.0	2.6	0.3	*	0.2	0.4	0.6	1.0	1.4	2.9	5.6
1999.....	70.4	2.4	0.3	*	0.2	0.3	0.6	0.8	1.5	3.2	5.5

Non-Hispanic White, male

2007 5/.....	41.9	3.1	*	*	*	0.4	0.5	0.7	1.1	2.7	5.2
2006 5/.....	42.6	2.6	*	*	*	*	0.6	0.7	1.5	2.5	5.1
2005 5/.....	44.8	2.8	0.4	*	*	*	0.4	0.8	1.6	2.5	5.2
2004 5/.....	47.0	2.9	*	*	*	0.3	0.6	1.0	1.2	2.7	5.8
2003 5/.....	49.7	1.9	*	*	*	0.4	0.7	0.8	1.7	2.7	5.5
2002.....	51.3	2.7	*	*	*	*	0.7	0.8	1.3	2.6	5.2
2001.....	52.0	2.2	*	*	*	0.3	0.6	0.8	1.7	2.9	5.6
2000.....	53.9	3.3	*	*	*	0.5	0.6	1.1	1.4	2.9	5.7
1999.....	53.9	3.1	*	*	*	0.3	0.5	0.8	1.5	3.1	5.7

Non-Hispanic White, female

2007 5/.....	64.7	2.4	*	*	*	*	0.5	0.7	1.4	2.6	4.5
2006 5/.....	65.5	2.2	0.4	*	*	*	0.4	0.7	1.3	2.4	4.8
2005 5/.....	69.6	2.0	*	*	*	0.3	0.5	0.7	1.4	2.6	5.0
2004 5/.....	73.7	2.2	*	*	*	*	0.6	0.8	1.6	2.9	5.0
2003 5/.....	78.2	2.1	*	*	*	0.4	0.4	0.7	1.5	2.5	5.2
2002.....	82.1	*	*	*	*	*	0.5	0.7	1.7	3.0	5.5
2001.....	82.6	*	*	*	*	0.3	0.4	0.9	1.6	3.0	5.1
2000.....	85.5	1.9	*	*	*	*	0.6	0.9	1.4	3.0	5.5
1999.....	86.1	*	*	*	*	*	0.7	0.8	1.4	3.2	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

Non-Hispanic Black, both sexes

2007 5/.....	44.4	5.2	*	*	*	*	0.9	1.5	4.2	8.1	15.8
2006 5/.....	44.7	7.7	*	*	*	*	1.0	1.8	3.9	8.8	16.4
2005 5/.....	46.5	6.2	*	*	*	*	1.1	2.4	3.8	7.1	17.6
2004 5/.....	48.7	6.2	*	*	*	*	1.0	2.3	4.3	7.3	17.8
2003 5/.....	51.1	4.9	*	*	*	0.7	1.3	2.1	4.3	9.1	19.2
2002.....	51.7	5.6	*	*	*	*	0.9	2.0	4.4	8.5	18.0
2001.....	52.9	6.9	*	*	*	*	1.4	2.3	3.6	8.3	18.9
2000.....	54.3	8.0	*	*	*	*	1.1	1.6	4.5	9.9	20.2
1999.....	53.9	4.7	*	*	*	*	0.8	1.9	3.8	9.4	20.3

Non-Hispanic Black, male

2007 5/.....	41.1	7.4	*	*	*	*	*	*	4.3	9.9	16.0
2006 5/.....	40.8	10.2	*	*	*	*	*	2.1	3.9	10.3	16.1
2005 5/.....	41.8	6.8	*	*	*	*	*	2.7	4.3	7.4	18.0
2004 5/.....	43.1	7.1	*	*	*	*	*	1.8	4.3	8.9	17.5
2003 5/.....	44.9	*	*	*	*	*	*	2.6	4.3	10.1	19.5
2002.....	45.1	*	*	*	*	*	*	1.7	4.1	7.8	17.4
2001.....	46.2	8.3	*	*	*	*	*	2.8	3.4	8.9	18.2
2000.....	47.7	9.2	*	*	*	*	*	*	4.2	9.5	21.0
1999.....	47.5	*	*	*	*	*	*	2.0	3.8	9.2	21.1

Non-Hispanic Black, female

2007 5/.....	47.4	*	*	*	*	*	*	1.6	4.0	6.4	15.6
2006 5/.....	48.3	*	*	*	*	*	*	1.6	3.8	7.4	16.7
2005 5/.....	50.8	*	*	*	*	*	*	2.0	3.4	6.8	17.3
2004 5/.....	53.8	*	*	*	*	*	*	2.7	4.3	5.9	18.0
2003 5/.....	56.7	*	*	*	*	*	1.7	1.6	4.3	8.3	18.9
2002.....	57.7	*	*	*	*	*	*	2.3	4.6	9.2	18.4
2001.....	59.0	*	*	*	*	*	1.8	1.8	3.9	7.7	19.6
2000.....	60.3	*	*	*	*	*	*	1.7	4.7	10.2	19.5
1999.....	59.8	*	*	*	*	*	*	1.8	3.7	9.6	19.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of circulatory system (I71-I78)

All origins 3/, both sexes

2007.....	7.3	*	*	*	*	0.1	0.2	0.2	0.5	0.9	1.4
2006.....	7.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.2
2005.....	8.0	*	*	*	*	*	0.2	0.3	0.5	0.8	1.3
2004.....	8.1	*	*	*	*	*	0.2	0.3	0.5	0.8	1.2
2003.....	8.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.2
2002.....	8.6	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.3
2001.....	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9	1.3
2000.....	9.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3
1999.....	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8	1.4

All origins 3/, male

2007.....	7.7	*	*	*	*	*	0.3	0.3	0.6	1.3	2.0
2006.....	8.0	*	*	*	*	*	0.3	0.4	0.6	1.1	1.8
2005.....	8.5	*	*	*	*	*	0.2	0.3	0.7	1.1	1.8
2004.....	8.5	*	*	*	*	*	0.3	0.5	0.6	1.1	1.7
2003.....	9.2	*	*	*	*	*	0.3	0.4	0.6	1.1	1.7
2002.....	9.4	*	*	*	*	*	0.2	0.3	0.7	0.7	1.8
2001.....	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2	1.8
2000.....	10.2	*	*	*	*	*	0.2	0.4	0.6	0.8	1.7
1999.....	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0	1.6

All origins 3/, female

2007.....	6.8	*	*	*	*	*	*	*	0.3	0.4	0.8
2006.....	7.0	*	*	*	*	*	*	*	0.2	0.5	0.7
2005.....	7.4	*	*	*	*	*	*	0.3	0.3	0.4	0.9
2004.....	7.6	*	*	*	*	*	*	*	0.4	0.6	0.8
2003.....	7.8	*	*	*	*	*	*	*	0.3	0.5	0.8
2002.....	7.9	*	*	*	*	*	*	0.2	0.4	0.7	0.7
2001.....	8.2	*	*	*	*	*	*	0.3	0.3	0.6	0.9
2000.....	8.3	*	*	*	*	*	*	*	0.4	0.6	0.9
1999.....	8.4	*	*	*	*	*	*	*	0.2	0.5	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of circulatory system (I71-I78)

Non-Hispanic 4/, both sexes

2007.....	8.2	*	*	*	*	*	0.2	0.2	0.5	1.0	1.4
2006.....	8.4	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.2
2005.....	8.9	*	*	*	*	*	0.2	0.3	0.5	0.8	1.4
2004.....	9.0	*	*	*	*	*	0.2	0.4	0.5	0.9	1.3
2003.....	9.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.3
2002.....	9.6	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.3
2001.....	9.9	*	*	*	*	*	0.2	0.4	0.5	0.9	1.4
2000.....	10.2	*	*	*	*	*	0.2	0.2	0.5	0.7	1.3
1999.....	10.2	*	*	*	*	*	0.2	0.3	0.5	0.8	1.4

Non-Hispanic 4/, male

2007.....	8.8	*	*	*	*	*	0.3	0.3	0.6	1.4	2.0
2006.....	9.1	*	*	*	*	*	0.3	0.4	0.6	1.2	1.9
2005.....	9.5	*	*	*	*	*	*	0.4	0.7	1.2	1.8
2004.....	9.6	*	*	*	*	*	0.3	0.5	0.7	1.1	1.8
2003.....	10.4	*	*	*	*	*	0.3	0.5	0.6	1.1	1.7
2002.....	10.5	*	*	*	*	*	0.2	0.3	0.7	0.8	1.9
2001.....	10.7	*	*	*	*	*	0.3	0.5	0.7	1.3	1.8
2000.....	11.3	*	*	*	*	*	0.3	0.4	0.7	0.9	1.7
1999.....	11.2	*	*	*	*	*	*	0.4	0.7	1.1	1.7

Non-Hispanic 4/, female

2007.....	7.7	*	*	*	*	*	*	*	0.3	0.5	0.9
2006.....	7.9	*	*	*	*	*	*	*	0.3	0.5	0.6
2005.....	8.3	*	*	*	*	*	*	*	0.3	0.4	0.9
2004.....	8.5	*	*	*	*	*	*	*	0.4	0.6	0.9
2003.....	8.6	*	*	*	*	*	*	*	0.3	0.5	0.8
2002.....	8.8	*	*	*	*	*	*	*	0.4	0.7	0.7
2001.....	9.1	*	*	*	*	*	*	0.3	0.3	0.6	0.9
2000.....	9.2	*	*	*	*	*	*	*	0.4	0.5	1.0
1999.....	9.3	*	*	*	*	*	*	*	0.2	0.6	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Aortic aneurysm and dissection (I71)

All origins 3/, both sexes

2007.....	4.3	*	*	*	*	*	0.1	0.1	0.4	0.7	1.0
2006.....	4.4	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0
2005.....	4.7	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2004.....	4.7	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
2003.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8
2002.....	5.1	*	*	*	*	0.1	*	0.2	0.4	0.5	0.9
2001.....	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9
2000.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9
1999.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9

All origins 3/, male

2007.....	5.1	*	*	*	*	*	0.2	0.2	0.5	1.0	1.5
2006.....	5.2	*	*	*	*	*	0.2	0.4	0.5	0.9	1.5
2005.....	5.6	*	*	*	*	*	*	0.3	0.5	0.9	1.5
2004.....	5.6	*	*	*	*	*	0.3	0.4	0.5	0.9	1.3
2003.....	6.3	*	*	*	*	*	0.2	0.3	0.5	0.9	1.3
2002.....	6.4	*	*	*	*	*	*	0.3	0.5	0.6	1.4
2001.....	6.5	*	*	*	*	*	*	0.4	0.5	0.9	1.4
2000.....	7.0	*	*	*	*	*	*	0.3	0.5	0.6	1.3
1999.....	7.0	*	*	*	*	*	*	0.3	0.5	0.9	1.2

All origins 3/, female

2007.....	3.6	*	*	*	*	*	*	*	*	0.3	0.5
2006.....	3.6	*	*	*	*	*	*	*	*	0.2	0.4
2005.....	3.8	*	*	*	*	*	*	*	*	0.2	0.4
2004.....	3.8	*	*	*	*	*	*	*	0.2	0.3	0.4
2003.....	3.9	*	*	*	*	*	*	*	*	0.3	0.4
2002.....	4.0	*	*	*	*	*	*	*	0.2	0.3	0.3
2001.....	4.2	*	*	*	*	*	*	*	*	0.3	0.3
2000.....	4.3	*	*	*	*	*	*	*	0.2	0.3	0.5
1999.....	4.4	*	*	*	*	*	*	*	*	0.3	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Aortic aneurysm and dissection (I71)

Non-Hispanic White, both sexes

2007 5/.....	5.4	*	*	*	*	*	*	*	0.3	0.6	0.8
2006 5/.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6	0.8
2005 5/.....	5.8	*	*	*	*	*	*	0.2	0.3	0.4	0.8
2004 5/.....	5.9	*	*	*	*	*	0.2	0.3	0.3	0.5	0.7
2003 5/.....	6.4	*	*	*	*	*	*	0.2	0.3	0.5	0.7
2002.....	6.5	*	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	6.7	*	*	*	*	*	0.2	0.3	0.3	0.6	0.7
2000.....	7.0	*	*	*	*	*	*	*	0.3	0.4	0.7
1999.....	7.1	*	*	*	*	*	*	0.2	0.3	0.6	0.8

Non-Hispanic White, male

2007 5/.....	6.4	*	*	*	*	*	*	*	0.4	1.0	1.2
2006 5/.....	6.6	*	*	*	*	*	*	0.3	0.4	0.9	1.3
2005 5/.....	7.0	*	*	*	*	*	*	*	0.5	0.7	1.2
2004 5/.....	7.0	*	*	*	*	*	*	0.4	0.4	0.8	1.1
2003 5/.....	7.9	*	*	*	*	*	*	*	0.4	0.8	1.2
2002.....	8.0	*	*	*	*	*	*	*	0.4	0.6	1.2
2001.....	8.2	*	*	*	*	*	*	0.4	0.5	0.9	1.1
2000.....	8.8	*	*	*	*	*	*	*	0.5	0.6	1.1
1999.....	8.9	*	*	*	*	*	*	*	0.5	0.9	1.0

Non-Hispanic White, female

2007 5/.....	4.5	*	*	*	*	*	*	*	*	*	0.5
2006 5/.....	4.5	*	*	*	*	*	*	*	*	0.3	0.3
2005 5/.....	4.7	*	*	*	*	*	*	*	*	*	0.4
2004 5/.....	4.7	*	*	*	*	*	*	*	*	*	0.3
2003 5/.....	4.9	*	*	*	*	*	*	*	*	*	0.3
2002.....	4.9	*	*	*	*	*	*	*	*	0.4	0.3
2001.....	5.2	*	*	*	*	*	*	*	*	*	0.3
2000.....	5.2	*	*	*	*	*	*	*	*	*	0.4
1999.....	5.3	*	*	*	*	*	*	*	*	0.3	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes

2007.....	3.0	*	*	*	*	*	*	*	0.1	0.2	0.4
2006.....	3.1	*	*	*	*	*	*	*	0.1	0.2	0.3
2005.....	3.3	*	*	*	*	*	*	*	0.1	0.2	0.4
2004.....	3.4	*	*	*	*	*	*	*	0.1	0.2	0.4
2003.....	3.4	*	*	*	*	*	*	*	0.1	0.2	0.4
2002.....	3.5	*	*	*	*	*	*	0.1	0.1	0.2	0.4
2001.....	3.5	*	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	3.6	*	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	3.6	*	*	*	*	*	*	*	0.1	0.2	0.5

All origins 3/, male

2007.....	2.6	*	*	*	*	*	*	*	*	0.2	0.4
2006.....	2.8	*	*	*	*	*	*	*	*	*	0.3
2005.....	2.9	*	*	*	*	*	*	*	*	0.2	0.3
2004.....	2.9	*	*	*	*	*	*	*	*	*	0.4
2003.....	3.0	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	3.0	*	*	*	*	*	*	*	*	*	0.4
2001.....	3.1	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	3.2	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	3.1	*	*	*	*	*	*	*	*	*	0.4

All origins 3/, female

2007.....	3.3	*	*	*	*	*	*	*	*	*	0.3
2006.....	3.4	*	*	*	*	*	*	*	*	0.2	0.2
2005.....	3.7	*	*	*	*	*	*	*	*	0.2	0.5
2004.....	3.9	*	*	*	*	*	*	*	*	0.3	0.5
2003.....	3.8	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	4.0	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	4.0	*	*	*	*	*	*	*	*	0.3	0.5
2000.....	4.0	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	4.1	*	*	*	*	*	*	*	*	0.2	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 4/, both sexes

2007.....	3.3	*	*	*	*	*	*	*	*	0.2	0.4
2006.....	3.5	*	*	*	*	*	*	*	*	0.2	0.2
2005.....	3.7	*	*	*	*	*	*	*	0.1	0.2	0.4
2004.....	3.8	*	*	*	*	*	*	*	0.1	0.2	0.5
2003.....	3.8	*	*	*	*	*	*	*	*	0.2	0.4
2002.....	3.9	*	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	3.9	*	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	4.0	*	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	3.9	*	*	*	*	*	*	*	0.2	0.2	0.5

Non-Hispanic 4/, male

2007.....	3.0	*	*	*	*	*	*	*	*	0.2	0.4
2006.....	3.1	*	*	*	*	*	*	*	*	*	0.3
2005.....	3.2	*	*	*	*	*	*	*	*	0.2	0.3
2004.....	3.2	*	*	*	*	*	*	*	*	*	0.4
2003.....	3.3	*	*	*	*	*	*	*	*	*	0.4
2002.....	3.4	*	*	*	*	*	*	*	*	*	0.4
2001.....	3.4	*	*	*	*	*	*	*	*	0.3	0.4
2000.....	3.5	*	*	*	*	*	*	*	*	*	0.3
1999.....	3.4	*	*	*	*	*	*	*	*	*	0.4

Non-Hispanic 4/, female

2007.....	3.7	*	*	*	*	*	*	*	*	*	0.3
2006.....	3.8	*	*	*	*	*	*	*	*	0.2	*
2005.....	4.1	*	*	*	*	*	*	*	*	0.2	0.5
2004.....	4.3	*	*	*	*	*	*	*	*	0.3	0.5
2003.....	4.2	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	4.3	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	4.4	*	*	*	*	*	*	*	*	0.3	0.6
2000.....	4.4	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	4.5	*	*	*	*	*	*	*	*	0.2	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other disorders of circulatory system (I80-I99)

All origins 3/, both sexes

2007.....	1.4	1.2	*	*	*	*	0.1	0.2	0.4	0.6	0.9
2006.....	1.3	0.8	*	*	*	*	0.2	0.2	0.4	0.5	0.7
2005.....	1.6	0.6	*	*	*	*	0.1	0.2	0.4	0.6	0.9
2004.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.5	0.9
2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6	0.9
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6	0.8
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4	0.8
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6	0.9
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4	0.8

All origins 3/, male

2007.....	1.3	1.4	*	*	*	*	*	0.2	0.5	0.7	1.0
2006.....	1.2	*	*	*	*	*	*	*	0.4	0.6	0.8
2005.....	1.5	*	*	*	*	*	*	0.2	0.4	0.5	1.0
2004.....	1.4	*	*	*	*	*	*	0.3	0.4	0.5	1.0
2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5	1.0
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6	0.8
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6	1.1
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5	0.9

All origins 3/, female

2007.....	1.4	1.1	*	*	*	*	*	0.3	0.2	0.6	0.7
2006.....	1.5	*	*	*	*	*	0.2	*	0.4	0.5	0.7
2005.....	1.8	*	*	*	*	*	*	0.3	0.4	0.7	0.9
2004.....	1.7	*	*	*	*	*	*	0.3	0.4	0.5	0.8
2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6	0.9
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4	0.8
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5	0.7
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other disorders of circulatory system (I80-I99)

Non-Hispanic 4/, both sexes

2007.....	1.5	1.1	*	*	*	*	*	0.2	0.4	0.7	1.0
2006.....	1.5	0.8	*	*	*	*	0.2	0.2	0.4	0.6	0.8
2005.....	1.8	0.8	*	*	*	*	0.1	0.3	0.4	0.6	1.0
2004.....	1.8	0.8	*	*	*	*	0.1	0.3	0.5	0.6	1.0
2003.....	1.8	0.8	*	*	*	*	0.2	0.3	0.4	0.6	1.0
2002.....	1.8	0.6	*	*	*	*	0.1	0.2	0.5	0.6	0.8
2001.....	1.8	*	*	*	*	*	0.2	0.3	0.4	0.5	0.9
2000.....	1.8	0.7	*	*	*	*	0.2	0.4	0.4	0.6	0.9
1999.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5	0.8

Non-Hispanic 4/, male

2007.....	1.4	1.3	*	*	*	*	*	*	0.5	0.8	1.2
2006.....	1.3	*	*	*	*	*	*	*	0.4	0.7	0.8
2005.....	1.6	*	*	*	*	*	*	0.2	0.4	0.5	1.0
2004.....	1.6	*	*	*	*	*	*	0.3	0.5	0.6	1.0
2003.....	1.5	*	*	*	*	*	*	0.3	0.4	0.6	1.0
2002.....	1.5	*	*	*	*	*	*	*	0.6	0.7	0.8
2001.....	1.5	*	*	*	*	*	*	*	0.4	0.5	0.9
2000.....	1.5	*	*	*	*	*	*	*	0.4	0.6	1.1
1999.....	1.4	*	*	*	*	*	*	*	0.3	0.5	1.0

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	*	*	0.3	0.3	0.6	0.8
2006.....	1.6	*	*	*	*	*	*	*	0.5	0.5	0.7
2005.....	1.9	*	*	*	*	*	*	0.3	0.5	0.6	0.9
2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.6	0.9
2003.....	2.1	*	*	*	*	*	*	0.3	0.5	0.7	1.0
2002.....	2.1	*	*	*	*	*	*	0.3	0.5	0.6	0.8
2001.....	2.0	*	*	*	*	*	*	0.3	0.5	0.5	0.8
2000.....	2.0	*	*	*	*	*	*	0.5	0.4	0.5	0.6
1999.....	1.8	*	*	*	*	*	*	*	0.3	0.4	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other disorders of circulatory system (I80-I99)

Non-Hispanic White, both sexes

2007 5/.....	1.5	1.0	*	*	*	*	*	0.2	0.4	0.5	0.8
2006 5/.....	1.5	0.9	*	*	*	*	*	*	0.4	0.5	0.6
2005 5/.....	1.8	*	*	*	*	*	*	0.2	0.4	0.5	0.7
2004 5/.....	1.8	*	*	*	*	*	*	0.2	0.4	0.4	0.7
2003 5/.....	1.8	*	*	*	*	*	*	0.3	0.3	0.5	0.8
2002.....	1.8	*	*	*	*	*	*	*	0.4	0.5	0.7
2001.....	1.8	*	*	*	*	*	*	0.2	0.3	0.4	0.7
2000.....	1.8	*	*	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	1.6	*	*	*	*	*	*	0.2	0.3	0.4	0.7

Non-Hispanic White, male

2007 5/.....	1.4	*	*	*	*	*	*	*	0.4	0.6	1.1
2006 5/.....	1.3	*	*	*	*	*	*	*	0.4	0.6	0.7
2005 5/.....	1.6	*	*	*	*	*	*	*	0.4	0.5	0.8
2004 5/.....	1.6	*	*	*	*	*	*	*	0.4	0.4	0.8
2003 5/.....	1.5	*	*	*	*	*	*	*	0.3	0.5	0.9
2002.....	1.5	*	*	*	*	*	*	*	0.4	0.5	0.6
2001.....	1.5	*	*	*	*	*	*	*	0.3	0.4	0.8
2000.....	1.5	*	*	*	*	*	*	*	*	0.5	1.0
1999.....	1.4	*	*	*	*	*	*	*	*	0.4	0.8

Non-Hispanic White, female

2007 5/.....	1.7	*	*	*	*	*	*	*	*	0.5	0.6
2006 5/.....	1.7	*	*	*	*	*	*	*	0.4	0.4	0.5
2005 5/.....	2.0	*	*	*	*	*	*	*	0.3	0.5	0.7
2004 5/.....	2.0	*	*	*	*	*	*	*	0.4	0.5	0.6
2003 5/.....	2.1	*	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	2.2	*	*	*	*	*	*	*	0.3	0.4	0.7
2001.....	2.1	*	*	*	*	*	*	*	0.4	0.4	0.7
2000.....	2.0	*	*	*	*	*	*	0.4	0.3	0.4	0.5
1999.....	1.8	*	*	*	*	*	*	*	0.3	0.3	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Influenza and pneumonia (J09-J18)

All origins 3/, both sexes

2007.....	17.5	5.2	0.7	0.2	0.3	0.2	0.5	0.7	0.9	1.4	2.3
2006.....	18.8	6.4	0.8	0.2	0.1	0.3	0.5	0.7	1.0	1.4	2.4
2005.....	21.3	6.5	0.7	0.3	0.3	0.3	0.5	0.7	1.0	1.5	2.7
2004.....	20.3	6.7	0.7	0.2	0.2	0.3	0.6	0.5	1.0	1.5	2.5
2003.....	22.4	8.0	1.0	0.4	0.3	0.4	0.7	0.8	1.1	1.8	2.7
2002.....	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7	2.6
2001.....	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8	2.6
2000.....	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8	3.0
1999.....	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0	2.7

All origins 3/, male

2007.....	16.2	6.2	0.7	0.2	0.3	0.2	0.6	0.8	1.0	1.4	2.6
2006.....	17.4	7.7	0.8	*	*	0.3	0.6	0.8	0.9	1.5	2.7
2005.....	19.2	7.0	0.7	0.3	0.3	0.5	0.6	0.8	1.2	1.8	3.1
2004.....	18.6	8.0	0.6	*	0.3	0.4	0.7	0.7	1.2	1.7	2.7
2003.....	20.1	9.4	1.1	0.4	0.3	0.5	0.7	0.9	1.3	1.9	3.1
2002.....	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8	3.1
2001.....	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0	3.0
2000.....	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9	3.8
1999.....	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2	3.1

All origins 3/, female

2007.....	18.7	4.2	0.7	0.2	0.3	0.3	0.4	0.6	0.8	1.3	1.9
2006.....	20.2	5.0	0.8	0.2	*	0.4	0.5	0.6	1.0	1.3	2.2
2005.....	23.2	5.8	0.7	0.3	0.3	*	0.4	0.7	0.9	1.2	2.3
2004.....	22.0	5.4	0.9	*	0.2	0.2	0.4	0.4	0.7	1.2	2.3
2003.....	24.6	6.6	0.9	0.4	0.4	0.3	0.7	0.6	0.8	1.6	2.3
2002.....	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6	2.2
2001.....	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6	2.2
2000.....	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7	2.2
1999.....	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Influenza and pneumonia (J09-J18)

Non-Hispanic 4/, both sexes

2007.....	19.5	5.4	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.4	2.4
2006.....	20.9	6.3	0.8	0.2	0.1	0.4	0.5	0.7	1.1	1.5	2.6
2005.....	23.6	6.2	0.6	0.3	0.3	0.4	0.6	0.8	1.1	1.6	2.8
2004.....	22.4	6.9	0.8	0.2	0.2	0.3	0.6	0.6	1.0	1.5	2.6
2003.....	24.7	8.1	1.0	0.4	0.4	0.4	0.7	0.8	1.1	1.9	2.8
2002.....	25.1	6.7	0.7	0.2	0.2	0.4	0.5	0.8	1.1	1.7	2.8
2001.....	23.8	7.5	0.6	0.2	0.2	0.3	0.6	0.7	1.2	1.9	2.7
2000.....	25.4	7.5	0.7	0.2	0.2	0.4	0.7	0.8	1.1	1.8	3.0
1999.....	25.0	8.4	0.9	0.2	0.2	0.4	0.6	0.8	1.1	2.0	2.8

Non-Hispanic 4/, male

2007.....	18.1	6.5	0.7	0.3	0.2	0.2	0.7	1.0	1.1	1.5	2.7
2006.....	19.4	7.8	0.8	*	*	0.3	0.6	0.8	1.0	1.6	2.8
2005.....	21.4	6.7	0.6	0.3	0.3	0.5	0.7	0.8	1.3	1.9	3.2
2004.....	20.6	8.0	0.6	*	0.3	0.4	0.7	0.8	1.2	1.8	2.8
2003.....	22.3	9.4	1.1	0.4	0.3	0.5	0.7	1.0	1.4	2.2	3.2
2002.....	22.5	7.8	0.8	*	*	0.4	0.6	0.7	1.2	1.9	3.2
2001.....	21.5	8.2	0.6	0.2	0.2	0.3	0.8	0.7	1.4	2.1	3.1
2000.....	22.7	8.5	0.7	0.2	*	0.4	0.7	0.9	1.2	1.9	3.8
1999.....	22.2	8.4	0.9	0.3	0.3	0.3	0.6	0.9	1.3	2.1	3.2

Non-Hispanic 4/, female

2007.....	20.8	4.3	0.7	*	0.3	0.3	0.4	0.7	0.9	1.4	2.0
2006.....	22.3	4.8	0.8	*	*	0.4	0.4	0.6	1.2	1.4	2.3
2005.....	25.7	5.7	0.6	0.3	0.3	*	0.4	0.7	1.0	1.3	2.4
2004.....	24.2	5.9	0.9	*	*	0.2	0.4	0.3	0.8	1.3	2.4
2003.....	27.1	6.8	1.0	0.4	0.4	0.3	0.7	0.7	0.9	1.7	2.4
2002.....	27.5	5.4	0.7	*	0.3	0.4	0.4	0.8	0.9	1.6	2.4
2001.....	26.1	6.7	0.6	0.3	0.2	0.3	0.5	0.7	0.9	1.7	2.3
2000.....	27.9	6.5	0.6	*	*	0.3	0.6	0.6	1.0	1.8	2.2
1999.....	27.6	8.4	0.9	*	*	0.4	0.5	0.7	0.9	2.0	2.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Influenza and pneumonia (J09-J18)

Non-Hispanic White, both sexes

2007 5/.....	21.5	3.8	0.5	0.2	0.3	0.3	0.6	0.7	0.8	1.3	2.2
2006 5/.....	23.1	4.5	0.6	0.2	*	0.4	0.5	0.7	0.9	1.4	2.4
2005 5/.....	26.2	4.1	0.5	0.3	0.3	0.4	0.6	0.8	1.1	1.4	2.5
2004 5/.....	24.8	4.8	0.6	*	0.2	0.3	0.5	0.5	0.9	1.3	2.3
2003 5/.....	27.4	6.2	0.9	0.3	0.4	0.4	0.7	0.7	1.0	1.7	2.4
2002.....	27.9	4.9	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.5	2.4
2001.....	26.2	4.9	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6	2.2
2000.....	27.9	5.0	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.5
1999.....	27.5	5.3	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.8	2.2

Non-Hispanic White, male

2007 5/.....	19.6	4.8	0.4	*	*	*	0.7	0.8	0.9	1.3	2.4
2006 5/.....	21.2	5.8	0.6	*	*	0.3	0.6	0.7	0.9	1.4	2.7
2005 5/.....	23.3	5.1	0.5	*	*	0.5	0.7	0.8	1.1	1.7	2.8
2004 5/.....	22.4	5.5	0.5	*	*	0.4	0.7	0.7	1.1	1.5	2.5
2003 5/.....	24.1	7.4	0.9	*	0.3	0.4	0.7	0.9	1.2	2.0	2.7
2002.....	24.6	6.1	0.5	*	*	0.4	0.6	0.7	1.1	1.5	2.8
2001.....	23.1	5.8	0.6	*	*	*	0.7	0.6	1.2	1.7	2.5
2000.....	24.5	5.8	0.5	*	*	0.4	0.7	0.7	1.0	1.7	3.2
1999.....	23.9	5.5	0.5	*	*	0.3	0.5	0.7	1.1	1.8	2.3

Non-Hispanic White, female

2007 5/.....	23.3	2.7	0.7	*	0.3	*	0.5	0.6	0.8	1.2	2.0
2006 5/.....	25.0	3.0	0.6	*	*	0.4	0.4	0.6	1.0	1.4	2.1
2005 5/.....	28.9	3.0	0.5	*	0.3	*	0.4	0.8	1.0	1.2	2.2
2004 5/.....	27.1	4.2	0.7	*	*	*	0.3	0.3	0.8	1.1	2.2
2003 5/.....	30.6	5.0	0.8	*	0.4	*	0.7	0.5	0.8	1.5	2.2
2002.....	31.1	3.6	0.5	*	0.3	*	0.4	0.8	0.8	1.4	2.1
2001.....	29.2	3.9	0.5	*	*	0.3	0.4	0.7	0.8	1.5	2.0
2000.....	31.3	4.2	0.6	*	*	*	0.5	0.6	1.0	1.7	1.9
1999.....	31.0	5.1	0.8	*	*	0.4	0.4	0.5	0.7	1.7	2.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J09-J18)

Non-Hispanic Black, both sexes

2007 5/.....	13.3	11.2	1.1	*	*	*	*	1.6	1.8	2.6	3.8
2006 5/.....	13.9	13.7	1.4	*	*	*	1.0	1.1	1.8	2.5	4.2
2005 5/.....	15.3	15.1	1.1	*	*	*	*	1.0	1.9	2.7	5.1
2004 5/.....	15.2	14.7	1.4	*	*	*	0.9	0.8	1.6	3.0	5.0
2003 5/.....	15.9	14.4	1.1	*	*	*	0.8	1.7	2.2	3.4	5.6
2002.....	16.1	13.3	1.7	*	*	*	0.7	1.1	2.0	3.6	5.3
2001.....	16.0	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	6.0
2000.....	16.9	17.5	1.2	*	*	*	1.1	1.4	2.1	2.9	6.3
1999.....	16.7	20.0	1.7	*	*	*	1.3	1.8	2.3	4.1	7.1

Non-Hispanic Black, male

2007 5/.....	13.5	12.5	*	*	*	*	*	1.9	2.0	2.8	5.1
2006 5/.....	13.6	16.3	*	*	*	*	*	1.5	*	3.0	4.4
2005 5/.....	15.1	14.0	*	*	*	*	*	*	2.4	3.5	6.4
2004 5/.....	15.5	17.0	*	*	*	*	*	*	2.0	3.8	5.9
2003 5/.....	16.5	16.1	*	*	*	*	*	1.7	2.9	3.7	7.2
2002.....	15.9	14.2	1.8	*	*	*	*	*	2.4	4.1	6.2
2001.....	16.3	16.6	*	*	*	*	*	*	2.8	4.7	7.1
2000.....	17.3	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.6
1999.....	16.9	20.1	2.1	*	*	*	*	2.0	2.6	4.3	9.2

Non-Hispanic Black, female

2007 5/.....	13.2	9.9	*	*	*	*	*	*	1.7	2.5	2.7
2006 5/.....	14.1	10.9	*	*	*	*	*	*	2.3	2.1	4.0
2005 5/.....	15.5	16.2	*	*	*	*	*	*	1.5	1.9	4.0
2004 5/.....	15.0	12.2	1.8	*	*	*	*	*	*	2.3	4.3
2003 5/.....	15.4	12.6	*	*	*	*	*	1.7	1.5	3.2	4.2
2002.....	16.2	12.3	*	*	*	*	*	*	1.6	3.2	4.5
2001.....	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.9
2000.....	16.5	16.2	*	*	*	*	*	*	1.7	2.7	4.3
1999.....	16.5	20.0	*	*	*	*	*	1.6	1.9	3.8	5.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pneumonia (J12-J18)

All origins 3/, both sexes

2007.....	17.3	4.9	0.5	0.1	0.2	0.2	0.5	0.7	0.9	1.3	2.2
2006.....	18.5	5.9	0.7	0.1	*	0.3	0.5	0.7	1.0	1.4	2.4
2005.....	20.6	6.0	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.5	2.6
2004.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4	2.5
2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7	2.6
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7	2.6
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8	2.6
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7	2.9
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0	2.7

All origins 3/, male

2007.....	16.1	5.8	0.5	*	0.2	0.2	0.6	0.8	1.0	1.4	2.6
2006.....	17.1	7.2	0.6	*	*	0.3	0.6	0.8	0.9	1.5	2.7
2005.....	18.8	6.6	0.6	0.2	0.2	0.4	0.6	0.8	1.1	1.8	3.0
2004.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7	2.7
2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9	3.0
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8	3.0
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0	3.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8	3.7
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2	3.1

All origins 3/, female

2007.....	18.6	4.0	0.6	*	*	0.2	0.4	0.6	0.8	1.2	1.8
2006.....	19.9	4.6	0.7	*	*	0.3	0.4	0.6	1.0	1.3	2.1
2005.....	22.5	5.3	0.6	0.2	0.2	*	0.3	0.7	0.9	1.2	2.3
2004.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2	2.3
2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5	2.2
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5	2.2
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6	2.2
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7	2.1
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pneumonia (J12-J18)

Non-Hispanic 4/, both sexes

2007.....	19.3	5.2	0.5	0.1	0.2	0.3	0.5	0.8	1.0	1.4	2.3
2006.....	20.6	5.9	0.7	*	*	0.3	0.5	0.7	1.0	1.5	2.5
2005.....	22.9	5.9	0.5	0.2	0.2	0.3	0.5	0.8	1.1	1.5	2.8
2004.....	22.0	6.6	0.6	0.2	0.2	0.3	0.5	0.6	1.0	1.5	2.6
2003.....	24.0	7.4	0.6	0.2	0.3	0.3	0.6	0.8	1.1	1.9	2.8
2002.....	24.8	6.5	0.7	0.2	0.2	0.3	0.5	0.7	1.1	1.7	2.7
2001.....	23.8	7.2	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.9	2.7
2000.....	24.7	7.3	0.6	0.2	0.1	0.3	0.7	0.7	1.1	1.8	2.9
1999.....	24.3	8.1	0.8	0.2	0.2	0.3	0.6	0.8	1.0	2.0	2.8

Non-Hispanic 4/, male

2007.....	18.0	6.2	0.5	*	*	0.2	0.7	0.9	1.1	1.5	2.7
2006.....	19.1	7.4	0.7	*	*	0.3	0.6	0.8	0.9	1.5	2.8
2005.....	20.9	6.5	0.5	*	*	0.5	0.7	0.8	1.2	1.8	3.1
2004.....	20.3	7.5	0.5	*	*	0.4	0.7	0.8	1.2	1.8	2.8
2003.....	21.7	8.7	0.7	0.3	0.2	0.4	0.7	0.9	1.4	2.1	3.2
2002.....	22.3	7.6	0.8	*	*	0.4	0.6	0.7	1.2	1.9	3.1
2001.....	21.4	7.8	0.6	*	0.2	0.3	0.7	0.7	1.3	2.1	3.0
2000.....	22.2	8.2	0.7	*	*	0.4	0.7	0.8	1.2	1.9	3.7
1999.....	21.7	8.0	0.8	*	0.3	0.3	0.6	0.9	1.2	2.1	3.2

Non-Hispanic 4/, female

2007.....	20.6	4.1	0.6	*	*	0.3	0.4	0.6	0.9	1.3	2.0
2006.....	21.9	4.4	0.7	*	*	0.3	0.4	0.6	1.2	1.4	2.3
2005.....	24.8	5.2	0.5	*	0.3	*	0.4	0.7	1.0	1.3	2.4
2004.....	23.7	5.6	0.8	*	*	*	0.4	0.3	0.7	1.2	2.4
2003.....	26.3	6.1	0.6	*	0.3	*	0.6	0.7	0.8	1.6	2.3
2002.....	27.2	5.3	0.6	*	*	0.3	0.4	0.8	0.9	1.6	2.3
2001.....	26.0	6.6	0.6	*	*	0.3	0.5	0.7	0.9	1.7	2.3
2000.....	27.1	6.4	0.5	*	*	0.3	0.6	0.6	1.0	1.8	2.2
1999.....	26.8	8.2	0.9	*	*	0.3	0.5	0.6	0.9	1.9	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pneumonia (J12-J18)

Non-Hispanic White, both sexes

2007 5/.....	21.3	3.6	0.4	*	0.2	0.2	0.6	0.7	0.8	1.3	2.1
2006 5/.....	22.7	4.1	0.6	*	*	0.4	0.4	0.6	0.9	1.3	2.4
2005 5/.....	25.3	3.8	0.4	0.2	0.2	0.3	0.5	0.8	1.0	1.4	2.5
2004 5/.....	24.3	4.6	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2	2.3
2003 5/.....	26.6	5.7	0.6	0.2	0.3	0.3	0.6	0.7	0.9	1.7	2.4
2002.....	27.6	4.7	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.5	2.4
2001.....	26.1	4.6	0.5	0.2	0.2	0.3	0.5	0.6	1.0	1.6	2.2
2000.....	27.1	4.8	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.4
1999.....	26.8	5.2	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.7	2.1

Non-Hispanic White, male

2007 5/.....	19.4	4.6	*	*	*	*	0.7	0.8	0.9	1.3	2.3
2006 5/.....	20.9	5.4	0.6	*	*	0.3	0.6	0.7	0.9	1.4	2.6
2005 5/.....	22.7	4.7	*	*	*	0.5	0.7	0.8	1.1	1.6	2.7
2004 5/.....	22.0	5.3	*	*	*	0.4	0.7	0.7	1.1	1.5	2.4
2003 5/.....	23.5	6.8	0.6	*	*	0.4	0.7	0.8	1.1	1.9	2.6
2002.....	24.3	6.0	0.5	*	*	0.4	0.6	0.6	1.1	1.5	2.8
2001.....	23.0	5.3	0.6	*	*	*	0.7	0.5	1.1	1.7	2.5
2000.....	23.8	5.4	0.4	*	*	0.3	0.7	0.6	1.0	1.7	3.1
1999.....	23.4	5.4	*	*	*	0.3	0.5	0.7	1.0	1.8	2.3

Non-Hispanic White, female

2007 5/.....	23.1	2.6	0.6	*	*	*	0.4	0.6	0.7	1.2	2.0
2006 5/.....	24.6	2.8	0.6	*	*	0.4	0.3	0.6	1.0	1.3	2.1
2005 5/.....	27.9	2.8	0.4	*	*	*	0.4	0.7	0.9	1.2	2.2
2004 5/.....	26.5	3.8	0.6	*	*	*	*	0.3	0.7	1.0	2.2
2003 5/.....	29.6	4.5	0.5	*	0.3	*	0.6	0.5	0.7	1.4	2.1
2002.....	30.7	3.4	0.5	*	*	*	0.4	0.8	0.8	1.4	2.1
2001.....	29.1	3.8	0.4	*	*	0.3	0.4	0.7	0.8	1.5	1.9
2000.....	30.3	4.2	0.5	*	*	*	0.5	0.6	0.9	1.6	1.8
1999.....	30.0	4.9	0.7	*	*	*	*	0.5	0.7	1.7	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pneumonia (J12-J18)

Non-Hispanic Black, both sexes

2007 5/.....	13.3	10.6	0.9	*	*	*	*	1.6	1.8	2.6	3.8
2006 5/.....	13.8	12.9	1.2	*	*	*	0.9	1.1	1.7	2.5	4.2
2005 5/.....	15.1	14.3	1.0	*	*	*	*	1.0	1.9	2.5	5.0
2004 5/.....	15.1	14.0	1.2	*	*	*	0.9	0.8	1.5	3.0	5.0
2003 5/.....	15.6	13.3	0.9	*	*	*	0.7	1.6	2.2	3.4	5.6
2002.....	16.0	13.1	1.5	*	*	*	0.7	1.1	2.0	3.6	5.3
2001.....	15.9	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	5.9
2000.....	16.6	17.2	1.2	*	*	*	1.1	1.4	2.1	2.9	6.2
1999.....	16.5	19.3	1.7	*	*	*	1.3	1.8	2.3	4.0	7.0

Non-Hispanic Black, male

2007 5/.....	13.5	11.6	*	*	*	*	*	1.9	2.0	2.7	5.1
2006 5/.....	13.5	15.7	*	*	*	*	*	1.5	*	3.0	4.4
2005 5/.....	15.0	14.0	*	*	*	*	*	*	2.4	3.2	6.4
2004 5/.....	15.4	15.8	*	*	*	*	*	*	1.9	3.8	5.9
2003 5/.....	16.2	15.2	*	*	*	*	*	1.6	2.9	3.7	7.2
2002.....	15.8	13.9	1.6	*	*	*	*	*	2.4	4.1	6.2
2001.....	16.3	16.6	*	*	*	*	*	*	2.7	4.7	7.1
2000.....	17.1	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.3
1999.....	16.8	19.1	2.0	*	*	*	*	2.0	2.6	4.3	9.1

Non-Hispanic Black, female

2007 5/.....	13.1	9.6	*	*	*	*	*	*	1.6	2.4	2.7
2006 5/.....	14.0	10.0	*	*	*	*	*	*	2.2	2.1	3.9
2005 5/.....	15.3	14.6	*	*	*	*	*	*	1.5	1.9	3.9
2004 5/.....	14.8	12.2	1.7	*	*	*	*	*	*	2.3	4.2
2003 5/.....	15.0	11.3	*	*	*	*	*	1.6	1.4	3.2	4.2
2002.....	16.1	12.3	*	*	*	*	*	*	1.6	3.2	4.5
2001.....	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.8
2000.....	16.3	15.5	*	*	*	*	*	*	1.7	2.7	4.3
1999.....	16.3	19.6	*	*	*	*	*	1.5	1.9	3.7	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes

2007.....	42.4	1.0	0.3	0.3	0.3	0.3	0.4	0.6	0.7	1.1	2.6
2006.....	41.6	0.7	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.0	2.8
2005.....	44.2	0.8	0.3	0.3	0.3	0.3	0.4	0.5	0.8	1.2	2.8
2004.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3	2.6
2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4	2.8
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6	2.9
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6	2.7
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4	2.8
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3	2.6

All origins 3/, male

2007.....	41.2	1.2	0.4	0.4	0.4	0.4	0.5	0.7	0.7	1.0	2.6
2006.....	40.2	0.9	0.3	0.3	0.4	0.4	0.5	0.6	0.7	1.0	2.6
2005.....	42.8	*	0.4	0.3	0.3	0.4	0.5	0.6	0.7	1.2	2.7
2004.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2	2.6
2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3	2.5
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5	3.0
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5	2.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3	2.6
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2	2.4

All origins 3/, female

2007.....	43.6	*	0.3	*	0.2	0.3	0.2	0.5	0.6	1.1	2.6
2006.....	43.0	*	*	*	*	0.2	0.4	0.5	0.7	1.1	2.9
2005.....	45.5	*	0.3	0.2	0.2	*	0.4	0.4	0.9	1.3	2.8
2004.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4	2.7
2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5	3.1
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7	2.8
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7	2.9
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4	2.9
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4	2.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 4/, both sexes

2007.....	48.5	1.0	0.4	0.3	0.4	0.3	0.4	0.6	0.8	1.2	2.9
2006.....	47.5	0.8	0.3	0.3	0.3	0.3	0.5	0.6	0.8	1.1	3.0
2005.....	50.2	0.8	0.4	0.3	0.3	0.3	0.5	0.6	0.9	1.4	3.0
2004.....	47.0	0.8	0.3	0.3	0.4	0.5	0.5	0.6	0.8	1.5	2.8
2003.....	49.0	0.8	0.3	0.2	0.4	0.4	0.6	0.7	0.9	1.5	3.0
2002.....	48.6	1.1	0.4	0.2	0.5	0.5	0.5	0.7	0.9	1.7	3.0
2001.....	48.4	1.1	0.3	0.2	0.3	0.4	0.6	0.6	1.0	1.7	2.9
2000.....	48.3	1.0	0.3	0.2	0.5	0.5	0.6	0.7	0.9	1.4	2.9
1999.....	49.4	1.0	0.4	0.2	0.5	0.6	0.6	0.8	1.0	1.3	2.7

Non-Hispanic 4/, male

2007.....	47.3	*	0.4	0.4	0.5	0.4	0.5	0.7	0.8	1.2	2.9
2006.....	46.1	*	0.3	0.3	0.5	0.4	0.6	0.7	0.9	1.0	2.9
2005.....	48.8	*	0.5	0.3	0.3	0.4	0.6	0.6	0.8	1.3	2.8
2004.....	46.2	*	0.4	0.3	0.5	0.6	0.5	0.7	0.9	1.4	2.6
2003.....	48.1	*	0.5	0.3	0.5	0.5	0.7	0.8	1.0	1.4	2.7
2002.....	48.4	1.2	0.6	0.2	0.5	0.7	0.5	0.8	0.9	1.6	3.1
2001.....	48.0	*	0.4	0.3	0.4	0.4	0.7	0.6	1.0	1.7	2.7
2000.....	48.7	*	0.4	0.3	0.5	0.6	0.7	0.8	1.0	1.4	2.8
1999.....	50.8	*	0.4	0.3	0.6	0.6	0.8	0.8	0.9	1.3	2.5

Non-Hispanic 4/, female

2007.....	49.6	*	0.4	*	*	0.3	0.3	0.5	0.7	1.3	2.9
2006.....	48.8	*	*	*	*	0.2	0.4	0.5	0.8	1.2	3.2
2005.....	51.5	*	0.3	*	*	*	0.4	0.5	0.9	1.5	3.1
2004.....	47.8	*	*	*	0.3	0.3	0.4	0.6	0.6	1.6	2.9
2003.....	49.9	*	*	*	0.3	0.3	0.5	0.6	0.8	1.6	3.4
2002.....	48.9	*	*	*	0.5	0.3	0.5	0.6	0.9	1.8	2.9
2001.....	48.7	*	*	*	0.3	0.3	0.5	0.6	1.0	1.7	3.0
2000.....	48.0	*	*	*	0.4	0.3	0.5	0.6	0.8	1.5	3.0
1999.....	48.0	*	*	*	0.3	0.5	0.5	0.8	1.1	1.4	2.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic White, both sexes

2007 5/.....	56.9	*	*	*	0.2	0.2	0.2	0.5	0.6	1.1	2.7
2006 5/.....	55.6	*	*	*	*	*	0.3	0.4	0.6	0.9	2.7
2005 5/.....	58.6	*	0.2	0.2	*	0.2	0.3	0.4	0.6	1.2	2.7
2004 5/.....	54.9	*	*	*	0.2	0.3	0.3	0.4	0.6	1.3	2.5
2003 5/.....	57.0	*	*	*	0.2	0.2	0.4	0.5	0.6	1.3	2.7
2002.....	56.4	0.9	0.2	*	0.2	0.3	0.4	0.5	0.7	1.4	2.6
2001.....	55.9	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	2.6
2000.....	55.7	*	*	*	0.2	0.3	0.4	0.5	0.6	1.1	2.4
1999.....	56.6	*	0.2	0.2	0.2	0.5	0.4	0.5	0.6	1.0	2.3

Non-Hispanic White, male

2007 5/.....	54.7	*	*	*	*	*	0.3	0.5	0.6	1.0	2.6
2006 5/.....	53.0	*	*	*	*	*	0.4	0.4	0.5	0.8	2.6
2005 5/.....	56.0	*	*	*	*	*	0.3	0.4	0.5	1.1	2.5
2004 5/.....	53.0	*	*	*	*	0.3	0.4	0.3	0.7	1.1	2.4
2003 5/.....	54.9	*	*	*	*	*	0.5	0.4	0.6	1.2	2.3
2002.....	55.1	*	*	*	*	0.4	0.4	0.5	0.8	1.1	2.7
2001.....	54.6	*	*	*	*	*	0.4	0.4	0.6	1.3	2.4
2000.....	55.1	*	*	*	*	0.3	0.5	0.5	0.6	0.9	2.4
1999.....	57.3	*	*	*	0.3	0.5	0.5	0.4	0.6	1.0	2.0

Non-Hispanic White, female

2007 5/.....	59.1	*	*	*	*	*	*	0.4	0.6	1.2	2.8
2006 5/.....	58.0	*	*	*	*	*	*	0.3	0.7	1.0	2.8
2005 5/.....	61.1	*	*	*	*	*	*	0.5	0.7	1.3	2.9
2004 5/.....	56.7	*	*	*	*	*	*	0.5	0.5	1.4	2.7
2003 5/.....	59.0	*	*	*	*	*	0.3	0.5	0.7	1.4	3.0
2002.....	57.7	*	*	*	*	*	0.5	0.5	0.7	1.6	2.5
2001.....	57.2	*	*	*	*	*	*	0.3	0.8	1.2	2.8
2000.....	56.2	*	*	*	*	*	0.4	0.5	0.7	1.3	2.5
1999.....	55.9	*	*	*	*	0.4	0.4	0.5	0.7	1.1	2.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic Black, both sexes

2007 5/.....	20.5	*	1.2	0.8	1.1	0.8	1.2	1.5	2.0	2.4	4.7
2006 5/.....	20.3	*	*	0.9	1.2	1.1	1.3	1.6	2.4	2.7	5.4
2005 5/.....	21.8	*	1.2	0.7	0.9	0.8	1.6	1.3	2.4	2.7	5.2
2004 5/.....	19.9	*	0.8	0.9	1.1	1.3	1.2	1.8	2.1	3.1	4.9
2003 5/.....	20.9	*	1.0	0.7	1.2	1.3	1.4	1.8	2.2	3.0	5.8
2002.....	21.4	*	1.2	0.7	1.6	1.4	1.2	1.7	2.2	3.9	6.5
2001.....	21.1	*	0.9	0.7	0.9	1.0	1.7	1.9	3.1	4.7	5.1
2000.....	21.5	*	0.9	0.7	1.4	1.5	1.4	1.9	2.5	3.6	6.3
1999.....	22.6	*	*	*	1.5	1.2	1.6	2.6	2.8	3.3	6.2

Non-Hispanic Black, male

2007 5/.....	22.9	*	*	*	1.5	*	1.7	1.8	2.5	2.6	5.1
2006 5/.....	22.7	*	*	*	1.6	1.4	1.4	1.8	3.0	2.6	5.3
2005 5/.....	24.7	*	1.6	*	1.3	1.2	1.8	1.6	2.5	2.8	5.2
2004 5/.....	22.8	*	*	1.4	1.3	1.7	1.4	2.5	2.7	3.3	5.0
2003 5/.....	24.2	*	*	*	1.6	1.6	1.6	2.5	3.3	3.0	5.2
2002.....	24.9	*	1.6	*	1.6	2.3	1.6	2.3	2.2	4.6	6.9
2001.....	24.5	*	*	*	*	1.4	1.9	2.2	3.4	4.4	5.1
2000.....	25.1	*	*	*	1.8	2.1	1.6	2.4	3.3	3.8	6.1
1999.....	27.0	*	*	*	2.1	1.5	2.3	2.8	2.6	3.3	6.2

Non-Hispanic Black, female

2007 5/.....	18.4	*	*	*	*	*	*	*	1.5	2.3	4.2
2006 5/.....	18.1	*	*	*	*	*	*	1.4	1.9	2.8	5.6
2005 5/.....	19.1	*	*	*	*	*	1.5	*	2.3	2.5	5.3
2004 5/.....	17.2	*	*	*	*	*	*	*	1.4	2.9	4.8
2003 5/.....	17.9	*	*	*	*	*	*	*	*	3.0	6.4
2002.....	18.2	*	*	*	1.5	*	*	*	2.2	3.3	6.2
2001.....	18.0	*	*	*	*	*	1.5	1.6	2.7	4.9	5.2
2000.....	18.2	*	*	*	*	*	*	1.5	1.7	3.4	6.4
1999.....	18.6	*	*	*	*	*	*	2.4	2.9	3.3	6.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Asthma (J45-J46)

All origins 3/, both sexes

2007.....	1.1	*	0.2	0.2	0.3	0.3	0.3	0.5	0.5	0.6	0.9
2006.....	1.2	*	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.7	1.0
2005.....	1.3	*	0.2	0.2	0.3	0.2	0.4	0.5	0.6	0.7	1.0
2004.....	1.3	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7	0.9
2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7	1.1
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9	1.2
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8	1.2
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9	1.1

All origins 3/, male

2007.....	0.9	*	0.2	0.3	0.4	0.4	0.5	0.6	0.6	0.5	0.7
2006.....	0.9	*	*	0.3	0.4	0.3	0.4	0.5	0.5	0.6	0.8
2005.....	0.9	*	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5	0.8
2004.....	1.0	*	*	0.2	0.4	0.5	0.5	0.5	0.6	0.6	0.8
2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6	0.9
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8	0.9
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8	0.9
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7	1.0
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8	0.8

All origins 3/, female

2007.....	1.4	*	0.2	*	0.2	0.2	0.2	0.4	0.4	0.7	1.0
2006.....	1.5	*	*	*	*	0.2	0.3	0.4	0.5	0.7	1.2
2005.....	1.7	*	*	*	*	*	0.3	0.4	0.7	0.8	1.2
2004.....	1.6	*	*	*	0.2	0.3	0.3	0.5	0.4	0.9	1.1
2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9	1.4
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0	1.4
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0	1.3
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9	1.4
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Asthma (J45-J46)

Non-Hispanic 4/, both sexes

2007.....	1.3	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.7	1.0
2006.....	1.3	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6	0.7	1.1
2005.....	1.4	*	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	1.1
2004.....	1.4	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.8	1.0
2003.....	1.5	*	0.2	0.2	0.4	0.4	0.5	0.6	0.7	0.8	1.2
2002.....	1.6	*	0.3	0.2	0.4	0.5	0.5	0.6	0.7	1.0	1.2
2001.....	1.6	*	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.0	1.1
2000.....	1.7	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.3
1999.....	1.8	*	0.2	0.2	0.4	0.5	0.5	0.7	0.7	0.9	1.1

Non-Hispanic 4/, male

2007.....	0.9	*	*	0.3	0.4	0.4	0.5	0.6	0.7	0.6	0.8
2006.....	1.0	*	*	0.3	0.4	0.3	0.5	0.6	0.6	0.6	0.9
2005.....	1.0	*	0.4	0.3	0.3	0.4	0.5	0.6	0.6	0.6	0.8
2004.....	1.0	*	*	0.3	0.4	0.5	0.5	0.6	0.6	0.7	0.8
2003.....	1.1	*	0.4	0.3	0.4	0.4	0.5	0.7	0.8	0.6	0.9
2002.....	1.2	*	0.4	*	0.5	0.7	0.5	0.7	0.7	0.9	1.0
2001.....	1.1	*	*	0.3	0.3	0.4	0.5	0.6	0.7	0.9	1.0
2000.....	1.2	*	*	0.2	0.5	0.5	0.6	0.7	0.8	0.8	1.1
1999.....	1.3	*	*	0.3	0.6	0.6	0.7	0.7	0.7	0.9	0.8

Non-Hispanic 4/, female

2007.....	1.6	*	*	*	*	*	*	0.4	0.5	0.8	1.1
2006.....	1.6	*	*	*	*	*	0.4	0.4	0.6	0.8	1.3
2005.....	1.9	*	*	*	*	*	0.4	0.5	0.7	0.9	1.3
2004.....	1.7	*	*	*	0.3	0.3	0.4	0.5	0.4	1.0	1.1
2003.....	1.9	*	*	*	0.3	0.3	0.4	0.5	0.6	0.9	1.5
2002.....	2.0	*	*	*	0.4	0.2	0.4	0.5	0.7	1.1	1.4
2001.....	2.1	*	*	*	0.3	0.3	0.4	0.5	0.8	1.1	1.3
2000.....	2.1	*	*	*	0.4	0.3	0.5	0.5	0.7	1.0	1.5
1999.....	2.2	*	*	*	0.3	0.5	0.4	0.6	0.8	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Asthma (J45-J46)

Non-Hispanic White, both sexes

2007 5/.....	1.1	*	*	*	0.2	0.2	0.2	0.4	0.4	0.5	0.7
2006 5/.....	1.1	*	*	*	*	*	0.3	0.3	0.3	0.5	0.8
2005 5/.....	1.2	*	*	*	*	0.2	0.2	0.4	0.4	0.5	0.8
2004 5/.....	1.2	*	*	*	0.2	0.2	0.3	0.3	0.3	0.6	0.7
2003 5/.....	1.3	*	*	*	0.2	0.2	0.3	0.4	0.5	0.5	0.9
2002.....	1.4	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7	0.7
2001.....	1.4	*	*	*	0.2	0.2	0.3	0.3	0.5	0.6	0.8
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.4	0.5	0.5	0.9
1999.....	1.5	*	*	*	0.2	0.4	0.4	0.4	0.5	0.6	0.8

Non-Hispanic White, male

2007 5/.....	0.7	*	*	*	*	*	*	0.4	0.4	0.3	0.6
2006 5/.....	0.7	*	*	*	*	*	*	0.4	*	0.4	0.7
2005 5/.....	0.8	*	*	*	*	*	0.3	0.4	0.3	0.4	0.6
2004 5/.....	0.8	*	*	*	*	*	0.3	*	0.4	0.5	0.5
2003 5/.....	0.9	*	*	*	*	*	0.4	0.4	0.4	0.4	0.6
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5	0.6
2001.....	0.9	*	*	*	*	*	*	0.3	0.3	0.5	0.6
2000.....	1.0	*	*	*	*	*	0.4	0.4	0.5	0.4	0.7
1999.....	1.0	*	*	*	0.3	0.4	0.4	0.3	0.4	0.5	0.4

Non-Hispanic White, female

2007 5/.....	1.4	*	*	*	*	*	*	0.3	0.4	0.6	0.8
2006 5/.....	1.5	*	*	*	*	*	*	*	0.4	0.6	1.0
2005 5/.....	1.7	*	*	*	*	*	*	0.5	0.5	0.7	1.0
2004 5/.....	1.6	*	*	*	*	*	*	0.4	*	0.8	0.8
2003 5/.....	1.7	*	*	*	*	*	*	0.4	0.5	0.7	1.1
2002.....	1.8	*	*	*	*	*	0.4	0.4	0.6	0.9	0.9
2001.....	1.9	*	*	*	*	*	*	*	0.6	0.6	1.0
2000.....	1.9	*	*	*	*	*	0.3	*	0.5	0.7	1.0
1999.....	2.1	*	*	*	*	0.4	*	0.4	0.5	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Asthma (J45-J46)

Non-Hispanic Black, both sexes

2007 5/.....	2.3	*	1.0	0.8	1.1	0.7	1.2	1.2	1.6	1.8	2.7
2006 5/.....	2.5	*	*	0.8	1.1	1.1	1.2	1.4	2.0	2.2	2.8
2005 5/.....	2.7	*	1.1	0.7	0.9	0.7	1.5	1.3	1.9	1.9	2.9
2004 5/.....	2.7	*	*	0.8	1.0	1.3	1.1	1.7	1.8	2.1	2.9
2003 5/.....	2.8	*	0.8	0.7	1.1	1.1	1.3	1.5	2.0	2.1	3.5
2002.....	3.0	*	0.9	*	1.6	1.3	1.2	1.6	1.6	3.0	4.1
2001.....	3.1	*	*	0.7	0.9	0.9	1.4	1.8	2.7	3.5	3.1
2000.....	3.3	*	*	0.7	1.4	1.3	1.2	1.8	2.1	2.7	3.7
1999.....	3.3	*	*	*	1.5	1.1	1.5	2.4	2.4	2.7	3.3

Non-Hispanic Black, male

2007 5/.....	2.2	*	*	*	1.4	*	1.7	1.6	2.3	1.9	2.4
2006 5/.....	2.3	*	*	*	1.5	1.4	1.4	1.6	2.5	2.0	2.8
2005 5/.....	2.3	*	1.6	*	1.3	*	1.6	1.6	1.9	1.8	2.6
2004 5/.....	2.6	*	*	*	1.3	1.6	1.3	2.4	2.4	1.9	2.7
2003 5/.....	2.5	*	*	*	1.5	1.3	1.4	2.2	2.9	2.0	2.7
2002.....	2.8	*	*	*	1.6	2.2	1.6	2.3	1.7	3.5	3.8
2001.....	2.7	*	*	*	*	*	1.6	2.0	3.0	3.1	2.9
2000.....	2.8	*	*	*	1.7	1.8	*	2.3	2.7	2.9	3.2
1999.....	2.9	*	*	*	2.0	1.3	2.1	2.8	2.6	2.7	2.9

Non-Hispanic Black, female

2007 5/.....	2.5	*	*	*	*	*	*	*	*	1.7	2.9
2006 5/.....	2.7	*	*	*	*	*	*	*	1.6	2.4	2.8
2005 5/.....	3.1	*	*	*	*	*	1.4	*	2.0	1.9	3.2
2004 5/.....	2.8	*	*	*	*	*	*	*	*	2.3	3.1
2003 5/.....	3.0	*	*	*	*	*	*	*	*	2.2	4.2
2002.....	3.1	*	*	*	1.5	*	*	*	1.5	2.4	4.3
2001.....	3.4	*	*	*	*	*	*	1.5	2.4	3.9	3.3
2000.....	3.7	*	*	*	*	*	*	*	1.5	2.7	4.1
1999.....	3.6	*	*	*	*	*	*	2.1	2.3	2.7	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other chronic lower respiratory diseases (J44,J47)

All origins 3/, both sexes

2007.....	36.8	*	*	*	*	*	*	0.1	0.1	0.4	1.4
2006.....	36.0	*	*	*	*	*	*	*	0.2	0.3	1.4
2005.....	37.8	*	*	*	*	*	*	*	0.2	0.5	1.5
2004.....	35.3	*	*	*	*	*	*	*	0.2	0.5	1.4
2003.....	36.6	*	*	*	*	*	*	*	0.1	0.6	1.4
2002.....	36.1	*	*	*	*	*	*	*	0.2	0.5	1.4
2001.....	35.7	*	*	*	*	*	*	*	0.2	0.5	1.3
2000.....	35.4	*	*	*	*	*	*	*	*	0.4	1.2
1999.....	36.0	*	*	*	*	*	*	0.1	0.1	0.3	1.2

All origins 3/, male

2007.....	35.7	*	*	*	*	*	*	*	*	0.4	1.5
2006.....	34.8	*	*	*	*	*	*	*	*	0.3	1.3
2005.....	36.6	*	*	*	*	*	*	*	*	0.5	1.4
2004.....	34.5	*	*	*	*	*	*	*	0.2	0.5	1.5
2003.....	35.9	*	*	*	*	*	*	*	*	0.6	1.3
2002.....	35.7	*	*	*	*	*	*	*	0.2	0.4	1.6
2001.....	35.3	*	*	*	*	*	*	*	0.2	0.5	1.3
2000.....	35.6	*	*	*	*	*	*	*	*	0.4	1.2
1999.....	37.1	*	*	*	*	*	*	*	*	0.3	1.2

All origins 3/, female

2007.....	37.9	*	*	*	*	*	*	*	*	0.4	1.3
2006.....	37.1	*	*	*	*	*	*	*	*	0.3	1.4
2005.....	39.0	*	*	*	*	*	*	*	*	0.5	1.5
2004.....	36.2	*	*	*	*	*	*	*	*	0.4	1.4
2003.....	37.4	*	*	*	*	*	*	*	*	0.5	1.4
2002.....	36.5	*	*	*	*	*	*	*	0.2	0.5	1.3
2001.....	36.0	*	*	*	*	*	*	*	*	0.5	1.4
2000.....	35.2	*	*	*	*	*	*	*	*	0.5	1.2
1999.....	35.0	*	*	*	*	*	*	*	*	0.3	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes

2007.....	9.5	7.5	0.6	0.2	0.2	0.3	0.4	0.5	0.6	0.9	1.4
2006.....	9.2	8.1	0.6	0.2	0.2	0.2	0.2	0.4	0.6	0.9	1.5
2005.....	9.1	7.4	0.6	0.2	0.1	0.2	0.3	0.3	0.6	0.7	1.4
2004.....	9.3	6.5	0.7	0.2	0.1	0.3	0.3	0.4	0.5	0.9	1.5
2003.....	8.8	7.0	0.5	0.1	0.2	0.3	0.4	0.4	0.6	0.9	1.3
2002.....	8.7	6.9	0.5	0.2	0.2	0.3	0.4	0.4	0.5	0.9	1.5
2001.....	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.5
2000.....	9.1	7.2	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.7
1999.....	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.6

All origins 3/, male

2007.....	9.6	8.7	0.8	*	*	0.3	0.5	0.5	0.6	0.9	1.5
2006.....	9.3	9.2	0.7	0.2	*	0.3	0.3	0.4	0.5	1.0	1.5
2005.....	9.0	8.9	0.7	*	*	0.3	0.3	0.4	0.5	0.7	1.4
2004.....	9.3	6.6	0.7	*	*	0.2	0.3	0.4	0.5	0.9	1.5
2003.....	8.7	8.0	0.6	*	0.2	0.3	0.4	0.5	0.6	0.9	1.4
2002.....	8.5	7.9	0.7	0.2	*	0.3	0.4	0.3	0.5	0.9	1.6
2001.....	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9	1.8
2000.....	9.0	7.9	0.6	0.2	*	0.3	0.4	0.5	0.8	1.2	1.8
1999.....	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1	1.7

All origins 3/, female

2007.....	9.3	6.2	0.4	0.2	*	0.2	0.4	0.4	0.6	0.9	1.4
2006.....	9.2	6.9	0.5	*	*	0.2	*	0.4	0.7	0.8	1.5
2005.....	9.2	5.8	0.5	*	*	*	0.3	0.3	0.6	0.8	1.4
2004.....	9.3	6.4	0.6	*	*	0.3	0.3	0.4	0.5	0.9	1.5
2003.....	8.9	5.9	0.5	*	*	0.3	0.3	0.3	0.6	0.9	1.3
2002.....	8.8	5.8	0.4	0.2	0.2	0.2	0.4	0.4	0.6	0.8	1.4
2001.....	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0	1.3
2000.....	9.2	6.4	0.6	*	0.2	0.3	0.3	0.5	0.7	0.9	1.6
1999.....	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 4/, both sexes

2007.....	10.4	8.1	0.6	0.2	0.2	0.3	0.4	0.5	0.6	1.0	1.5
2006.....	10.2	8.6	0.7	0.2	0.2	0.2	0.2	0.4	0.7	1.0	1.6
2005.....	10.0	7.8	0.6	0.2	0.1	0.2	0.3	0.3	0.6	0.8	1.5
2004.....	10.2	6.7	0.7	0.2	0.1	0.3	0.3	0.4	0.6	0.9	1.6
2003.....	9.6	7.0	0.6	*	0.1	0.3	0.4	0.5	0.6	0.9	1.4
2002.....	9.5	7.3	0.5	0.2	0.2	0.3	0.4	0.4	0.6	0.9	1.5
2001.....	10.0	7.8	0.6	0.2	0.2	0.3	0.4	0.6	0.6	1.1	1.6
2000.....	9.8	7.5	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.8
1999.....	9.7	6.9	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.1	1.6

Non-Hispanic 4/, male

2007.....	10.7	9.0	0.8	*	*	0.4	0.5	0.6	0.6	1.0	1.5
2006.....	10.4	9.8	0.8	*	*	0.3	0.3	0.4	0.6	1.0	1.5
2005.....	10.0	9.6	0.8	*	*	0.3	0.3	0.3	0.5	0.7	1.4
2004.....	10.3	6.9	0.8	*	*	0.2	0.4	0.4	0.6	1.0	1.6
2003.....	9.5	8.3	0.6	*	*	0.3	0.4	0.6	0.7	0.9	1.4
2002.....	9.4	8.7	0.7	*	*	0.3	0.4	0.3	0.5	1.0	1.7
2001.....	9.8	10.0	0.6	*	0.2	0.3	0.5	0.5	0.6	1.0	1.8
2000.....	9.8	8.4	0.6	0.2	*	0.3	0.5	0.5	0.8	1.1	1.9
1999.....	9.6	7.9	0.8	0.2	*	0.4	0.4	0.5	0.9	1.2	1.7

Non-Hispanic 4/, female

2007.....	10.2	7.1	0.4	*	*	0.3	0.4	0.5	0.7	1.0	1.4
2006.....	10.0	7.3	0.5	*	*	*	*	0.4	0.8	0.9	1.6
2005.....	10.1	5.9	0.4	*	*	*	0.3	0.3	0.7	0.8	1.5
2004.....	10.1	6.5	0.6	*	*	0.3	0.3	0.4	0.5	0.9	1.6
2003.....	9.7	5.7	0.5	*	*	0.3	0.3	0.3	0.6	0.9	1.4
2002.....	9.6	5.9	0.4	*	0.3	0.2	0.4	0.4	0.6	0.9	1.4
2001.....	10.2	5.5	0.6	*	0.2	0.3	0.4	0.7	0.6	1.1	1.4
2000.....	9.9	6.6	0.5	*	*	0.3	0.3	0.6	0.7	0.9	1.6
1999.....	9.8	5.9	0.5	*	0.2	*	0.3	0.6	0.8	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic White, both sexes

2007 5/.....	11.7	5.9	0.5	*	*	0.3	0.4	0.5	0.6	0.8	1.3
2006 5/.....	11.4	5.4	0.6	0.2	*	0.2	0.2	0.5	0.6	0.9	1.4
2005 5/.....	11.1	6.0	0.5	0.2	*	0.2	0.3	0.3	0.5	0.7	1.3
2004 5/.....	11.4	4.5	0.6	*	*	0.3	0.3	0.3	0.5	0.7	1.4
2003 5/.....	10.7	4.9	0.4	*	*	0.3	0.3	0.4	0.6	0.8	1.1
2002.....	10.5	4.7	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.4
2001.....	11.0	4.8	0.5	0.2	0.2	0.3	0.4	0.5	0.5	0.8	1.4
2000.....	10.8	5.4	0.5	0.2	0.2	0.3	0.3	0.5	0.7	0.8	1.6
1999.....	10.6	4.9	0.6	0.2	0.2	0.3	0.3	0.4	0.7	0.9	1.3

Non-Hispanic White, male

2007 5/.....	11.9	6.8	0.6	*	*	0.3	0.5	0.5	0.5	0.8	1.4
2006 5/.....	11.6	5.6	0.7	*	*	*	*	0.4	0.5	1.0	1.3
2005 5/.....	11.0	7.3	0.7	*	*	*	*	0.3	0.4	0.7	1.3
2004 5/.....	11.4	4.6	0.6	*	*	*	0.4	*	0.5	0.7	1.4
2003 5/.....	10.6	5.5	0.4	*	*	*	0.3	0.5	0.6	0.8	1.2
2002.....	10.3	5.6	0.5	*	*	*	0.4	*	0.4	0.8	1.4
2001.....	10.8	6.2	0.4	*	*	0.3	0.4	0.4	0.4	0.8	1.7
2000.....	10.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9	1.7
1999.....	10.5	5.6	0.6	*	*	0.4	0.4	0.5	0.8	1.0	1.5

Non-Hispanic White, female

2007 5/.....	11.4	4.9	*	*	*	*	0.3	0.4	0.6	0.8	1.3
2006 5/.....	11.2	5.2	*	*	*	*	*	0.5	0.8	0.8	1.4
2005 5/.....	11.2	4.8	*	*	*	*	*	*	0.6	0.7	1.3
2004 5/.....	11.3	4.4	0.5	*	*	*	*	0.4	0.4	0.8	1.4
2003 5/.....	10.8	4.4	*	*	*	*	*	*	0.5	0.7	1.0
2002.....	10.7	3.7	*	*	*	*	*	0.4	0.6	0.8	1.3
2001.....	11.3	3.4	0.5	*	*	*	0.3	0.6	0.5	0.8	1.1
2000.....	11.0	4.1	0.5	*	*	*	*	0.5	0.6	0.7	1.5
1999.....	10.7	4.0	0.5	*	*	*	*	0.4	0.7	0.8	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic Black, both sexes

2007 5/.....	6.7	16.4	1.2	*	*	*	0.7	0.9	1.2	1.8	2.4
2006 5/.....	6.4	20.3	0.9	*	*	*	*	*	1.1	1.5	2.9
2005 5/.....	6.6	15.7	0.9	*	*	*	*	*	1.2	1.5	2.8
2004 5/.....	6.4	15.3	1.1	*	*	*	*	*	1.1	2.3	3.1
2003 5/.....	6.0	15.3	1.1	*	*	*	0.9	0.8	1.2	1.9	3.1
2002.....	6.2	17.3	1.2	*	*	*	0.9	*	1.1	1.8	2.9
2001.....	6.9	19.0	1.1	*	*	*	0.8	1.2	1.4	2.6	3.3
2000.....	6.7	17.7	0.8	*	*	*	0.8	0.9	1.4	2.4	3.1
1999.....	6.7	15.8	1.1	*	*	*	*	1.1	1.7	2.4	3.4

Non-Hispanic Black, male

2007 5/.....	6.8	18.1	1.7	*	*	*	*	*	*	2.1	2.5
2006 5/.....	6.4	24.7	*	*	*	*	*	*	*	1.7	3.0
2005 5/.....	6.7	20.5	*	*	*	*	*	*	*	*	2.8
2004 5/.....	6.5	16.4	*	*	*	*	*	*	*	2.9	3.4
2003 5/.....	6.2	18.6	*	*	*	*	*	*	*	2.0	2.6
2002.....	6.3	20.3	*	*	*	*	*	*	*	2.0	3.6
2001.....	6.6	24.3	*	*	*	*	*	*	1.6	2.0	3.2
2000.....	6.8	17.1	*	*	*	*	*	*	*	3.0	3.5
1999.....	6.6	18.0	*	*	*	*	*	*	1.8	2.5	3.2

Non-Hispanic Black, female

2007 5/.....	6.6	14.5	*	*	*	*	*	*	*	1.6	2.3
2006 5/.....	6.4	15.8	*	*	*	*	*	*	*	*	2.9
2005 5/.....	6.5	10.7	*	*	*	*	*	*	*	1.8	2.9
2004 5/.....	6.3	14.2	*	*	*	*	*	*	*	1.8	2.9
2003 5/.....	5.9	11.9	*	*	*	*	*	*	*	1.8	3.6
2002.....	6.2	14.2	*	*	*	*	*	*	*	1.7	2.3
2001.....	7.2	13.4	*	*	*	*	*	*	*	3.0	3.4
2000.....	6.6	18.3	*	*	*	*	*	*	*	1.9	2.7
1999.....	6.9	13.6	*	*	*	*	*	1.8	1.6	2.3	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes

2007.....	9.7	*	*	*	*	*	0.1	0.5	1.4	3.5	8.3
2006.....	9.2	*	*	*	*	*	0.1	0.3	1.3	3.3	8.2
2005.....	9.3	*	*	*	*	*	*	0.4	1.2	3.4	8.6
2004.....	9.2	*	*	*	*	*	*	0.3	1.3	3.7	8.7
2003.....	9.5	*	*	*	*	*	*	0.4	1.4	4.1	9.3
2002.....	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2	9.7
2001.....	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0	9.8
2000.....	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8	10.2
1999.....	9.4	*	*	*	*	*	*	0.4	1.6	4.8	9.9

All origins 3/, male

2007.....	12.9	*	*	*	*	*	*	0.6	1.9	4.6	11.2
2006.....	12.1	*	*	*	*	*	*	0.4	1.7	4.3	10.8
2005.....	12.3	*	*	*	*	*	*	0.5	1.4	4.5	11.5
2004.....	12.1	*	*	*	*	*	*	0.3	1.8	4.9	12.2
2003.....	12.5	*	*	*	*	*	*	0.6	1.9	5.4	12.8
2002.....	12.3	*	*	*	*	*	*	0.5	1.8	5.5	12.9
2001.....	12.4	*	*	*	*	*	*	0.6	2.0	6.4	13.2
2000.....	12.5	*	*	*	*	*	*	0.6	2.1	6.3	14.4
1999.....	12.5	*	*	*	*	*	*	0.4	2.1	6.3	14.5

All origins 3/, female

2007.....	6.5	*	*	*	*	*	*	0.4	0.8	2.4	5.4
2006.....	6.4	*	*	*	*	*	*	0.2	0.8	2.4	5.6
2005.....	6.4	*	*	*	*	*	*	0.3	0.9	2.3	5.7
2004.....	6.3	*	*	*	*	*	*	*	0.7	2.5	5.4
2003.....	6.5	*	*	*	*	*	*	0.3	0.8	2.9	5.9
2002.....	6.7	*	*	*	*	*	*	0.3	1.0	3.0	6.5
2001.....	6.7	*	*	*	*	*	*	0.3	1.0	3.5	6.4
2000.....	6.5	*	*	*	*	*	*	0.3	1.1	3.2	6.1
1999.....	6.4	*	*	*	*	*	*	0.3	1.2	3.4	5.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 4/, both sexes

2007	9.8	*	*	*	*	*	0.1	0.5	1.2	3.4	8.1
2006	9.4	*	*	*	*	*	0.1	0.3	1.1	3.1	8.0
2005	9.4	*	*	*	*	*	*	0.3	1.1	3.2	8.3
2004	9.3	*	*	*	*	*	*	0.3	1.2	3.6	8.3
2003	9.6	*	*	*	*	*	*	0.4	1.1	4.0	9.0
2002	9.5	*	*	*	*	*	*	0.4	1.3	4.1	9.2
2001	9.5	*	*	*	*	*	*	0.4	1.4	4.8	9.2
2000	9.4	*	*	*	*	*	0.1	0.5	1.4	4.6	9.7
1999	9.5	*	*	*	*	*	*	0.4	1.5	4.5	9.3

Non-Hispanic 4/, male

2007	13.0	*	*	*	*	*	*	0.6	1.7	4.2	10.4
2006	12.3	*	*	*	*	*	*	0.4	1.4	3.8	10.2
2005	12.3	*	*	*	*	*	*	0.4	1.2	4.0	10.7
2004	12.3	*	*	*	*	*	*	0.3	1.6	4.7	11.3
2003	12.6	*	*	*	*	*	*	0.5	1.5	4.9	12.1
2002	12.2	*	*	*	*	*	*	0.4	1.6	5.1	11.8
2001	12.3	*	*	*	*	*	*	0.5	1.8	5.9	12.1
2000	12.4	*	*	*	*	*	*	0.6	1.7	5.9	13.3
1999	12.5	*	*	*	*	*	*	0.4	1.8	5.6	13.4

Non-Hispanic 4/, female

2007	6.8	*	*	*	*	*	*	0.4	0.8	2.5	5.8
2006	6.6	*	*	*	*	*	*	*	0.8	2.5	5.8
2005	6.6	*	*	*	*	*	*	*	1.0	2.4	5.8
2004	6.5	*	*	*	*	*	*	*	0.7	2.6	5.5
2003	6.7	*	*	*	*	*	*	0.3	0.7	3.1	6.0
2002	6.9	*	*	*	*	*	*	0.3	1.1	3.0	6.7
2001	6.9	*	*	*	*	*	*	0.3	1.0	3.7	6.5
2000	6.7	*	*	*	*	*	*	0.3	1.2	3.3	6.1
1999	6.6	*	*	*	*	*	*	0.3	1.2	3.4	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic White, both sexes

2007 5/.....	10.7	*	*	*	*	*	*	0.5	1.3	3.6	8.4
2006 5/.....	10.3	*	*	*	*	*	*	0.3	1.1	3.2	8.2
2005 5/.....	10.2	*	*	*	*	*	*	0.3	1.0	3.4	8.6
2004 5/.....	10.1	*	*	*	*	*	*	0.2	1.2	3.6	8.5
2003 5/.....	10.3	*	*	*	*	*	*	0.3	1.1	4.0	9.2
2002.....	10.2	*	*	*	*	*	*	0.3	1.3	4.1	9.4
2001.....	10.1	*	*	*	*	*	*	0.3	1.4	4.8	9.0
2000.....	10.0	*	*	*	*	*	*	0.3	1.3	4.4	9.4
1999.....	9.9	*	*	*	*	*	*	0.4	1.4	4.4	8.8

Non-Hispanic White, male

2007 5/.....	14.2	*	*	*	*	*	*	0.5	1.8	4.5	10.8
2006 5/.....	13.4	*	*	*	*	*	*	0.4	1.4	4.0	10.5
2005 5/.....	13.3	*	*	*	*	*	*	0.4	1.1	4.2	11.1
2004 5/.....	13.2	*	*	*	*	*	*	*	1.6	4.6	11.5
2003 5/.....	13.6	*	*	*	*	*	*	0.4	1.5	5.0	12.5
2002.....	13.2	*	*	*	*	*	*	0.4	1.6	5.1	12.2
2001.....	13.0	*	*	*	*	*	*	0.4	1.8	5.8	11.7
2000.....	13.1	*	*	*	*	*	*	0.5	1.6	5.7	13.1
1999.....	13.1	*	*	*	*	*	*	0.5	1.8	5.5	12.7

Non-Hispanic White, female

2007 5/.....	7.4	*	*	*	*	*	*	0.4	0.8	2.8	6.0
2006 5/.....	7.2	*	*	*	*	*	*	*	0.7	2.4	6.0
2005 5/.....	7.2	*	*	*	*	*	*	*	0.8	2.5	6.1
2004 5/.....	7.1	*	*	*	*	*	*	*	0.7	2.7	5.5
2003 5/.....	7.2	*	*	*	*	*	*	*	0.7	3.1	6.0
2002.....	7.5	*	*	*	*	*	*	*	0.9	3.1	6.7
2001.....	7.4	*	*	*	*	*	*	*	1.0	3.8	6.3
2000.....	7.0	*	*	*	*	*	*	*	1.0	3.1	5.8
1999.....	6.9	*	*	*	*	*	*	*	1.1	3.3	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcoholic liver disease (K70)

All origins 3/, both sexes

2007.....	4.8	*	*	*	*	*	*	0.4	1.1	2.6	5.8
2006.....	4.4	*	*	*	*	*	*	0.3	1.0	2.4	5.5
2005.....	4.4	*	*	*	*	*	*	0.3	0.8	2.4	5.7
2004.....	4.3	*	*	*	*	*	*	0.2	0.9	2.5	5.8
2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9	5.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8	5.9
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.1
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3	6.6
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.4

All origins 3/, male

2007.....	7.1	*	*	*	*	*	*	0.5	1.5	3.4	7.8
2006.....	6.4	*	*	*	*	*	*	0.4	1.4	3.1	7.1
2005.....	6.5	*	*	*	*	*	*	0.4	1.0	3.2	7.5
2004.....	6.3	*	*	*	*	*	*	0.3	1.3	3.3	8.0
2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7	8.0
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6	8.0
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4	8.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3	9.4
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3	9.4

All origins 3/, female

2007.....	2.5	*	*	*	*	*	*	0.3	0.6	1.8	3.9
2006.....	2.4	*	*	*	*	*	*	*	0.6	1.7	3.8
2005.....	2.3	*	*	*	*	*	*	*	0.6	1.6	3.9
2004.....	2.3	*	*	*	*	*	*	*	0.5	1.8	3.6
2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0	3.8
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0	3.9
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2	3.9
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcoholic liver disease (K70)

Hispanic, both sexes

2007.....	4.7	*	*	*	*	*	*	0.6	1.4	3.0	6.7
2006.....	4.3	*	*	*	*	*	*	*	1.6	3.0	6.5
2005.....	4.2	*	*	*	*	*	*	*	1.2	3.2	6.9
2004.....	4.1	*	*	*	*	*	*	*	1.3	3.1	7.8
2003.....	4.3	*	*	*	*	*	*	*	1.6	3.8	7.7
2002.....	4.3	*	*	*	*	*	*	*	1.1	3.5	7.9
2001.....	4.7	*	*	*	*	*	*	*	1.5	4.1	8.8
2000.....	4.8	*	*	*	*	*	*	*	1.7	4.3	10.0
1999.....	4.6	*	*	*	*	*	*	*	1.5	4.7	9.7

Hispanic, male

2007.....	7.5	*	*	*	*	*	*	*	2.1	4.5	10.9
2006.....	6.9	*	*	*	*	*	*	*	2.3	4.5	9.6
2005.....	6.8	*	*	*	*	*	*	*	1.8	5.0	10.5
2004.....	6.6	*	*	*	*	*	*	*	2.1	4.6	12.4
2003.....	6.9	*	*	*	*	*	*	*	2.3	6.3	11.8
2002.....	7.1	*	*	*	*	*	*	*	1.8	5.7	13.0
2001.....	7.7	*	*	*	*	*	*	*	2.4	6.9	13.8
2000.....	7.9	*	*	*	*	*	*	*	3.0	6.4	16.0
1999.....	7.4	*	*	*	*	*	*	*	2.3	7.1	15.6

Hispanic, female

2007.....	1.6	*	*	*	*	*	*	*	*	1.2	2.0
2006.....	1.6	*	*	*	*	*	*	*	*	1.3	3.0
2005.....	1.4	*	*	*	*	*	*	*	*	1.3	2.9
2004.....	1.4	*	*	*	*	*	*	*	*	1.3	2.8
2003.....	1.5	*	*	*	*	*	*	*	*	*	3.4
2002.....	1.4	*	*	*	*	*	*	*	*	*	2.5
2001.....	1.5	*	*	*	*	*	*	*	*	*	3.6
2000.....	1.4	*	*	*	*	*	*	*	*	2.1	3.8
1999.....	1.5	*	*	*	*	*	*	*	*	2.1	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcoholic liver disease (K70)

Non-Hispanic 4/, both sexes

2007	4.8	*	*	*	*	*	*	0.4	1.0	2.5	5.7
2006	4.4	*	*	*	*	*	*	0.3	0.8	2.2	5.3
2005	4.4	*	*	*	*	*	*	0.3	0.7	2.3	5.5
2004	4.3	*	*	*	*	*	*	0.2	0.8	2.4	5.5
2003	4.2	*	*	*	*	*	*	0.3	0.8	2.7	5.6
2002	4.2	*	*	*	*	*	*	0.2	0.9	2.6	5.6
2001	4.2	*	*	*	*	*	*	0.2	1.0	3.2	5.8
2000	4.2	*	*	*	*	*	*	0.3	1.0	3.1	6.2
1999	4.2	*	*	*	*	*	*	0.2	1.0	3.1	6.0

Non-Hispanic 4/, male

2007	7.0	*	*	*	*	*	*	0.4	1.3	3.2	7.2
2006	6.3	*	*	*	*	*	*	0.4	1.1	2.7	6.6
2005	6.4	*	*	*	*	*	*	0.4	0.8	2.8	6.9
2004	6.3	*	*	*	*	*	*	0.3	1.2	3.0	7.4
2003	6.2	*	*	*	*	*	*	0.4	1.2	3.3	7.5
2002	6.0	*	*	*	*	*	*	0.3	1.1	3.2	7.3
2001	6.2	*	*	*	*	*	*	*	1.2	4.0	7.7
2000	6.2	*	*	*	*	*	*	0.4	1.2	4.0	8.6
1999	6.4	*	*	*	*	*	*	0.2	1.2	3.9	8.6

Non-Hispanic 4/, female

2007	2.7	*	*	*	*	*	*	0.3	0.6	1.9	4.2
2006	2.5	*	*	*	*	*	*	*	0.6	1.8	4.0
2005	2.5	*	*	*	*	*	*	*	0.6	1.7	4.0
2004	2.4	*	*	*	*	*	*	*	0.5	1.9	3.7
2003	2.3	*	*	*	*	*	*	*	0.4	2.1	3.9
2002	2.4	*	*	*	*	*	*	*	0.7	2.0	4.1
2001	2.4	*	*	*	*	*	*	*	0.7	2.4	3.9
2000	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9
1999	2.2	*	*	*	*	*	*	*	0.8	2.4	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcoholic liver disease (K70)

Non-Hispanic White, both sexes

2007 5/.....	5.2	*	*	*	*	*	*	0.4	1.0	2.8	5.9
2006 5/.....	4.7	*	*	*	*	*	*	0.2	0.8	2.3	5.6
2005 5/.....	4.7	*	*	*	*	*	*	0.2	0.6	2.4	5.6
2004 5/.....	4.6	*	*	*	*	*	*	0.2	0.9	2.5	5.7
2003 5/.....	4.5	*	*	*	*	*	*	0.2	0.8	2.7	5.8
2002.....	4.4	*	*	*	*	*	*	*	0.9	2.6	5.8
2001.....	4.4	*	*	*	*	*	*	*	1.0	3.2	5.7
2000.....	4.3	*	*	*	*	*	*	0.2	0.9	3.0	6.0
1999.....	4.3	*	*	*	*	*	*	0.2	0.9	3.0	5.7

Non-Hispanic White, male

2007 5/.....	7.6	*	*	*	*	*	*	0.4	1.3	3.4	7.5
2006 5/.....	6.9	*	*	*	*	*	*	0.3	1.1	2.9	7.1
2005 5/.....	6.8	*	*	*	*	*	*	0.3	0.7	3.0	7.2
2004 5/.....	6.7	*	*	*	*	*	*	*	1.2	2.9	7.6
2003 5/.....	6.6	*	*	*	*	*	*	0.4	1.1	3.3	7.7
2002.....	6.4	*	*	*	*	*	*	*	1.1	3.1	7.4
2001.....	6.4	*	*	*	*	*	*	*	1.2	4.0	7.6
2000.....	6.5	*	*	*	*	*	*	0.4	1.2	3.8	8.3
1999.....	6.5	*	*	*	*	*	*	*	1.1	3.8	8.1

Non-Hispanic White, female

2007 5/.....	2.9	*	*	*	*	*	*	0.3	0.6	2.1	4.4
2006 5/.....	2.7	*	*	*	*	*	*	*	0.5	1.7	4.1
2005 5/.....	2.6	*	*	*	*	*	*	*	0.5	1.7	4.0
2004 5/.....	2.5	*	*	*	*	*	*	*	0.5	2.0	3.8
2003 5/.....	2.4	*	*	*	*	*	*	*	0.5	2.0	4.0
2002.....	2.4	*	*	*	*	*	*	*	0.6	2.1	4.2
2001.....	2.4	*	*	*	*	*	*	*	0.7	2.5	3.9
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2	3.7
1999.....	2.2	*	*	*	*	*	*	*	0.7	2.3	3.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/, both sexes

2007.....	4.9	*	*	*	*	*	*	0.1	0.3	0.9	2.5
2006.....	4.8	*	*	*	*	*	*	*	0.3	1.0	2.7
2005.....	4.9	*	*	*	*	*	*	*	0.4	1.0	2.9
2004.....	4.9	*	*	*	*	*	*	*	0.3	1.2	2.9
2003.....	5.2	*	*	*	*	*	*	0.2	0.4	1.3	3.4
2002.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5	3.7
2001.....	5.2	*	*	*	*	*	*	0.2	0.4	1.6	3.6
2000.....	5.1	*	*	*	*	*	0.1	0.2	0.5	1.5	3.6
1999.....	5.1	*	*	*	*	*	*	0.2	0.6	1.5	3.5

All origins 3/, male

2007.....	5.8	*	*	*	*	*	*	*	0.5	1.2	3.4
2006.....	5.7	*	*	*	*	*	*	*	0.3	1.2	3.7
2005.....	5.8	*	*	*	*	*	*	*	0.4	1.3	4.0
2004.....	5.8	*	*	*	*	*	*	*	0.5	1.7	4.1
2003.....	6.2	*	*	*	*	*	*	*	0.5	1.6	4.8
2002.....	6.1	*	*	*	*	*	*	*	0.6	1.9	4.9
2001.....	6.1	*	*	*	*	*	*	0.2	0.6	2.0	4.8
2000.....	6.0	*	*	*	*	*	*	*	0.6	2.0	5.0
1999.....	6.0	*	*	*	*	*	*	*	0.7	2.0	5.2

All origins 3/, female

2007.....	4.0	*	*	*	*	*	*	*	0.2	0.6	1.6
2006.....	4.0	*	*	*	*	*	*	*	0.2	0.7	1.8
2005.....	4.0	*	*	*	*	*	*	*	0.4	0.7	1.9
2004.....	4.1	*	*	*	*	*	*	*	0.2	0.7	1.8
2003.....	4.3	*	*	*	*	*	*	*	0.3	0.9	2.0
2002.....	4.5	*	*	*	*	*	*	0.2	0.4	1.0	2.6
2001.....	4.4	*	*	*	*	*	*	*	0.3	1.3	2.5
2000.....	4.3	*	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	4.3	*	*	*	*	*	*	*	0.4	1.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 4/, both sexes

2007.....	5.1	*	*	*	*	*	*	*	0.3	0.8	2.4
2006.....	5.0	*	*	*	*	*	*	*	0.3	0.9	2.7
2005.....	5.1	*	*	*	*	*	*	*	0.4	1.0	2.8
2004.....	5.0	*	*	*	*	*	*	*	0.3	1.2	2.8
2003.....	5.4	*	*	*	*	*	*	0.2	0.3	1.3	3.3
2002.....	5.4	*	*	*	*	*	*	0.2	0.4	1.4	3.6
2001.....	5.3	*	*	*	*	*	*	0.2	0.4	1.5	3.4
2000.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5	3.5
1999.....	5.2	*	*	*	*	*	*	0.2	0.5	1.4	3.3

Non-Hispanic 4/, male

2007.....	6.0	*	*	*	*	*	*	*	0.4	1.0	3.2
2006.....	6.0	*	*	*	*	*	*	*	0.3	1.1	3.5
2005.....	6.0	*	*	*	*	*	*	*	0.4	1.2	3.8
2004.....	6.0	*	*	*	*	*	*	*	0.4	1.7	3.9
2003.....	6.4	*	*	*	*	*	*	*	0.4	1.6	4.6
2002.....	6.2	*	*	*	*	*	*	*	0.5	1.9	4.5
2001.....	6.2	*	*	*	*	*	*	0.3	0.6	1.9	4.4
2000.....	6.1	*	*	*	*	*	*	*	0.5	1.9	4.7
1999.....	6.1	*	*	*	*	*	*	*	0.6	1.7	4.8

Non-Hispanic 4/, female

2007.....	4.1	*	*	*	*	*	*	*	*	0.6	1.6
2006.....	4.1	*	*	*	*	*	*	*	*	0.7	1.9
2005.....	4.1	*	*	*	*	*	*	*	0.4	0.7	1.8
2004.....	4.1	*	*	*	*	*	*	*	0.2	0.7	1.8
2003.....	4.4	*	*	*	*	*	*	*	0.3	1.0	2.1
2002.....	4.6	*	*	*	*	*	*	*	0.3	1.0	2.6
2001.....	4.5	*	*	*	*	*	*	*	0.3	1.2	2.5
2000.....	4.4	*	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	4.4	*	*	*	*	*	*	*	0.4	1.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic White, both sexes

2007 5/.....	5.6	*	*	*	*	*	*	*	0.3	0.9	2.5
2006 5/.....	5.5	*	*	*	*	*	*	*	0.3	0.9	2.7
2005 5/.....	5.5	*	*	*	*	*	*	*	0.3	1.0	3.0
2004 5/.....	5.5	*	*	*	*	*	*	*	0.3	1.2	2.7
2003 5/.....	5.9	*	*	*	*	*	*	*	0.3	1.4	3.4
2002.....	5.9	*	*	*	*	*	*	*	0.4	1.5	3.6
2001.....	5.8	*	*	*	*	*	*	*	0.5	1.6	3.3
2000.....	5.7	*	*	*	*	*	*	*	0.4	1.4	3.4
1999.....	5.6	*	*	*	*	*	*	*	0.5	1.4	3.2

Non-Hispanic White, male

2007 5/.....	6.6	*	*	*	*	*	*	*	0.4	1.1	3.3
2006 5/.....	6.5	*	*	*	*	*	*	*	0.3	1.1	3.4
2005 5/.....	6.5	*	*	*	*	*	*	*	0.3	1.3	4.0
2004 5/.....	6.5	*	*	*	*	*	*	*	0.4	1.7	3.8
2003 5/.....	6.9	*	*	*	*	*	*	*	0.4	1.7	4.8
2002.....	6.7	*	*	*	*	*	*	*	0.5	2.0	4.8
2001.....	6.6	*	*	*	*	*	*	*	0.6	1.8	4.2
2000.....	6.6	*	*	*	*	*	*	*	0.4	1.9	4.8
1999.....	6.6	*	*	*	*	*	*	*	0.6	1.7	4.6

Non-Hispanic White, female

2007 5/.....	4.6	*	*	*	*	*	*	*	*	0.6	1.6
2006 5/.....	4.6	*	*	*	*	*	*	*	*	0.7	2.0
2005 5/.....	4.6	*	*	*	*	*	*	*	0.3	0.8	2.0
2004 5/.....	4.5	*	*	*	*	*	*	*	*	0.7	1.7
2003 5/.....	4.8	*	*	*	*	*	*	*	*	1.0	2.0
2002.....	5.0	*	*	*	*	*	*	*	*	1.0	2.5
2001.....	5.0	*	*	*	*	*	*	*	0.4	1.3	2.4
2000.....	4.8	*	*	*	*	*	*	*	0.4	1.0	2.1
1999.....	4.8	*	*	*	*	*	*	*	0.4	1.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes

2007.....	15.4	3.4	0.1	*	*	0.1	0.3	0.5	0.8	1.3	2.2
2006.....	15.1	3.9	*	*	*	0.1	0.3	0.4	1.0	1.3	2.3
2005.....	14.8	3.9	*	*	*	0.2	0.3	0.6	0.9	1.2	2.1
2004.....	14.5	4.3	*	*	*	0.1	0.3	0.5	0.7	1.4	2.2
2003.....	14.6	4.5	*	*	*	0.1	0.3	0.5	0.9	1.3	2.3
2002.....	14.2	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3	2.1
2001.....	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3	2.1
2000.....	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2	2.1
1999.....	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0

All origins 3/, male

2007.....	15.2	3.6	*	*	*	*	0.3	0.5	0.9	1.4	2.5
2006.....	15.0	4.8	*	*	*	*	0.4	0.4	1.2	1.4	2.7
2005.....	14.6	4.2	*	*	*	*	0.3	0.5	1.0	1.3	2.5
2004.....	14.1	5.3	*	*	*	*	0.3	0.5	0.6	1.5	2.6
2003.....	14.3	5.2	*	*	*	*	0.4	0.5	0.9	1.4	2.7
2002.....	13.9	5.3	*	*	*	*	0.3	0.5	0.8	1.4	2.3
2001.....	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4	2.3
2000.....	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3	2.5
1999.....	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3	2.3

All origins 3/, female

2007.....	15.6	3.2	*	*	*	*	0.3	0.5	0.7	1.2	1.8
2006.....	15.3	3.0	*	*	*	*	0.3	0.4	0.8	1.1	1.9
2005.....	15.0	3.5	*	*	*	0.2	0.3	0.6	0.8	1.1	1.7
2004.....	14.8	3.2	*	*	*	*	0.2	0.5	0.8	1.3	1.8
2003.....	14.9	3.8	*	*	*	*	0.3	0.5	0.8	1.1	1.9
2002.....	14.5	3.2	*	*	*	*	0.3	0.7	0.7	1.1	1.9
2001.....	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3	1.9
2000.....	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1	1.6
1999.....	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 4/, both sexes

2007.....	17.1	3.3	0.2	*	*	*	0.3	0.6	0.9	1.5	2.3
2006.....	16.7	4.0	*	*	*	0.1	0.4	0.5	1.0	1.4	2.4
2005.....	16.4	3.9	*	*	*	0.2	0.3	0.6	0.9	1.3	2.2
2004.....	15.9	4.3	*	*	*	0.1	0.3	0.5	0.8	1.5	2.3
2003.....	16.0	4.6	*	*	*	0.1	0.3	0.6	1.0	1.3	2.3
2002.....	15.6	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3	2.1
2001.....	15.1	3.5	*	*	*	*	0.4	0.5	0.7	1.4	2.1
2000.....	14.4	4.4	*	*	*	0.1	0.3	0.5	0.8	1.3	2.1
1999.....	13.8	4.8	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0

Non-Hispanic 4/, male

2007.....	17.0	3.3	*	*	*	*	0.3	0.6	1.0	1.5	2.7
2006.....	16.7	5.1	*	*	*	*	0.4	0.4	1.3	1.6	2.9
2005.....	16.2	4.2	*	*	*	*	0.3	0.6	1.0	1.4	2.6
2004.....	15.6	5.2	*	*	*	*	0.3	0.5	0.8	1.6	2.6
2003.....	15.8	5.4	*	*	*	*	0.4	0.6	1.0	1.5	2.8
2002.....	15.4	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.3
2001.....	14.8	3.7	*	*	*	*	0.3	0.5	0.7	1.5	2.2
2000.....	14.1	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.5
1999.....	13.5	4.8	*	*	*	*	0.3	0.5	0.9	1.4	2.3

Non-Hispanic 4/, female

2007.....	17.2	3.4	*	*	*	*	0.3	0.5	0.8	1.4	1.8
2006.....	16.8	2.9	*	*	*	*	0.3	0.5	0.8	1.2	2.0
2005.....	16.5	3.5	*	*	*	*	0.3	0.6	0.8	1.2	1.8
2004.....	16.2	3.4	*	*	*	*	0.2	0.5	0.9	1.4	2.0
2003.....	16.2	3.6	*	*	*	*	0.3	0.6	0.9	1.2	1.9
2002.....	15.8	3.2	*	*	*	*	0.3	0.7	0.7	1.2	1.9
2001.....	15.4	3.3	*	*	*	*	0.4	0.4	0.8	1.3	1.9
2000.....	14.7	3.4	*	*	*	*	0.4	0.4	0.7	1.2	1.7
1999.....	14.1	4.8	*	*	*	*	0.3	0.6	0.8	1.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic White, both sexes

2007 5/.....	17.0	2.7	*	*	*	*	0.2	0.3	0.6	0.9	1.5
2006 5/.....	16.5	3.4	*	*	*	*	0.3	0.3	0.7	0.9	1.5
2005 5/.....	16.2	2.7	*	*	*	*	0.2	0.4	0.6	0.9	1.3
2004 5/.....	15.8	2.6	*	*	*	*	0.2	0.4	0.5	1.0	1.6
2003 5/.....	15.8	3.5	*	*	*	*	0.2	0.4	0.6	0.8	1.4
2002.....	15.4	3.3	*	*	*	*	0.2	0.4	0.6	0.9	1.4
2001.....	14.9	2.4	*	*	*	*	0.3	0.3	0.5	0.9	1.3
2000.....	14.2	2.7	*	*	*	*	0.3	0.4	0.5	0.8	1.4
1999.....	13.4	3.5	*	*	*	*	0.2	0.3	0.5	0.8	1.2

Non-Hispanic White, male

2007 5/.....	17.1	2.4	*	*	*	*	*	0.4	0.7	1.0	1.8
2006 5/.....	16.7	4.6	*	*	*	*	0.4	*	1.0	1.0	1.8
2005 5/.....	16.2	2.7	*	*	*	*	*	0.4	0.7	1.0	1.7
2004 5/.....	15.6	3.1	*	*	*	*	*	0.4	0.4	1.0	1.9
2003 5/.....	15.7	4.4	*	*	*	*	*	0.4	0.6	0.8	1.8
2002.....	15.3	4.1	*	*	*	*	*	0.3	0.7	0.9	1.5
2001.....	14.8	2.5	*	*	*	*	*	*	0.4	1.2	1.5
2000.....	14.1	3.5	*	*	*	*	*	0.4	0.6	0.9	1.7
1999.....	13.3	3.9	*	*	*	*	*	0.3	0.6	0.9	1.4

Non-Hispanic White, female

2007 5/.....	16.9	3.0	*	*	*	*	*	*	0.5	0.7	1.2
2006 5/.....	16.4	2.1	*	*	*	*	*	*	0.5	0.7	1.1
2005 5/.....	16.2	2.7	*	*	*	*	*	0.5	0.5	0.8	1.0
2004 5/.....	16.0	2.1	*	*	*	*	*	*	0.7	1.0	1.2
2003 5/.....	15.9	2.5	*	*	*	*	*	0.4	0.6	0.7	1.0
2002.....	15.6	2.5	*	*	*	*	*	0.4	0.5	0.8	1.3
2001.....	15.0	2.3	*	*	*	*	0.3	*	0.6	0.7	1.2
2000.....	14.2	1.8	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	13.5	3.1	*	*	*	*	*	0.4	0.5	0.6	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic Black, both sexes

2007 5/.....	21.8	6.7	*	*	*	*	0.7	1.6	2.5	5.1	7.1
2006 5/.....	22.1	6.8	*	*	*	*	0.8	1.4	2.6	4.2	7.9
2005 5/.....	21.5	9.1	*	*	*	*	*	1.7	2.9	3.7	7.6
2004 5/.....	21.0	10.4	*	*	*	*	0.8	1.4	2.5	4.3	7.1
2003 5/.....	21.4	9.5	*	*	*	*	0.8	1.5	3.1	4.6	8.2
2002.....	20.5	8.5	*	*	*	*	0.9	1.9	2.3	4.1	6.8
2001.....	20.2	8.5	*	*	*	*	0.8	1.4	2.2	4.1	6.9
2000.....	19.5	12.0	*	*	*	*	*	1.1	2.7	4.5	6.5
1999.....	19.2	10.2	*	*	*	*	0.8	1.5	2.4	4.2	7.2

Non-Hispanic Black, male

2007 5/.....	20.5	7.7	*	*	*	*	*	1.7	2.9	4.9	9.0
2006 5/.....	21.0	7.4	*	*	*	*	*	*	3.2	4.6	9.4
2005 5/.....	20.3	11.2	*	*	*	*	*	1.8	3.0	3.9	8.8
2004 5/.....	19.9	12.4	*	*	*	*	*	*	3.0	5.4	7.4
2003 5/.....	20.8	9.9	*	*	*	*	*	*	3.7	5.6	9.1
2002.....	19.7	10.6	*	*	*	*	*	1.7	2.3	4.5	7.8
2001.....	18.6	8.9	*	*	*	*	*	1.8	2.6	3.4	7.2
2000.....	18.3	13.6	*	*	*	*	*	*	3.0	5.3	8.2
1999.....	18.1	8.7	*	*	*	*	*	*	2.4	5.0	8.5

Non-Hispanic Black, female

2007 5/.....	23.0	*	*	*	*	*	*	1.5	2.2	5.3	5.4
2006 5/.....	23.0	*	*	*	*	*	*	1.5	2.2	3.8	6.7
2005 5/.....	22.5	6.8	*	*	*	*	*	1.6	2.7	3.6	6.5
2004 5/.....	22.0	8.4	*	*	*	*	*	1.6	2.2	3.3	6.9
2003 5/.....	21.9	9.0	*	*	*	*	*	*	2.5	3.7	7.4
2002.....	21.2	6.3	*	*	*	*	*	2.1	2.2	3.7	6.0
2001.....	21.7	8.1	*	*	*	*	*	*	1.9	4.6	6.7
2000.....	20.7	10.2	*	*	*	*	*	*	2.5	3.9	5.0
1999.....	20.2	11.8	*	*	*	*	*	1.7	2.4	3.4	6.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Renal failure (N17-N19)

All origins 3/, both sexes

2007.....	14.3	3.2	*	*	*	*	0.3	0.4	0.7	1.2	2.0
2006.....	14.5	3.7	*	*	*	0.1	0.3	0.4	0.9	1.2	2.2
2005.....	14.5	3.7	*	*	*	0.1	0.2	0.5	0.8	1.2	2.0
2004.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3	2.1
2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2	2.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2	2.0
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2	2.0
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1	1.9

All origins 3/, male

2007.....	14.2	3.4	*	*	*	*	0.3	0.5	0.8	1.4	2.3
2006.....	14.3	4.4	*	*	*	*	0.3	0.3	1.0	1.3	2.6
2005.....	14.2	4.1	*	*	*	*	0.3	0.5	0.9	1.2	2.3
2004.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4	2.5
2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3	2.6
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3	2.2
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3	2.2
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3	2.4
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3	2.2

All origins 3/, female

2007.....	14.5	3.1	*	*	*	*	0.3	0.4	0.7	1.1	1.6
2006.....	14.6	3.0	*	*	*	*	0.3	0.4	0.7	1.1	1.8
2005.....	14.7	3.2	*	*	*	*	0.2	0.5	0.7	1.1	1.7
2004.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2	1.8
2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1	1.8
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1	1.8
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2	1.8
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0	1.6
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Renal failure (N17-N19)

Non-Hispanic 4/, both sexes

2007.....	15.9	3.2	*	*	*	*	0.3	0.5	0.8	1.4	2.1
2006.....	16.0	3.7	*	*	*	0.1	0.3	0.4	0.9	1.3	2.4
2005.....	16.0	3.7	*	*	*	0.1	0.2	0.6	0.9	1.2	2.1
2004.....	15.7	4.1	*	*	*	0.1	0.3	0.5	0.8	1.4	2.2
2003.....	15.7	4.3	*	*	*	*	0.3	0.5	0.9	1.3	2.3
2002.....	15.3	4.1	*	*	*	*	0.3	0.6	0.7	1.3	2.1
2001.....	14.9	3.3	*	*	*	*	0.3	0.4	0.7	1.3	2.0
2000.....	14.1	4.2	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0
1999.....	13.5	4.7	*	*	*	0.1	0.2	0.5	0.8	1.1	1.9

Non-Hispanic 4/, male

2007.....	15.8	3.2	*	*	*	*	0.3	0.6	0.9	1.5	2.5
2006.....	15.9	4.6	*	*	*	*	0.4	0.4	1.2	1.4	2.8
2005.....	15.8	4.1	*	*	*	*	0.2	0.5	1.0	1.3	2.5
2004.....	15.3	5.1	*	*	*	*	0.3	0.5	0.7	1.6	2.5
2003.....	15.5	5.0	*	*	*	*	0.3	0.5	1.0	1.4	2.7
2002.....	15.1	5.1	*	*	*	*	0.3	0.5	0.8	1.4	2.2
2001.....	14.6	3.5	*	*	*	*	0.3	0.5	0.7	1.4	2.2
2000.....	13.8	5.2	*	*	*	*	*	0.4	0.8	1.4	2.4
1999.....	13.2	4.6	*	*	*	*	*	0.4	0.8	1.3	2.2

Non-Hispanic 4/, female

2007.....	16.0	3.2	*	*	*	*	0.3	0.4	0.8	1.2	1.7
2006.....	16.1	2.8	*	*	*	*	0.3	0.5	0.7	1.1	1.9
2005.....	16.2	3.2	*	*	*	*	0.2	0.6	0.8	1.2	1.8
2004.....	16.0	3.1	*	*	*	*	*	0.4	0.8	1.3	1.9
2003.....	15.9	3.5	*	*	*	*	0.3	0.5	0.8	1.2	1.9
2002.....	15.5	2.9	*	*	*	*	0.3	0.6	0.7	1.1	1.9
2001.....	15.2	3.1	*	*	*	*	0.3	0.4	0.7	1.3	1.8
2000.....	14.4	3.1	*	*	*	*	0.3	0.4	0.7	1.1	1.7
1999.....	13.8	4.8	*	*	*	*	0.3	0.5	0.7	0.9	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Renal failure (N17-N19)

Non-Hispanic White, both sexes

2007 5/.....	15.8	2.5	*	*	*	*	0.2	0.3	0.6	0.8	1.4
2006 5/.....	15.8	3.1	*	*	*	*	0.3	0.3	0.7	0.8	1.4
2005 5/.....	15.8	2.5	*	*	*	*	0.2	0.4	0.5	0.8	1.3
2004 5/.....	15.5	2.5	*	*	*	*	0.2	0.3	0.4	1.0	1.5
2003 5/.....	15.5	3.2	*	*	*	*	0.2	0.4	0.6	0.7	1.4
2002.....	15.2	3.1	*	*	*	*	0.2	0.3	0.5	0.8	1.3
2001.....	14.6	2.2	*	*	*	*	0.2	0.3	0.4	0.9	1.3
2000.....	13.9	2.5	*	*	*	*	0.2	0.3	0.4	0.8	1.3
1999.....	13.1	3.5	*	*	*	*	*	0.3	0.5	0.7	1.2

Non-Hispanic White, male

2007 5/.....	15.9	2.1	*	*	*	*	*	0.3	0.6	1.0	1.6
2006 5/.....	15.9	4.1	*	*	*	*	0.4	*	0.9	0.9	1.8
2005 5/.....	15.8	2.6	*	*	*	*	*	0.3	0.6	0.9	1.6
2004 5/.....	15.3	3.0	*	*	*	*	*	0.4	0.3	1.0	1.9
2003 5/.....	15.4	4.0	*	*	*	*	*	0.4	0.6	0.8	1.7
2002.....	15.0	3.9	*	*	*	*	*	*	0.6	0.9	1.4
2001.....	14.5	2.3	*	*	*	*	*	*	0.4	1.1	1.5
2000.....	13.7	3.3	*	*	*	*	*	0.3	0.5	0.8	1.6
1999.....	13.0	3.8	*	*	*	*	*	*	0.5	0.8	1.4

Non-Hispanic White, female

2007 5/.....	15.8	2.9	*	*	*	*	*	*	0.5	0.6	1.1
2006 5/.....	15.7	2.1	*	*	*	*	*	*	0.4	0.7	1.1
2005 5/.....	15.9	2.3	*	*	*	*	*	0.5	0.5	0.8	1.0
2004 5/.....	15.7	2.0	*	*	*	*	*	*	0.6	1.0	1.1
2003 5/.....	15.6	2.3	*	*	*	*	*	0.4	0.6	0.7	1.0
2002.....	15.3	2.2	*	*	*	*	*	0.4	0.4	0.7	1.2
2001.....	14.7	2.1	*	*	*	*	*	*	0.5	0.7	1.1
2000.....	14.0	*	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	13.2	3.1	*	*	*	*	*	*	0.5	0.6	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Renal failure (N17-N19)

Non-Hispanic Black, both sexes

2007 5/.....	20.3	6.5	*	*	*	*	0.7	1.4	2.4	4.8	6.6
2006 5/.....	21.1	6.6	*	*	*	*	0.7	1.2	2.4	3.9	7.7
2005 5/.....	21.0	8.7	*	*	*	*	*	1.5	2.7	3.6	7.3
2004 5/.....	20.7	10.1	*	*	*	*	0.7	1.2	2.4	4.1	6.8
2003 5/.....	21.1	9.3	*	*	*	*	0.7	1.4	2.9	4.6	8.1
2002.....	20.2	8.0	*	*	*	*	0.8	1.8	2.1	4.0	6.8
2001.....	20.0	8.5	*	*	*	*	0.8	1.3	2.2	3.9	6.8
2000.....	19.2	11.4	*	*	*	*	*	1.0	2.6	4.2	6.3
1999.....	18.8	10.2	*	*	*	*	*	1.5	2.3	3.9	6.9

Non-Hispanic Black, male

2007 5/.....	19.1	7.7	*	*	*	*	*	1.6	2.6	4.9	8.3
2006 5/.....	20.0	7.1	*	*	*	*	*	*	2.7	4.4	9.3
2005 5/.....	19.8	11.2	*	*	*	*	*	*	2.9	3.7	8.3
2004 5/.....	19.6	12.1	*	*	*	*	*	*	2.9	5.3	7.1
2003 5/.....	20.5	9.6	*	*	*	*	*	*	3.6	5.6	9.0
2002.....	19.4	10.0	*	*	*	*	*	*	2.1	4.5	7.6
2001.....	18.4	8.9	*	*	*	*	*	1.7	2.6	3.3	7.0
2000.....	17.9	13.6	*	*	*	*	*	*	2.8	5.1	8.0
1999.....	17.7	8.7	*	*	*	*	*	*	2.3	4.7	8.3

Non-Hispanic Black, female

2007 5/.....	21.4	*	*	*	*	*	*	*	2.2	4.7	5.1
2006 5/.....	22.1	*	*	*	*	*	*	1.4	2.1	3.5	6.3
2005 5/.....	22.1	*	*	*	*	*	*	1.5	2.6	3.6	6.4
2004 5/.....	21.7	8.0	*	*	*	*	*	*	2.0	3.1	6.6
2003 5/.....	21.7	9.0	*	*	*	*	*	*	2.4	3.7	7.2
2002.....	20.9	*	*	*	*	*	*	2.0	2.1	3.6	6.0
2001.....	21.4	8.1	*	*	*	*	*	*	1.8	4.5	6.6
2000.....	20.3	9.2	*	*	*	*	*	*	2.4	3.5	4.9
1999.....	19.8	11.8	*	*	*	*	*	1.7	2.3	3.3	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/, both sexes

2007.....	0.3	*	0.2	0.6	0.8	0.9	0.7	0.3
2006.....	0.3	*	0.2	0.7	0.8	0.8	0.7	0.2
2005.....	0.3	*	0.2	0.7	0.8	0.8	0.6	0.3
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.5	0.3
2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5	0.3
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4	0.1
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3	0.1
2000.....	0.1	*	0.4	0.4	0.4	0.5	0.4	0.2
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4	0.1

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	0.5	*	0.4	1.1	1.5	1.7	1.5	0.6
2006.....	0.5	*	0.3	1.4	1.7	1.6	1.3	0.5
2005.....	0.5	*	0.4	1.4	1.6	1.6	1.3	0.6
2004.....	0.5	*	0.4	1.3	1.5	1.4	0.9	0.5
2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0	0.5
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7	0.2
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7	0.3
2000.....	0.3	*	0.8	0.8	0.9	0.9	0.8	0.3
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Hispanic, both sexes

2007.....	0.3	*	*	0.6	0.6	0.8	0.7	*
2006.....	0.3	*	*	0.8	0.8	0.9	0.6	*
2005.....	0.3	*	*	0.6	0.6	1.0	0.6	*
2004.....	0.3	*	*	*	0.7	0.6	*	*
2003.....	0.3	*	*	*	0.8	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	0.6	*	*	*
2000.....	0.2	*	*	0.6	0.6	*	*	*
1999.....	0.2	*	*	*	*	*	0.8	*

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	0.6	*	*	1.2	1.5	1.7	1.5	*
2006.....	0.7	*	*	1.8	1.8	2.0	1.3	*
2005.....	0.6	*	*	1.4	1.3	2.1	1.4	*
2004.....	0.6	*	*	*	1.5	1.3	*	*
2003.....	0.5	*	*	*	1.8	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	1.3	*	*	*
2000.....	0.5	*	*	1.4	1.3	*	*	*
1999.....	0.4	*	*	*	*	*	1.6	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	0.2	0.5	0.8	0.9	0.7	0.3
2006.....	0.2	*	0.1	0.7	0.8	0.7	0.7	0.2
2005.....	0.2	*	0.2	0.7	0.8	0.7	0.6	0.3
2004.....	0.2	*	0.1	0.7	0.7	0.7	0.4	0.3
2003.....	0.2	*	*	0.4	0.5	0.6	0.5	0.2
2002.....	0.1	*	0.1	0.3	0.5	0.4	0.3	0.1
2001.....	0.1	*	0.2	0.3	0.5	0.5	0.3	0.1
2000.....	0.1	*	*	0.4	0.4	0.5	0.3	0.2
1999.....	0.1	*	0.2	0.3	0.4	0.5	0.3	0.1

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	0.5	*	0.4	1.1	1.6	1.7	1.5	0.5
2006.....	0.5	*	0.3	1.4	1.7	1.4	1.3	0.4
2005.....	0.5	*	0.3	1.4	1.6	1.5	1.3	0.6
2004.....	0.4	*	0.3	1.3	1.5	1.5	0.9	0.5
2003.....	0.3	*	*	0.8	1.0	1.2	1.0	0.5
2002.....	0.2	*	0.3	0.6	0.9	0.8	0.7	0.2
2001.....	0.3	*	0.3	0.7	0.9	0.9	0.6	0.3
2000.....	0.3	*	*	0.7	0.8	0.9	0.7	0.3
1999.....	0.3	*	0.3	0.6	0.9	1.0	0.7	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic White, both sexes

2007 5/.....	0.2	*	*	0.4	0.5	0.6	0.5	0.2
2006 5/.....	0.1	*	*	0.5	0.5	0.5	0.4	0.2
2005 5/.....	0.2	*	*	0.5	0.5	0.5	0.4	0.2
2004 5/.....	0.2	*	*	0.4	0.4	0.6	0.3	0.2
2003 5/.....	0.1	*	*	0.3	0.4	0.3	0.3	0.2
2002.....	0.1	*	*	*	0.3	0.3	0.2	*
2001.....	0.1	*	*	0.2	0.3	0.3	0.2	*
2000.....	0.1	*	*	0.2	0.3	0.3	0.2	*
1999.....	0.1	*	*	*	0.2	0.3	0.2	*

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	0.3	*	*	0.7	1.0	1.3	1.1	0.4
2006 5/.....	0.3	*	*	1.0	1.0	0.9	0.8	0.3
2005 5/.....	0.3	*	*	1.0	0.9	1.1	0.9	0.5
2004 5/.....	0.3	*	*	0.7	0.9	1.1	0.7	0.4
2003 5/.....	0.2	*	*	0.5	0.8	0.7	0.5	0.3
2002.....	0.1	*	*	*	0.5	0.6	0.3	*
2001.....	0.2	*	*	0.4	0.5	0.7	0.3	*
2000.....	0.2	*	*	0.4	0.5	0.6	0.5	*
1999.....	0.2	*	*	*	0.5	0.7	0.4	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic Black, both sexes

2007 5/.....	0.7	*	0.6	1.6	1.9	2.0	1.6	*
2006 5/.....	0.7	*	*	1.6	2.3	1.9	1.9	*
2005 5/.....	0.7	*	0.6	1.6	2.4	1.9	1.6	0.9
2004 5/.....	0.7	*	*	2.0	2.3	1.8	1.1	*
2003 5/.....	0.5	*	*	1.0	1.3	1.9	1.5	*
2002.....	0.4	*	*	1.0	1.5	1.0	1.2	*
2001.....	0.4	*	*	1.1	1.4	1.1	1.0	*
2000.....	0.4	*	*	1.1	1.2	1.3	0.9	*
1999.....	0.5	*	*	1.1	1.4	1.2	1.0	*

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	1.2	*	1.2	3.2	3.8	3.8	3.1	*
2006 5/.....	1.3	*	*	3.2	4.6	3.7	3.6	*
2005 5/.....	1.4	*	1.3	3.3	4.7	3.7	2.9	1.6
2004 5/.....	1.3	*	*	4.1	4.4	3.4	2.1	*
2003 5/.....	1.0	*	*	2.0	2.4	3.6	2.8	*
2002.....	0.8	*	*	2.0	2.8	1.9	2.3	*
2001.....	0.8	*	*	2.2	2.7	2.2	1.9	*
2000.....	0.8	*	*	2.2	2.3	2.5	1.7	*
1999.....	0.9	*	*	2.1	2.7	2.4	1.8	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy with abortive outcome (000-007)

All origins 3/, both sexes

2007.....	0.0	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	0.0	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy with abortive outcome (000-007)

Hispanic, both sexes

2007.....	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 4/, both sexes

2007	0.0	*	*	*	*	*	*	*
2006	0.0	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*

Non-Hispanic 4/, male

2007
2006
2005
2004
2003
2002
2001
2000
1999

Non-Hispanic 4/, female

2007	0.0	*	*	*	*	*	*	*
2006	0.0	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy with abortive outcome (000-007)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy with abortive outcome (000-007)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

All origins 3/, both sexes

2007.....	0.2	*	0.2	0.5	0.7	0.8	0.7	0.3
2006.....	0.2	*	0.2	0.7	0.8	0.7	0.6	0.2
2005.....	0.2	*	0.2	0.7	0.7	0.8	0.6	0.3
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.4	0.3
2003.....	0.2	*	0.1	0.4	0.5	0.5	0.5	0.2
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
2001.....	0.1	*	0.2	0.3	0.5	0.4	0.3	0.1
2000.....	0.1	*	0.4	0.4	0.4	0.4	0.3	0.1
1999.....	0.1	*	0.1	0.3	0.4	0.4	0.4	0.1

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	0.5	*	0.4	1.1	1.5	1.6	1.4	0.6
2006.....	0.5	*	0.3	1.4	1.6	1.5	1.3	0.5
2005.....	0.5	*	0.4	1.4	1.5	1.5	1.2	0.6
2004.....	0.4	*	0.4	1.2	1.4	1.4	0.9	0.5
2003.....	0.3	*	0.2	0.8	1.1	1.1	0.9	0.5
2002.....	0.2	*	0.3	0.6	0.8	0.8	0.7	0.2
2001.....	0.3	*	0.3	0.7	0.9	0.9	0.6	0.3
2000.....	0.3	*	0.8	0.8	0.8	0.9	0.7	0.3
1999.....	0.3	*	0.3	0.6	0.9	0.8	0.7	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Hispanic, both sexes

2007.....	0.3	*	*	0.5	0.6	0.8	0.7	*
2006.....	0.3	*	*	0.8	0.8	0.9	0.6	*
2005.....	0.3	*	*	0.6	0.6	0.9	0.6	*
2004.....	0.3	*	*	*	0.7	0.6	*	*
2003.....	0.2	*	*	*	0.7	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	0.6	*	*	*
2000.....	0.2	*	*	0.6	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	0.6	*	*	1.2	1.4	1.7	1.5	*
2006.....	0.7	*	*	1.7	1.8	1.9	1.3	*
2005.....	0.6	*	*	1.4	1.3	2.1	1.4	*
2004.....	0.6	*	*	*	1.5	1.3	*	*
2003.....	0.5	*	*	*	1.6	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	1.3	*	*	*
2000.....	0.5	*	*	1.3	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic 4/, both sexes

2007.....	0.2	*	0.2	0.5	0.8	0.8	0.7	0.3
2006.....	0.2	*	0.1	0.6	0.8	0.7	0.7	0.2
2005.....	0.2	*	0.2	0.7	0.8	0.7	0.6	0.3
2004.....	0.2	*	0.1	0.6	0.7	0.7	0.4	0.2
2003.....	0.2	*	*	0.4	0.5	0.5	0.5	0.2
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
2001.....	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
2000.....	0.1	*	*	0.3	0.4	0.4	0.3	0.1
1999.....	0.1	*	0.2	0.3	0.4	0.4	0.3	0.1

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	0.5	*	0.4	1.1	1.5	1.6	1.4	0.5
2006.....	0.4	*	0.3	1.3	1.6	1.4	1.3	0.4
2005.....	0.5	*	0.3	1.4	1.5	1.4	1.1	0.6
2004.....	0.4	*	0.3	1.2	1.4	1.4	0.8	0.5
2003.....	0.3	*	*	0.8	1.0	1.1	0.9	0.4
2002.....	0.2	*	0.3	0.5	0.9	0.8	0.6	0.2
2001.....	0.2	*	0.3	0.6	0.9	0.8	0.6	0.2
2000.....	0.2	*	*	0.7	0.7	0.8	0.6	0.3
1999.....	0.2	*	0.3	0.6	0.8	0.8	0.6	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic White, both sexes

2007 5/.....	0.2	*	*	0.4	0.5	0.6	0.5	0.2
2006 5/.....	0.1	*	*	0.5	0.5	0.4	0.4	0.2
2005 5/.....	0.2	*	*	0.5	0.5	0.5	0.4	0.2
2004 5/.....	0.1	*	*	0.3	0.4	0.5	0.3	0.2
2003 5/.....	0.1	*	*	0.3	0.3	0.3	0.3	0.2
2002.....	0.1	*	*	*	0.3	0.3	0.2	*
2001.....	0.1	*	*	0.2	0.2	0.3	0.1	*
2000.....	0.1	*	*	0.2	0.2	0.3	0.2	*
1999.....	0.1	*	*	*	0.2	0.3	0.2	*

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	0.3	*	*	0.7	1.0	1.2	1.1	0.4
2006 5/.....	0.3	*	*	1.0	0.9	0.9	0.8	0.3
2005 5/.....	0.3	*	*	1.0	0.9	1.0	0.8	0.5
2004 5/.....	0.3	*	*	0.7	0.9	1.1	0.7	0.4
2003 5/.....	0.2	*	*	0.5	0.7	0.6	0.5	0.3
2002.....	0.1	*	*	*	0.5	0.5	0.3	*
2001.....	0.1	*	*	0.3	0.5	0.6	0.3	*
2000.....	0.1	*	*	0.4	0.5	0.6	0.4	*
1999.....	0.1	*	*	*	0.5	0.6	0.4	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic Black, both sexes

2007 5/.....	0.6	*	*	1.4	1.8	1.8	1.5	*
2006 5/.....	0.6	*	*	1.5	2.2	1.8	1.9	*
2005 5/.....	0.7	*	*	1.6	2.2	1.8	1.4	0.9
2004 5/.....	0.6	*	*	1.8	2.1	1.7	0.9	*
2003 5/.....	0.5	*	*	0.9	1.3	1.7	1.4	*
2002.....	0.4	*	*	0.9	1.4	0.9	1.1	*
2001.....	0.4	*	*	1.0	1.3	1.0	0.9	*
2000.....	0.3	*	*	1.0	1.0	1.1	0.7	*
1999.....	0.4	*	*	1.0	1.3	1.1	0.8	*

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	1.2	*	*	2.9	3.5	3.5	2.9	*
2006 5/.....	1.2	*	*	3.1	4.4	3.4	3.5	*
2005 5/.....	1.3	*	*	3.3	4.3	3.5	2.7	1.6
2004 5/.....	1.2	*	*	3.6	4.1	3.2	1.8	*
2003 5/.....	0.9	*	*	1.8	2.4	3.2	2.6	*
2002.....	0.7	*	*	1.9	2.7	1.8	2.0	*
2001.....	0.7	*	*	2.0	2.5	1.9	1.6	*
2000.....	0.7	*	*	1.9	1.9	2.2	1.4	*
1999.....	0.8	*	*	2.0	2.4	2.1	1.6	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/, both sexes

2007.....	3.5	135.9	3.3	1.0	0.9	0.9	1.0	1.1	1.0	1.1	1.3
2006.....	3.5	140.9	3.2	0.9	0.8	1.1	1.1	1.1	1.0	1.1	1.1
2005.....	3.5	135.2	3.2	1.0	1.0	1.2	1.2	1.2	1.0	1.1	1.2
2004.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0	1.2
2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1	1.4
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2	1.3
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2	1.4
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2	1.3
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2	1.2

All origins 3/, male

2007.....	3.7	140.7	3.4	1.0	0.9	1.0	1.2	1.2	1.1	1.1	1.3
2006.....	3.8	146.6	2.9	0.9	0.9	1.3	1.3	1.4	1.2	1.3	1.2
2005.....	3.7	141.1	3.2	0.9	1.0	1.6	1.4	1.3	1.1	1.2	1.3
2004.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2	1.3
2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2	1.5
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2	1.4
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2	1.6
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4	1.5
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3	1.3

All origins 3/, female

2007.....	3.2	130.8	3.2	1.0	0.8	0.7	0.8	0.9	0.9	1.1	1.3
2006.....	3.2	134.9	3.4	1.0	0.7	0.8	0.9	0.9	0.9	0.9	1.0
2005.....	3.3	129.0	3.3	1.2	0.9	0.8	1.0	1.0	0.9	1.0	1.2
2004.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8	1.2
2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1	1.3
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1	1.3
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 4/, both sexes

2007.....	3.3	131.5	3.3	1.0	0.9	0.9	1.1	1.2	1.0	1.2	1.4
2006.....	3.3	136.8	3.1	1.0	0.8	1.0	1.2	1.2	1.1	1.2	1.1
2005.....	3.3	130.4	3.2	1.0	0.9	1.2	1.2	1.2	1.1	1.2	1.3
2004.....	3.4	135.9	3.4	1.1	0.9	1.3	1.1	1.1	1.1	1.1	1.3
2003.....	3.5	136.0	3.4	0.9	1.0	1.1	1.2	1.1	1.2	1.2	1.4
2002.....	3.6	134.9	3.3	1.0	1.1	1.2	1.3	1.2	1.3	1.3	1.4
2001.....	3.5	132.8	3.7	0.9	0.9	1.3	1.3	1.3	1.2	1.2	1.5
2000.....	3.6	148.8	3.1	1.0	1.0	1.2	1.2	1.2	1.4	1.3	1.4
1999.....	3.6	143.1	3.6	1.0	1.1	1.1	1.2	1.1	1.3	1.3	1.3

Non-Hispanic 4/, male

2007.....	3.5	137.2	3.2	1.0	1.0	1.0	1.2	1.3	1.2	1.2	1.4
2006.....	3.6	143.6	2.7	0.9	0.8	1.2	1.3	1.5	1.2	1.4	1.3
2005.....	3.6	137.1	3.1	0.8	1.0	1.7	1.5	1.4	1.3	1.3	1.3
2004.....	3.6	139.1	3.7	1.0	0.8	1.4	1.3	1.3	1.3	1.3	1.4
2003.....	3.7	144.2	3.3	0.9	1.0	1.4	1.3	1.3	1.2	1.3	1.6
2002.....	3.8	138.1	3.5	1.0	1.1	1.4	1.4	1.6	1.5	1.3	1.5
2001.....	3.7	139.4	3.6	1.0	1.0	1.6	1.5	1.3	1.4	1.3	1.6
2000.....	3.9	154.4	3.1	1.0	1.0	1.4	1.4	1.3	1.7	1.4	1.5
1999.....	3.8	148.4	3.9	1.0	1.1	1.4	1.4	1.2	1.4	1.4	1.4

Non-Hispanic 4/, female

2007.....	3.1	125.6	3.3	1.0	0.9	0.8	0.9	1.0	0.9	1.2	1.4
2006.....	3.0	129.7	3.5	1.0	0.7	0.9	1.0	0.9	1.0	1.0	1.0
2005.....	3.1	123.3	3.3	1.2	0.9	0.8	1.0	1.1	1.0	1.0	1.3
2004.....	3.2	132.6	3.2	1.1	0.9	1.2	0.9	0.9	1.0	0.8	1.2
2003.....	3.3	127.6	3.4	0.9	0.9	0.8	1.0	0.8	1.2	1.2	1.3
2002.....	3.4	131.5	3.2	1.0	1.0	1.0	1.2	0.9	1.2	1.3	1.3
2001.....	3.3	125.8	3.7	0.9	0.9	0.9	1.1	1.2	1.1	1.2	1.4
2000.....	3.4	142.8	3.0	0.9	1.1	1.0	1.0	1.0	1.1	1.1	1.2
1999.....	3.4	137.4	3.3	1.1	1.1	0.9	1.0	1.1	1.2	1.2	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic White, both sexes

2007 5/.....	3.1	123.6	2.9	0.9	0.8	0.8	1.1	1.2	1.0	1.2	1.5
2006 5/.....	3.2	129.5	2.8	1.0	0.7	1.0	1.1	1.2	1.2	1.2	1.2
2005 5/.....	3.2	123.0	2.9	0.9	0.8	1.1	1.2	1.2	1.2	1.2	1.3
2004 5/.....	3.3	130.5	2.9	0.9	0.8	1.2	1.0	1.2	1.2	1.1	1.4
2003 5/.....	3.3	129.8	3.0	0.8	0.9	1.1	1.1	1.1	1.2	1.3	1.5
2002.....	3.4	131.2	3.0	1.0	1.0	1.2	1.3	1.3	1.4	1.3	1.4
2001.....	3.4	127.4	3.1	0.9	0.9	1.2	1.2	1.3	1.3	1.2	1.5
2000.....	3.4	141.2	2.6	0.9	1.0	1.0	1.2	1.2	1.5	1.3	1.3
1999.....	3.4	136.8	3.0	0.9	1.0	1.1	1.1	1.2	1.3	1.3	1.3

Non-Hispanic White, male

2007 5/.....	3.3	131.0	2.9	0.9	0.9	0.8	1.2	1.3	1.1	1.2	1.5
2006 5/.....	3.4	135.2	2.6	1.0	0.8	1.1	1.2	1.5	1.3	1.4	1.3
2005 5/.....	3.4	129.9	3.0	0.7	0.9	1.6	1.5	1.3	1.3	1.3	1.3
2004 5/.....	3.4	132.8	3.1	1.0	0.8	1.3	1.1	1.5	1.3	1.2	1.5
2003 5/.....	3.4	136.1	2.9	0.8	1.0	1.4	1.3	1.3	1.2	1.4	1.7
2002.....	3.6	137.2	3.2	0.9	1.0	1.3	1.4	1.6	1.6	1.3	1.6
2001.....	3.6	135.5	3.1	1.0	1.0	1.5	1.3	1.5	1.5	1.3	1.6
2000.....	3.6	147.9	2.6	0.8	0.9	1.1	1.5	1.3	1.8	1.4	1.5
1999.....	3.6	144.4	3.2	0.8	1.0	1.4	1.4	1.3	1.4	1.5	1.4

Non-Hispanic White, female

2007 5/.....	2.9	115.9	2.9	0.8	0.8	0.7	1.0	1.0	1.0	1.2	1.5
2006 5/.....	2.9	123.5	2.9	0.9	0.5	0.8	1.0	0.8	1.1	0.9	1.1
2005 5/.....	3.0	115.8	2.9	1.1	0.8	0.7	0.9	1.1	1.0	1.2	1.4
2004 5/.....	3.2	128.0	2.8	0.9	0.9	1.1	1.0	0.9	1.2	0.9	1.2
2003 5/.....	3.2	123.2	3.0	0.8	0.9	0.8	0.9	0.9	1.3	1.2	1.4
2002.....	3.3	125.0	2.9	1.0	1.1	1.0	1.2	1.0	1.2	1.3	1.3
2001.....	3.2	118.8	3.2	0.8	0.7	0.9	1.0	1.1	1.2	1.1	1.4
2000.....	3.2	134.2	2.6	0.9	1.0	0.9	1.0	1.0	1.2	1.1	1.2
1999.....	3.3	128.9	2.9	1.0	1.0	0.8	0.9	1.2	1.2	1.2	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic Black, both sexes

2007 5/.....	4.6	166.3	4.6	1.4	1.1	1.3	1.2	1.4	1.7	1.7	1.3
2006 5/.....	4.6	174.3	4.2	1.2	1.0	1.4	1.3	1.6	1.1	1.7	1.1
2005 5/.....	4.4	163.4	4.1	1.2	1.3	1.7	1.5	1.7	1.4	1.3	1.4
2004 5/.....	4.6	164.3	5.1	1.5	1.1	1.9	1.7	1.2	1.1	1.3	1.2
2003 5/.....	4.6	160.5	4.8	1.2	1.1	1.3	1.8	1.3	1.6	1.3	1.2
2002.....	4.7	156.9	4.7	1.2	1.3	1.7	1.6	1.4	1.5	1.4	1.9
2001.....	4.7	163.5	5.4	1.0	1.2	2.0	1.9	1.4	1.2	1.5	1.7
2000.....	5.0	184.1	5.2	1.3	1.4	1.9	1.3	1.4	1.4	1.6	1.7
1999.....	4.8	174.8	5.8	1.5	1.8	1.4	1.5	1.1	1.2	1.5	1.3

Non-Hispanic Black, male

2007 5/.....	5.0	167.8	4.4	1.3	*	1.6	1.6	1.8	2.3	1.8	*
2006 5/.....	5.1	182.2	3.2	*	*	1.7	1.6	1.9	*	1.8	*
2005 5/.....	4.8	167.1	3.6	*	1.3	2.2	1.8	2.2	*	1.9	*
2004 5/.....	5.2	170.5	5.9	*	*	2.1	2.7	*	1.6	1.8	*
2003 5/.....	5.2	173.1	5.0	1.3	*	1.7	1.9	*	*	*	*
2002.....	5.2	153.3	4.8	1.4	1.5	2.1	1.8	1.8	1.7	*	2.1
2001.....	5.1	167.5	5.5	*	*	2.7	2.3	*	*	*	2.2
2000.....	5.5	186.0	5.6	1.7	1.3	2.6	*	1.8	1.7	1.7	2.0
1999.....	5.3	179.5	6.6	1.4	1.8	1.7	1.9	*	*	*	1.8

Non-Hispanic Black, female

2007 5/.....	4.3	164.7	4.8	1.4	*	*	*	*	*	1.5	*
2006 5/.....	4.2	166.0	5.3	1.4	*	*	*	*	*	1.6	*
2005 5/.....	4.1	159.6	4.7	1.4	1.4	*	*	*	1.5	*	1.3
2004 5/.....	4.1	157.8	4.3	1.8	*	1.7	*	*	*	*	*
2003 5/.....	4.0	147.3	4.7	*	*	*	1.6	*	1.6	*	*
2002.....	4.4	160.6	4.7	*	*	*	1.4	*	*	1.5	1.7
2001.....	4.4	159.2	5.4	*	1.4	1.4	1.5	1.8	*	1.7	*
2000.....	4.5	182.2	4.8	*	1.5	*	*	*	*	1.5	*
1999.....	4.4	170.0	5.0	1.5	1.8	*	*	*	*	1.5	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes

2007.....	11.1	85.0	1.4	0.2	0.3	1.0	1.8	2.1	2.2	2.8	3.8
2006.....	10.6	83.8	1.5	0.3	0.3	0.9	2.2	2.4	2.6	3.2	4.2
2005.....	10.8	87.4	1.4	0.3	0.5	1.2	2.5	2.8	3.1	4.1	5.2
2004.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6	4.9
2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8	5.9
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6	6.0
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2	7.7
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7	7.6
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1	5.1

All origins 3/, male

2007.....	9.6	96.3	1.6	0.2	0.4	1.3	2.4	2.8	2.8	3.4	4.8
2006.....	9.5	95.2	1.6	0.4	0.4	1.1	2.9	3.2	3.7	4.0	5.5
2005.....	10.0	101.8	1.6	0.3	0.6	1.6	3.4	3.9	4.0	5.1	6.4
2004.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6	6.5
2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4	7.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1	7.6
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0	9.9
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6	10.2
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3	6.7

All origins 3/, female

2007.....	12.6	73.1	1.3	0.2	0.3	0.6	1.1	1.4	1.6	2.2	2.7
2006.....	11.7	71.9	1.3	0.3	0.2	0.7	1.5	1.5	1.5	2.4	3.0
2005.....	11.5	72.3	1.3	0.3	0.3	0.8	1.4	1.7	2.1	3.1	4.1
2004.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7	3.3
2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3	4.4
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1	4.5
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4	5.5
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7	5.0
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0	3.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, both sexes

2007.....	4.3	54.0	1.4	*	*	1.0	2.0	2.1	1.5	2.0	2.9
2006.....	4.0	52.5	1.2	*	*	0.8	1.8	1.8	2.1	2.5	2.5
2005.....	4.4	59.5	1.2	*	*	0.9	2.0	2.1	2.6	2.9	3.4
2004.....	3.9	50.4	1.1	*	*	1.2	2.0	2.0	2.2	2.8	3.5
2003.....	4.1	56.8	0.9	*	*	1.4	1.4	2.4	2.2	3.0	3.2
2002.....	4.0	56.6	1.3	*	*	1.1	1.6	1.8	2.3	2.6	3.6
2001.....	5.9	60.7	1.6	*	0.6	2.3	3.2	3.7	4.7	5.9	6.1
2000.....	5.9	65.6	1.7	*	*	1.6	2.6	3.5	4.5	6.3	7.2
1999.....	3.8	54.9	1.0	*	*	1.0	1.6	1.8	2.6	2.9	3.1

Hispanic, male

2007.....	4.8	61.3	1.4	*	*	1.7	3.1	2.9	2.2	3.0	4.0
2006.....	4.5	62.1	1.1	*	*	1.3	2.5	2.7	3.1	3.8	4.0
2005.....	5.1	68.9	1.3	*	*	*	2.8	3.3	3.8	4.0	4.7
2004.....	4.5	60.0	1.2	*	*	1.5	3.0	2.4	3.2	3.8	5.2
2003.....	4.8	69.9	*	*	*	1.6	1.8	3.5	3.2	4.3	4.4
2002.....	4.6	66.4	1.6	*	*	1.2	2.1	2.5	3.4	3.6	4.4
2001.....	7.4	67.2	1.8	*	*	3.4	4.3	5.4	6.6	8.3	8.6
2000.....	7.5	78.6	2.1	*	*	1.7	3.9	5.1	6.5	9.3	10.6
1999.....	4.4	61.3	*	*	*	1.3	2.4	2.4	3.5	4.2	4.5

Hispanic, female

2007.....	3.7	46.4	1.4	*	*	*	*	1.1	*	*	1.7
2006.....	3.4	42.5	1.3	*	*	*	*	*	*	*	*
2005.....	3.5	49.8	1.2	*	*	*	*	*	1.3	1.6	2.1
2004.....	3.3	40.4	*	*	*	*	*	1.5	*	1.5	1.7
2003.....	3.2	43.2	*	*	*	*	*	*	*	1.5	1.8
2002.....	3.3	46.4	*	*	*	*	*	*	*	1.4	2.8
2001.....	4.3	53.9	*	*	*	*	1.8	1.7	2.5	3.2	3.6
2000.....	4.1	52.1	1.4	*	*	1.4	*	1.6	2.3	3.0	3.6
1999.....	3.1	48.2	*	*	*	*	*	*	1.6	1.6	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 4/, both sexes

2007.....	12.3	94.5	1.4	0.2	0.3	0.9	1.7	2.1	2.4	3.0	3.9
2006.....	11.7	92.7	1.5	0.4	0.3	0.9	2.3	2.5	2.8	3.3	4.4
2005.....	11.8	94.5	1.5	0.3	0.4	1.3	2.5	2.9	3.2	4.3	5.4
2004.....	11.2	90.9	1.2	0.3	0.3	1.0	2.2	2.8	2.9	3.8	5.0
2003.....	11.8	89.4	1.8	0.4	0.5	1.5	2.8	3.4	3.7	5.1	6.2
2002.....	11.3	92.8	1.7	0.3	0.4	1.3	2.5	3.2	3.9	4.9	6.3
2001.....	12.1	92.6	1.8	0.4	0.5	1.7	2.9	3.4	4.1	6.2	7.8
2000.....	12.0	102.4	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.5	7.5
1999.....	10.3	101.9	1.6	0.3	0.4	1.2	2.1	2.4	3.2	4.2	5.2

Non-Hispanic 4/, male

2007.....	10.5	107.2	1.6	*	0.4	1.2	2.2	2.8	3.0	3.5	4.9
2006.....	10.3	104.3	1.7	0.4	0.4	1.1	2.9	3.3	3.9	4.1	5.7
2005.....	10.8	110.2	1.7	0.3	0.6	1.7	3.5	4.0	4.1	5.2	6.6
2004.....	10.2	106.6	1.2	0.3	0.3	1.2	2.9	3.9	3.6	4.8	6.6
2003.....	11.0	102.9	2.0	0.3	0.6	1.9	3.8	4.7	4.8	6.7	7.7
2002.....	10.9	106.2	1.9	0.4	0.4	1.7	3.4	4.5	5.1	6.5	8.0
2001.....	11.8	106.2	2.1	0.4	0.5	2.2	3.9	4.6	5.3	7.9	9.9
2000.....	11.9	118.0	1.7	0.5	0.5	1.9	3.9	4.5	5.6	7.2	10.0
1999.....	10.0	116.7	1.9	0.3	0.4	1.5	2.9	3.0	4.3	5.3	6.9

Non-Hispanic 4/, female

2007.....	14.0	81.2	1.3	*	0.3	0.7	1.2	1.4	1.7	2.4	2.9
2006.....	13.0	80.5	1.3	0.3	0.3	0.7	1.6	1.7	1.7	2.6	3.3
2005.....	12.7	78.0	1.3	0.3	0.3	0.8	1.5	1.9	2.2	3.4	4.3
2004.....	12.1	74.3	1.2	*	0.4	0.7	1.4	1.7	2.1	2.8	3.5
2003.....	12.6	75.3	1.6	0.5	0.4	1.0	1.8	2.1	2.5	3.6	4.7
2002.....	11.7	78.7	1.4	0.3	0.3	0.8	1.7	2.0	2.6	3.4	4.6
2001.....	12.3	78.3	1.5	0.3	0.4	1.1	1.9	2.2	2.9	4.6	5.7
2000.....	12.1	86.1	1.5	0.3	0.4	1.2	1.9	2.2	2.9	3.8	5.1
1999.....	10.5	86.3	1.4	0.3	0.3	0.8	1.3	1.7	2.2	3.1	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic White, both sexes

2007 5/.....	13.0	79.1	1.3	0.2	0.3	0.9	1.6	1.9	2.2	2.7	3.7
2006 5/.....	12.4	77.1	1.2	0.3	0.3	0.9	2.3	2.6	2.7	3.3	4.2
2005 5/.....	12.3	77.9	1.2	0.2	0.3	1.2	2.5	2.9	3.2	4.2	5.3
2004 5/.....	11.7	71.6	1.0	0.2	0.3	0.9	2.2	2.8	2.8	3.7	4.8
2003 5/.....	12.3	72.6	1.5	0.4	0.4	1.5	2.8	3.2	3.6	5.0	5.9
2002.....	11.6	76.0	1.3	0.2	0.3	1.2	2.5	3.1	3.7	4.7	6.0
2001.....	12.4	73.1	1.5	0.3	0.4	1.7	2.8	3.3	3.9	6.0	7.7
2000.....	12.2	79.7	1.4	0.3	0.4	1.5	2.8	3.1	4.0	5.4	7.2
1999.....	10.2	79.3	1.4	0.3	0.3	1.0	1.8	2.0	3.0	3.8	4.5

Non-Hispanic White, male

2007 5/.....	10.7	92.3	1.6	*	0.3	1.1	2.1	2.6	2.7	3.0	4.7
2006 5/.....	10.5	89.1	1.4	0.4	0.4	1.0	3.0	3.4	3.7	4.1	5.3
2005 5/.....	10.9	94.6	1.4	*	0.5	1.6	3.5	3.9	4.1	5.1	6.5
2004 5/.....	10.3	83.1	1.0	*	*	1.2	2.9	4.0	3.5	4.7	6.3
2003 5/.....	11.0	87.4	1.6	0.3	0.4	2.0	3.9	4.4	4.8	6.4	7.3
2002.....	10.7	88.2	1.4	*	0.4	1.6	3.4	4.4	4.9	6.2	7.6
2001.....	11.8	85.2	1.7	*	0.5	2.1	3.8	4.6	5.1	7.5	9.9
2000.....	11.8	91.8	1.7	0.4	0.4	1.9	3.8	4.2	5.4	7.2	9.8
1999.....	9.6	92.3	1.7	0.3	0.4	1.3	2.6	2.7	3.8	4.9	6.0

Non-Hispanic White, female

2007 5/.....	15.3	65.3	1.1	*	*	0.7	1.2	1.2	1.6	2.5	2.7
2006 5/.....	14.2	64.5	1.1	*	*	0.7	1.5	1.7	1.6	2.6	3.2
2005 5/.....	13.7	60.4	1.0	*	*	0.7	1.5	1.8	2.2	3.3	4.0
2004 5/.....	13.0	59.4	1.0	*	0.3	0.7	1.3	1.5	2.0	2.7	3.3
2003 5/.....	13.5	57.1	1.4	0.5	0.3	1.0	1.7	2.1	2.4	3.5	4.5
2002.....	12.5	63.2	1.3	*	*	0.8	1.5	1.8	2.5	3.2	4.4
2001.....	13.1	60.4	1.3	*	*	1.2	1.9	1.9	2.8	4.5	5.4
2000.....	12.6	66.9	1.2	*	0.4	1.0	1.7	2.0	2.6	3.5	4.7
1999.....	10.9	65.5	1.0	0.4	*	0.7	1.0	1.4	2.1	2.6	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic Black, both sexes

2007 5/.....	12.0	161.6	1.9	*	*	1.2	2.4	3.6	4.1	4.9	6.1
2006 5/.....	11.4	161.4	2.8	*	*	1.2	2.6	3.0	4.0	4.4	7.0
2005 5/.....	12.4	170.4	2.5	0.8	0.9	1.8	3.2	4.2	4.2	6.0	7.9
2004 5/.....	11.9	174.5	2.0	*	0.7	1.3	2.4	4.0	4.3	5.5	7.7
2003 5/.....	12.7	161.6	3.3	*	1.1	1.5	3.5	5.1	5.2	7.3	9.9
2002.....	12.6	165.4	2.7	0.8	0.8	1.9	3.3	4.8	5.9	7.8	9.8
2001.....	12.9	176.9	3.0	0.6	0.7	1.7	3.8	4.8	5.5	8.5	10.1
2000.....	13.9	205.6	2.3	0.6	0.8	2.2	4.3	5.5	6.4	7.3	11.3
1999.....	13.0	202.0	2.9	*	*	1.8	3.8	4.4	5.5	7.9	10.5

Non-Hispanic Black, male

2007 5/.....	12.2	173.4	1.9	*	*	1.8	3.0	4.6	5.5	6.5	7.7
2006 5/.....	12.0	172.0	3.1	*	*	1.5	3.2	3.7	5.8	5.3	9.3
2005 5/.....	13.3	187.0	2.7	*	*	2.4	4.6	5.6	5.1	7.1	9.1
2004 5/.....	12.8	208.3	2.2	*	*	1.6	3.1	4.9	5.6	6.6	9.6
2003 5/.....	13.8	171.9	3.6	*	1.3	1.8	4.4	7.6	6.4	9.7	12.6
2002.....	14.4	186.3	3.6	*	*	2.5	3.7	6.0	8.2	10.6	12.7
2001.....	14.2	195.9	3.7	*	*	2.4	5.3	5.6	6.6	10.9	12.0
2000.....	15.7	236.8	2.2	*	*	2.1	5.7	7.3	8.2	8.5	13.3
1999.....	14.9	228.0	2.9	*	*	2.5	4.9	5.5	8.4	9.5	13.8

Non-Hispanic Black, female

2007 5/.....	11.8	149.2	1.9	*	*	*	1.7	2.7	2.9	3.5	4.7
2006 5/.....	10.8	150.2	2.4	*	*	*	1.9	2.2	2.4	3.6	5.0
2005 5/.....	11.5	153.1	2.3	*	*	1.3	1.8	2.9	3.3	5.0	6.8
2004 5/.....	11.0	139.4	1.7	*	*	*	1.6	3.0	3.1	4.4	6.1
2003 5/.....	11.6	150.9	3.0	*	*	*	2.7	2.9	4.0	5.1	7.5
2002.....	10.9	143.6	1.9	*	*	*	2.9	3.7	3.8	5.3	7.2
2001.....	11.7	156.8	2.2	*	*	*	2.4	4.1	4.5	6.4	8.4
2000.....	12.4	173.4	2.4	*	*	2.2	3.0	3.8	4.8	6.3	9.4
1999.....	11.2	175.3	2.8	*	*	*	2.7	3.5	2.9	6.5	7.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/, both sexes

2007.....	79.0	31.4	3.6	1.9	2.4	3.6	5.6	6.9	8.7	12.6	20.0
2006.....	79.3	28.4	3.4	1.8	2.2	3.9	6.1	7.4	9.2	13.4	23.0
2005.....	73.4	29.2	3.8	1.9	2.4	4.0	5.8	7.0	8.7	13.9	22.8
2004.....	68.8	29.2	3.8	1.9	2.4	3.8	5.6	7.1	9.0	14.3	21.9
2003.....	69.3	29.7	4.0	2.1	2.4	4.1	5.8	7.4	9.3	14.5	23.1
2002.....	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1	23.5
2001.....	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1	21.7
2000.....	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9	22.1
1999.....	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0	21.6

All origins 3/, male

2007.....	64.9	35.8	3.9	2.2	2.8	4.2	6.3	7.8	9.6	14.2	23.0
2006.....	65.2	32.5	3.6	1.9	2.3	4.6	7.0	8.6	10.5	15.4	27.2
2005.....	60.9	34.8	4.3	2.0	2.5	5.0	6.5	7.9	9.9	16.1	27.6
2004.....	57.2	33.8	4.0	2.0	2.6	4.4	6.4	8.2	10.1	16.6	26.0
2003.....	57.3	34.8	4.2	2.3	2.8	4.6	6.6	8.2	10.4	16.5	27.8
2002.....	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4	28.2
2001.....	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8	25.7
2000.....	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5	26.3
1999.....	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8	26.6

All origins 3/, female

2007.....	92.7	26.9	3.4	1.6	2.0	3.0	4.8	6.0	7.7	10.9	17.1
2006.....	93.0	24.1	3.1	1.7	2.0	3.1	5.2	6.3	8.0	11.5	19.0
2005.....	85.6	23.2	3.3	1.7	2.2	2.9	5.0	6.1	7.5	11.7	17.9
2004.....	79.9	24.5	3.6	1.8	2.2	3.1	4.8	6.0	7.8	12.0	17.9
2003.....	81.0	24.5	3.8	1.8	2.1	3.4	4.9	6.5	8.1	12.5	18.5
2002.....	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9	18.9
2001.....	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5	17.7
2000.....	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3	17.9
1999.....	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3	16.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Hispanic, both sexes

2007.....	25.7	28.8	3.1	1.7	2.3	2.9	4.9	4.9	5.2	8.9	13.4
2006.....	26.4	24.6	3.3	1.3	2.3	3.7	5.2	5.8	6.5	10.0	18.2
2005.....	24.0	29.1	4.1	1.4	1.8	4.1	4.0	5.5	6.8	9.6	17.5
2004.....	22.3	26.1	3.3	1.5	2.3	3.2	4.1	4.4	6.5	9.8	16.8
2003.....	23.1	27.1	4.3	1.8	2.5	4.2	4.0	5.2	7.2	11.1	17.4
2002.....	21.6	25.1	3.8	2.0	2.2	3.1	4.2	4.6	6.6	9.7	17.0
2001.....	21.2	22.9	3.7	1.7	1.8	3.2	4.1	4.6	5.8	9.7	15.4
2000.....	20.7	25.4	3.7	1.6	1.8	2.6	3.9	5.0	6.8	11.0	17.1
1999.....	21.5	26.5	3.7	1.4	1.9	3.1	3.8	5.1	8.0	11.2	17.4

Hispanic, male

2007.....	23.5	33.3	3.1	1.9	2.5	3.3	4.9	5.3	6.0	10.4	16.5
2006.....	24.9	28.0	3.3	1.5	2.3	4.8	5.8	6.8	8.0	11.7	24.4
2005.....	23.2	34.6	4.2	1.5	1.7	4.7	5.0	6.1	8.4	11.2	22.8
2004.....	21.3	28.7	3.5	1.6	2.6	3.9	4.9	5.0	7.6	13.4	21.5
2003.....	22.2	32.6	4.5	1.9	2.7	4.5	4.3	6.1	8.7	12.6	21.6
2002.....	21.1	27.2	4.1	2.5	2.3	3.2	4.7	5.3	7.9	12.1	23.1
2001.....	20.2	25.4	4.3	1.6	2.0	3.9	4.2	5.0	7.1	12.6	20.0
2000.....	20.2	25.3	3.6	1.6	2.1	3.2	4.6	5.5	8.8	14.3	23.1
1999.....	21.0	32.1	3.4	1.6	1.7	3.6	3.9	5.9	10.6	15.3	23.2

Hispanic, female

2007.....	28.1	24.2	3.2	1.6	2.1	2.4	4.9	4.5	4.2	7.3	10.1
2006.....	28.1	21.1	3.4	1.2	2.3	2.5	4.5	4.5	4.7	8.0	11.2
2005.....	24.9	23.3	3.9	1.2	2.0	3.5	2.9	4.7	5.0	7.8	11.6
2004.....	23.4	23.5	3.1	1.5	2.1	2.4	3.1	3.7	5.2	5.7	11.8
2003.....	24.1	21.5	4.0	1.7	2.2	3.9	3.6	4.0	5.3	9.4	12.8
2002.....	22.1	22.8	3.5	1.5	2.0	3.0	3.5	3.8	5.1	7.1	10.4
2001.....	22.3	20.2	3.1	1.8	1.5	2.5	3.9	4.0	4.3	6.5	10.6
2000.....	21.4	25.5	3.8	1.6	1.4	1.9	3.0	4.4	4.6	7.4	10.9
1999.....	22.0	20.6	4.0	1.3	2.1	2.5	3.7	4.2	5.0	6.7	11.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 4/, both sexes

2007.....	88.3	32.1	3.8	2.0	2.5	3.8	5.7	7.4	9.5	13.3	21.1
2006.....	88.3	29.5	3.4	1.9	2.1	3.9	6.3	7.8	9.9	14.1	23.8
2005.....	81.6	28.9	3.7	2.0	2.5	4.0	6.2	7.4	9.1	14.7	23.5
2004.....	76.2	30.0	3.9	2.0	2.4	3.9	5.9	7.8	9.5	15.1	22.6
2003.....	76.5	30.1	4.0	2.1	2.4	4.0	6.1	7.9	9.7	15.0	23.8
2002.....	74.4	29.0	4.3	1.9	2.3	3.6	5.9	7.1	9.4	14.8	24.3
2001.....	71.8	29.0	4.3	2.0	2.3	3.9	5.4	6.9	9.5	14.8	22.3
2000.....	69.3	29.6	4.0	1.9	2.6	3.8	5.8	6.8	9.3	14.3	22.5
1999.....	66.8	30.8	4.3	2.3	2.5	4.0	5.6	6.5	9.0	14.3	21.9

Non-Hispanic 4/, male

2007.....	72.5	36.4	4.1	2.3	2.9	4.4	6.7	8.4	10.6	15.1	24.1
2006.....	72.4	33.8	3.7	2.0	2.3	4.6	7.2	9.0	11.1	16.1	27.5
2005.....	67.4	34.5	4.2	2.2	2.7	5.0	6.9	8.4	10.3	17.0	28.2
2004.....	63.3	35.0	4.1	2.2	2.6	4.5	6.7	9.0	10.6	17.1	26.6
2003.....	63.0	35.0	4.2	2.4	2.8	4.6	7.1	8.7	10.7	17.1	28.5
2002.....	61.2	34.9	4.6	2.1	2.5	4.2	6.6	7.9	11.2	17.1	28.7
2001.....	58.8	32.7	4.6	1.8	2.6	4.5	6.4	7.8	10.4	17.4	26.2
2000.....	56.9	33.6	4.4	1.9	2.7	4.3	6.9	7.9	10.7	16.8	26.5
1999.....	55.2	35.1	4.7	2.3	2.6	4.9	6.3	7.6	10.3	16.8	26.7

Non-Hispanic 4/, female

2007.....	103.4	27.6	3.5	1.7	2.0	3.1	4.8	6.3	8.5	11.5	18.1
2006.....	103.5	25.0	3.0	1.8	1.9	3.3	5.3	6.7	8.7	12.1	20.1
2005.....	95.1	23.0	3.1	1.8	2.2	2.8	5.5	6.4	8.0	12.4	18.8
2004.....	88.5	24.7	3.7	1.9	2.2	3.2	5.1	6.5	8.4	13.0	18.7
2003.....	89.4	25.1	3.7	1.8	2.0	3.3	5.1	7.0	8.6	12.9	19.2
2002.....	86.8	22.7	4.0	1.8	2.1	3.0	5.2	6.2	7.7	12.5	19.9
2001.....	84.2	25.2	4.0	2.1	2.1	3.2	4.5	6.0	8.6	12.1	18.5
2000.....	81.0	25.4	3.5	1.8	2.4	3.3	4.6	5.6	7.9	11.8	18.6
1999.....	77.9	26.2	3.9	2.2	2.4	3.2	4.8	5.6	7.7	11.8	17.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic White, both sexes

2007 5/.....	97.1	23.8	3.1	1.8	2.3	3.6	5.5	6.8	8.5	12.0	19.3
2006 5/.....	96.9	23.3	3.0	1.8	2.0	3.6	6.1	7.4	9.0	12.7	21.9
2005 5/.....	88.9	23.2	3.3	1.8	2.1	3.8	5.9	6.8	8.2	13.2	21.9
2004 5/.....	82.7	24.8	3.5	1.9	2.3	3.6	5.5	6.9	8.1	13.5	20.9
2003 5/.....	83.0	24.6	3.5	2.0	2.2	3.8	5.8	7.0	8.4	13.2	21.8
2002.....	80.5	24.7	3.8	1.8	2.2	3.5	5.2	6.3	8.3	13.2	22.1
2001.....	77.3	23.2	3.9	2.0	2.2	3.6	5.0	6.0	8.2	13.0	19.6
2000.....	74.3	24.0	3.4	1.7	2.4	3.7	5.4	5.8	8.1	12.3	19.9
1999.....	71.5	24.2	4.0	2.1	2.3	3.9	5.0	6.0	8.0	12.1	19.0

Non-Hispanic White, male

2007 5/.....	78.4	26.6	3.5	2.1	2.7	4.4	6.5	8.1	9.8	14.0	22.6
2006 5/.....	78.2	26.8	3.3	1.8	2.2	4.3	7.5	9.0	10.8	15.0	26.0
2005 5/.....	72.3	27.7	3.8	2.0	2.4	5.0	6.9	8.0	9.8	16.0	27.2
2004 5/.....	67.4	29.6	3.7	2.1	2.5	4.3	6.5	8.7	9.2	15.9	25.6
2003 5/.....	66.9	30.4	3.6	2.5	2.5	4.7	7.3	8.2	9.7	15.5	26.9
2002.....	64.8	31.4	4.0	2.0	2.3	4.2	6.2	7.3	10.3	16.0	27.0
2001.....	61.9	25.3	4.4	1.9	2.4	4.4	6.2	7.0	9.3	15.8	23.5
2000.....	59.6	26.7	3.8	1.9	2.7	4.3	6.6	7.3	9.7	14.9	23.9
1999.....	57.6	27.4	4.5	2.1	2.3	5.0	5.9	7.2	9.4	14.8	24.0

Non-Hispanic White, female

2007 5/.....	115.1	20.9	2.8	1.6	1.9	2.8	4.4	5.5	7.2	10.1	15.9
2006 5/.....	114.9	19.5	2.7	1.7	1.7	2.8	4.6	5.8	7.2	10.4	17.9
2005 5/.....	105.0	18.4	2.8	1.6	1.8	2.5	5.0	5.6	6.5	10.3	16.5
2004 5/.....	97.5	19.8	3.3	1.6	2.1	2.9	4.4	5.2	7.0	11.0	16.1
2003 5/.....	98.4	18.6	3.5	1.6	1.9	2.9	4.3	5.8	7.1	10.8	16.7
2002.....	95.5	17.7	3.6	1.7	2.0	2.6	4.2	5.3	6.3	10.3	17.1
2001.....	92.1	21.0	3.4	2.1	2.0	2.8	3.8	5.0	7.2	10.2	15.8
2000.....	88.3	21.2	2.9	1.6	2.1	3.0	4.1	4.3	6.6	9.7	15.9
1999.....	84.9	20.9	3.5	2.1	2.3	2.9	4.1	4.7	6.5	9.4	14.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic Black, both sexes

2007 5/.....	68.5	61.9	6.0	2.8	3.2	4.8	7.7	11.7	16.9	23.5	36.2
2006 5/.....	68.7	53.8	4.6	2.3	2.9	5.5	8.1	11.7	16.7	23.2	38.5
2005 5/.....	65.9	51.0	4.9	2.9	3.9	5.2	8.3	11.6	16.2	25.5	37.7
2004 5/.....	63.1	50.5	5.4	2.7	3.0	5.2	8.8	13.5	18.4	27.0	36.5
2003 5/.....	63.0	52.6	5.5	2.3	3.3	5.0	8.3	13.8	17.5	28.1	39.8
2002.....	61.7	46.1	6.1	2.3	3.0	4.4	9.8	12.7	17.2	26.5	42.8
2001.....	61.1	53.8	5.6	2.0	3.0	5.4	8.0	12.0	17.3	27.4	42.3
2000.....	60.1	51.3	6.1	2.4	3.5	4.9	8.6	13.4	17.8	27.8	42.7
1999.....	58.0	58.4	5.5	3.0	3.2	4.9	9.3	10.9	16.0	29.5	43.4

Non-Hispanic Black, male

2007 5/.....	61.4	71.1	6.1	3.3	4.0	4.8	8.2	12.1	17.0	25.1	38.8
2006 5/.....	61.5	61.0	5.0	2.8	3.2	6.2	7.5	11.6	15.5	24.6	41.5
2005 5/.....	58.9	63.8	5.8	3.1	4.3	6.0	8.2	12.0	16.0	25.8	40.6
2004 5/.....	58.1	56.9	5.8	2.2	3.1	6.0	8.3	13.0	20.6	27.3	38.5
2003 5/.....	57.6	56.8	6.3	1.9	4.1	4.8	7.3	13.4	17.8	29.9	43.4
2002.....	56.6	50.6	6.6	2.3	3.4	4.3	9.8	13.0	18.5	27.2	45.2
2001.....	56.4	61.3	5.2	1.6	3.4	5.6	8.1	12.9	18.2	30.5	48.2
2000.....	56.1	61.4	6.4	2.1	3.2	5.1	9.4	13.6	18.9	30.9	48.8
1999.....	54.4	66.5	5.6	3.3	3.9	5.0	10.1	11.0	16.7	32.5	48.5

Non-Hispanic Black, female

2007 5/.....	75.0	52.3	5.9	2.3	2.4	4.7	7.1	11.3	16.8	22.1	33.9
2006 5/.....	75.2	46.3	4.1	1.8	2.6	4.8	8.7	11.9	17.8	22.0	35.9
2005 5/.....	72.3	37.6	4.0	2.8	3.4	4.4	8.4	11.2	16.3	25.3	35.1
2004 5/.....	67.6	43.8	5.0	3.2	2.9	4.4	9.4	13.8	16.5	26.6	34.8
2003 5/.....	67.9	48.4	4.6	2.7	2.6	5.3	9.4	14.1	17.3	26.5	36.7
2002.....	66.3	41.3	5.6	2.3	2.5	4.5	9.9	12.4	16.0	25.9	40.7
2001.....	65.2	46.0	6.0	2.5	2.7	5.2	8.0	11.2	16.5	24.6	37.2
2000.....	63.7	40.9	5.8	2.7	3.7	4.8	7.8	13.2	16.8	25.0	37.3
1999.....	61.3	50.0	5.4	2.7	2.5	4.7	8.6	10.8	15.3	26.9	38.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes

2007.....	41.0	30.2	9.6	4.9	6.0	30.2	44.7	38.6	35.0	36.5	41.8
2006.....	40.6	27.8	9.9	5.3	5.9	31.2	45.3	38.8	35.1	36.9	43.3
2005.....	39.7	26.4	10.3	5.5	6.4	31.4	43.4	36.0	33.8	35.3	41.6
2004.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8	39.7
2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0	40.4
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9	39.4
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1	36.7
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9	35.4
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2	35.4

All origins 3/, male

2007.....	53.7	32.8	11.7	5.6	7.4	41.3	67.8	58.0	51.9	52.4	58.7
2006.....	53.5	31.6	11.1	6.1	7.3	43.0	69.1	59.2	52.1	52.7	60.8
2005.....	52.3	27.5	12.4	6.3	8.1	43.0	66.2	54.8	49.7	50.6	58.7
2004.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7	56.0
2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2	58.0
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5	56.8
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5	54.1
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9	53.1
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4	53.1

All origins 3/, female

2007.....	28.7	27.4	7.5	4.1	4.6	18.6	20.1	18.4	17.7	20.5	25.0
2006.....	28.1	23.8	8.6	4.4	4.4	18.9	19.9	17.6	17.6	20.8	25.8
2005.....	27.5	25.2	8.0	4.7	4.7	19.2	19.2	16.3	17.5	19.8	24.6
2004.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7	23.6
2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7	23.0
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2	22.4
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7	19.6
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0	18.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0	18.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, both sexes

2007.....	25.8	16.2	9.0	4.7	4.7	24.9	42.7	30.8	25.9	26.6	30.9
2006.....	27.2	19.5	9.7	4.3	6.0	29.2	42.8	30.9	28.4	28.9	34.6
2005.....	26.9	18.5	10.6	4.6	4.9	30.4	43.3	31.1	27.6	28.4	31.9
2004.....	25.2	17.5	9.0	4.6	5.6	28.7	37.6	29.7	24.9	28.7	30.6
2003.....	26.1	15.1	9.9	5.7	5.8	31.0	36.3	31.8	26.8	29.9	33.9
2002.....	26.1	15.2	10.9	5.3	5.4	30.8	39.6	28.8	26.4	28.5	35.6
2001.....	25.8	18.8	10.3	5.0	5.7	27.8	37.0	30.2	26.1	31.6	33.1
2000.....	25.0	15.3	10.4	5.7	5.6	27.9	35.5	27.9	26.6	30.3	33.1
1999.....	25.5	17.7	11.9	4.9	6.4	26.1	35.3	28.7	28.0	31.0	34.6

Hispanic, male

2007.....	37.6	17.4	11.3	5.7	6.0	37.0	66.8	46.3	39.6	39.9	46.5
2006.....	39.7	20.8	11.3	5.4	7.3	43.8	66.3	46.5	43.1	43.8	52.8
2005.....	39.0	18.3	13.3	5.2	5.9	43.4	66.6	47.3	42.3	42.3	47.7
2004.....	36.5	18.8	10.5	5.1	7.2	43.0	57.2	45.1	37.8	43.8	46.0
2003.....	38.3	17.4	11.9	7.0	8.1	44.9	55.4	48.8	41.9	45.4	51.1
2002.....	38.5	19.5	13.5	6.3	6.8	45.8	60.0	44.5	41.3	42.9	55.3
2001.....	37.6	18.0	12.7	6.0	6.4	40.2	56.9	47.0	40.2	48.9	51.7
2000.....	36.9	17.0	11.6	5.7	6.5	40.8	54.8	44.0	41.7	46.3	53.9
1999.....	37.7	20.7	13.5	6.0	7.8	37.3	54.5	44.7	43.4	48.7	54.7

Hispanic, female

2007.....	13.1	14.9	6.5	3.7	3.4	12.2	14.9	11.2	9.6	11.4	13.6
2006.....	13.8	18.1	8.0	3.3	4.6	13.7	14.8	11.2	10.8	11.9	14.5
2005.....	13.8	18.6	7.7	4.0	3.8	16.6	15.3	10.7	10.4	12.3	14.8
2004.....	13.1	16.2	7.5	4.0	4.0	13.4	13.6	10.6	9.8	11.6	13.8
2003.....	13.1	12.7	7.8	4.3	3.5	15.9	12.7	10.9	9.3	12.4	15.4
2002.....	12.8	10.8	8.2	4.3	3.8	14.4	14.5	9.7	9.2	12.4	14.5
2001.....	13.2	19.7	7.8	4.0	5.0	14.0	12.6	10.2	10.0	12.6	13.5
2000.....	12.4	13.5	9.2	5.6	4.7	13.1	11.9	9.0	9.4	12.9	11.3
1999.....	12.6	14.6	10.3	3.7	4.9	13.4	12.2	9.9	10.3	11.6	13.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 4/, both sexes

2007.....	43.6	34.6	9.8	4.9	6.4	31.3	45.0	40.4	37.3	38.5	43.5
2006.....	42.8	30.2	9.8	5.5	5.9	31.6	45.8	40.7	36.6	38.3	44.5
2005.....	41.8	28.5	10.2	5.7	6.8	31.6	43.3	37.1	35.0	36.5	42.8
2004.....	40.1	28.1	10.6	6.0	7.6	33.7	41.8	35.7	31.7	35.8	40.8
2003.....	39.2	25.9	11.1	5.5	7.5	33.3	42.0	32.9	30.8	35.7	41.1
2002.....	38.5	25.5	10.4	6.0	7.6	35.7	41.1	33.0	31.5	35.7	39.7
2001.....	36.9	25.4	11.3	6.6	7.7	33.6	39.8	31.5	28.8	34.3	36.9
2000.....	36.0	25.0	12.2	7.0	8.1	34.3	39.3	30.8	28.8	33.1	35.4
1999.....	36.2	23.4	12.4	7.5	8.3	34.5	37.8	30.1	29.2	32.2	35.3

Non-Hispanic 4/, male

2007.....	56.5	37.5	11.8	5.6	7.8	42.1	68.0	61.0	55.2	55.0	60.8
2006.....	55.8	34.7	10.9	6.3	7.2	42.6	69.5	62.5	54.4	54.5	61.9
2005.....	54.5	30.0	12.1	6.5	8.6	42.9	65.9	56.6	51.3	52.1	60.2
2004.....	51.9	31.8	12.1	7.2	9.4	44.8	62.8	54.0	46.4	50.7	57.3
2003.....	50.9	28.7	13.2	6.2	9.6	44.6	63.1	49.3	45.6	50.9	58.6
2002.....	50.3	28.6	12.7	7.2	9.6	47.8	62.7	50.6	46.8	51.5	56.7
2001.....	48.5	28.0	13.4	7.9	10.0	46.1	61.4	48.1	43.4	49.3	54.0
2000.....	47.3	27.7	14.5	8.3	10.5	46.6	59.6	47.4	43.1	49.0	52.5
1999.....	47.4	26.6	14.5	9.0	10.8	46.1	57.3	46.1	43.6	46.9	52.5

Non-Hispanic 4/, female

2007.....	31.2	31.5	7.7	4.1	4.9	20.0	21.0	19.8	19.5	22.1	26.7
2006.....	30.4	25.5	8.7	4.7	4.4	19.9	20.9	18.9	19.0	22.4	27.4
2005.....	29.6	26.9	8.1	4.8	4.9	19.7	20.0	17.5	18.9	21.1	25.8
2004.....	28.8	24.2	9.0	4.8	5.7	21.9	20.2	17.4	17.1	21.1	24.8
2003.....	28.1	22.9	8.9	4.7	5.2	21.4	20.4	16.5	16.2	20.8	23.9
2002.....	27.3	22.3	8.1	4.8	5.6	23.0	19.2	15.5	16.3	20.2	23.2
2001.....	26.0	22.6	9.2	5.3	5.3	20.5	18.1	15.1	14.4	19.6	20.2
2000.....	25.2	22.2	9.7	5.6	5.6	21.5	18.9	14.3	14.6	17.5	18.6
1999.....	25.5	20.0	10.2	6.0	5.8	22.4	17.9	14.3	14.9	17.6	18.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic White, both sexes

2007 5/.....	47.0	27.6	9.2	4.4	6.1	34.6	49.3	44.6	40.8	41.0	46.2
2006 5/.....	45.7	24.3	8.9	4.8	5.8	34.5	50.5	44.3	39.3	40.5	46.4
2005 5/.....	44.4	24.5	9.8	5.1	6.6	34.3	47.2	40.2	37.0	38.3	44.7
2004 5/.....	42.8	24.1	10.2	5.3	7.4	37.3	45.7	38.0	33.6	37.4	42.7
2003 5/.....	41.6	21.6	10.7	5.0	7.2	36.5	45.7	34.9	32.3	37.0	42.3
2002.....	40.6	19.8	9.4	5.1	7.3	39.0	44.2	34.3	33.2	36.4	40.8
2001.....	38.5	20.9	10.9	5.9	7.3	36.0	42.2	32.9	29.6	34.9	36.8
2000.....	37.3	20.1	11.1	6.3	8.0	37.3	41.4	31.1	29.1	33.4	35.0
1999.....	37.2	19.0	11.1	6.6	8.1	37.2	39.1	30.8	29.1	32.1	34.8

Non-Hispanic White, male

2007 5/.....	60.0	30.7	11.1	5.0	7.5	45.9	74.7	66.8	59.5	57.7	63.4
2006 5/.....	58.6	27.6	9.8	5.4	7.0	45.8	76.9	67.4	57.3	56.5	63.9
2005 5/.....	57.0	25.6	11.8	6.0	8.2	46.1	71.9	60.9	53.4	53.9	62.3
2004 5/.....	54.6	28.3	11.8	6.3	9.3	49.2	68.9	57.0	48.3	52.6	59.4
2003 5/.....	53.1	24.1	12.9	5.6	9.2	48.4	68.2	52.0	47.4	51.9	60.2
2002.....	52.0	23.3	11.1	6.1	9.2	51.6	67.2	51.9	48.9	51.9	57.9
2001.....	49.7	22.9	13.2	6.7	9.5	48.8	65.0	49.7	44.2	49.7	53.7
2000.....	48.1	22.9	13.4	7.5	10.2	50.0	62.6	47.2	43.4	49.3	51.5
1999.....	47.8	22.0	12.9	7.7	10.4	48.8	59.1	46.9	43.0	46.6	51.5

Non-Hispanic White, female

2007 5/.....	34.5	24.4	7.1	3.8	4.6	22.6	22.5	22.1	21.8	24.1	28.9
2006 5/.....	33.3	20.8	8.0	4.2	4.4	22.6	22.7	20.8	21.0	24.3	29.0
2005 5/.....	32.3	23.4	7.6	4.3	4.8	21.8	21.4	19.1	20.5	22.5	27.2
2004 5/.....	31.4	19.7	8.5	4.3	5.4	24.8	21.5	18.7	18.6	22.0	26.1
2003 5/.....	30.5	19.0	8.4	4.4	5.1	24.0	22.4	17.4	17.1	21.9	24.4
2002.....	29.6	16.0	7.5	4.1	5.4	25.7	20.7	16.4	17.2	20.9	23.8
2001.....	27.8	18.7	8.5	5.0	5.1	22.5	18.9	15.8	14.7	20.0	19.9
2000.....	26.9	17.1	8.7	5.0	5.6	24.0	19.6	14.7	14.6	17.5	18.5
1999.....	27.0	15.9	9.3	5.4	5.7	25.1	18.6	14.5	15.0	17.6	18.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic Black, both sexes

2007 5/.....	34.9	64.0	12.6	7.1	7.5	21.4	31.9	31.7	33.1	37.9	40.5
2006 5/.....	36.2	54.8	14.2	8.3	6.5	22.1	31.9	34.9	36.4	39.2	45.4
2005 5/.....	35.9	48.6	12.6	8.0	7.9	23.3	32.5	30.5	35.9	37.0	42.0
2004 5/.....	33.7	47.0	12.8	8.8	8.7	22.2	30.6	33.4	31.7	36.0	39.4
2003 5/.....	33.3	44.7	13.7	7.7	8.4	21.3	31.7	30.6	31.1	36.7	42.7
2002.....	34.0	50.1	14.6	9.6	9.2	24.4	32.7	34.1	30.9	39.1	40.8
2001.....	34.4	44.5	14.2	9.8	9.7	25.3	34.2	31.3	30.9	36.5	45.3
2000.....	34.4	46.5	16.8	10.0	9.1	24.3	35.5	34.2	32.8	36.7	44.6
1999.....	36.1	44.1	18.2	11.8	9.9	25.8	35.8	32.5	34.0	37.4	44.3

Non-Hispanic Black, male

2007 5/.....	50.0	65.2	14.5	8.6	9.0	30.9	47.7	50.3	52.8	58.5	62.1
2006 5/.....	52.5	64.1	16.3	10.0	8.3	33.5	47.4	56.0	59.6	60.8	67.2
2005 5/.....	51.3	52.3	14.1	8.8	10.7	33.3	49.7	48.9	56.3	56.9	61.6
2004 5/.....	47.9	49.2	14.4	10.8	10.5	31.9	45.5	53.5	50.6	53.1	58.0
2003 5/.....	47.4	49.0	15.9	9.1	11.0	30.9	49.3	47.9	48.9	56.7	62.0
2002.....	49.0	52.1	18.9	11.4	12.4	35.1	50.7	55.8	48.8	60.3	59.1
2001.....	49.5	50.4	15.8	13.0	13.2	37.0	52.9	49.5	48.0	54.3	68.4
2000.....	50.3	49.5	19.4	12.2	12.5	35.3	54.5	55.9	51.1	56.9	69.9
1999.....	52.3	48.5	21.8	15.0	13.5	37.9	56.5	52.7	53.4	56.5	70.0

Non-Hispanic Black, female

2007 5/.....	21.2	62.8	10.6	5.6	6.0	11.7	15.8	13.8	15.3	19.4	21.5
2006 5/.....	21.5	45.0	12.0	6.6	4.7	10.4	16.0	15.0	15.5	20.1	26.2
2005 5/.....	21.8	44.8	11.1	7.1	5.0	13.0	15.0	13.2	17.5	19.2	24.8
2004 5/.....	20.8	44.8	11.1	6.8	6.9	12.2	15.6	14.7	14.6	20.8	23.0
2003 5/.....	20.5	40.3	11.4	6.1	5.7	11.5	14.2	14.7	15.1	18.9	25.7
2002.....	20.3	48.0	10.1	7.7	5.9	13.3	15.2	14.4	14.9	20.3	24.6
2001.....	20.6	38.3	12.4	6.4	6.0	13.3	16.3	14.6	15.5	20.7	24.9
2000.....	20.1	43.3	14.2	7.8	5.5	13.0	17.5	14.6	16.5	18.8	22.3
1999.....	21.4	39.6	14.5	8.4	6.3	13.4	16.3	14.3	16.6	20.5	21.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes

2007.....	15.5	3.0	3.5	2.8	4.0	22.6	28.9	20.2	16.3	15.7	15.9
2006.....	16.2	3.4	3.8	3.1	4.1	23.9	29.8	20.9	17.4	16.4	16.8
2005.....	16.3	3.6	4.0	3.3	4.3	24.4	29.3	20.5	17.6	16.3	17.0
2004.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4	16.4
2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7	17.5
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4	17.3
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1	16.8
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9	16.6

All origins 3/, male

2007.....	22.5	2.7	3.9	3.1	4.8	29.4	43.4	30.4	24.6	23.5	23.5
2006.....	23.1	3.3	4.0	3.5	4.7	31.2	44.6	31.8	25.9	23.7	24.2
2005.....	23.3	3.5	4.5	3.7	5.1	31.8	44.0	31.2	26.0	23.5	24.8
2004.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0	24.1
2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1	25.5
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7	24.5
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7	25.4
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0	24.0

All origins 3/, female

2007.....	8.8	3.3	3.1	2.4	3.3	15.5	13.4	9.6	7.8	7.8	8.4
2006.....	9.4	3.6	3.6	2.8	3.3	16.2	14.0	9.5	8.6	9.1	9.4
2005.....	9.5	3.6	3.5	3.0	3.5	16.5	13.6	9.3	8.9	8.9	9.2
2004.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8	8.9
2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3	9.5
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6	9.9
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2	9.4
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5	9.0
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8	9.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

Hispanic, both sexes											
2007.....	13.4	3.0	4.3	3.2	3.4	18.5	31.4	19.6	14.4	13.4	13.5
2006.....	14.8	4.6	4.7	2.8	4.5	23.4	32.1	20.0	15.5	14.9	15.9
2005.....	14.9	4.7	5.3	3.4	3.9	24.3	32.3	20.5	15.9	14.8	14.9
2004.....	14.5	4.4	4.7	3.5	4.4	23.8	28.5	20.4	15.7	15.1	14.6
2003.....	15.2	4.5	4.2	3.8	4.4	25.2	27.1	21.6	16.5	15.9	17.1
2002.....	15.3	3.1	5.8	3.7	3.9	24.6	30.2	20.2	16.2	15.2	17.0
2001.....	15.6	5.3	5.3	3.3	4.3	22.7	29.0	21.6	15.9	17.3	17.3
2000.....	14.8	5.4	5.2	4.0	4.3	21.9	27.0	19.4	15.0	15.6	16.9
1999.....	14.4	5.3	5.5	3.1	4.7	20.0	25.9	19.6	15.9	14.4	16.6

Hispanic, male											
2007.....	19.7	*	4.8	3.8	4.3	26.6	47.7	28.8	21.6	19.3	19.0
2006.....	21.7	4.6	5.3	3.4	5.2	34.2	49.1	29.7	23.0	21.6	23.3
2005.....	21.7	4.4	6.5	3.9	4.6	33.4	48.7	30.8	23.7	21.0	21.6
2004.....	21.3	*	5.4	3.9	5.4	35.1	42.8	30.4	23.3	22.8	21.5
2003.....	22.2	5.7	4.6	4.8	6.1	35.1	40.7	32.7	25.5	23.2	24.7
2002.....	22.4	*	7.1	4.2	5.1	35.4	44.7	30.3	24.5	22.4	25.6
2001.....	22.8	*	6.0	3.8	4.9	31.7	44.1	33.2	24.1	25.8	26.0
2000.....	21.6	6.1	5.6	3.8	5.2	31.3	40.4	30.2	22.1	22.9	25.6
1999.....	20.8	5.7	5.8	3.6	5.6	27.7	38.5	29.7	23.5	21.6	25.2

Hispanic, female											
2007.....	6.7	*	3.7	2.6	2.4	9.9	12.5	8.1	5.7	6.7	7.5
2006.....	7.4	4.6	4.0	2.2	3.7	12.0	11.8	7.7	6.5	7.2	7.6
2005.....	7.6	5.0	4.1	2.9	3.2	14.6	12.6	7.6	6.8	7.8	7.6
2004.....	7.3	5.0	4.1	3.1	3.4	11.7	11.0	8.1	6.9	6.3	7.1
2003.....	7.6	*	3.8	2.8	2.6	14.4	10.3	8.0	6.1	7.7	8.9
2002.....	7.7	*	4.5	3.2	2.7	12.8	12.4	8.0	6.6	7.2	7.8
2001.....	8.0	6.7	4.6	2.7	3.7	12.7	10.5	7.8	6.4	7.9	8.2
2000.....	7.6	*	4.8	4.1	3.3	11.3	10.7	6.8	6.8	7.8	7.7
1999.....	7.6	*	5.2	2.5	3.8	11.3	10.7	7.9	7.2	6.5	7.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	15.9	3.0	3.3	2.7	4.2	23.5	28.3	20.3	16.8	16.1	16.3
2006.....	16.4	3.1	3.5	3.2	3.9	23.9	29.3	21.1	17.8	16.7	16.8
2005.....	16.5	3.2	3.6	3.3	4.4	24.3	28.6	20.4	17.9	16.5	17.2
2004.....	16.6	3.3	3.9	3.4	5.2	26.4	28.5	21.0	16.9	16.7	16.6
2003.....	16.7	3.5	4.0	3.4	5.2	26.9	29.0	19.7	16.5	16.8	17.5
2002.....	16.9	3.2	3.7	3.5	5.1	29.0	29.7	20.5	17.3	17.3	17.1
2001.....	16.7	3.4	4.2	3.9	5.2	27.2	29.2	20.7	17.1	17.4	17.2
2000.....	16.8	4.3	4.3	4.1	5.4	27.9	29.7	20.5	17.0	17.3	16.7
1999.....	16.9	4.9	4.4	4.6	5.4	28.3	28.5	20.0	17.2	17.2	16.5

Non-Hispanic 4/, male

2007.....	22.9	2.6	3.6	2.9	4.9	30.0	42.3	30.7	25.3	24.4	24.3
2006.....	23.3	2.9	3.5	3.5	4.6	30.5	43.5	32.3	26.6	24.0	24.2
2005.....	23.6	3.3	3.9	3.6	5.2	31.4	42.8	31.1	26.5	24.0	25.2
2004.....	23.3	3.9	4.1	3.9	6.1	33.3	41.7	31.3	24.6	24.2	24.4
2003.....	23.4	3.5	4.5	3.7	6.3	34.5	42.6	28.9	24.6	24.1	25.6
2002.....	23.8	3.4	4.2	3.9	6.0	37.2	44.1	31.1	25.5	24.9	24.3
2001.....	23.5	3.2	4.7	4.4	6.5	36.0	43.7	30.9	25.6	24.4	25.2
2000.....	23.4	4.5	4.8	4.5	6.6	36.2	43.9	30.8	25.1	24.9	24.5
1999.....	23.3	5.1	4.9	5.3	6.6	36.0	42.3	29.7	24.8	24.3	23.7

Non-Hispanic 4/, female

2007.....	9.1	3.4	3.0	2.4	3.5	16.6	13.6	9.9	8.3	8.0	8.5
2006.....	9.8	3.2	3.5	2.9	3.3	17.0	14.4	9.9	9.1	9.4	9.6
2005.....	9.8	3.2	3.3	3.0	3.5	16.9	13.8	9.6	9.4	9.1	9.5
2004.....	10.3	2.6	3.8	3.0	4.3	19.1	14.8	10.6	9.3	9.2	9.1
2003.....	10.3	3.5	3.6	3.1	4.0	18.9	15.1	10.4	8.4	9.5	9.6
2002.....	10.4	2.9	3.1	3.1	4.1	20.4	15.0	10.0	9.1	9.9	10.1
2001.....	10.2	3.6	3.6	3.4	3.8	18.0	14.4	10.5	8.6	10.5	9.5
2000.....	10.5	4.2	3.8	3.7	4.2	19.3	15.3	10.4	9.0	9.8	9.1
1999.....	10.7	4.7	3.9	3.9	4.2	20.3	14.5	10.3	9.6	10.2	9.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

Non-Hispanic White, both sexes

2007 5/.....	16.4	2.7	2.9	2.6	4.3	25.9	29.7	21.1	16.9	16.2	16.8
2006 5/.....	16.9	2.6	3.0	3.0	4.0	26.3	31.3	21.4	18.1	17.1	17.3
2005 5/.....	17.1	2.5	3.3	3.2	4.4	26.6	30.3	21.1	18.2	16.8	17.7
2004 5/.....	17.2	2.9	3.7	3.3	5.4	29.4	30.3	21.3	17.0	16.9	17.0
2003 5/.....	17.2	3.2	3.8	3.3	5.4	29.7	30.7	20.1	16.6	17.0	17.7
2002.....	17.5	2.5	3.3	3.3	5.2	32.1	31.4	20.5	17.7	17.3	17.4
2001.....	17.0	3.1	3.8	3.6	5.2	29.6	30.3	21.3	17.1	17.5	17.3
2000.....	17.1	3.5	3.8	4.0	5.6	30.9	30.9	20.2	17.0	17.4	16.6
1999.....	17.1	4.3	3.7	4.4	5.6	31.1	29.5	20.2	17.2	16.9	16.5

Non-Hispanic White, male

2007 5/.....	23.6	2.7	3.3	2.8	4.9	32.5	44.6	31.6	25.0	24.2	24.6
2006 5/.....	23.8	2.6	2.9	3.1	4.6	33.0	46.5	32.6	26.4	24.2	24.8
2005 5/.....	24.2	2.8	3.5	3.6	5.2	33.9	45.4	31.8	26.3	23.9	25.7
2004 5/.....	23.9	3.3	3.8	3.8	6.4	36.6	44.7	31.4	24.1	24.4	24.7
2003 5/.....	23.9	3.3	4.4	3.5	6.5	37.6	44.7	29.2	24.6	24.0	25.7
2002.....	24.3	2.9	3.6	3.7	6.1	40.8	46.6	30.7	25.8	24.6	24.5
2001.....	23.8	2.9	4.4	4.0	6.4	38.6	45.5	31.5	25.4	24.2	25.0
2000.....	23.7	3.8	4.3	4.5	6.8	39.6	45.6	29.7	24.8	24.8	24.1
1999.....	23.5	4.9	4.0	4.9	6.8	39.0	43.7	29.8	24.3	23.7	23.5

Non-Hispanic White, female

2007 5/.....	9.5	2.7	2.5	2.3	3.6	18.9	14.0	10.5	8.7	8.2	8.9
2006 5/.....	10.2	2.6	3.2	2.8	3.4	19.2	15.3	10.0	9.7	10.0	9.9
2005 5/.....	10.3	2.3	3.2	2.9	3.6	18.8	14.5	10.1	10.0	9.6	9.7
2004 5/.....	10.7	2.4	3.6	2.8	4.3	21.7	15.4	11.0	9.8	9.4	9.2
2003 5/.....	10.7	3.0	3.2	3.1	4.2	21.3	16.4	10.7	8.5	9.9	9.6
2002.....	10.9	2.1	2.9	2.8	4.2	22.9	15.8	10.2	9.5	10.0	10.3
2001.....	10.6	3.3	3.2	3.2	3.8	20.0	14.8	10.9	8.7	10.8	9.6
2000.....	10.9	3.2	3.4	3.5	4.4	21.8	15.9	10.5	9.0	10.0	9.1
1999.....	11.0	3.6	3.4	3.7	4.4	22.8	15.0	10.5	9.9	10.2	9.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

Non-Hispanic Black, both sexes

2007 5/.....	15.1	4.2	4.6	3.4	4.1	16.1	24.1	20.6	19.7	19.5	17.1
2006 5/.....	15.6	4.9	5.2	4.1	3.9	16.0	23.5	22.8	21.9	18.4	17.5
2005 5/.....	15.4	5.7	4.7	3.8	4.4	17.6	24.1	19.9	20.8	18.3	17.2
2004 5/.....	15.6	5.0	5.2	4.0	4.9	17.0	23.1	23.1	20.2	18.0	16.8
2003 5/.....	15.6	4.4	5.1	3.8	4.7	16.7	24.0	21.0	19.1	18.5	19.2
2002.....	15.7	5.6	5.0	4.7	4.7	18.3	25.1	23.9	18.4	19.7	17.3
2001.....	16.3	*	5.4	5.5	5.4	19.3	26.7	21.6	19.6	18.5	19.8
2000.....	16.4	6.9	5.9	4.6	5.0	17.8	27.7	24.4	20.0	18.4	19.7
1999.....	16.7	7.7	7.1	6.0	5.2	19.2	26.6	22.0	19.2	20.5	18.4

Non-Hispanic Black, male

2007 5/.....	23.4	*	4.8	4.0	4.8	22.2	36.5	33.4	32.6	32.2	28.3
2006 5/.....	23.6	*	5.8	4.9	4.8	22.8	34.6	36.3	36.5	30.0	26.8
2005 5/.....	23.4	*	5.1	3.9	5.7	24.0	36.8	32.3	34.8	30.0	27.0
2004 5/.....	23.4	6.2	5.4	4.3	5.4	23.5	33.9	37.2	33.0	28.5	26.2
2003 5/.....	23.3	*	5.1	4.7	5.8	23.4	37.6	32.9	30.7	29.4	28.9
2002.....	23.6	*	6.5	5.0	5.8	24.7	37.8	39.0	29.5	30.5	25.6
2001.....	24.5	*	5.8	6.9	7.3	27.2	40.5	34.1	31.1	27.9	31.5
2000.....	24.8	7.2	6.4	5.0	6.7	24.6	41.7	39.5	31.7	29.5	31.6
1999.....	24.9	*	8.0	7.3	6.6	26.7	40.3	35.5	30.5	30.7	28.8

Non-Hispanic Black, female

2007 5/.....	7.5	*	4.5	2.8	3.4	9.7	11.5	8.3	8.1	8.2	7.1
2006 5/.....	8.2	*	4.6	3.3	3.0	9.0	12.1	10.1	8.6	8.0	9.4
2005 5/.....	8.1	6.5	4.3	3.7	3.2	11.0	11.2	8.2	8.2	7.9	8.7
2004 5/.....	8.5	*	5.0	3.8	4.3	10.3	12.3	10.0	8.6	8.8	8.6
2003 5/.....	8.5	*	5.2	2.9	3.5	9.7	10.6	10.0	8.7	8.8	10.7
2002.....	8.5	*	3.6	4.3	3.6	11.6	12.7	10.2	8.5	10.2	10.1
2001.....	8.9	*	5.0	4.1	3.4	11.1	13.5	10.1	9.2	10.2	9.5
2000.....	8.8	*	5.4	4.2	3.2	10.8	14.4	10.7	9.6	8.6	9.2
1999.....	9.4	9.3	6.2	4.6	3.7	11.7	13.7	9.8	9.0	11.5	9.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes

2007.....	14.6	2.9	3.3	2.6	3.8	22.0	27.8	19.4	15.4	14.7	14.8
2006.....	15.1	3.4	3.6	2.9	3.7	23.2	28.8	19.9	16.4	15.2	15.4
2005.....	15.3	3.6	3.8	3.2	4.0	23.6	28.2	19.6	16.5	15.1	15.6
2004.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2	15.1
2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4	16.0
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8	15.7
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9	15.5
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6	15.1
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1	14.8

All origins 3/, male

2007.....	20.9	2.6	3.7	2.9	4.5	28.5	41.7	29.0	23.1	21.8	21.7
2006.....	21.4	3.3	3.8	3.2	4.2	30.2	42.9	30.1	24.4	21.7	21.9
2005.....	21.7	3.5	4.2	3.5	4.7	30.7	42.3	29.6	24.2	21.8	22.6
2004.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9	21.9
2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9	23.1
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4	22.1
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3	22.5
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1	21.9
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2	21.0

All origins 3/, female

2007.....	8.4	3.2	3.0	2.3	3.0	15.2	12.9	9.3	7.4	7.5	7.9
2006.....	9.0	3.5	3.4	2.6	3.2	15.7	13.7	9.2	8.1	8.7	8.9
2005.....	9.1	3.6	3.4	2.9	3.2	16.2	13.2	9.1	8.6	8.4	8.7
2004.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3	8.3
2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8	9.0
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2	9.4
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6	8.7
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1	8.4
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0	8.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Hispanic, both sexes

2007.....	12.8	2.9	4.1	3.2	3.2	18.2	30.5	19.0	13.8	12.7	12.7
2006.....	14.2	4.6	4.6	2.7	4.2	22.9	31.4	19.3	14.9	14.0	14.6
2005.....	14.3	4.7	5.2	3.3	3.7	23.8	31.4	19.9	15.4	14.0	14.3
2004.....	13.9	4.4	4.6	3.5	4.2	23.2	27.8	19.6	14.7	14.2	13.7
2003.....	14.4	4.5	4.1	3.7	4.2	24.6	26.2	20.3	15.6	15.0	15.9
2002.....	14.6	3.1	5.6	3.5	3.7	23.9	29.0	19.6	15.6	14.5	16.1
2001.....	14.3	4.9	4.7	3.0	4.0	22.1	27.7	20.2	14.5	15.4	15.2
2000.....	13.9	5.2	4.9	3.7	4.0	21.2	26.0	18.5	14.0	14.4	15.2
1999.....	13.3	5.3	5.2	2.9	4.4	19.1	24.5	18.2	14.5	13.0	15.0

Hispanic, male

2007.....	18.7	*	4.6	3.7	4.2	26.0	46.4	27.9	20.5	17.9	17.7
2006.....	20.7	4.6	5.3	3.3	4.8	33.5	48.0	28.7	22.1	20.2	21.1
2005.....	20.7	4.4	6.3	3.7	4.3	32.5	47.2	29.6	22.8	19.7	20.7
2004.....	20.2	*	5.3	3.8	5.0	34.3	41.7	29.0	21.6	21.4	20.0
2003.....	21.1	5.7	4.3	4.6	5.7	34.3	39.5	30.9	24.1	21.6	22.6
2002.....	21.3	*	6.8	3.9	4.9	34.3	42.7	29.3	23.4	21.3	24.2
2001.....	21.0	*	5.3	3.6	4.5	30.8	42.1	31.1	22.2	22.9	23.4
2000.....	20.1	6.1	5.2	3.5	4.8	30.2	38.7	28.6	20.7	20.6	22.9
1999.....	19.1	5.7	5.6	3.3	5.0	26.4	36.4	27.3	21.4	19.4	22.7

Hispanic, female

2007.....	6.5	*	3.6	2.6	2.2	9.8	12.2	7.9	5.6	6.6	7.1
2006.....	7.2	4.6	3.9	2.1	3.6	11.7	11.5	7.3	6.4	7.0	7.5
2005.....	7.4	5.0	4.1	2.9	3.0	14.5	12.4	7.6	6.7	7.5	7.3
2004.....	7.1	5.0	3.9	3.1	3.3	11.3	10.8	7.9	6.6	5.9	6.8
2003.....	7.3	*	3.8	2.7	2.6	14.1	9.8	7.4	5.9	7.4	8.7
2002.....	7.4	*	4.4	3.0	2.4	12.5	12.2	7.8	6.6	7.0	7.5
2001.....	7.2	6.2	4.1	2.5	3.5	12.5	10.1	7.1	5.8	7.1	6.6
2000.....	7.2	*	4.6	4.0	3.1	10.9	10.4	6.7	6.4	7.6	7.1
1999.....	7.2	*	4.7	2.4	3.7	10.9	10.3	7.5	6.6	6.0	7.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic 4/, both sexes

2007.....	14.8	2.9	3.1	2.4	3.9	22.8	27.2	19.4	15.7	15.1	15.1
2006.....	15.3	3.0	3.3	3.0	3.6	23.1	28.1	20.0	16.7	15.4	15.4
2005.....	15.4	3.2	3.4	3.2	4.0	23.5	27.5	19.4	16.7	15.3	15.8
2004.....	15.5	3.2	3.8	3.2	4.9	25.5	27.3	19.6	15.7	15.3	15.2
2003.....	15.5	3.4	3.9	3.2	4.8	25.9	27.7	18.6	15.3	15.4	16.0
2002.....	15.9	3.0	3.5	3.3	4.6	28.3	28.7	19.6	16.1	15.9	15.6
2001.....	15.5	3.2	3.9	3.7	4.7	26.3	28.0	19.4	15.7	16.0	15.5
2000.....	15.6	4.2	4.1	3.8	5.0	26.8	28.3	19.2	15.6	15.7	15.0
1999.....	15.4	4.7	4.0	4.1	4.9	27.0	26.8	18.3	15.5	15.3	14.7

Non-Hispanic 4/, male

2007.....	21.3	2.5	3.4	2.7	4.6	29.0	40.6	29.2	23.7	22.6	22.4
2006.....	21.5	2.9	3.3	3.2	4.0	29.4	41.6	30.4	25.0	21.9	21.9
2005.....	21.8	3.3	3.6	3.5	4.8	30.3	41.1	29.5	24.5	22.1	22.9
2004.....	21.5	3.8	3.9	3.6	5.7	32.1	40.0	29.2	22.6	22.0	22.1
2003.....	21.4	3.4	4.3	3.4	5.9	33.0	40.5	27.3	22.7	21.9	23.1
2002.....	21.9	3.2	4.0	3.7	5.4	36.2	42.4	29.6	23.5	22.5	21.8
2001.....	21.5	3.1	4.3	4.1	5.9	34.5	41.9	29.0	23.2	22.1	22.3
2000.....	21.4	4.2	4.5	4.2	6.0	34.5	41.6	28.5	22.8	22.3	21.6
1999.....	21.0	4.8	4.4	4.6	5.8	34.2	39.6	27.1	22.3	21.4	20.7

Non-Hispanic 4/, female

2007.....	8.7	3.3	2.8	2.2	3.2	16.3	13.1	9.6	7.8	7.6	8.0
2006.....	9.3	3.1	3.3	2.8	3.1	16.5	14.0	9.6	8.5	9.0	9.0
2005.....	9.4	3.2	3.2	2.9	3.2	16.5	13.4	9.4	8.9	8.5	8.9
2004.....	9.8	2.6	3.7	2.8	4.0	18.7	14.3	10.0	8.9	8.7	8.5
2003.....	9.8	3.3	3.4	2.9	3.7	18.4	14.6	9.9	8.0	9.0	9.0
2002.....	10.1	2.9	3.0	2.8	3.8	20.0	14.7	9.7	8.7	9.5	9.6
2001.....	9.8	3.4	3.5	3.2	3.5	17.6	14.0	9.9	8.3	9.9	8.9
2000.....	10.0	4.2	3.6	3.4	3.9	18.7	14.9	10.0	8.5	9.3	8.5
1999.....	10.0	4.7	3.6	3.5	3.8	19.5	13.9	9.7	8.8	9.4	8.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic White, both sexes

2007 5/.....	15.4	2.7	2.7	2.4	4.0	25.1	28.6	20.2	15.8	15.1	15.6
2006 5/.....	15.7	2.5	2.8	2.7	3.6	25.4	30.0	20.3	16.9	15.8	15.8
2005 5/.....	16.0	2.5	3.1	3.1	4.0	25.7	29.2	20.1	16.9	15.5	16.2
2004 5/.....	16.0	2.8	3.5	3.1	5.0	28.5	29.1	19.9	15.7	15.5	15.5
2003 5/.....	15.9	3.1	3.6	3.0	4.9	28.5	29.3	18.9	15.3	15.4	16.0
2002.....	16.3	2.4	3.0	3.1	4.8	31.3	30.2	19.5	16.4	15.8	15.9
2001.....	15.8	3.0	3.5	3.4	4.7	28.5	29.0	19.9	15.7	16.0	15.5
2000.....	15.8	3.3	3.7	3.7	5.1	29.7	29.4	18.7	15.5	15.8	14.8
1999.....	15.6	4.1	3.4	3.8	5.0	29.7	27.7	18.4	15.4	15.0	14.6

Non-Hispanic White, male

2007 5/.....	21.8	2.7	3.0	2.6	4.7	31.5	42.8	30.0	23.3	22.3	22.7
2006 5/.....	22.0	2.6	2.7	2.8	3.9	31.8	44.4	30.7	24.7	22.0	22.3
2005 5/.....	22.4	2.8	3.2	3.4	4.7	32.7	43.7	30.1	24.2	22.0	23.2
2004 5/.....	22.0	3.3	3.6	3.5	6.0	35.4	42.9	29.3	22.0	22.1	22.4
2003 5/.....	21.8	3.2	4.2	3.2	6.0	35.9	42.4	27.5	22.6	21.6	23.1
2002.....	22.3	2.7	3.3	3.5	5.6	39.7	44.6	29.0	23.6	22.1	22.0
2001.....	21.7	2.9	3.9	3.7	5.9	36.9	43.5	29.3	22.9	21.8	22.0
2000.....	21.5	3.4	4.1	4.2	6.1	37.9	43.0	27.2	22.4	22.1	21.1
1999.....	21.1	4.7	3.6	4.3	5.9	37.0	40.8	26.9	21.8	20.7	20.4

Non-Hispanic White, female

2007 5/.....	9.1	2.6	2.3	2.2	3.2	18.4	13.5	10.1	8.3	7.9	8.5
2006 5/.....	9.7	2.5	3.0	2.6	3.2	18.7	14.8	9.8	8.9	9.5	9.2
2005 5/.....	9.8	2.2	3.1	2.7	3.2	18.2	14.0	9.8	9.5	9.0	9.2
2004 5/.....	10.2	2.3	3.5	2.6	4.0	21.2	14.8	10.3	9.3	8.8	8.6
2003 5/.....	10.2	2.9	3.0	2.9	3.7	20.7	15.8	10.1	8.0	9.2	9.0
2002.....	10.5	2.1	2.8	2.6	3.9	22.5	15.5	9.8	9.0	9.5	9.8
2001.....	10.1	3.1	3.1	3.0	3.5	19.6	14.2	10.3	8.4	10.2	8.9
2000.....	10.3	3.2	3.2	3.2	4.1	21.1	15.4	10.1	8.5	9.5	8.5
1999.....	10.3	3.5	3.1	3.3	4.0	22.0	14.3	9.8	9.0	9.3	8.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic Black, both sexes

2007 5/.....	14.2	4.1	4.5	3.1	3.9	15.4	23.1	19.8	18.9	18.6	16.0
2006 5/.....	14.7	4.7	5.0	3.9	3.8	15.5	23.0	21.7	20.9	17.1	16.3
2005 5/.....	14.4	5.7	4.5	3.7	4.2	17.1	22.9	19.1	19.8	17.4	16.1
2004 5/.....	14.6	4.9	5.0	3.9	4.7	16.2	22.2	21.9	19.2	17.0	15.6
2003 5/.....	14.7	4.4	5.0	3.6	4.5	15.9	23.1	20.2	18.1	17.3	17.9
2002.....	14.8	5.4	4.9	4.3	4.3	17.7	24.5	23.3	17.6	18.4	16.1
2001.....	15.2	*	5.3	5.1	4.9	18.6	26.1	20.7	18.5	17.2	18.1
2000.....	15.5	6.9	5.6	4.3	4.7	16.9	26.9	23.7	18.6	16.9	18.3
1999.....	15.5	7.6	6.5	5.5	4.8	18.3	25.6	20.9	17.6	18.8	16.9

Non-Hispanic Black, male

2007 5/.....	21.9	*	4.6	3.6	4.5	21.3	35.0	32.1	31.3	30.6	26.5
2006 5/.....	22.2	*	5.7	4.7	4.5	22.1	33.8	34.2	34.9	27.7	24.9
2005 5/.....	21.8	*	4.9	3.7	5.2	23.2	34.8	31.1	33.0	28.2	25.1
2004 5/.....	21.7	*	5.2	4.1	5.3	22.2	32.2	34.8	31.2	26.5	23.9
2003 5/.....	21.7	*	5.0	4.3	5.4	22.2	35.8	31.6	28.9	27.3	26.9
2002.....	22.0	*	6.3	4.8	5.1	23.9	36.9	37.7	28.0	28.2	23.3
2001.....	22.7	*	5.6	6.5	6.5	26.1	39.4	32.7	29.3	25.6	28.2
2000.....	23.1	7.2	6.0	4.5	6.4	23.2	40.3	38.0	29.3	26.9	29.0
1999.....	22.8	*	7.0	6.5	6.0	25.3	38.8	33.6	28.0	27.8	26.0

Non-Hispanic Black, female

2007 5/.....	7.2	*	4.4	2.6	3.3	9.4	10.9	8.1	7.6	8.0	6.7
2006 5/.....	7.9	*	4.3	3.2	3.0	8.8	11.9	9.8	8.3	7.8	8.8
2005 5/.....	7.7	6.5	4.1	3.6	3.1	11.0	10.8	7.9	7.9	7.7	8.1
2004 5/.....	8.1	*	4.9	3.6	4.0	10.1	12.2	9.8	8.4	8.5	8.2
2003 5/.....	8.3	*	4.9	2.9	3.5	9.4	10.5	9.6	8.4	8.5	10.1
2002.....	8.3	*	3.5	3.9	3.4	11.4	12.5	10.1	8.2	9.7	9.7
2001.....	8.5	*	4.9	3.7	3.2	10.8	13.4	9.7	8.8	9.7	9.1
2000.....	8.5	*	5.2	4.1	3.1	10.5	14.2	10.6	9.0	8.2	8.8
1999.....	8.9	9.3	5.9	4.5	3.6	11.2	13.1	9.4	8.4	10.8	8.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins 3/, both sexes

2007.....	0.4	*	*	*	0.2	0.3	0.5	0.3	0.3	0.4	0.5
2006.....	0.4	*	*	*	0.2	0.4	0.4	0.4	0.3	0.5	0.5
2005.....	0.4	*	0.2	*	0.2	0.4	0.5	0.4	0.4	0.5	0.5
2004.....	0.4	*	*	0.1	0.2	0.5	0.4	0.5	0.4	0.4	0.5
2003.....	0.5	*	*	0.1	0.2	0.6	0.7	0.5	0.5	0.5	0.6
2002.....	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5	0.5
2001.....	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6
2000.....	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7	0.7
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9	0.9

All origins 3/, male

2007.....	0.6	*	*	*	*	0.5	0.7	0.6	0.5	0.6	0.7
2006.....	0.6	*	*	*	0.3	0.5	0.7	0.6	0.4	0.8	0.9
2005.....	0.7	*	*	*	0.3	0.7	0.7	0.7	0.6	0.7	0.8
2004.....	0.6	*	*	*	0.2	0.7	0.7	0.8	0.7	0.7	0.8
2003.....	0.7	*	*	*	0.2	0.9	1.0	0.8	0.8	0.8	0.9
2002.....	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8	0.9
2001.....	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1	1.0
2000.....	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1	1.2
1999.....	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4	1.5

All origins 3/, female

2007.....	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*
2006.....	0.2	*	*	*	*	0.2	*	*	*	*	*
2005.....	0.2	*	*	*	*	*	*	*	*	0.2	0.2
2004.....	0.2	*	*	*	*	0.2	*	*	*	*	0.2
2003.....	0.2	*	*	*	*	0.3	0.2	0.2	*	0.2	0.3
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.2
2001.....	0.2	*	*	*	*	0.2	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2	0.3
1999.....	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic 4/, both sexes

2007.....	0.4	*	*	*	0.2	0.3	0.5	0.3	0.3	0.4	0.4
2006.....	0.4	*	*	*	0.2	0.4	0.4	0.4	0.3	0.5	0.5
2005.....	0.4	*	0.2	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5
2004.....	0.4	*	*	0.1	0.1	0.5	0.5	0.5	0.4	0.4	0.5
2003.....	0.5	*	*	*	0.2	0.7	0.7	0.5	0.5	0.5	0.6
2002.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.4	0.5	0.5
2001.....	0.5	*	*	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6
2000.....	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.6	0.6	0.7
1999.....	0.7	*	0.3	0.3	0.3	0.9	1.0	0.7	0.7	0.9	0.9

Non-Hispanic 4/, male

2007.....	0.6	*	*	*	*	0.5	0.7	0.6	0.5	0.6	0.7
2006.....	0.6	*	*	*	0.2	0.5	0.7	0.6	0.4	0.8	0.8
2005.....	0.7	*	*	*	0.3	0.7	0.7	0.6	0.7	0.7	0.8
2004.....	0.6	*	*	*	*	0.7	0.8	0.7	0.6	0.6	0.8
2003.....	0.7	*	*	*	0.2	1.0	1.1	0.7	0.8	0.8	0.8
2002.....	0.6	*	*	*	0.4	0.4	0.7	0.5	0.6	0.8	0.9
2001.....	0.7	*	*	*	0.4	0.7	0.8	0.6	0.8	1.0	1.0
2000.....	0.8	*	*	*	0.4	1.0	1.0	0.8	0.8	1.0	1.2
1999.....	1.1	*	0.4	0.4	0.5	1.2	1.7	1.2	1.0	1.3	1.4

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	*	0.2	0.2	*	*	*	*
2006.....	0.2	*	*	*	*	*	*	*	*	*	*
2005.....	0.2	*	*	*	*	*	*	*	*	0.2	0.3
2004.....	0.2	*	*	*	*	*	*	*	*	*	0.2
2003.....	0.2	*	*	*	*	0.4	0.3	*	*	0.2	0.3
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	0.3	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	0.3	*	*	0.3	0.3	0.2
1999.....	0.4	*	*	*	*	0.6	0.3	0.3	0.5	0.4	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic White, both sexes

2007 5/.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.3	0.4
2006 5/.....	0.4	*	*	*	0.2	0.4	0.5	0.3	0.4	0.4	0.5
2005 5/.....	0.4	*	*	*	0.2	0.5	0.4	0.3	0.4	0.5	0.5
2004 5/.....	0.4	*	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5
2003 5/.....	0.5	*	*	*	0.2	0.8	0.7	0.5	0.5	0.5	0.5
2002.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.4	0.5	0.4
2001.....	0.5	*	*	*	0.3	0.5	0.5	0.4	0.4	0.5	0.5
2000.....	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.5	0.6	0.6
1999.....	0.8	*	0.3	0.3	0.4	0.9	1.0	0.8	0.7	0.9	0.9

Non-Hispanic White, male

2007 5/.....	0.5	*	*	*	*	0.5	0.8	0.6	0.5	0.5	0.6
2006 5/.....	0.6	*	*	*	*	0.6	0.7	0.5	0.5	0.7	0.8
2005 5/.....	0.6	*	*	*	*	0.6	0.6	0.6	0.6	0.7	0.7
2004 5/.....	0.6	*	*	*	*	0.7	0.7	0.6	0.6	0.6	0.8
2003 5/.....	0.7	*	*	*	*	1.1	1.2	0.7	0.7	0.7	0.9
2002.....	0.6	*	*	*	0.4	0.5	0.8	0.5	0.6	0.8	0.7
2001.....	0.7	*	*	*	0.3	0.7	0.8	0.7	0.7	0.9	0.8
2000.....	0.8	*	*	*	0.4	1.1	1.0	0.8	0.8	1.0	1.1
1999.....	1.1	*	*	0.4	0.5	1.2	1.6	1.2	1.0	1.3	1.4

Non-Hispanic White, female

2007 5/.....	0.1	*	*	*	*	*	*	*	*	*	*
2006 5/.....	0.2	*	*	*	*	*	*	*	*	*	*
2005 5/.....	0.2	*	*	*	*	*	*	*	*	*	0.3
2004 5/.....	0.2	*	*	*	*	*	*	*	*	*	0.2
2003 5/.....	0.2	*	*	*	*	0.4	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	0.4	*	*	*	*	*
1999.....	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins 3/, both sexes

2007.....	0.6	*	*	*	*	0.3	0.6	0.5	0.7	0.6	0.7
2006.....	0.6	*	*	0.1	0.2	0.3	0.6	0.7	0.7	0.7	0.9
2005.....	0.6	*	*	*	0.2	0.3	0.6	0.5	0.7	0.7	0.8
2004.....	0.7	*	*	*	0.1	0.4	0.6	0.8	0.7	0.9	0.8
2003.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.6	0.9	0.9
2002.....	0.7	*	*	0.1	0.2	0.4	0.6	0.5	0.7	0.9	0.9
2001.....	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9	1.2
2000.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9	1.0
1999.....	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9	0.9

All origins 3/, male

2007.....	1.0	*	*	*	*	0.4	0.9	0.8	1.0	1.1	1.1
2006.....	1.0	*	*	*	0.3	0.5	1.0	1.1	1.0	1.2	1.4
2005.....	1.0	*	*	*	*	0.4	0.9	0.9	1.1	1.1	1.4
2004.....	1.1	*	*	*	*	0.5	0.9	1.2	1.2	1.4	1.3
2003.....	1.1	*	*	*	0.2	0.6	1.0	0.8	1.0	1.4	1.6
2002.....	1.1	*	*	*	0.2	0.6	0.9	0.9	1.2	1.5	1.5
2001.....	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4	1.8
2000.....	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5	1.7
1999.....	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4	1.5

All origins 3/, female

2007.....	0.2	*	*	*	*	*	0.2	0.2	0.3	*	0.3
2006.....	0.3	*	*	*	*	0.2	0.3	*	0.3	0.2	0.4
2005.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3	0.3
2004.....	0.3	*	*	*	*	0.2	0.3	0.3	0.2	0.3	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.3	0.2	0.3	0.3
2002.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3	0.3
2001.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4	0.5
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2	0.3
1999.....	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	0.7	*	*	*	*	0.3	0.6	0.6	0.7	0.7	0.7
2006.....	0.7	*	*	0.1	0.2	0.4	0.7	0.7	0.8	0.8	0.9
2005.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.8	0.8	0.9
2004.....	0.7	*	*	*	0.2	0.4	0.7	0.9	0.8	1.0	0.9
2003.....	0.7	*	*	0.2	0.2	0.4	0.6	0.6	0.7	0.9	1.0
2002.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.8	0.9	0.9
2001.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.9	0.8	1.1
2000.....	0.7	*	*	0.1	0.2	0.5	0.7	0.8	0.9	0.9	1.0
1999.....	0.7	*	*	0.2	0.2	0.4	0.7	0.9	0.9	1.0	1.0

Non-Hispanic 4/, male

2007.....	1.1	*	*	*	*	0.5	1.0	0.9	1.1	1.2	1.2
2006.....	1.1	*	*	*	0.3	0.5	1.2	1.3	1.1	1.3	1.5
2005.....	1.1	*	*	*	*	0.5	1.0	1.0	1.3	1.2	1.6
2004.....	1.2	*	*	*	*	0.5	1.0	1.3	1.4	1.6	1.4
2003.....	1.2	*	*	*	*	0.6	1.1	0.9	1.1	1.5	1.6
2002.....	1.2	*	*	*	*	0.6	1.0	1.0	1.3	1.6	1.6
2001.....	1.2	*	*	*	*	0.8	1.0	1.3	1.5	1.3	1.9
2000.....	1.2	*	*	*	*	0.7	1.3	1.5	1.5	1.6	1.8
1999.....	1.2	*	*	*	0.4	0.7	1.0	1.5	1.5	1.5	1.6

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	*	*	0.3	*	0.3	*	0.3
2006.....	0.3	*	*	*	*	0.2	0.3	*	0.4	0.3	0.4
2005.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3
2004.....	0.3	*	*	*	*	0.2	0.3	0.4	*	0.3	0.4
2003.....	0.3	*	*	*	*	*	*	0.3	0.3	0.3	0.3
2002.....	0.2	*	*	*	*	0.3	*	*	0.3	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.3	0.4	*	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic White, both sexes

2007 5/.....	0.7	*	*	*	*	0.4	0.7	0.6	0.8	0.8	0.8
2006 5/.....	0.8	*	*	0.2	0.2	0.4	0.8	0.8	0.8	0.9	1.1
2005 5/.....	0.8	*	*	*	0.2	0.4	0.7	0.7	0.9	0.8	1.0
2004 5/.....	0.8	*	*	*	0.2	0.4	0.8	1.0	0.9	1.1	1.0
2003 5/.....	0.8	*	*	0.2	0.2	0.4	0.7	0.7	0.8	1.0	1.1
2002.....	0.8	*	*	*	0.2	0.5	0.7	0.7	0.9	1.0	1.1
2001.....	0.8	*	*	*	0.2	0.6	0.8	1.0	1.0	0.9	1.3
2000.....	0.8	*	*	*	0.2	0.5	0.9	1.0	0.9	1.0	1.2
1999.....	0.8	*	*	0.2	0.3	0.5	0.8	1.0	1.0	1.1	1.1

Non-Hispanic White, male

2007 5/.....	1.2	*	*	*	*	0.5	1.0	1.0	1.2	1.4	1.3
2006 5/.....	1.3	*	*	*	0.4	0.6	1.4	1.4	1.2	1.4	1.6
2005 5/.....	1.3	*	*	*	*	0.5	1.1	1.2	1.5	1.3	1.8
2004 5/.....	1.3	*	*	*	*	0.5	1.1	1.5	1.6	1.7	1.5
2003 5/.....	1.3	*	*	*	*	0.6	1.1	1.0	1.3	1.7	1.8
2002.....	1.3	*	*	*	*	0.7	1.2	1.1	1.5	1.8	1.8
2001.....	1.3	*	*	*	*	1.0	1.2	1.5	1.7	1.5	2.1
2000.....	1.4	*	*	*	*	0.7	1.6	1.7	1.6	1.7	2.0
1999.....	1.3	*	*	*	0.4	0.7	1.2	1.7	1.6	1.6	1.7

Non-Hispanic White, female

2007 5/.....	0.3	*	*	*	*	*	*	*	0.4	*	0.3
2006 5/.....	0.3	*	*	*	*	*	*	*	0.4	0.4	0.5
2005 5/.....	0.3	*	*	*	*	*	*	*	*	0.4	0.3
2004 5/.....	0.3	*	*	*	*	0.3	0.4	0.5	*	0.4	0.4
2003 5/.....	0.3	*	*	*	*	*	*	*	0.3	0.4	0.4
2002.....	0.3	*	*	*	*	0.3	*	*	*	0.3	0.3
2001.....	0.3	*	*	*	*	*	0.3	0.5	*	0.4	0.4
2000.....	0.3	*	*	*	*	0.3	*	*	*	*	0.4
1999.....	0.3	*	*	*	*	*	0.3	0.4	0.4	0.5	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes

2007.....	25.5	27.2	6.1	2.1	2.0	7.6	15.8	18.4	18.7	20.8	25.9
2006.....	24.4	24.3	6.1	2.2	1.8	7.3	15.5	17.9	17.7	20.4	26.5
2005.....	23.4	22.8	6.3	2.2	2.1	7.1	14.2	15.5	16.2	19.1	24.6
2004.....	21.8	22.2	6.1	2.3	2.2	6.9	12.6	13.7	13.8	18.3	23.3
2003.....	21.0	19.9	6.8	2.0	2.1	6.3	12.3	12.7	13.7	18.3	22.9
2002.....	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8	22.3
2001.....	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7	19.4
2000.....	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8	18.6
1999.....	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3	18.8

All origins 3/, male

2007.....	31.2	30.2	7.8	2.5	2.7	11.9	24.5	27.6	27.3	28.9	35.2
2006.....	30.4	28.3	7.2	2.7	2.5	11.7	24.5	27.3	26.3	29.1	36.6
2005.....	29.0	24.0	7.9	2.6	3.0	11.2	22.2	23.6	23.7	27.1	33.9
2004.....	26.8	25.1	7.4	2.9	3.1	11.0	19.8	21.1	20.5	25.7	31.9
2003.....	26.0	22.4	8.5	2.5	3.0	10.0	19.4	19.6	20.2	26.2	32.4
2002.....	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9	32.3
2001.....	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8	28.7
2000.....	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2	28.3
1999.....	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4	29.1

All origins 3/, female

2007.....	19.9	24.1	4.3	1.6	1.3	3.2	6.6	8.7	9.9	12.7	16.6
2006.....	18.6	20.2	5.0	1.7	1.1	2.7	5.9	8.1	8.9	11.7	16.4
2005.....	18.0	21.5	4.5	1.7	1.2	2.7	5.6	7.0	8.6	10.9	15.3
2004.....	16.9	19.3	4.8	1.6	1.3	2.6	4.9	6.0	7.0	10.9	14.7
2003.....	16.2	17.3	5.0	1.6	1.2	2.3	4.9	5.5	7.0	10.4	13.5
2002.....	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7	12.5
2001.....	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5	10.2
2000.....	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4	9.0
1999.....	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2	8.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

Hispanic, both sexes

2007.....	12.3	13.2	4.7	1.5	1.4	6.5	11.4	11.1	11.5	13.2	17.3
2006.....	12.4	14.8	5.0	1.5	1.5	5.8	10.7	10.9	12.9	14.0	18.7
2005.....	12.0	13.7	5.2	1.2	1.0	6.1	11.0	10.5	11.7	13.5	17.1
2004.....	10.6	13.1	4.3	1.1	1.2	4.9	9.1	9.3	9.2	13.7	16.0
2003.....	11.0	10.6	5.6	1.9	1.4	5.8	9.2	10.2	10.3	14.0	16.8
2002.....	10.8	12.1	5.1	1.6	1.4	6.2	9.4	8.6	10.2	13.2	18.6
2001.....	10.1	13.6	5.0	1.8	1.4	5.1	8.0	8.6	10.3	14.4	15.8
2000.....	10.2	9.9	5.2	1.7	1.4	6.0	8.5	8.4	11.6	14.7	16.2
1999.....	11.1	12.4	6.4	1.8	1.7	6.0	9.4	9.1	12.1	16.6	18.0

Hispanic, male

2007.....	17.9	14.6	6.5	1.9	1.7	10.4	19.1	17.5	18.0	20.6	27.4
2006.....	18.0	16.1	6.0	2.0	2.1	9.6	17.2	16.8	20.1	22.1	29.4
2005.....	17.3	13.9	6.8	1.3	1.4	10.0	18.0	16.5	18.7	21.4	26.1
2004.....	15.2	14.9	5.1	1.3	1.8	7.8	14.4	14.8	14.5	21.0	24.5
2003.....	16.1	11.8	7.3	2.2	2.0	9.7	14.7	16.1	16.4	22.2	26.4
2002.....	16.1	15.7	6.4	2.1	1.7	10.5	15.3	14.2	16.8	20.5	29.7
2001.....	14.8	14.2	6.7	2.3	1.5	8.5	12.8	13.8	16.1	23.1	25.7
2000.....	15.3	10.9	5.9	1.8	1.3	9.5	14.5	13.8	19.5	23.4	28.3
1999.....	16.9	15.0	7.7	2.4	2.2	9.6	15.9	15.1	20.0	27.1	29.5

Hispanic, female

2007.....	6.4	11.7	2.8	1.2	*	2.3	2.5	3.1	3.9	4.7	6.1
2006.....	6.4	13.5	4.0	1.1	*	1.7	3.0	3.5	4.3	4.7	6.9
2005.....	6.2	13.6	3.6	1.1	*	2.0	2.7	3.1	3.6	4.5	7.2
2004.....	5.8	11.3	3.4	*	*	1.7	2.6	2.5	3.0	5.3	6.7
2003.....	5.4	9.4	3.9	1.5	*	1.5	2.4	2.9	3.2	4.7	6.5
2002.....	5.2	8.3	3.7	1.2	1.2	1.6	2.1	1.7	2.6	5.2	6.8
2001.....	5.2	13.0	3.3	1.3	1.3	*	2.1	2.3	3.6	4.7	5.3
2000.....	4.9	8.8	4.4	1.5	1.4	1.9	*	2.2	2.6	5.1	3.6
1999.....	5.0	9.8	5.2	1.2	*	2.0	1.4	2.1	3.1	5.1	6.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	27.7	31.6	6.5	2.2	2.2	7.9	16.7	20.1	20.5	22.4	27.2
2006.....	26.4	27.1	6.3	2.3	1.9	7.7	16.5	19.6	18.8	21.7	27.6
2005.....	25.2	25.3	6.6	2.4	2.4	7.3	14.8	16.7	17.2	20.0	25.6
2004.....	23.5	24.8	6.6	2.5	2.4	7.3	13.3	14.7	14.8	19.1	24.2
2003.....	22.5	22.4	7.1	2.1	2.3	6.3	13.0	13.2	14.3	18.9	23.6
2002.....	21.6	22.3	6.7	2.5	2.6	6.7	11.4	12.5	14.2	18.4	22.7
2001.....	20.2	22.0	7.2	2.7	2.6	6.4	10.7	10.9	11.8	16.9	19.7
2000.....	19.2	20.7	7.9	2.9	2.7	6.4	9.7	10.3	11.8	15.8	18.7
1999.....	19.3	18.5	8.0	2.9	2.9	6.1	9.3	10.1	12.0	15.0	18.7

Non-Hispanic 4/, male

2007.....	33.6	34.9	8.2	2.7	2.9	12.2	25.6	30.3	29.8	30.7	36.5
2006.....	32.6	31.8	7.4	2.8	2.6	12.2	26.1	30.2	27.8	30.4	37.7
2005.....	30.9	26.8	8.3	2.9	3.4	11.5	23.1	25.5	24.8	28.1	35.0
2004.....	28.7	27.9	8.0	3.3	3.3	11.5	21.1	22.7	21.9	26.5	32.9
2003.....	27.5	25.2	8.7	2.5	3.2	10.0	20.5	20.4	20.9	26.7	33.1
2002.....	26.5	25.2	8.5	3.3	3.6	10.5	18.5	19.5	21.3	26.6	32.4
2001.....	25.0	24.9	8.7	3.5	3.6	10.1	17.7	17.2	17.8	24.8	28.8
2000.....	23.9	23.2	9.7	3.7	3.8	10.4	15.7	16.7	18.0	24.1	28.0
1999.....	24.0	21.5	9.6	3.7	4.2	10.0	15.1	16.4	18.8	22.7	28.8

Non-Hispanic 4/, female

2007.....	22.1	28.1	4.8	1.7	1.4	3.3	7.4	9.9	11.3	14.2	18.2
2006.....	20.6	22.2	5.2	1.8	1.1	2.9	6.5	9.0	9.9	13.0	17.8
2005.....	19.8	23.7	4.8	1.8	1.4	2.8	6.2	7.8	9.6	12.0	16.4
2004.....	18.6	21.6	5.2	1.8	1.4	2.8	5.3	6.7	7.8	11.9	15.7
2003.....	17.8	19.5	5.3	1.6	1.3	2.5	5.3	6.0	7.7	11.3	14.3
2002.....	16.9	19.4	4.9	1.7	1.5	2.7	4.2	5.5	7.2	10.3	13.2
2001.....	15.7	19.1	5.6	1.9	1.5	2.5	3.6	4.6	5.8	9.1	10.7
2000.....	14.7	18.0	6.0	1.9	1.4	2.2	3.6	3.9	5.6	7.7	9.6
1999.....	14.8	15.2	6.3	2.1	1.6	2.1	3.4	3.9	5.3	7.4	8.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

Non-Hispanic White, both sexes

2007 5/.....	30.6	24.9	6.3	1.8	1.8	8.7	19.6	23.5	23.8	24.7	29.4
2006 5/.....	28.8	21.7	5.9	1.8	1.8	8.3	19.2	22.9	21.2	23.4	29.1
2005 5/.....	27.3	21.9	6.4	1.9	2.2	7.7	16.9	19.1	18.9	21.5	27.0
2004 5/.....	25.6	21.2	6.5	2.0	2.0	7.9	15.3	16.8	16.5	20.4	25.7
2003 5/.....	24.4	18.4	6.9	1.7	1.8	6.8	14.9	14.8	15.7	20.0	24.6
2002.....	23.1	17.2	6.1	1.9	2.1	6.9	12.8	13.8	15.5	19.1	23.4
2001.....	21.5	17.8	7.1	2.3	2.2	6.5	11.9	11.6	12.5	17.4	19.5
2000.....	20.1	16.6	7.3	2.3	2.3	6.4	10.5	10.9	12.1	16.0	18.4
1999.....	20.0	14.7	7.4	2.2	2.5	6.1	9.6	10.6	11.9	15.2	18.3

Non-Hispanic White, male

2007 5/.....	36.5	28.0	7.9	2.2	2.6	13.4	30.0	35.2	34.5	33.5	38.8
2006 5/.....	34.8	25.0	6.9	2.3	2.4	12.8	30.4	34.8	30.9	32.4	39.2
2005 5/.....	32.7	22.8	8.4	2.4	3.0	12.2	26.5	29.0	27.1	30.0	36.5
2004 5/.....	30.7	25.0	8.0	2.5	2.8	12.5	24.2	25.6	24.2	28.2	34.7
2003 5/.....	29.2	20.8	8.5	2.1	2.7	10.8	23.5	22.7	22.8	27.8	34.5
2002.....	27.8	20.4	7.6	2.4	3.0	10.8	20.6	21.2	23.1	27.2	33.4
2001.....	25.9	20.0	8.8	2.8	3.0	10.2	19.6	18.2	18.9	25.6	28.7
2000.....	24.5	19.1	9.1	3.0	3.4	10.4	17.0	17.5	18.6	24.5	27.4
1999.....	24.3	17.1	8.9	2.7	3.6	9.8	15.4	17.1	18.7	22.9	28.0

Non-Hispanic White, female

2007 5/.....	24.9	21.6	4.6	1.5	1.0	3.8	8.6	11.6	13.1	15.9	20.0
2006 5/.....	23.1	18.2	4.8	1.3	1.0	3.5	7.5	10.8	11.4	14.3	19.1
2005 5/.....	22.0	21.1	4.4	1.4	1.3	3.0	6.9	9.0	10.5	12.9	17.5
2004 5/.....	20.7	17.3	4.9	1.4	1.1	3.1	6.1	7.7	8.7	12.7	16.8
2003 5/.....	19.7	16.0	5.1	1.3	1.0	2.7	6.1	6.7	8.6	12.1	14.7
2002.....	18.6	13.9	4.6	1.2	1.2	2.8	4.8	6.2	7.8	10.9	13.5
2001.....	17.2	15.5	5.3	1.8	1.2	2.5	4.1	4.9	6.0	9.2	10.3
2000.....	16.0	14.0	5.4	1.5	1.2	2.3	3.8	4.2	5.6	7.5	9.3
1999.....	16.0	12.3	5.9	1.7	1.4	2.2	3.6	4.0	5.1	7.4	8.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

Non-Hispanic Black, both sexes

2007 5/.....	19.9	59.8	8.0	3.7	3.4	5.4	7.8	11.1	13.4	18.3	23.5
2006 5/.....	20.7	49.9	9.0	4.2	2.6	6.1	8.4	12.1	14.5	20.9	27.8
2005 5/.....	20.5	42.9	7.9	4.2	3.5	5.7	8.4	10.6	15.0	18.7	24.8
2004 5/.....	18.2	42.0	7.6	4.8	3.9	5.2	7.4	10.3	11.5	18.0	22.6
2003 5/.....	17.7	40.3	8.5	3.8	3.7	4.7	7.6	9.7	12.0	18.2	23.5
2002.....	18.3	44.5	9.5	4.9	4.5	6.1	7.6	10.2	12.5	19.4	23.4
2001.....	18.1	41.6	8.7	4.3	4.3	6.1	7.5	9.7	11.3	18.0	25.5
2000.....	18.0	39.5	10.9	5.4	4.1	6.5	7.8	9.8	12.8	18.3	24.9
1999.....	19.4	36.4	11.1	5.8	4.8	6.5	9.2	10.5	14.8	16.9	25.9

Non-Hispanic Black, male

2007 5/.....	26.6	61.9	9.8	4.6	4.1	8.7	11.3	16.9	20.2	26.3	33.8
2006 5/.....	28.8	59.5	10.5	5.1	3.5	10.8	12.8	19.7	23.0	30.8	40.5
2005 5/.....	27.9	47.3	9.0	4.9	5.1	9.3	12.9	16.5	21.4	26.9	34.6
2004 5/.....	24.6	43.0	9.1	6.5	5.1	8.4	11.5	16.2	17.6	24.6	31.8
2003 5/.....	24.1	45.3	10.8	4.4	5.2	7.5	11.6	15.0	18.2	27.3	33.1
2002.....	25.5	46.7	12.5	6.4	6.7	10.4	12.9	16.8	19.2	29.8	33.5
2001.....	25.0	48.2	10.0	6.2	5.9	9.8	12.4	15.4	16.9	26.4	36.9
2000.....	25.5	42.3	12.9	7.2	5.8	10.7	12.8	16.4	19.4	27.5	38.2
1999.....	27.5	42.3	13.7	7.7	6.8	11.2	16.2	17.2	22.8	25.9	41.2

Non-Hispanic Black, female

2007 5/.....	13.7	57.6	6.1	2.8	2.6	1.9	4.3	5.6	7.2	11.2	14.4
2006 5/.....	13.3	39.9	7.5	3.4	1.6	1.4	3.9	4.9	6.8	12.1	16.7
2005 5/.....	13.7	38.3	6.8	3.4	1.8	2.1	3.8	4.9	9.3	11.3	16.1
2004 5/.....	12.3	40.9	6.1	3.0	2.6	1.9	3.3	4.7	6.0	12.0	14.4
2003 5/.....	12.0	35.1	6.2	3.2	2.2	1.8	3.6	4.7	6.5	10.2	15.1
2002.....	11.8	42.3	6.5	3.4	2.4	1.7	2.5	4.2	6.4	10.1	14.6
2001.....	11.8	34.6	7.4	2.3	2.6	2.2	2.8	4.5	6.3	10.5	15.4
2000.....	11.3	36.6	8.8	3.6	2.4	2.2	3.1	3.9	6.9	10.2	13.0
1999.....	12.0	30.3	8.3	3.9	2.6	1.7	2.6	4.5	7.6	9.0	12.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Falls (W00-W19)

All origins 3/, both sexes

2007.....	7.5	0.6	0.2	*	0.1	0.4	0.7	0.8	0.8	1.2	1.6
2006.....	7.0	0.6	0.2	*	0.1	0.4	0.8	0.7	0.9	1.1	1.8
2005.....	6.6	*	0.2	*	*	0.4	0.8	0.7	0.8	1.1	1.7
2004.....	6.4	0.6	0.3	*	0.1	0.4	0.7	0.8	0.8	1.1	1.8
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2	1.7
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2	1.8
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2	1.6
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0	1.6
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2	1.7

All origins 3/, male

2007.....	7.8	*	0.3	*	*	0.6	1.2	1.4	1.4	1.9	2.5
2006.....	7.3	*	0.3	*	*	0.6	1.3	1.1	1.5	1.8	2.8
2005.....	7.0	*	0.2	*	*	0.6	1.2	1.2	1.3	1.8	2.6
2004.....	6.8	*	0.4	*	*	0.7	1.3	1.3	1.4	1.8	2.9
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0	2.7
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7	2.9
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1	2.8
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8	2.7
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0	2.8

All origins 3/, female

2007.....	7.2	*	*	*	*	*	*	*	0.3	0.4	0.6
2006.....	6.6	*	*	*	*	*	*	*	0.3	0.4	0.7
2005.....	6.3	*	*	*	*	*	0.3	*	0.3	0.4	0.7
2004.....	6.0	*	*	*	*	*	*	0.2	0.2	0.5	0.7
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3	0.6
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6	0.7
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3	0.6
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Falls (W00-W19)

Non-Hispanic 4/, both sexes

2007.....	8.3	0.6	0.2	*	*	0.4	0.7	0.7	0.7	1.1	1.6
2006.....	7.7	*	0.2	*	*	0.4	0.7	0.6	0.8	1.0	1.7
2005.....	7.3	*	0.2	*	*	0.4	0.7	0.6	0.7	1.1	1.7
2004.....	7.0	*	0.3	*	0.1	0.4	0.7	0.7	0.8	1.1	1.8
2003.....	6.5	*	0.3	*	0.1	0.5	0.7	0.7	0.7	1.2	1.6
2002.....	6.2	*	0.2	*	0.1	0.4	0.8	0.6	0.8	1.1	1.7
2001.....	5.7	*	0.2	0.1	0.2	0.4	0.9	0.8	0.8	1.2	1.6
2000.....	5.1	*	0.2	*	*	0.5	0.7	0.6	0.7	1.0	1.6
1999.....	5.1	*	0.4	0.1	0.2	0.6	0.6	0.7	0.9	1.2	1.7

Non-Hispanic 4/, male

2007.....	8.6	*	*	*	*	0.6	1.1	1.3	1.2	1.8	2.5
2006.....	8.0	*	0.3	*	*	0.6	1.2	0.9	1.3	1.6	2.7
2005.....	7.6	*	*	*	*	0.6	1.1	1.1	1.2	1.7	2.7
2004.....	7.4	*	0.4	*	*	0.6	1.2	1.1	1.3	1.6	2.9
2003.....	6.8	*	0.5	*	*	0.7	1.1	1.1	1.1	2.1	2.5
2002.....	6.5	*	0.3	*	*	0.6	1.3	1.2	1.3	1.7	2.8
2001.....	6.2	*	0.3	*	0.2	0.7	1.5	1.4	1.3	2.0	2.6
2000.....	5.5	*	*	*	*	0.8	1.1	1.1	1.3	1.7	2.6
1999.....	5.5	*	0.5	*	*	1.0	1.1	1.3	1.4	1.9	2.7

Non-Hispanic 4/, female

2007.....	8.1	*	*	*	*	*	*	*	0.3	0.5	0.7
2006.....	7.4	*	*	*	*	*	*	*	0.3	0.4	0.7
2005.....	7.0	*	*	*	*	*	0.3	*	0.3	0.4	0.7
2004.....	6.7	*	*	*	*	*	*	*	0.3	0.5	0.7
2003.....	6.2	*	*	*	*	*	0.3	*	0.3	0.4	0.7
2002.....	5.9	*	*	*	*	*	*	*	0.3	0.6	0.7
2001.....	5.3	*	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	4.7	*	*	*	*	*	0.3	*	*	0.3	0.7
1999.....	4.6	*	*	*	*	*	*	*	0.3	0.5	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Falls (W00-W19)

Non-Hispanic White, both sexes

2007 5/.....	9.8	*	*	*	*	0.5	0.7	0.8	0.9	1.2	1.6
2006 5/.....	9.0	*	*	*	*	0.4	0.8	0.6	0.9	1.1	1.8
2005 5/.....	8.5	*	*	*	*	0.4	0.9	0.7	0.8	1.1	1.8
2004 5/.....	8.2	*	0.3	*	0.2	0.5	0.8	0.7	0.9	1.1	1.8
2003 5/.....	7.5	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2	1.7
2002.....	7.1	*	*	*	*	0.5	0.9	0.8	0.9	1.2	1.8
2001.....	6.6	*	0.2	*	0.2	0.5	1.0	0.9	0.9	1.2	1.5
2000.....	5.8	*	*	*	*	0.6	0.8	0.7	0.8	1.0	1.7
1999.....	5.7	*	0.4	*	0.2	0.7	0.7	0.8	0.8	1.3	1.7

Non-Hispanic White, male

2007 5/.....	10.0	*	*	*	*	0.6	1.2	1.4	1.4	1.9	2.5
2006 5/.....	9.2	*	*	*	*	0.7	1.3	1.0	1.4	1.7	2.8
2005 5/.....	8.7	*	*	*	*	0.7	1.4	1.3	1.2	1.8	2.9
2004 5/.....	8.5	*	*	*	*	0.7	1.4	1.1	1.5	1.7	3.0
2003 5/.....	7.7	*	0.5	*	*	0.8	1.3	1.3	1.2	2.1	2.6
2002.....	7.3	*	*	*	*	0.7	1.5	1.4	1.4	1.7	2.8
2001.....	7.0	*	*	*	*	0.8	1.9	1.6	1.4	2.0	2.5
2000.....	6.2	*	*	*	*	1.0	1.3	1.3	1.3	1.7	2.7
1999.....	6.1	*	0.5	*	*	1.1	1.2	1.4	1.4	2.0	2.7

Non-Hispanic White, female

2007 5/.....	9.7	*	*	*	*	*	*	*	0.4	0.5	0.7
2006 5/.....	8.8	*	*	*	*	*	*	*	0.3	0.4	0.8
2005 5/.....	8.3	*	*	*	*	*	0.3	*	*	0.4	0.8
2004 5/.....	7.9	*	*	*	*	*	*	*	*	0.4	0.7
2003 5/.....	7.3	*	*	*	*	*	*	*	0.4	0.4	0.7
2002.....	7.0	*	*	*	*	*	*	*	*	0.7	0.7
2001.....	6.2	*	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	5.5	*	*	*	*	*	*	*	*	0.3	0.7
1999.....	5.4	*	*	*	*	*	*	*	0.3	0.5	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental drowning and submersion (W65-W74)

All origins 3/, both sexes

2007.....	1.1	1.3	2.8	0.6	0.5	1.5	1.5	1.0	0.9	0.8	1.1
2006.....	1.2	1.2	2.8	0.7	0.6	1.5	1.4	1.2	0.9	1.0	1.1
2005.....	1.2	1.6	3.0	0.6	0.6	1.5	1.6	1.0	0.9	1.0	1.3
2004.....	1.1	1.5	2.7	0.7	0.7	1.5	1.3	1.1	0.8	0.9	1.0
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0	1.1
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2	1.1
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0	1.1
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1	1.0
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1	1.0

All origins 3/, male

2007.....	1.8	1.4	3.7	0.8	0.7	2.5	2.6	1.7	1.5	1.4	1.7
2006.....	1.9	1.3	3.5	1.0	0.8	2.6	2.5	2.2	1.6	1.6	1.8
2005.....	1.9	1.3	4.1	0.8	1.0	2.6	2.8	1.7	1.5	1.7	2.0
2004.....	1.8	1.8	3.2	0.9	1.0	2.6	2.3	1.9	1.4	1.6	1.7
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7	1.8
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0	1.9
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6	1.7
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9	1.7
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9	1.7

All origins 3/, female

2007.....	0.5	1.3	1.8	0.4	0.3	0.4	0.3	0.2	0.3	0.3	0.5
2006.....	0.5	1.2	2.1	0.4	0.2	0.3	0.4	0.3	*	0.4	0.5
2005.....	0.5	1.8	2.0	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.5
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.2	0.3	0.2	0.2	0.4
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental drowning and submersion (W65-W74)

Non-Hispanic 4/, both sexes

2007.....	1.2	1.4	3.0	0.7	0.5	1.4	1.3	1.0	0.9	0.9	1.1
2006.....	1.2	1.1	2.9	0.8	0.6	1.4	1.3	1.2	0.9	1.1	1.2
2005.....	1.2	1.5	3.1	0.7	0.7	1.4	1.4	1.0	0.9	1.0	1.3
2004.....	1.1	1.4	2.8	0.8	0.7	1.5	1.2	1.1	0.8	0.9	1.0
2003.....	1.1	1.5	3.0	0.6	0.7	1.3	1.3	0.9	0.8	1.0	1.1
2002.....	1.2	1.4	3.1	0.9	0.8	1.5	1.4	1.2	1.0	1.2	1.2
2001.....	1.1	1.6	3.1	0.8	0.9	1.6	1.3	1.1	0.7	1.0	1.1
2000.....	1.2	2.1	3.4	1.0	0.9	1.7	1.3	1.1	1.0	1.1	1.1
1999.....	1.2	1.7	3.3	1.0	0.9	1.7	1.4	1.1	1.1	1.1	1.0

Non-Hispanic 4/, male

2007.....	1.8	1.4	3.9	0.9	0.7	2.4	2.3	1.7	1.4	1.5	1.7
2006.....	1.9	*	3.5	1.1	0.8	2.4	2.1	2.1	1.7	1.7	1.8
2005.....	1.9	1.3	4.1	0.9	1.1	2.5	2.4	1.7	1.5	1.8	2.1
2004.....	1.8	1.8	3.4	1.0	1.0	2.7	2.1	1.9	1.3	1.6	1.7
2003.....	1.8	1.7	3.7	0.9	1.0	2.4	2.3	1.6	1.4	1.8	1.8
2002.....	1.9	1.6	4.1	1.3	1.2	2.5	2.6	2.0	1.7	2.1	1.9
2001.....	1.8	1.3	3.8	1.1	1.2	2.9	2.4	2.0	1.3	1.6	1.8
2000.....	2.0	2.4	4.3	1.4	1.4	3.1	2.2	1.9	1.8	1.9	1.8
1999.....	2.0	1.7	4.2	1.4	1.3	3.0	2.4	1.9	1.9	1.9	1.8

Non-Hispanic 4/, female

2007.....	0.5	1.3	2.0	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.5
2006.....	0.6	*	2.2	0.4	0.3	0.3	0.4	0.3	*	0.4	0.5
2005.....	0.5	1.8	2.0	0.5	0.3	0.2	0.3	0.3	0.3	0.3	0.5
2004.....	0.5	*	2.2	0.5	0.3	0.3	0.3	0.4	0.3	0.3	0.4
2003.....	0.5	*	2.3	0.4	0.3	*	0.3	*	*	0.3	0.4
2002.....	0.5	*	2.1	0.4	0.5	0.3	*	0.3	0.3	0.4	0.4
2001.....	0.5	1.9	2.4	0.5	0.5	0.3	0.3	0.3	*	0.3	0.4
2000.....	0.5	1.9	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	*	0.3	0.3	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental drowning and submersion (W65-W74)

Non-Hispanic White, both sexes

2007 5/.....	1.1	1.2	3.3	0.6	0.3	1.2	1.2	0.9	0.8	0.9	1.2
2006 5/.....	1.2	1.3	3.1	0.6	0.4	1.1	1.2	1.1	0.8	1.1	1.3
2005 5/.....	1.2	1.8	3.4	0.5	0.5	1.1	1.3	1.0	0.9	0.9	1.3
2004 5/.....	1.1	1.5	3.2	0.5	0.4	1.2	1.1	1.0	0.8	0.9	1.0
2003 5/.....	1.1	1.4	3.3	0.5	0.4	1.2	1.2	0.8	0.7	1.0	1.1
2002.....	1.1	1.4	3.3	0.6	0.5	1.2	1.3	1.0	1.0	1.2	1.2
2001.....	1.1	1.5	3.5	0.6	0.5	1.3	1.3	1.0	0.7	0.9	1.0
2000.....	1.1	1.7	3.7	0.8	0.6	1.4	1.1	1.0	1.0	1.1	1.0
1999.....	1.1	1.3	3.6	0.6	0.5	1.4	1.2	0.9	1.0	1.1	1.1

Non-Hispanic White, male

2007 5/.....	1.7	*	4.2	0.8	0.4	1.9	2.1	1.4	1.3	1.3	1.7
2006 5/.....	1.8	*	3.6	0.8	0.6	1.8	1.9	2.0	1.6	1.6	1.9
2005 5/.....	1.8	*	4.6	0.6	0.7	2.0	2.3	1.7	1.4	1.6	2.0
2004 5/.....	1.7	1.8	3.8	0.8	0.6	2.1	2.0	1.7	1.3	1.5	1.8
2003 5/.....	1.7	*	4.0	0.7	0.6	2.1	2.1	1.5	1.2	1.7	1.8
2002.....	1.7	*	4.3	0.8	0.7	2.0	2.3	1.8	1.6	1.9	1.9
2001.....	1.6	*	4.3	0.6	0.7	2.2	2.2	1.7	1.1	1.4	1.7
2000.....	1.8	1.9	4.7	1.0	0.9	2.5	1.9	1.7	1.7	1.9	1.7
1999.....	1.8	*	4.5	0.8	0.7	2.4	2.0	1.6	1.7	1.8	1.7

Non-Hispanic White, female

2007 5/.....	0.5	*	2.5	*	*	0.4	0.3	*	*	0.4	0.6
2006 5/.....	0.6	*	2.5	*	*	0.4	0.4	0.3	*	0.5	0.6
2005 5/.....	0.5	2.3	2.3	0.4	*	*	*	*	0.4	*	0.6
2004 5/.....	0.5	*	2.5	*	*	*	*	0.4	0.3	0.3	0.3
2003 5/.....	0.5	*	2.7	*	*	*	*	*	*	0.4	0.3
2002.....	0.5	*	2.2	*	*	*	*	*	*	0.4	0.5
2001.....	0.5	1.7	2.6	0.5	*	*	0.4	*	*	0.4	0.4
2000.....	0.5	*	2.6	0.5	*	*	0.4	*	*	0.4	0.3
1999.....	0.5	*	2.6	0.5	*	*	*	*	0.3	0.4	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes

2007.....	1.1	0.9	1.2	0.7	0.4	0.4	0.5	0.6	0.5	0.6	0.9
2006.....	1.0	0.7	1.2	0.6	0.3	0.4	0.5	0.5	0.5	0.6	0.7
2005.....	1.1	0.8	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.7	0.9
2004.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7	0.7
2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8	1.0
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7	0.9
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9	1.1
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8	1.0
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9	1.0

All origins 3/, male

2007.....	1.3	0.9	1.5	0.7	0.4	0.5	0.7	0.6	0.7	0.6	1.2
2006.....	1.2	*	1.4	0.6	0.3	0.4	0.6	0.7	0.6	0.8	0.9
2005.....	1.3	*	1.5	0.8	0.4	0.4	0.6	0.7	0.6	0.9	1.2
2004.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9	1.0
2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1	1.2
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9	1.3
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2	1.6
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2	1.5

All origins 3/, female

2007.....	0.9	*	0.9	0.6	0.4	0.3	0.4	0.5	0.3	0.5	0.5
2006.....	0.8	*	1.0	0.6	0.3	0.3	0.3	0.3	0.4	0.5	0.5
2005.....	0.9	*	1.0	0.6	0.4	0.2	0.4	0.4	0.5	0.4	0.5
2004.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4	0.5
2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5	0.7
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4	0.6
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5	0.7
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5	0.7
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 4/, both sexes

2007.....	1.2	1.0	1.3	0.7	0.4	0.4	0.5	0.6	0.6	0.6	0.9
2006.....	1.1	0.8	1.3	0.6	0.3	0.4	0.5	0.5	0.5	0.7	0.8
2005.....	1.2	0.8	1.4	0.8	0.5	0.3	0.5	0.6	0.6	0.7	0.9
2004.....	1.2	0.7	1.6	1.0	0.5	0.4	0.6	0.7	0.6	0.7	0.8
2003.....	1.3	0.8	1.6	0.7	0.4	0.4	0.7	0.7	0.7	0.8	1.1
2002.....	1.2	1.0	1.5	0.8	0.5	0.4	0.6	0.7	0.8	0.7	1.0
2001.....	1.3	1.3	1.7	0.9	0.5	0.4	0.7	0.7	0.7	0.9	1.1
2000.....	1.3	1.0	2.2	1.0	0.5	0.4	0.7	0.7	0.7	0.9	1.0
1999.....	1.3	1.2	2.1	0.9	0.5	0.4	0.7	0.8	0.7	0.9	1.0

Non-Hispanic 4/, male

2007.....	1.4	*	1.6	0.8	0.5	0.5	0.7	0.7	0.8	0.7	1.3
2006.....	1.3	*	1.5	0.6	0.3	0.4	0.6	0.7	0.6	0.8	1.0
2005.....	1.4	*	1.6	0.9	0.5	0.4	0.6	0.8	0.7	1.0	1.2
2004.....	1.4	*	1.7	1.3	0.5	0.4	0.7	1.0	0.8	1.0	1.1
2003.....	1.5	*	1.9	0.8	0.4	0.4	0.8	0.8	0.9	1.2	1.3
2002.....	1.5	*	1.7	1.0	0.6	0.6	0.7	0.9	1.0	0.9	1.4
2001.....	1.6	1.5	2.0	1.1	0.5	0.5	1.0	0.8	0.9	1.3	1.6
2000.....	1.6	*	2.6	1.2	0.4	0.4	0.9	0.9	0.9	1.3	1.2
1999.....	1.6	*	2.3	1.0	0.6	0.6	0.8	1.1	1.1	1.3	1.5

Non-Hispanic 4/, female

2007.....	1.0	*	1.0	0.6	0.4	0.3	0.4	0.5	0.4	0.5	0.6
2006.....	0.9	*	1.2	0.7	0.4	0.3	0.4	0.4	0.4	0.5	0.6
2005.....	0.9	*	1.1	0.7	0.4	0.2	0.5	0.5	0.5	0.5	0.5
2004.....	1.0	*	1.5	0.7	0.5	0.3	0.6	0.4	0.4	0.5	0.5
2003.....	1.0	1.3	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.5	0.8
2002.....	0.9	*	1.2	0.7	0.4	0.3	0.4	0.4	0.6	0.5	0.6
2001.....	1.0	*	1.3	0.6	0.4	0.4	0.4	0.5	0.5	0.6	0.7
2000.....	1.0	*	1.7	0.8	0.5	0.3	0.5	0.5	0.5	0.5	0.7
1999.....	1.0	*	1.8	0.8	0.4	0.3	0.5	0.6	0.4	0.6	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic White, both sexes

2007 5/.....	1.1	*	0.8	0.5	0.3	0.4	0.6	0.5	0.6	0.5	0.8
2006 5/.....	1.1	*	0.9	0.5	0.2	0.4	0.5	0.5	0.5	0.6	0.8
2005 5/.....	1.1	*	1.0	0.6	0.4	0.3	0.6	0.6	0.5	0.7	0.9
2004 5/.....	1.1	*	1.3	0.7	0.4	0.4	0.6	0.6	0.6	0.7	0.7
2003 5/.....	1.2	*	1.3	0.5	0.3	0.4	0.7	0.7	0.7	0.9	1.0
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.6	0.8	0.7	1.0
2001.....	1.1	1.1	1.3	0.7	0.4	0.4	0.7	0.5	0.7	0.8	1.0
2000.....	1.1	*	1.5	0.7	0.4	0.4	0.6	0.6	0.6	0.8	0.9
1999.....	1.1	0.9	1.5	0.6	0.4	0.4	0.6	0.8	0.7	0.8	1.0

Non-Hispanic White, male

2007 5/.....	1.3	*	1.1	0.5	0.3	0.5	0.7	0.7	0.8	0.6	1.1
2006 5/.....	1.3	*	1.2	0.4	*	0.5	0.6	0.6	0.6	0.7	0.9
2005 5/.....	1.3	*	1.4	0.7	0.4	0.4	0.6	0.7	0.7	1.0	1.1
2004 5/.....	1.3	*	1.4	0.8	0.4	0.4	0.7	0.9	0.8	0.9	0.9
2003 5/.....	1.4	*	1.6	0.5	*	0.4	1.0	0.8	0.9	1.2	1.4
2002.....	1.3	*	1.1	0.6	0.5	0.5	0.7	0.8	1.0	0.8	1.4
2001.....	1.4	*	1.7	0.9	0.4	0.5	1.0	0.8	0.9	1.1	1.5
2000.....	1.4	*	1.8	0.8	0.3	0.4	0.9	0.7	0.8	1.1	1.1
1999.....	1.4	*	1.7	0.6	0.5	0.6	0.8	1.1	1.1	1.1	1.4

Non-Hispanic White, female

2007 5/.....	0.9	*	0.5	0.5	*	0.3	0.4	0.4	0.3	0.5	0.6
2006 5/.....	0.9	*	0.7	0.5	*	0.3	*	0.3	0.4	0.5	0.6
2005 5/.....	0.9	*	0.7	0.4	*	*	0.5	0.5	0.4	0.5	0.6
2004 5/.....	0.9	*	1.2	0.5	0.3	*	0.5	0.4	0.4	0.6	0.5
2003 5/.....	1.0	*	1.0	0.5	*	*	0.5	0.6	0.6	0.5	0.7
2002.....	0.8	*	0.8	0.4	*	*	0.3	*	0.6	0.5	0.6
2001.....	0.9	*	0.9	0.5	0.4	0.3	0.3	*	0.5	0.5	0.6
2000.....	0.9	*	1.1	0.6	0.4	*	0.4	0.4	0.4	0.4	0.6
1999.....	0.8	*	1.3	0.6	0.3	*	0.4	0.5	0.3	0.5	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic Black, both sexes

2007 5/.....	2.0	*	3.2	1.7	1.1	*	*	1.0	0.9	1.1	1.6
2006 5/.....	1.8	*	3.1	1.6	*	*	0.7	0.9	0.9	1.1	1.0
2005 5/.....	2.0	*	2.8	1.7	0.9	*	*	0.9	1.0	1.1	1.2
2004 5/.....	2.1	*	2.9	2.4	1.0	*	0.8	1.2	0.8	0.9	1.4
2003 5/.....	2.1	*	3.1	2.0	1.0	*	*	1.0	0.8	1.0	1.4
2002.....	2.1	*	3.7	2.1	0.9	0.7	0.9	1.2	1.1	1.0	1.1
2001.....	2.3	*	3.4	1.8	0.8	*	1.0	1.4	1.0	1.6	2.1
2000.....	2.6	*	5.1	2.2	0.8	*	1.0	1.2	1.3	1.6	2.0
1999.....	2.6	*	4.7	2.2	0.9	*	0.9	1.1	0.9	1.9	1.7

Non-Hispanic Black, male

2007 5/.....	2.5	*	3.5	2.1	*	*	*	*	*	*	2.5
2006 5/.....	2.1	*	3.1	1.5	*	*	*	*	*	*	*
2005 5/.....	2.4	*	2.8	1.8	*	*	*	*	*	*	2.0
2004 5/.....	2.7	*	3.0	3.3	*	*	*	1.6	*	*	2.0
2003 5/.....	2.5	*	3.4	2.0	*	*	*	*	*	*	1.6
2002.....	2.6	*	4.3	2.3	*	*	*	*	*	1.6	1.8
2001.....	3.0	*	3.7	2.3	*	*	*	*	*	2.6	3.0
2000.....	3.2	*	5.8	2.7	*	*	*	*	1.6	2.3	2.6
1999.....	3.1	*	5.3	2.5	*	*	*	*	*	2.8	2.5

Non-Hispanic Black, female

2007 5/.....	1.6	*	2.8	1.3	*	*	*	*	*	*	*
2006 5/.....	1.6	*	3.1	1.6	*	*	*	*	*	*	*
2005 5/.....	1.5	*	2.7	1.7	*	*	*	*	*	*	*
2004 5/.....	1.7	*	2.9	1.5	*	*	*	*	*	*	*
2003 5/.....	1.8	*	2.8	1.9	*	*	*	*	*	*	*
2002.....	1.6	*	3.0	1.8	*	*	*	*	*	*	*
2001.....	1.8	*	3.1	1.3	*	*	*	1.5	*	*	*
2000.....	2.0	*	4.3	1.7	*	*	*	*	*	*	1.4
1999.....	2.1	*	4.1	1.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes

2007.....	9.9	*	0.2	*	0.3	3.9	11.0	13.9	14.2	15.8	19.2
2006.....	9.2	*	0.2	*	0.2	3.5	10.4	13.0	13.0	15.1	19.3
2005.....	8.0	0.5	0.1	*	0.2	3.0	8.8	10.6	11.3	13.5	17.1
2004.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8	16.3
2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5	15.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9	14.8
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4	12.0
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5	11.2
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1	11.1

All origins 3/, male

2007.....	13.2	*	*	*	0.4	5.8	16.5	20.3	20.0	21.0	24.8
2006.....	12.6	*	*	*	0.2	5.2	16.0	19.2	18.6	20.5	25.4
2005.....	10.9	*	*	*	0.2	4.5	13.4	15.5	15.7	18.0	22.1
2004.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8	20.9
2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8	20.7
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5	20.3
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6	16.8
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7	16.4
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3	16.6

All origins 3/, female

2007.....	6.7	*	*	*	0.3	1.9	5.2	7.1	8.3	10.6	13.8
2006.....	5.9	*	*	*	*	1.6	4.4	6.5	7.3	9.6	13.4
2005.....	5.1	*	*	*	*	1.5	3.9	5.4	6.7	8.9	12.0
2004.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7	11.7
2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1	10.3
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2	9.4
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1	7.3
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3	6.0
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9	5.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental poisoning and exposure to noxious substances (X40-X49)

Hispanic, both sexes

2007.....	5.4	*	*	*	*	2.7	5.6	6.6	7.1	9.1	11.3
2006.....	5.2	*	*	*	*	2.0	4.6	5.8	8.1	9.0	12.5
2005.....	4.7	*	*	*	*	1.8	4.8	5.4	6.9	8.8	11.2
2004.....	4.2	*	*	*	*	1.7	3.9	4.8	5.0	8.3	10.2
2003.....	4.5	*	*	*	*	2.1	4.0	5.2	6.3	9.7	11.1
2002.....	4.3	*	*	*	*	1.8	3.8	4.3	5.7	8.3	12.2
2001.....	3.6	*	*	*	*	1.5	2.7	4.3	5.5	8.4	9.1
2000.....	3.7	*	*	*	*	1.2	2.8	3.9	6.8	9.5	9.8
1999.....	4.4	*	*	*	*	1.3	3.2	4.8	7.5	11.4	12.4

Hispanic, male

2007.....	8.3	*	*	*	*	3.8	9.0	10.1	10.4	13.7	17.4
2006.....	7.8	*	*	*	*	2.8	6.7	8.5	12.2	13.5	18.6
2005.....	7.3	*	*	*	*	2.6	7.4	8.2	10.5	13.5	16.6
2004.....	6.2	*	*	*	*	2.4	5.6	7.6	7.6	12.0	15.3
2003.....	6.9	*	*	*	*	3.3	6.0	7.7	9.7	14.9	16.3
2002.....	6.6	*	*	*	*	2.6	5.9	6.8	8.9	12.4	19.0
2001.....	5.6	*	*	*	*	2.1	3.9	6.8	8.1	13.1	14.2
2000.....	6.0	*	*	*	*	1.8	4.4	6.1	11.0	14.5	17.1
1999.....	7.3	*	*	*	*	1.7	5.3	7.6	12.2	18.4	20.5

Hispanic, female

2007.....	2.2	*	*	*	*	1.5	1.5	2.1	3.0	4.0	4.5
2006.....	2.4	*	*	*	*	*	2.0	2.4	3.3	3.9	5.8
2005.....	2.0	*	*	*	*	*	1.7	2.0	2.8	3.4	5.3
2004.....	1.9	*	*	*	*	*	1.8	1.4	1.9	4.1	4.7
2003.....	1.9	*	*	*	*	*	1.6	2.1	2.4	3.8	5.4
2002.....	1.7	*	*	*	*	*	1.3	*	2.0	3.7	5.0
2001.....	1.5	*	*	*	*	*	1.3	1.4	2.5	3.2	3.7
2000.....	1.3	*	*	*	*	*	*	1.3	1.9	3.9	2.2
1999.....	1.4	*	*	*	*	*	*	1.5	2.0	3.8	4.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 4/, both sexes

2007.....	10.7	*	0.2	*	0.4	4.2	12.2	15.6	16.1	17.1	20.5
2006.....	9.8	*	0.2	*	0.2	3.8	11.7	14.8	14.2	16.2	20.4
2005.....	8.5	*	*	*	0.2	3.3	9.6	11.8	12.2	14.3	17.9
2004.....	7.6	*	*	*	0.2	3.4	8.5	9.9	10.2	13.6	17.1
2003.....	7.0	*	0.3	*	0.2	2.6	7.8	8.5	9.8	12.9	16.0
2002.....	6.3	0.6	0.2	*	0.1	2.5	6.3	7.5	9.2	12.4	15.1
2001.....	5.1	*	0.2	*	0.1	2.1	5.3	6.0	7.1	10.6	12.2
2000.....	4.6	*	0.2	*	0.1	1.8	4.6	5.3	6.8	9.4	11.2
1999.....	4.3	*	0.2	*	0.1	1.3	3.9	4.8	6.6	8.7	10.8

Non-Hispanic 4/, male

2007.....	14.1	*	*	*	0.5	6.2	18.1	23.0	22.7	22.5	26.0
2006.....	13.4	*	*	*	0.3	5.7	18.1	22.2	20.3	21.9	26.4
2005.....	11.5	*	*	*	*	4.9	14.7	17.5	17.0	18.9	22.9
2004.....	10.2	*	*	*	0.3	5.0	13.2	14.8	14.2	17.7	21.6
2003.....	9.5	*	0.3	*	0.3	3.9	12.1	12.7	13.8	17.0	21.2
2002.....	8.8	*	*	*	*	3.7	9.8	11.3	13.1	17.1	20.3
2001.....	7.2	*	*	*	*	3.1	8.5	9.1	10.1	14.7	17.0
2000.....	6.6	*	*	*	*	2.8	7.2	8.3	9.8	13.5	16.2
1999.....	6.3	*	*	*	*	1.9	6.0	7.6	9.9	12.5	15.9

Non-Hispanic 4/, female

2007.....	7.4	*	*	*	0.3	2.0	6.0	8.2	9.5	11.8	15.2
2006.....	6.4	*	*	*	*	1.7	4.9	7.4	8.1	10.7	14.5
2005.....	5.6	*	*	*	*	1.5	4.3	6.1	7.5	9.8	12.9
2004.....	5.1	*	*	*	*	1.6	3.7	5.0	6.1	9.5	12.6
2003.....	4.6	*	*	*	*	1.3	3.4	4.4	5.9	8.8	10.9
2002.....	4.0	*	*	*	*	1.3	2.8	3.8	5.2	7.7	9.9
2001.....	3.1	*	*	*	*	1.0	2.1	2.9	4.1	6.5	7.6
2000.....	2.7	*	*	*	*	0.8	1.9	2.2	3.8	5.5	6.3
1999.....	2.4	*	*	*	*	0.6	1.8	2.1	3.4	5.0	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic White, both sexes

2007 5/.....	11.7	*	*	*	0.3	5.1	14.8	19.1	19.2	19.5	22.8
2006 5/.....	10.5	*	*	*	0.2	4.7	14.3	17.9	16.5	17.8	21.6
2005 5/.....	9.0	*	*	*	0.2	4.0	11.8	14.1	14.0	15.8	19.1
2004 5/.....	8.2	*	*	*	0.2	4.2	10.4	12.0	11.7	14.9	18.4
2003 5/.....	7.4	*	*	*	0.2	3.2	9.7	10.1	11.1	13.8	16.9
2002.....	6.6	*	*	*	0.2	3.0	7.6	8.8	10.3	13.1	15.7
2001.....	5.2	*	*	*	*	2.4	6.2	6.9	7.7	11.0	12.3
2000.....	4.6	*	*	*	*	2.2	5.4	5.9	7.2	9.7	11.0
1999.....	4.2	*	*	*	*	1.5	4.3	5.3	6.7	8.8	10.3

Non-Hispanic White, male

2007 5/.....	15.4	*	*	*	0.4	7.8	22.2	28.1	27.0	25.4	28.8
2006 5/.....	14.2	*	*	*	*	7.2	22.3	26.6	23.2	23.7	27.7
2005 5/.....	12.2	*	*	*	*	6.1	18.1	20.8	19.4	20.8	24.3
2004 5/.....	10.9	*	*	*	*	6.3	16.1	17.7	16.4	19.5	23.1
2003 5/.....	10.1	*	*	*	*	4.8	14.9	15.1	15.4	18.0	22.4
2002.....	9.1	*	*	*	*	4.5	11.8	13.0	14.7	18.1	21.0
2001.....	7.3	*	*	*	*	3.8	9.9	10.3	11.2	15.3	17.2
2000.....	6.6	*	*	*	*	3.3	8.6	9.2	10.4	14.0	15.8
1999.....	6.1	*	*	*	*	2.2	6.6	8.3	10.0	12.6	15.3

Non-Hispanic White, female

2007 5/.....	8.1	*	*	*	*	2.3	7.1	10.0	11.2	13.6	16.8
2006 5/.....	7.0	*	*	*	*	2.1	5.9	9.2	9.6	11.9	15.5
2005 5/.....	6.1	*	*	*	*	1.8	5.1	7.3	8.5	10.7	13.8
2004 5/.....	5.5	*	*	*	*	1.9	4.5	6.1	6.9	10.2	13.7
2003 5/.....	4.9	*	*	*	*	1.5	4.3	5.1	6.6	9.5	11.5
2002.....	4.3	*	*	*	*	1.5	3.4	4.5	5.8	8.2	10.4
2001.....	3.1	*	*	*	*	1.0	2.5	3.4	4.2	6.7	7.4
2000.....	2.7	*	*	*	*	1.0	2.0	2.5	3.9	5.4	6.2
1999.....	2.4	*	*	*	*	0.7	2.0	2.3	3.3	4.9	5.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic Black, both sexes

2007 5/.....	8.4	*	*	*	*	1.0	3.7	5.8	8.7	12.4	15.5
2006 5/.....	9.2	*	*	*	*	0.8	3.7	7.0	9.6	15.0	20.7
2005 5/.....	7.8	*	*	*	*	0.7	3.0	5.8	8.9	11.9	17.2
2004 5/.....	6.7	*	*	*	*	0.8	2.9	4.6	6.3	11.5	15.1
2003 5/.....	6.4	*	*	*	*	*	2.3	4.1	7.5	12.1	15.7
2002.....	6.4	*	*	*	*	*	2.3	4.5	7.0	12.1	16.1
2001.....	6.0	*	*	*	*	0.9	2.4	4.1	6.4	11.2	15.6
2000.....	5.7	*	*	*	*	*	2.2	4.3	7.0	10.9	15.6
1999.....	5.8	*	*	*	*	*	2.9	4.3	8.2	10.4	16.3

Non-Hispanic Black, male

2007 5/.....	11.6	*	*	*	*	1.5	4.9	8.4	12.6	16.8	20.1
2006 5/.....	13.3	*	*	*	*	1.2	5.1	11.4	14.9	21.2	28.3
2005 5/.....	11.0	*	*	*	*	*	4.2	8.8	12.1	15.7	22.1
2004 5/.....	9.1	*	*	*	*	*	4.3	6.8	8.8	14.1	19.8
2003 5/.....	9.1	*	*	*	*	*	3.7	5.8	10.7	17.1	21.2
2002.....	9.3	*	*	*	*	*	3.7	7.2	10.1	17.3	22.2
2001.....	8.6	*	*	*	*	*	3.9	6.8	8.4	15.5	20.9
2000.....	8.4	*	*	*	*	*	2.8	7.1	9.8	15.0	23.2
1999.....	8.8	*	*	*	*	*	4.8	6.9	11.7	14.9	24.4

Non-Hispanic Black, female

2007 5/.....	5.6	*	*	*	*	*	2.5	3.3	5.1	8.5	11.4
2006 5/.....	5.4	*	*	*	*	*	2.2	2.8	4.9	9.5	14.1
2005 5/.....	4.9	*	*	*	*	*	1.7	2.9	6.1	8.5	12.9
2004 5/.....	4.4	*	*	*	*	*	1.5	2.7	4.1	9.2	11.0
2003 5/.....	4.1	*	*	*	*	*	*	2.5	4.7	7.7	10.9
2002.....	3.8	*	*	*	*	*	*	2.1	4.2	7.5	10.7
2001.....	3.6	*	*	*	*	*	*	1.7	4.7	7.4	10.9
2000.....	3.2	*	*	*	*	*	1.5	1.8	4.4	7.2	8.8
1999.....	3.1	*	*	*	*	*	*	2.0	5.0	6.4	9.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes

2007.....	5.7	23.9	1.6	0.6	0.6	1.1	1.7	1.9	2.1	2.2	2.9
2006.....	5.8	21.5	1.6	0.6	0.6	1.2	1.9	2.1	2.2	2.5	3.3
2005.....	6.3	19.5	1.5	0.6	0.7	1.4	2.0	2.3	2.4	2.6	3.5
2004.....	5.8	19.2	1.6	0.6	0.6	1.2	1.9	1.9	2.2	2.6	3.2
2003.....	5.9	16.9	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.6	3.4
2002.....	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6	3.3
2001.....	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9	3.2
2000.....	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0	3.4
1999.....	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7	3.7

All origins 3/, male

2007.....	6.7	26.5	1.9	0.7	0.8	1.8	2.9	3.1	3.4	3.6	4.8
2006.....	7.0	25.2	1.7	0.8	0.8	2.0	3.2	3.5	3.5	4.2	5.4
2005.....	7.4	21.1	1.8	0.8	1.0	2.3	3.3	3.8	4.0	4.2	5.6
2004.....	6.8	21.7	2.0	0.8	1.0	2.0	3.1	3.1	3.7	4.1	5.0
2003.....	6.9	19.5	2.2	0.6	0.9	1.8	3.3	3.3	3.4	4.0	5.5
2002.....	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3	5.4
2001.....	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8	5.3
2000.....	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1	5.6
1999.....	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5	6.1

All origins 3/, female

2007.....	4.6	21.2	1.2	0.5	0.3	0.4	0.5	0.6	0.7	0.8	1.1
2006.....	4.7	17.6	1.5	0.4	0.3	0.4	0.6	0.7	0.8	0.8	1.3
2005.....	5.1	17.9	1.2	0.5	0.3	0.6	0.7	0.7	0.8	0.9	1.5
2004.....	4.8	16.5	1.1	0.4	0.3	0.4	0.5	0.6	0.7	1.0	1.4
2003.....	4.9	14.1	1.3	0.3	0.2	0.3	0.7	0.7	0.6	1.1	1.4
2002.....	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0	1.3
2001.....	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1	1.2
2000.....	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8	1.3
1999.....	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	6.2	28.1	1.7	0.6	0.6	1.1	1.7	1.8	2.1	2.3	2.9
2006.....	6.4	24.2	1.6	0.6	0.6	1.3	1.9	2.2	2.1	2.5	3.4
2005.....	6.8	21.9	1.6	0.7	0.8	1.5	2.0	2.2	2.4	2.6	3.6
2004.....	6.3	21.9	1.7	0.6	0.7	1.2	1.8	1.9	2.2	2.6	3.3
2003.....	6.4	19.2	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.7	3.5
2002.....	6.5	18.9	1.6	0.6	0.8	1.4	1.8	2.0	2.1	2.7	3.4
2001.....	6.7	18.3	1.9	0.6	0.7	1.3	2.0	1.9	2.2	2.9	3.3
2000.....	6.7	16.9	1.8	0.6	0.8	1.4	1.9	2.2	2.3	3.0	3.4
1999.....	7.1	15.0	1.9	0.7	0.9	1.5	2.0	2.2	2.4	2.7	3.9

Non-Hispanic 4/, male

2007.....	7.3	31.0	2.0	0.7	0.9	1.8	2.8	3.1	3.4	3.7	4.7
2006.....	7.5	28.6	1.7	0.8	0.9	2.1	3.1	3.6	3.5	4.2	5.5
2005.....	8.0	23.8	1.9	0.8	1.1	2.3	3.3	3.7	3.9	4.3	5.7
2004.....	7.4	24.6	2.1	0.8	1.1	2.0	3.0	3.3	3.8	4.1	5.2
2003.....	7.5	22.2	2.3	0.6	1.0	1.8	3.2	3.3	3.4	4.2	5.6
2002.....	7.5	21.0	2.0	0.7	1.2	2.1	3.1	3.3	3.6	4.3	5.4
2001.....	7.7	21.1	2.1	0.8	1.1	1.9	3.4	3.2	3.8	4.8	5.3
2000.....	7.7	19.0	2.2	0.8	1.3	2.2	3.4	3.8	3.8	5.1	5.6
1999.....	8.1	17.9	2.3	0.9	1.4	2.4	3.5	3.8	4.0	4.5	6.3

Non-Hispanic 4/, female

2007.....	5.1	25.0	1.3	0.5	0.3	0.4	0.5	0.6	0.8	0.9	1.2
2006.....	5.2	19.6	1.5	0.4	0.3	0.4	0.6	0.7	0.8	0.9	1.4
2005.....	5.7	19.9	1.3	0.5	0.4	0.6	0.7	0.7	0.9	1.0	1.6
2004.....	5.3	19.0	1.2	0.4	0.3	0.4	0.5	0.6	0.7	1.1	1.5
2003.....	5.4	16.0	1.3	0.3	0.3	0.3	0.8	0.7	0.7	1.2	1.4
2002.....	5.6	16.8	1.3	0.5	0.4	0.5	0.5	0.8	0.7	1.1	1.4
2001.....	5.8	15.4	1.6	0.4	0.3	0.6	0.6	0.6	0.6	1.2	1.3
2000.....	5.7	14.7	1.4	0.4	0.3	0.6	0.5	0.6	0.8	0.9	1.3
1999.....	6.2	12.0	1.5	0.5	0.5	0.5	0.5	0.7	0.8	1.0	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic White, both sexes

2007 5/.....	6.7	22.1	1.7	0.6	0.6	1.3	1.9	2.0	2.3	2.4	2.8
2006 5/.....	6.9	18.9	1.5	0.6	0.6	1.3	2.1	2.4	2.3	2.7	3.5
2005 5/.....	7.2	18.6	1.6	0.7	0.8	1.5	2.1	2.5	2.5	2.7	3.6
2004 5/.....	6.9	18.4	1.6	0.6	0.7	1.4	1.9	2.2	2.3	2.6	3.5
2003 5/.....	6.9	15.5	1.7	0.5	0.7	1.1	2.1	2.1	2.2	2.7	3.6
2002.....	7.0	14.5	1.5	0.5	0.8	1.5	2.0	2.3	2.3	2.7	3.5
2001.....	7.2	14.5	1.7	0.7	0.7	1.4	2.2	2.0	2.2	3.1	3.2
2000.....	7.1	13.7	1.7	0.6	0.8	1.4	2.0	2.3	2.4	3.0	3.4
1999.....	7.6	12.1	1.7	0.8	1.0	1.6	2.2	2.3	2.4	2.9	3.9

Non-Hispanic White, male

2007 5/.....	7.7	25.0	2.1	0.7	0.9	2.1	3.2	3.3	3.7	3.9	4.3
2006 5/.....	8.0	22.0	1.7	0.8	1.0	2.1	3.6	4.0	3.7	4.3	5.5
2005 5/.....	8.4	20.1	2.0	0.8	1.2	2.4	3.4	4.1	4.1	4.4	5.7
2004 5/.....	7.9	22.0	2.2	0.8	1.1	2.2	3.3	3.6	3.9	4.2	5.4
2003 5/.....	7.9	18.0	2.2	0.6	1.1	1.9	3.5	3.5	3.6	4.2	5.8
2002.....	7.9	17.0	1.8	0.7	1.1	2.4	3.5	3.7	3.8	4.3	5.7
2001.....	8.1	16.8	2.0	0.9	1.1	2.1	3.7	3.3	3.8	5.1	5.2
2000.....	8.0	16.0	2.1	0.8	1.3	2.3	3.5	4.0	4.0	5.2	5.5
1999.....	8.4	14.2	2.0	1.0	1.4	2.6	3.7	4.1	4.0	4.8	6.3

Non-Hispanic White, female

2007 5/.....	5.6	19.0	1.3	0.5	*	0.4	0.5	0.6	0.8	0.9	1.2
2006 5/.....	5.8	15.6	1.4	0.4	*	0.4	0.6	0.7	0.8	1.0	1.4
2005 5/.....	6.2	17.0	1.2	0.5	0.4	0.6	0.7	0.8	0.8	0.9	1.5
2004 5/.....	5.9	14.7	0.9	0.5	*	0.4	0.5	0.7	0.8	1.1	1.6
2003 5/.....	6.0	12.9	1.2	*	*	0.4	0.7	0.7	0.8	1.3	1.4
2002.....	6.1	11.9	1.2	0.4	0.5	0.5	0.5	0.9	0.7	1.1	1.2
2001.....	6.4	12.1	1.5	0.5	0.3	0.7	0.7	0.6	0.7	1.1	1.2
2000.....	6.3	11.2	1.3	0.4	*	0.5	0.5	0.7	0.7	0.9	1.3
1999.....	6.8	9.8	1.5	0.5	0.5	0.6	0.6	0.6	0.8	1.0	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes

2007.....	11.5	*	0.9	6.9	12.6	12.7	13.3	14.3	16.8
2006.....	11.1	*	1.0	7.3	12.5	12.4	12.3	14.7	15.5
2005.....	11.0	*	1.3	7.7	12.4	11.8	13.0	13.8	16.0
2004.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0	16.0
2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8	15.9
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3	16.1
2001 6/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3	15.2
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7	15.4
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4	14.3

All origins 3/, male

2007.....	18.3	*	1.2	11.1	20.8	20.3	21.1	22.1	25.5
2006.....	17.8	*	1.4	11.5	20.8	19.7	19.7	22.7	23.6
2005.....	17.7	*	1.9	12.1	20.2	19.2	20.6	21.4	24.6
2004.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7	24.2
2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9	24.5
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7	24.7
2001 6/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7	23.6
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7	23.9
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7	22.1

All origins 3/, female

2007.....	4.8	*	0.5	2.5	3.9	4.7	5.3	6.3	8.2
2006.....	4.6	*	0.6	2.8	3.6	4.7	4.7	6.6	7.4
2005.....	4.5	*	0.7	3.0	4.0	4.1	5.3	6.1	7.4
2004.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3	7.9
2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7	7.4
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0	7.7
2001 6/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9	6.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8	7.0
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1	6.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Hispanic, both sexes

2007.....	5.4	*	*	5.7	8.5	8.2	7.7	6.7	7.9
2006.....	4.9	*	0.6	5.5	9.2	7.6	6.6	6.4	6.9
2005.....	5.1	*	1.0	6.1	9.2	7.0	6.6	7.2	7.6
2004.....	5.3	*	1.0	6.4	9.5	7.6	6.7	7.0	6.6
2003.....	5.0	*	1.2	5.9	8.0	6.6	6.9	6.4	7.3
2002.....	5.0	*	1.0	5.7	7.6	6.6	6.6	7.0	7.0
2001 6/.....	5.0	*	1.1	5.2	7.0	7.3	7.0	7.5	7.0
2000.....	5.1	*	0.9	5.6	7.9	6.7	6.2	6.9	8.3
1999.....	5.0	*	1.0	5.7	6.7	6.7	6.7	8.4	6.4

Hispanic, male

2007.....	8.8	*	*	9.4	13.6	12.0	12.3	10.4	12.7
2006.....	7.9	*	*	7.9	15.0	11.5	10.3	10.2	11.0
2005.....	8.3	*	1.4	9.5	14.3	11.3	10.1	11.3	12.4
2004.....	8.6	*	1.2	9.9	15.3	11.9	10.5	11.1	10.5
2003.....	8.3	*	1.7	9.2	12.8	10.5	11.1	10.3	11.8
2002.....	8.3	*	1.1	9.1	11.9	10.8	10.6	11.3	11.1
2001 6/.....	8.3	*	1.5	7.8	11.0	11.6	11.1	12.8	11.8
2000.....	8.4	*	1.2	8.5	13.0	11.1	10.0	10.7	13.7
1999.....	8.2	*	1.4	8.9	10.8	10.6	10.9	13.2	10.6

Hispanic, female

2007.....	1.8	*	*	1.9	2.5	3.4	2.1	2.5	2.6
2006.....	1.7	*	*	2.9	2.3	2.7	2.2	2.1	2.3
2005.....	1.7	*	*	2.4	3.0	1.7	2.5	2.5	2.3
2004.....	1.8	*	*	2.7	2.4	2.2	2.3	2.4	2.5
2003.....	1.5	*	*	2.4	2.0	1.8	2.1	2.0	2.3
2002.....	1.6	*	*	2.0	2.3	1.5	1.9	2.3	2.6
2001 6/.....	1.5	*	*	2.5	2.2	2.2	2.3	1.7	1.9
2000.....	1.5	*	*	2.4	1.6	1.5	1.8	2.7	2.6
1999.....	1.6	*	*	2.0	1.6	2.2	1.8	3.1	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007	12.5	*	1.0	7.1	13.5	13.8	14.8	15.8	18.3
2006	12.2	*	1.1	7.6	13.1	13.6	13.7	16.3	16.8
2005	12.0	*	1.4	8.0	13.0	13.0	14.5	15.0	17.2
2004	11.9	*	1.4	8.5	13.1	13.5	14.3	15.2	17.3
2003	11.7	*	1.1	7.5	12.9	13.1	14.8	15.0	17.0
2002	11.8	*	1.3	7.7	13.3	14.2	13.7	15.5	17.3
2001 6/	11.6	*	1.3	8.4	13.0	13.8	14.1	15.3	16.1
2000	11.1	*	1.6	8.4	13.5	13.1	12.9	14.6	16.1
1999	11.2	*	1.2	8.4	13.5	13.3	14.2	15.1	15.1

Non-Hispanic 4/, male

2007	20.1	*	1.4	11.4	22.4	22.6	23.5	24.7	27.7
2006	19.6	*	1.6	12.3	22.0	22.0	22.2	25.4	25.7
2005	19.3	*	2.0	12.5	21.5	21.4	23.2	23.4	26.5
2004	19.2	*	1.8	13.1	22.1	22.4	22.9	23.6	26.1
2003	19.1	*	1.7	12.1	21.9	21.8	23.9	23.9	26.2
2002	19.4	*	1.9	12.7	22.8	23.4	22.2	24.6	26.5
2001 6/	19.0	*	2.0	13.8	22.7	23.3	22.9	24.2	24.9
2000	18.3	*	2.4	13.8	23.3	21.9	20.9	23.2	24.9
1999	18.4	*	1.9	13.8	23.1	22.3	23.0	24.0	23.2

Non-Hispanic 4/, female

2007	5.3	*	0.6	2.6	4.2	5.0	6.1	7.0	9.1
2006	5.1	*	0.7	2.8	3.8	5.1	5.3	7.4	8.1
2005	4.9	*	0.7	3.1	4.1	4.6	5.9	6.7	8.1
2004	5.0	*	1.0	3.7	3.8	4.6	5.9	6.9	8.6
2003	4.6	*	0.5	2.7	3.6	4.4	5.8	6.3	8.0
2002	4.6	*	0.6	2.4	3.7	5.1	5.2	6.5	8.3
2001 6/	4.5	*	0.6	2.7	3.3	4.4	5.3	6.5	7.5
2000	4.3	*	0.6	2.8	3.5	4.4	5.0	6.1	7.5
1999	4.3	*	0.5	2.8	3.7	4.4	5.6	6.4	7.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic White, both sexes

2007 5/.....	14.4	*	0.9	7.8	14.5	15.4	16.9	18.1	21.2
2006 5/.....	13.9	*	1.1	8.3	14.1	15.1	15.5	18.8	19.4
2005 5/.....	13.6	*	1.3	8.8	14.0	14.1	16.5	17.2	19.7
2004 5/.....	13.5	*	1.4	9.2	14.1	14.6	16.1	17.4	19.9
2003 5/.....	13.3	*	1.1	8.3	13.6	14.3	16.6	17.0	19.4
2002.....	13.4	*	1.3	8.6	14.5	15.5	15.3	17.5	19.7
2001 6/.....	13.0	*	1.3	9.3	13.9	15.1	15.7	17.2	18.2
2000.....	12.5	*	1.7	9.0	14.3	14.0	14.4	16.4	18.0
1999.....	12.4	*	1.3	9.0	14.2	14.2	15.6	16.7	17.0

Non-Hispanic White, male

2007 5/.....	22.9	*	1.2	12.5	24.0	25.1	26.6	28.0	31.7
2006 5/.....	22.3	*	1.6	13.3	23.7	24.2	24.7	28.7	29.2
2005 5/.....	22.0	*	2.0	14.0	23.0	23.2	25.9	26.6	29.9
2004 5/.....	21.6	*	1.8	14.2	23.8	24.1	25.6	26.9	29.7
2003 5/.....	21.6	*	1.7	13.3	23.3	23.4	26.4	26.6	29.7
2002.....	21.9	*	2.0	14.2	24.7	24.7	24.4	27.3	30.0
2001 6/.....	21.4	*	2.1	15.4	24.1	25.3	25.2	26.8	27.9
2000.....	20.4	*	2.6	14.8	24.8	23.2	22.9	25.7	27.6
1999.....	20.3	*	2.1	14.7	24.3	23.5	25.0	25.9	26.0

Non-Hispanic White, female

2007 5/.....	6.1	*	0.5	2.9	4.5	5.5	7.0	8.1	10.7
2006 5/.....	5.9	*	0.7	3.0	4.0	6.0	6.2	8.9	9.6
2005 5/.....	5.6	*	0.6	3.3	4.6	5.0	6.9	7.7	9.5
2004 5/.....	5.7	*	1.0	3.9	4.1	4.8	6.5	7.9	10.1
2003 5/.....	5.3	*	0.5	3.0	3.6	5.0	6.6	7.3	9.2
2002.....	5.3	*	0.6	2.7	4.1	6.1	6.0	7.6	9.5
2001 6/.....	5.0	*	0.6	3.0	3.6	4.8	6.1	7.6	8.5
2000.....	4.9	*	0.7	3.0	3.7	4.7	5.7	7.1	8.5
1999.....	4.9	*	0.5	3.0	3.9	4.8	6.2	7.5	8.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic Black, both sexes

2007 5/.....	5.0	*	1.2	4.1	8.4	8.5	9.0	8.3	7.1
2006 5/.....	5.1	*	1.1	4.3	8.7	8.9	8.5	8.5	6.5
2005 5/.....	5.2	*	1.2	4.5	9.4	9.2	8.7	7.5	7.7
2004 5/.....	5.3	*	1.4	4.8	10.0	10.6	8.9	6.9	6.5
2003 5/.....	5.3	*	1.2	3.9	10.8	9.8	9.4	7.7	7.1
2002.....	5.2	*	1.0	4.0	9.4	10.8	8.5	8.3	7.2
2001 6/.....	5.3	*	1.2	4.5	10.5	9.7	8.7	8.1	6.9
2000.....	5.5	*	1.2	5.7	11.4	10.2	7.9	7.2	8.2
1999.....	5.5	*	0.9	5.9	11.1	10.0	9.0	8.6	7.0

Non-Hispanic Black, male

2007 5/.....	8.6	*	1.7	6.9	14.6	14.5	16.2	14.0	12.0
2006 5/.....	9.1	*	1.7	7.2	15.0	16.3	16.0	15.4	11.5
2005 5/.....	8.9	*	1.5	7.4	16.7	15.8	15.2	12.5	13.4
2004 5/.....	9.2	*	1.7	7.5	17.5	18.8	14.8	11.6	11.2
2003 5/.....	9.0	*	1.9	6.7	18.5	18.1	16.2	13.4	11.6
2002.....	9.3	*	1.6	6.9	16.6	20.5	15.6	14.7	11.7
2001 6/.....	9.3	*	1.7	7.5	20.0	17.7	15.5	14.2	10.8
2000.....	9.6	*	2.1	9.9	20.1	18.2	14.1	12.9	13.7
1999.....	9.9	*	1.4	10.1	20.3	18.7	16.3	16.1	11.1

Non-Hispanic Black, female

2007 5/.....	1.7	*	*	1.3	2.1	2.6	2.5	3.2	2.8
2006 5/.....	1.4	*	*	1.3	2.3	2.0	1.7	2.3	2.2
2005 5/.....	1.8	*	*	1.4	2.0	3.0	2.8	2.9	2.7
2004 5/.....	1.8	*	*	1.9	2.5	3.0	3.6	2.8	2.5
2003 5/.....	1.8	*	*	*	3.2	2.2	3.2	2.6	3.1
2002.....	1.6	*	*	*	2.4	2.1	2.1	2.6	3.3
2001 6/.....	1.7	*	*	*	*	2.4	2.5	2.6	3.4
2000.....	1.7	*	*	1.5	3.1	3.0	2.3	2.2	3.3
1999.....	1.6	*	*	1.6	2.3	2.1	2.6	1.9	3.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes

2007	5.8	*	0.3	2.9	6.0	5.5	5.9	6.0	7.3
2006	5.6	*	0.3	3.3	6.0	5.4	5.2	6.2	6.7
2005	5.7	*	0.4	3.5	5.8	5.6	5.7	5.9	7.0
2004	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9	7.1
2003	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0	7.2
2002	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6	7.4
2001	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4	7.0
2000	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7	7.4
1999	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1	6.7

All origins 3/, male

2007	10.2	*	0.4	5.2	10.6	9.6	10.1	10.1	12.0
2006	10.0	*	0.5	5.8	10.6	9.2	9.2	10.4	11.2
2005	10.2	*	0.7	6.1	10.3	9.7	9.7	10.1	11.8
2004	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8	11.7
2003	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2	12.1
2002	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3	12.6
2001	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8	12.0
2000	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1	12.6
1999	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8	11.2

All origins 3/, female

2007	1.4	*	*	0.6	1.1	1.2	1.6	1.9	2.6
2006	1.4	*	*	0.6	1.2	1.4	1.2	1.9	2.3
2005	1.4	*	*	0.8	1.1	1.3	1.6	1.7	2.3
2004	1.5	*	*	1.0	1.1	1.3	1.7	1.9	2.6
2003	1.4	*	*	0.8	1.2	1.3	1.5	1.8	2.3
2002	1.4	*	*	0.7	1.2	1.6	1.7	2.0	2.3
2001	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1	2.2
2000	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3	2.4
1999	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic 4/, both sexes

2007	6.4	*	0.3	3.0	6.6	6.0	6.7	6.8	8.1
2006	6.3	*	0.3	3.5	6.5	6.1	5.9	7.0	7.4
2005	6.4	*	0.5	3.7	6.2	6.3	6.5	6.6	7.6
2004	6.3	*	0.3	3.9	6.6	6.3	6.5	6.5	7.7
2003	6.4	*	0.4	3.8	7.0	6.5	6.9	6.6	7.8
2002	6.5	*	0.4	3.8	7.3	7.2	6.3	7.3	8.1
2001	6.5	*	0.5	4.5	7.2	7.4	6.9	7.0	7.6
2000	6.4	*	0.6	4.7	7.9	7.0	6.6	7.3	7.9
1999	6.4	*	0.5	5.2	7.9	7.3	7.1	7.5	7.1

Non-Hispanic 4/, male

2007	11.4	*	0.5	5.4	11.7	10.7	11.7	11.5	13.3
2006	11.2	*	0.5	6.2	11.5	10.6	10.5	11.8	12.4
2005	11.4	*	0.8	6.5	11.0	11.1	11.2	11.3	13.0
2004	11.1	*	0.5	6.7	11.9	11.2	11.0	11.0	12.6
2003	11.5	*	0.6	6.6	12.6	11.4	12.0	11.3	13.1
2002	11.7	*	0.7	6.6	13.1	12.7	10.7	12.4	13.7
2001	11.6	*	0.7	7.8	13.2	13.1	12.0	11.8	12.9
2000	11.4	*	0.9	8.2	14.3	12.3	11.2	12.1	13.3
1999	11.5	*	0.8	9.0	14.3	12.8	12.1	12.7	12.0

Non-Hispanic 4/, female

2007	1.6	*	*	0.6	1.3	1.4	1.8	2.1	3.0
2006	1.6	*	*	0.7	1.3	1.5	1.4	2.2	2.6
2005	1.6	*	*	0.8	1.2	1.5	1.8	1.9	2.5
2004	1.7	*	*	1.1	1.2	1.4	2.0	2.1	2.8
2003	1.6	*	*	0.8	1.3	1.5	1.8	2.0	2.6
2002	1.6	*	*	0.8	1.3	1.8	1.9	2.2	2.5
2001	1.6	*	0.2	1.0	1.1	1.7	1.9	2.3	2.4
2000	1.6	*	0.2	1.1	1.4	1.6	2.1	2.6	2.6
1999	1.6	*	*	1.1	1.5	1.7	2.1	2.5	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic White, both sexes

2007 5/.....	7.5	*	0.3	3.3	7.1	6.8	7.8	8.0	9.5
2006 5/.....	7.3	*	0.4	3.8	6.9	6.6	6.7	8.2	8.7
2005 5/.....	7.4	*	0.5	4.2	6.6	7.0	7.4	7.6	8.9
2004 5/.....	7.3	*	0.4	4.4	7.1	6.8	7.3	7.6	8.9
2003 5/.....	7.4	*	0.4	4.3	7.4	7.0	7.7	7.6	9.1
2002.....	7.5	*	0.5	4.1	7.8	7.8	7.2	8.3	9.4
2001.....	7.4	*	0.5	5.0	7.5	8.1	7.9	8.0	8.6
2000.....	7.3	*	0.7	5.0	8.5	7.4	7.5	8.3	9.0
1999.....	7.3	*	0.6	5.6	8.2	7.8	8.0	8.4	8.1

Non-Hispanic White, male

2007 5/.....	13.3	*	0.6	5.7	12.4	12.0	13.3	13.3	15.5
2006 5/.....	13.0	*	0.7	6.7	12.2	11.4	11.8	13.5	14.2
2005 5/.....	13.2	*	0.9	7.2	11.7	12.2	12.5	12.8	14.8
2004 5/.....	12.8	*	0.6	7.5	12.8	12.0	12.2	12.7	14.5
2003 5/.....	13.2	*	0.7	7.4	13.3	12.1	13.3	12.8	15.2
2002.....	13.4	*	0.8	7.2	14.0	13.3	12.0	14.0	15.8
2001.....	13.2	*	0.8	8.6	13.6	14.2	13.4	13.2	14.5
2000.....	12.9	*	1.2	8.6	15.3	12.9	12.5	13.5	15.0
1999.....	12.9	*	1.0	9.7	14.7	13.6	13.3	13.8	13.5

Non-Hispanic White, female

2007 5/.....	1.9	*	*	0.7	1.4	1.5	2.2	2.6	3.6
2006 5/.....	1.9	*	*	0.7	1.4	1.8	1.7	2.8	3.1
2005 5/.....	1.8	*	*	1.0	1.3	1.7	2.2	2.4	3.0
2004 5/.....	1.9	*	*	1.2	1.3	1.6	2.4	2.5	3.4
2003 5/.....	1.8	*	*	0.9	1.3	1.8	2.1	2.4	3.0
2002.....	1.8	*	*	0.9	1.5	2.3	2.3	2.6	2.9
2001.....	1.9	*	*	1.2	1.2	1.9	2.3	2.7	2.7
2000.....	1.9	*	*	1.2	1.5	1.8	2.3	3.0	3.0
1999.....	1.9	*	*	1.3	1.5	2.0	2.5	2.9	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes

2007.....	5.7	*	0.6	4.0	6.6	7.2	7.4	8.2	9.5
2006.....	5.5	*	0.7	4.0	6.4	6.9	7.1	8.5	8.7
2005.....	5.3	*	0.9	4.2	6.5	6.2	7.3	7.8	9.0
2004.....	5.3	*	1.1	4.4	6.2	6.6	7.3	8.1	8.9
2003.....	5.0	*	0.8	3.7	5.6	6.1	7.4	7.8	8.7
2002.....	5.0	*	0.8	3.8	5.7	6.4	6.9	7.7	8.7
2001 6/.....	4.8	*	0.9	3.8	5.4	5.9	6.7	7.9	8.2
2000.....	4.5	*	0.9	3.6	5.3	5.8	6.0	7.0	7.9
1999.....	4.5	*	0.7	3.2	5.1	5.7	6.7	7.3	7.7

All origins 3/, male

2007.....	8.1	*	0.8	5.9	10.2	10.8	11.0	12.0	13.5
2006.....	7.8	*	0.9	5.7	10.2	10.5	10.5	12.3	12.4
2005.....	7.5	*	1.2	6.0	10.0	9.5	10.9	11.3	12.8
2004.....	7.6	*	1.2	6.2	9.7	10.3	11.0	11.9	12.5
2003.....	7.3	*	1.2	5.4	8.8	9.3	11.0	11.7	12.4
2002.....	7.3	*	1.2	5.8	8.9	9.7	10.6	11.4	12.1
2001 6/.....	7.1	*	1.3	5.8	8.7	9.3	10.2	11.9	11.6
2000.....	6.6	*	1.4	5.4	8.5	8.9	9.1	10.6	11.3
1999.....	6.6	*	1.1	4.6	8.0	8.8	10.2	10.9	10.8

All origins 3/, female

2007.....	3.4	*	0.4	1.9	2.8	3.5	3.8	4.4	5.6
2006.....	3.2	*	0.5	2.2	2.4	3.3	3.5	4.7	5.1
2005.....	3.1	*	0.6	2.2	2.9	2.8	3.6	4.4	5.1
2004.....	3.1	*	0.9	2.5	2.4	2.8	3.5	4.4	5.3
2003.....	2.8	*	0.4	1.9	2.2	2.7	3.7	4.0	5.1
2002.....	2.9	*	0.4	1.6	2.3	2.9	3.1	4.0	5.4
2001 6/.....	2.6	*	0.4	1.7	2.0	2.5	3.1	3.9	4.8
2000.....	2.5	*	0.4	1.7	1.9	2.6	2.7	3.4	4.6
1999.....	2.5	*	0.3	1.6	2.0	2.5	3.2	3.7	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Hispanic, both sexes

2007	3.4	*	*	3.3	5.2	5.1	5.0	4.4	5.3
2006	3.1	*	*	3.2	5.3	4.8	4.3	4.2	4.6
2005	3.2	*	0.9	3.8	5.3	4.2	4.2	4.8	4.8
2004	3.2	*	1.0	3.5	5.0	4.5	4.4	4.9	3.9
2003	2.9	*	0.9	3.4	4.1	3.4	4.5	4.0	4.7
2002	2.9	*	0.7	2.9	3.8	3.6	4.2	4.5	4.5
2001 6/	2.8	*	1.0	3.1	3.5	3.7	4.1	5.0	4.6
2000	2.8	*	0.6	2.8	3.9	3.6	3.7	4.6	4.9
1999	2.7	*	*	2.6	2.8	3.6	4.1	5.0	3.8

Hispanic, male

2007	5.2	*	*	5.1	8.0	6.8	7.8	6.6	8.1
2006	4.6	*	*	3.7	8.1	7.1	6.2	6.4	6.9
2005	4.9	*	1.1	5.6	7.7	6.6	6.3	7.1	7.8
2004	4.9	*	1.1	4.8	7.7	7.0	6.7	7.5	5.9
2003	4.5	*	1.3	4.9	6.2	5.1	6.9	6.2	7.3
2002	4.4	*	*	4.2	5.5	5.9	6.4	7.0	7.0
2001 6/	4.4	*	1.3	4.3	5.1	5.5	6.2	8.3	7.4
2000	4.3	*	*	3.8	6.2	5.9	5.9	7.0	7.7
1999	4.2	*	*	4.0	4.3	5.4	6.5	7.9	6.0

Hispanic, female

2007	1.4	*	*	1.4	1.9	3.0	1.7	2.0	2.0
2006	1.4	*	*	2.7	1.8	1.8	1.9	1.8	2.0
2005	1.4	*	*	1.9	2.4	1.2	1.8	2.2	1.5
2004	1.4	*	*	2.2	1.7	1.3	1.8	2.0	1.8
2003	1.2	*	*	1.7	1.5	1.3	1.7	1.5	2.0
2002	1.2	*	*	1.5	1.7	*	1.7	1.7	1.9
2001 6/	1.2	*	*	1.9	1.4	1.4	1.8	*	*
2000	1.1	*	*	1.7	*	*	*	1.9	1.9
1999	1.1	*	*	*	*	1.4	1.4	1.9	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007	6.1	*	0.7	4.1	6.9	7.8	8.0	9.0	10.2
2006	5.9	*	0.8	4.2	6.6	7.5	7.8	9.4	9.4
2005	5.6	*	0.9	4.2	6.8	6.7	8.0	8.4	9.6
2004	5.7	*	1.1	4.5	6.4	7.1	7.9	8.7	9.6
2003	5.3	*	0.8	3.7	5.9	6.7	7.9	8.4	9.2
2002	5.3	*	0.8	3.9	6.1	7.0	7.4	8.2	9.2
2001 6/	5.1	*	0.9	3.9	5.8	6.4	7.1	8.3	8.5
2000	4.8	*	1.0	3.7	5.5	6.2	6.3	7.3	8.2
1999	4.7	*	0.7	3.3	5.6	6.0	7.1	7.6	8.0

Non-Hispanic 4/, male

2007	8.7	*	0.8	6.1	10.7	11.9	11.9	13.2	14.4
2006	8.4	*	1.0	6.1	10.6	11.4	11.7	13.6	13.3
2005	8.0	*	1.2	6.0	10.5	10.3	12.0	12.1	13.5
2004	8.1	*	1.3	6.4	10.2	11.2	11.9	12.6	13.5
2003	7.7	*	1.1	5.4	9.3	10.4	11.8	12.6	13.1
2002	7.7	*	1.2	6.1	9.7	10.7	11.5	12.1	12.7
2001 6/	7.5	*	1.3	6.0	9.5	10.2	10.9	12.4	12.0
2000	6.9	*	1.5	5.6	9.0	9.6	9.7	11.1	11.6
1999	6.9	*	1.1	4.8	8.9	9.4	10.9	11.3	11.3

Non-Hispanic 4/, female

2007	3.7	*	0.5	2.0	3.0	3.6	4.2	4.9	6.2
2006	3.5	*	0.6	2.1	2.5	3.6	3.9	5.2	5.5
2005	3.4	*	0.5	2.3	3.0	3.2	4.0	4.7	5.7
2004	3.4	*	0.9	2.6	2.6	3.1	3.9	4.8	5.8
2003	3.1	*	0.4	1.9	2.3	2.9	4.0	4.3	5.4
2002	3.1	*	0.4	1.6	2.4	3.3	3.3	4.4	5.8
2001 6/	2.9	*	0.4	1.7	2.2	2.7	3.4	4.2	5.1
2000	2.7	*	0.4	1.7	2.1	2.8	2.9	3.6	4.9
1999	2.7	*	0.3	1.7	2.2	2.6	3.4	3.9	4.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic White, both sexes

2007 5/.....	6.9	*	0.5	4.5	7.5	8.6	9.0	10.1	11.6
2006 5/.....	6.6	*	0.8	4.5	7.1	8.5	8.8	10.7	10.7
2005 5/.....	6.2	*	0.8	4.6	7.4	7.2	9.1	9.6	10.7
2004 5/.....	6.3	*	1.0	4.8	7.0	7.7	8.8	9.8	10.9
2003 5/.....	5.9	*	0.7	4.1	6.2	7.3	8.9	9.3	10.3
2002.....	6.0	*	0.9	4.5	6.7	7.6	8.1	9.2	10.4
2001 6/.....	5.6	*	0.8	4.4	6.5	7.0	7.8	9.2	9.6
2000.....	5.2	*	0.9	4.0	5.8	6.6	6.9	8.1	9.1
1999.....	5.2	*	0.7	3.4	6.0	6.4	7.7	8.3	8.9

Non-Hispanic White, male

2007 5/.....	9.6	*	0.7	6.8	11.6	13.1	13.2	14.6	16.2
2006 5/.....	9.3	*	0.9	6.6	11.4	12.8	12.9	15.2	15.0
2005 5/.....	8.8	*	1.1	6.8	11.3	11.0	13.4	13.7	15.0
2004 5/.....	8.9	*	1.2	6.7	11.0	12.2	13.4	14.2	15.2
2003 5/.....	8.4	*	1.0	5.9	10.0	11.4	13.2	13.8	14.4
2002.....	8.5	*	1.2	7.0	10.7	11.4	12.4	13.4	14.2
2001 6/.....	8.2	*	1.2	6.8	10.5	11.0	11.9	13.6	13.4
2000.....	7.5	*	1.4	6.2	9.5	10.3	10.4	12.1	12.6
1999.....	7.4	*	1.1	5.0	9.6	9.9	11.6	12.1	12.6

Non-Hispanic White, female

2007 5/.....	4.2	*	0.4	2.2	3.1	4.0	4.8	5.6	7.1
2006 5/.....	4.0	*	0.6	2.3	2.6	4.1	4.6	6.1	6.4
2005 5/.....	3.7	*	0.5	2.3	3.3	3.3	4.7	5.4	6.5
2004 5/.....	3.7	*	0.8	2.7	2.8	3.2	4.1	5.4	6.7
2003 5/.....	3.4	*	0.3	2.1	2.3	3.2	4.5	4.9	6.2
2002.....	3.5	*	0.5	1.8	2.6	3.8	3.8	5.0	6.6
2001 6/.....	3.2	*	0.4	1.7	2.4	2.9	3.8	4.8	5.8
2000.....	3.0	*	0.4	1.7	2.1	2.9	3.4	4.0	5.5
1999.....	3.0	*	*	1.7	2.3	2.8	3.7	4.5	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic Black, both sexes

2007 5/.....	2.5	*	1.0	1.8	3.3	4.1	4.2	4.6	4.3
2006 5/.....	2.4	*	1.0	1.9	3.3	3.6	4.1	4.9	3.5
2005 5/.....	2.6	*	1.0	2.0	3.9	4.4	4.2	3.9	4.5
2004 5/.....	2.6	*	1.2	2.5	4.0	4.9	4.5	3.9	4.0
2003 5/.....	2.5	*	1.0	1.8	4.1	4.0	4.4	4.7	4.6
2002.....	2.4	*	0.7	1.5	3.4	4.4	4.4	4.1	4.2
2001 6/.....	2.3	*	0.8	1.5	3.1	3.8	4.5	4.3	3.6
2000.....	2.4	*	1.1	1.8	4.3	4.0	3.7	3.7	4.8
1999.....	2.3	*	0.8	2.0	3.2	4.1	4.7	4.3	4.2

Non-Hispanic Black, male

2007 5/.....	3.9	*	1.3	2.6	5.2	6.6	7.2	7.2	6.8
2006 5/.....	4.0	*	1.6	2.9	5.2	6.2	7.7	8.5	5.7
2005 5/.....	3.9	*	1.3	3.0	6.5	6.9	6.5	5.5	7.1
2004 5/.....	4.1	*	1.3	3.8	6.6	8.0	6.6	5.9	6.2
2003 5/.....	3.9	*	1.6	2.8	6.2	6.5	6.8	7.6	7.3
2002.....	3.8	*	1.2	2.5	5.2	7.3	7.6	7.0	6.4
2001 6/.....	3.7	*	1.2	2.3	5.6	6.8	7.4	7.0	5.2
2000.....	3.9	*	1.9	2.6	7.1	6.4	6.6	6.3	7.4
1999.....	3.8	*	*	3.0	5.5	7.1	7.9	7.7	5.8

Non-Hispanic Black, female

2007 5/.....	1.2	*	*	*	1.4	1.7	1.5	2.2	2.1
2006 5/.....	1.0	*	*	*	1.4	*	*	1.7	1.7
2005 5/.....	1.4	*	*	*	*	2.0	2.1	2.4	2.1
2004 5/.....	1.3	*	*	*	1.4	2.1	2.6	2.1	2.0
2003 5/.....	1.3	*	*	*	2.0	1.7	2.3	2.2	2.3
2002.....	1.1	*	*	*	1.6	1.7	1.4	1.6	2.3
2001 6/.....	1.1	*	*	*	*	*	1.9	1.9	2.2
2000.....	1.1	*	*	*	1.7	1.9	*	1.4	2.4
1999.....	1.0	*	*	*	*	*	1.9	*	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes

2007.....	6.1	8.3	2.4	0.7	1.0	10.4	15.8	12.9	10.4	7.6	6.5
2006.....	6.2	8.1	2.2	0.8	1.2	10.7	16.2	13.5	9.8	7.6	6.3
2005.....	6.1	7.5	2.3	0.6	1.1	9.9	16.1	13.6	10.0	7.5	6.7
2004.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3	6.3
2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6	6.4
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9	6.5
2001 6/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4	8.6
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6	6.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7	6.6

All origins 3/, male

2007.....	9.8	9.5	2.5	0.7	1.4	17.6	26.7	21.4	17.3	11.8	9.4
2006.....	10.0	9.4	2.5	0.8	1.6	18.2	27.5	22.4	16.2	11.4	9.4
2005.....	9.8	8.2	2.6	0.7	1.3	16.8	27.2	22.8	16.4	11.2	10.0
2004.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2	9.0
2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4	9.5
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8	9.6
2001 6/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3	12.6
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9	9.7
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0	9.5

All origins 3/, female

2007.....	2.5	7.0	2.3	0.7	0.7	2.8	4.2	4.0	3.4	3.5	3.6
2006.....	2.5	6.8	2.0	0.7	0.7	2.9	4.2	4.2	3.3	3.6	3.3
2005.....	2.5	6.6	2.0	0.6	0.8	2.5	4.3	4.0	3.5	3.9	3.4
2004.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3	3.6
2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9	3.4
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1	3.5
2001 6/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5	4.7
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4	3.6
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Hispanic, both sexes

2007	7.6	7.8	2.0	0.5	1.3	14.0	20.9	13.9	11.2	7.8	6.8
2006	8.0	8.3	1.7	0.7	1.7	15.8	20.5	14.5	9.9	9.2	6.8
2005	8.2	6.4	2.3	*	1.1	15.7	20.4	15.8	11.8	9.1	7.0
2004	7.9	8.1	2.3	0.7	1.1	14.8	19.7	15.2	9.9	8.2	7.2
2003	8.4	6.9	2.5	*	1.0	14.5	21.8	15.6	10.6	9.3	7.5
2002	8.1	6.2	2.6	0.8	1.0	14.8	20.2	14.8	9.8	8.7	7.2
2001 6/	9.0	6.7	2.3	0.5	1.1	13.8	22.3	16.1	11.7	11.0	9.6
2000	8.3	7.0	1.9	0.7	1.6	15.0	19.1	13.7	11.1	8.8	8.6
1999	8.4	7.0	1.8	0.8	2.0	15.7	19.6	14.8	10.5	9.1	8.4

Hispanic, male

2007	12.4	8.3	1.5	*	1.9	24.6	35.2	21.9	17.9	12.1	10.1
2006	13.1	10.1	1.8	*	2.5	28.0	33.8	23.5	15.7	14.3	10.4
2005	13.6	6.3	2.5	*	1.6	27.8	33.9	25.3	18.9	13.9	11.3
2004	13.0	9.1	2.5	*	1.4	25.8	32.8	24.5	15.7	13.1	10.4
2003	13.6	7.9	2.1	*	1.5	25.1	34.5	25.2	16.9	14.6	11.0
2002	13.2	6.6	3.1	*	1.2	25.5	32.9	23.9	15.4	13.0	11.1
2001 6/	14.5	7.9	2.5	*	1.5	23.9	36.1	25.3	17.6	16.5	15.2
2000	13.4	6.6	2.0	*	2.3	25.7	30.9	22.2	17.5	13.5	13.6
1999	13.8	6.7	1.7	*	2.8	26.0	31.8	24.3	16.6	14.1	13.6

Hispanic, female

2007	2.5	7.3	2.4	*	*	2.8	4.3	3.8	3.3	2.8	3.2
2006	2.4	6.5	1.5	*	*	3.0	4.7	3.3	3.1	3.4	2.8
2005	2.5	6.6	2.0	*	*	2.9	4.3	3.7	3.6	3.7	2.4
2004	2.4	7.0	2.1	*	*	2.9	3.7	3.6	3.3	2.7	3.7
2003	2.8	5.9	2.9	*	*	3.1	5.9	3.8	3.3	3.4	3.8
2002	2.6	5.9	2.1	*	*	3.1	4.5	3.6	3.2	3.8	3.1
2001 6/	3.2	5.5	2.1	*	*	2.6	5.3	5.2	4.9	5.0	3.7
2000	2.8	7.4	1.9	*	*	2.8	4.6	3.7	3.8	3.6	3.5
1999	2.8	7.3	1.9	*	*	4.0	4.9	3.6	3.4	3.5	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	5.8	8.3	2.5	0.7	1.0	9.5	14.6	12.6	10.2	7.6	6.4
2006.....	5.9	8.0	2.4	0.8	1.0	9.7	15.2	13.2	9.7	7.2	6.2
2005.....	5.7	7.7	2.3	0.7	1.0	8.7	15.1	13.0	9.6	7.2	6.6
2004.....	5.5	7.8	2.4	0.6	1.0	8.2	13.9	12.7	9.0	7.0	6.1
2003.....	5.7	8.9	2.3	0.7	0.9	8.4	15.3	12.8	9.1	7.3	6.2
2002.....	5.7	7.7	2.7	0.7	1.0	8.2	15.5	12.7	9.3	7.7	6.4
2001 6/.....	6.7	8.4	2.8	0.7	0.9	8.5	16.0	14.4	11.3	10.1	8.4
2000.....	5.6	9.6	2.4	0.7	1.0	8.4	15.1	11.5	8.6	7.4	6.4
1999.....	5.7	8.9	2.6	0.9	1.1	9.4	14.7	11.5	8.7	7.4	6.3

Non-Hispanic 4/, male

2007.....	9.2	9.6	2.8	0.7	1.2	16.0	24.6	21.2	17.0	11.6	9.2
2006.....	9.3	9.1	2.7	0.8	1.4	16.2	25.9	22.0	16.2	10.8	9.1
2005.....	9.1	8.8	2.6	0.7	1.2	14.6	25.4	22.0	15.7	10.5	9.7
2004.....	8.7	7.5	2.5	0.6	1.2	13.2	23.6	21.5	14.5	10.8	8.8
2003.....	9.0	10.5	2.6	0.8	1.3	14.0	25.8	21.5	14.3	10.7	9.2
2002.....	8.9	8.0	2.9	0.6	1.2	13.3	26.2	21.3	14.3	11.5	9.3
2001 6/.....	10.4	9.8	3.2	0.7	1.0	14.0	26.8	23.4	17.5	14.9	12.1
2000.....	8.6	11.2	2.6	0.7	1.3	13.4	25.5	18.9	13.2	10.4	9.2
1999.....	8.6	10.1	2.6	0.9	1.3	15.1	24.3	18.3	13.1	10.4	9.0

Non-Hispanic 4/, female

2007.....	2.5	6.9	2.2	0.7	0.7	2.8	4.2	4.0	3.4	3.6	3.7
2006.....	2.5	6.8	2.1	0.7	0.7	2.9	4.0	4.4	3.3	3.7	3.3
2005.....	2.5	6.6	2.0	0.6	0.9	2.5	4.3	4.0	3.5	3.9	3.5
2004.....	2.5	8.1	2.3	0.6	0.7	3.0	4.0	3.9	3.6	3.4	3.6
2003.....	2.5	7.2	2.1	0.6	0.6	2.5	4.5	4.0	4.0	3.9	3.3
2002.....	2.7	7.4	2.6	0.8	0.8	2.8	4.6	4.3	4.2	4.1	3.5
2001 6/.....	3.2	6.9	2.4	0.7	0.7	2.7	5.0	5.4	5.2	5.5	4.7
2000.....	2.7	7.9	2.2	0.7	0.8	3.0	4.7	4.2	4.0	4.5	3.6
1999.....	2.9	7.6	2.5	0.9	0.9	3.4	5.1	4.7	4.4	4.4	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic White, both sexes

2007 5/.....	2.7	6.5	1.6	0.6	0.4	2.6	4.5	4.5	4.2	3.7	3.9
2006 5/.....	2.7	6.3	1.5	0.6	0.6	2.6	4.3	4.1	3.6	3.5	3.7
2005 5/.....	2.7	6.0	1.5	0.4	0.5	2.5	4.3	4.3	3.7	3.6	3.9
2004 5/.....	2.7	5.9	1.4	0.4	0.6	2.6	4.4	4.2	3.8	3.7	3.8
2003 5/.....	2.7	6.8	1.3	0.4	0.5	2.5	4.5	4.0	3.8	3.9	3.7
2002.....	2.8	5.0	1.8	0.5	0.6	2.8	4.7	4.1	3.8	4.2	4.0
2001 6/.....	4.0	5.8	1.7	0.7	0.6	2.9	5.5	6.5	6.6	7.0	6.2
2000.....	2.8	6.2	1.6	0.6	0.7	2.7	4.4	4.0	3.9	4.1	3.8
1999.....	2.9	6.3	1.7	0.6	0.7	3.1	4.7	4.2	4.4	4.3	3.9

Non-Hispanic White, male

2007 5/.....	3.7	7.2	1.9	0.6	0.4	3.4	6.3	6.4	6.0	5.0	5.1
2006 5/.....	3.6	6.5	1.6	0.5	0.6	3.4	6.1	5.8	5.1	4.6	5.1
2005 5/.....	3.6	6.9	1.6	0.6	0.4	3.5	6.0	5.9	5.0	4.5	5.1
2004 5/.....	3.6	6.1	1.6	0.4	0.7	3.3	6.1	5.8	5.1	5.2	4.9
2003 5/.....	3.6	8.0	1.3	0.4	0.7	3.6	6.5	5.4	5.1	5.0	4.8
2002.....	3.8	5.8	1.9	0.3	0.6	3.9	6.6	5.6	5.0	5.6	5.7
2001 6/.....	5.6	6.8	2.0	0.6	0.6	3.9	7.5	9.3	9.4	10.3	8.7
2000.....	3.6	8.3	1.8	0.6	0.8	3.5	6.0	5.2	5.2	5.1	5.3
1999.....	3.9	8.0	1.7	0.6	0.8	4.2	6.1	5.7	5.7	5.6	5.2

Non-Hispanic White, female

2007 5/.....	1.8	5.7	1.4	0.6	0.4	1.8	2.6	2.5	2.4	2.4	2.7
2006 5/.....	1.8	6.2	1.3	0.7	0.5	1.7	2.4	2.5	2.0	2.4	2.3
2005 5/.....	1.8	5.0	1.4	*	0.6	1.5	2.5	2.7	2.4	2.7	2.6
2004 5/.....	1.8	5.8	1.3	0.4	0.4	1.8	2.7	2.6	2.4	2.3	2.7
2003 5/.....	1.8	5.5	1.4	*	*	1.4	2.4	2.6	2.4	2.8	2.5
2002.....	1.9	4.1	1.8	0.6	0.6	1.6	2.7	2.5	2.7	2.8	2.3
2001 6/.....	2.4	4.7	1.5	0.7	0.6	1.9	3.5	3.6	3.8	3.7	3.7
2000.....	1.9	4.1	1.3	0.5	0.6	1.9	2.8	2.8	2.6	3.1	2.3
1999.....	2.0	4.4	1.6	0.6	0.7	2.0	3.3	2.8	3.1	3.0	2.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic Black, both sexes

2007 5/.....	22.9	16.1	6.3	1.2	3.2	38.5	61.0	51.9	40.7	29.0	21.3
2006 5/.....	23.6	15.9	6.0	1.7	3.0	38.8	64.7	56.4	41.4	27.9	21.2
2005 5/.....	22.8	14.5	5.5	1.5	3.1	34.6	63.4	55.1	40.4	27.5	22.2
2004 5/.....	21.7	15.8	6.2	1.4	2.6	32.8	58.1	55.1	36.8	26.1	20.6
2003 5/.....	22.7	16.8	6.4	1.8	2.8	34.2	64.5	56.9	37.7	26.3	22.1
2002.....	22.5	17.6	6.5	1.4	2.6	31.9	65.6	56.5	38.1	27.8	20.9
2001 6/.....	22.7	19.5	7.1	0.9	1.9	33.6	64.9	54.6	36.5	29.0	22.0
2000.....	22.1	24.1	6.1	1.2	2.6	34.1	66.4	50.4	34.3	26.6	22.3
1999.....	21.7	20.0	6.4	2.0	2.6	37.7	63.0	48.8	32.6	25.7	21.5

Non-Hispanic Black, male

2007 5/.....	41.1	19.9	6.9	1.4	4.6	69.1	109.3	94.5	76.4	50.6	35.5
2006 5/.....	42.0	20.3	6.4	2.2	4.3	68.8	116.4	101.9	76.9	47.9	34.7
2005 5/.....	40.9	15.9	6.9	1.6	4.1	62.0	114.0	102.7	75.1	46.3	37.8
2004 5/.....	38.3	13.9	6.3	1.4	3.5	56.9	105.9	102.9	67.0	44.6	33.6
2003 5/.....	40.1	18.6	7.8	2.1	3.8	60.7	115.5	106.2	66.6	45.4	37.9
2002.....	39.4	16.4	7.3	1.5	3.5	54.8	119.6	104.5	67.2	46.6	33.1
2001 6/.....	39.3	21.7	8.4	*	2.8	59.7	120.3	99.7	64.5	44.2	34.7
2000.....	38.3	24.9	6.1	*	3.3	59.0	122.1	93.5	59.9	43.8	34.8
1999.....	37.2	20.1	6.2	2.0	3.5	64.6	116.0	87.2	56.2	41.1	34.6

Non-Hispanic Black, female

2007 5/.....	6.4	12.1	5.8	*	1.8	7.0	11.5	11.1	8.4	9.7	8.8
2006 5/.....	6.8	11.3	5.6	*	1.7	7.9	11.8	13.5	9.4	10.1	9.3
2005 5/.....	6.3	13.0	4.1	1.4	2.0	6.4	11.9	10.2	9.2	10.6	8.5
2004 5/.....	6.6	17.7	6.1	1.5	1.6	7.9	10.2	10.6	9.5	9.5	9.1
2003 5/.....	6.8	14.8	4.9	1.4	1.7	7.0	14.0	11.4	11.8	9.4	8.2
2002.....	7.2	18.9	5.7	1.4	1.6	8.2	13.2	12.7	11.9	11.1	10.2
2001 6/.....	7.6	17.1	5.7	*	*	6.6	11.8	13.6	11.4	15.6	10.7
2000.....	7.4	23.3	6.0	1.5	1.8	8.6	13.7	11.4	11.3	11.3	11.2
1999.....	7.7	20.0	6.7	1.9	1.6	10.1	12.8	14.1	11.4	12.1	10.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All origins 3/, both sexes											
2007.....	4.2	*	0.3	0.2	0.8	8.8	13.2	10.4	8.0	5.3	4.2
2006.....	4.3	*	0.3	0.3	0.8	9.1	13.7	10.8	7.7	5.2	3.9
2005.....	4.2	*	0.2	0.2	0.7	8.3	13.1	11.0	7.8	5.1	4.1
2004.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9	3.8
2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0	3.8
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2	3.9
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1	3.7
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7	3.8
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6	3.7
All origins 3/, male											
2007.....	7.2	*	0.3	0.2	1.1	15.4	23.1	18.0	14.0	8.7	6.5
2006.....	7.4	*	0.3	0.3	1.3	16.0	24.1	18.7	13.4	8.4	6.2
2005.....	7.2	*	0.3	0.3	1.0	14.7	23.1	19.3	13.5	8.1	6.6
2004.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3	5.9
2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0	5.9
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4	6.2
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0	5.6
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4	5.8
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1	5.5
All origins 3/, female											
2007.....	1.2	*	0.3	0.2	0.4	1.9	2.6	2.3	2.0	1.8	1.8
2006.....	1.3	*	*	0.3	0.4	1.8	2.6	2.5	1.9	2.0	1.6
2005.....	1.2	*	*	*	0.4	1.5	2.4	2.3	2.1	2.1	1.7
2004.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4	1.6
2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9	1.7
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0	1.7
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2	1.7
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9	1.7
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Hispanic, both sexes

2007.....	5.2	*	*	*	1.0	11.4	16.9	10.3	8.0	5.2	4.2
2006.....	5.6	*	*	*	1.3	13.4	16.7	11.1	7.5	5.8	4.4
2005.....	5.7	*	*	*	0.8	13.2	15.6	12.2	8.9	6.1	4.4
2004.....	5.4	*	*	*	0.7	12.4	15.4	11.3	7.3	5.4	4.6
2003.....	5.8	*	*	*	0.7	11.9	17.6	11.9	7.7	6.1	4.4
2002.....	5.6	*	*	*	0.7	12.6	15.9	11.6	6.8	5.4	4.6
2001.....	5.7	*	*	*	0.7	11.7	17.1	11.7	7.2	5.8	5.4
2000.....	5.5	*	*	*	1.0	12.6	14.8	10.3	7.4	4.8	4.9
1999.....	5.7	*	*	*	1.4	13.4	15.3	10.5	6.8	5.4	5.0

Hispanic, male

2007.....	9.1	*	*	*	1.6	20.2	29.2	16.9	13.3	8.4	6.5
2006.....	9.7	*	*	*	2.1	24.0	28.7	18.3	12.4	9.3	6.9
2005.....	10.0	*	*	*	1.3	24.1	26.8	20.2	14.8	9.3	7.2
2004.....	9.5	*	*	*	1.1	22.0	26.3	19.3	12.1	9.4	7.0
2003.....	10.0	*	*	*	1.1	21.1	29.0	20.1	13.1	9.7	6.7
2002.....	9.7	*	*	*	*	22.3	27.0	19.3	11.6	8.7	7.3
2001.....	9.9	*	*	*	*	20.4	28.8	19.8	12.0	9.4	8.8
2000.....	9.6	*	*	*	1.7	21.9	24.9	17.5	12.3	7.5	8.1
1999.....	9.8	*	*	*	2.1	22.9	25.4	18.1	11.1	9.1	7.9

Hispanic, female

2007.....	1.1	*	*	*	*	2.0	2.7	2.2	1.7	1.4	1.5
2006.....	1.2	*	*	*	*	2.1	2.4	2.0	1.6	1.8	1.5
2005.....	1.2	*	*	*	*	1.6	2.2	2.1	2.1	2.4	*
2004.....	1.1	*	*	*	*	2.1	2.0	1.4	1.8	*	2.0
2003.....	1.3	*	*	*	*	2.0	3.3	1.9	1.5	1.9	1.9
2002.....	1.2	*	*	*	*	2.0	2.2	2.2	1.4	1.7	1.8
2001.....	1.3	*	*	*	*	2.1	2.8	2.0	1.6	1.8	1.9
2000.....	1.3	*	*	*	*	2.0	2.5	2.0	1.8	1.9	*
1999.....	1.4	*	*	*	*	2.6	3.2	1.7	1.8	*	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic 4/, both sexes

2007.....	4.0	*	0.3	0.3	0.7	8.3	12.3	10.3	8.0	5.3	4.2
2006.....	4.0	*	0.3	0.3	0.7	8.2	12.9	10.7	7.8	5.1	3.8
2005.....	3.9	*	0.2	0.2	0.7	7.3	12.5	10.6	7.5	4.8	4.0
2004.....	3.7	*	0.2	0.2	0.6	6.7	11.4	10.3	6.9	4.7	3.6
2003.....	3.8	*	0.2	0.3	0.7	6.9	12.7	10.5	6.9	4.7	3.7
2002.....	3.8	*	0.3	0.3	0.7	6.7	13.0	10.3	6.9	5.1	3.8
2001.....	3.7	*	0.3	0.3	0.5	6.7	12.7	9.8	6.5	5.0	3.4
2000.....	3.6	*	0.2	0.2	0.6	6.7	12.2	9.0	6.0	4.6	3.6
1999.....	3.6	*	0.4	0.3	0.7	7.6	11.9	8.8	5.9	4.5	3.5

Non-Hispanic 4/, male

2007.....	6.9	*	0.3	0.3	1.0	14.3	21.5	18.3	14.0	8.7	6.5
2006.....	6.9	*	*	0.3	1.1	14.3	22.8	18.7	13.7	8.1	6.0
2005.....	6.7	*	*	0.3	0.9	12.7	22.2	18.9	13.0	7.8	6.4
2004.....	6.4	*	*	0.2	1.0	11.4	20.3	18.5	11.8	8.0	5.7
2003.....	6.5	*	*	0.3	1.0	12.1	22.4	18.7	11.7	7.7	5.8
2002.....	6.5	*	0.3	*	0.9	11.4	22.9	18.2	11.5	8.3	6.0
2001.....	6.3	*	0.3	0.4	0.8	11.7	22.6	17.1	10.8	7.8	5.2
2000.....	6.0	*	*	0.2	0.9	11.5	21.6	15.7	10.0	7.3	5.5
1999.....	6.0	*	0.4	0.4	1.0	12.9	20.9	15.3	9.8	6.7	5.2

Non-Hispanic 4/, female

2007.....	1.2	*	0.3	*	0.4	1.9	2.6	2.4	2.0	1.9	1.9
2006.....	1.3	*	*	0.3	0.4	1.8	2.6	2.6	2.0	2.1	1.7
2005.....	1.2	*	*	*	0.4	1.5	2.4	2.4	2.1	2.0	1.8
2004.....	1.2	*	*	*	0.3	1.7	2.3	2.1	2.1	1.5	1.6
2003.....	1.2	*	*	*	0.3	1.4	2.7	2.2	2.2	1.9	1.7
2002.....	1.3	*	*	0.3	0.5	1.8	3.0	2.4	2.2	2.0	1.7
2001.....	1.2	*	0.3	0.3	0.3	1.4	2.6	2.6	2.1	2.3	1.7
2000.....	1.2	*	*	*	0.3	1.7	2.7	2.3	2.1	2.0	1.8
1999.....	1.3	*	0.4	*	0.4	2.0	2.8	2.5	2.1	2.2	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic White, both sexes

2007 5/.....	1.5	*	*	0.2	0.3	1.8	3.1	3.1	2.9	2.3	2.3
2006 5/.....	1.4	*	*	0.2	0.3	1.7	2.7	2.6	2.4	2.2	2.1
2005 5/.....	1.4	*	*	*	0.3	1.7	2.6	2.8	2.4	2.2	2.3
2004 5/.....	1.5	*	*	*	0.3	1.6	2.7	2.7	2.6	2.2	2.1
2003 5/.....	1.4	*	*	*	0.3	1.7	3.0	2.6	2.4	2.3	2.0
2002.....	1.5	*	0.2	0.2	0.3	1.9	3.1	2.7	2.5	2.5	2.3
2001.....	1.6	*	0.2	0.3	0.4	1.8	3.2	2.8	2.6	2.7	2.1
2000.....	1.4	*	*	0.2	0.3	1.7	2.7	2.5	2.4	2.4	2.1
1999.....	1.5	*	0.2	0.2	0.4	2.0	2.9	2.7	2.6	2.5	2.1

Non-Hispanic White, male

2007 5/.....	2.2	*	*	*	*	2.5	4.5	4.7	4.3	3.4	3.3
2006 5/.....	2.0	*	*	*	0.4	2.6	4.2	4.0	3.6	3.1	3.1
2005 5/.....	2.1	*	*	*	0.3	2.5	4.0	4.1	3.5	2.9	3.1
2004 5/.....	2.1	*	*	*	0.5	2.2	4.0	4.2	3.8	3.5	2.9
2003 5/.....	2.1	*	*	*	0.4	2.6	4.8	3.9	3.5	3.1	2.8
2002.....	2.2	*	*	*	0.4	2.8	4.7	4.0	3.5	3.5	3.4
2001.....	2.2	*	*	0.4	0.5	2.6	4.7	4.3	3.6	3.9	2.9
2000.....	2.0	*	*	*	0.5	2.5	4.0	3.5	3.4	3.4	3.0
1999.....	2.2	*	*	*	0.5	3.0	4.2	4.0	3.8	3.4	2.9

Non-Hispanic White, female

2007 5/.....	0.9	*	*	*	*	1.1	1.6	1.4	1.5	1.2	1.3
2006 5/.....	0.8	*	*	*	*	0.9	1.2	1.3	1.1	1.4	1.1
2005 5/.....	0.8	*	*	*	*	0.9	1.1	1.5	1.4	1.5	1.4
2004 5/.....	0.8	*	*	*	*	1.0	1.3	1.1	1.3	1.0	1.4
2003 5/.....	0.8	*	*	*	*	0.7	1.2	1.2	1.4	1.5	1.2
2002.....	0.9	*	*	*	*	1.0	1.5	1.4	1.4	1.5	1.2
2001.....	0.9	*	*	*	*	0.9	1.6	1.4	1.6	1.5	1.3
2000.....	0.9	*	*	*	*	0.9	1.5	1.4	1.4	1.5	1.3
1999.....	0.9	*	*	*	0.3	0.9	1.5	1.3	1.4	1.6	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic Black, both sexes

2007 5/.....	18.0	*	*	*	2.4	35.3	55.0	45.5	34.7	22.0	15.0
2006 5/.....	18.6	*	*	*	2.5	34.7	59.2	49.5	35.9	21.4	14.2
2005 5/.....	17.7	*	*	*	2.1	31.0	57.1	48.8	34.6	20.1	14.9
2004 5/.....	16.6	*	*	*	1.9	28.8	52.4	48.5	30.4	19.1	13.1
2003 5/.....	17.3	*	*	0.8	2.1	30.0	57.1	50.1	31.3	18.8	14.1
2002.....	17.1	*	*	*	2.0	28.1	59.0	48.9	31.0	20.1	13.3
2001.....	16.3	*	*	*	1.4	29.5	57.6	45.3	27.9	19.1	11.8
2000.....	16.0	*	*	*	1.8	29.5	57.3	42.5	26.6	17.8	12.9
1999.....	15.9	*	*	0.7	1.9	33.0	55.8	40.7	24.3	16.3	12.5

Non-Hispanic Black, male

2007 5/.....	34.0	*	*	*	3.6	64.6	101.0	85.6	67.2	40.6	26.5
2006 5/.....	35.0	*	*	*	3.8	63.2	108.4	92.1	68.9	39.2	24.8
2005 5/.....	33.6	*	*	*	3.2	56.9	105.6	93.4	66.2	37.0	26.9
2004 5/.....	31.4	*	*	*	3.0	51.8	97.8	93.4	57.3	35.9	24.0
2003 5/.....	32.6	*	*	*	3.0	55.0	105.3	96.7	58.8	35.3	25.1
2002.....	31.9	*	*	*	2.8	49.8	109.9	94.2	58.1	36.9	23.1
2001.....	30.6	*	*	*	2.2	54.2	110.3	86.3	53.2	32.6	20.7
2000.....	30.1	*	*	*	2.8	53.3	109.1	82.3	49.4	32.6	22.0
1999.....	29.3	*	*	*	2.9	58.3	105.7	77.2	45.3	28.0	21.0

Non-Hispanic Black, female

2007 5/.....	3.4	*	*	*	*	5.2	8.0	7.1	5.1	5.5	5.0
2006 5/.....	3.7	*	*	*	*	5.6	8.9	9.3	6.1	5.6	4.8
2005 5/.....	3.2	*	*	*	*	4.4	7.9	6.7	6.1	5.0	4.3
2004 5/.....	3.1	*	*	*	*	5.1	6.9	6.7	6.1	4.2	3.5
2003 5/.....	3.4	*	*	*	1.2	4.2	9.4	7.2	6.5	4.2	4.6
2002.....	3.7	*	*	*	*	5.6	9.5	7.6	6.8	5.2	4.7
2001.....	3.2	*	*	*	*	4.0	7.2	8.0	5.2	7.0	4.0
2000.....	3.3	*	*	*	*	5.2	8.4	6.5	6.2	4.6	4.9
1999.....	3.7	*	*	*	*	7.2	8.5	7.8	5.5	5.9	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes

2007.....	1.9	7.9	2.1	0.4	0.3	1.5	2.6	2.5	2.4	2.4	2.3
2006.....	1.9	8.0	2.0	0.4	0.3	1.6	2.6	2.7	2.0	2.3	2.4
2005.....	1.9	7.3	2.1	0.4	0.4	1.6	3.0	2.6	2.2	2.5	2.5
2004.....	2.0	7.8	2.1	0.4	0.3	1.7	2.9	2.7	2.2	2.4	2.5
2003.....	2.0	8.3	2.1	0.4	0.3	1.7	2.9	2.6	2.3	2.7	2.6
2002.....	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7	2.6
2001 6/.....	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2	4.9
2000.....	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9	2.9
1999.....	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0	2.9

All origins 3/, male

2007.....	2.5	8.9	2.3	0.4	0.3	2.1	3.6	3.4	3.3	3.1	2.9
2006.....	2.6	9.3	2.2	0.5	0.3	2.2	3.5	3.7	2.7	3.1	3.2
2005.....	2.6	7.9	2.3	0.4	0.3	2.1	4.0	3.5	3.0	3.1	3.4
2004.....	2.5	7.9	2.3	0.4	0.3	2.2	4.0	3.5	2.9	2.9	3.1
2003.....	2.6	9.7	2.3	0.4	0.4	2.3	3.8	3.3	2.8	3.4	3.6
2002.....	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4	3.4
2001 6/.....	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3	7.0
2000.....	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5	3.9
1999.....	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9	4.0

All origins 3/, female

2007.....	1.3	6.8	2.0	0.5	0.3	0.9	1.6	1.7	1.5	1.7	1.8
2006.....	1.3	6.6	1.8	0.4	0.3	1.1	1.6	1.7	1.3	1.6	1.6
2005.....	1.3	6.6	1.8	0.4	0.4	1.0	1.9	1.7	1.5	1.8	1.7
2004.....	1.4	7.7	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9	1.9
2003.....	1.4	6.9	2.0	0.3	0.2	1.1	2.0	1.8	1.8	1.9	1.6
2002.....	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1	1.8
2001 6/.....	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2	2.9
2000.....	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4	1.8
1999.....	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Hispanic, both sexes

2007	2.4	7.3	1.8	*	*	2.6	4.0	3.5	3.2	2.6	2.7
2006	2.4	8.2	1.4	*	*	2.5	3.8	3.4	2.5	3.4	2.4
2005	2.5	6.3	2.1	*	*	2.5	4.8	3.6	2.9	3.0	2.6
2004	2.5	7.7	2.1	*	*	2.4	4.4	3.8	2.6	2.8	2.6
2003	2.6	6.8	2.2	*	*	2.6	4.2	3.7	2.8	3.2	3.1
2002	2.5	6.0	2.2	*	*	2.2	4.3	3.2	2.9	3.2	2.6
2001 6/	3.3	6.2	1.9	*	*	2.1	5.2	4.4	4.5	5.3	4.2
2000	2.7	6.9	1.8	*	*	2.4	4.3	3.3	3.7	4.0	3.7
1999	2.7	6.7	1.6	*	*	2.3	4.3	4.2	3.7	3.6	3.4

Hispanic, male

2007	3.4	7.4	1.4	*	*	4.4	6.0	5.0	4.5	3.7	3.6
2006	3.4	10.1	1.5	*	*	4.0	5.1	5.1	3.3	4.9	3.5
2005	3.6	6.1	2.3	*	*	3.7	7.1	5.2	4.1	4.6	4.1
2004	3.5	8.9	2.3	*	*	3.9	6.5	5.2	3.6	3.7	3.4
2003	3.6	7.7	1.8	*	*	4.0	5.5	5.1	3.8	4.8	4.3
2002	3.5	6.6	2.6	*	*	3.2	5.9	4.7	3.9	4.3	3.7
2001 6/	4.5	7.2	2.2	*	*	3.5	7.4	5.4	5.6	7.2	6.5
2000	3.8	6.6	1.8	*	*	3.8	6.0	4.7	5.2	6.0	5.4
1999	4.0	6.2	1.5	*	*	3.2	6.4	6.2	5.5	5.1	5.7

Hispanic, female

2007	1.3	7.1	2.2	*	*	*	1.6	1.6	1.6	1.4	1.6
2006	1.3	6.3	1.3	*	*	*	2.3	1.2	1.4	1.6	*
2005	1.3	6.6	1.9	*	*	1.3	2.1	1.6	1.5	1.3	*
2004	1.4	6.5	1.9	*	*	*	1.8	2.1	1.5	1.8	1.7
2003	1.5	5.9	2.6	*	*	*	2.6	1.9	1.8	1.5	1.9
2002	1.4	5.4	1.8	*	*	*	2.2	1.4	1.8	2.1	*
2001 6/	1.9	5.2	1.6	*	*	*	2.5	3.2	3.2	3.2	1.8
2000	1.6	7.2	1.7	*	*	*	2.1	1.7	2.0	1.8	2.0
1999	1.4	7.3	1.8	*	*	1.4	1.7	1.9	1.6	2.1	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	1.8	8.0	2.2	0.4	0.3	1.3	2.3	2.3	2.2	2.3	2.3
2006.....	1.8	7.8	2.2	0.4	0.3	1.5	2.3	2.5	1.9	2.1	2.4
2005.....	1.8	7.6	2.1	0.4	0.4	1.4	2.6	2.4	2.0	2.3	2.5
2004.....	1.8	7.7	2.1	0.4	0.3	1.6	2.5	2.4	2.1	2.3	2.5
2003.....	1.9	8.6	2.1	0.4	0.3	1.5	2.6	2.3	2.2	2.5	2.5
2002.....	1.9	7.5	2.4	0.4	0.3	1.5	2.5	2.4	2.4	2.6	2.5
2001 6/.....	3.1	8.2	2.4	0.4	0.3	1.8	3.3	4.6	4.9	5.1	4.9
2000.....	2.0	9.2	2.2	0.4	0.4	1.7	3.0	2.6	2.6	2.8	2.8
1999.....	2.1	8.7	2.2	0.6	0.4	1.8	2.8	2.6	2.8	2.9	2.8

Non-Hispanic 4/, male

2007.....	2.4	9.3	2.5	0.4	0.2	1.6	3.1	2.9	3.0	2.9	2.7
2006.....	2.4	9.0	2.4	0.5	0.3	1.8	3.1	3.2	2.6	2.6	3.1
2005.....	2.4	8.5	2.3	0.5	0.3	1.8	3.2	3.1	2.7	2.8	3.3
2004.....	2.3	7.4	2.3	0.4	0.2	1.9	3.3	3.0	2.7	2.7	3.1
2003.....	2.4	10.1	2.4	0.5	0.3	1.9	3.4	2.8	2.5	3.1	3.5
2002.....	2.5	7.8	2.5	0.4	0.3	1.9	3.4	3.1	2.8	3.2	3.4
2001 6/.....	4.2	9.5	2.8	0.4	0.2	2.2	4.2	6.3	6.7	7.1	6.9
2000.....	2.6	10.6	2.4	0.4	0.3	1.9	3.9	3.2	3.2	3.1	3.8
1999.....	2.6	10.0	2.3	0.5	0.3	2.3	3.3	3.0	3.3	3.7	3.8

Non-Hispanic 4/, female

2007.....	1.3	6.7	1.9	0.4	0.3	0.9	1.5	1.7	1.4	1.7	1.8
2006.....	1.3	6.6	1.9	0.4	0.3	1.1	1.4	1.8	1.3	1.6	1.7
2005.....	1.3	6.6	1.8	0.4	0.5	0.9	1.9	1.7	1.4	1.9	1.7
2004.....	1.4	8.0	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9	2.0
2003.....	1.4	7.1	1.8	0.3	0.3	1.1	1.8	1.8	1.8	2.0	1.6
2002.....	1.4	7.1	2.3	0.5	0.3	1.1	1.7	1.8	2.0	2.1	1.7
2001 6/.....	2.0	6.8	2.0	0.4	0.4	1.3	2.4	2.8	3.1	3.2	3.0
2000.....	1.5	7.8	2.0	0.5	0.5	1.4	2.0	1.9	1.9	2.5	1.8
1999.....	1.6	7.3	2.2	0.7	0.4	1.4	2.3	2.2	2.4	2.1	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic White, both sexes

2007 5/.....	1.2	6.3	1.4	0.4	*	0.8	1.4	1.4	1.3	1.4	1.5
2006 5/.....	1.3	6.3	1.3	0.3	0.2	0.8	1.6	1.5	1.2	1.3	1.6
2005 5/.....	1.2	5.9	1.3	0.3	0.2	0.9	1.7	1.5	1.3	1.4	1.6
2004 5/.....	1.3	5.9	1.4	0.3	0.2	1.0	1.8	1.6	1.2	1.5	1.7
2003 5/.....	1.2	6.8	1.2	0.2	0.2	0.8	1.5	1.4	1.4	1.6	1.6
2002.....	1.3	5.0	1.6	0.3	0.2	0.9	1.6	1.4	1.4	1.7	1.7
2001 6/.....	2.4	5.6	1.5	0.3	0.3	1.2	2.4	3.6	4.0	4.3	4.1
2000.....	1.3	6.0	1.4	0.3	0.4	1.0	1.7	1.6	1.5	1.7	1.7
1999.....	1.4	6.2	1.4	0.4	0.3	1.2	1.9	1.6	1.8	1.8	1.8

Non-Hispanic White, male

2007 5/.....	1.5	7.0	1.6	0.4	*	0.9	1.8	1.6	1.8	1.7	1.7
2006 5/.....	1.6	6.5	1.5	*	*	0.8	1.9	1.8	1.5	1.6	1.9
2005 5/.....	1.5	6.7	1.4	0.3	*	1.1	1.9	1.8	1.5	1.6	2.0
2004 5/.....	1.5	6.1	1.5	*	*	1.1	2.1	1.6	1.3	1.7	2.0
2003 5/.....	1.5	8.0	1.2	*	*	1.0	1.7	1.4	1.6	1.9	2.1
2002.....	1.6	5.8	1.6	*	*	1.1	2.0	1.6	1.5	2.1	2.3
2001 6/.....	3.4	6.7	1.8	*	*	1.3	2.8	5.0	5.7	6.4	5.8
2000.....	1.6	8.1	1.7	0.4	0.3	1.0	2.0	1.7	1.8	1.8	2.3
1999.....	1.7	7.8	1.5	0.4	*	1.3	1.9	1.7	1.9	2.2	2.3

Non-Hispanic White, female

2007 5/.....	0.9	5.6	1.2	0.4	*	0.7	1.0	1.1	0.9	1.2	1.4
2006 5/.....	1.0	6.1	1.1	0.4	*	0.8	1.2	1.2	0.9	1.0	1.2
2005 5/.....	0.9	5.0	1.2	*	0.3	0.6	1.4	1.2	1.1	1.2	1.2
2004 5/.....	1.0	5.8	1.2	*	*	0.9	1.4	1.5	1.1	1.3	1.3
2003 5/.....	1.0	5.4	1.2	*	*	0.7	1.2	1.4	1.1	1.4	1.2
2002.....	1.0	4.1	1.6	0.4	*	0.6	1.3	1.1	1.3	1.4	1.1
2001 6/.....	1.5	4.5	1.2	0.4	0.4	1.0	1.9	2.2	2.3	2.2	2.4
2000.....	1.0	3.9	1.2	*	0.4	1.0	1.4	1.4	1.2	1.7	1.0
1999.....	1.1	4.4	1.4	0.5	0.4	1.1	1.8	1.4	1.7	1.4	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic Black, both sexes

2007 5/.....	4.9	15.3	5.6	0.7	0.8	3.2	6.0	6.4	6.1	6.9	6.3
2006 5/.....	5.0	15.3	5.2	1.0	*	4.0	5.5	6.9	5.5	6.5	7.0
2005 5/.....	5.1	14.1	5.0	1.0	0.9	3.6	6.2	6.3	5.8	7.4	7.3
2004 5/.....	5.1	15.3	5.4	0.9	0.7	4.0	5.7	6.6	6.4	7.0	7.5
2003 5/.....	5.4	15.8	5.6	1.0	0.6	4.3	7.4	6.8	6.5	7.5	7.9
2002.....	5.4	16.7	5.9	1.0	0.6	3.8	6.6	7.6	7.0	7.6	7.6
2001 6/.....	6.4	19.1	6.3	*	*	4.1	7.3	9.3	8.6	10.0	10.2
2000.....	6.1	22.9	5.7	0.9	0.8	4.6	9.0	7.8	7.7	8.8	9.4
1999.....	5.8	19.7	5.7	1.3	0.7	4.7	7.2	8.1	8.3	9.4	9.1

Non-Hispanic Black, male

2007 5/.....	7.1	19.0	6.2	*	*	4.4	8.3	8.9	9.2	10.0	9.0
2006 5/.....	7.1	19.7	5.6	1.5	*	5.7	8.1	9.8	8.0	8.7	9.9
2005 5/.....	7.4	15.2	6.2	*	*	5.2	8.4	9.3	8.9	9.3	10.8
2004 5/.....	6.9	13.6	5.6	*	*	5.1	8.1	9.5	9.7	8.8	9.7
2003 5/.....	7.6	16.8	6.9	*	*	5.6	10.2	9.5	7.8	10.1	12.9
2002.....	7.6	15.8	6.6	*	*	5.0	9.7	10.3	9.1	9.7	10.1
2001 6/.....	8.7	21.4	7.7	*	*	5.6	10.0	13.4	11.3	11.6	14.1
2000.....	8.2	22.5	5.7	*	*	5.7	13.0	11.2	10.5	11.2	12.8
1999.....	7.9	19.7	5.6	1.2	*	6.3	10.3	10.0	10.9	13.0	13.6

Non-Hispanic Black, female

2007 5/.....	3.0	11.5	4.9	*	*	1.9	3.6	4.1	3.3	4.2	3.8
2006 5/.....	3.1	10.6	4.9	*	*	2.4	2.9	4.2	3.3	4.4	4.4
2005 5/.....	3.1	13.0	3.7	*	*	1.9	4.0	3.5	3.1	5.7	4.3
2004 5/.....	3.5	17.1	5.3	*	*	2.8	3.3	3.9	3.5	5.3	5.6
2003 5/.....	3.4	14.8	4.3	*	*	2.8	4.6	4.2	5.2	5.1	3.6
2002.....	3.5	17.7	5.2	*	*	2.6	3.6	5.1	5.1	5.9	5.5
2001 6/.....	4.4	16.8	5.0	*	*	2.6	4.6	5.6	6.2	8.6	6.7
2000.....	4.1	23.3	5.8	*	*	3.4	5.3	4.8	5.1	6.7	6.4
1999.....	4.0	19.6	5.7	1.4	*	2.9	4.4	6.3	5.9	6.2	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	1.8	2.1	0.3	*	0.2	0.8	1.9	2.1	2.2	2.6	3.2
2006.....	1.7	2.2	0.3	0.1	0.2	0.8	2.0	2.0	2.1	2.6	3.2
2005.....	1.6	2.5	0.4	0.1	0.2	0.7	1.6	1.8	1.9	2.3	3.1
2004.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9	3.4
2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2	3.9
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3	3.4
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0	3.2
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6	3.1
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8	2.9

All origins 3/, male

2007.....	2.3	2.5	0.3	*	0.2	1.3	2.7	2.8	2.7	3.4	3.9
2006.....	2.2	2.7	0.3	*	0.3	1.2	2.9	2.8	2.7	3.2	3.8
2005.....	2.0	3.2	0.5	*	0.3	1.0	2.3	2.4	2.3	2.7	3.6
2004.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8	4.1
2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2	4.9
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5	4.3
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9	4.4
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7	4.2
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9	4.0

All origins 3/, female

2007.....	1.3	1.6	0.3	*	*	0.3	1.0	1.4	1.7	1.9	2.4
2006.....	1.2	1.7	0.3	*	*	0.5	1.0	1.1	1.4	2.0	2.6
2005.....	1.3	1.8	0.3	*	*	0.4	0.9	1.2	1.5	1.9	2.6
2004.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9	2.7
2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2	2.9
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1	2.6
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6	1.9
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	2.0	2.1	0.3	*	0.2	0.9	2.1	2.4	2.6	3.0	3.5
2006.....	1.9	2.3	0.3	0.1	0.2	0.9	2.2	2.3	2.4	2.8	3.5
2005.....	1.7	2.7	0.5	*	0.2	0.7	1.7	2.1	2.1	2.5	3.4
2004.....	1.8	2.8	0.3	0.1	0.2	0.8	1.8	2.1	2.4	3.2	3.7
2003.....	1.9	2.2	0.2	*	0.2	0.9	2.1	2.0	2.5	3.5	4.2
2002.....	1.8	2.7	0.4	0.1	0.2	0.7	1.8	2.1	2.5	3.6	3.7
2001.....	1.6	2.4	0.3	*	0.2	0.7	1.6	1.8	2.2	3.2	3.3
2000.....	1.5	1.9	0.3	*	0.2	0.6	1.5	1.7	1.9	2.8	3.3
1999.....	1.5	1.6	0.3	*	0.2	0.8	1.5	1.7	2.2	3.0	3.1

Non-Hispanic 4/, male

2007.....	2.5	2.8	0.3	*	*	1.4	3.0	3.2	3.3	3.8	4.3
2006.....	2.4	2.7	0.4	*	0.3	1.3	3.3	3.4	3.1	3.4	4.2
2005.....	2.1	3.6	0.6	*	0.4	1.0	2.5	2.8	2.6	2.9	3.9
2004.....	2.4	3.3	0.3	*	0.3	1.2	2.7	3.0	3.3	4.2	4.5
2003.....	2.5	3.0	*	*	0.3	1.3	3.1	2.9	3.4	4.6	5.3
2002.....	2.3	2.8	0.3	*	0.3	1.1	2.6	3.0	3.4	4.9	4.6
2001.....	2.1	2.7	*	*	0.3	1.0	2.5	2.7	2.9	4.2	4.5
2000.....	2.0	1.6	0.4	*	0.4	1.0	2.2	2.5	2.7	3.9	4.5
1999.....	2.1	1.9	*	*	0.3	1.3	2.4	2.5	3.0	4.2	4.2

Non-Hispanic 4/, female

2007.....	1.5	1.3	*	*	*	0.3	1.0	1.6	2.0	2.1	2.7
2006.....	1.4	1.9	0.3	*	*	0.5	1.1	1.3	1.6	2.3	2.9
2005.....	1.4	1.9	0.4	*	*	0.4	0.8	1.3	1.7	2.1	2.9
2004.....	1.3	2.3	0.3	*	*	0.4	0.9	1.2	1.6	2.2	2.9
2003.....	1.3	1.3	*	*	*	0.5	1.1	1.2	1.7	2.4	3.2
2002.....	1.3	2.6	0.4	*	*	0.4	1.1	1.2	1.6	2.3	2.8
2001.....	1.0	2.0	0.4	*	*	0.3	0.7	1.0	1.5	2.3	2.2
2000.....	0.9	2.2	*	*	*	0.3	0.7	0.9	1.2	1.7	2.1
1999.....	0.9	*	*	*	*	0.3	0.7	0.9	1.5	1.9	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic White, both sexes

2007 5/.....	2.0	1.5	0.3	*	0.2	0.8	2.3	2.8	3.0	3.2	3.7
2006 5/.....	1.9	2.0	0.2	*	0.2	1.0	2.5	2.6	2.7	3.0	3.7
2005 5/.....	1.8	2.2	0.4	*	0.2	0.8	1.9	2.3	2.4	2.7	3.4
2004 5/.....	1.9	2.3	0.3	*	0.2	0.8	2.1	2.4	2.5	3.4	3.9
2003 5/.....	2.0	2.0	*	*	0.2	1.0	2.4	2.3	2.8	3.7	4.4
2002.....	1.8	1.8	0.4	*	0.2	0.8	2.1	2.3	2.6	3.7	3.8
2001.....	1.6	1.7	0.3	*	0.2	0.8	1.8	2.0	2.4	3.3	3.4
2000.....	1.4	1.5	0.2	*	0.2	0.7	1.5	1.8	2.0	2.8	3.2
1999.....	1.4	1.3	*	*	0.2	0.8	1.6	1.7	2.3	3.0	3.0

Non-Hispanic White, male

2007 5/.....	2.5	2.2	*	*	*	1.3	3.4	3.7	3.7	4.1	4.4
2006 5/.....	2.4	2.4	*	*	*	1.5	3.7	3.7	3.5	3.4	4.2
2005 5/.....	2.1	3.0	0.4	*	0.4	1.1	2.8	3.1	2.8	3.1	3.9
2004 5/.....	2.4	2.7	*	*	*	1.2	3.2	3.5	3.3	4.3	4.6
2003 5/.....	2.5	2.4	*	*	*	1.5	3.3	3.1	3.7	4.9	5.4
2002.....	2.3	1.9	*	*	0.3	1.1	2.9	3.2	3.5	4.9	4.7
2001.....	2.1	2.0	*	*	0.3	1.2	2.6	2.8	3.1	4.1	4.4
2000.....	2.0	*	*	*	0.4	1.1	2.2	2.6	2.7	3.9	4.3
1999.....	2.0	*	*	*	0.4	1.4	2.4	2.4	3.0	3.9	4.0

Non-Hispanic White, female

2007 5/.....	1.6	*	*	*	*	0.4	1.2	1.8	2.3	2.3	3.0
2006 5/.....	1.5	*	*	*	*	0.6	1.2	1.6	1.9	2.5	3.2
2005 5/.....	1.5	*	*	*	*	0.5	0.9	1.6	2.0	2.3	2.9
2004 5/.....	1.4	2.0	*	*	*	0.4	1.0	1.4	1.7	2.4	3.2
2003 5/.....	1.4	*	*	*	*	0.6	1.4	1.5	2.0	2.6	3.5
2002.....	1.3	1.8	*	*	*	0.5	1.2	1.3	1.7	2.5	2.9
2001.....	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.5	2.4
2000.....	0.9	*	*	*	*	0.4	0.8	1.1	1.3	1.8	2.0
1999.....	1.0	*	*	*	*	*	0.7	1.1	1.6	2.0	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic Black, both sexes

2007 5/.....	2.0	4.2	*	*	*	1.0	1.2	1.5	1.7	2.7	3.5
2006 5/.....	1.9	3.5	0.8	*	*	*	1.4	1.6	1.7	3.0	3.7
2005 5/.....	1.8	4.9	0.9	*	*	0.6	1.2	1.5	1.9	2.6	4.2
2004 5/.....	1.9	4.9	*	*	*	0.7	1.1	1.0	2.7	3.3	3.6
2003 5/.....	1.8	3.3	*	*	*	*	1.2	1.4	2.2	3.3	4.3
2002.....	2.0	5.7	*	*	*	*	1.3	2.0	2.5	4.1	4.5
2001.....	1.8	5.9	*	*	*	*	1.1	1.5	2.0	3.9	4.0
2000.....	1.9	*	*	*	*	*	1.4	1.7	2.2	3.4	4.9
1999.....	2.0	*	*	*	*	0.8	1.7	2.0	2.6	4.0	4.7

Non-Hispanic Black, male

2007 5/.....	2.8	*	*	*	*	1.8	1.9	2.2	2.1	3.8	5.1
2006 5/.....	2.8	*	*	*	*	*	2.1	2.7	2.4	4.2	5.6
2005 5/.....	2.5	*	*	*	*	*	1.6	2.4	2.4	3.3	5.5
2004 5/.....	2.7	*	*	*	*	*	1.6	*	4.2	4.8	4.8
2003 5/.....	2.8	*	*	*	*	*	2.0	2.5	3.5	4.5	6.3
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.7	5.6
2001.....	2.8	6.4	*	*	*	*	2.0	2.5	2.8	6.0	6.6
2000.....	2.9	*	*	*	*	*	2.6	2.8	3.5	5.0	7.3
1999.....	3.1	*	*	*	*	1.3	2.8	3.6	3.8	6.4	6.7

Non-Hispanic Black, female

2007 5/.....	1.2	*	*	*	*	*	*	*	*	1.8	2.0
2006 5/.....	1.1	*	*	*	*	*	*	*	*	1.9	2.1
2005 5/.....	1.2	*	*	*	*	*	*	*	1.5	2.0	3.1
2004 5/.....	1.1	*	*	*	*	*	*	*	1.4	1.9	2.5
2003 5/.....	1.0	*	*	*	*	*	*	*	*	2.3	2.6
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9	3.4
2001.....	0.9	*	*	*	*	*	*	*	*	2.1	1.8
2000.....	1.0	*	*	*	*	*	*	*	*	2.0	2.8
1999.....	0.9	*	*	*	*	*	*	*	1.5	1.8	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	1.7	2.1	0.3	*	0.1	0.6	1.7	2.0	2.1	2.6	3.1
2006.....	1.6	2.2	0.3	0.1	0.2	0.7	1.8	1.8	2.0	2.5	3.2
2005.....	1.5	2.5	0.4	*	0.2	0.6	1.3	1.7	1.8	2.3	3.0
2004.....	1.6	2.6	0.3	0.1	0.2	0.6	1.5	1.8	2.0	2.8	3.3
2003.....	1.7	1.9	0.2	*	0.1	0.7	1.7	1.7	2.3	3.1	3.8
2002.....	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2	3.3
2001.....	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9	3.1
2000.....	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6	3.0
1999.....	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7	2.8

All origins 3/, male

2007.....	2.1	2.5	0.3	*	*	1.0	2.5	2.5	2.6	3.2	3.8
2006.....	2.1	2.7	0.3	*	0.2	0.9	2.6	2.6	2.6	3.0	3.8
2005.....	1.8	3.2	0.5	*	0.2	0.8	1.9	2.2	2.2	2.6	3.5
2004.....	2.1	3.1	0.3	*	0.2	0.9	2.1	2.4	2.7	3.7	3.9
2003.....	2.2	2.4	*	*	0.2	0.9	2.4	2.4	3.0	4.1	4.8
2002.....	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3	4.1
2001.....	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8	4.2
2000.....	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6	4.1
1999.....	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7	3.8

All origins 3/, female

2007.....	1.3	1.6	0.3	*	*	0.3	0.9	1.3	1.6	1.9	2.4
2006.....	1.2	1.7	0.3	*	*	0.4	1.0	1.1	1.3	2.0	2.6
2005.....	1.2	1.8	0.3	*	*	0.4	0.8	1.1	1.5	1.9	2.6
2004.....	1.2	2.0	0.3	*	*	0.3	0.8	1.1	1.3	1.9	2.6
2003.....	1.2	1.3	*	*	*	0.4	1.0	1.1	1.5	2.1	2.8
2002.....	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0	2.6
2001.....	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6	1.9
1999.....	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	1.9	2.1	0.3	*	0.1	0.7	1.9	2.2	2.5	2.9	3.4
2006.....	1.8	2.3	0.3	0.1	0.2	0.8	2.0	2.2	2.3	2.7	3.5
2005.....	1.7	2.7	0.4	*	0.2	0.6	1.4	1.9	2.1	2.4	3.3
2004.....	1.7	2.8	0.3	*	0.2	0.7	1.6	2.0	2.3	3.1	3.5
2003.....	1.8	2.2	0.2	*	0.1	0.7	1.9	1.9	2.4	3.4	4.1
2002.....	1.7	2.7	0.3	0.1	0.2	0.6	1.6	2.0	2.4	3.4	3.6
2001.....	1.5	2.4	0.3	*	0.2	0.6	1.4	1.6	2.1	3.1	3.2
2000.....	1.4	1.9	0.3	*	0.2	0.5	1.2	1.5	1.8	2.7	3.2
1999.....	1.4	1.6	0.3	*	0.1	0.5	1.2	1.5	2.1	2.9	3.0

Non-Hispanic 4/, male

2007.....	2.3	2.8	0.3	*	*	1.0	2.8	2.9	3.2	3.7	4.1
2006.....	2.3	2.7	0.3	*	0.2	1.0	3.0	3.1	3.0	3.2	4.1
2005.....	2.0	3.6	0.5	*	0.3	0.8	2.1	2.6	2.5	2.8	3.8
2004.....	2.2	3.3	0.3	*	0.2	1.0	2.4	2.8	3.1	4.1	4.3
2003.....	2.3	3.0	*	*	*	1.0	2.7	2.7	3.3	4.5	5.2
2002.....	2.2	2.8	0.3	*	0.3	0.8	2.2	2.8	3.2	4.6	4.4
2001.....	2.0	2.7	*	*	0.2	0.8	2.0	2.3	2.8	4.0	4.4
2000.....	1.9	1.6	0.4	*	0.3	0.7	1.8	2.2	2.5	3.8	4.4
1999.....	1.9	1.9	*	*	*	0.7	1.8	2.1	2.8	3.9	4.0

Non-Hispanic 4/, female

2007.....	1.4	1.3	*	*	*	0.3	1.0	1.6	1.9	2.1	2.7
2006.....	1.3	1.9	0.3	*	*	0.5	1.0	1.3	1.6	2.2	2.9
2005.....	1.3	1.9	0.3	*	*	0.4	0.8	1.3	1.7	2.1	2.8
2004.....	1.3	2.3	0.3	*	*	0.3	0.9	1.2	1.5	2.1	2.8
2003.....	1.3	1.3	*	*	*	0.5	1.1	1.2	1.6	2.3	3.1
2002.....	1.2	2.6	0.4	*	*	0.4	1.0	1.1	1.5	2.2	2.8
2001.....	1.0	2.0	0.4	*	*	0.3	0.7	0.9	1.5	2.2	2.1
2000.....	0.9	2.2	*	*	*	0.3	0.6	0.9	1.1	1.7	2.0
1999.....	0.9	*	*	*	*	*	0.6	0.9	1.4	1.8	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic White, both sexes

2007 5/.....	2.0	1.5	0.3	*	*	0.8	2.2	2.6	3.0	3.1	3.6
2006 5/.....	1.9	2.0	*	*	0.2	0.9	2.3	2.5	2.6	2.8	3.6
2005 5/.....	1.7	2.2	0.4	*	0.2	0.7	1.6	2.2	2.3	2.6	3.3
2004 5/.....	1.8	2.3	0.3	*	0.2	0.7	1.9	2.4	2.4	3.3	3.7
2003 5/.....	1.9	2.0	*	*	*	0.9	2.2	2.2	2.7	3.6	4.3
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	2.2	2.5	3.6	3.7
2001.....	1.5	1.7	0.3	*	0.2	0.7	1.6	1.8	2.3	3.1	3.3
2000.....	1.4	1.5	0.2	*	0.2	0.6	1.3	1.7	1.9	2.7	3.1
1999.....	1.3	1.3	*	*	*	0.6	1.3	1.6	2.2	2.8	2.8

Non-Hispanic White, male

2007 5/.....	2.4	2.2	*	*	*	1.2	3.1	3.4	3.6	3.9	4.2
2006 5/.....	2.3	2.4	*	*	*	1.3	3.4	3.5	3.4	3.2	4.1
2005 5/.....	2.0	3.0	0.4	*	*	0.9	2.4	2.9	2.7	3.0	3.7
2004 5/.....	2.3	2.7	*	*	*	1.1	2.8	3.3	3.2	4.2	4.4
2003 5/.....	2.4	2.4	*	*	*	1.2	3.1	3.0	3.5	4.7	5.2
2002.....	2.2	1.9	*	*	*	0.9	2.6	3.0	3.3	4.7	4.5
2001.....	2.0	2.0	*	*	*	1.0	2.2	2.5	3.0	3.9	4.3
2000.....	1.8	*	*	*	0.3	0.9	1.9	2.3	2.5	3.8	4.1
1999.....	1.8	*	*	*	*	0.9	1.9	2.1	2.8	3.7	3.8

Non-Hispanic White, female

2007 5/.....	1.6	*	*	*	*	0.3	1.1	1.8	2.3	2.3	2.9
2006 5/.....	1.4	*	*	*	*	0.5	1.2	1.5	1.8	2.4	3.2
2005 5/.....	1.4	*	*	*	*	0.4	0.9	1.5	1.9	2.2	2.9
2004 5/.....	1.4	2.0	*	*	*	0.3	1.0	1.4	1.6	2.3	3.1
2003 5/.....	1.4	*	*	*	*	0.6	1.3	1.5	1.9	2.5	3.4
2002.....	1.3	1.8	*	*	*	0.5	1.1	1.3	1.7	2.5	2.8
2001.....	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.3	2.3
2000.....	0.9	*	*	*	*	0.3	0.7	1.0	1.3	1.7	2.0
1999.....	0.9	*	*	*	*	*	0.6	1.1	1.5	1.9	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic Black, both sexes

2007 5/.....	1.8	4.2	*	*	*	*	0.9	1.3	1.6	2.7	3.5
2006 5/.....	1.8	3.5	0.8	*	*	*	1.0	1.2	1.6	2.8	3.6
2005 5/.....	1.7	4.9	0.8	*	*	*	0.9	1.3	1.9	2.5	4.1
2004 5/.....	1.8	4.9	*	*	*	*	0.9	0.8	2.6	3.2	3.5
2003 5/.....	1.8	3.3	*	*	*	*	0.8	1.3	2.0	3.3	4.3
2002.....	1.9	5.7	*	*	*	*	1.1	1.8	2.4	4.0	4.4
2001.....	1.7	5.9	*	*	*	*	0.8	1.3	2.0	3.9	4.0
2000.....	1.7	*	*	*	*	*	0.9	1.4	2.0	3.3	4.8
1999.....	1.8	*	*	*	*	*	1.2	1.6	2.4	3.9	4.6

Non-Hispanic Black, male

2007 5/.....	2.6	*	*	*	*	*	1.7	1.8	1.8	3.7	5.1
2006 5/.....	2.5	*	*	*	*	*	1.3	1.9	2.3	3.8	5.5
2005 5/.....	2.4	*	*	*	*	*	*	2.1	2.4	3.1	5.4
2004 5/.....	2.5	*	*	*	*	*	*	*	3.9	4.7	4.7
2003 5/.....	2.6	*	*	*	*	*	1.4	2.1	3.3	4.5	6.2
2002.....	2.7	*	*	*	*	*	1.4	2.7	3.3	6.4	5.5
2001.....	2.6	6.4	*	*	*	*	1.5	2.3	2.7	6.0	6.5
2000.....	2.6	*	*	*	*	*	1.6	2.2	3.1	4.7	7.2
1999.....	2.8	*	*	*	*	*	1.8	2.9	3.5	6.2	6.6

Non-Hispanic Black, female

2007 5/.....	1.1	*	*	*	*	*	*	*	*	1.8	2.0
2006 5/.....	1.0	*	*	*	*	*	*	*	*	1.9	2.0
2005 5/.....	1.1	*	*	*	*	*	*	*	1.5	2.0	3.0
2004 5/.....	1.1	*	*	*	*	*	*	*	*	1.9	2.5
2003 5/.....	1.0	*	*	*	*	*	*	*	*	2.2	2.6
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.8	3.4
2001.....	0.9	*	*	*	*	*	*	*	*	2.1	1.7
2000.....	0.9	*	*	*	*	*	*	*	*	2.0	2.6
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.8	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/, both sexes

2007.....	0.9	0.5	0.1	*	*	*	*	0.2	0.2	0.3	0.4
2006.....	0.8	0.6	0.1	*	*	0.1	0.1	0.1	0.2	0.2	0.4
2005.....	0.9	*	*	*	*	*	*	0.2	0.2	0.3	0.4
2004.....	1.0	*	*	*	*	0.1	0.1	0.2	0.2	0.3	0.4
2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4	0.5
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3	0.4
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3	0.4
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3	0.4
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3	0.4

All origins 3/, male

2007.....	0.8	*	*	*	*	*	*	*	*	0.2	0.4
2006.....	0.8	*	*	*	*	*	*	*	*	*	0.3
2005.....	0.8	*	*	*	*	*	*	*	*	0.2	0.4
2004.....	0.9	*	*	*	*	*	*	*	*	0.2	0.4
2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3	0.4
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3	0.4
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3	0.4

All origins 3/, female

2007.....	0.9	*	*	*	*	*	*	*	0.2	0.3	0.4
2006.....	0.9	*	*	*	*	*	*	0.2	*	0.2	0.4
2005.....	1.0	*	*	*	*	*	*	*	0.3	0.3	0.4
2004.....	1.0	*	*	*	*	*	*	*	0.3	0.3	0.5
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3	0.4
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

All origins 3/, both sexes

2007.....	7.7	*	*	*	*	0.3	0.7	1.3	2.4	4.7	9.7
2006.....	7.4	*	*	*	*	0.2	0.5	1.0	2.2	4.6	10.1
2005.....	7.3	*	*	*	*	0.2	0.5	1.0	1.7	4.7	10.1
2004.....	7.2	*	*	*	*	0.2	0.5	0.9	2.1	5.0	10.0
2003.....	7.1	*	*	*	*	0.2	0.4	0.9	2.1	5.3	10.6
2002.....	7.0	*	*	*	*	0.2	0.4	0.8	2.1	5.4	10.6
2001.....	7.1	*	*	*	*	0.2	0.3	0.9	2.3	6.0	10.6
2000.....	7.0	*	*	*	*	0.2	0.3	0.9	2.3	6.0	11.1
1999.....	7.0	*	*	*	*	0.1	0.4	0.8	2.4	6.2	10.9

All origins 3/, male

2007.....	11.7	*	*	*	*	0.3	1.0	1.8	3.6	6.7	13.6
2006.....	11.2	*	*	*	*	0.4	0.7	1.7	3.2	6.3	14.1
2005.....	11.1	*	*	*	*	0.3	0.9	1.5	2.5	6.6	14.4
2004.....	11.0	*	*	*	*	0.2	0.8	1.4	3.2	7.1	14.6
2003.....	10.9	*	*	*	*	0.4	0.6	1.4	3.2	7.4	15.2
2002.....	10.8	*	*	*	*	0.3	0.6	1.4	3.1	7.6	15.3
2001.....	10.8	*	*	*	*	0.3	0.6	1.3	3.3	8.5	15.3
2000.....	10.9	*	*	*	*	0.3	0.5	1.5	3.5	8.7	16.4
1999.....	10.9	*	*	*	*	0.2	0.6	1.2	3.3	8.7	16.6

All origins 3/, female

2007.....	3.8	*	*	*	*	*	0.3	0.6	1.2	2.7	5.8
2006.....	3.7	*	*	*	*	*	0.2	0.4	1.1	2.8	6.2
2005.....	3.6	*	*	*	*	*	*	0.4	1.0	2.7	5.8
2004.....	3.5	*	*	*	*	*	*	0.4	1.0	2.9	5.3
2003.....	3.4	*	*	*	*	*	*	0.3	0.9	3.2	6.0
2002.....	3.4	*	*	*	*	*	*	0.3	1.1	3.3	6.1
2001.....	3.4	*	*	*	*	*	*	0.5	1.3	3.5	6.0
2000.....	3.2	*	*	*	*	*	*	0.3	1.1	3.2	5.8
1999.....	3.2	*	*	*	*	*	*	0.3	1.5	3.7	5.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic 4/, both sexes

2007.....	7.9	*	*	*	*	0.2	0.6	1.1	2.2	4.6	9.6
2006.....	7.5	*	*	*	*	0.2	0.5	0.9	1.9	4.4	9.9
2005.....	7.4	*	*	*	*	0.1	0.5	0.8	1.5	4.5	9.8
2004.....	7.4	*	*	*	*	0.2	0.4	0.8	2.0	4.9	9.8
2003.....	7.2	*	*	*	*	0.2	0.3	0.8	1.9	5.2	10.5
2002.....	7.1	*	*	*	*	0.1	0.3	0.7	2.0	5.3	10.4
2001.....	7.1	*	*	*	*	0.2	0.3	0.7	2.2	5.9	10.3
2000.....	7.0	*	*	*	*	0.2	0.2	0.8	2.2	5.7	10.6
1999.....	7.0	*	*	*	*	*	0.3	0.7	2.2	5.9	10.5

Non-Hispanic 4/, male

2007.....	11.8	*	*	*	*	0.3	0.9	1.6	3.1	6.4	13.0
2006.....	11.3	*	*	*	*	0.3	0.7	1.4	2.7	5.9	13.5
2005.....	11.2	*	*	*	*	*	0.7	1.2	2.1	6.2	13.7
2004.....	11.2	*	*	*	*	*	0.6	1.3	2.8	6.8	14.1
2003.....	11.0	*	*	*	*	0.3	0.5	1.3	3.0	6.9	14.9
2002.....	10.7	*	*	*	*	0.2	0.4	1.1	2.7	7.1	14.5
2001.....	10.8	*	*	*	*	0.2	0.5	1.1	3.0	8.1	14.6
2000.....	10.7	*	*	*	*	0.2	0.3	1.3	3.1	8.2	15.4
1999.....	10.8	*	*	*	*	*	0.5	1.1	2.9	8.1	15.7

Non-Hispanic 4/, female

2007.....	4.1	*	*	*	*	*	0.3	0.7	1.3	2.9	6.3
2006.....	3.9	*	*	*	*	*	0.3	0.4	1.1	2.9	6.4
2005.....	3.8	*	*	*	*	*	*	0.4	1.0	2.8	6.0
2004.....	3.7	*	*	*	*	*	*	0.4	1.2	3.1	5.6
2003.....	3.6	*	*	*	*	*	*	0.3	0.9	3.5	6.2
2002.....	3.6	*	*	*	*	*	*	0.3	1.2	3.5	6.4
2001.....	3.6	*	*	*	*	*	*	0.4	1.4	3.8	6.2
2000.....	3.4	*	*	*	*	*	*	0.3	1.3	3.3	6.0
1999.....	3.4	*	*	*	*	*	*	0.3	1.6	3.8	5.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic White, both sexes

2007 5/.....	8.4	*	*	*	*	0.3	0.7	1.1	2.3	4.9	10.1
2006 5/.....	8.0	*	*	*	*	0.2	0.5	0.8	1.9	4.4	10.3
2005 5/.....	7.9	*	*	*	*	*	0.4	0.8	1.5	4.6	10.0
2004 5/.....	7.7	*	*	*	*	0.2	0.4	0.8	2.0	5.1	10.1
2003 5/.....	7.5	*	*	*	*	0.2	0.3	0.8	1.9	5.2	10.7
2002.....	7.3	*	*	*	*	0.2	0.3	0.7	1.9	5.2	10.6
2001.....	7.2	*	*	*	*	*	0.2	0.6	2.1	5.9	10.0
2000.....	7.0	*	*	*	*	0.2	0.2	0.8	2.0	5.6	10.2
1999.....	6.9	*	*	*	*	*	0.3	0.7	2.1	5.6	9.9

Non-Hispanic White, male

2007 5/.....	12.6	*	*	*	*	0.4	0.9	1.6	3.3	6.7	13.5
2006 5/.....	12.0	*	*	*	*	0.4	0.7	1.3	2.8	6.0	14.0
2005 5/.....	11.8	*	*	*	*	*	0.7	1.2	2.0	6.4	14.0
2004 5/.....	11.7	*	*	*	*	*	0.6	1.1	2.8	6.8	14.5
2003 5/.....	11.4	*	*	*	*	*	0.5	1.3	2.9	6.9	15.2
2002.....	11.1	*	*	*	*	*	0.4	1.0	2.6	7.0	14.8
2001.....	10.8	*	*	*	*	*	0.4	1.0	2.9	7.8	13.9
2000.....	10.8	*	*	*	*	*	*	1.2	2.8	8.0	14.8
1999.....	10.7	*	*	*	*	*	0.5	1.1	2.7	7.6	14.9

Non-Hispanic White, female

2007 5/.....	4.4	*	*	*	*	*	0.4	0.7	1.2	3.2	6.7
2006 5/.....	4.2	*	*	*	*	*	*	0.4	1.0	2.8	6.6
2005 5/.....	4.0	*	*	*	*	*	*	0.5	0.9	2.8	6.0
2004 5/.....	3.9	*	*	*	*	*	*	0.4	1.1	3.3	5.7
2003 5/.....	3.8	*	*	*	*	*	*	*	1.0	3.4	6.2
2002.....	3.7	*	*	*	*	*	*	*	1.1	3.4	6.4
2001.....	3.6	*	*	*	*	*	*	*	1.3	3.9	6.1
2000.....	3.4	*	*	*	*	*	*	*	1.2	3.2	5.7
1999.....	3.3	*	*	*	*	*	*	*	1.5	3.5	4.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic Black, both sexes

2007 5/.....	5.8	*	*	*	*	*	*	*	1.4	2.9	6.7
2006 5/.....	5.6	*	*	*	*	*	*	0.7	1.3	3.4	7.4
2005 5/.....	6.1	*	*	*	*	*	*	*	1.2	3.1	8.2
2004 5/.....	6.3	*	*	*	*	*	*	*	1.4	3.7	7.8
2003 5/.....	6.5	*	*	*	*	*	*	0.9	1.4	5.0	9.0
2002.....	6.6	*	*	*	*	*	*	*	2.0	4.5	9.2
2001.....	7.5	*	*	*	*	*	*	0.8	1.9	5.3	12.5
2000.....	7.6	*	*	*	*	*	*	0.9	2.6	5.6	13.1
1999.....	8.0	*	*	*	*	*	*	*	2.2	7.3	13.2

Non-Hispanic Black, male

2007 5/.....	9.2	*	*	*	*	*	*	*	1.9	4.1	9.3
2006 5/.....	8.7	*	*	*	*	*	*	*	*	4.3	9.9
2005 5/.....	9.4	*	*	*	*	*	*	*	*	4.3	11.2
2004 5/.....	10.0	*	*	*	*	*	*	*	2.2	5.9	11.9
2003 5/.....	10.3	*	*	*	*	*	*	*	2.3	6.8	13.0
2002.....	10.3	*	*	*	*	*	*	*	3.0	6.3	12.9
2001.....	11.9	*	*	*	*	*	*	*	2.6	8.5	18.6
2000.....	11.7	*	*	*	*	*	*	1.6	3.5	8.3	19.3
1999.....	12.5	*	*	*	*	*	*	*	2.9	10.5	19.2

Non-Hispanic Black, female

2007 5/.....	2.7	*	*	*	*	*	*	*	*	1.7	4.3
2006 5/.....	2.9	*	*	*	*	*	*	*	*	2.6	5.1
2005 5/.....	3.1	*	*	*	*	*	*	*	*	2.2	5.6
2004 5/.....	2.9	*	*	*	*	*	*	*	*	1.8	4.2
2003 5/.....	3.0	*	*	*	*	*	*	*	*	3.3	5.5
2002.....	3.3	*	*	*	*	*	*	*	*	2.9	6.0
2001.....	3.5	*	*	*	*	*	*	*	*	2.5	7.1
2000.....	3.9	*	*	*	*	*	*	*	1.8	3.3	7.6
1999.....	4.0	*	*	*	*	*	*	*	1.6	4.5	7.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

All origins 3/, both sexes

2007	12.7	0.8	0.3	*	0.3	4.5	12.9	16.8	17.4	19.9	24.8
2006	12.8	1.1	0.2	*	0.2	4.1	13.1	16.5	17.1	20.2	26.2
2005	11.3	0.9	0.2	*	0.2	3.7	11.0	13.5	15.1	18.3	24.0
2004	10.5	0.7	0.2	*	0.2	3.9	10.0	12.2	13.1	18.4	23.2
2003	9.9	0.6	0.2	*	0.2	3.3	9.4	10.9	13.3	18.3	22.8
2002	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5	21.8
2001	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5	18.5
2000	7.0	*	*	*	0.2	2.2	5.9	7.5	10.1	14.4	17.5
1999	6.9	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2	17.4

All origins 3/, male

2007	16.1	*	0.2	*	0.3	6.6	18.8	23.6	23.4	25.3	30.3
2006	16.6	1.1	0.2	*	0.2	6.0	19.5	23.3	23.5	25.8	32.9
2005	14.5	*	0.3	*	*	5.4	16.0	18.9	20.0	23.0	29.9
2004	13.4	*	*	*	0.2	5.6	14.7	17.3	17.6	23.6	28.3
2003	12.9	*	0.3	*	*	4.5	13.9	15.4	17.7	23.8	29.1
2002	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1	28.0
2001	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7	24.6
2000	9.5	*	*	*	*	3.1	8.8	10.8	14.1	19.8	23.8
1999	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6	24.3

All origins 3/, female

2007	9.5	*	0.3	*	0.2	2.2	6.7	9.8	11.3	14.5	19.3
2006	9.1	1.0	*	*	*	2.2	6.3	9.3	10.7	14.5	19.6
2005	8.2	*	*	*	0.2	2.0	5.6	7.7	10.1	13.5	18.2
2004	7.6	*	*	*	0.3	2.1	5.0	6.7	8.5	13.2	18.2
2003	7.0	*	*	*	*	1.9	4.7	6.2	8.8	12.8	16.6
2002	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9	15.8
2001	5.1	*	*	*	*	1.6	3.2	4.4	6.7	10.3	12.6
2000	4.6	*	*	*	*	1.4	2.9	4.1	6.1	9.0	11.4
1999	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.8	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Hispanic, both sexes

2007	6.0	*	*	*	*	2.8	5.5	7.2	7.3	10.2	12.9
2006	6.5	*	*	*	*	2.1	5.2	6.6	9.6	11.0	15.7
2005	6.1	*	*	*	*	2.2	5.6	6.6	8.2	10.4	14.9
2004	5.5	*	*	*	*	2.0	4.5	6.0	6.7	10.7	13.7
2003	5.9	*	*	*	*	2.1	4.7	6.0	8.5	12.0	14.9
2002	5.5	*	*	*	*	1.8	4.4	5.0	7.5	11.3	15.8
2001	4.7	*	*	*	*	1.7	3.1	5.0	7.0	11.1	12.7
2000	4.8	*	*	*	*	1.3	3.2	5.0	8.9	12.2	13.3
1999	5.8	*	*	*	*	1.6	3.8	6.5	9.9	14.7	15.5

Hispanic, male

2007	8.7	*	*	*	*	3.8	8.1	10.5	10.1	14.2	18.8
2006	9.3	*	*	*	*	2.9	7.2	9.4	13.7	15.2	22.7
2005	8.9	*	*	*	*	2.9	8.2	9.7	11.9	14.9	21.4
2004	7.8	*	*	*	*	2.8	6.1	8.7	9.9	15.3	19.3
2003	8.7	*	*	*	*	3.0	6.9	8.8	12.3	17.7	21.8
2002	8.2	*	*	*	*	2.6	6.4	7.8	11.0	16.4	24.1
2001	7.0	*	*	*	*	2.3	4.1	7.6	9.8	16.6	19.3
2000	7.4	*	*	*	*	1.8	4.9	7.1	14.2	17.9	21.8
1999	9.2	*	*	*	*	2.1	6.1	9.9	15.8	22.5	25.4

Hispanic, female

2007	3.1	*	*	*	*	1.7	2.5	3.1	3.9	5.5	6.3
2006	3.4	*	*	*	*	1.2	2.8	3.1	4.8	6.1	7.9
2005	3.0	*	*	*	*	1.4	2.4	2.7	3.8	5.3	7.7
2004	2.9	*	*	*	*	*	2.4	2.6	2.9	5.5	7.6
2003	2.9	*	*	*	*	*	1.9	2.5	4.2	5.6	7.6
2002	2.6	*	*	*	*	*	1.9	1.7	3.4	5.8	7.0
2001	2.2	*	*	*	*	*	1.8	1.8	3.8	5.0	5.7
2000	2.1	*	*	*	*	*	*	2.6	2.9	6.0	4.4
1999	2.2	*	*	*	*	*	*	2.5	3.2	6.1	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic 4/, both sexes

2007	13.9	0.8	0.3	*	0.3	4.8	14.5	19.2	20.0	21.9	26.7
2006	13.9	1.1	0.3	*	0.2	4.5	14.8	18.9	18.9	22.0	27.7
2005	12.1	0.9	0.2	*	0.2	4.0	12.1	15.1	16.6	19.7	25.3
2004	11.2	0.9	0.2	*	0.3	4.3	11.2	13.7	14.5	19.7	24.5
2003	10.4	0.7	0.2	*	0.1	3.5	10.4	12.0	14.3	19.3	23.8
2002	9.5	0.8	0.2	*	0.2	3.3	8.6	10.9	13.3	18.4	22.5
2001	8.0	0.7	0.2	*	0.2	2.8	7.2	8.9	11.2	16.0	19.1
2000	7.2	*	*	*	0.2	2.4	6.4	7.9	10.2	14.6	17.9
1999	6.9	*	0.2	*	0.1	1.9	5.6	7.6	10.3	14.0	17.3

Non-Hispanic 4/, male

2007	17.4	*	0.3	*	0.4	7.2	21.1	27.1	27.1	27.8	32.3
2006	17.9	*	*	*	0.3	6.6	22.3	27.2	26.0	28.0	34.4
2005	15.4	*	0.3	*	*	5.9	17.7	21.4	21.9	24.6	31.1
2004	14.3	*	*	*	*	6.1	16.7	19.7	19.3	25.0	29.6
2003	13.5	*	*	*	*	4.8	15.5	17.1	18.8	24.7	30.0
2002	12.3	*	*	*	*	4.6	12.5	15.6	18.2	24.2	28.3
2001	10.6	*	*	*	*	3.9	10.8	12.9	15.2	21.2	25.0
2000	9.7	*	*	*	*	3.3	9.6	11.6	14.0	20.0	23.8
1999	9.3	*	*	*	*	2.5	8.0	10.9	14.3	19.0	23.8

Non-Hispanic 4/, female

2007	10.5	*	0.3	*	*	2.3	7.5	11.2	13.0	16.1	21.3
2006	10.1	*	*	*	*	2.4	7.0	10.7	11.9	16.1	21.3
2005	9.0	*	*	*	*	2.1	6.3	8.8	11.4	14.9	19.7
2004	8.3	*	*	*	0.3	2.3	5.5	7.6	9.6	14.4	19.6
2003	7.6	*	*	*	*	2.1	5.2	6.9	9.7	14.0	17.7
2002	6.9	*	*	*	*	1.9	4.6	6.2	8.6	12.8	16.8
2001	5.5	*	*	*	*	1.6	3.5	4.9	7.2	11.0	13.3
2000	4.9	*	*	*	*	1.5	3.2	4.3	6.6	9.4	12.1
1999	4.6	*	*	*	*	1.2	3.1	4.2	6.4	9.0	11.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic White, both sexes

2007 5/.....	15.3	*	*	*	0.3	6.0	17.7	23.5	24.0	24.8	29.6
2006 5/.....	14.9	*	0.2	*	0.3	5.7	18.2	23.0	22.0	24.2	29.5
2005 5/.....	13.1	0.9	*	*	0.2	4.9	14.9	17.9	19.2	21.8	27.0
2004 5/.....	12.1	*	*	*	0.3	5.3	13.7	16.4	16.6	21.5	26.5
2003 5/.....	11.2	*	*	*	*	4.3	12.8	14.2	16.1	20.7	25.0
2002.....	10.0	*	*	*	0.2	3.9	10.4	12.6	14.9	19.7	23.4
2001.....	8.3	*	*	*	0.2	3.3	8.6	10.3	12.2	16.8	19.3
2000.....	7.4	*	*	*	0.2	2.9	7.5	8.9	10.9	15.0	17.6
1999.....	6.9	*	*	*	*	2.2	6.3	8.2	10.7	14.2	16.7

Non-Hispanic White, male

2007 5/.....	19.0	*	*	*	0.4	8.9	25.9	33.2	32.4	31.1	35.5
2006 5/.....	18.9	*	*	*	*	8.3	27.6	32.7	29.6	30.3	36.1
2005 5/.....	16.4	*	*	*	*	7.3	21.9	25.3	25.2	27.0	33.0
2004 5/.....	15.3	*	*	*	*	7.6	20.5	23.6	22.1	27.3	31.5
2003 5/.....	14.3	*	*	*	*	6.0	19.1	20.4	21.2	26.3	31.4
2002.....	12.8	*	*	*	*	5.5	15.0	17.9	20.3	25.6	29.3
2001.....	10.8	*	*	*	*	4.8	12.7	14.7	16.7	22.1	25.2
2000.....	9.8	*	*	*	*	4.0	11.3	12.8	14.8	20.6	23.2
1999.....	9.2	*	*	*	*	2.9	9.0	11.5	14.8	19.1	22.8

Non-Hispanic White, female

2007 5/.....	11.7	*	*	*	*	2.9	8.9	13.7	15.6	18.5	23.7
2006 5/.....	11.1	*	*	*	*	2.9	8.4	13.2	14.2	18.0	23.0
2005 5/.....	9.8	*	*	*	*	2.4	7.5	10.4	13.0	16.5	21.1
2004 5/.....	9.1	*	*	*	*	2.8	6.7	9.1	10.9	15.7	21.5
2003 5/.....	8.2	*	*	*	*	2.5	6.3	7.9	10.9	15.0	18.7
2002.....	7.4	*	*	*	*	2.3	5.6	7.3	9.4	13.8	17.5
2001.....	5.8	*	*	*	*	1.8	4.3	5.8	7.6	11.5	13.5
2000.....	5.1	*	*	*	*	1.7	3.7	4.9	6.9	9.5	12.1
1999.....	4.8	*	*	*	*	1.4	3.5	4.7	6.6	9.2	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic Black, both sexes

2007 5/.....	10.8	*	*	*	*	1.0	4.2	7.3	10.4	16.4	21.2
2006 5/.....	12.5	*	*	*	*	0.9	4.5	8.5	12.6	19.5	27.9
2005 5/.....	10.8	*	*	*	*	1.0	3.4	7.7	11.8	16.5	24.2
2004 5/.....	9.7	*	*	*	*	1.0	3.5	6.4	9.9	16.8	21.7
2003 5/.....	9.5	*	*	*	*	0.7	3.0	5.8	10.7	18.4	24.2
2002.....	9.4	*	*	*	*	1.0	3.3	6.8	10.6	17.7	24.6
2001.....	8.7	*	*	*	*	0.9	3.1	5.9	9.7	17.0	23.9
2000.....	8.5	*	*	*	*	0.9	3.2	6.2	10.7	16.9	24.5
1999.....	8.7	*	*	*	*	0.8	3.9	7.1	11.5	16.7	26.1

Non-Hispanic Black, male

2007 5/.....	14.6	*	*	*	*	1.5	5.8	10.1	14.4	22.0	26.8
2006 5/.....	17.8	*	*	*	*	*	6.1	13.3	19.4	26.9	37.9
2005 5/.....	14.7	*	*	*	*	*	4.8	11.4	15.1	20.5	30.6
2004 5/.....	13.1	*	*	*	*	1.4	4.9	9.3	13.6	21.1	27.8
2003 5/.....	13.0	*	*	*	*	*	4.6	7.7	14.0	24.9	31.8
2002.....	13.2	*	*	*	*	1.4	4.8	10.1	14.3	24.7	31.8
2001.....	12.5	*	*	*	*	*	5.2	9.4	12.6	22.7	32.4
2000.....	12.3	*	*	*	*	*	4.7	9.9	14.7	23.2	34.5
1999.....	12.9	*	*	*	*	*	5.9	11.4	16.1	24.1	37.2

Non-Hispanic Black, female

2007 5/.....	7.3	*	*	*	*	*	2.5	4.7	6.8	11.5	16.3
2006 5/.....	7.6	*	*	*	*	*	2.9	4.0	6.5	12.9	19.2
2005 5/.....	7.2	*	*	*	*	*	2.0	4.2	8.9	13.0	18.6
2004 5/.....	6.6	*	*	*	*	*	2.1	3.8	6.6	13.0	16.3
2003 5/.....	6.3	*	*	*	*	*	1.5	4.1	7.7	12.7	17.6
2002.....	6.0	*	*	*	*	*	1.9	3.8	7.2	11.4	18.3
2001.....	5.3	*	*	*	*	*	*	2.7	7.2	11.9	16.4
2000.....	5.0	*	*	*	*	*	1.9	3.0	7.0	11.3	15.5
1999.....	4.9	*	*	*	*	*	2.0	3.2	7.5	10.2	16.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

All origins 3/, both sexes

2007.....	10.4	*	0.4	0.3	1.2	12.4	20.1	16.6	14.4	11.8	11.9
2006.....	10.3	*	0.3	0.4	1.3	13.2	20.6	17.0	13.5	11.8	11.1
2005.....	10.4	*	0.4	0.3	1.3	12.5	20.0	17.2	14.2	11.6	11.6
2004.....	10.1	*	0.3	0.3	1.1	12.0	19.3	16.9	13.3	11.2	11.4
2003.....	10.4	*	0.3	0.3	1.2	12.1	21.1	17.4	13.8	11.6	11.5
2002.....	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4	11.9
2001.....	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2	11.3
2000.....	10.2	*	0.3	0.3	1.5	12.9	20.9	16.3	12.8	12.0	11.8
1999.....	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3	10.9

All origins 3/, male

2007.....	18.2	*	0.5	0.4	1.9	21.8	35.3	28.9	24.9	19.7	19.3
2006.....	18.1	*	0.5	0.4	2.0	23.3	36.2	29.4	23.6	19.5	18.3
2005.....	18.3	*	0.5	0.4	2.0	22.0	35.3	30.2	24.3	19.3	19.2
2004.....	17.6	*	0.4	0.3	1.8	20.7	34.2	29.8	22.5	19.0	18.5
2003.....	18.3	*	0.3	0.4	2.0	21.2	37.1	30.7	23.7	19.3	18.9
2002.....	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6	19.7
2001.....	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0	18.6
2000.....	17.8	*	0.4	0.4	2.4	22.4	37.0	28.5	21.4	19.5	19.4
1999.....	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0	17.7

All origins 3/, female

2007.....	2.7	*	0.4	0.3	0.5	2.6	3.9	3.6	3.6	3.8	4.6
2006.....	2.8	*	*	0.4	0.5	2.5	3.8	4.1	3.2	4.1	4.1
2005.....	2.7	*	0.3	0.3	0.6	2.4	3.6	3.7	3.8	3.9	4.1
2004.....	2.7	*	0.3	0.3	0.4	2.9	3.5	3.4	3.9	3.3	4.4
2003.....	2.7	*	0.3	0.2	0.4	2.4	4.2	3.6	3.7	3.8	4.2
2002.....	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2	4.2
2001.....	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6	4.1
2000.....	2.8	*	*	0.2	0.5	2.9	4.2	3.9	4.1	4.5	4.3
1999.....	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6	4.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Hispanic, both sexes

2007.....	7.7	*	*	*	1.2	14.5	21.3	14.2	11.1	7.8	7.2
2006.....	7.8	*	*	*	1.5	16.5	21.5	14.6	10.5	8.4	7.2
2005.....	8.1	*	*	*	1.0	16.3	21.1	15.7	11.9	9.1	7.4
2004.....	7.9	*	*	*	0.9	15.8	21.0	15.1	10.2	7.9	7.5
2003.....	8.3	*	*	*	1.1	15.3	22.6	15.8	10.8	8.8	7.3
2002.....	8.1	*	*	*	1.1	16.1	20.7	15.2	9.7	8.5	7.2
2001.....	8.3	*	*	*	1.0	14.8	21.7	16.2	10.6	8.9	8.2
2000.....	8.2	*	*	*	1.4	16.1	19.7	13.9	10.2	7.5	8.7
1999.....	8.5	*	*	*	1.9	17.4	20.2	14.3	10.0	9.2	8.0

Hispanic, male

2007.....	13.4	*	*	*	1.9	25.9	36.9	23.4	18.6	12.9	11.8
2006.....	13.7	*	*	*	2.5	29.8	37.1	24.0	17.5	13.8	11.9
2005.....	14.2	*	*	*	1.7	29.5	36.1	26.0	19.6	14.5	12.2
2004.....	13.9	*	*	*	1.4	28.1	35.9	25.4	17.0	13.7	11.9
2003.....	14.6	*	*	*	1.8	27.0	37.6	26.7	18.5	14.3	12.0
2002.....	14.2	*	*	*	1.4	28.5	35.1	25.4	16.7	13.9	11.6
2001.....	14.6	*	*	*	1.5	25.5	36.4	27.3	17.8	14.8	14.0
2000.....	14.2	*	*	*	2.4	27.9	33.3	23.5	16.9	11.9	14.9
1999.....	14.6	*	*	*	2.9	29.5	33.6	24.3	16.7	15.2	13.2

Hispanic, female

2007.....	1.5	*	*	*	*	2.5	3.3	2.7	2.2	1.9	2.1
2006.....	1.5	*	*	*	*	2.3	3.0	2.8	2.1	2.2	2.0
2005.....	1.6	*	*	*	*	2.1	3.0	2.8	2.8	2.8	2.2
2004.....	1.5	*	*	*	*	2.7	2.6	2.4	2.3	1.3	2.7
2003.....	1.7	*	*	*	*	2.8	4.1	2.3	1.9	2.5	2.2
2002.....	1.6	*	*	*	*	2.6	2.9	2.8	1.7	2.4	2.6
2001.....	1.7	*	*	*	*	2.9	3.7	3.0	2.3	2.3	2.2
2000.....	1.8	*	*	*	*	2.7	3.1	2.6	2.5	2.8	2.2
1999.....	2.0	*	*	*	*	3.6	4.0	2.6	2.2	2.7	2.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic 4/, both sexes

2007	10.8	*	0.4	0.4	1.2	11.9	19.8	17.1	15.2	12.6	12.6
2006	10.7	*	0.4	0.4	1.2	12.5	20.3	17.5	14.2	12.5	11.7
2005	10.7	*	0.4	0.3	1.4	11.7	19.6	17.6	14.6	12.1	12.2
2004	10.4	*	0.4	0.3	1.2	11.3	18.8	17.3	13.9	11.8	11.9
2003	10.6	*	0.3	0.3	1.2	11.4	20.7	17.7	14.3	12.0	12.0
2002	10.8	*	0.4	0.3	1.3	11.3	21.3	18.3	13.8	13.0	12.4
2001	10.6	*	0.5	0.4	1.3	11.9	20.8	17.9	14.0	12.7	11.6
2000	10.4	*	0.3	0.3	1.5	12.2	21.0	16.8	13.2	12.5	12.0
1999	10.6	*	0.5	0.4	1.6	13.8	21.1	17.0	13.7	12.6	11.2

Non-Hispanic 4/, male

2007	19.0	*	0.5	0.5	1.8	20.9	34.8	30.4	26.5	21.2	20.5
2006	18.8	*	0.5	0.5	1.9	21.9	35.8	30.8	25.1	20.7	19.3
2005	18.9	*	0.6	0.4	2.1	20.4	35.0	31.2	25.3	20.1	20.2
2004	18.2	*	0.4	0.4	1.9	19.2	33.6	30.9	23.7	20.0	19.3
2003	18.8	*	0.4	0.4	2.0	20.0	36.8	31.6	24.8	20.1	19.8
2002	19.0	*	0.5	0.3	2.0	19.5	37.9	32.4	23.4	21.7	20.7
2001	18.7	*	0.5	0.5	1.9	20.9	37.7	31.6	23.8	20.7	19.0
2000	18.3	*	0.4	0.4	2.4	21.1	37.6	29.6	22.1	20.5	19.8
1999	18.5	*	0.5	0.5	2.4	23.8	37.5	29.7	23.2	20.5	18.1

Non-Hispanic 4/, female

2007	2.9	*	0.4	0.3	0.5	2.6	4.0	3.8	3.9	4.2	5.0
2006	2.9	*	*	0.4	0.5	2.5	4.0	4.3	3.5	4.4	4.3
2005	2.8	*	*	0.3	0.6	2.5	3.7	3.9	4.0	4.1	4.4
2004	2.9	*	*	*	0.4	2.9	3.6	3.6	4.2	3.7	4.6
2003	2.9	*	*	*	0.4	2.3	4.2	3.9	4.0	4.0	4.4
2002	3.0	*	*	0.4	0.7	2.7	4.5	4.4	4.3	4.4	4.4
2001	3.0	*	0.4	0.3	0.6	2.5	3.8	4.4	4.2	4.9	4.3
2000	3.0	*	*	0.2	0.5	2.9	4.4	4.1	4.3	4.7	4.5
1999	3.0	*	0.4	0.3	0.8	3.4	4.5	4.4	4.3	4.9	4.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic White, both sexes

2007 5/.....	9.4	*	0.3	0.3	0.8	5.5	10.7	10.4	11.2	10.8	12.3
2006 5/.....	9.1	*	*	0.4	0.9	6.1	10.3	9.8	9.6	10.9	11.3
2005 5/.....	9.2	*	0.3	0.2	1.1	6.3	10.0	10.3	10.3	10.3	11.7
2004 5/.....	9.1	*	*	0.2	0.9	6.6	10.5	10.0	10.3	10.4	11.6
2003 5/.....	9.3	*	*	0.2	0.9	6.5	11.2	10.1	10.6	10.6	11.7
2002.....	9.4	*	0.3	0.2	1.0	6.7	11.8	11.2	10.2	11.3	12.2
2001.....	9.4	*	0.3	0.4	1.1	7.4	11.5	11.7	11.1	11.5	11.3
2000.....	9.1	*	0.2	0.3	1.4	7.4	12.0	10.7	10.4	11.3	11.7
1999.....	9.3	*	0.3	0.3	1.5	8.4	12.1	11.3	11.2	11.5	10.8

Non-Hispanic White, male

2007 5/.....	16.1	*	*	*	1.2	8.9	17.9	17.6	18.3	17.6	19.6
2006 5/.....	15.6	*	*	0.4	1.3	10.3	17.5	16.4	16.2	17.3	18.2
2005 5/.....	15.9	*	*	*	1.7	10.5	17.2	17.2	16.7	16.6	18.8
2004 5/.....	15.6	*	*	*	1.4	10.8	17.9	17.0	16.8	17.0	18.4
2003 5/.....	16.0	*	*	*	1.5	11.1	19.4	17.0	17.6	17.1	18.9
2002.....	16.3	*	*	*	1.5	11.2	20.2	18.3	16.6	18.3	20.2
2001.....	16.3	*	*	0.5	1.7	12.3	19.9	19.7	18.1	18.3	18.5
2000.....	15.7	*	*	0.4	2.2	12.3	20.6	17.7	16.8	17.8	19.0
1999.....	15.9	*	*	*	2.2	14.2	20.6	18.9	18.2	18.3	17.4

Non-Hispanic White, female

2007 5/.....	2.9	*	*	*	0.4	1.9	3.1	3.0	3.9	4.0	5.1
2006 5/.....	2.8	*	*	*	0.3	1.7	2.7	3.2	2.9	4.4	4.4
2005 5/.....	2.8	*	*	*	0.5	1.9	2.5	3.3	3.7	4.0	4.6
2004 5/.....	2.9	*	*	*	0.3	2.3	2.7	2.8	3.8	3.6	4.9
2003 5/.....	2.7	*	*	*	*	1.7	2.7	3.1	3.6	4.1	4.4
2002.....	2.8	*	*	*	0.5	1.9	3.2	3.8	3.8	4.2	4.3
2001.....	2.9	*	*	0.4	0.5	2.2	2.9	3.5	4.0	4.6	4.2
2000.....	2.9	*	*	*	0.5	2.2	3.2	3.5	3.9	4.7	4.4
1999.....	2.9	*	*	*	0.7	2.3	3.3	3.5	4.0	4.7	4.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic Black, both sexes

2007 5/.....	21.3	*	1.0	0.8	2.8	39.4	62.3	51.9	40.3	26.5	18.3
2006 5/.....	22.0	*	1.1	0.7	2.8	38.9	66.5	56.7	41.3	25.6	17.7
2005 5/.....	21.1	*	1.1	0.8	2.5	34.7	64.1	54.7	40.8	24.7	18.7
2004 5/.....	19.9	*	1.1	0.8	2.3	31.9	60.1	55.7	35.8	22.8	16.2
2003 5/.....	20.7	*	0.9	0.9	2.6	33.2	66.0	58.1	37.2	22.3	17.1
2002.....	20.7	*	*	*	2.5	32.3	67.0	57.1	36.3	25.2	16.9
2001.....	19.8	*	1.0	*	1.9	33.8	66.7	52.3	32.6	23.3	15.7
2000.....	19.8	*	*	*	2.3	35.0	66.3	50.2	31.7	22.3	16.9
1999.....	20.0	*	1.1	0.9	2.4	39.2	66.3	48.2	29.8	21.5	16.0

Non-Hispanic Black, male

2007 5/.....	40.3	*	*	*	4.3	72.3	114.2	97.4	77.8	48.8	32.5
2006 5/.....	41.4	*	*	*	4.5	70.8	122.0	105.9	79.0	47.5	31.7
2005 5/.....	40.0	*	*	*	3.8	63.7	118.6	104.6	78.5	46.1	34.4
2004 5/.....	37.6	*	*	*	3.7	57.3	111.7	107.4	67.4	42.8	29.9
2003 5/.....	39.1	*	*	*	3.9	61.2	121.6	112.6	70.3	42.0	30.3
2002.....	38.9	*	*	*	3.7	57.5	125.0	111.0	68.1	46.3	29.6
2001.....	37.3	*	*	*	3.1	62.1	127.9	99.4	62.2	40.8	27.2
2000.....	37.1	*	*	*	3.8	63.5	125.7	97.1	58.5	41.1	29.2
1999.....	37.1	*	*	1.3	3.5	69.4	125.3	91.9	55.9	38.1	27.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic Black, female

2007 5/.....	4.0	*	*	*	1.3	5.6	9.1	8.2	6.2	6.6	5.7
2006 5/.....	4.2	*	*	*	*	6.1	9.9	10.2	7.1	6.2	5.4
2005 5/.....	3.8	*	*	*	*	5.0	8.8	7.8	6.8	5.5	4.9
2004 5/.....	3.8	*	*	*	*	5.9	8.3	7.6	7.2	4.9	4.1
2003 5/.....	4.0	*	*	*	1.3	4.4	10.9	7.8	7.5	4.9	5.6
2002.....	4.3	*	*	*	1.3	6.2	10.6	8.0	7.6	6.5	5.7
2001.....	4.0	*	*	*	*	4.6	8.0	9.4	6.0	7.8	5.6
2000.....	4.1	*	*	*	*	5.9	10.1	7.8	7.7	5.6	6.1
1999.....	4.5	*	*	*	*	8.3	10.3	8.7	6.5	6.7	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

All origins 3/, both sexes

2007.....	340.0	509.0	726.3	1,068.3	1,627.5	2,491.3	3,945.9	6,381.4	12,946.5
2006.....	347.7	516.4	730.1	1,110.2	1,656.6	2,554.8	4,033.4	6,524.0	13,253.1
2005.....	353.1	520.8	734.6	1,136.9	1,700.0	2,657.6	4,154.0	6,712.9	13,798.6
2004.....	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
2003.....	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6

All origins 3/, male

2007.....	423.8	646.6	922.2	1,328.4	2,002.2	3,046.9	4,817.2	7,758.7	14,006.4
2006.....	435.3	659.7	920.0	1,373.6	2,040.2	3,117.5	4,944.6	7,942.7	14,309.1
2005.....	443.7	666.0	925.7	1,410.0	2,084.2	3,267.0	5,103.4	8,147.4	14,889.4
2004.....	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
2003.....	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2

All origins 3/, female

2007.....	258.1	376.8	541.8	829.7	1,299.4	2,037.2	3,304.0	5,511.7	12,442.3
2006.....	262.2	378.9	551.0	869.7	1,321.4	2,097.0	3,366.6	5,637.2	12,759.0
2005.....	264.6	381.6	554.3	887.9	1,364.7	2,164.0	3,464.7	5,822.0	13,297.7
2004.....	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
2003.....	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

Hispanic, both sexes

2007	245.0	378.2	539.5	798.2	1,211.9	1,822.6	2,959.5	4,746.9	8,542.2
2006	253.6	385.2	549.3	813.9	1,229.6	1,922.0	3,052.6	4,794.7	9,022.8
2005	256.1	398.8	549.3	859.4	1,276.5	2,067.5	3,196.2	5,186.4	9,436.7
2004	258.4	385.6	557.4	862.6	1,289.9	2,017.4	3,218.2	5,038.4	9,483.8
2003	272.7	400.3	590.7	886.5	1,352.1	2,107.8	3,380.8	5,251.8	10,431.5
2002	266.6	393.2	583.0	905.8	1,378.9	2,179.7	3,411.7	5,365.5	10,679.0
2001	280.9	403.9	580.6	910.1	1,412.8	2,229.5	3,556.3	5,501.5	11,821.8
2000	281.1	394.3	612.1	913.8	1,425.9	2,278.4	3,543.6	5,777.6	11,822.9
1999	283.6	398.1	602.7	940.8	1,527.8	2,264.3	3,577.0	5,774.4	12,177.9

Hispanic, male

2007	323.0	500.0	694.7	1,028.9	1,556.5	2,278.5	3,609.0	5,546.1	8,953.7
2006	326.7	507.1	718.7	1,030.0	1,553.4	2,393.6	3,720.7	5,719.9	9,435.5
2005	339.5	522.2	712.8	1,118.0	1,596.8	2,611.5	3,987.7	6,265.6	10,140.5
2004	343.3	518.0	706.7	1,118.2	1,606.8	2,514.1	3,990.1	6,105.0	9,932.8
2003	365.3	532.3	763.9	1,146.5	1,701.1	2,660.5	4,124.7	6,357.9	11,029.2
2002	353.7	516.8	779.4	1,158.2	1,765.5	2,747.9	4,258.0	6,447.7	11,674.1
2001	379.7	529.6	751.2	1,158.6	1,817.5	2,791.5	4,456.8	6,783.5	12,888.3
2000	378.6	519.5	801.5	1,181.3	1,804.0	2,918.9	4,480.2	7,220.4	13,086.2
1999	383.9	525.5	788.0	1,215.5	1,942.0	2,931.5	4,571.8	7,005.8	13,485.9

Hispanic, female

2007	162.4	256.1	391.9	592.1	921.1	1,465.7	2,487.6	4,213.5	8,318.9
2006	177.0	263.9	389.3	622.4	957.8	1,554.8	2,569.3	4,178.9	8,803.5
2005	169.1	276.8	395.9	631.0	1,009.5	1,645.2	2,627.7	4,470.5	9,068.4
2004	170.7	256.1	417.9	639.0	1,027.7	1,634.2	2,669.9	4,334.3	9,253.0
2003	177.9	271.9	430.3	659.9	1,065.6	1,683.6	2,853.0	4,531.9	10,128.1
2002	178.1	273.2	402.6	685.7	1,063.3	1,742.1	2,807.4	4,674.8	10,186.0
2001	181.7	283.8	426.0	695.1	1,085.0	1,800.8	2,917.0	4,707.4	11,300.5
2000	183.8	274.9	440.6	683.1	1,119.7	1,793.2	2,880.3	4,899.7	11,202.8
1999	184.2	277.2	435.3	704.0	1,192.9	1,762.5	2,874.8	5,039.8	11,547.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

Non-Hispanic 4/, both sexes

2007	351.7	521.7	741.8	1,088.7	1,658.3	2,539.1	4,009.6	6,473.0	13,170.0
2006	358.3	528.6	744.2	1,132.1	1,686.9	2,597.2	4,092.4	6,612.2	13,450.1
2005	363.5	531.3	748.5	1,156.7	1,729.7	2,695.5	4,207.6	6,784.5	13,990.9
2004	361.6	522.0	747.3	1,158.9	1,756.4	2,715.6	4,243.0	6,789.9	13,997.7
2003	367.2	526.6	766.2	1,201.4	1,821.5	2,815.4	4,408.5	7,042.5	14,740.6
2002	365.0	521.3	782.3	1,204.9	1,859.4	2,879.6	4,490.9	7,161.0	14,958.7
2001	360.4	519.3	783.7	1,228.7	1,896.2	2,908.7	4,528.0	7,201.4	15,199.0
2000	357.3	515.8	802.9	1,259.9	1,931.9	2,956.6	4,586.6	7,375.0	15,619.8
1999	350.4	507.2	812.7	1,271.3	1,977.5	3,025.1	4,639.4	7,416.1	15,631.0

Non-Hispanic 4/, male

2007	436.4	660.8	940.6	1,349.8	2,033.5	3,099.0	4,893.4	7,887.9	14,289.1
2006	447.7	673.6	934.9	1,397.6	2,072.6	3,162.7	5,017.0	8,061.4	14,558.8
2005	454.9	678.1	940.7	1,429.2	2,116.2	3,306.7	5,163.4	8,239.6	15,121.2
2004	454.3	667.7	931.4	1,428.8	2,157.1	3,321.9	5,232.9	8,272.9	15,260.4
2003	466.2	669.7	952.7	1,479.9	2,246.3	3,468.2	5,458.2	8,599.8	15,984.2
2002	463.0	662.5	972.1	1,493.6	2,292.2	3,569.8	5,572.1	8,727.3	16,418.0
2001	460.5	653.3	969.6	1,514.1	2,341.5	3,621.9	5,644.0	8,842.8	16,866.3
2000	460.9	648.3	998.9	1,554.1	2,388.4	3,704.3	5,730.8	9,144.1	17,639.8
1999	450.9	637.0	1,014.4	1,577.2	2,466.2	3,815.8	5,842.6	9,272.9	17,710.4

Non-Hispanic 4/, female

2007	269.7	388.9	554.6	848.6	1,328.9	2,079.9	3,358.0	5,582.5	12,641.5
2006	271.9	389.9	564.3	889.0	1,348.6	2,135.6	3,415.2	5,709.7	12,934.8
2005	275.0	391.0	567.0	907.5	1,391.0	2,199.0	3,513.2	5,884.1	13,475.2
2004	271.9	382.9	573.6	912.5	1,406.4	2,226.0	3,531.1	5,879.4	13,433.1
2003	271.5	389.8	590.4	947.5	1,451.7	2,290.3	3,661.1	6,092.9	14,194.2
2002	270.3	386.3	603.7	942.0	1,484.2	2,326.6	3,727.8	6,210.5	14,329.1
2001	263.6	391.0	608.9	968.9	1,511.2	2,340.5	3,747.4	6,219.1	14,505.6
2000	257.3	388.8	619.1	992.4	1,539.1	2,366.3	3,790.4	6,332.6	14,802.9
1999	253.4	382.7	623.8	993.6	1,557.9	2,406.0	3,803.4	6,330.3	14,798.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

Non-Hispanic White, both sexes

2007 5/.....	327.7	483.6	693.4	1,039.6	1,608.0	2,499.8	3,995.4	6,515.4	13,413.3
2006 5/.....	331.4	487.4	696.1	1,081.7	1,634.0	2,559.2	4,077.7	6,640.0	13,688.2
2005 5/.....	333.9	486.6	699.3	1,101.0	1,675.1	2,652.5	4,179.8	6,803.0	14,222.4
2004 5/.....	331.1	477.6	696.6	1,105.0	1,700.5	2,669.8	4,207.6	6,807.8	14,191.1
2003 5/.....	333.6	478.6	716.6	1,142.1	1,762.4	2,761.7	4,371.2	7,056.2	14,944.8
2002.....	331.7	475.5	731.3	1,145.4	1,800.0	2,825.5	4,446.2	7,166.0	15,149.1
2001.....	324.7	474.0	730.8	1,169.7	1,827.4	2,849.2	4,476.8	7,196.1	15,371.2
2000.....	318.7	470.1	749.9	1,199.4	1,865.4	2,889.5	4,525.0	7,369.2	15,799.7
1999.....	310.1	461.9	754.0	1,206.2	1,911.7	2,949.5	4,570.3	7,406.6	15,795.3

Non-Hispanic White, male

2007 5/.....	409.6	613.1	875.5	1,281.9	1,962.5	3,037.2	4,859.8	7,942.6	14,588.3
2006 5/.....	417.1	621.0	869.7	1,327.5	2,000.1	3,100.3	4,983.5	8,103.3	14,841.1
2005 5/.....	420.7	622.6	872.9	1,354.3	2,042.8	3,242.9	5,122.3	8,261.3	15,401.3
2004 5/.....	418.2	612.7	863.8	1,355.2	2,082.3	3,257.1	5,178.6	8,308.7	15,489.2
2003 5/.....	426.7	609.7	888.4	1,402.1	2,169.8	3,393.7	5,403.0	8,607.0	16,234.4
2002.....	424.3	604.5	906.2	1,413.9	2,216.9	3,500.6	5,514.6	8,733.5	16,641.9
2001.....	417.3	596.5	901.6	1,439.8	2,256.8	3,543.1	5,580.2	8,828.0	17,055.3
2000.....	415.7	588.9	928.4	1,477.3	2,305.7	3,615.5	5,653.4	9,137.4	17,853.2
1999.....	399.6	578.8	939.4	1,494.4	2,384.2	3,724.5	5,759.4	9,266.7	17,871.2

Non-Hispanic White, female

2007 5/.....	246.7	356.7	517.6	811.8	1,289.9	2,049.4	3,344.3	5,605.4	12,856.7
2006 5/.....	246.7	356.7	528.4	851.6	1,306.2	2,107.8	3,400.4	5,718.8	13,150.7
2005 5/.....	248.0	353.5	531.6	864.3	1,346.2	2,162.2	3,481.1	5,892.3	13,683.1
2004 5/.....	244.9	345.4	535.3	871.8	1,360.0	2,185.0	3,496.3	5,878.7	13,609.6
2003 5/.....	241.4	350.2	551.1	900.2	1,400.2	2,242.3	3,624.5	6,103.0	14,377.2
2002.....	240.1	349.1	563.3	896.1	1,430.8	2,273.2	3,681.6	6,207.1	14,504.3
2001.....	233.1	353.9	567.1	919.2	1,448.1	2,285.2	3,695.5	6,211.0	14,670.6
2000.....	222.8	353.6	579.3	941.7	1,478.3	2,305.2	3,731.9	6,318.4	14,971.7
1999.....	221.7	347.5	576.9	939.4	1,497.5	2,331.8	3,735.9	6,309.1	14,967.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

Non-Hispanic Black, both sexes

2007 5/.....	565.0	883.6	1,243.2	1,738.9	2,433.4	3,373.8	4,924.3	7,076.4	12,468.1
2006 5/.....	590.6	914.1	1,260.3	1,789.0	2,454.3	3,432.1	4,987.2	7,380.4	12,655.4
2005 5/.....	616.8	935.4	1,281.3	1,846.9	2,511.2	3,572.9	5,236.9	7,641.2	13,239.1
2004 5/.....	621.7	928.8	1,296.4	1,845.3	2,554.0	3,637.4	5,336.6	7,587.7	13,492.2
2003 5/.....	647.8	959.3	1,321.4	1,920.6	2,640.7	3,810.9	5,490.4	7,836.9	14,107.7
2002.....	648.6	943.7	1,346.7	1,911.2	2,670.7	3,877.8	5,630.1	8,036.6	14,386.7
2001.....	656.8	946.1	1,351.9	1,920.5	2,777.9	3,951.1	5,712.5	8,098.0	14,660.8
2000.....	679.7	956.6	1,368.9	1,959.9	2,809.4	4,065.0	5,837.2	8,234.9	14,837.5
1999.....	681.6	945.4	1,422.8	1,998.3	2,851.2	4,191.4	5,927.6	8,223.1	14,839.2

Non-Hispanic Black, male

2007 5/.....	696.8	1,137.9	1,651.2	2,301.6	3,178.7	4,336.7	6,308.6	8,795.9	13,189.0
2006 5/.....	732.9	1,199.1	1,662.0	2,355.8	3,190.8	4,451.9	6,441.8	9,165.9	13,403.1
2005 5/.....	772.6	1,211.4	1,701.2	2,427.3	3,232.7	4,585.9	6,672.3	9,492.7	13,978.1
2004 5/.....	782.6	1,211.2	1,702.7	2,416.8	3,298.9	4,655.2	6,839.3	9,346.6	14,647.3
2003 5/.....	819.8	1,251.2	1,722.5	2,493.7	3,390.4	4,907.6	7,066.5	9,941.1	15,082.2
2002.....	818.5	1,229.3	1,751.3	2,503.4	3,402.3	4,983.1	7,157.7	10,054.7	15,776.9
2001.....	846.0	1,222.7	1,746.2	2,456.8	3,547.9	5,105.6	7,313.8	10,346.7	16,576.2
2000.....	869.4	1,248.2	1,802.9	2,498.8	3,559.1	5,258.8	7,462.1	10,491.3	16,944.8
1999.....	902.1	1,238.5	1,851.6	2,583.0	3,644.4	5,379.6	7,591.6	10,490.8	17,385.3

Non-Hispanic Black, female

2007 5/.....	450.5	668.4	909.1	1,298.9	1,886.2	2,710.8	4,091.7	6,173.3	12,169.3
2006 5/.....	467.0	673.7	929.8	1,348.2	1,911.3	2,732.5	4,118.7	6,438.9	12,350.5
2005 5/.....	481.1	702.6	935.2	1,395.5	1,977.0	2,880.6	4,382.4	6,664.8	12,941.6
2004 5/.....	481.7	690.6	962.5	1,400.6	2,001.4	2,947.5	4,447.1	6,662.3	13,035.6
2003 5/.....	498.1	712.7	992.0	1,474.3	2,084.8	3,071.6	4,554.5	6,735.8	13,730.3
2002.....	501.4	702.1	1,014.9	1,448.4	2,128.3	3,136.6	4,718.0	6,986.6	13,855.4
2001.....	493.1	710.9	1,028.9	1,498.8	2,209.4	3,182.0	4,756.8	6,952.7	13,951.4
2000.....	515.5	708.3	1,013.7	1,534.7	2,256.3	3,272.3	4,857.7	7,099.0	14,054.9
1999.....	491.7	697.1	1,073.0	1,539.8	2,263.4	3,402.8	4,925.0	7,099.0	13,887.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

All origins 3/, both sexes

2007.....	0.2	0.4	0.8	1.3	3.3	6.5	13.7	25.3	52.0
2006.....	0.2	0.4	0.9	1.5	3.3	6.7	13.5	26.2	50.5
2005.....	0.1	0.4	0.5	1.2	2.8	6.5	11.6	21.5	45.8
2004.....	0.2	0.3	0.6	1.3	2.2	4.6	9.3	17.5	34.3
2003.....	0.1	0.2	0.4	0.9	1.7	3.3	7.0	12.3	24.8
2002.....	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.7	21.0
2001.....	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.8
2000.....	*	*	0.2	0.3	0.9	1.5	2.7	5.3	11.8
1999.....	*	*	*	0.3	0.7	1.2	2.3	4.5	10.3

All origins 3/, male

2007.....	*	0.4	0.9	1.3	3.3	6.3	13.7	24.7	50.4
2006.....	0.2	0.4	0.9	1.8	3.7	6.3	13.9	25.3	50.0
2005.....	*	0.4	0.5	1.4	2.9	6.3	12.4	22.1	46.1
2004.....	*	0.3	0.6	1.3	2.2	4.3	9.3	17.8	32.0
2003.....	*	0.2	0.5	0.8	1.5	3.1	6.9	12.9	21.4
2002.....	*	*	0.4	0.8	1.5	3.1	5.1	10.1	20.4
2001.....	*	*	*	0.4	0.7	2.0	3.5	6.2	12.1
2000.....	*	*	0.3	*	1.0	1.6	2.9	5.8	11.4
1999.....	*	*	*	*	0.6	1.2	2.7	4.9	9.5

All origins 3/, female

2007.....	0.2	0.5	0.6	1.3	3.3	6.7	13.7	25.7	52.7
2006.....	*	0.4	0.8	1.3	3.0	6.9	13.3	26.8	50.8
2005.....	*	0.4	0.6	1.0	2.6	6.6	11.0	21.1	45.6
2004.....	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
2003.....	*	0.2	0.4	0.9	1.8	3.5	7.1	12.0	26.3
2002.....	*	*	0.4	0.6	1.3	2.3	5.0	9.5	21.3
2001.....	*	0.2	*	0.5	0.8	1.8	3.0	6.5	13.1
2000.....	*	*	*	*	0.8	1.5	2.6	4.9	11.9
1999.....	*	*	*	*	0.8	1.3	2.0	4.3	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 4/, both sexes

2007.....	0.1	0.4	0.8	1.4	3.3	6.8	14.0	25.9	52.9
2006.....	0.2	0.4	0.9	1.5	3.4	6.7	13.5	26.6	51.8
2005.....	0.1	0.4	0.6	1.3	2.8	6.7	11.9	21.8	46.7
2004.....	0.2	0.3	0.6	1.3	2.2	4.7	9.5	17.9	35.0
2003.....	*	0.2	0.5	0.9	1.7	3.4	7.2	12.7	25.2
2002.....	0.1	0.2	0.5	0.7	1.4	2.7	5.1	9.8	21.4
2001.....	*	0.2	0.2	0.5	0.8	1.9	3.2	6.4	12.8
2000.....	*	*	0.2	0.3	0.9	1.6	2.8	5.3	11.9
1999.....	*	*	*	0.3	0.7	1.3	2.3	4.7	10.4

Non-Hispanic 4/, male

2007.....	*	0.4	1.0	1.3	3.2	6.5	13.8	25.4	51.4
2006.....	0.3	0.4	0.9	1.7	3.8	6.3	14.1	25.7	51.2
2005.....	*	0.4	0.6	1.5	3.0	6.5	12.9	22.6	47.4
2004.....	*	0.3	0.6	1.4	2.2	4.4	9.5	18.3	32.4
2003.....	*	*	0.5	0.8	1.5	3.2	7.0	13.3	21.4
2002.....	*	*	0.4	0.7	1.6	3.2	5.1	10.2	21.0
2001.....	*	*	*	0.4	0.8	2.0	3.5	6.4	12.0
2000.....	*	*	*	*	1.0	1.5	2.9	5.9	11.5
1999.....	*	*	*	*	0.6	1.2	2.7	5.1	9.6

Non-Hispanic 4/, female

2007.....	*	0.5	0.6	1.4	3.3	7.0	14.2	26.2	53.6
2006.....	*	0.5	0.8	1.4	3.1	7.1	13.1	27.1	52.1
2005.....	*	0.4	0.5	1.0	2.7	6.8	11.2	21.4	46.4
2004.....	*	0.3	0.6	1.2	2.2	5.0	9.5	17.6	36.1
2003.....	*	0.2	0.5	0.9	1.9	3.6	7.3	12.3	26.8
2002.....	*	*	0.5	0.6	1.3	2.4	5.0	9.6	21.6
2001.....	*	0.2	*	0.6	0.8	1.8	3.0	6.4	13.2
2000.....	*	*	*	*	0.8	1.6	2.7	5.0	12.1
1999.....	*	*	*	*	0.8	1.3	2.1	4.4	10.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic White, both sexes

2007 5/.....	0.1	0.4	0.8	1.3	3.3	7.2	14.8	27.4	56.1
2006 5/.....	0.2	0.4	0.8	1.5	3.6	7.0	14.4	28.4	55.1
2005 5/.....	0.2	0.3	0.5	1.3	3.0	7.0	12.6	23.1	49.7
2004 5/.....	0.1	0.3	0.6	1.3	2.3	5.0	10.0	18.8	37.3
2003 5/.....	*	0.2	0.5	0.8	1.8	3.7	7.3	13.6	26.5
2002.....	0.1	0.2	0.4	0.6	1.6	2.8	5.4	10.4	22.5
2001.....	*	0.2	0.2	0.5	0.7	2.0	3.4	6.6	13.6
2000.....	*	*	0.2	0.3	0.9	1.6	2.9	5.7	12.3
1999.....	*	*	*	0.3	0.7	1.3	2.4	4.9	11.0

Non-Hispanic White, male

2007 5/.....	*	0.4	1.0	1.3	3.3	6.6	14.4	26.7	54.7
2006 5/.....	*	0.3	0.9	1.6	4.0	6.5	14.8	27.1	54.4
2005 5/.....	*	0.3	0.5	1.6	3.2	6.8	13.5	23.3	50.9
2004 5/.....	*	0.3	0.5	1.5	2.2	4.7	10.0	19.3	34.5
2003 5/.....	*	*	0.4	0.7	1.6	3.4	7.3	14.2	22.7
2002.....	*	*	0.4	0.7	1.6	3.1	5.3	10.9	22.1
2001.....	*	*	*	*	0.7	2.1	3.6	6.4	12.6
2000.....	*	*	*	*	1.0	1.6	3.1	6.2	12.0
1999.....	*	*	*	*	0.7	1.3	2.8	5.4	10.0

Non-Hispanic White, female

2007 5/.....	*	0.4	0.6	1.4	3.4	7.6	15.1	27.9	56.7
2006 5/.....	*	0.4	0.7	1.3	3.2	7.5	14.1	29.2	55.4
2005 5/.....	*	0.4	0.5	1.0	2.8	7.2	11.9	22.9	49.2
2004 5/.....	*	0.3	0.6	1.2	2.4	5.4	10.0	18.5	38.6
2003 5/.....	*	*	0.5	0.9	2.0	3.9	7.4	13.2	28.2
2002.....	*	*	0.4	0.6	1.5	2.5	5.4	10.2	22.7
2001.....	*	*	*	0.5	0.7	1.9	3.2	6.7	14.0
2000.....	*	*	*	*	0.8	1.7	2.8	5.4	12.5
1999.....	*	*	*	*	0.8	1.3	2.1	4.6	11.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

All origins 3/, both sexes

2007.....	0.1	0.2	0.2	0.3	0.4	0.7	0.9	1.5	1.6
2006.....	0.2	0.2	0.3	0.4	0.6	0.7	0.9	1.7	2.1
2005.....	0.2	0.2	0.2	0.4	0.5	0.7	1.3	1.3	2.2
2004.....	0.1	0.2	0.2	0.3	0.6	1.0	1.1	1.5	2.4
2003.....	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2

All origins 3/, male

2007.....	0.2	0.3	0.4	0.5	0.4	0.9	1.2	2.0	2.8
2006.....	0.3	0.4	0.4	0.7	0.8	0.9	1.4	2.5	3.1
2005.....	0.2	0.3	0.3	0.6	0.7	1.1	1.7	1.7	3.3
2004.....	0.2	0.3	0.3	0.6	0.8	1.4	1.2	2.2	3.8
2003.....	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1

All origins 3/, female

2007.....	*	*	*	*	*	0.5	0.6	1.3	1.1
2006.....	*	*	*	*	*	0.5	0.6	1.3	1.6
2005.....	*	*	*	*	0.4	0.4	0.9	1.0	1.7
2004.....	*	*	*	*	*	0.6	1.0	1.1	1.7
2003.....	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	0.4	0.7	1.5	1.7	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

Non-Hispanic 4/, both sexes

2007.....	0.1	0.2	0.2	0.3	0.3	0.7	0.7	1.5	1.5
2006.....	0.2	0.2	0.2	0.3	0.5	0.6	0.8	1.7	2.0
2005.....	0.1	0.2	0.2	0.3	0.5	0.7	1.3	1.2	2.2
2004.....	0.1	0.2	0.2	0.3	0.5	0.8	0.9	1.4	2.2
2003.....	0.2	0.3	0.3	0.3	0.5	0.7	1.3	2.0	2.2
2002.....	0.2	0.2	0.3	0.4	0.8	0.8	1.5	2.1	2.5
2001.....	0.2	0.2	0.4	0.4	0.6	1.0	1.3	1.9	2.7
2000.....	0.2	0.3	0.2	0.4	0.7	0.9	1.6	2.3	2.5
1999.....	0.2	0.3	0.5	0.5	0.7	1.1	1.6	2.2	3.9

Non-Hispanic 4/, male

2007.....	0.2	0.3	0.3	0.4	*	0.8	0.9	2.0	2.3
2006.....	0.4	0.3	0.4	0.6	0.8	0.9	1.3	2.4	2.8
2005.....	*	0.3	0.3	0.5	0.7	1.0	1.7	1.4	3.2
2004.....	*	0.3	0.3	0.5	0.8	1.2	1.0	1.9	3.5
2003.....	0.4	0.4	0.5	0.4	0.6	0.9	1.9	2.6	3.0
2002.....	*	0.3	0.5	0.6	1.1	1.2	1.8	3.0	3.8
2001.....	0.3	0.3	0.6	0.7	0.8	1.4	1.7	2.2	4.3
2000.....	0.3	0.5	0.4	0.6	1.0	1.0	2.0	3.6	4.3
1999.....	0.4	0.4	0.7	0.8	1.0	1.6	2.1	3.5	6.8

Non-Hispanic 4/, female

2007.....	*	*	*	*	*	0.6	0.5	1.2	1.1
2006.....	*	*	*	*	*	*	*	1.2	1.6
2005.....	*	*	*	*	*	*	0.9	1.0	1.7
2004.....	*	*	*	*	*	0.5	0.9	1.1	1.6
2003.....	*	*	*	*	*	0.5	1.0	1.6	1.8
2002.....	*	*	*	*	0.5	0.6	1.4	1.5	1.9
2001.....	*	*	*	*	*	0.7	1.0	1.7	2.1
2000.....	*	*	*	*	0.5	0.9	1.4	1.6	1.8
1999.....	*	*	*	*	*	0.7	1.3	1.5	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	0.2	0.4	0.4	0.9	1.4
2006 5/.....	*	*	*	*	0.3	0.3	0.6	1.2	1.5
2005 5/.....	*	*	*	0.2	0.4	0.3	0.8	0.8	1.6
2004 5/.....	*	*	*	*	0.3	0.5	0.6	1.0	1.5
2003 5/.....	*	*	0.2	*	*	0.6	0.9	1.6	1.7
2002.....	*	*	*	0.3	0.4	0.5	1.2	1.6	1.9
2001.....	*	*	0.2	0.3	0.4	0.6	1.0	1.3	2.0
2000.....	*	0.2	*	*	0.4	0.5	1.0	1.6	2.0
1999.....	*	*	0.3	0.3	0.5	0.6	1.1	1.7	3.0

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	*	1.1	2.1
2006 5/.....	*	*	*	*	*	*	0.9	1.6	1.7
2005 5/.....	*	*	*	*	*	*	1.0	*	2.3
2004 5/.....	*	*	*	*	*	0.7	0.8	1.2	2.4
2003 5/.....	*	*	0.4	*	*	0.8	1.1	2.0	2.4
2002.....	*	*	*	*	0.7	0.9	1.3	2.2	2.9
2001.....	*	*	*	*	*	0.6	1.3	1.5	2.9
2000.....	*	*	*	*	0.6	0.7	1.2	2.0	3.6
1999.....	*	*	0.5	*	0.7	1.0	1.3	2.8	4.9

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	0.8	1.0
2006 5/.....	*	*	*	*	*	*	*	0.9	1.3
2005 5/.....	*	*	*	*	*	*	0.6	0.7	1.3
2004 5/.....	*	*	*	*	*	*	0.6	0.9	1.1
2003 5/.....	*	*	*	*	*	*	0.8	1.3	1.4
2002.....	*	*	*	*	*	*	1.1	1.2	1.4
2001.....	*	*	*	*	*	0.5	0.8	1.1	1.7
2000.....	*	*	*	*	*	*	0.9	1.3	1.4
1999.....	*	*	*	*	*	*	0.9	1.1	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

All origins 3/, both sexes

2007.....	0.1	0.2	0.2	0.2	0.3	0.5	0.7	1.2	1.3
2006.....	0.1	0.2	0.2	0.4	0.4	0.5	0.7	1.4	1.6
2005.....	0.1	0.2	0.2	0.3	0.3	0.6	1.0	1.0	1.9
2004.....	0.1	0.2	0.2	0.3	0.4	0.8	0.8	1.1	2.2
2003.....	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7

All origins 3/, male

2007.....	*	0.2	0.2	0.4	*	0.6	1.1	1.5	2.4
2006.....	0.2	0.3	0.3	0.6	0.6	0.7	1.2	2.0	2.4
2005.....	*	0.3	0.2	0.4	0.5	0.9	1.2	1.4	2.7
2004.....	*	0.3	*	0.4	0.6	1.2	0.9	1.7	3.7
2003.....	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4

All origins 3/, female

2007.....	*	*	*	*	*	*	0.5	0.9	0.9
2006.....	*	*	*	*	*	*	*	1.0	1.2
2005.....	*	*	*	*	*	*	0.8	0.8	1.5
2004.....	*	*	*	*	*	0.5	0.7	0.8	1.5
2003.....	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	0.5	1.2	1.4	2.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

Non-Hispanic 4/, both sexes

2007	*	0.1	0.1	0.2	0.3	0.5	0.6	1.1	1.2
2006	0.2	0.1	0.2	0.3	0.3	0.5	0.6	1.4	1.5
2005	0.1	0.2	0.2	0.2	0.3	0.5	1.0	1.0	1.8
2004	0.1	0.2	0.1	0.2	0.4	0.7	0.6	1.0	2.0
2003	0.2	0.2	0.3	0.2	0.4	0.5	1.1	1.5	1.7
2002	0.1	0.2	0.2	0.3	0.6	0.6	1.2	1.8	2.2
2001	0.1	0.2	0.3	0.4	0.4	0.7	1.1	1.5	2.4
2000	0.2	0.2	0.2	0.3	0.5	0.7	1.3	2.0	2.2
1999	0.2	0.2	0.4	0.5	0.6	0.8	1.4	2.0	3.5

Non-Hispanic 4/, male

2007	*	0.2	*	0.4	*	0.6	0.8	1.4	2.0
2006	0.3	*	0.3	0.5	0.6	0.7	1.1	2.0	2.2
2005	*	0.2	*	0.4	0.5	0.9	1.2	1.3	2.7
2004	*	0.3	*	0.4	0.6	1.0	*	1.4	3.3
2003	0.3	0.3	0.4	*	0.5	0.8	1.6	2.0	2.4
2002	*	0.3	0.4	0.5	0.9	0.9	1.3	2.5	3.3
2001	*	0.3	0.5	0.7	0.6	1.0	1.4	1.7	3.8
2000	0.3	0.4	0.3	0.5	0.8	0.8	1.7	3.0	3.8
1999	0.3	0.3	0.6	0.8	0.9	1.3	1.8	3.2	6.1

Non-Hispanic 4/, female

2007	*	*	*	*	*	*	*	0.9	0.8
2006	*	*	*	*	*	*	*	1.0	1.2
2005	*	*	*	*	*	*	0.8	0.8	1.4
2004	*	*	*	*	*	0.5	0.6	0.8	1.3
2003	*	*	*	*	*	*	0.8	1.2	1.5
2002	*	*	*	*	*	*	1.1	1.4	1.7
2001	*	*	*	*	*	0.5	0.8	1.4	1.8
2000	*	*	*	*	*	0.6	1.1	1.4	1.5
1999	*	*	*	*	*	0.5	1.1	1.3	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	*	*	0.4	0.7	1.1
2006 5/.....	*	*	*	*	*	*	0.4	1.0	1.1
2005 5/.....	*	*	*	*	*	*	0.6	0.6	1.3
2004 5/.....	*	*	*	*	*	0.4	0.4	0.8	1.4
2003 5/.....	*	*	0.2	*	*	0.4	0.8	1.2	1.3
2002.....	*	*	*	*	0.3	0.4	0.9	1.3	1.6
2001.....	*	*	0.2	0.2	0.3	0.4	0.8	1.0	1.7
2000.....	*	0.2	*	*	0.3	0.4	0.9	1.3	1.8
1999.....	*	*	0.2	0.3	0.4	0.5	0.9	1.5	2.8

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	*	*	1.8
2006 5/.....	*	*	*	*	*	*	*	1.5	*
2005 5/.....	*	*	*	*	*	*	*	*	1.8
2004 5/.....	*	*	*	*	*	*	*	*	2.2
2003 5/.....	*	*	*	*	*	*	0.9	1.6	1.8
2002.....	*	*	*	*	0.6	0.7	0.9	1.7	2.7
2001.....	*	*	*	*	*	*	1.1	*	2.4
2000.....	*	*	*	*	*	*	1.1	1.6	3.2
1999.....	*	*	*	*	0.6	0.8	1.1	2.5	4.8

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	*	0.8
2006 5/.....	*	*	*	*	*	*	*	0.7	1.0
2005 5/.....	*	*	*	*	*	*	*	*	1.0
2004 5/.....	*	*	*	*	*	*	*	0.7	1.0
2003 5/.....	*	*	*	*	*	*	0.7	1.0	1.1
2002.....	*	*	*	*	*	*	0.9	1.0	1.2
2001.....	*	*	*	*	*	*	0.6	0.9	1.4
2000.....	*	*	*	*	*	*	0.7	1.2	1.2
1999.....	*	*	*	*	*	*	0.7	1.0	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

All origins 3/, both sexes

2007.....	4.4	6.8	10.7	15.7	26.0	41.3	63.2	101.3	174.4
2006.....	3.9	6.6	10.3	16.2	25.3	40.3	66.0	103.7	177.3
2005.....	4.0	6.6	10.5	16.1	25.7	40.8	63.6	104.8	187.3
2004.....	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
2003.....	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7

All origins 3/, male

2007.....	4.8	7.4	11.9	16.6	28.7	46.9	70.8	116.4	189.3
2006.....	4.2	7.7	11.7	17.4	27.6	43.9	75.3	117.7	194.8
2005.....	4.4	7.1	11.5	17.8	28.6	45.9	71.7	119.8	201.9
2004.....	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
2003.....	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5

All origins 3/, female

2007.....	4.0	6.2	9.7	14.9	23.6	36.7	57.6	91.8	167.3
2006.....	3.7	5.6	8.9	15.1	23.2	37.4	59.2	94.9	169.2
2005.....	3.6	6.1	9.5	14.5	23.2	36.7	57.8	95.4	180.6
2004.....	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
2003.....	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

Hispanic, both sexes

2007	3.2	4.7	8.4	10.9	20.1	34.1	46.9	84.7	124.1
2006	3.0	5.3	7.3	13.6	19.1	32.4	54.3	81.5	122.9
2005	2.8	4.8	7.7	12.5	20.5	31.9	46.5	83.4	134.8
2004	3.5	4.8	7.8	12.4	19.1	32.8	51.2	83.7	138.0
2003	3.4	4.5	7.4	11.5	19.0	26.0	50.2	81.2	151.8
2002	2.7	5.1	6.2	11.2	20.1	39.9	54.9	78.4	156.3
2001	3.6	4.3	5.4	11.8	20.5	31.5	54.1	77.0	171.3
2000	2.9	5.2	6.4	11.2	16.2	28.9	48.1	79.6	144.7
1999	2.1	4.4	6.9	9.4	20.6	31.9	52.3	78.7	180.3

Hispanic, male

2007	4.1	5.3	9.3	13.8	23.2	42.5	56.0	101.5	113.8
2006	3.5	6.8	9.1	14.7	24.5	38.5	64.4	95.8	138.7
2005	3.3	5.7	9.3	12.7	22.1	39.7	56.2	103.3	137.0
2004	4.5	6.3	8.1	13.5	22.8	33.1	56.4	91.5	147.1
2003	4.4	5.7	9.1	10.3	22.7	28.6	56.5	85.2	160.1
2002	3.8	6.9	6.4	12.6	20.7	43.9	61.4	91.5	164.0
2001	3.8	6.0	6.4	14.7	24.5	36.6	68.9	89.6	160.0
2000	4.1	6.3	8.1	12.4	22.4	32.1	58.3	94.2	157.2
1999	2.6	5.4	8.6	12.2	25.9	34.1	70.2	84.9	181.2

Hispanic, female

2007	2.2	4.1	7.5	8.3	17.5	27.6	40.3	73.4	129.6
2006	2.4	3.8	5.7	12.5	14.4	27.6	47.0	71.9	114.4
2005	2.3	4.0	6.2	12.4	19.2	25.9	39.6	70.3	133.6
2004	2.4	3.3	7.6	11.4	16.1	32.6	47.5	78.5	133.2
2003	2.4	3.3	5.9	12.4	16.0	23.9	45.7	78.5	147.7
2002	*	3.2	6.1	10.1	19.6	36.8	50.3	70.1	152.5
2001	3.4	2.6	4.5	9.4	17.2	27.6	43.6	69.2	176.9
2000	*	4.2	4.8	10.2	11.2	26.5	40.8	70.8	138.5
1999	*	3.5	5.4	6.9	16.3	30.3	39.6	75.0	179.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

Non-Hispanic 4/, both sexes

2007.....	4.5	7.0	10.9	16.1	26.4	41.7	64.2	102.2	176.8
2006.....	4.1	6.8	10.5	16.4	25.7	40.8	66.7	104.7	179.7
2005.....	4.1	6.7	10.7	16.3	26.0	41.3	64.5	105.8	189.6
2004.....	4.1	7.1	10.5	16.5	25.3	41.7	65.8	104.5	188.7
2003.....	4.3	6.7	10.5	17.0	25.6	41.7	69.3	108.7	204.1
2002.....	4.4	6.3	10.3	16.4	27.4	43.2	70.6	110.5	204.2
2001.....	4.1	6.3	10.1	15.6	25.6	41.3	67.8	104.7	206.8
2000.....	4.0	6.1	9.5	15.4	24.6	38.8	64.4	105.6	217.5
1999.....	3.9	5.7	9.1	14.7	24.8	38.5	64.1	103.4	221.2

Non-Hispanic 4/, male

2007.....	4.8	7.6	12.1	16.8	29.1	47.1	71.7	117.3	193.2
2006.....	4.3	7.7	11.9	17.5	27.8	44.2	76.0	118.8	197.5
2005.....	4.5	7.2	11.6	18.1	28.9	46.1	72.4	120.5	205.3
2004.....	4.4	7.6	11.6	17.3	27.1	47.5	75.4	122.4	206.3
2003.....	4.7	7.2	11.4	18.2	29.0	46.0	77.7	124.0	215.9
2002.....	4.8	6.8	10.8	18.1	29.5	48.3	81.2	128.1	215.6
2001.....	4.3	6.7	10.9	16.6	28.0	47.1	78.8	123.1	226.8
2000.....	4.5	7.0	10.2	16.9	26.9	43.6	74.3	122.0	224.8
1999.....	4.3	6.2	9.7	16.0	27.7	44.1	74.5	121.5	234.5

Non-Hispanic 4/, female

2007.....	4.2	6.4	9.9	15.4	24.0	37.3	58.7	92.7	169.0
2006.....	3.8	5.8	9.1	15.3	23.9	38.1	59.9	95.9	171.4
2005.....	3.8	6.3	9.8	14.6	23.4	37.4	58.8	96.7	182.5
2004.....	3.8	6.6	9.4	15.8	23.7	37.0	58.9	93.5	180.8
2003.....	3.9	6.2	9.7	15.9	22.6	38.3	63.3	99.4	199.0
2002.....	4.0	5.8	9.7	14.8	25.5	39.1	63.0	99.7	199.3
2001.....	3.8	6.0	9.3	14.8	23.5	36.6	60.2	93.6	198.5
2000.....	3.5	5.2	9.0	14.0	22.5	35.0	57.6	95.9	214.5
1999.....	3.4	5.2	8.5	13.5	22.3	34.1	56.8	92.9	215.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

Non-Hispanic White, both sexes

2007 5/.....	3.6	5.6	9.2	14.1	23.1	37.1	59.7	97.0	169.3
2006 5/.....	3.2	5.4	8.7	14.0	22.7	36.5	62.2	98.9	173.2
2005 5/.....	3.3	5.4	9.0	14.0	22.9	37.8	59.4	99.0	179.2
2004 5/.....	3.2	5.8	8.9	13.9	22.0	37.8	61.2	98.1	179.6
2003 5/.....	3.4	5.2	8.8	14.5	21.8	37.5	64.3	103.1	192.4
2002.....	3.6	4.9	8.5	13.6	23.7	38.9	64.6	104.0	192.6
2001.....	3.1	5.0	8.2	13.2	22.3	36.9	62.4	99.1	195.6
2000.....	3.0	4.8	7.9	12.7	20.9	34.8	58.2	97.7	204.3
1999.....	2.8	4.5	7.2	12.3	21.2	33.5	57.6	95.7	206.0

Non-Hispanic White, male

2007 5/.....	3.8	5.9	10.0	14.4	25.1	41.4	67.0	113.5	185.4
2006 5/.....	3.5	6.2	9.7	14.7	24.8	39.9	71.1	113.1	193.3
2005 5/.....	3.6	5.7	9.6	15.9	25.3	41.8	67.0	113.4	193.6
2004 5/.....	3.3	6.3	9.7	14.4	23.6	43.0	70.4	114.6	198.5
2003 5/.....	3.8	5.6	9.3	15.6	24.6	41.5	71.9	118.1	204.8
2002.....	4.0	5.2	9.1	15.4	25.2	43.8	73.6	119.9	203.9
2001.....	3.3	5.2	8.6	14.2	24.2	41.6	72.4	116.6	213.6
2000.....	3.4	5.3	8.4	14.1	22.9	39.7	66.7	114.0	211.2
1999.....	3.2	4.9	7.5	12.8	23.4	38.3	67.8	112.1	218.1

Non-Hispanic White, female

2007 5/.....	3.3	5.3	8.4	13.8	21.2	33.4	54.1	86.5	161.7
2006 5/.....	2.9	4.7	7.7	13.4	20.9	33.5	55.6	90.0	163.9
2005 5/.....	3.0	5.2	8.4	12.3	20.8	34.5	53.8	90.0	172.6
2004 5/.....	3.0	5.3	8.0	13.4	20.5	33.5	54.5	87.8	171.1
2003 5/.....	3.1	4.8	8.4	13.4	19.4	34.2	58.8	93.9	187.0
2002.....	3.3	4.6	7.9	11.9	22.3	34.9	58.2	94.2	187.7
2001.....	2.9	4.7	7.9	12.3	20.6	33.1	55.2	88.6	188.1
2000.....	2.5	4.2	7.3	11.4	19.1	30.8	52.1	88.1	201.6
1999.....	2.5	4.2	6.9	11.8	19.4	29.7	50.4	86.1	201.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

Non-Hispanic Black, both sexes

2007 5/.....	11.0	17.4	26.1	36.9	61.7	88.5	125.3	183.4	299.4
2006 5/.....	10.3	17.2	26.4	39.3	56.7	87.2	127.7	189.4	290.9
2005 5/.....	10.3	17.0	26.0	39.3	56.8	82.6	128.0	206.2	340.8
2004 5/.....	10.5	17.5	26.0	40.7	59.0	86.9	129.4	199.5	324.7
2003 5/.....	10.8	18.0	25.8	42.2	62.0	89.2	135.4	196.5	373.9
2002.....	9.6	17.4	27.6	41.8	63.3	92.0	148.3	210.3	370.8
2001.....	10.9	17.1	27.6	38.9	58.2	90.9	141.2	194.2	368.6
2000.....	11.3	17.3	24.7	40.5	60.8	86.7	148.0	218.1	395.9
1999.....	11.3	15.5	25.8	37.0	58.8	95.6	143.9	211.6	414.2

Non-Hispanic Black, male

2007 5/.....	12.0	20.6	31.1	43.3	74.0	107.4	143.6	197.6	336.8
2006 5/.....	10.4	20.9	32.2	46.5	62.6	96.2	157.0	214.2	300.5
2005 5/.....	11.2	20.2	31.4	44.4	69.1	98.8	146.5	241.0	379.7
2004 5/.....	12.0	19.0	31.2	45.4	66.3	105.1	156.7	248.9	346.5
2003 5/.....	12.1	20.3	32.7	46.8	76.2	101.1	162.6	237.7	409.1
2002.....	10.5	20.6	29.3	46.2	73.2	106.7	185.4	260.5	399.3
2001.....	12.0	19.1	34.0	41.8	67.5	112.6	173.0	237.9	430.8
2000.....	12.8	21.9	27.9	45.9	69.8	96.7	185.0	251.5	432.1
1999.....	13.2	18.1	30.1	47.7	72.8	116.0	165.8	270.1	451.6

Non-Hispanic Black, female

2007 5/.....	10.2	14.7	22.0	31.9	52.7	75.5	114.3	175.9	283.9
2006 5/.....	10.2	14.0	21.7	33.6	52.4	81.0	110.2	176.3	287.0
2005 5/.....	9.5	14.4	21.6	35.3	47.7	71.6	117.0	187.8	325.2
2004 5/.....	9.1	16.1	21.7	37.0	53.6	74.6	113.2	173.5	316.0
2003 5/.....	9.6	15.9	20.2	38.5	51.4	81.1	119.3	174.9	360.3
2002.....	8.9	14.8	26.3	38.3	56.0	82.1	126.1	184.2	359.9
2001.....	10.0	15.5	22.4	36.6	51.4	76.4	122.2	172.0	345.6
2000.....	10.0	13.4	22.0	36.2	54.1	80.1	125.7	201.3	382.5
1999.....	9.6	13.3	22.2	28.7	48.4	82.0	130.8	182.6	400.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute poliomyelitis (A80)

Non-Hispanic White, both sexes

Table with 10 columns (Year, 45-49, 50-54, 55-59, 60-64, 65-69, 70-74, 75-79, 80-84, 85+) and 8 rows of data for years 1999-2007, all cells containing an asterisk.

Non-Hispanic White, male

Table with 10 columns (Year, 45-49, 50-54, 55-59, 60-64, 65-69, 70-74, 75-79, 80-84, 85+) and 8 rows of data for years 1999-2007, all cells containing an asterisk.

Non-Hispanic White, female

Table with 10 columns (Year, 45-49, 50-54, 55-59, 60-64, 65-69, 70-74, 75-79, 80-84, 85+) and 8 rows of data for years 1999-2007, all cells containing an asterisk.

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

All origins 3/, both sexes

2007.....	4.5	8.5	9.0	5.3	4.3	4.7	4.3	4.4	2.9
2006.....	4.8	8.9	8.0	5.0	4.5	4.5	5.0	4.0	3.1
2005.....	4.2	6.8	5.3	3.5	3.2	3.8	3.6	3.8	2.4
2004.....	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
2003.....	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7

All origins 3/, male

2007.....	6.3	12.5	13.9	7.7	5.4	4.9	4.4	5.1	3.0
2006.....	6.8	13.4	11.9	7.0	5.2	4.7	5.1	3.8	3.0
2005.....	6.3	10.2	8.2	4.7	3.5	4.0	3.3	4.2	2.6
2004.....	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
2003.....	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9

All origins 3/, female

2007.....	2.7	4.8	4.4	3.2	3.3	4.6	4.3	4.0	2.9
2006.....	2.8	4.6	4.2	3.3	3.8	4.3	5.0	4.1	3.1
2005.....	2.2	3.6	2.7	2.3	2.9	3.7	3.8	3.6	2.3
2004.....	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
2003.....	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Hispanic, both sexes

2007	6.5	10.6	14.9	11.1	10.6	11.4	8.8	9.3	*
2006	7.0	12.5	14.5	10.7	10.7	10.9	11.2	8.5	8.6
2005	4.8	9.6	9.3	7.4	8.1	9.2	7.2	*	*
2004	5.8	8.2	7.4	8.1	7.3	6.7	7.7	*	*
2003	5.7	8.6	7.7	6.6	8.6	8.7	9.8	8.2	*
2002	6.8	9.4	8.1	8.6	8.8	8.4	11.5	8.9	*
2001	8.1	7.5	11.1	9.5	8.2	12.1	12.2	*	*
2000	6.7	7.3	7.0	10.7	10.3	12.2	12.9	13.4	*
1999	5.9	8.1	7.6	8.0	11.7	13.3	13.6	13.5	*

Hispanic, male

2007	9.9	14.7	22.8	16.0	12.2	13.6	*	*	*
2006	10.0	19.2	20.2	13.7	12.1	11.8	10.3	*	*
2005	7.5	14.0	12.8	9.4	10.0	*	*	*	*
2004	8.6	12.3	11.3	9.1	9.6	*	*	*	*
2003	8.8	13.5	11.5	8.9	9.1	*	*	*	*
2002	10.8	13.5	11.9	11.0	7.4	*	12.8	*	*
2001	12.4	12.1	17.0	10.3	*	*	*	*	*
2000	10.3	11.1	9.4	14.7	10.4	11.7	*	*	*
1999	9.1	12.3	9.9	9.5	12.4	13.7	*	*	*

Hispanic, female

2007	2.8	6.4	7.3	6.7	9.3	9.8	9.7	*	*
2006	3.8	5.8	9.2	8.0	9.5	10.1	11.9	10.9	*
2005	1.9	5.2	5.9	5.6	6.4	10.5	7.8	*	*
2004	2.8	4.2	3.7	7.3	5.4	7.3	9.8	*	*
2003	2.5	3.8	4.1	4.7	8.3	10.8	10.8	*	*
2002	2.8	5.5	4.6	6.5	10.0	8.8	10.5	*	*
2001	3.8	3.2	5.8	8.7	10.0	15.0	13.6	*	*
2000	3.2	3.6	4.8	7.2	10.3	12.5	13.1	*	*
1999	2.7	4.1	5.4	6.7	11.3	13.0	15.4	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Non-Hispanic 4/, both sexes

2007.....	4.2	8.3	8.4	4.8	3.8	4.2	4.0	4.1	2.8
2006.....	4.5	8.5	7.4	4.5	3.9	3.9	4.6	3.7	2.8
2005.....	4.1	6.5	5.0	3.1	2.8	3.4	3.3	3.7	2.2
2004.....	4.1	6.3	4.6	3.2	3.0	3.4	3.3	3.4	2.6
2003.....	4.7	6.1	4.2	2.8	3.0	3.9	4.0	3.9	2.3
2002.....	5.4	5.7	3.9	3.3	3.5	4.2	4.4	4.2	3.7
2001.....	5.0	5.0	3.5	3.4	4.0	4.4	4.7	4.4	3.6
2000.....	5.0	4.6	3.4	3.3	3.9	4.5	5.1	4.9	3.8
1999.....	4.3	3.4	3.0	2.9	4.0	4.7	4.9	5.1	4.7

Non-Hispanic 4/, male

2007.....	5.8	12.2	13.0	7.0	4.9	4.3	4.2	4.9	2.9
2006.....	6.3	12.7	11.2	6.4	4.7	4.1	4.8	3.7	3.0
2005.....	6.1	9.8	7.7	4.3	3.0	3.8	3.1	4.1	2.4
2004.....	6.1	9.6	7.0	4.2	3.7	3.6	3.8	3.2	3.0
2003.....	6.9	9.3	6.2	3.5	3.3	4.7	4.5	4.9	2.7
2002.....	8.1	9.1	5.6	4.1	4.1	4.5	5.3	4.7	4.8
2001.....	7.7	7.8	5.0	4.0	4.6	4.7	4.8	5.3	4.3
2000.....	7.7	7.0	4.8	3.9	4.2	4.6	6.2	5.9	5.4
1999.....	6.5	5.3	4.3	3.4	4.4	4.9	5.9	7.4	6.8

Non-Hispanic 4/, female

2007.....	2.6	4.6	4.1	2.8	2.8	4.1	3.9	3.7	2.7
2006.....	2.7	4.5	3.8	2.8	3.3	3.8	4.5	3.8	2.7
2005.....	2.2	3.4	2.4	2.1	2.6	3.2	3.5	3.5	2.2
2004.....	2.2	3.2	2.4	2.2	2.3	3.2	3.0	3.5	2.4
2003.....	2.6	3.0	2.3	2.2	2.7	3.4	3.7	3.2	2.2
2002.....	2.8	2.6	2.3	2.6	2.9	4.1	3.8	3.9	3.2
2001.....	2.4	2.3	2.2	2.8	3.4	4.3	4.6	3.9	3.3
2000.....	2.4	2.3	2.0	2.7	3.6	4.4	4.3	4.3	3.2
1999.....	2.1	1.5	1.8	2.5	3.6	4.5	4.2	3.7	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Non-Hispanic White, both sexes

2007 5/.....	4.2	8.2	7.7	4.2	3.3	3.6	3.5	3.7	2.5
2006 5/.....	4.5	8.3	6.4	3.9	3.1	3.4	4.0	3.2	2.5
2005 5/.....	4.1	6.3	4.5	2.6	2.2	2.7	3.0	3.5	2.0
2004 5/.....	4.1	5.9	3.9	2.7	2.6	2.9	3.0	2.9	2.4
2003 5/.....	4.7	5.7	3.4	2.3	2.4	3.5	3.6	3.6	2.2
2002.....	5.2	5.4	3.3	2.7	2.9	3.8	4.1	4.0	3.5
2001.....	4.8	4.4	3.0	2.9	3.4	3.9	4.4	4.2	3.3
2000.....	4.7	3.9	2.7	2.7	3.3	4.1	4.9	4.6	3.7
1999.....	3.9	2.9	2.4	2.5	3.4	4.1	4.5	5.0	4.5

Non-Hispanic White, male

2007 5/.....	5.8	12.1	11.9	6.0	4.4	3.5	3.8	4.4	2.8
2006 5/.....	6.4	12.4	9.8	5.5	3.7	3.6	4.2	3.5	3.1
2005 5/.....	6.1	9.4	7.0	3.5	2.4	3.1	3.1	3.8	2.2
2004 5/.....	6.0	9.0	5.9	3.5	3.2	3.2	3.6	2.6	2.7
2003 5/.....	6.9	8.6	5.1	2.9	2.8	4.1	3.9	4.7	2.7
2002.....	7.9	8.5	4.7	3.3	3.7	4.1	5.1	4.7	4.6
2001.....	7.4	7.0	4.3	3.6	4.3	4.4	4.9	5.4	4.4
2000.....	7.3	6.1	3.9	3.3	3.6	4.2	6.0	5.6	5.4
1999.....	5.7	4.5	3.6	3.1	4.0	4.6	5.6	7.7	6.5

Non-Hispanic White, female

2007 5/.....	2.7	4.4	3.6	2.4	2.3	3.7	3.3	3.2	2.4
2006 5/.....	2.6	4.2	3.2	2.4	2.6	3.3	3.8	3.0	2.2
2005 5/.....	2.1	3.2	2.0	1.7	2.0	2.3	2.9	3.3	1.9
2004 5/.....	2.2	2.8	2.0	1.9	2.1	2.6	2.6	3.1	2.3
2003 5/.....	2.5	2.8	1.8	1.7	2.1	2.9	3.4	2.9	2.0
2002.....	2.6	2.4	2.0	2.2	2.2	3.5	3.4	3.6	3.1
2001.....	2.2	1.8	1.7	2.3	2.7	3.4	4.1	3.5	2.9
2000.....	2.1	1.8	1.5	2.3	3.2	4.0	4.0	4.0	3.1
1999.....	2.1	1.3	1.4	1.9	2.9	3.7	3.8	3.4	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Non-Hispanic Black, both sexes

2007 5/.....	4.4	10.7	14.5	9.0	5.7	6.0	5.3	6.0	*
2006 5/.....	4.9	11.3	14.8	9.2	9.1	5.0	6.7	6.8	*
2005 5/.....	4.9	9.9	9.3	7.4	5.9	6.5	5.1	5.0	*
2004 5/.....	5.2	10.5	10.4	6.6	4.4	5.0	3.8	6.8	*
2003 5/.....	5.3	10.3	9.6	6.4	5.2	5.6	6.3	5.4	*
2002.....	7.3	9.2	7.8	6.3	6.1	5.8	6.0	*	*
2001.....	6.8	9.6	7.3	6.2	5.1	6.2	6.2	*	*
2000.....	7.4	8.8	6.9	4.5	5.3	6.3	5.8	7.2	*
1999.....	7.2	6.8	5.7	4.3	6.7	6.9	6.5	*	*

Non-Hispanic Black, male

2007 5/.....	6.0	15.9	23.2	14.7	7.8	8.4	*	*	*
2006 5/.....	6.4	16.7	23.8	14.2	11.6	6.3	9.2	*	*
2005 5/.....	6.9	14.5	14.4	10.7	6.9	7.8	*	*	*
2004 5/.....	8.2	16.4	17.0	10.0	7.0	*	*	*	*
2003 5/.....	7.8	16.5	15.3	7.7	5.6	*	*	*	*
2002.....	10.6	15.1	12.6	8.1	7.9	*	*	*	*
2001.....	10.1	14.3	10.1	7.5	*	*	*	*	*
2000.....	11.2	13.8	10.3	5.3	6.9	7.5	*	*	*
1999.....	12.2	11.4	7.7	4.6	5.6	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	3.1	6.3	7.4	4.5	4.2	4.3	*	*	*
2006 5/.....	3.5	6.8	7.3	5.3	7.3	*	*	7.8	*
2005 5/.....	3.1	5.9	5.1	4.8	5.3	5.6	6.8	*	*
2004 5/.....	2.6	5.5	4.9	3.9	*	*	*	*	*
2003 5/.....	3.2	5.1	4.9	5.4	4.9	5.0	5.5	*	*
2002.....	4.5	4.3	3.9	4.9	4.7	5.7	5.6	*	*
2001.....	3.9	5.6	4.9	5.1	5.4	6.7	7.4	*	*
2000.....	4.2	4.5	4.0	3.9	4.1	5.4	*	*	*
1999.....	2.9	2.9	4.1	4.1	7.5	7.6	6.7	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes

2007.....	9.9	9.0	6.0	4.3	2.8	1.7	1.0	0.6	*
2006.....	10.9	9.3	6.5	4.1	3.1	1.8	1.0	0.6	*
2005.....	11.3	9.9	6.3	4.0	2.8	1.8	1.0	0.5	*
2004.....	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
2003.....	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All origins 3/, male

2007.....	14.4	13.6	9.5	7.3	5.0	3.0	2.1	1.0	*
2006.....	16.1	14.4	10.3	6.9	5.0	3.1	1.9	1.1	*
2005.....	17.2	15.4	10.5	6.6	4.8	3.1	1.8	*	*
2004.....	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
2003.....	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All origins 3/, female

2007.....	5.6	4.5	2.8	1.6	0.8	0.7	*	*	*
2006.....	5.7	4.3	2.8	1.5	1.4	0.8	*	*	*
2005.....	5.6	4.6	2.4	1.6	1.1	0.7	*	*	*
2004.....	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
2003.....	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, both sexes

2007	10.8	10.8	7.2	7.0	6.2	*	*	*	*
2006	11.8	11.5	7.8	7.4	5.2	4.0	*	*	*
2005	13.8	12.0	8.6	5.2	5.0	3.5	*	*	*
2004	14.3	13.9	9.6	6.6	6.4	*	*	*	*
2003	17.0	14.4	12.3	6.2	6.2	4.1	*	*	*
2002	15.4	13.7	10.8	7.8	4.3	*	*	*	*
2001	16.4	14.2	10.5	7.3	6.0	*	*	*	*
2000	18.1	15.4	10.0	8.9	4.7	*	*	*	*
1999	17.1	13.1	9.8	6.6	6.0	*	*	*	*

Hispanic, male

2007	16.4	17.5	10.6	12.1	11.3	*	*	*	*
2006	18.9	18.2	12.8	13.1	8.2	7.6	*	*	*
2005	22.5	18.5	14.3	10.1	8.3	*	*	*	*
2004	22.1	21.9	15.6	12.6	9.6	*	*	*	*
2003	27.9	23.7	20.2	8.6	10.1	*	*	*	*
2002	24.0	21.8	17.4	11.5	8.5	*	*	*	*
2001	25.0	24.8	17.6	11.1	11.3	*	*	*	*
2000	28.4	26.2	17.3	12.7	8.2	*	*	*	*
1999	26.8	20.8	17.6	12.5	10.8	*	*	*	*

Hispanic, female

2007	4.9	4.1	4.0	*	*	*	*	*	*
2006	4.3	4.8	3.1	*	*	*	*	*	*
2005	4.8	5.6	3.2	*	*	*	*	*	*
2004	6.3	6.0	4.0	*	*	*	*	*	*
2003	5.8	5.4	5.1	*	*	*	*	*	*
2002	6.7	5.9	4.7	4.5	*	*	*	*	*
2001	7.8	4.1	4.2	*	*	*	*	*	*
2000	7.9	5.0	*	5.7	*	*	*	*	*
1999	7.5	5.8	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 4/, both sexes

2007	9.7	8.7	5.9	4.0	2.5	1.6	0.9	0.6	*
2006	10.6	9.0	6.3	3.7	2.9	1.7	0.9	0.5	*
2005	11.0	9.5	6.1	3.8	2.7	1.6	1.0	0.4	*
2004	11.1	9.0	5.9	3.9	2.8	1.4	0.9	0.4	*
2003	11.5	8.9	5.9	3.9	2.5	1.7	0.9	0.4	*
2002	12.1	9.0	5.5	3.6	2.7	1.4	0.9	0.4	*
2001	11.5	8.1	5.3	4.1	2.4	1.3	0.8	*	*
2000	12.0	8.4	5.5	3.7	2.4	1.6	0.8	*	*
1999	11.7	8.1	5.5	3.1	2.4	1.4	0.7	*	*

Non-Hispanic 4/, male

2007	14.0	13.1	9.3	6.8	4.5	2.8	1.9	*	*
2006	15.6	13.9	10.1	6.3	4.7	2.8	1.7	*	*
2005	16.4	14.9	10.1	6.2	4.6	2.8	1.7	*	*
2004	16.8	13.9	9.9	6.5	4.6	2.3	1.5	*	*
2003	17.5	14.3	9.7	6.7	4.1	3.0	1.7	*	*
2002	18.6	14.9	9.1	6.3	4.8	2.4	1.7	*	*
2001	18.0	13.2	9.1	6.6	3.6	2.5	1.6	*	*
2000	19.2	13.8	9.3	6.5	4.1	2.7	1.6	*	*
1999	18.9	13.6	9.5	5.2	4.3	2.5	1.2	*	*

Non-Hispanic 4/, female

2007	5.6	4.5	2.6	1.5	0.8	0.6	*	*	*
2006	5.8	4.3	2.8	1.4	1.3	0.7	*	*	*
2005	5.6	4.4	2.3	1.6	1.0	0.6	*	*	*
2004	5.6	4.3	2.1	1.5	1.2	0.6	0.5	*	*
2003	5.6	3.7	2.2	1.4	1.2	0.7	*	*	*
2002	5.8	3.4	2.0	1.2	1.0	0.6	*	*	*
2001	5.1	3.2	1.8	1.8	1.4	*	*	*	*
2000	5.1	3.2	1.9	1.2	0.9	0.7	*	*	*
1999	4.8	2.8	1.9	1.2	0.9	0.6	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic White, both sexes

2007 5/.....	4.3	3.5	2.5	1.6	1.2	0.7	0.5	*	*
2006 5/.....	4.6	3.9	2.6	1.7	1.3	0.6	0.4	*	*
2005 5/.....	4.8	4.3	2.6	1.6	1.1	0.7	0.5	*	*
2004 5/.....	4.9	3.7	2.7	1.6	1.1	0.6	0.4	*	*
2003 5/.....	5.2	3.7	2.6	1.7	0.9	0.8	0.5	*	*
2002.....	5.5	3.5	2.6	1.7	1.0	0.5	0.4	*	*
2001.....	5.0	3.4	2.6	1.5	0.8	0.6	0.3	*	*
2000.....	5.5	3.7	2.7	1.5	1.0	0.6	0.4	*	*
1999.....	5.2	3.9	2.4	1.4	1.1	0.5	0.3	*	*

Non-Hispanic White, male

2007 5/.....	7.1	5.9	4.3	2.7	2.4	1.4	1.1	*	*
2006 5/.....	8.1	6.6	4.5	3.2	2.3	1.2	0.8	*	*
2005 5/.....	8.1	7.1	4.7	2.7	2.0	1.4	0.9	*	*
2004 5/.....	8.3	6.4	4.9	2.9	2.1	1.2	0.8	*	*
2003 5/.....	8.9	6.6	4.6	3.2	1.5	1.5	1.1	*	*
2002.....	9.5	6.3	4.7	3.2	1.9	1.0	*	*	*
2001.....	8.8	6.2	4.8	2.6	1.4	1.4	*	*	*
2000.....	9.6	6.5	4.9	2.8	2.0	0.9	0.8	*	*
1999.....	9.0	7.2	4.4	2.6	2.1	0.9	*	*	*

Non-Hispanic White, female

2007 5/.....	1.6	1.2	0.7	0.5	*	*	*	*	*
2006 5/.....	1.2	1.3	0.7	*	*	*	*	*	*
2005 5/.....	1.5	1.4	0.7	0.5	*	*	*	*	*
2004 5/.....	1.5	1.0	0.5	*	*	*	*	*	*
2003 5/.....	1.6	0.9	0.7	*	*	*	*	*	*
2002.....	1.6	0.8	0.6	*	*	*	*	*	*
2001.....	1.2	0.6	0.6	0.4	*	*	*	*	*
2000.....	1.4	0.8	0.7	*	*	*	*	*	*
1999.....	1.3	0.8	0.5	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic Black, both sexes

2007 5/.....	45.1	45.6	32.3	26.7	14.6	10.2	6.0	*	*
2006 5/.....	50.4	45.7	36.7	22.0	17.0	10.8	5.9	*	*
2005 5/.....	52.2	48.5	34.6	24.4	16.8	10.4	6.2	*	*
2004 5/.....	53.0	48.2	33.6	24.8	17.5	8.6	6.4	*	*
2003 5/.....	54.5	47.5	34.4	23.9	17.1	10.9	4.9	*	*
2002.....	58.4	50.5	30.8	20.5	17.8	10.6	7.2	*	*
2001.....	57.6	45.0	29.3	26.3	16.2	8.5	6.4	*	*
2000.....	59.5	46.9	29.6	22.2	14.7	11.5	6.0	*	*
1999.....	59.6	42.8	33.0	18.4	14.9	11.2	4.9	*	*

Non-Hispanic Black, male

2007 5/.....	62.3	68.0	51.3	48.7	26.2	18.4	12.3	*	*
2006 5/.....	69.6	71.1	58.8	37.2	27.8	19.3	12.4	*	*
2005 5/.....	76.4	76.4	59.1	41.6	30.1	18.2	11.9	*	*
2004 5/.....	78.3	73.9	56.8	42.4	29.4	15.3	9.8	*	*
2003 5/.....	81.2	76.3	58.5	41.7	29.6	19.5	*	*	*
2002.....	87.2	84.9	50.9	36.7	32.0	19.5	15.0	*	*
2001.....	88.7	72.3	50.7	43.7	25.8	15.5	12.8	*	*
2000.....	93.8	77.5	51.3	40.2	25.3	21.9	11.6	*	*
1999.....	96.6	71.1	57.7	30.5	26.3	21.1	*	*	*

Non-Hispanic Black, female

2007 5/.....	30.2	26.6	16.8	9.5	6.1	4.5	*	*	*
2006 5/.....	33.8	24.2	18.5	10.2	9.1	5.0	*	*	*
2005 5/.....	31.2	24.9	14.4	10.9	6.9	5.1	*	*	*
2004 5/.....	31.1	26.6	14.6	11.1	8.7	*	*	*	*
2003 5/.....	31.3	23.2	14.6	10.0	7.9	5.0	*	*	*
2002.....	33.4	21.4	14.2	8.0	7.2	4.6	*	*	*
2001.....	30.6	21.7	11.7	12.7	9.0	*	*	*	*
2000.....	29.8	20.8	11.8	8.1	6.9	4.5	*	*	*
1999.....	27.7	18.8	12.8	8.9	6.5	4.6	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/, both sexes

2007.....	0.9	1.4	2.2	3.5	4.8	6.7	9.5	13.5	20.0
2006.....	1.0	1.6	2.3	3.3	4.8	7.1	9.8	13.8	20.4
2005.....	2.4	4.2	3.9	4.5	5.9	8.3	12.0	15.8	21.2
2004.....	2.5	3.5	4.2	4.4	6.0	8.3	11.2	15.9	22.5
2003.....	2.4	3.5	3.7	4.5	5.9	8.9	12.3	14.8	22.5
2002.....	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001.....	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000.....	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
1999.....	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0

All origins 3/, male

2007.....	1.1	1.9	2.5	4.0	5.2	8.4	10.9	17.2	25.3
2006.....	1.2	2.1	2.8	3.8	5.6	8.1	11.6	17.2	25.1
2005.....	3.3	5.8	5.2	5.3	7.0	10.0	14.1	19.6	23.6
2004.....	3.3	4.9	5.6	5.2	7.2	9.9	12.9	18.4	29.6
2003.....	3.3	4.7	4.9	5.4	7.0	10.3	13.9	18.8	26.5
2002.....	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001.....	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000.....	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
1999.....	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9

All origins 3/, female

2007.....	0.8	1.0	2.0	3.1	4.5	5.4	8.4	11.2	17.5
2006.....	0.8	1.2	1.8	2.9	4.2	6.3	8.4	11.7	18.2
2005.....	1.5	2.5	2.7	3.8	5.0	7.0	10.5	13.4	20.1
2004.....	1.6	2.2	2.8	3.6	5.0	7.0	10.0	14.4	19.3
2003.....	1.6	2.2	2.6	3.7	5.0	7.8	11.1	12.4	20.8
2002.....	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001.....	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000.....	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
1999.....	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, both sexes

2007.....	0.9	2.0	2.1	4.1	4.7	5.6	7.9	13.3	15.4
2006.....	1.1	1.3	1.7	2.7	3.7	7.2	7.1	9.2	10.2
2005.....	3.2	7.0	5.5	5.6	7.2	10.2	13.1	13.6	15.4
2004.....	3.5	5.3	6.6	5.2	6.1	9.6	14.4	15.0	18.6
2003.....	3.3	5.9	4.7	6.8	8.4	8.5	15.1	15.6	13.0
2002.....	2.4	4.9	4.6	3.9	6.2	6.5	8.8	15.6	17.7
2001.....	2.1	2.0	3.2	2.8	4.9	7.1	10.5	9.9	15.0
2000.....	1.7	2.3	3.1	3.2	4.5	7.3	9.5	14.5	19.2
1999.....	2.0	2.7	3.0	4.8	8.1	7.2	13.9	14.7	26.0

Hispanic, male

2007.....	*	2.9	2.6	4.2	*	*	*	16.3	*
2006.....	1.6	2.0	*	*	5.6	*	*	*	*
2005.....	4.4	10.5	7.7	6.6	8.0	9.9	13.5	*	*
2004.....	5.0	7.4	8.8	7.7	7.7	10.3	16.1	19.8	*
2003.....	4.9	9.4	7.5	7.9	12.0	*	16.4	*	*
2002.....	3.4	7.6	6.2	*	8.1	*	*	*	*
2001.....	2.7	2.9	5.0	*	*	*	5.0	*	*
2000.....	*	*	4.6	*	*	*	*	*	*
1999.....	*	3.7	*	6.0	9.3	*	16.4	*	*

Hispanic, female

2007.....	*	*	*	4.0	5.2	*	8.3	11.4	13.2
2006.....	*	*	*	*	*	7.4	*	*	*
2005.....	1.9	3.7	3.5	4.8	6.6	10.5	12.8	14.5	17.6
2004.....	2.0	3.3	4.6	*	*	8.9	13.1	*	19.4
2003.....	*	2.6	*	5.8	5.3	9.8	14.2	14.9	*
2002.....	*	*	*	*	*	*	*	14.6	*
2001.....	*	*	*	*	*	*	9.7	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	7.2	8.0	12.1	*	28.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 4/, both sexes

2007.....	0.9	1.4	2.2	3.5	4.8	6.8	9.6	13.5	20.3
2006.....	1.0	1.6	2.3	3.4	4.9	7.1	9.9	14.0	20.9
2005.....	2.3	3.9	3.8	4.4	5.8	8.2	12.0	15.9	21.5
2004.....	2.3	3.4	4.0	4.3	6.0	8.2	11.0	15.9	22.6
2003.....	2.3	3.2	3.6	4.3	5.7	8.9	12.1	14.8	22.9
2002.....	1.8	2.5	2.8	3.8	5.4	8.2	11.7	13.7	21.8
2001.....	1.4	1.9	2.3	4.0	5.7	6.9	9.3	12.4	20.3
2000.....	1.3	1.7	2.3	3.6	4.7	7.0	9.9	12.8	21.2
1999.....	2.0	2.6	3.3	4.4	6.7	9.0	12.5	16.7	27.0

Non-Hispanic 4/, male

2007.....	1.1	1.8	2.5	4.0	5.3	8.5	11.1	17.2	25.6
2006.....	1.2	2.0	2.9	3.9	5.6	8.2	11.8	17.3	25.8
2005.....	3.1	5.3	4.9	5.2	6.9	10.0	14.1	20.0	24.2
2004.....	3.1	4.7	5.3	5.0	7.1	9.9	12.8	18.2	30.2
2003.....	3.1	4.3	4.7	5.2	6.6	10.5	13.7	18.9	27.0
2002.....	2.6	3.4	3.6	4.7	6.4	9.8	14.1	17.7	26.8
2001.....	2.0	2.3	2.9	4.6	6.2	7.3	11.4	16.0	24.0
2000.....	1.6	2.2	2.6	4.0	5.9	8.8	12.1	15.9	27.8
1999.....	2.4	2.9	3.5	5.1	7.9	10.7	15.2	20.6	32.3

Non-Hispanic 4/, female

2007.....	0.8	1.0	2.0	3.0	4.4	5.4	8.4	11.1	17.7
2006.....	0.9	1.2	1.8	2.9	4.3	6.2	8.6	12.0	18.6
2005.....	1.5	2.4	2.7	3.7	4.9	6.7	10.4	13.4	20.2
2004.....	1.6	2.1	2.7	3.6	5.0	6.9	9.8	14.5	19.2
2003.....	1.6	2.2	2.7	3.5	5.0	7.6	10.9	12.2	21.2
2002.....	1.1	1.6	2.1	3.0	4.5	6.9	10.0	11.3	19.7
2001.....	0.9	1.5	1.8	3.4	5.2	6.5	7.9	10.2	18.8
2000.....	1.1	1.2	2.1	3.2	3.8	5.6	8.4	10.9	18.6
1999.....	1.6	2.3	3.0	3.7	5.7	7.6	10.7	14.4	24.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic White, both sexes

2007 5/.....	0.7	1.3	2.1	3.3	4.7	6.8	9.4	13.5	20.8
2006 5/.....	0.8	1.5	2.2	3.2	4.8	7.1	10.0	14.0	21.1
2005 5/.....	2.1	3.7	3.6	4.1	5.8	7.8	11.6	15.7	21.5
2004 5/.....	2.3	3.3	3.7	4.0	5.7	7.8	10.7	15.6	22.6
2003 5/.....	2.2	3.0	3.3	4.0	5.6	8.9	11.5	14.6	23.4
2002.....	1.7	2.2	2.4	3.5	5.3	8.0	11.4	13.7	22.2
2001.....	1.3	1.7	2.0	3.7	5.5	6.6	9.2	12.2	20.2
2000.....	1.2	1.5	2.2	3.2	4.6	6.8	9.8	12.6	21.0
1999.....	1.5	2.1	2.9	3.9	6.3	8.6	11.9	16.1	26.4

Non-Hispanic White, male

2007 5/.....	0.8	1.6	2.4	3.8	4.9	8.4	11.0	17.3	26.1
2006 5/.....	1.0	1.8	2.6	3.5	5.6	8.1	11.6	17.3	26.5
2005 5/.....	2.9	5.3	4.6	4.9	6.8	9.8	13.7	19.8	24.4
2004 5/.....	3.1	4.5	4.9	4.6	7.0	9.3	12.4	17.9	30.8
2003 5/.....	3.0	3.9	4.3	4.8	6.4	10.2	13.2	18.6	27.6
2002.....	2.4	3.0	2.9	4.3	6.4	9.6	13.8	17.5	27.2
2001.....	1.8	2.1	2.5	4.3	5.9	7.1	11.3	16.0	24.2
2000.....	1.5	1.9	2.5	3.6	5.6	8.6	11.6	15.8	27.0
1999.....	2.0	2.4	3.3	4.4	7.3	10.4	14.7	18.9	30.8

Non-Hispanic White, female

2007 5/.....	0.6	0.9	1.9	2.8	4.5	5.4	8.2	11.0	18.3
2006 5/.....	0.7	1.2	1.8	2.9	4.1	6.2	8.8	11.9	18.6
2005 5/.....	1.3	2.2	2.6	3.5	4.8	6.2	9.9	13.1	20.1
2004 5/.....	1.5	2.1	2.6	3.4	4.5	6.7	9.4	14.2	18.9
2003 5/.....	1.4	2.0	2.4	3.3	4.9	7.8	10.3	12.2	21.6
2002.....	1.0	1.4	1.9	2.8	4.3	6.7	9.8	11.4	20.0
2001.....	0.7	1.4	1.5	3.1	5.1	6.2	7.7	10.0	18.6
2000.....	1.0	1.0	1.9	2.8	3.7	5.3	8.5	10.8	18.6
1999.....	1.1	1.9	2.5	3.4	5.4	7.2	10.0	14.4	24.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic Black, both sexes

2007 5/.....	2.1	2.4	3.2	6.0	6.7	6.9	11.3	14.8	17.4
2006 5/.....	2.1	3.3	4.2	4.8	6.7	8.4	11.0	14.3	22.6
2005 5/.....	3.5	5.0	5.4	6.7	6.4	9.6	13.7	17.4	22.7
2004 5/.....	3.0	4.0	5.9	6.9	7.4	10.5	15.1	19.2	24.8
2003 5/.....	3.6	5.3	6.1	6.3	5.8	8.4	16.5	16.1	19.9
2002.....	3.2	4.9	6.2	5.5	5.4	9.9	15.3	11.5	21.8
2001.....	2.6	3.5	4.5	6.2	7.1	9.8	9.9	13.5	20.9
2000.....	2.7	3.5	4.0	6.0	6.2	10.1	11.6	16.1	24.2
1999.....	5.2	6.0	6.5	8.2	10.9	13.3	19.1	23.8	34.3

Non-Hispanic Black, male

2007 5/.....	2.4	3.3	3.3	6.8	9.9	9.7	12.3	16.6	24.4
2006 5/.....	2.5	4.8	5.8	6.6	6.3	10.1	15.7	15.4	21.1
2005 5/.....	4.4	6.1	8.0	8.4	8.3	10.7	19.3	20.1	23.8
2004 5/.....	4.2	6.0	8.9	9.4	7.5	15.0	19.1	22.0	30.5
2003 5/.....	4.7	7.1	7.8	8.5	8.6	13.1	16.8	22.6	23.4
2002.....	4.5	7.0	9.5	7.5	6.7	12.3	18.3	*	28.4
2001.....	3.7	4.4	5.5	7.5	9.6	10.0	10.4	16.4	*
2000.....	3.3	4.8	4.6	7.4	8.8	12.6	16.9	19.7	31.7
1999.....	6.0	6.6	5.9	10.4	12.5	16.3	22.2	42.5	46.4

Non-Hispanic Black, female

2007 5/.....	1.8	1.6	3.1	5.3	4.3	4.9	10.7	13.8	14.5
2006 5/.....	1.8	2.0	2.9	3.4	6.9	7.2	8.2	13.7	23.2
2005 5/.....	2.7	4.1	3.3	5.4	4.9	8.9	10.4	15.9	22.2
2004 5/.....	1.9	2.3	3.5	4.9	7.4	7.4	12.7	17.8	22.5
2003 5/.....	2.6	3.7	4.6	4.5	3.8	5.2	16.4	12.6	18.6
2002.....	2.0	3.2	3.4	4.0	4.6	8.3	13.5	9.7	19.2
2001.....	1.6	2.8	3.7	5.1	5.2	9.6	9.6	12.1	20.5
2000.....	2.1	2.3	3.5	4.9	4.3	8.4	8.4	14.3	21.4
1999.....	4.5	5.5	7.0	6.5	9.6	11.3	17.2	14.5	29.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Malignant neoplasms (C00-C97)

All origins 3/, both sexes

2007.....	82.1	149.4	249.4	398.6	610.3	847.0	1,133.0	1,414.7	1,590.2
2006.....	84.1	152.1	252.7	414.6	615.1	863.3	1,144.9	1,418.8	1,606.1
2005.....	86.5	154.8	255.6	422.1	622.2	886.3	1,158.4	1,427.7	1,637.7
2004.....	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
2003.....	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8

All origins 3/, male

2007.....	78.8	160.0	279.6	459.3	720.6	1,027.9	1,431.9	1,878.7	2,249.2
2006.....	80.7	162.1	282.3	476.3	731.6	1,045.7	1,442.3	1,901.9	2,248.7
2005.....	83.4	164.9	284.8	484.5	743.6	1,091.8	1,475.1	1,901.6	2,319.3
2004.....	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
2003.....	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2

All origins 3/, female

2007.....	85.5	139.2	220.9	342.8	513.8	699.2	912.8	1,121.7	1,276.7
2006.....	87.5	142.6	224.8	358.2	513.3	714.8	927.2	1,116.9	1,305.5
2005.....	89.6	145.1	228.1	365.2	516.2	719.8	928.5	1,133.4	1,324.6
2004.....	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
2003.....	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

Hispanic, both sexes

2007.....	56.6	98.3	162.0	248.1	392.5	530.6	744.8	965.5	1,026.6
2006.....	54.3	101.3	159.3	255.2	386.1	560.2	773.4	950.9	1,077.5
2005.....	56.3	106.1	160.2	265.3	400.4	585.9	795.9	1,017.1	1,128.1
2004.....	54.9	102.5	166.7	270.3	397.8	567.8	784.8	1,009.2	1,131.2
2003.....	60.7	107.1	177.1	267.0	408.6	582.3	819.1	1,005.4	1,234.9
2002.....	57.4	105.5	173.2	271.9	412.8	600.5	831.8	1,000.4	1,308.8
2001.....	61.7	110.3	172.3	291.3	413.5	616.9	846.0	1,033.1	1,363.7
2000.....	64.3	105.4	180.7	272.5	422.5	633.8	851.8	1,138.5	1,383.5
1999.....	65.0	110.9	179.3	282.8	448.9	611.2	834.8	1,094.3	1,399.6

Hispanic, male

2007.....	52.2	103.3	173.1	292.4	487.4	659.3	959.8	1,249.0	1,417.9
2006.....	49.1	101.3	183.0	287.2	476.3	708.0	980.6	1,285.7	1,440.1
2005.....	52.6	107.2	182.5	317.8	495.0	749.6	1,028.9	1,377.3	1,601.5
2004.....	52.3	111.2	181.5	322.8	481.2	726.1	1,052.8	1,373.1	1,508.8
2003.....	58.9	111.1	194.1	321.9	509.5	759.7	1,064.0	1,334.8	1,668.6
2002.....	54.2	111.0	204.8	323.5	518.7	756.5	1,111.5	1,332.6	1,869.0
2001.....	60.1	109.9	189.4	340.6	519.1	802.4	1,171.7	1,423.2	1,967.4
2000.....	60.9	104.2	199.7	323.3	521.3	820.6	1,140.5	1,637.2	2,049.7
1999.....	64.4	111.7	208.3	345.2	567.6	802.5	1,120.3	1,520.0	1,945.3

Hispanic, female

2007.....	61.2	93.3	151.4	208.5	312.4	429.9	588.7	776.3	814.2
2006.....	59.7	101.4	136.9	226.9	310.4	445.1	623.4	728.1	884.9
2005.....	60.2	105.1	139.3	219.0	321.6	458.9	628.5	778.2	880.4
2004.....	57.5	93.9	152.9	224.3	328.9	445.6	594.4	768.9	937.0
2003.....	62.4	103.1	161.4	219.2	325.7	446.1	645.4	791.1	1,014.8
2002.....	60.7	100.3	144.2	227.0	326.2	480.3	632.0	788.4	1,031.2
2001.....	63.3	110.7	156.7	248.7	328.0	475.3	614.9	791.4	1,068.6
2000.....	67.6	106.5	163.5	228.8	342.4	492.3	647.3	835.0	1,056.5
1999.....	65.7	110.1	153.2	228.9	352.8	467.3	633.2	840.3	1,136.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

Non-Hispanic 4/, both sexes

2007.....	85.4	154.8	257.2	410.6	627.3	870.5	1,158.9	1,440.6	1,619.0
2006.....	87.7	157.2	260.6	427.2	632.4	884.8	1,168.3	1,443.4	1,630.6
2005.....	90.0	159.4	263.5	434.2	638.8	906.6	1,180.2	1,448.1	1,660.5
2004.....	90.3	160.5	269.0	439.1	655.1	913.3	1,193.3	1,443.5	1,674.3
2003.....	92.8	163.5	274.6	453.4	664.3	929.0	1,214.2	1,468.1	1,714.7
2002.....	93.8	166.2	284.5	459.2	680.4	952.6	1,224.0	1,474.9	1,736.8
2001.....	95.3	169.1	288.6	464.3	690.0	960.9	1,226.9	1,480.7	1,776.6
2000.....	96.1	170.7	296.5	476.3	706.7	967.3	1,236.6	1,515.9	1,831.1
1999.....	96.1	171.2	303.1	484.0	714.7	980.0	1,237.8	1,503.6	1,816.1

Non-Hispanic 4/, male

2007.....	82.3	166.1	289.2	472.4	738.3	1,054.0	1,463.2	1,916.5	2,296.0
2006.....	84.6	168.2	290.6	490.8	750.2	1,068.5	1,471.1	1,935.4	2,289.8
2005.....	87.1	170.4	293.2	496.9	761.6	1,114.2	1,501.6	1,928.5	2,354.9
2004.....	89.9	173.4	298.3	504.4	775.9	1,114.3	1,527.7	1,918.9	2,387.3
2003.....	94.0	174.7	303.4	520.4	791.7	1,138.5	1,556.5	1,965.5	2,444.0
2002.....	93.3	176.8	316.2	532.8	814.2	1,180.0	1,573.7	1,974.9	2,513.2
2001.....	94.7	179.2	320.0	538.6	832.2	1,196.9	1,584.2	1,998.0	2,634.8
2000.....	96.4	179.1	329.4	549.5	852.4	1,207.8	1,602.6	2,061.5	2,732.4
1999.....	97.2	179.5	338.6	563.0	863.4	1,231.0	1,616.8	2,060.5	2,747.5

Non-Hispanic 4/, female

2007.....	88.4	143.9	227.1	353.8	529.8	719.9	934.6	1,141.1	1,299.3
2006.....	90.7	146.7	232.4	369.0	529.1	734.9	946.6	1,136.9	1,324.2
2005.....	92.8	148.9	235.5	376.8	531.2	738.0	946.7	1,150.8	1,343.6
2004.....	90.6	148.1	241.4	379.5	549.6	751.0	952.9	1,151.7	1,355.6
2003.....	91.7	152.7	247.5	392.3	553.5	760.5	970.5	1,164.9	1,394.3
2002.....	94.3	156.0	254.6	392.3	564.4	770.5	977.1	1,171.5	1,401.8
2001.....	95.8	159.5	259.0	396.6	567.0	773.0	976.9	1,171.2	1,419.6
2000.....	95.8	162.7	265.6	409.7	581.3	777.5	981.8	1,194.4	1,466.6
1999.....	95.0	163.1	269.9	412.2	587.0	783.4	974.5	1,178.0	1,443.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

Non-Hispanic White, both sexes

2007 5/.....	81.4	147.9	248.1	403.5	624.8	874.5	1,165.6	1,451.4	1,631.9
2006 5/.....	83.6	150.5	253.4	420.7	630.5	889.2	1,176.2	1,452.4	1,643.1
2005 5/.....	85.1	151.8	254.6	426.1	635.6	910.3	1,187.2	1,453.4	1,668.6
2004 5/.....	85.4	152.5	260.8	433.3	651.5	913.4	1,196.0	1,447.9	1,680.9
2003 5/.....	87.3	155.1	267.4	447.1	661.9	928.9	1,213.9	1,468.8	1,717.8
2002.....	89.0	158.8	276.0	451.8	676.5	950.9	1,222.3	1,473.3	1,737.9
2001.....	90.2	162.2	279.7	458.2	682.7	955.9	1,222.7	1,473.0	1,775.2
2000.....	90.3	163.6	288.2	470.0	701.3	959.9	1,227.7	1,512.6	1,826.6
1999.....	90.4	163.5	293.3	474.6	707.7	971.9	1,226.4	1,498.7	1,808.3

Non-Hispanic White, male

2007 5/.....	78.1	157.7	275.1	459.3	727.4	1,047.1	1,459.5	1,920.4	2,300.4
2006 5/.....	80.9	159.5	278.9	478.0	739.5	1,061.5	1,468.4	1,937.1	2,300.2
2005 5/.....	82.6	161.5	279.2	481.7	749.5	1,105.9	1,498.8	1,921.9	2,358.3
2004 5/.....	84.5	163.2	285.6	491.2	764.3	1,100.3	1,514.8	1,911.4	2,387.1
2003 5/.....	87.9	163.8	292.1	506.7	780.1	1,128.6	1,541.2	1,955.2	2,441.7
2002.....	88.6	167.2	303.2	515.9	803.1	1,167.1	1,556.2	1,957.1	2,507.7
2001.....	89.1	169.4	305.9	525.6	815.0	1,178.6	1,562.9	1,968.7	2,616.8
2000.....	90.1	168.5	314.5	535.3	837.1	1,182.9	1,574.8	2,036.1	2,714.1
1999.....	89.9	168.3	323.3	544.9	846.1	1,209.4	1,586.9	2,037.9	2,721.3

Non-Hispanic White, female

2007 5/.....	84.6	138.3	222.0	351.1	532.7	729.8	944.2	1,152.4	1,315.2
2006 5/.....	86.2	141.7	228.7	367.2	532.8	745.4	957.8	1,147.3	1,336.7
2005 5/.....	87.6	142.4	231.0	374.2	533.8	747.9	956.3	1,160.9	1,353.2
2004 5/.....	86.3	142.0	236.9	379.4	550.9	759.0	962.4	1,161.0	1,364.5
2003 5/.....	86.7	146.5	243.7	391.6	556.9	764.8	977.1	1,169.9	1,399.1
2002.....	89.5	150.7	249.8	392.4	564.4	774.0	983.3	1,177.4	1,405.4
2001.....	91.3	155.1	254.6	395.8	565.8	775.0	981.8	1,173.8	1,425.1
2000.....	90.5	158.8	263.2	409.5	582.0	780.3	983.7	1,201.6	1,468.7
1999.....	91.0	158.8	264.6	409.6	586.3	782.7	973.5	1,180.6	1,444.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

Non-Hispanic Black, both sexes

2007 5/.....	122.1	227.7	370.9	559.2	794.6	1,008.9	1,310.5	1,549.0	1,696.8
2006 5/.....	126.9	229.8	367.1	573.9	786.1	1,021.8	1,306.2	1,581.1	1,700.4
2005 5/.....	132.3	236.9	383.0	585.2	797.8	1,043.3	1,317.3	1,607.0	1,764.4
2004 5/.....	132.9	240.8	385.1	579.8	817.5	1,076.5	1,373.5	1,600.7	1,794.3
2003 5/.....	141.5	248.2	390.9	600.8	816.7	1,097.7	1,405.0	1,662.0	1,867.5
2002.....	138.6	243.5	407.7	606.9	845.5	1,143.2	1,443.0	1,680.9	1,904.3
2001.....	141.4	246.5	412.8	598.1	871.7	1,172.3	1,446.2	1,735.8	1,947.5
2000.....	148.6	257.2	415.1	616.2	874.9	1,193.3	1,497.2	1,718.5	2,015.1
1999.....	147.7	260.2	437.3	644.3	891.9	1,206.6	1,516.9	1,715.6	2,024.0

Non-Hispanic Black, male

2007 5/.....	120.4	257.3	455.8	709.1	1,028.1	1,338.9	1,806.1	2,230.2	2,691.9
2006 5/.....	123.6	263.4	450.9	720.7	1,027.7	1,377.1	1,816.6	2,319.6	2,577.4
2005 5/.....	130.0	264.4	470.6	745.1	1,044.7	1,419.2	1,831.8	2,385.2	2,720.4
2004 5/.....	137.6	280.7	466.2	742.0	1,054.9	1,467.0	1,952.1	2,376.3	2,781.1
2003 5/.....	149.1	288.5	475.6	763.8	1,065.1	1,477.8	2,004.0	2,453.4	2,859.6
2002.....	140.8	277.8	494.5	790.8	1,094.5	1,559.5	2,071.8	2,539.6	3,008.4
2001.....	147.2	286.7	504.5	756.4	1,154.2	1,619.6	2,100.9	2,672.1	3,238.5
2000.....	155.3	299.9	521.3	783.3	1,154.5	1,675.4	2,183.1	2,709.5	3,290.1
1999.....	164.2	305.9	543.1	830.7	1,174.9	1,663.5	2,224.9	2,656.8	3,361.1

Non-Hispanic Black, female

2007 5/.....	123.6	202.7	301.4	442.0	623.2	781.6	1,012.5	1,191.3	1,284.3
2006 5/.....	129.8	201.4	298.1	459.7	607.9	778.1	1,001.4	1,191.6	1,342.7
2005 5/.....	134.3	213.6	310.7	460.7	615.0	786.4	1,011.1	1,196.5	1,379.5
2004 5/.....	128.9	207.2	318.5	453.6	641.5	811.8	1,031.0	1,192.7	1,404.3
2003 5/.....	134.9	214.2	321.3	473.8	632.4	841.5	1,049.3	1,247.9	1,483.2
2002.....	136.7	214.5	336.6	463.2	660.9	863.9	1,067.5	1,234.1	1,482.4
2001.....	136.3	212.4	337.8	473.7	663.1	874.2	1,055.5	1,258.9	1,469.3
2000.....	142.8	220.9	328.1	484.4	668.7	873.2	1,083.7	1,219.7	1,541.6
1999.....	133.4	221.4	351.1	498.2	682.1	903.4	1,090.3	1,249.0	1,524.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/, both sexes

2007.....	1.7	3.5	5.4	7.1	9.3	10.7	12.7	14.2	17.1
2006.....	1.8	3.5	5.2	6.6	8.7	10.6	13.0	14.1	16.3
2005.....	1.9	3.5	5.6	7.5	9.2	10.6	11.8	14.2	17.0
2004.....	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
2003.....	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3

All origins 3/, male

2007.....	2.6	5.5	8.6	11.9	14.9	16.2	19.7	20.6	24.0
2006.....	2.9	5.6	8.4	10.6	13.9	16.0	19.7	21.4	21.9
2005.....	2.8	5.7	9.1	11.9	14.8	16.0	17.9	20.4	22.0
2004.....	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
2003.....	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1

All origins 3/, female

2007.....	0.9	1.7	2.4	2.8	4.3	6.1	7.6	10.1	13.8
2006.....	0.8	1.4	2.1	2.9	4.2	6.3	8.1	9.6	13.7
2005.....	1.0	1.4	2.3	3.4	4.3	6.2	7.3	10.4	14.7
2004.....	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
2003.....	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 4/, both sexes

2007.....	1.8	3.8	5.6	7.5	9.5	10.9	13.0	14.4	17.4
2006.....	2.0	3.6	5.4	6.9	8.8	10.8	13.3	14.3	16.5
2005.....	2.0	3.7	5.8	7.7	9.5	10.9	12.0	14.1	17.6
2004.....	2.1	4.0	5.3	7.7	9.6	11.3	13.0	15.1	18.2
2003.....	2.1	3.8	5.7	8.2	10.0	11.0	13.0	13.7	18.7
2002.....	2.1	3.7	6.0	8.4	9.1	12.4	13.3	14.3	18.7
2001.....	2.1	3.6	6.3	8.1	9.5	11.8	13.6	15.4	18.4
2000.....	2.1	3.9	5.8	8.3	10.0	11.7	12.8	14.1	19.5
1999.....	2.4	3.8	5.9	8.2	9.5	11.5	13.6	15.4	19.4

Non-Hispanic 4/, male

2007.....	2.7	5.9	9.0	12.3	15.1	16.6	20.2	21.1	24.4
2006.....	3.1	5.9	8.7	11.1	14.0	16.2	19.9	21.6	22.3
2005.....	3.0	6.0	9.4	12.2	15.2	16.3	18.4	20.1	22.8
2004.....	3.3	6.4	8.9	12.1	15.2	16.6	19.8	21.9	24.9
2003.....	3.4	6.5	9.3	13.0	15.2	16.8	19.1	20.2	27.1
2002.....	3.4	6.0	9.7	13.4	14.4	18.3	19.3	20.6	26.7
2001.....	3.3	6.0	10.4	13.3	14.5	18.2	19.5	22.4	25.8
2000.....	3.3	6.2	9.6	13.0	15.3	17.1	19.0	20.1	26.5
1999.....	3.7	6.0	9.1	13.4	14.3	17.5	20.0	22.7	27.3

Non-Hispanic 4/, female

2007.....	1.0	1.8	2.4	3.0	4.5	6.3	7.8	10.2	14.2
2006.....	0.9	1.5	2.2	3.1	4.3	6.5	8.5	9.7	13.8
2005.....	1.0	1.5	2.3	3.6	4.6	6.5	7.4	10.5	15.2
2004.....	1.0	1.6	1.8	3.6	4.8	6.9	8.2	11.0	15.1
2003.....	0.9	1.2	2.2	3.8	5.4	6.4	8.6	9.8	15.0
2002.....	1.0	1.5	2.6	3.8	4.6	7.7	9.1	10.5	15.3
2001.....	1.0	1.4	2.5	3.4	5.3	6.8	9.4	11.3	15.3
2000.....	1.0	1.7	2.2	4.0	5.5	7.5	8.4	10.5	16.6
1999.....	1.1	1.7	2.9	3.6	5.3	6.8	9.1	11.1	16.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic White, both sexes

2007 5/.....	1.6	3.5	5.2	7.2	9.1	10.9	13.0	14.7	17.9
2006 5/.....	1.8	3.4	5.0	6.5	8.4	10.6	13.3	14.5	17.1
2005 5/.....	1.8	3.3	5.3	7.3	9.0	10.8	12.0	14.1	17.9
2004 5/.....	1.7	3.5	4.8	7.1	9.2	11.2	13.0	15.2	18.7
2003 5/.....	1.9	3.3	5.4	7.5	9.5	10.7	12.7	13.5	19.1
2002.....	1.8	3.2	5.5	7.8	8.6	12.7	13.5	14.4	18.9
2001.....	1.8	3.3	5.7	7.5	8.9	11.6	13.6	15.7	18.9
2000.....	1.8	3.3	5.1	7.4	9.5	11.3	12.7	14.1	19.8
1999.....	1.8	3.1	5.1	7.6	9.0	11.1	13.3	15.5	19.9

Non-Hispanic White, male

2007 5/.....	2.4	5.4	8.3	12.0	14.3	16.0	19.9	21.3	24.8
2006 5/.....	2.8	5.5	8.0	10.3	13.0	15.5	19.6	21.5	23.0
2005 5/.....	2.6	5.3	8.3	11.4	14.1	15.6	17.7	19.7	22.9
2004 5/.....	2.6	5.6	8.1	11.1	14.2	15.9	19.3	21.8	24.8
2003 5/.....	2.8	5.8	8.7	11.5	14.3	15.7	18.2	19.4	27.4
2002.....	2.9	5.1	8.8	12.2	13.3	18.4	19.5	20.5	26.8
2001.....	2.7	5.4	9.2	11.9	13.4	17.4	19.2	22.5	26.4
2000.....	2.8	5.0	8.3	11.4	14.1	16.1	18.3	19.7	27.4
1999.....	2.8	4.9	7.7	12.3	13.5	16.9	19.1	22.3	27.5

Non-Hispanic White, female

2007 5/.....	0.8	1.6	2.2	2.8	4.4	6.7	7.9	10.4	14.7
2006 5/.....	0.8	1.3	2.1	3.0	4.3	6.5	8.7	10.1	14.4
2005 5/.....	0.9	1.3	2.3	3.4	4.5	6.7	7.8	10.7	15.6
2004 5/.....	0.8	1.5	1.6	3.5	4.8	7.4	8.4	11.1	16.0
2003 5/.....	0.9	0.9	2.2	3.9	5.3	6.6	8.7	9.9	15.4
2002.....	0.8	1.3	2.3	3.7	4.5	8.0	9.1	10.7	15.5
2001.....	0.8	1.2	2.3	3.3	5.0	6.9	9.7	11.6	15.7
2000.....	0.9	1.6	2.0	3.7	5.5	7.4	8.7	10.7	16.8
1999.....	0.9	1.3	2.6	3.4	5.1	6.6	9.2	11.4	16.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic Black, both sexes

2007 5/.....	3.1	5.6	9.4	10.8	14.5	13.1	14.9	12.2	14.6
2006 5/.....	3.3	5.7	8.7	11.3	13.5	14.4	14.5	12.9	10.1
2005 5/.....	3.7	6.7	9.5	12.9	14.3	14.0	13.4	14.2	12.8
2004 5/.....	4.7	7.9	9.8	13.6	14.1	14.0	15.1	15.7	12.7
2003 5/.....	4.1	6.6	9.1	14.8	15.8	15.5	15.1	16.3	15.5
2002.....	4.3	7.3	10.4	15.0	13.5	12.2	11.6	12.8	16.0
2001.....	4.4	6.4	12.5	13.4	15.5	16.0	14.2	13.5	13.1
2000.....	4.1	8.5	12.3	15.6	15.0	16.5	14.5	16.1	14.9
1999.....	6.0	9.0	12.7	14.7	14.5	16.6	17.4	15.8	14.5

Non-Hispanic Black, male

2007 5/.....	4.5	9.5	16.0	17.9	26.5	25.5	26.9	20.1	24.4
2006 5/.....	5.2	9.4	15.5	20.3	25.6	25.3	25.8	26.0	*
2005 5/.....	6.3	11.9	18.1	22.9	26.9	26.3	27.6	28.1	21.9
2004 5/.....	7.9	14.1	17.4	24.3	26.7	27.6	26.1	26.4	27.4
2003 5/.....	7.3	11.6	16.8	28.6	28.1	30.3	27.1	30.9	25.6
2002.....	7.0	13.2	18.7	27.5	24.6	20.9	16.9	21.8	26.3
2001.....	7.1	11.2	22.7	25.5	25.5	29.6	24.6	21.3	*
2000.....	6.3	16.3	22.0	28.1	27.7	29.7	28.9	29.2	*
1999.....	10.4	14.8	21.8	25.7	24.2	27.0	33.4	29.2	25.0

Non-Hispanic Black, female

2007 5/.....	1.8	2.3	4.0	5.3	5.7	4.5	7.7	8.0	10.5
2006 5/.....	1.6	2.6	3.1	4.3	4.6	6.9	7.7	*	8.6
2005 5/.....	1.5	2.3	2.4	5.1	4.9	5.6	*	*	9.2
2004 5/.....	1.9	2.7	3.5	5.4	4.8	4.7	8.6	10.0	*
2003 5/.....	*	2.4	2.7	4.0	6.8	5.5	8.0	8.7	11.6
2002.....	2.0	2.3	3.5	5.3	5.3	6.4	8.4	8.1	12.1
2001.....	2.0	2.3	4.1	3.9	8.1	6.9	7.9	9.6	10.7
2000.....	2.2	*	4.3	5.7	5.7	7.7	5.8	9.5	14.4
1999.....	2.2	4.0	5.3	6.1	7.3	9.6	7.9	9.2	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

All origins 3/, both sexes

2007.....	2.4	4.3	8.3	12.9	16.9	22.2	24.4	27.9	25.0
2006.....	2.3	4.8	8.4	13.2	17.4	22.0	26.9	27.0	25.6
2005.....	2.3	4.9	8.4	12.6	17.5	22.4	27.6	27.3	24.9
2004.....	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
2003.....	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8

All origins 3/, male

2007.....	4.1	7.6	14.5	22.8	30.2	38.6	44.2	52.8	48.4
2006.....	4.1	8.1	15.0	23.2	30.8	38.7	47.0	51.3	48.5
2005.....	3.9	8.8	14.8	22.2	31.1	39.6	49.6	52.4	46.0
2004.....	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
2003.....	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0

All origins 3/, female

2007.....	0.7	1.2	2.4	3.7	5.2	8.9	9.9	12.1	13.9
2006.....	0.7	1.7	2.2	4.1	5.8	8.4	12.2	11.8	14.9
2005.....	0.7	1.2	2.5	3.9	5.6	8.5	11.6	11.6	15.3
2004.....	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
2003.....	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

Non-Hispanic 4/, both sexes

2007.....	2.5	4.6	8.7	13.5	17.7	23.1	25.2	28.5	25.7
2006.....	2.5	5.1	8.7	13.7	18.2	22.6	27.6	27.6	26.1
2005.....	2.5	5.2	8.9	13.1	18.0	23.2	28.4	27.9	25.2
2004.....	2.5	5.0	8.5	13.6	18.1	22.0	26.7	28.2	26.2
2003.....	2.6	4.9	9.1	13.4	18.2	22.8	26.1	28.1	25.4
2002.....	2.5	5.0	9.0	13.3	18.8	23.1	26.3	28.4	24.5
2001.....	2.5	5.1	8.2	13.3	19.3	23.5	26.7	26.5	26.0
2000.....	2.5	5.2	9.1	13.7	18.5	22.5	24.7	27.4	26.1
1999.....	2.5	4.9	9.0	13.4	18.7	22.7	24.5	24.8	26.8

Non-Hispanic 4/, male

2007.....	4.3	8.0	15.2	23.8	31.6	40.0	45.6	54.1	49.7
2006.....	4.4	8.5	15.6	24.1	32.0	39.8	48.3	52.6	49.8
2005.....	4.2	9.3	15.5	23.0	31.9	40.8	50.9	53.9	46.3
2004.....	4.3	8.6	14.8	23.6	31.5	38.1	48.0	52.7	49.4
2003.....	4.4	8.6	15.9	23.7	31.7	39.5	46.8	51.9	46.3
2002.....	4.3	8.7	15.8	22.9	32.4	41.4	47.8	51.6	43.5
2001.....	4.3	9.0	14.3	23.2	33.8	40.8	47.6	48.9	48.4
2000.....	4.5	9.1	15.9	23.8	31.5	38.6	44.2	49.3	47.8
1999.....	4.4	8.7	15.8	22.4	32.6	40.4	44.4	45.3	48.0

Non-Hispanic 4/, female

2007.....	0.8	1.2	2.6	4.0	5.4	9.3	10.2	12.4	14.3
2006.....	0.7	1.8	2.3	4.2	6.1	8.6	12.5	12.0	15.1
2005.....	0.8	1.3	2.5	4.1	5.8	8.9	11.9	11.8	15.5
2004.....	0.7	1.5	2.5	4.5	6.4	8.9	11.4	13.2	15.8
2003.....	0.8	1.3	2.7	3.9	6.5	9.4	11.4	13.7	16.1
2002.....	0.8	1.4	2.6	4.5	6.9	8.4	11.1	14.2	16.3
2001.....	0.8	1.5	2.4	4.3	6.8	9.7	12.1	13.1	16.7
2000.....	0.7	1.5	2.8	4.5	7.3	9.8	11.1	14.5	17.3
1999.....	0.6	1.3	2.7	5.3	6.8	8.9	10.7	12.8	18.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

Non-Hispanic White, both sexes

2007 5/.....	2.6	4.5	8.5	13.7	18.1	23.8	26.0	29.0	26.5
2006 5/.....	2.5	4.9	8.7	13.9	18.4	23.2	28.2	28.5	27.0
2005 5/.....	2.4	5.2	8.8	13.0	18.1	23.7	28.8	28.7	26.2
2004 5/.....	2.4	4.9	8.3	13.3	18.1	21.9	27.2	28.6	26.5
2003 5/.....	2.4	4.6	8.8	12.9	17.7	23.2	26.5	28.3	26.1
2002.....	2.4	4.9	8.5	12.8	18.2	23.0	26.3	28.7	25.1
2001.....	2.4	4.8	7.4	12.6	18.5	22.7	26.8	26.8	26.5
2000.....	2.4	4.8	8.4	12.9	17.9	22.2	24.2	27.4	26.7
1999.....	2.3	4.5	8.3	12.5	17.9	22.2	23.9	25.0	27.2

Non-Hispanic White, male

2007 5/.....	4.5	8.0	15.0	24.2	32.4	41.0	46.6	55.2	51.3
2006 5/.....	4.5	8.4	15.7	24.4	32.4	40.6	49.3	55.0	51.3
2005 5/.....	4.3	9.5	15.5	22.8	32.0	41.8	52.0	55.3	49.0
2004 5/.....	4.2	8.6	14.6	23.3	31.6	38.3	48.9	54.0	50.1
2003 5/.....	4.2	8.1	15.5	23.1	30.6	40.2	47.7	53.2	47.9
2002.....	4.3	8.6	15.0	22.3	31.8	41.3	48.3	52.3	44.7
2001.....	4.1	8.3	13.1	22.1	32.5	39.8	48.1	49.5	49.0
2000.....	4.4	8.4	14.7	22.8	30.8	38.2	43.4	50.1	49.4
1999.....	4.1	8.0	14.6	21.0	31.7	39.5	43.4	46.1	48.0

Non-Hispanic White, female

2007 5/.....	0.7	1.1	2.3	3.8	5.2	9.3	10.4	12.4	14.7
2006 5/.....	0.6	1.6	1.8	4.0	5.9	8.6	12.5	11.8	15.7
2005 5/.....	0.6	1.0	2.3	3.8	5.7	8.7	11.7	12.0	15.8
2004 5/.....	0.5	1.3	2.2	4.0	6.1	8.4	11.3	12.8	16.0
2003 5/.....	0.7	1.1	2.3	3.4	6.2	9.2	11.2	13.0	16.5
2002.....	0.6	1.2	2.3	4.0	6.1	8.0	10.5	14.3	16.6
2001.....	0.6	1.3	1.9	3.7	6.2	8.8	11.8	13.2	17.1
2000.....	0.4	1.2	2.4	3.7	6.6	9.3	10.7	13.8	17.5
1999.....	0.5	1.1	2.3	4.6	5.9	8.3	10.3	12.6	18.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

Non-Hispanic Black, both sexes

2007 5/.....	2.7	6.2	12.4	17.0	20.2	23.7	24.5	27.5	19.2
2006 5/.....	3.1	7.6	11.9	16.9	21.9	23.2	25.5	24.0	18.3
2005 5/.....	3.3	6.9	12.2	17.9	21.9	24.2	31.0	22.9	17.5
2004 5/.....	4.1	6.7	12.3	20.6	23.5	27.4	27.7	26.8	21.9
2003 5/.....	4.2	8.5	14.3	21.2	26.8	25.2	27.3	30.1	21.4
2002.....	3.7	7.0	15.3	20.8	28.1	29.8	31.9	26.7	20.6
2001.....	3.9	8.8	16.3	22.2	30.1	36.1	31.2	28.1	24.0
2000.....	4.2	10.0	17.4	23.7	27.5	29.9	36.4	30.5	22.6
1999.....	4.5	9.4	17.4	23.9	29.1	31.6	36.0	24.1	23.6

Non-Hispanic Black, male

2007 5/.....	3.9	10.6	21.0	29.8	36.9	42.0	46.0	49.9	41.0
2006 5/.....	5.0	12.5	19.3	29.2	38.9	42.7	45.1	40.0	35.7
2005 5/.....	5.1	11.6	20.5	32.2	40.7	41.6	54.2	48.9	26.6
2004 5/.....	6.5	10.9	20.3	34.1	40.7	44.2	49.9	44.1	39.6
2003 5/.....	7.3	14.2	23.7	37.1	47.8	41.8	48.1	44.5	38.4
2002.....	5.8	11.7	27.5	35.0	46.3	54.4	54.0	49.1	38.3
2001.....	6.2	15.7	28.3	39.1	52.6	60.9	55.9	54.8	47.4
2000.....	6.5	17.8	31.3	39.1	45.6	50.2	66.0	46.3	38.7
1999.....	7.7	16.5	30.8	40.9	46.8	56.8	68.2	37.2	49.9

Non-Hispanic Black, female

2007 5/.....	1.6	2.6	5.4	6.9	8.0	11.2	11.5	15.7	10.1
2006 5/.....	1.4	3.5	5.8	7.4	9.4	9.8	13.7	15.6	11.2
2005 5/.....	1.7	3.0	5.3	6.7	8.0	12.2	17.2	9.1	13.8
2004 5/.....	2.0	3.1	5.7	10.0	10.7	16.0	14.6	17.8	14.9
2003 5/.....	1.6	3.6	6.5	8.8	11.3	14.1	15.0	22.5	14.9
2002.....	1.9	3.1	5.3	9.7	14.6	13.3	18.8	15.0	13.8
2001.....	1.8	2.9	6.5	8.9	13.5	19.5	16.4	14.6	15.4
2000.....	2.2	3.4	6.0	11.6	14.2	16.3	18.6	22.5	16.6
1999.....	1.7	3.4	6.4	10.6	15.9	14.9	16.6	17.6	13.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

All origins 3/, both sexes

2007.....	2.2	3.1	4.6	7.2	11.0	15.9	21.9	29.0	35.4
2006.....	2.1	3.3	5.0	7.2	11.0	15.5	22.3	29.9	35.8
2005.....	2.1	3.1	4.8	7.3	11.8	16.7	23.5	30.6	38.5
2004.....	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
2003.....	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1

All origins 3/, male

2007.....	2.5	4.1	6.3	10.1	16.1	23.1	30.8	41.4	51.1
2006.....	2.6	4.4	7.0	10.1	15.4	21.4	31.3	44.7	50.2
2005.....	2.5	4.3	6.6	10.4	16.4	23.8	32.5	44.3	54.2
2004.....	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
2003.....	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3

All origins 3/, female

2007.....	1.8	2.2	3.0	4.5	6.6	10.1	15.3	21.1	27.9
2006.....	1.7	2.3	3.1	4.5	7.2	10.6	15.8	20.7	29.1
2005.....	1.8	2.0	3.0	4.5	7.7	11.0	17.0	22.1	31.2
2004.....	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
2003.....	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

Hispanic, both sexes

2007	3.2	4.5	5.9	10.1	18.9	20.6	31.1	48.1	59.7
2006	3.1	4.9	9.1	11.7	14.8	27.5	35.5	39.9	47.0
2005	3.9	5.5	8.0	11.2	16.0	26.9	39.3	51.3	50.9
2004	3.6	6.0	10.1	13.9	19.1	31.2	38.8	52.2	62.8
2003	4.1	6.4	9.3	14.4	19.8	28.9	42.9	44.7	62.7
2002	3.4	5.2	8.5	14.5	16.9	28.3	43.2	53.9	72.1
2001	4.1	7.0	8.8	14.6	21.4	32.7	40.2	55.1	63.8
2000	4.1	5.7	7.1	13.2	23.2	35.0	41.0	66.8	65.7
1999	3.0	5.5	9.1	15.0	23.2	29.9	44.2	59.9	73.7

Hispanic, male

2007	3.1	5.1	7.1	16.2	28.9	27.5	45.1	63.5	84.4
2006	3.6	5.6	11.1	16.0	20.3	36.2	43.8	58.1	55.3
2005	3.6	7.0	10.0	15.1	20.4	37.3	60.0	75.3	76.0
2004	3.8	6.0	13.2	17.2	26.9	38.9	58.7	80.2	84.1
2003	4.6	7.0	11.5	18.0	25.0	44.0	62.0	66.5	73.9
2002	4.0	5.5	10.8	17.7	24.0	44.3	55.7	72.1	107.1
2001	4.5	8.5	10.2	21.5	30.5	45.7	51.8	76.6	95.2
2000	3.9	6.5	9.4	19.3	34.7	54.5	53.9	104.5	98.8
1999	4.2	6.6	12.7	21.1	33.2	38.7	67.9	84.9	75.5

Hispanic, female

2007	3.3	3.9	4.9	4.6	10.5	15.2	20.9	37.8	46.3
2006	2.6	4.2	7.1	8.0	10.2	20.8	29.4	27.8	42.5
2005	4.2	4.1	6.1	7.7	12.3	18.8	24.4	35.4	37.8
2004	3.4	6.0	7.1	11.0	12.8	25.2	24.5	33.6	51.9
2003	3.5	5.7	7.3	11.4	15.5	17.4	29.3	30.5	57.0
2002	2.8	5.0	6.4	11.6	11.1	15.9	34.3	42.3	54.7
2001	3.7	5.7	7.6	8.7	14.1	22.7	32.0	41.8	48.4
2000	4.3	5.0	5.0	7.9	13.9	20.3	31.9	43.9	49.5
1999	*	4.6	5.8	9.8	15.0	23.4	27.5	45.0	72.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

Non-Hispanic 4/, both sexes

2007.....	2.0	3.0	4.5	6.9	10.3	15.6	21.2	27.8	34.1
2006.....	2.0	3.2	4.6	6.8	10.7	14.5	21.4	29.3	35.2
2005.....	1.9	2.8	4.5	7.0	11.4	15.9	22.5	29.5	37.8
2004.....	1.9	3.3	4.9	7.5	11.5	17.3	22.4	29.1	41.2
2003.....	2.1	3.5	4.9	8.1	11.8	17.7	23.1	31.5	42.7
2002.....	1.9	3.4	5.4	8.0	11.9	18.2	24.6	32.0	46.3
2001.....	2.0	3.4	5.0	8.7	13.2	18.6	26.2	33.9	44.7
2000.....	2.2	3.4	5.4	9.4	13.1	18.7	26.4	35.6	52.0
1999.....	2.2	3.7	5.9	9.4	13.6	19.5	27.4	35.9	49.2

Non-Hispanic 4/, male

2007.....	2.4	4.0	6.3	9.5	15.0	22.7	29.8	40.1	49.2
2006.....	2.4	4.3	6.6	9.6	15.0	20.3	30.4	43.9	49.9
2005.....	2.4	3.9	6.3	10.1	16.1	22.8	30.8	42.5	53.0
2004.....	2.4	4.5	7.0	10.7	16.2	24.0	32.0	40.5	61.7
2003.....	2.6	4.7	7.2	11.8	17.3	25.4	32.6	43.4	59.2
2002.....	2.6	4.7	7.4	11.5	17.6	26.4	35.3	45.8	65.0
2001.....	2.6	4.6	7.2	13.1	18.7	26.3	38.3	48.2	61.0
2000.....	2.8	4.6	7.6	13.6	18.4	26.8	38.6	49.3	74.5
1999.....	3.0	5.2	8.5	13.1	19.4	27.8	40.3	53.7	71.0

Non-Hispanic 4/, female

2007.....	1.6	2.0	2.8	4.5	6.3	9.7	14.9	20.2	27.0
2006.....	1.5	2.1	2.7	4.2	6.9	9.9	14.8	20.3	28.4
2005.....	1.5	1.8	2.8	4.3	7.4	10.3	16.5	21.4	30.9
2004.....	1.4	2.1	3.0	4.6	7.4	11.9	15.5	22.1	32.1
2003.....	1.5	2.3	2.8	4.8	7.1	11.5	16.3	24.2	35.5
2002.....	1.3	2.1	3.5	4.9	7.0	11.6	17.1	23.6	38.3
2001.....	1.3	2.2	2.9	4.8	8.4	12.5	17.8	25.3	37.9
2000.....	1.5	2.2	3.4	5.6	8.6	12.3	17.8	27.6	42.9
1999.....	1.4	2.2	3.5	6.1	8.6	13.1	18.4	25.5	40.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

Non-Hispanic White, both sexes

2007 5/.....	1.5	2.4	3.5	5.6	8.7	13.0	18.1	23.9	30.3
2006 5/.....	1.5	2.7	3.8	5.5	8.9	12.2	18.3	25.2	31.5
2005 5/.....	1.4	2.4	3.7	5.7	9.3	13.6	19.1	25.6	33.8
2004 5/.....	1.4	2.6	3.9	6.2	9.3	14.7	19.6	25.8	37.6
2003 5/.....	1.5	2.8	4.1	6.9	10.2	15.3	20.0	27.8	37.5
2002.....	1.6	2.6	4.5	6.7	9.9	15.5	21.3	28.0	40.7
2001.....	1.6	2.6	4.3	7.3	10.8	16.3	23.0	29.8	40.0
2000.....	1.6	2.8	4.6	8.1	11.0	16.0	22.6	31.2	46.3
1999.....	1.8	2.9	4.8	7.7	12.0	16.8	23.8	31.8	43.5

Non-Hispanic White, male

2007 5/.....	1.8	3.2	5.1	7.6	12.8	19.3	25.4	33.5	44.0
2006 5/.....	2.0	3.7	5.6	7.8	12.7	17.2	26.5	37.9	45.2
2005 5/.....	1.7	3.3	5.3	8.0	13.0	19.4	26.4	37.0	47.2
2004 5/.....	1.8	3.7	5.7	8.8	13.3	20.0	28.0	35.5	56.9
2003 5/.....	2.0	3.7	6.1	9.9	14.6	21.9	28.2	38.7	52.3
2002.....	2.1	3.7	6.4	9.6	14.9	22.7	31.2	40.7	58.0
2001.....	2.1	3.6	6.1	10.9	15.1	23.0	34.2	42.4	55.7
2000.....	2.1	4.0	6.5	11.9	15.7	23.2	33.5	43.8	67.3
1999.....	2.5	4.1	7.1	10.7	17.3	24.3	35.2	47.6	62.6

Non-Hispanic White, female

2007 5/.....	1.3	1.6	2.0	3.7	5.1	7.8	12.5	17.8	23.9
2006 5/.....	1.1	1.7	2.1	3.4	5.5	7.9	12.2	17.2	25.2
2005 5/.....	1.1	1.5	2.2	3.6	6.1	8.8	13.7	18.5	27.7
2004 5/.....	1.0	1.5	2.2	3.8	5.8	10.3	13.5	19.8	28.9
2003 5/.....	1.1	1.8	2.2	4.2	6.2	9.9	14.0	21.2	31.0
2002.....	1.0	1.5	2.7	3.9	5.4	9.6	14.2	20.2	33.3
2001.....	1.0	1.7	2.4	3.9	7.1	10.9	15.0	22.2	33.5
2000.....	1.1	1.7	2.8	4.5	6.9	10.2	15.0	23.8	37.9
1999.....	1.1	1.6	2.6	4.9	7.4	10.9	15.7	22.6	35.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

Non-Hispanic Black, both sexes

2007 5/.....	3.8	6.5	9.7	15.6	21.6	30.1	44.2	59.5	65.5
2006 5/.....	3.6	5.5	8.5	15.2	21.3	29.1	42.7	65.2	68.2
2005 5/.....	4.1	5.3	9.2	14.7	24.1	30.6	49.5	57.1	75.4
2004 5/.....	4.0	7.0	10.8	15.8	27.9	34.4	44.2	60.8	71.4
2003 5/.....	4.5	7.5	9.9	17.0	22.6	33.7	49.6	63.0	86.0
2002.....	3.7	7.7	10.7	16.4	25.0	39.7	53.7	63.8	100.1
2001.....	3.8	7.5	9.3	18.0	27.8	35.0	51.7	70.6	89.3
2000.....	4.8	7.2	11.2	17.6	27.0	38.7	57.3	75.6	102.7
1999.....	4.0	8.6	13.3	19.4	24.0	40.7	59.2	74.6	103.1

Non-Hispanic Black, male

2007 5/.....	5.3	9.9	13.5	24.2	34.0	45.5	69.3	95.0	96.0
2006 5/.....	4.8	7.9	12.8	22.6	31.7	38.9	68.6	102.5	85.2
2005 5/.....	5.6	7.7	13.8	24.9	35.8	49.4	68.4	77.7	117.0
2004 5/.....	5.4	9.6	15.7	25.2	41.6	56.9	63.9	94.7	99.6
2003 5/.....	6.4	11.2	15.2	26.6	37.0	55.3	76.6	86.8	107.6
2002.....	5.2	11.3	15.7	24.6	37.1	60.9	74.6	91.3	136.7
2001.....	5.2	11.6	13.8	28.0	44.8	55.1	76.8	107.1	106.3
2000.....	6.7	9.9	16.6	26.4	38.6	57.7	83.3	104.7	138.6
1999.....	5.8	12.4	19.0	28.7	35.6	60.6	90.9	121.3	149.8

Non-Hispanic Black, female

2007 5/.....	2.6	3.6	6.6	8.8	12.5	19.5	29.1	40.8	52.8
2006 5/.....	2.5	3.4	4.9	9.4	13.7	22.4	27.2	45.5	61.3
2005 5/.....	2.7	3.2	5.4	6.7	15.4	17.8	38.3	46.3	58.6
2004 5/.....	2.8	4.7	6.8	8.5	17.8	19.2	32.6	42.9	60.2
2003 5/.....	2.8	4.4	5.5	9.6	12.0	19.1	33.6	50.5	77.6
2002.....	2.4	4.6	6.7	10.0	15.9	25.5	41.2	49.5	86.1
2001.....	2.6	4.1	5.5	10.2	15.2	21.6	36.8	52.1	83.0
2000.....	3.1	4.8	6.8	10.7	18.5	26.1	41.5	60.9	89.4
1999.....	2.4	5.4	8.7	12.1	15.4	27.6	40.2	51.4	85.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes

2007.....	8.0	14.1	22.4	34.3	50.4	72.3	100.7	141.1	200.8
2006.....	7.6	13.5	21.8	34.8	52.0	73.9	104.4	146.8	207.5
2005.....	7.4	13.4	21.4	34.2	52.7	75.8	105.7	149.2	216.9
2004.....	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
2003.....	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7

All origins 3/, male

2007.....	8.7	16.3	26.8	42.9	62.0	90.6	123.5	173.0	223.5
2006.....	8.5	15.1	25.6	43.1	64.9	91.9	127.3	181.3	234.6
2005.....	8.2	15.2	25.9	42.4	67.7	94.4	131.8	180.2	245.9
2004.....	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
2003.....	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0

All origins 3/, female

2007.....	7.3	12.1	18.3	26.5	40.1	57.4	83.9	121.0	190.0
2006.....	6.7	11.9	18.2	27.2	40.8	59.2	87.6	125.3	194.9
2005.....	6.6	11.6	17.1	26.7	39.7	60.8	86.8	130.0	203.6
2004.....	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
2003.....	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

Hispanic, both sexes

2007	5.9	10.8	17.7	26.4	35.1	51.0	74.9	102.4	133.4
2006	5.2	10.5	17.1	25.2	38.4	55.1	82.4	110.9	142.4
2005	5.0	9.8	15.2	23.1	37.6	58.0	84.7	104.4	146.7
2004	4.9	10.1	16.8	27.3	37.7	60.2	75.8	105.8	147.5
2003	6.1	11.0	20.6	27.8	41.9	59.5	83.2	115.6	144.6
2002	5.0	11.3	17.3	30.7	41.9	63.0	90.1	104.7	165.7
2001	4.6	10.6	16.7	30.1	40.4	66.7	89.8	108.3	193.8
2000	6.3	11.2	19.2	28.1	40.4	60.3	85.4	121.4	181.1
1999	6.7	10.3	14.5	29.2	46.8	67.8	88.4	123.9	176.8

Hispanic, male

2007	6.2	13.0	21.2	32.3	49.1	66.7	90.7	117.8	145.3
2006	5.7	10.6	22.4	30.1	51.9	71.7	101.5	131.0	157.9
2005	6.1	10.6	19.1	29.5	47.5	77.5	101.5	127.8	176.8
2004	6.2	11.6	20.1	36.0	48.1	74.5	104.8	129.3	172.3
2003	7.3	13.7	24.6	37.2	54.3	78.6	102.1	151.8	172.4
2002	6.3	13.4	24.8	40.0	56.9	80.3	118.4	125.8	174.0
2001	4.6	11.1	19.8	39.7	53.2	83.7	113.9	122.1	257.1
2000	6.0	13.1	22.4	36.0	54.8	74.9	109.3	175.2	235.8
1999	7.0	11.8	16.6	33.7	58.7	91.1	114.7	172.9	211.4

Hispanic, female

2007	5.5	8.6	14.3	21.1	23.2	38.7	63.4	92.1	126.9
2006	4.8	10.3	12.2	20.9	27.0	42.2	68.6	97.6	134.2
2005	4.0	9.1	11.5	17.4	29.3	42.9	72.6	88.9	131.0
2004	3.6	8.6	13.8	19.7	29.1	49.2	55.2	90.3	134.7
2003	4.9	8.5	16.8	19.7	31.7	44.9	69.8	92.0	130.5
2002	3.7	9.2	10.5	22.6	29.6	49.7	69.9	91.2	161.6
2001	4.6	10.2	13.9	21.9	30.1	53.7	72.6	99.7	162.9
2000	6.5	9.3	16.3	21.3	28.7	49.3	68.5	88.7	154.3
1999	6.4	8.8	12.5	25.4	37.2	50.2	69.9	94.7	160.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 4/, both sexes

2007	8.2	14.5	22.8	34.9	51.5	73.8	102.4	143.3	204.3
2006	7.9	13.7	22.2	35.6	53.1	75.1	105.8	148.7	210.6
2005	7.7	13.7	21.9	35.0	53.8	76.9	106.8	151.5	220.1
2004	7.7	13.8	22.8	36.5	54.3	79.4	113.2	154.4	228.8
2003	8.1	14.5	23.1	39.3	60.4	84.8	118.7	160.7	243.2
2002	8.6	15.3	25.2	41.4	62.9	87.4	120.2	165.3	244.0
2001	8.8	15.6	26.0	39.9	64.4	89.9	122.4	167.9	257.4
2000	8.4	14.9	25.6	43.0	63.3	94.1	128.6	176.5	273.1
1999	8.4	14.7	26.0	44.5	64.8	93.0	130.5	174.2	274.3

Non-Hispanic 4/, male

2007	9.0	16.6	27.2	43.7	62.9	92.2	125.6	176.3	228.0
2006	8.9	15.5	25.8	44.1	65.9	93.0	128.9	184.1	238.4
2005	8.4	15.7	26.4	43.4	69.1	95.4	133.3	182.9	249.2
2004	8.6	16.5	27.6	44.3	67.6	99.3	140.8	185.5	258.5
2003	9.2	16.8	27.8	47.7	76.3	104.8	145.0	198.2	277.3
2002	9.4	17.8	30.5	52.3	79.1	108.1	148.8	200.3	279.2
2001	9.9	17.6	31.5	48.4	80.5	111.7	151.0	204.3	299.6
2000	9.3	17.3	31.1	52.0	78.7	117.9	161.4	214.0	318.6
1999	9.2	16.9	31.5	54.2	79.0	115.4	163.0	216.7	327.5

Non-Hispanic 4/, female

2007	7.5	12.5	18.7	26.8	41.5	58.7	85.3	122.5	193.0
2006	7.0	12.1	18.7	27.8	41.9	60.5	88.8	126.6	197.7
2005	6.9	11.9	17.6	27.4	40.5	62.0	87.6	132.0	206.8
2004	6.8	11.3	18.4	29.4	42.6	63.2	93.3	135.3	215.6
2003	7.1	12.4	18.8	31.7	46.6	68.7	100.0	137.8	228.1
2002	7.8	12.9	20.2	31.5	48.9	70.9	99.9	144.1	228.8
2001	7.6	13.6	20.9	32.2	50.5	72.6	102.4	146.1	239.9
2000	7.5	12.6	20.4	34.8	50.1	75.4	105.7	154.4	254.7
1999	7.5	12.5	20.9	35.7	52.7	75.4	108.0	149.4	253.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic White, both sexes

2007 5/.....	7.6	13.4	21.0	32.7	49.6	71.8	100.4	142.4	204.3
2006 5/.....	7.3	12.7	21.0	33.4	50.9	73.7	103.0	146.2	210.5
2005 5/.....	7.0	12.4	20.3	32.8	51.3	75.5	104.1	149.4	218.8
2004 5/.....	7.0	12.9	21.6	34.7	52.1	77.1	111.4	152.8	228.5
2003 5/.....	7.4	13.1	21.5	37.6	59.0	81.9	116.6	159.8	242.1
2002.....	7.7	14.5	23.7	39.4	61.1	85.8	118.0	162.3	243.2
2001.....	8.0	14.5	24.4	38.4	62.2	87.8	119.2	165.7	256.9
2000.....	7.5	13.9	23.9	41.4	61.5	92.3	126.6	173.7	271.9
1999.....	7.8	13.7	24.5	42.8	62.9	91.0	128.3	171.9	273.4

Non-Hispanic White, male

2007 5/.....	8.3	15.4	24.9	41.2	60.4	89.8	123.4	176.1	227.6
2006 5/.....	8.1	14.3	24.6	41.6	63.4	90.5	125.8	179.6	236.7
2005 5/.....	7.8	14.3	24.5	40.8	66.0	93.7	129.4	182.1	248.8
2004 5/.....	8.0	15.4	26.4	42.2	65.1	96.6	137.5	183.8	259.6
2003 5/.....	8.4	15.2	26.2	45.4	74.7	102.1	142.3	197.5	276.8
2002.....	8.6	16.8	28.9	49.8	77.2	106.6	145.8	195.9	277.6
2001.....	9.1	16.8	29.9	47.0	78.0	109.8	148.2	201.9	298.2
2000.....	8.5	16.3	29.0	49.9	76.3	115.6	159.5	209.2	315.8
1999.....	8.6	16.0	29.6	52.3	77.1	114.2	159.7	214.9	328.2

Non-Hispanic White, female

2007 5/.....	7.0	11.4	17.2	24.6	40.0	56.8	83.0	120.9	193.3
2006 5/.....	6.5	11.2	17.4	25.8	39.7	59.7	86.0	125.2	198.2
2005 5/.....	6.2	10.6	16.2	25.4	38.2	60.3	85.3	129.1	205.1
2004 5/.....	6.1	10.4	17.0	27.6	40.6	61.1	92.3	133.6	214.6
2003 5/.....	6.4	10.9	16.9	30.3	45.0	65.4	97.9	136.7	226.8
2002.....	6.9	12.2	18.8	29.8	46.9	68.7	98.2	141.8	228.3
2001.....	6.9	12.3	19.2	30.4	48.2	69.9	98.6	143.8	239.8
2000.....	6.6	11.7	19.1	33.5	48.4	73.5	103.5	152.6	254.1
1999.....	6.9	11.4	19.6	34.1	50.4	72.5	106.3	146.5	251.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic Black, both sexes

2007 5/.....	12.9	24.6	39.3	59.8	77.1	103.1	140.0	181.5	229.9
2006 5/.....	12.9	22.9	34.7	58.5	80.6	102.9	150.4	200.5	244.7
2005 5/.....	12.9	24.4	37.6	58.5	84.3	102.7	150.1	200.5	264.1
2004 5/.....	12.4	22.4	36.9	58.0	81.4	115.5	150.3	192.9	262.5
2003 5/.....	13.5	26.0	40.1	60.7	82.5	127.2	158.1	200.1	282.9
2002.....	14.9	24.1	40.9	63.1	88.2	116.3	162.7	221.3	278.3
2001.....	14.8	25.2	44.0	59.0	92.8	122.1	174.5	213.3	289.0
2000.....	15.6	25.4	40.7	63.9	88.5	127.9	169.6	228.8	312.9
1999.....	13.6	23.9	42.8	67.1	91.4	126.5	174.1	227.5	308.8

Non-Hispanic Black, male

2007 5/.....	14.8	29.3	48.1	76.0	100.2	133.0	177.8	224.0	273.1
2006 5/.....	15.1	26.9	40.7	72.1	102.2	139.9	192.0	271.8	314.3
2005 5/.....	13.9	27.9	47.2	75.4	108.3	127.1	199.3	236.0	297.8
2004 5/.....	13.3	27.1	43.1	71.0	102.7	149.6	201.9	245.3	277.4
2003 5/.....	15.6	30.8	46.7	78.0	104.8	159.4	199.5	245.3	321.8
2002.....	16.1	28.7	48.6	81.5	110.8	139.2	212.2	288.6	332.6
2001.....	16.8	27.0	52.3	68.3	117.0	148.7	209.5	264.9	352.3
2000.....	17.0	29.9	52.3	78.0	111.1	162.6	211.9	302.1	381.6
1999.....	15.4	27.4	54.9	82.1	111.6	148.9	229.6	275.4	351.8

Non-Hispanic Black, female

2007 5/.....	11.3	20.7	32.1	47.1	60.2	82.6	117.3	159.1	211.9
2006 5/.....	11.0	19.6	29.7	48.0	64.7	77.5	125.6	162.9	216.4
2005 5/.....	12.0	21.4	29.6	45.3	66.6	86.0	120.8	181.8	250.5
2004 5/.....	11.6	18.5	31.9	47.8	65.5	92.4	119.8	165.4	256.6
2003 5/.....	11.7	22.0	34.7	47.2	65.9	105.4	133.4	176.5	267.8
2002.....	13.8	20.1	34.5	48.7	71.3	100.9	133.2	186.3	257.5
2001.....	13.1	23.7	37.2	51.6	75.0	104.4	153.6	187.0	265.6
2000.....	14.4	21.6	31.2	52.7	71.8	104.8	144.0	191.8	287.4
1999.....	12.0	20.9	33.0	55.3	76.4	111.7	140.7	203.7	292.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes

2007.....	3.4	8.2	12.3	13.3	17.1	23.8	30.2	36.0	32.0
2006.....	3.2	8.2	11.5	13.1	17.2	23.9	30.2	34.1	33.5
2005.....	3.7	8.5	10.2	12.9	16.6	24.3	30.2	34.4	32.5
2004.....	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
2003.....	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6

All origins 3/, male

2007.....	5.0	13.7	20.3	20.9	24.8	33.8	43.0	50.3	43.3
2006.....	5.0	13.3	18.5	19.6	25.7	33.7	42.2	48.0	43.8
2005.....	5.9	13.8	16.7	19.5	24.2	34.8	42.8	48.2	44.8
2004.....	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
2003.....	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1

All origins 3/, female

2007.....	1.8	3.0	4.9	6.4	10.4	15.6	20.8	26.9	26.7
2006.....	1.5	3.3	4.9	7.1	9.8	15.9	21.4	25.4	28.7
2005.....	1.5	3.4	4.1	6.8	9.9	15.8	21.0	25.9	26.8
2004.....	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
2003.....	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Hispanic, both sexes

2007.....	4.4	9.7	16.3	19.6	30.2	35.4	47.4	63.3	41.5
2006.....	4.1	11.5	15.4	19.8	27.2	37.2	50.6	53.0	55.2
2005.....	4.9	12.1	16.6	20.6	27.1	42.4	51.0	53.8	54.8
2004.....	5.1	9.9	13.6	20.8	25.3	37.9	46.2	52.5	50.9
2003.....	5.1	10.7	14.4	18.6	30.0	35.6	48.7	51.6	51.8
2002.....	4.6	7.7	12.8	19.6	30.1	39.1	44.0	44.6	59.9
2001.....	5.6	8.1	12.1	16.9	24.9	33.7	50.7	54.6	52.5
2000.....	5.2	6.2	11.6	16.4	24.2	39.8	45.9	62.4	58.4
1999.....	4.5	7.3	9.8	17.2	29.2	33.9	43.6	61.1	53.3

Hispanic, male

2007.....	5.5	16.6	25.9	31.5	44.3	45.5	64.0	85.2	52.9
2006.....	6.1	18.6	26.1	29.7	40.1	53.4	64.9	76.2	62.0
2005.....	7.7	19.3	27.2	31.1	36.9	60.4	65.9	72.7	71.0
2004.....	7.2	16.5	23.3	33.7	36.1	50.5	58.7	69.8	71.4
2003.....	8.2	18.3	21.1	29.0	38.7	53.8	69.3	60.3	55.4
2002.....	7.1	13.5	20.9	28.5	44.0	55.3	62.1	56.0	75.3
2001.....	10.0	12.6	19.2	24.7	35.4	48.5	68.2	67.5	64.8
2000.....	8.1	10.4	18.0	23.9	35.4	54.9	65.0	94.2	72.6
1999.....	6.9	13.3	16.0	26.8	36.3	51.4	56.2	77.0	66.9

Hispanic, female

2007.....	3.2	2.8	7.1	8.9	18.2	27.6	35.3	48.6	35.3
2006.....	2.0	4.5	5.3	11.1	16.3	24.6	40.2	37.6	51.5
2005.....	1.9	5.0	6.6	11.4	18.9	28.4	40.3	41.2	46.3
2004.....	2.8	3.5	4.6	9.6	16.3	28.1	37.2	41.1	40.3
2003.....	2.0	3.2	8.2	9.4	22.9	21.6	34.1	46.0	50.0
2002.....	2.0	*	5.4	11.9	18.8	26.7	31.1	37.2	52.2
2001.....	*	3.7	5.6	10.1	16.4	22.3	38.2	46.6	46.5
2000.....	2.3	*	5.8	9.9	15.1	28.4	32.4	43.0	51.4
1999.....	*	*	4.2	9.0	23.4	20.7	34.7	51.6	46.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 4/, both sexes

2007.....	3.2	8.1	11.9	12.8	16.0	22.8	29.0	34.3	31.5
2006.....	3.1	7.8	11.1	12.5	16.4	22.9	28.8	33.0	32.3
2005.....	3.5	8.1	9.6	12.2	15.7	22.9	28.8	33.3	31.3
2004.....	3.8	7.3	9.0	11.8	15.9	22.7	27.7	33.6	31.5
2003.....	3.6	7.1	8.0	11.1	15.4	22.7	28.5	31.2	32.6
2002.....	3.3	6.5	7.6	10.9	15.9	21.5	26.7	30.9	33.9
2001.....	3.6	5.6	7.1	11.1	15.7	20.6	25.7	31.0	33.2
2000.....	3.5	5.3	7.4	10.8	15.4	20.5	26.2	30.0	31.7
1999.....	3.3	4.7	6.8	10.4	15.5	21.1	25.3	29.8	29.7

Non-Hispanic 4/, male

2007.....	4.9	13.4	19.7	20.0	23.2	32.8	41.5	48.1	42.6
2006.....	4.8	12.6	17.8	18.8	24.5	32.2	40.6	46.2	42.6
2005.....	5.7	13.1	15.7	18.6	23.2	33.0	41.3	46.6	43.4
2004.....	6.0	12.3	14.3	17.3	22.6	32.8	37.6	47.1	40.0
2003.....	5.7	11.6	12.4	16.3	22.0	32.0	42.3	45.3	47.1
2002.....	5.5	10.8	11.6	16.3	23.3	31.3	38.1	44.0	44.8
2001.....	5.6	9.4	10.6	16.6	22.6	29.4	37.6	42.9	49.7
2000.....	5.8	8.3	11.2	15.9	22.7	29.5	38.0	44.6	43.4
1999.....	5.4	7.1	10.2	14.8	22.8	30.7	35.7	41.6	38.7

Non-Hispanic 4/, female

2007.....	1.7	3.0	4.7	6.2	9.7	14.6	19.7	25.7	26.2
2006.....	1.4	3.1	4.8	6.7	9.3	15.2	20.1	24.8	27.6
2005.....	1.4	3.2	3.8	6.4	9.2	14.8	19.8	25.0	25.8
2004.....	1.6	2.5	4.1	6.7	10.1	14.5	20.7	25.3	27.7
2003.....	1.5	2.8	3.9	6.4	9.7	15.1	18.8	22.6	26.2
2002.....	1.2	2.4	3.9	6.0	9.4	13.6	18.6	23.0	29.2
2001.....	1.7	2.0	3.7	6.1	9.7	13.7	17.4	23.9	26.4
2000.....	1.4	2.4	3.9	6.2	9.1	13.4	18.0	21.4	27.0
1999.....	1.2	2.3	3.7	6.4	9.1	13.5	18.0	22.9	26.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic White, both sexes

2007 5/.....	2.8	7.0	10.0	10.9	14.5	21.1	27.6	32.8	30.4
2006 5/.....	2.5	6.6	9.3	10.8	14.6	21.4	27.2	31.5	31.7
2005 5/.....	3.0	6.7	8.0	10.4	14.1	20.8	27.4	31.5	30.4
2004 5/.....	3.0	6.2	7.4	10.3	14.1	20.6	25.9	32.5	30.1
2003 5/.....	2.8	5.7	6.8	9.5	13.8	20.6	27.0	29.8	31.1
2002.....	2.7	5.2	6.2	9.7	14.4	19.7	25.6	29.8	32.6
2001.....	2.9	4.6	5.7	9.9	14.2	19.2	24.6	29.2	32.4
2000.....	2.8	4.2	6.1	9.4	13.8	18.8	24.7	29.2	30.8
1999.....	2.5	3.9	5.5	9.0	14.1	19.6	24.2	28.8	28.9

Non-Hispanic White, male

2007 5/.....	4.1	11.4	16.2	16.5	20.7	30.0	39.3	46.7	41.0
2006 5/.....	3.8	10.6	14.4	15.7	21.4	30.1	38.7	44.8	42.3
2005 5/.....	4.8	10.6	12.8	15.7	20.6	29.9	39.6	44.0	43.4
2004 5/.....	4.7	10.5	11.3	14.8	20.1	29.5	36.0	45.7	38.0
2003 5/.....	4.5	9.2	10.4	13.9	19.8	29.5	40.1	43.8	45.0
2002.....	4.5	8.5	9.3	14.3	21.2	28.4	36.5	41.9	43.6
2001.....	4.3	7.5	8.4	14.9	21.0	27.6	35.9	41.3	48.9
2000.....	4.4	6.4	8.9	13.6	20.3	27.1	36.5	43.5	41.6
1999.....	4.0	5.6	8.1	12.8	20.8	28.8	34.1	40.8	39.0

Non-Hispanic White, female

2007 5/.....	1.4	2.7	3.9	5.7	9.0	13.6	18.8	24.0	25.4
2006 5/.....	1.1	2.7	4.3	6.2	8.5	14.1	18.6	23.1	26.8
2005 5/.....	1.1	2.9	3.3	5.5	8.3	13.2	18.4	23.7	24.5
2004 5/.....	1.3	2.1	3.6	6.0	8.8	13.2	18.5	24.3	26.5
2003 5/.....	1.2	2.4	3.3	5.3	8.4	13.2	17.5	21.2	25.0
2002.....	0.9	2.0	3.2	5.3	8.4	12.5	17.8	22.5	27.8
2001.....	1.4	1.7	3.2	5.2	8.2	12.3	16.6	22.0	25.5
2000.....	1.2	2.0	3.4	5.5	8.0	12.1	16.3	20.8	26.5
1999.....	1.0	2.1	3.0	5.6	8.1	12.3	17.2	21.7	24.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic Black, both sexes

2007 5/.....	4.4	13.6	23.1	23.6	22.5	27.7	34.1	35.1	34.8
2006 5/.....	5.4	14.7	23.1	21.0	22.6	27.0	31.2	36.6	31.6
2005 5/.....	5.7	14.8	19.3	22.3	21.7	30.1	29.5	39.5	28.9
2004 5/.....	7.2	12.8	18.3	16.9	23.7	29.1	34.0	35.7	36.6
2003 5/.....	6.2	14.5	14.9	18.7	18.9	28.5	30.5	33.4	41.9
2002.....	5.8	12.6	15.7	14.1	19.6	28.0	30.0	32.3	38.1
2001.....	7.3	10.9	15.6	14.5	20.1	24.1	26.2	41.7	35.3
2000.....	7.1	10.9	13.0	16.2	20.7	27.3	29.2	29.0	31.8
1999.....	6.7	8.3	12.2	14.9	21.4	28.4	26.9	34.3	34.9

Non-Hispanic Black, male

2007 5/.....	6.8	23.8	41.6	42.6	37.3	42.7	51.5	46.5	50.6
2006 5/.....	8.0	25.1	43.4	36.3	37.7	43.1	46.0	45.6	34.8
2005 5/.....	9.2	27.3	35.9	37.1	36.3	45.5	45.9	59.0	31.4
2004 5/.....	12.0	23.0	32.7	27.3	37.9	45.9	39.6	51.4	47.8
2003 5/.....	10.2	26.1	24.5	29.2	26.8	43.1	43.9	46.0	63.9
2002.....	9.9	22.3	25.6	22.8	32.3	46.2	43.2	51.5	41.6
2001.....	11.8	19.1	26.1	21.7	28.1	33.1	38.4	49.9	42.7
2000.....	12.6	18.1	22.7	25.9	33.6	42.0	37.1	46.3	48.1
1999.....	12.1	14.5	19.0	22.0	32.4	42.6	40.6	43.4	30.9

Non-Hispanic Black, female

2007 5/.....	2.3	5.0	8.0	8.7	11.6	17.4	23.6	29.1	28.2
2006 5/.....	3.1	5.9	6.4	9.1	11.4	16.1	22.3	31.8	30.3
2005 5/.....	2.7	4.3	5.6	10.8	10.9	19.6	19.7	29.2	28.0
2004 5/.....	3.2	4.3	6.5	8.7	13.1	17.8	30.6	27.4	32.1
2003 5/.....	2.8	4.7	7.1	10.5	13.0	18.6	22.5	26.9	33.4
2002.....	2.3	4.4	7.5	7.3	10.2	15.8	22.1	22.3	36.8
2001.....	3.3	3.9	7.0	8.9	14.2	18.1	19.0	37.5	32.5
2000.....	2.3	4.8	5.1	8.6	11.2	17.5	24.4	20.3	25.7
1999.....	2.1	3.0	6.7	9.4	13.2	19.1	18.6	29.9	36.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

All origins 3/, both sexes

2007.....	4.1	9.0	15.6	25.3	38.0	53.4	70.5	90.4	94.6
2006.....	4.4	8.5	15.3	26.2	37.7	53.4	70.2	89.7	96.9
2005.....	4.5	8.7	15.7	25.1	37.6	52.6	69.6	88.4	97.4
2004.....	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
2003.....	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7

All origins 3/, male

2007.....	4.8	11.2	19.3	31.3	45.7	61.3	79.2	100.6	104.9
2006.....	5.2	10.7	19.2	31.9	45.3	61.2	77.3	99.0	102.1
2005.....	5.2	10.6	19.6	30.6	43.9	61.6	78.7	96.5	102.8
2004.....	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
2003.....	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5

All origins 3/, female

2007.....	3.3	6.8	12.1	19.9	31.2	46.8	64.1	83.9	89.8
2006.....	3.5	6.3	11.7	21.0	31.1	47.0	64.9	83.9	94.4
2005.....	3.8	6.9	12.0	20.0	32.1	45.3	63.0	83.4	94.9
2004.....	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
2003.....	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

Hispanic, both sexes

2007	3.0	5.8	11.7	16.5	25.8	41.2	52.0	74.1	71.5
2006	3.2	5.8	9.5	19.9	28.8	40.1	57.9	62.8	77.1
2005	3.2	5.8	11.0	21.8	31.1	42.2	56.7	74.7	78.7
2004	2.4	6.4	10.6	19.9	32.4	38.9	54.8	75.4	69.9
2003	2.9	5.6	11.5	18.3	29.2	42.7	55.7	67.6	79.8
2002	2.8	5.6	11.2	17.4	26.8	44.7	53.9	58.4	79.3
2001	2.4	5.4	11.6	17.8	29.2	46.5	55.5	71.5	83.8
2000	3.0	5.5	9.6	18.9	29.4	46.3	56.6	71.3	82.9
1999	3.7	5.8	10.4	17.0	26.8	40.0	55.5	58.7	84.2

Hispanic, male

2007	3.4	7.1	13.9	20.6	30.8	45.5	49.6	60.4	77.2
2006	3.4	7.1	11.1	21.7	35.6	45.4	65.4	74.5	65.4
2005	3.8	7.2	13.7	27.3	37.5	44.5	62.1	71.8	77.2
2004	3.2	6.8	12.1	25.1	37.0	49.7	58.2	84.9	57.4
2003	3.6	6.9	15.1	21.7	33.1	49.1	63.8	72.8	70.8
2002	3.7	7.3	13.0	19.5	32.8	51.3	54.4	58.3	60.2
2001	2.9	6.2	13.6	19.8	36.1	50.3	64.1	77.9	62.9
2000	4.2	6.5	12.7	21.6	39.5	52.5	57.6	73.6	80.6
1999	5.1	6.6	13.4	17.6	32.8	49.4	53.1	62.9	86.3

Hispanic, female

2007	2.5	4.4	9.6	12.8	21.6	37.9	53.7	83.3	68.4
2006	3.0	4.6	8.0	18.3	23.2	35.9	52.5	55.0	83.3
2005	2.6	4.5	8.6	17.0	25.8	40.4	52.8	76.7	79.5
2004	*	5.9	9.2	15.3	28.6	30.7	52.4	69.2	76.3
2003	2.3	4.4	8.2	15.4	25.9	37.7	50.0	64.3	84.4
2002	1.9	4.0	9.4	15.7	21.8	39.5	53.5	58.4	88.7
2001	*	4.6	9.8	16.0	23.5	43.6	49.4	67.6	94.0
2000	*	4.6	6.7	16.6	21.1	41.6	55.9	69.9	84.1
1999	2.4	5.0	7.7	16.4	21.9	32.9	57.3	56.3	83.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

Non-Hispanic 4/, both sexes

2007.....	4.2	9.3	15.9	26.0	38.9	54.3	71.7	91.3	95.7
2006.....	4.5	8.7	15.8	26.7	38.4	54.3	70.8	91.1	97.7
2005.....	4.6	9.0	16.1	25.3	38.1	53.2	70.3	89.1	98.1
2004.....	4.4	9.0	15.6	26.2	37.6	53.3	69.6	85.7	98.9
2003.....	4.2	8.9	15.0	25.1	36.9	53.6	68.1	86.9	95.7
2002.....	4.3	8.4	15.8	24.4	37.9	53.7	70.9	82.6	94.9
2001.....	4.2	8.8	15.1	25.3	37.4	53.0	67.6	84.6	100.6
2000.....	4.2	8.9	15.5	25.2	37.7	51.6	70.9	85.9	96.6
1999.....	4.6	8.9	16.0	25.5	38.7	53.0	67.5	82.6	100.1

Non-Hispanic 4/, male

2007.....	5.0	11.6	19.7	32.1	46.8	62.5	81.2	103.1	106.2
2006.....	5.4	11.1	19.9	32.6	46.0	62.2	77.9	100.2	104.0
2005.....	5.3	11.0	20.2	30.8	44.4	62.7	79.7	97.9	104.1
2004.....	5.7	11.4	19.1	31.8	45.7	62.3	78.7	95.1	104.1
2003.....	5.4	11.0	18.4	30.3	43.2	61.5	76.6	100.5	96.5
2002.....	5.5	10.8	19.6	30.0	44.0	61.3	80.1	93.2	102.4
2001.....	5.1	11.1	18.9	31.1	43.9	59.3	77.7	95.5	106.4
2000.....	5.5	11.4	18.9	30.6	44.9	57.2	81.4	95.5	103.5
1999.....	6.0	11.3	19.4	30.6	45.6	61.2	76.5	92.9	108.2

Non-Hispanic 4/, female

2007.....	3.4	7.1	12.3	20.4	32.0	47.5	64.7	83.9	90.8
2006.....	3.6	6.5	12.0	21.3	31.7	47.8	65.7	85.4	94.8
2005.....	3.9	7.1	12.2	20.3	32.6	45.6	63.5	83.7	95.4
2004.....	3.1	6.7	12.2	21.1	30.6	46.1	63.1	79.9	96.5
2003.....	3.1	6.9	11.8	20.4	31.5	47.3	62.1	78.6	95.3
2002.....	3.2	6.1	12.2	19.4	32.5	47.6	64.5	76.1	91.7
2001.....	3.2	6.6	11.6	20.0	31.8	48.0	60.5	78.0	98.2
2000.....	3.1	6.5	12.2	20.3	31.4	47.3	63.7	80.3	93.8
1999.....	3.2	6.6	12.7	20.8	32.8	46.5	61.2	76.5	96.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

Non-Hispanic White, both sexes

2007 5/.....	4.0	8.7	15.3	25.8	38.4	53.3	71.2	90.3	94.6
2006 5/.....	4.3	8.4	15.4	26.0	37.9	53.4	70.6	90.8	96.9
2005 5/.....	4.5	8.4	15.6	24.8	37.5	52.7	69.4	88.3	97.6
2004 5/.....	4.2	8.5	14.9	25.4	37.0	52.0	69.0	85.3	98.1
2003 5/.....	4.0	8.3	14.5	24.2	36.4	52.7	67.6	85.8	94.9
2002.....	4.2	7.9	15.3	23.8	36.7	52.3	69.8	82.2	94.6
2001.....	3.9	8.6	14.6	25.3	36.7	51.8	66.7	83.6	99.4
2000.....	4.1	8.6	14.9	24.6	36.6	50.5	69.6	84.7	94.7
1999.....	4.5	8.4	15.1	24.8	37.8	51.9	66.0	81.1	98.9

Non-Hispanic White, male

2007 5/.....	4.7	11.0	18.9	31.6	47.0	61.9	81.3	103.5	106.6
2006 5/.....	5.3	10.7	19.4	32.1	45.8	61.9	78.4	101.5	105.3
2005 5/.....	5.2	10.2	19.4	30.5	43.7	63.1	79.0	97.5	104.3
2004 5/.....	5.6	10.9	18.2	31.1	44.9	60.5	78.6	96.1	105.0
2003 5/.....	5.0	10.3	17.8	29.4	43.3	60.6	76.5	100.2	96.8
2002.....	5.3	10.3	19.0	29.5	43.3	60.1	80.0	93.6	102.9
2001.....	4.9	10.8	18.6	31.0	43.1	58.3	77.6	95.1	108.0
2000.....	5.3	11.0	18.2	30.1	44.5	56.0	80.9	94.5	103.0
1999.....	5.8	10.5	18.2	29.9	44.6	60.4	75.3	91.7	108.1

Non-Hispanic White, female

2007 5/.....	3.4	6.6	11.9	20.3	30.8	46.1	63.5	81.9	88.9
2006 5/.....	3.3	6.2	11.5	20.4	30.8	46.4	64.7	84.1	93.0
2005 5/.....	3.8	6.6	11.9	19.4	31.9	44.1	62.3	82.6	94.5
2004 5/.....	2.9	6.2	11.7	20.1	30.0	45.0	61.9	78.7	95.1
2003 5/.....	3.0	6.4	11.2	19.4	30.2	46.2	61.2	76.9	94.0
2002.....	3.1	5.6	11.8	18.4	30.9	45.9	62.6	75.2	91.0
2001.....	3.0	6.4	10.9	19.9	31.0	46.6	58.9	76.7	95.8
2000.....	3.0	6.3	11.7	19.6	29.8	46.0	61.6	78.9	91.4
1999.....	3.1	6.4	12.1	20.1	31.8	45.1	59.5	74.9	95.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

Non-Hispanic Black, both sexes

2007 5/.....	5.5	14.9	24.1	33.1	51.7	70.2	89.9	112.5	116.1
2006 5/.....	6.4	13.0	22.2	38.1	50.4	68.5	86.7	102.1	114.7
2005 5/.....	6.4	14.0	22.6	34.5	50.1	67.3	88.4	106.8	111.7
2004 5/.....	6.1	13.6	22.9	37.0	49.0	71.7	85.8	93.9	115.1
2003 5/.....	6.6	14.8	21.1	36.5	48.2	69.3	84.3	108.6	113.3
2002.....	5.8	12.6	22.9	35.2	51.3	74.8	89.3	93.2	108.3
2001.....	6.5	12.5	21.1	31.9	48.9	71.5	85.2	106.0	122.0
2000.....	5.9	13.5	23.0	33.9	51.1	70.6	94.9	105.2	121.2
1999.....	6.4	14.4	25.6	32.7	51.9	70.8	88.3	105.7	119.0

Non-Hispanic Black, male

2007 5/.....	7.1	18.6	31.8	44.5	57.9	78.8	98.0	106.8	109.1
2006 5/.....	6.8	17.3	29.3	46.9	59.9	76.9	88.8	100.4	92.5
2005 5/.....	7.4	17.8	30.7	40.7	60.7	72.2	97.4	119.4	101.8
2004 5/.....	7.7	16.4	30.1	45.6	61.8	89.8	90.9	91.8	102.6
2003 5/.....	9.2	19.1	26.8	43.3	51.4	82.6	89.7	114.0	106.5
2002.....	7.4	16.8	29.6	42.0	57.6	85.5	92.0	106.1	102.8
2001.....	8.0	15.6	24.6	38.9	60.2	78.8	94.8	110.4	100.5
2000.....	7.4	18.0	29.5	41.0	56.8	80.0	102.6	117.6	110.4
1999.....	8.6	20.0	34.8	40.5	61.1	78.6	101.5	118.7	115.3

Non-Hispanic Black, female

2007 5/.....	4.2	11.8	17.7	24.3	47.2	64.3	85.0	115.4	119.0
2006 5/.....	6.0	9.3	16.3	31.2	43.3	62.8	85.5	102.9	123.7
2005 5/.....	5.5	10.8	16.0	29.7	42.3	64.0	83.1	100.2	115.7
2004 5/.....	4.7	11.2	16.9	30.3	39.4	59.5	82.8	95.0	120.1
2003 5/.....	4.3	11.2	16.4	31.2	45.8	60.4	81.0	105.8	116.0
2002.....	4.3	9.1	17.5	29.9	46.7	67.7	87.7	86.4	110.4
2001.....	5.2	9.8	18.2	26.5	40.6	66.5	79.5	103.7	130.0
2000.....	4.5	9.7	17.7	28.3	47.0	64.4	90.3	98.9	125.2
1999.....	4.4	9.6	18.1	26.6	45.1	65.7	80.4	99.2	120.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

All origins 3/, both sexes

2007.....	0.6	1.4	2.3	3.6	4.9	5.3	6.6	6.8	6.0
2006.....	0.7	1.4	2.6	3.8	5.2	6.2	6.9	7.4	6.3
2005.....	0.7	1.3	2.4	3.9	5.8	6.7	6.8	7.3	6.3
2004.....	0.7	1.4	2.6	4.1	5.3	6.2	7.1	6.0	6.5
2003.....	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0

All origins 3/, male

2007.....	1.1	2.4	4.1	6.1	8.4	8.8	12.4	13.6	13.9
2006.....	1.2	2.3	4.4	6.2	9.2	10.5	12.5	14.7	14.8
2005.....	1.0	2.1	4.0	6.7	10.2	11.4	12.2	14.5	15.7
2004.....	1.2	2.4	4.4	6.8	9.4	11.0	13.3	12.4	15.6
2003.....	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8

All origins 3/, female

2007.....	0.2	0.5	0.6	1.2	1.8	2.5	2.3	2.5	2.3
2006.....	0.3	0.5	1.0	1.5	1.7	2.7	2.8	2.8	2.3
2005.....	0.4	0.6	0.8	1.3	1.9	3.0	2.8	2.8	1.9
2004.....	0.2	0.5	0.9	1.6	1.7	2.2	2.6	2.1	2.4
2003.....	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

Non-Hispanic 4/, both sexes

2007.....	0.7	1.5	2.4	3.7	5.1	5.4	6.7	7.0	6.1
2006.....	0.8	1.4	2.7	3.9	5.4	6.4	7.0	7.3	6.3
2005.....	0.7	1.4	2.5	4.0	6.0	6.8	6.7	7.4	6.2
2004.....	0.8	1.5	2.6	4.1	5.2	6.3	7.0	6.0	6.5
2003.....	0.8	1.4	2.6	4.7	6.0	6.8	6.6	7.3	6.5
2002.....	0.8	1.5	2.8	4.4	5.4	6.7	8.0	7.3	5.4
2001.....	0.7	1.6	3.0	4.6	5.5	7.4	7.5	7.0	6.7
2000.....	0.8	1.6	3.0	4.7	6.1	7.4	8.2	8.0	6.5
1999.....	0.8	1.5	3.2	4.8	6.2	7.5	7.2	7.3	6.8

Non-Hispanic 4/, male

2007.....	1.1	2.6	4.2	6.2	8.6	8.8	12.5	13.9	14.0
2006.....	1.2	2.4	4.5	6.4	9.4	10.7	12.5	14.4	14.7
2005.....	1.1	2.2	4.2	6.9	10.5	11.4	11.9	14.6	15.7
2004.....	1.3	2.5	4.5	6.8	9.2	11.1	13.2	12.3	15.6
2003.....	1.2	2.5	4.3	8.0	10.4	11.6	12.7	15.3	15.0
2002.....	1.3	2.5	4.7	7.4	9.1	11.3	14.9	15.1	13.2
2001.....	1.2	2.6	5.0	7.6	9.4	13.0	14.5	14.0	18.4
2000.....	1.3	2.6	5.0	8.1	10.0	13.1	15.1	16.7	17.3
1999.....	1.3	2.7	5.5	8.1	10.4	12.6	13.0	15.6	17.3

Non-Hispanic 4/, female

2007.....	0.3	0.5	0.7	1.3	2.0	2.7	2.4	2.6	2.3
2006.....	0.3	0.5	1.0	1.6	1.8	2.9	3.0	2.9	2.4
2005.....	0.4	0.6	0.9	1.4	2.0	3.1	2.9	2.9	1.9
2004.....	0.3	0.6	0.9	1.7	1.8	2.4	2.6	2.1	2.5
2003.....	0.3	0.4	1.1	1.7	2.1	2.9	2.3	2.5	2.7
2002.....	0.3	0.5	1.0	1.6	2.3	3.0	3.1	2.5	2.1
2001.....	0.3	0.5	1.1	1.8	2.3	3.0	2.6	2.8	1.8
2000.....	0.4	0.6	1.0	1.5	2.7	2.9	3.4	2.8	2.1
1999.....	0.2	0.5	1.1	1.8	2.5	3.4	3.1	2.5	2.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

Non-Hispanic White, both sexes

2007 5/.....	0.6	1.4	2.1	3.3	4.7	5.1	6.3	6.8	5.9
2006 5/.....	0.7	1.3	2.5	3.7	4.9	6.2	6.9	7.1	6.2
2005 5/.....	0.7	1.2	2.2	3.5	5.5	6.5	6.7	7.2	5.9
2004 5/.....	0.7	1.3	2.4	3.9	4.9	6.0	6.8	5.7	6.4
2003 5/.....	0.6	1.2	2.4	4.2	5.6	6.4	6.4	7.0	6.1
2002.....	0.6	1.3	2.5	3.9	5.2	6.4	7.5	7.1	5.5
2001.....	0.5	1.3	2.6	4.2	5.2	6.8	7.0	6.8	6.5
2000.....	0.7	1.3	2.5	4.0	5.7	6.8	7.6	7.8	6.5
1999.....	0.6	1.3	2.8	4.3	5.7	7.2	7.0	7.2	6.5

Non-Hispanic White, male

2007 5/.....	1.0	2.2	3.7	5.5	7.8	8.1	11.7	13.1	13.4
2006 5/.....	1.0	2.1	4.1	6.0	8.4	10.1	12.0	13.9	14.5
2005 5/.....	0.9	1.8	3.6	5.9	9.6	10.4	11.8	14.0	14.8
2004 5/.....	1.2	2.0	3.9	6.2	8.5	10.4	12.6	11.5	15.0
2003 5/.....	1.1	2.0	3.7	7.2	9.4	10.6	11.9	14.3	14.1
2002.....	1.1	2.1	4.1	6.4	8.5	10.6	13.8	14.8	13.3
2001.....	0.9	2.1	4.3	7.0	8.6	11.9	13.3	13.3	18.3
2000.....	1.1	2.1	4.1	6.9	9.2	11.9	14.1	16.3	17.4
1999.....	1.0	2.3	4.7	7.3	9.5	11.9	12.7	15.0	16.0

Non-Hispanic White, female

2007 5/.....	0.2	0.6	0.7	1.2	2.0	2.5	2.3	2.7	2.3
2006 5/.....	0.3	0.5	0.9	1.6	1.8	3.0	3.1	2.8	2.3
2005 5/.....	0.4	0.5	0.8	1.4	1.9	3.1	2.9	3.0	1.8
2004 5/.....	0.2	0.6	0.9	1.7	1.8	2.3	2.6	2.1	2.5
2003 5/.....	*	0.4	1.0	1.5	2.2	2.9	2.4	2.5	2.6
2002.....	*	0.4	1.0	1.6	2.3	2.9	3.0	2.3	2.2
2001.....	*	0.5	0.9	1.6	2.1	2.7	2.6	2.9	1.6
2000.....	0.3	0.6	0.9	1.4	2.6	2.7	3.1	2.8	2.1
1999.....	*	0.3	1.0	1.6	2.4	3.4	3.1	2.6	2.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

Non-Hispanic Black, both sexes

2007 5/.....	1.3	3.2	4.8	7.9	9.6	10.2	13.4	9.3	8.4
2006 5/.....	1.6	2.9	5.5	6.8	11.2	9.9	10.5	9.7	8.2
2005 5/.....	1.3	3.4	5.7	9.3	11.8	11.5	8.9	11.2	10.4
2004 5/.....	1.3	3.6	5.6	7.9	9.1	11.7	11.4	11.4	8.1
2003 5/.....	1.7	3.6	5.9	10.0	10.7	12.8	11.3	12.7	11.3
2002.....	1.9	3.7	6.3	9.6	9.4	12.1	15.6	11.2	*
2001.....	2.0	3.9	7.1	8.9	9.6	13.9	13.6	9.4	8.4
2000.....	2.1	4.1	7.7	11.0	10.7	15.5	15.9	11.8	7.0
1999.....	2.2	4.0	7.7	10.0	11.5	11.9	10.4	9.4	8.4

Non-Hispanic Black, male

2007 5/.....	2.3	6.3	10.0	14.8	19.1	17.1	28.3	23.6	23.6
2006 5/.....	2.9	5.2	9.6	12.6	22.2	19.9	23.0	21.8	19.2
2005 5/.....	2.2	5.2	10.6	18.9	22.8	22.1	17.4	26.6	29.5
2004 5/.....	2.2	6.8	11.2	14.6	18.5	22.6	24.7	26.4	23.4
2003 5/.....	2.8	6.9	11.0	18.1	22.8	25.6	26.2	31.7	27.7
2002.....	3.3	6.2	11.8	19.3	17.4	22.9	33.8	22.6	*
2001.....	3.7	7.1	12.9	15.9	17.7	26.2	31.3	22.9	*
2000.....	3.3	7.7	14.7	21.0	20.0	30.4	29.9	27.5	*
1999.....	4.0	7.1	14.8	18.1	21.3	23.2	20.3	26.6	25.0

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	5.4	*	*	*
2006 5/.....	*	*	2.0	*	*	*	*	*	*
2005 5/.....	*	1.8	*	*	3.6	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	3.7	*	*	*	*	*
2002.....	*	*	*	*	*	4.8	*	*	*
2001.....	*	*	*	3.5	*	5.7	*	*	*
2000.....	*	*	*	*	3.9	5.7	7.6	*	*
1999.....	*	*	*	3.8	4.3	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/, both sexes

2007.....	19.0	38.7	69.2	128.4	213.6	292.9	363.2	381.8	299.8
2006.....	19.9	39.4	71.8	136.5	216.9	297.1	366.8	382.3	300.5
2005.....	20.4	40.1	73.8	142.7	218.7	308.3	372.3	380.0	302.3
2004.....	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
2003.....	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0

All origins 3/, male

2007.....	20.3	45.1	85.0	156.0	265.6	368.1	475.3	530.8	453.0
2006.....	21.6	47.5	88.3	167.8	272.4	378.8	486.0	544.1	457.7
2005.....	22.8	49.0	91.0	177.3	275.5	399.2	497.5	540.5	468.0
2004.....	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
2003.....	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2

All origins 3/, female

2007.....	17.8	32.5	54.4	103.1	168.1	231.5	280.6	287.7	227.0
2006.....	18.3	31.6	56.2	107.8	168.4	230.7	279.5	281.3	226.9
2005.....	18.1	31.7	57.6	111.2	169.1	234.7	281.3	280.3	226.2
2004.....	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
2003.....	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Hispanic, both sexes

2007	5.3	12.6	23.4	43.5	87.4	116.4	159.8	178.0	169.8
2006	5.6	12.1	23.0	43.4	87.0	114.0	159.6	180.6	160.8
2005	6.5	13.5	24.7	51.5	80.8	130.7	173.0	199.4	177.1
2004	6.3	13.5	26.5	48.0	82.4	128.6	173.2	187.3	193.1
2003	6.6	15.0	27.2	53.2	88.6	127.6	177.3	194.3	189.2
2002	6.0	15.0	32.0	53.1	87.4	139.2	181.6	187.2	189.0
2001	6.6	16.0	30.9	54.6	89.4	131.8	177.2	191.2	200.7
2000	7.0	16.0	31.6	49.2	90.9	135.4	182.1	228.9	213.7
1999	6.6	15.4	33.2	58.0	96.1	134.2	178.5	204.4	227.3

Hispanic, male

2007	6.0	16.0	29.4	60.4	122.9	165.0	229.5	272.0	256.1
2006	6.3	16.1	31.6	64.5	122.7	163.6	236.9	281.7	263.9
2005	7.5	17.9	34.9	78.1	118.1	196.2	259.8	317.0	269.0
2004	7.8	19.0	36.6	69.5	122.5	190.8	274.1	288.8	287.2
2003	7.0	20.1	39.9	77.8	137.5	193.9	272.2	285.9	290.9
2002	6.8	20.3	47.0	79.8	133.1	198.8	296.3	304.3	307.9
2001	8.6	21.6	43.8	80.9	130.8	206.3	281.7	316.8	335.2
2000	9.1	22.3	46.0	75.1	130.9	214.9	304.1	374.0	368.8
1999	8.8	21.4	50.6	90.5	152.2	205.7	294.1	360.0	362.3

Hispanic, female

2007	4.6	9.3	17.7	28.3	57.4	78.4	109.2	115.3	123.0
2006	4.8	8.2	14.9	24.7	57.1	75.4	103.6	113.4	106.0
2005	5.4	9.3	15.1	28.0	49.7	79.9	110.6	121.4	129.0
2004	4.7	8.1	17.0	29.1	49.3	80.5	101.5	120.3	144.8
2003	6.2	10.0	15.5	31.7	48.5	76.7	109.9	134.7	137.5
2002	5.2	10.0	18.2	29.7	50.0	93.3	99.6	112.4	130.1
2001	4.6	10.7	19.2	31.7	56.0	75.0	103.1	113.4	135.0
2000	4.8	10.0	18.5	26.8	58.6	75.1	95.7	140.7	137.5
1999	4.4	9.6	17.5	30.0	50.6	80.4	96.9	111.6	162.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 4/, both sexes

2007.....	20.8	41.5	73.5	135.4	223.6	306.3	377.0	393.8	306.5
2006.....	21.7	42.2	76.0	144.0	227.0	310.5	380.2	393.3	306.9
2005.....	22.1	42.8	78.0	149.9	229.3	320.6	384.5	389.2	307.9
2004.....	22.2	43.0	82.1	152.2	237.2	323.3	383.7	384.0	301.6
2003.....	22.8	43.1	86.1	156.2	240.0	325.7	386.5	388.2	303.0
2002.....	21.9	43.9	90.5	160.2	243.4	331.9	386.3	386.0	301.0
2001.....	21.9	44.4	93.2	161.7	249.8	334.1	382.4	375.1	305.8
2000.....	22.3	45.9	96.8	167.6	255.5	335.4	376.0	379.5	304.5
1999.....	21.1	46.6	97.2	167.7	252.4	331.5	368.7	367.5	298.8

Non-Hispanic 4/, male

2007.....	22.3	48.3	90.1	163.8	276.6	382.8	491.8	546.6	464.2
2006.....	23.6	50.8	93.2	175.9	283.6	393.9	502.0	558.8	467.4
2005.....	24.7	52.1	95.7	184.8	287.2	412.8	512.0	552.3	477.9
2004.....	25.3	52.8	100.5	187.7	297.3	414.2	521.4	542.1	474.8
2003.....	26.7	52.7	106.1	191.5	302.3	424.0	526.5	553.1	483.1
2002.....	25.3	54.1	111.7	200.8	308.3	436.2	525.7	556.7	488.9
2001.....	25.9	54.6	114.3	203.5	324.7	447.4	533.1	549.4	518.4
2000.....	27.2	55.6	120.7	210.9	334.0	451.9	525.2	564.0	526.2
1999.....	25.7	56.8	122.7	215.3	332.1	450.2	521.5	557.1	534.6

Non-Hispanic 4/, female

2007.....	19.4	35.0	57.8	109.3	177.0	243.4	292.4	297.6	232.1
2006.....	19.9	34.0	59.8	114.8	177.4	242.5	290.9	290.2	232.3
2005.....	19.6	33.9	61.3	117.9	178.6	245.8	292.0	288.2	230.3
2004.....	19.2	33.7	64.7	119.9	184.6	250.0	284.8	287.0	224.2
2003.....	18.9	33.9	67.3	124.0	185.8	246.6	286.8	287.7	223.9
2002.....	18.5	34.1	70.7	123.3	187.2	248.2	287.9	282.4	220.0
2001.....	17.9	34.6	73.3	123.7	185.0	243.9	276.9	270.8	217.4
2000.....	17.5	36.6	74.4	128.2	188.0	243.4	272.1	270.8	214.8
1999.....	16.7	36.8	73.3	124.5	183.9	238.6	262.5	256.5	204.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic White, both sexes

2007 5/.....	20.1	40.4	72.8	136.1	227.9	314.9	386.4	403.1	310.7
2006 5/.....	21.1	40.4	75.4	144.8	232.0	318.8	390.7	400.4	313.0
2005 5/.....	21.4	41.5	76.5	150.5	234.1	328.9	394.1	395.4	311.3
2004 5/.....	21.5	41.5	80.5	153.8	241.0	331.6	390.5	388.4	306.3
2003 5/.....	21.5	41.6	85.3	157.8	244.7	332.7	392.4	392.9	306.7
2002.....	21.0	42.4	89.4	160.6	247.9	339.0	392.3	392.1	304.8
2001.....	20.7	43.2	92.7	163.9	253.2	339.3	386.7	377.9	308.6
2000.....	20.8	44.4	96.3	169.9	258.9	339.0	378.9	385.7	306.8
1999.....	19.6	45.0	96.5	168.4	254.2	334.6	370.3	370.4	300.5

Non-Hispanic White, male

2007 5/.....	20.7	45.8	86.6	160.8	276.1	386.0	496.9	552.8	464.7
2006 5/.....	22.4	47.6	89.8	172.3	283.6	396.7	507.6	564.4	474.2
2005 5/.....	23.4	49.8	91.0	181.1	287.6	416.1	518.6	556.2	480.2
2004 5/.....	23.6	49.3	96.2	184.0	297.2	416.1	522.4	542.5	480.3
2003 5/.....	24.8	49.5	102.1	188.4	302.0	426.0	525.8	553.6	486.8
2002.....	23.5	50.8	107.5	195.6	307.6	438.5	525.3	557.9	493.3
2001.....	23.8	51.4	110.5	201.8	322.2	446.0	530.5	545.9	520.0
2000.....	25.0	51.8	116.5	209.8	331.3	448.7	520.4	565.8	527.3
1999.....	23.2	52.9	118.9	211.0	328.0	446.4	517.2	553.9	536.9

Non-Hispanic White, female

2007 5/.....	19.5	35.0	59.5	112.9	184.6	255.3	303.2	307.7	237.7
2006 5/.....	19.7	33.4	61.5	119.1	185.8	253.8	303.2	297.2	237.8
2005 5/.....	19.3	33.3	62.5	121.9	186.3	256.5	301.9	295.0	234.1
2004 5/.....	19.3	33.8	65.5	125.5	190.9	261.8	293.9	293.0	228.4
2003 5/.....	18.3	33.8	69.1	129.4	193.8	255.9	295.8	294.2	227.4
2002.....	18.5	34.2	72.1	128.1	194.9	257.7	297.2	290.6	223.4
2001.....	17.7	35.1	75.5	128.8	192.3	252.7	284.9	276.5	220.6
2000.....	16.7	37.2	77.0	133.0	195.2	250.6	279.5	278.7	218.0
1999.....	16.1	37.2	75.1	128.9	189.4	245.5	267.2	262.1	206.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic Black, both sexes

2007 5/.....	30.3	58.9	97.8	166.6	251.3	307.5	363.6	362.7	287.9
2006 5/.....	31.2	64.1	100.2	177.7	250.8	311.7	361.4	384.1	268.4
2005 5/.....	31.7	62.0	110.3	184.2	253.1	322.1	368.2	380.9	294.1
2004 5/.....	31.8	64.2	116.3	182.5	262.0	314.7	387.9	393.4	281.5
2003 5/.....	36.2	63.4	117.7	186.9	261.7	334.4	393.2	383.7	283.5
2002.....	33.3	66.0	123.3	197.4	270.6	336.0	395.6	370.2	274.9
2001.....	33.6	64.6	121.4	185.0	277.6	353.4	401.1	375.0	281.6
2000.....	36.4	70.5	124.8	192.2	284.2	363.5	399.7	347.2	291.0
1999.....	35.8	71.8	127.4	202.3	286.3	356.2	406.5	365.7	293.3

Non-Hispanic Black, male

2007 5/.....	37.0	77.2	143.4	240.3	368.6	450.3	537.0	581.8	514.8
2006 5/.....	36.9	84.5	144.3	259.1	371.5	461.2	553.8	606.1	448.1
2005 5/.....	39.3	81.3	159.9	268.1	371.1	478.6	554.3	608.7	503.4
2004 5/.....	42.1	91.5	164.8	276.5	375.6	476.8	609.0	636.7	488.8
2003 5/.....	46.3	88.7	172.0	274.1	383.7	503.2	627.5	640.7	509.3
2002.....	44.0	93.7	175.7	297.6	403.1	515.9	638.5	626.3	490.1
2001.....	46.2	93.9	177.0	271.9	426.0	561.5	668.4	667.2	550.9
2000.....	48.8	102.3	187.8	276.6	438.7	568.2	664.3	623.1	567.1
1999.....	49.8	104.8	189.6	307.4	434.8	564.3	662.7	669.5	580.0

Non-Hispanic Black, female

2007 5/.....	24.6	43.5	60.5	108.9	165.2	209.1	259.4	247.6	193.9
2006 5/.....	26.2	46.9	63.9	114.4	161.8	209.1	246.6	267.0	195.1
2005 5/.....	25.2	45.8	69.4	119.0	165.8	215.1	257.5	260.7	209.9
2004 5/.....	22.9	41.3	76.3	109.4	177.8	204.9	257.0	265.4	199.6
2003 5/.....	27.4	42.1	73.1	119.0	171.2	220.6	254.1	249.2	196.0
2002.....	24.1	42.5	80.4	119.2	172.3	215.4	250.6	237.0	192.7
2001.....	22.6	39.8	75.8	116.7	168.1	214.7	241.6	226.1	181.8
2000.....	25.7	43.4	73.3	125.6	170.3	227.5	240.2	208.2	188.4
1999.....	23.7	43.9	76.8	120.0	176.2	218.0	252.2	215.1	186.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

All origins 3/, both sexes

2007.....	2.2	3.2	4.8	6.0	7.8	10.4	14.1	17.5	19.9
2006.....	2.2	3.2	4.5	6.2	8.2	10.8	14.0	17.8	20.1
2005.....	2.2	3.5	4.6	5.9	8.1	10.9	15.1	16.2	19.8
2004.....	2.3	3.3	4.7	6.1	8.1	10.6	13.6	16.3	17.7
2003.....	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4

All origins 3/, male

2007.....	2.7	4.4	6.7	8.9	11.6	16.0	22.1	28.6	35.8
2006.....	2.8	4.3	6.2	9.0	11.9	15.9	22.5	29.9	36.7
2005.....	2.7	4.6	6.4	7.9	11.9	16.2	23.5	26.0	34.9
2004.....	2.9	4.5	6.6	8.9	11.8	16.1	21.8	26.7	31.4
2003.....	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5

All origins 3/, female

2007.....	1.7	2.0	3.0	3.3	4.5	5.9	8.3	10.6	12.4
2006.....	1.7	2.1	2.9	3.6	5.0	6.7	7.8	10.2	12.4
2005.....	1.8	2.4	2.9	4.2	4.8	6.6	9.0	10.1	12.9
2004.....	1.7	2.2	3.0	3.6	4.9	6.2	7.7	9.9	11.5
2003.....	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

Non-Hispanic 4/, both sexes

2007.....	2.4	3.4	5.1	6.4	8.2	11.0	14.8	18.3	20.6
2006.....	2.4	3.5	4.8	6.5	8.7	11.4	14.6	18.4	20.8
2005.....	2.4	3.7	4.9	6.3	8.6	11.4	15.8	16.9	20.5
2004.....	2.5	3.5	5.0	6.5	8.6	11.2	14.2	16.9	18.3
2003.....	2.7	3.5	5.2	6.7	7.7	10.7	14.5	16.0	18.1
2002.....	2.5	3.6	4.7	6.3	8.3	10.5	13.3	15.3	18.2
2001.....	2.8	3.9	4.9	6.5	8.2	11.2	12.7	15.1	18.3
2000.....	2.8	4.0	4.9	6.6	8.3	10.3	12.3	15.3	19.3
1999.....	2.8	3.8	4.9	6.4	8.6	10.1	13.2	14.6	16.7

Non-Hispanic 4/, male

2007.....	3.0	4.8	7.3	9.5	12.2	16.9	23.3	29.9	37.2
2006.....	3.0	4.7	6.6	9.6	12.7	16.8	23.7	31.1	38.0
2005.....	2.9	4.9	6.8	8.3	12.5	17.0	24.6	27.2	36.2
2004.....	3.2	4.8	7.0	9.4	12.5	17.0	22.7	27.8	32.7
2003.....	3.4	4.8	7.3	9.5	10.9	15.8	23.4	25.4	30.9
2002.....	3.5	4.7	6.5	8.9	12.0	15.6	20.6	24.0	30.2
2001.....	3.5	5.1	6.6	9.3	12.5	17.0	20.2	24.8	31.0
2000.....	3.4	5.5	6.7	9.2	11.9	15.2	19.0	25.2	32.6
1999.....	3.7	4.8	7.0	8.8	12.4	15.2	20.4	23.3	28.2

Non-Hispanic 4/, female

2007.....	1.9	2.2	3.2	3.5	4.7	6.2	8.6	11.0	12.7
2006.....	1.9	2.3	3.1	3.7	5.2	7.0	8.0	10.5	12.8
2005.....	1.9	2.6	3.1	4.5	5.2	6.8	9.4	10.5	13.4
2004.....	1.9	2.4	3.1	3.8	5.2	6.5	8.0	10.2	11.9
2003.....	2.0	2.3	3.2	4.2	4.8	6.6	8.1	10.2	12.4
2002.....	1.5	2.6	3.0	3.8	5.1	6.4	8.1	10.0	12.9
2001.....	2.1	2.8	3.2	3.9	4.6	6.6	7.5	9.2	12.9
2000.....	2.2	2.6	3.1	4.3	5.2	6.5	7.7	9.4	13.9
1999.....	2.0	2.7	2.9	4.2	5.4	6.1	8.1	9.5	12.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

Non-Hispanic White, both sexes

2007 5/.....	2.9	4.2	6.1	7.4	9.5	12.6	16.5	20.2	22.4
2006 5/.....	3.0	4.1	5.6	7.5	10.2	13.0	16.3	20.2	22.5
2005 5/.....	2.9	4.5	5.7	7.2	9.9	13.0	17.7	18.4	22.2
2004 5/.....	3.1	4.2	5.9	7.5	9.8	12.7	15.8	18.3	19.9
2003 5/.....	3.2	4.2	6.0	7.7	8.8	12.1	15.9	17.5	19.6
2002.....	3.0	4.3	5.4	7.2	9.5	11.7	14.7	16.8	19.6
2001.....	3.4	4.6	5.7	7.5	9.3	12.5	14.0	16.3	19.9
2000.....	3.4	4.7	5.6	7.6	9.4	11.4	13.5	16.4	20.8
1999.....	3.4	4.4	5.7	7.3	9.8	11.2	14.4	15.7	17.9

Non-Hispanic White, male

2007 5/.....	3.5	5.7	8.5	10.9	13.9	19.3	25.8	33.2	40.6
2006 5/.....	3.7	5.5	7.7	10.9	14.7	19.1	26.1	34.3	41.3
2005 5/.....	3.5	5.9	7.8	9.5	14.4	19.3	27.3	29.7	39.7
2004 5/.....	3.9	5.6	8.1	10.8	14.3	19.3	25.2	30.2	35.8
2003 5/.....	4.0	5.6	8.4	10.8	12.4	17.9	25.7	27.8	33.8
2002.....	4.2	5.5	7.4	10.2	13.6	17.2	22.7	26.3	32.9
2001.....	4.2	6.0	7.6	10.7	14.0	18.8	22.2	27.0	34.1
2000.....	4.1	6.4	7.7	10.6	13.3	16.8	20.8	27.1	35.7
1999.....	4.4	5.6	8.1	10.0	14.0	16.9	22.2	25.2	30.5

Non-Hispanic White, female

2007 5/.....	2.2	2.7	3.8	4.0	5.4	7.0	9.6	11.9	13.8
2006 5/.....	2.3	2.7	3.6	4.4	6.1	8.0	8.9	11.4	13.8
2005 5/.....	2.3	3.1	3.7	5.1	5.9	7.7	10.6	11.4	14.3
2004 5/.....	2.3	2.8	3.7	4.4	5.8	7.4	9.0	11.0	12.8
2003 5/.....	2.5	2.8	3.7	4.9	5.5	7.4	8.8	11.2	13.3
2002.....	1.8	3.1	3.4	4.4	5.8	7.1	9.0	10.9	13.8
2001.....	2.5	3.3	3.8	4.5	5.1	7.3	8.3	9.9	13.9
2000.....	2.7	3.0	3.6	4.9	6.0	7.1	8.4	10.0	14.7
1999.....	2.4	3.2	3.4	4.8	6.1	6.7	8.9	10.2	12.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

All origins 3/, both sexes

2007.....	11.0	16.6	23.4	31.0	39.4	45.8	60.3	81.8	116.6
2006.....	11.6	17.0	24.7	32.7	38.0	47.1	63.1	82.7	116.7
2005.....	11.8	17.9	25.0	33.1	38.7	49.3	63.5	82.9	122.7
2004.....	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
2003.....	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4

All origins 3/, male

2007.....	*	0.2	0.3	0.6	0.8	1.3	1.6	2.4	3.4
2006.....	*	0.3	0.4	0.8	0.8	1.2	1.8	2.6	3.0
2005.....	*	0.3	0.4	0.7	1.0	1.7	1.8	2.5	2.4
2004.....	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
2003.....	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5

All origins 3/, female

2007.....	21.6	32.3	45.1	58.9	73.2	82.3	103.5	131.9	170.4
2006.....	22.7	33.0	47.5	61.9	70.4	84.4	108.0	132.9	169.9
2005.....	23.2	34.9	48.3	62.6	71.6	87.9	108.3	132.8	177.9
2004.....	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
2003.....	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Hispanic, both sexes

2007	7.6	12.0	15.8	20.6	24.8	29.5	35.5	45.9	57.2
2006	7.9	12.1	16.0	23.4	23.4	30.1	38.5	44.8	58.7
2005	7.6	13.0	16.7	20.4	25.6	26.6	38.2	51.7	62.5
2004	7.3	13.2	18.8	22.0	29.1	29.8	38.8	45.0	72.3
2003	8.4	13.3	19.8	21.4	27.5	26.5	42.6	41.0	77.7
2002	8.9	13.7	15.9	20.0	26.3	28.6	36.8	55.3	71.0
2001	9.6	15.2	20.5	24.1	25.1	31.7	38.8	40.7	68.8
2000	10.4	14.5	21.2	20.7	30.5	32.3	41.3	53.5	71.7
1999	9.5	15.7	18.0	21.6	25.2	31.5	37.1	55.2	86.3

Hispanic, male

2007	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Hispanic, female

2007	15.7	24.0	30.5	38.6	45.3	51.9	60.5	75.0	88.3
2006	16.2	24.0	31.1	43.9	42.6	53.2	65.2	74.1	89.9
2005	15.4	25.8	32.0	38.4	46.7	47.2	65.6	84.8	95.1
2004	14.6	25.9	36.1	41.1	53.1	50.8	66.3	73.5	108.8
2003	16.9	26.2	37.9	40.1	50.1	46.2	72.9	67.7	114.8
2002	17.9	26.4	30.5	37.3	47.6	50.0	63.1	89.1	105.3
2001	19.2	29.5	39.1	44.7	44.2	55.5	64.9	64.3	102.4
2000	20.7	28.0	39.9	38.2	54.7	55.2	69.5	84.2	105.8
1999	18.8	30.4	33.6	40.3	45.0	54.0	62.8	87.2	127.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Non-Hispanic 4/, both sexes

2007	11.4	17.1	24.1	31.8	40.6	47.1	62.0	83.8	119.6
2006	12.0	17.5	25.4	33.5	39.1	48.3	64.7	84.8	119.4
2005	12.3	18.4	25.7	34.1	39.7	50.8	65.0	84.4	125.4
2004	12.2	19.2	26.7	33.9	41.7	50.9	66.1	84.1	126.3
2003	13.0	19.6	27.1	34.4	40.7	53.3	67.1	89.2	134.7
2002	13.3	20.2	26.9	34.7	42.2	53.7	68.6	91.0	136.8
2001	13.8	21.1	27.6	34.7	42.7	55.3	70.4	89.8	136.4
2000	13.9	21.3	28.6	35.9	44.1	56.4	72.0	92.1	149.7
1999	13.6	21.4	28.8	35.6	44.5	57.1	71.3	93.5	146.2

Non-Hispanic 4/, male

2007	*	0.2	0.3	0.6	0.9	1.3	1.7	2.4	3.6
2006	*	0.3	0.5	0.8	0.8	1.3	1.8	2.7	3.2
2005	*	0.3	0.4	0.7	1.0	1.8	1.9	2.6	2.6
2004	*	*	0.4	0.6	1.0	1.2	2.2	2.1	3.3
2003	*	0.4	0.4	0.8	1.0	1.3	1.6	3.0	3.8
2002	*	0.3	0.4	0.7	0.8	1.3	2.0	2.8	3.5
2001	*	0.4	0.5	0.8	1.2	1.8	2.3	2.4	3.7
2000	*	0.3	0.7	0.8	1.1	1.7	2.2	2.4	4.2
1999	*	0.4	0.6	0.7	1.1	1.4	1.9	2.4	4.7

Non-Hispanic 4/, female

2007	22.3	33.1	46.5	60.6	75.5	84.6	106.4	135.1	174.4
2006	23.5	33.9	48.9	63.4	72.6	86.7	110.8	135.9	173.5
2005	24.1	35.7	49.7	64.6	73.5	90.7	110.8	135.0	181.5
2004	23.9	37.3	51.6	64.2	77.2	91.0	112.1	134.5	181.3
2003	25.5	37.9	52.3	65.1	75.3	95.2	113.7	141.7	192.2
2002	26.0	39.3	51.9	65.7	78.2	95.8	115.6	144.6	194.4
2001	27.0	40.9	53.0	65.7	78.7	97.9	118.0	142.1	191.6
2000	27.1	41.3	54.8	67.8	81.1	99.6	120.6	145.0	208.6
1999	26.7	41.6	55.3	67.3	81.8	100.8	119.4	146.9	202.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Non-Hispanic White, both sexes

2007 5/.....	10.0	15.8	22.5	30.7	40.0	47.0	61.8	84.3	120.8
2006 5/.....	10.7	16.1	23.9	32.6	38.9	48.1	65.0	85.1	119.8
2005 5/.....	10.7	16.7	24.0	32.9	39.0	50.9	65.2	85.2	126.3
2004 5/.....	10.9	17.5	25.6	32.8	41.5	50.8	66.7	85.3	126.8
2003 5/.....	11.8	18.1	25.6	33.4	40.1	53.2	67.2	89.6	134.7
2002.....	12.0	18.8	25.7	34.1	41.8	53.4	69.1	90.3	137.7
2001.....	12.6	19.8	26.2	33.9	41.9	54.9	70.8	89.9	137.1
2000.....	12.5	20.1	27.2	35.3	44.0	56.8	72.3	92.9	149.6
1999.....	12.3	20.1	27.4	34.6	43.8	56.8	71.0	94.0	147.8

Non-Hispanic White, male

2007 5/.....	*	*	*	0.6	0.8	1.4	1.6	2.5	3.6
2006 5/.....	*	*	0.4	0.8	0.8	1.2	1.8	2.6	3.3
2005 5/.....	*	0.3	0.4	0.7	0.9	1.7	1.7	2.7	2.6
2004 5/.....	*	*	0.4	0.6	1.0	1.1	2.3	2.0	2.9
2003 5/.....	*	0.4	0.4	0.8	1.0	1.2	1.6	3.0	3.9
2002.....	*	*	0.4	0.6	0.6	1.3	1.8	2.7	3.4
2001.....	*	0.3	0.5	0.8	1.1	1.7	2.1	2.4	3.5
2000.....	*	0.3	0.6	0.8	1.1	1.7	2.2	2.2	4.0
1999.....	*	0.3	0.5	0.6	1.0	1.2	1.8	2.3	4.5

Non-Hispanic White, female

2007 5/.....	19.9	31.0	43.9	59.1	75.3	85.2	107.2	136.5	176.3
2006 5/.....	21.2	31.7	46.6	62.3	72.9	87.3	112.2	137.0	174.1
2005 5/.....	21.3	32.8	46.8	63.0	73.2	91.7	112.3	136.7	182.9
2004 5/.....	21.6	34.4	49.9	62.7	77.6	91.9	113.8	136.8	182.4
2003 5/.....	23.3	35.5	50.0	63.7	74.8	96.0	114.6	142.8	192.3
2002.....	23.7	37.0	50.0	65.2	78.3	96.0	117.2	143.9	195.8
2001.....	24.9	38.8	50.8	64.6	78.0	98.1	119.4	142.8	192.6
2000.....	24.7	39.4	52.6	67.2	81.7	101.1	121.5	146.9	208.3
1999.....	24.3	39.5	53.0	66.0	81.3	101.1	119.5	148.1	204.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Non-Hispanic Black, both sexes

2007 5/.....	21.2	28.8	40.4	47.6	56.2	59.8	79.4	101.5	138.8
2006 5/.....	22.0	28.3	40.1	48.1	50.5	59.8	79.5	106.7	144.7
2005 5/.....	23.6	32.8	44.5	48.8	54.8	62.3	76.4	92.9	146.1
2004 5/.....	21.7	32.7	41.0	49.7	52.1	63.9	77.3	91.4	145.6
2003 5/.....	23.1	32.0	44.2	48.7	55.3	66.2	81.1	106.5	163.3
2002.....	23.4	32.9	41.2	47.0	55.4	68.2	79.9	118.8	154.0
2001.....	23.2	33.7	43.4	48.5	58.9	71.7	83.8	107.3	153.9
2000.....	24.3	33.7	44.9	48.2	53.5	66.5	85.5	101.4	178.7
1999.....	24.6	35.2	45.2	49.6	59.0	71.1	89.0	106.3	149.1

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	39.3	52.8	72.6	84.4	95.9	100.1	125.3	153.3	194.6
2006 5/.....	40.8	51.4	72.4	84.6	86.6	99.2	125.9	160.7	202.5
2005 5/.....	43.9	60.2	80.7	85.7	93.4	102.7	118.6	141.5	204.1
2004 5/.....	40.5	59.7	74.3	87.2	89.7	105.5	122.0	137.5	199.6
2003 5/.....	42.9	58.7	79.9	85.6	95.2	109.3	128.5	160.7	224.5
2002.....	43.4	60.1	74.3	82.3	94.3	112.9	125.3	178.6	210.7
2001.....	42.9	61.5	78.3	85.6	101.0	117.2	131.3	160.3	208.3
2000.....	45.0	61.9	80.5	84.9	92.4	108.9	135.6	150.4	241.6
1999.....	45.6	64.2	81.0	86.9	101.6	116.0	140.9	156.7	201.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes

2007.....	1.8	2.4	2.7	2.8	3.1	3.2	3.4	4.5	4.7
2006.....	2.1	2.4	2.5	2.7	3.1	3.4	3.6	4.1	5.2
2005.....	2.2	2.3	2.6	2.6	3.0	3.1	3.5	4.6	5.4
2004.....	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
2003.....	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	3.5	4.7	5.2	5.3	5.9	5.8	5.9	7.3	7.0
2006.....	4.1	4.7	4.9	5.2	5.7	6.1	6.2	6.6	7.7
2005.....	4.3	4.6	5.1	5.0	5.7	5.7	6.0	7.5	7.8
2004.....	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
2003.....	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Hispanic, both sexes

2007.....	2.3	3.1	4.0	3.9	3.2	4.0	5.0	*	*
2006.....	2.3	3.6	3.3	3.2	3.9	5.0	4.8	9.5	*
2005.....	2.4	3.4	3.6	4.1	4.3	5.6	*	*	*
2004.....	2.2	2.6	3.3	5.6	3.9	4.7	4.8	*	*
2003.....	2.5	3.3	3.5	2.6	3.1	4.1	5.5	8.2	*
2002.....	2.0	2.8	3.5	3.7	3.7	5.9	6.4	*	*
2001.....	3.1	2.8	3.9	4.3	*	4.6	5.9	*	*
2000.....	2.6	2.9	4.2	5.1	5.0	6.3	7.3	*	13.9
1999.....	3.3	3.4	3.7	4.1	4.7	5.2	6.5	*	*

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	4.8	6.2	7.8	7.4	5.9	7.2	8.6	*	*
2006.....	4.6	7.1	6.5	6.0	7.1	8.9	8.2	15.8	*
2005.....	4.9	6.7	6.9	7.7	7.9	9.9	*	*	*
2004.....	4.5	5.1	6.4	10.6	7.1	8.3	8.2	*	*
2003.....	5.0	6.6	6.7	4.9	5.6	7.2	9.5	13.5	*
2002.....	4.1	5.6	6.7	6.9	6.6	10.5	11.0	*	*
2001.....	6.3	5.6	7.4	8.0	*	8.0	10.2	*	*
2000.....	5.3	5.7	7.9	9.4	9.1	11.0	12.5	*	20.8
1999.....	6.5	6.7	7.1	7.7	8.4	9.2	11.0	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 4/, both sexes

2007.....	1.7	2.3	2.6	2.7	3.1	3.1	3.3	4.4	4.7
2006.....	2.0	2.3	2.5	2.7	3.0	3.2	3.5	3.7	5.2
2005.....	2.2	2.2	2.5	2.5	2.9	2.9	3.5	4.6	5.3
2004.....	2.0	2.0	2.6	2.7	3.1	3.2	3.2	4.2	5.0
2003.....	2.0	2.3	2.8	2.9	3.2	2.9	3.6	5.0	5.7
2002.....	2.1	2.4	2.6	2.7	3.3	3.5	4.0	4.0	6.1
2001.....	2.2	2.6	2.5	3.1	3.0	3.4	3.9	5.4	7.0
2000.....	2.2	2.6	3.0	2.9	3.5	3.7	4.1	5.5	7.1
1999.....	2.3	2.6	2.9	3.5	3.6	3.8	4.6	4.5	7.9

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	3.4	4.6	5.0	5.1	5.9	5.7	5.7	7.2	6.9
2006.....	4.0	4.5	4.8	5.1	5.6	5.9	6.0	6.1	7.6
2005.....	4.3	4.3	4.9	4.8	5.5	5.3	6.0	7.4	7.7
2004.....	4.0	3.9	5.1	5.2	5.9	5.8	5.4	6.8	7.2
2003.....	3.8	4.4	5.4	5.5	5.9	5.2	6.1	8.0	8.2
2002.....	4.0	4.7	5.0	5.2	6.1	6.2	6.8	6.5	8.7
2001.....	4.3	5.0	4.9	5.9	5.5	6.1	6.7	8.6	10.0
2000.....	4.3	5.1	5.8	5.5	6.5	6.7	7.0	8.7	9.9
1999.....	4.5	5.1	5.7	6.6	6.7	6.8	7.8	7.1	11.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic White, both sexes

2007 5/.....	1.5	2.1	2.3	2.3	2.7	2.7	2.7	4.0	3.9
2006 5/.....	1.9	2.0	2.3	2.4	2.5	2.6	2.9	3.3	4.1
2005 5/.....	1.8	2.0	2.1	2.1	2.5	2.6	2.9	3.9	4.7
2004 5/.....	1.8	1.7	2.4	2.3	2.6	2.6	2.7	3.6	3.9
2003 5/.....	1.8	2.0	2.5	2.5	2.7	2.3	2.7	4.4	4.8
2002.....	1.8	2.1	2.4	2.2	2.7	2.7	3.5	3.4	4.9
2001.....	2.0	2.3	2.2	2.7	2.4	2.9	3.3	4.5	6.1
2000.....	1.9	2.3	2.6	2.4	2.8	3.1	3.6	4.8	6.0
1999.....	2.1	2.4	2.5	2.9	3.0	2.9	3.9	4.0	6.7

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	3.0	4.1	4.5	4.5	5.0	5.0	4.7	6.5	5.8
2006 5/.....	3.7	4.0	4.6	4.6	4.8	4.8	5.1	5.3	6.1
2005 5/.....	3.7	3.9	4.2	4.1	4.8	4.7	5.1	6.4	6.8
2004 5/.....	3.6	3.3	4.7	4.5	4.9	4.8	4.7	5.8	5.7
2003 5/.....	3.5	3.9	4.9	4.9	5.1	4.2	4.6	7.1	7.0
2002.....	3.6	4.2	4.6	4.3	5.0	5.0	5.9	5.4	7.0
2001.....	4.0	4.5	4.2	5.2	4.4	5.3	5.6	7.3	8.6
2000.....	3.8	4.5	5.1	4.6	5.2	5.5	6.1	7.6	8.4
1999.....	4.2	4.7	5.0	5.5	5.7	5.3	6.6	6.3	9.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic Black, both sexes

2007 5/.....	3.0	4.3	4.2	5.7	7.2	6.6	9.4	9.6	13.0
2006 5/.....	3.2	4.1	3.4	5.0	6.9	8.8	8.4	9.2	17.5
2005 5/.....	4.2	3.4	5.2	5.3	6.7	6.3	9.6	12.4	13.4
2004 5/.....	3.6	4.0	4.8	6.3	7.3	8.2	8.1	10.4	16.4
2003 5/.....	3.6	3.9	5.2	6.3	7.2	7.9	11.5	11.7	14.9
2002.....	3.9	4.6	4.3	6.2	8.4	10.3	9.5	12.5	17.5
2001.....	3.3	4.5	4.7	6.1	8.0	7.8	9.7	15.5	19.0
2000.....	3.8	5.0	6.2	6.6	9.6	10.0	10.9	13.8	19.7
1999.....	3.1	4.2	6.3	8.3	8.7	13.3	12.4	9.7	21.7

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	5.5	7.9	7.6	10.1	12.5	11.2	15.1	14.6	18.4
2006 5/.....	6.0	7.6	6.2	8.9	11.9	14.8	13.5	14.1	24.7
2005 5/.....	7.8	6.3	9.5	9.5	11.6	10.7	15.3	19.0	18.8
2004 5/.....	6.8	7.4	8.7	11.2	12.8	13.7	13.0	15.8	22.9
2003 5/.....	6.7	7.2	9.5	11.1	12.6	13.2	18.3	17.8	20.6
2002.....	7.2	8.4	7.8	11.0	14.6	17.2	15.1	19.1	24.2
2001.....	6.1	8.3	8.6	10.8	13.8	13.1	15.6	23.3	26.1
2000.....	7.0	9.3	11.4	11.7	16.7	16.6	17.4	20.7	27.0
1999.....	5.7	7.7	11.4	14.9	15.2	22.1	19.8	14.5	29.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins 3/, both sexes

2007.....	0.9	1.8	3.7	6.3	9.7	11.4	13.5	17.0	20.1
2006.....	1.0	1.9	3.8	6.2	8.8	11.6	14.3	17.6	21.6
2005.....	0.9	1.8	3.7	6.4	8.7	10.8	13.1	17.4	22.1
2004.....	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
2003.....	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	1.8	3.5	7.1	12.1	18.1	20.8	23.5	27.8	29.6
2006.....	1.9	3.6	7.4	11.9	16.4	21.0	24.7	28.6	31.6
2005.....	1.8	3.5	7.2	12.1	16.4	19.6	22.6	28.2	32.2
2004.....	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
2003.....	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Hispanic, both sexes

2007.....	0.9	1.3	2.6	5.0	7.0	10.0	8.6	8.1	8.2
2006.....	1.0	1.4	2.9	4.4	5.7	7.7	11.5	11.8	15.3
2005.....	1.0	2.2	2.4	4.8	7.2	8.5	10.2	14.7	11.1
2004.....	*	*	2.3	5.2	8.9	6.5	11.0	9.8	9.5
2003.....	*	1.7	2.5	3.6	6.3	7.8	8.8	11.5	20.7
2002.....	*	1.3	3.3	3.1	8.2	9.9	10.7	11.6	20.5
2001.....	*	2.2	3.0	4.9	6.2	9.1	9.6	15.4	18.8
2000.....	*	1.8	3.2	4.1	5.8	8.4	10.7	*	19.9
1999.....	*	1.8	3.1	4.5	7.8	9.0	9.7	12.3	15.4

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	1.9	2.5	5.1	9.4	13.0	17.8	14.8	13.4	12.7
2006.....	2.1	2.9	5.6	8.3	10.4	13.7	19.7	19.6	23.4
2005.....	2.1	4.3	4.7	9.1	13.3	15.1	17.5	24.4	16.9
2004.....	*	*	4.5	9.8	16.3	11.5	18.8	16.2	14.4
2003.....	*	3.3	4.8	6.6	11.5	13.8	15.1	18.9	31.3
2002.....	*	2.6	6.4	5.8	14.9	17.6	18.3	19.0	30.7
2001.....	*	4.2	5.8	9.2	11.2	16.1	16.5	24.9	27.9
2000.....	*	3.4	6.2	7.7	10.6	14.7	18.3	*	29.7
1999.....	*	3.5	5.8	8.5	14.1	15.7	16.5	19.7	22.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 4/, both sexes

2007.....	0.9	1.8	3.8	6.4	9.8	11.5	13.8	17.5	20.7
2006.....	1.0	1.9	3.9	6.4	9.0	11.9	14.4	17.9	21.9
2005.....	0.9	1.8	3.8	6.4	8.8	11.0	13.3	17.5	22.6
2004.....	0.9	1.9	3.5	5.8	8.8	11.8	14.0	18.1	22.8
2003.....	0.8	1.9	3.6	6.3	9.2	11.5	13.9	17.3	22.9
2002.....	0.9	2.1	3.5	5.9	8.4	11.9	14.3	17.2	24.6
2001.....	0.9	2.0	3.2	5.9	8.4	11.5	14.7	19.3	24.8
2000.....	1.0	1.8	3.6	6.1	8.6	11.2	14.2	18.8	24.9
1999.....	1.0	1.7	3.3	5.6	8.8	11.0	14.6	19.6	24.3

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	1.8	3.6	7.3	12.4	18.5	21.0	24.0	28.6	30.5
2006.....	1.9	3.7	7.5	12.2	16.8	21.6	25.0	29.0	32.0
2005.....	1.8	3.4	7.4	12.3	16.6	19.9	22.9	28.4	32.9
2004.....	1.8	3.7	6.8	11.1	16.4	21.4	24.0	29.3	33.0
2003.....	1.5	3.8	7.0	12.1	17.2	20.8	23.9	27.9	33.0
2002.....	1.7	4.0	6.8	11.3	15.8	21.4	24.4	27.7	35.3
2001.....	1.8	3.9	6.2	11.2	15.6	20.6	25.0	30.9	35.1
2000.....	1.9	3.5	6.9	11.6	15.9	20.0	24.0	29.9	34.9
1999.....	1.9	3.3	6.3	10.7	16.4	19.7	24.8	31.1	34.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic White, both sexes

2007 5/.....	0.9	1.8	3.6	5.8	8.4	10.2	12.6	16.5	20.6
2006 5/.....	0.9	1.8	3.5	5.6	7.7	10.5	13.4	17.2	22.0
2005 5/.....	0.9	1.7	3.5	5.5	8.0	9.9	12.1	17.1	22.2
2004 5/.....	0.9	1.7	3.3	5.1	7.5	10.5	13.1	17.8	23.0
2003 5/.....	0.7	1.9	3.5	5.5	8.0	10.7	13.2	16.4	22.8
2002.....	0.8	1.9	3.1	5.4	7.1	10.8	13.2	16.5	23.8
2001.....	0.9	1.9	3.1	5.2	7.3	10.5	13.9	18.7	24.5
2000.....	1.0	1.7	3.4	5.4	7.4	9.9	13.1	17.7	24.5
1999.....	0.9	1.6	3.0	5.2	7.6	10.3	13.9	19.0	23.6

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	1.7	3.5	7.0	11.3	15.8	18.7	22.1	27.0	30.3
2006 5/.....	1.7	3.7	6.9	10.8	14.6	19.3	23.5	28.0	32.2
2005 5/.....	1.7	3.3	6.9	10.7	15.1	18.2	21.1	27.7	32.4
2004 5/.....	1.9	3.3	6.5	9.9	14.3	19.1	22.7	28.8	33.3
2003 5/.....	1.5	3.7	6.8	10.7	15.1	19.4	22.7	26.5	32.8
2002.....	1.6	3.8	6.2	10.3	13.4	19.7	22.6	26.6	34.1
2001.....	1.7	3.7	6.1	10.0	13.7	19.0	23.8	30.0	34.7
2000.....	1.9	3.4	6.7	10.5	13.8	17.9	22.3	28.2	34.3
1999.....	1.9	3.2	5.9	10.0	14.2	18.5	23.6	30.3	33.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic Black, both sexes

2007 5/.....	1.2	2.6	5.6	13.2	24.9	25.4	29.3	32.5	27.4
2006 5/.....	1.5	2.1	7.1	14.3	21.0	25.9	27.9	31.0	26.3
2005 5/.....	1.4	2.5	6.3	15.3	18.4	23.2	27.6	27.3	32.0
2004 5/.....	1.0	3.1	5.2	12.1	20.6	27.0	27.1	25.8	26.5
2003 5/.....	1.0	2.6	5.3	14.3	20.4	22.8	25.4	31.9	30.3
2002.....	1.1	3.0	6.8	11.0	20.4	25.5	28.1	28.8	39.6
2001.....	1.4	3.0	4.6	11.7	19.1	24.6	26.6	31.7	32.8
2000.....	1.0	2.4	5.3	12.3	20.0	26.5	28.3	35.6	33.7
1999.....	*	2.3	5.9	10.8	21.1	22.0	25.8	30.2	35.9

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	2.3	4.7	10.2	23.6	43.2	42.9	46.9	49.5	38.7
2006 5/.....	2.7	3.9	13.0	25.4	36.5	43.7	44.5	47.4	37.0
2005 5/.....	2.6	4.6	11.6	27.3	32.1	39.1	44.0	41.7	44.8
2004 5/.....	1.9	5.7	9.5	21.6	35.9	45.3	43.1	39.4	36.9
2003 5/.....	1.9	4.8	9.6	25.4	35.5	38.2	40.5	48.6	42.1
2002.....	2.0	5.5	12.3	19.6	35.5	42.7	44.9	43.8	54.8
2001.....	2.7	5.6	8.3	20.9	33.3	41.1	42.4	47.9	44.9
2000.....	2.0	4.5	9.7	22.0	34.8	44.0	45.3	53.6	46.2
1999.....	*	4.2	10.8	19.3	36.8	36.5	41.4	45.2	49.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

All origins 3/, both sexes

2007.....	2.4	4.7	7.5	11.2	16.3	21.6	26.6	35.0	37.6
2006.....	2.8	5.1	7.7	12.4	16.3	22.0	28.3	34.8	37.8
2005.....	2.8	5.3	8.4	12.2	16.5	22.3	28.0	34.7	38.1
2004.....	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
2003.....	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6

All origins 3/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All origins 3/, female

2007.....	4.8	9.1	14.6	21.6	30.6	39.3	46.3	57.1	55.4
2006.....	5.5	9.9	14.9	23.7	30.6	39.9	49.1	56.5	55.5
2005.....	5.5	10.3	16.3	23.4	30.9	40.4	48.3	56.3	55.6
2004.....	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
2003.....	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Hispanic, both sexes

2007.....	2.2	3.3	4.7	8.2	11.9	12.4	21.5	21.7	22.9
2006.....	1.9	3.5	4.1	8.8	10.2	16.2	24.9	22.6	27.8
2005.....	2.0	4.5	5.4	9.5	11.5	15.1	16.7	24.8	27.4
2004.....	2.5	3.1	5.5	7.9	11.2	15.1	19.4	23.3	28.1
2003.....	2.4	3.2	5.8	7.3	10.8	15.0	20.9	23.8	28.5
2002.....	2.2	4.5	4.1	7.6	10.8	14.1	24.0	24.5	23.8
2001.....	2.0	3.5	5.4	9.3	14.1	14.8	16.4	26.3	30.6
2000.....	2.4	3.8	5.7	7.5	12.5	16.6	24.5	25.6	32.5
1999.....	1.7	4.0	4.5	8.1	8.8	16.4	19.0	24.1	23.9

Hispanic, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2007.....	4.5	6.7	9.1	15.5	21.9	22.1	37.1	36.2	35.3
2006.....	3.8	7.0	8.0	16.7	18.7	28.8	42.9	37.6	42.5
2005.....	4.0	9.1	10.5	18.0	21.2	26.8	28.7	41.2	41.7
2004.....	5.1	6.1	10.7	14.8	20.4	26.8	33.1	38.6	42.5
2003.....	4.8	6.3	11.1	13.7	19.7	26.6	35.8	39.2	43.0
2002.....	4.4	8.8	7.9	14.3	19.6	25.0	41.1	40.1	35.6
2001.....	4.1	6.9	10.3	17.4	25.5	26.2	28.1	42.6	45.6
2000.....	4.7	7.5	10.9	13.9	22.6	29.1	41.8	41.2	48.5
1999.....	3.4	7.7	8.6	15.1	15.9	28.7	32.5	38.4	35.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Non-Hispanic 4/, both sexes

2007.....	2.4	4.8	7.8	11.5	16.7	22.3	27.0	35.8	38.3
2006.....	2.9	5.2	8.0	12.7	16.8	22.4	28.5	35.4	38.3
2005.....	2.9	5.3	8.7	12.4	16.9	22.8	28.7	35.2	38.6
2004.....	3.0	5.4	8.1	12.2	18.3	23.0	29.5	37.0	38.1
2003.....	2.9	5.4	8.5	13.4	17.8	22.3	29.9	37.5	38.8
2002.....	3.3	5.6	8.7	13.3	17.7	22.8	30.6	38.4	38.7
2001.....	3.0	5.7	8.5	13.2	17.8	23.0	30.9	37.8	39.8
2000.....	3.1	5.6	8.5	12.6	17.8	22.1	31.2	36.8	41.2
1999.....	3.3	5.6	8.5	12.9	17.5	22.6	29.5	35.7	38.1

Non-Hispanic 4/, male

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2007.....	4.8	9.4	15.1	22.1	31.3	40.7	46.9	58.3	56.4
2006.....	5.6	10.2	15.5	24.3	31.5	40.7	49.5	57.4	56.0
2005.....	5.7	10.5	16.8	23.8	31.6	41.4	49.5	57.0	56.2
2004.....	5.9	10.5	15.7	23.2	34.3	41.6	50.6	59.8	55.1
2003.....	5.8	10.6	16.5	25.7	33.2	40.3	51.2	60.4	55.9
2002.....	6.5	10.9	16.8	25.4	33.1	41.1	52.2	61.6	55.3
2001.....	5.9	11.1	16.4	25.2	33.3	41.4	52.4	60.4	56.4
2000.....	6.1	11.0	16.5	24.0	33.1	39.5	52.9	58.4	57.8
1999.....	6.4	11.0	16.5	24.5	32.6	40.3	50.0	56.6	53.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Non-Hispanic White, both sexes

2007 5/.....	2.6	4.9	8.1	11.9	17.5	23.2	28.0	37.0	39.7
2006 5/.....	3.0	5.4	8.3	13.1	17.5	23.7	29.3	36.7	39.3
2005 5/.....	3.0	5.6	9.0	12.9	17.4	23.3	29.6	36.1	40.0
2004 5/.....	3.0	5.5	8.2	12.7	18.8	23.6	30.4	38.5	39.1
2003 5/.....	3.0	5.6	8.9	13.9	18.4	22.8	30.8	38.7	39.5
2002.....	3.4	5.9	9.0	13.9	18.5	23.5	31.4	39.3	39.6
2001.....	3.2	5.9	8.7	13.4	18.3	23.8	31.8	39.0	40.6
2000.....	3.3	5.8	8.9	13.1	18.3	22.7	32.3	37.9	42.2
1999.....	3.4	5.8	8.9	13.4	18.1	23.1	30.0	36.7	39.2

Non-Hispanic White, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2007 5/.....	5.1	9.8	15.9	23.1	33.3	42.7	49.1	60.6	58.5
2006 5/.....	5.9	10.6	16.3	25.5	33.1	43.4	51.1	59.8	57.6
2005 5/.....	5.9	11.1	17.8	25.0	32.9	42.7	51.5	58.6	58.2
2004 5/.....	6.0	10.8	16.2	24.5	35.5	43.1	52.6	62.4	56.5
2003 5/.....	6.0	11.0	17.5	26.9	34.8	41.6	53.0	62.6	56.8
2002.....	6.8	11.6	17.6	26.9	34.8	42.6	53.9	63.4	56.7
2001.....	6.4	11.8	17.1	25.9	34.4	43.2	54.3	62.5	57.5
2000.....	6.6	11.4	17.5	25.2	34.4	41.0	54.9	60.4	59.2
1999.....	6.8	11.4	17.3	25.8	33.9	41.5	51.0	58.3	54.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Non-Hispanic Black, both sexes

2007 5/.....	2.0	4.2	6.8	10.8	14.4	19.7	23.8	28.4	27.9
2006 5/.....	2.7	4.5	6.6	11.5	15.8	16.5	25.8	25.2	33.7
2005 5/.....	2.3	4.4	7.1	11.2	15.9	22.6	25.7	31.8	31.1
2004 5/.....	3.1	5.1	7.6	10.4	18.7	22.8	26.9	29.6	32.5
2003 5/.....	2.8	5.2	6.9	11.3	15.4	20.9	26.8	29.3	37.2
2002.....	2.9	4.2	7.8	11.1	15.1	20.3	26.9	33.1	35.4
2001.....	2.0	4.5	8.1	12.8	16.6	20.8	26.4	28.4	36.5
2000.....	2.2	4.8	6.6	11.2	16.9	19.9	26.6	27.9	34.7
1999.....	2.4	5.2	7.4	11.2	15.3	22.9	28.9	28.5	29.1

Non-Hispanic Black, male

2007 5/.....
2006 5/.....
2005 5/.....
2004 5/.....
2003 5/.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2007 5/.....	3.7	7.8	12.4	19.2	25.0	33.2	38.1	43.3	39.4
2006 5/.....	5.1	8.3	12.1	20.4	27.4	27.8	41.2	38.5	47.5
2005 5/.....	4.3	8.0	12.9	20.0	27.8	38.0	41.0	48.6	43.7
2004 5/.....	5.9	9.4	13.9	18.6	32.6	38.3	42.8	45.2	45.4
2003 5/.....	5.3	9.6	12.6	20.1	26.8	35.0	42.7	44.6	51.6
2002.....	5.4	7.8	14.2	19.7	26.4	34.0	42.9	50.3	48.9
2001.....	3.7	8.3	14.8	22.8	28.9	34.6	42.2	42.9	50.0
2000.....	4.1	9.0	11.9	20.0	29.3	33.1	42.7	41.9	47.5
1999.....	4.5	9.6	13.4	20.0	26.6	38.1	46.3	42.6	40.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins 3/, both sexes

2007.....	2.1	3.7	6.3	9.9	14.7	18.4	24.4	30.7	33.0
2006.....	2.0	3.7	6.5	9.6	14.8	18.2	24.0	28.9	34.3
2005.....	2.0	4.1	6.4	10.8	14.4	19.9	24.3	30.3	32.6
2004.....	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
2003.....	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4

All origins 3/, male

2007.....	2.9	5.4	9.2	14.6	21.2	25.4	35.1	45.1	50.9
2006.....	2.8	5.4	9.5	14.0	21.6	25.0	34.5	40.8	52.1
2005.....	2.9	5.8	9.3	15.7	20.7	28.9	34.1	43.3	50.6
2004.....	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
2003.....	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4

All origins 3/, female

2007.....	1.2	2.0	3.7	5.5	9.0	12.6	16.6	21.6	24.5
2006.....	1.2	2.1	3.6	5.6	8.8	12.6	16.4	21.5	25.9
2005.....	1.2	2.4	3.6	6.4	8.8	12.5	17.2	22.2	24.3
2004.....	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
2003.....	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Hispanic, both sexes

2007	2.3	3.2	6.2	8.6	12.0	16.7	24.0	33.5	25.4
2006	1.5	3.4	6.5	7.4	10.6	16.5	20.5	24.5	23.5
2005	1.9	4.0	4.8	8.4	13.8	18.1	21.7	26.2	32.9
2004	2.3	4.2	6.1	11.2	13.3	15.0	19.4	29.6	28.1
2003	1.7	2.7	6.3	10.8	12.9	20.4	24.2	22.5	26.9
2002	2.1	3.8	6.6	10.2	11.9	14.1	23.5	25.8	26.1
2001	2.6	4.3	4.6	10.0	12.2	17.0	24.4	21.4	31.9
2000	2.4	3.4	6.2	10.0	10.8	14.0	21.4	30.6	25.9
1999	1.9	4.1	6.7	9.4	13.3	14.6	21.0	25.3	28.8

Hispanic, male

2007	3.4	5.1	8.6	13.6	16.7	24.9	38.7	52.7	38.6
2006	2.1	4.9	8.9	10.8	14.7	26.7	30.9	34.4	32.7
2005	2.9	4.0	7.4	12.2	19.5	26.6	29.7	38.5	46.1
2004	3.4	5.9	8.3	15.4	23.8	22.8	29.4	43.4	43.4
2003	2.6	3.2	8.2	15.0	18.8	26.9	34.6	28.1	38.5
2002	2.6	5.8	9.9	13.3	16.9	20.6	41.0	35.5	40.2
2001	3.8	6.0	6.6	14.4	17.7	26.1	35.5	35.1	57.1
2000	3.4	4.5	10.1	13.2	14.9	21.9	34.7	45.6	40.3
1999	3.3	5.6	8.8	15.8	18.9	23.9	29.6	*	*

Hispanic, female

2007	*	*	3.9	4.2	8.0	10.3	13.3	20.7	18.2
2006	*	*	4.3	4.4	7.1	8.6	13.0	18.0	18.6
2005	*	3.9	*	5.0	9.1	11.4	15.9	18.0	26.1
2004	*	2.5	4.0	7.5	*	8.9	12.3	20.6	20.2
2003	*	*	4.4	7.1	8.0	15.4	16.8	18.9	21.1
2002	*	*	3.5	7.4	7.7	9.1	11.0	19.7	19.1
2001	*	2.6	*	6.1	7.7	10.1	16.5	*	19.5
2000	*	*	*	7.2	7.5	8.1	12.0	21.5	*
1999	*	*	4.8	*	8.8	7.7	14.9	22.5	23.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 4/, both sexes

2007.....	2.0	3.7	6.3	10.0	14.9	18.5	24.4	30.5	33.4
2006.....	2.1	3.7	6.4	9.8	15.1	18.3	24.2	29.2	34.8
2005.....	2.0	4.1	6.5	11.0	14.3	20.0	24.5	30.5	32.5
2004.....	2.1	4.0	6.8	11.3	14.6	19.4	24.8	29.7	32.9
2003.....	2.0	4.1	7.2	11.4	14.6	19.4	24.8	30.3	33.4
2002.....	2.4	4.3	7.2	10.9	16.3	19.9	24.6	28.8	32.7
2001.....	2.4	4.4	7.3	12.0	14.8	21.0	24.4	29.3	32.3
2000.....	2.2	4.6	7.7	10.9	15.2	19.5	24.6	29.5	33.9
1999.....	2.2	4.3	7.4	10.9	14.4	19.4	23.1	26.9	31.4

Non-Hispanic 4/, male

2007.....	2.9	5.4	9.2	14.6	21.5	25.4	34.7	44.6	51.6
2006.....	2.9	5.4	9.5	14.3	22.1	24.8	34.6	41.1	53.2
2005.....	2.8	6.0	9.4	15.9	20.6	29.1	34.4	43.5	50.7
2004.....	2.9	5.9	10.0	16.7	20.4	28.1	34.5	44.0	47.1
2003.....	3.1	6.0	10.6	17.3	20.3	27.6	36.0	45.6	49.0
2002.....	3.5	6.5	10.6	16.2	23.3	28.0	36.3	40.0	50.4
2001.....	3.4	6.5	10.8	17.4	21.2	30.1	34.3	42.4	52.6
2000.....	3.1	6.8	11.2	15.8	21.7	27.5	34.5	44.1	55.4
1999.....	3.2	6.1	10.8	15.8	20.4	27.5	33.8	39.2	49.8

Non-Hispanic 4/, female

2007.....	1.2	2.1	3.7	5.6	9.1	12.8	16.8	21.7	24.8
2006.....	1.2	2.1	3.5	5.7	8.9	12.9	16.6	21.7	26.2
2005.....	1.3	2.3	3.7	6.5	8.8	12.6	17.2	22.4	24.2
2004.....	1.3	2.2	3.8	6.3	9.6	12.3	17.9	21.0	26.5
2003.....	1.1	2.3	3.9	6.0	9.6	12.8	16.9	21.0	26.6
2002.....	1.5	2.3	4.0	6.1	10.1	13.5	16.4	22.0	25.1
2001.....	1.4	2.4	4.0	7.1	9.2	13.7	17.4	21.5	23.8
2000.....	1.4	2.4	4.4	6.4	9.6	13.3	17.7	20.9	25.2
1999.....	1.2	2.5	4.2	6.4	9.4	13.1	15.6	19.7	24.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic White, both sexes

2007 5/.....	2.0	3.7	6.4	10.1	15.3	19.1	24.8	31.2	33.9
2006 5/.....	2.1	3.7	6.7	10.2	15.6	18.8	25.1	29.9	35.5
2005 5/.....	2.0	4.2	6.7	11.2	14.8	20.7	25.0	31.3	33.0
2004 5/.....	2.2	4.1	7.1	11.7	15.1	19.7	25.2	30.5	33.8
2003 5/.....	2.0	4.3	7.2	11.6	15.1	20.1	25.4	31.1	34.0
2002.....	2.5	4.4	7.4	11.2	16.9	20.5	25.0	29.3	33.5
2001.....	2.4	4.4	7.5	12.2	15.4	21.5	24.7	29.7	32.8
2000.....	2.3	4.7	7.9	11.2	15.5	19.9	24.9	29.9	34.6
1999.....	2.1	4.3	7.5	11.3	14.8	19.7	23.3	27.3	31.5

Non-Hispanic White, male

2007 5/.....	2.7	5.3	9.2	14.7	21.8	25.8	35.5	45.4	53.0
2006 5/.....	2.9	5.5	9.6	14.7	22.6	25.5	35.8	42.2	54.6
2005 5/.....	2.8	6.0	9.7	15.9	21.2	29.7	35.0	44.5	52.1
2004 5/.....	3.0	6.1	10.3	17.3	20.8	28.6	34.8	45.4	48.6
2003 5/.....	3.1	6.1	10.8	17.6	20.9	28.2	36.5	46.9	50.4
2002.....	3.6	6.6	10.8	16.6	24.2	28.6	37.1	40.3	52.4
2001.....	3.6	6.5	11.1	17.3	22.0	30.8	34.3	42.9	54.3
2000.....	3.2	6.9	11.4	16.2	21.7	27.8	34.9	44.5	56.9
1999.....	3.1	6.0	10.9	16.4	20.7	27.7	34.1	40.4	50.2

Non-Hispanic White, female

2007 5/.....	1.3	2.0	3.7	5.9	9.5	13.5	16.6	22.1	24.8
2006 5/.....	1.3	2.0	3.8	6.0	9.4	13.2	17.1	22.1	26.6
2005 5/.....	1.3	2.4	3.8	6.8	9.1	13.2	17.6	23.0	24.3
2004 5/.....	1.3	2.2	3.9	6.5	10.1	12.4	18.1	21.3	27.2
2003 5/.....	0.9	2.5	3.8	6.2	9.8	13.5	17.4	21.3	26.8
2002.....	1.4	2.2	4.1	6.2	10.4	13.9	16.3	22.6	25.3
2001.....	1.3	2.4	4.0	7.5	9.6	14.0	17.9	21.7	23.8
2000.....	1.4	2.5	4.6	6.4	10.1	13.5	17.9	21.3	25.7
1999.....	1.2	2.6	4.4	6.5	9.7	13.3	15.7	19.5	24.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic Black, both sexes

2007 5/.....	2.7	4.4	7.1	11.0	14.1	16.9	26.2	28.9	30.9
2006 5/.....	2.5	4.9	6.7	8.6	13.6	16.9	19.8	26.4	33.7
2005 5/.....	2.4	4.5	6.9	12.3	13.7	17.8	22.8	26.8	30.9
2004 5/.....	2.4	4.1	6.4	10.7	13.3	19.5	26.9	25.6	28.8
2003 5/.....	2.6	4.2	8.4	11.5	13.6	16.3	22.5	26.7	34.2
2002.....	2.9	4.5	7.2	10.9	14.7	17.2	26.2	27.0	28.1
2001.....	2.6	5.1	7.5	12.5	12.0	19.4	24.3	28.7	30.0
2000.....	2.5	4.6	7.5	10.7	15.2	18.7	23.9	29.0	29.6
1999.....	3.0	5.1	7.7	9.9	13.8	19.5	23.4	24.7	31.7

Non-Hispanic Black, male

2007 5/.....	4.2	6.6	10.6	18.1	22.7	24.9	33.3	43.0	42.8
2006 5/.....	4.0	6.9	11.3	13.9	22.7	23.1	28.1	40.0	49.5
2005 5/.....	3.7	7.2	10.4	19.9	20.1	29.9	33.1	40.3	40.0
2004 5/.....	3.3	6.3	9.5	16.4	19.7	29.9	39.2	36.7	39.6
2003 5/.....	3.7	6.6	11.7	17.5	19.5	25.9	35.5	40.0	44.7
2002.....	3.6	6.6	12.1	17.3	20.5	24.3	35.2	39.0	39.4
2001.....	3.5	8.0	10.8	21.1	17.5	29.6	39.3	43.3	46.2
2000.....	3.3	7.1	11.5	15.7	25.6	28.4	32.8	48.1	49.3
1999.....	4.3	8.2	12.5	14.1	20.5	28.4	34.8	28.3	49.9

Non-Hispanic Black, female

2007 5/.....	1.4	2.5	4.2	5.5	7.8	11.4	21.9	21.5	26.0
2006 5/.....	*	3.3	3.0	4.5	6.9	12.6	14.8	19.3	27.3
2005 5/.....	*	2.2	4.0	6.4	8.9	9.6	16.7	19.7	27.2
2004 5/.....	1.6	2.3	3.8	6.3	8.5	12.4	19.6	19.7	24.5
2003 5/.....	1.5	2.2	5.7	6.8	9.2	9.8	14.7	19.7	30.1
2002.....	2.3	2.6	3.1	5.9	10.4	12.4	20.7	20.7	23.8
2001.....	1.9	2.7	4.9	5.8	7.9	12.6	15.3	21.2	24.0
2000.....	1.9	2.5	4.3	6.7	7.5	12.3	18.6	19.4	22.2
1999.....	1.9	2.4	3.8	6.5	8.9	13.6	16.6	22.8	24.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2007.....	0.9	1.8	3.5	6.3	11.4	18.5	31.7	47.1	68.6
2006.....	0.8	1.7	3.5	7.0	11.2	19.2	30.0	46.9	67.4
2005.....	0.9	1.6	3.5	6.8	11.5	18.9	30.0	46.7	68.3
2004.....	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
2003.....	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All origins 3/, male									
2007.....	1.2	2.7	5.4	10.1	18.5	30.4	54.1	84.3	132.7
2006.....	1.1	2.6	5.5	11.1	17.7	31.4	51.4	85.8	131.6
2005.....	1.2	2.2	5.3	10.8	18.7	31.2	50.4	86.3	132.1
2004.....	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
2003.....	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All origins 3/, female									
2007.....	0.6	0.9	1.8	2.9	5.1	8.9	15.3	23.6	38.1
2006.....	0.5	0.8	1.6	3.3	5.6	9.4	14.4	22.6	37.4
2005.....	0.6	1.0	1.7	3.1	5.2	8.9	15.2	22.1	39.0
2004.....	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
2003.....	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

Non-Hispanic 4/, both sexes

2007.....	0.9	1.9	3.7	6.5	11.8	19.2	32.5	48.5	70.5
2006.....	0.8	1.8	3.6	7.3	11.6	20.0	30.9	48.1	69.0
2005.....	1.0	1.7	3.6	7.0	11.8	19.6	30.8	47.7	69.6
2004.....	1.0	1.8	3.7	7.0	11.9	19.6	31.9	46.5	69.4
2003.....	0.9	1.8	3.6	7.2	12.2	19.0	30.8	45.7	65.6
2002.....	0.9	1.8	3.9	6.5	12.4	20.8	32.8	44.8	66.9
2001.....	0.8	1.8	3.6	7.3	12.4	19.5	30.0	45.5	69.6
2000.....	0.8	1.7	3.7	7.1	12.3	20.1	29.7	46.2	70.3
1999.....	0.8	1.9	3.9	7.2	12.1	20.7	30.3	44.5	69.3

Non-Hispanic 4/, male

2007.....	1.2	2.9	5.7	10.3	19.2	31.4	55.0	86.9	137.1
2006.....	1.2	2.8	5.8	11.5	18.4	32.7	52.9	88.0	135.5
2005.....	1.3	2.3	5.6	11.1	19.2	32.2	51.9	88.4	134.6
2004.....	1.4	2.9	5.8	11.2	19.2	31.7	54.9	83.7	136.5
2003.....	1.5	2.8	5.7	11.3	19.6	30.6	53.1	81.6	127.2
2002.....	1.3	2.5	6.1	10.4	19.6	33.2	55.7	78.2	129.8
2001.....	1.2	2.8	5.4	11.6	20.1	32.6	52.2	82.2	136.4
2000.....	1.2	2.6	5.7	11.1	19.9	33.0	51.5	83.5	141.3
1999.....	1.2	3.0	6.0	11.2	19.7	33.6	53.5	81.0	136.6

Non-Hispanic 4/, female

2007.....	0.6	0.9	1.9	3.0	5.4	9.1	15.9	24.3	39.1
2006.....	0.5	0.9	1.6	3.4	5.6	9.6	14.8	23.2	38.1
2005.....	0.6	1.0	1.8	3.3	5.2	9.4	15.5	22.5	40.0
2004.....	0.6	0.8	1.7	3.2	5.4	9.8	15.4	23.7	39.4
2003.....	0.3	0.9	1.7	3.4	5.7	9.7	14.8	23.8	38.6
2002.....	0.5	1.1	1.9	3.1	6.2	10.9	16.6	24.5	39.8
2001.....	0.4	0.9	1.8	3.3	5.7	9.1	14.5	23.6	41.9
2000.....	0.5	0.8	1.7	3.4	5.7	9.9	14.4	24.2	41.6
1999.....	0.3	0.9	1.8	3.6	5.6	10.6	14.1	23.2	42.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

Non-Hispanic White, both sexes

2007 5/.....	0.9	1.9	3.8	6.8	12.5	20.5	33.8	50.6	73.3
2006 5/.....	0.9	1.9	3.8	7.7	12.2	20.8	32.3	50.0	71.5
2005 5/.....	1.0	1.7	3.8	7.4	12.4	20.7	32.1	49.8	71.5
2004 5/.....	1.0	1.9	3.9	7.6	12.7	20.6	33.6	48.5	71.1
2003 5/.....	0.9	1.9	3.8	7.6	12.7	20.2	31.8	46.9	67.6
2002.....	1.0	1.9	4.0	7.0	13.0	21.6	34.0	46.0	68.0
2001.....	0.8	1.9	3.7	7.6	13.0	20.4	31.0	46.4	71.5
2000.....	0.9	1.8	3.9	7.5	12.9	20.8	30.4	47.1	72.8
1999.....	0.7	2.0	3.9	7.5	12.5	21.3	31.2	45.8	70.7

Non-Hispanic White, male

2007 5/.....	1.3	3.0	5.8	10.8	20.3	33.6	57.5	92.3	145.0
2006 5/.....	1.3	3.0	6.0	12.1	19.6	34.7	56.0	92.5	142.3
2005 5/.....	1.5	2.4	6.0	11.7	20.5	34.0	54.2	92.9	140.2
2004 5/.....	1.4	3.0	6.2	12.0	21.0	33.6	57.9	88.5	143.1
2003 5/.....	1.5	3.0	5.9	12.2	20.4	32.7	55.4	84.9	133.6
2002.....	1.4	2.7	6.5	11.1	20.7	34.9	57.9	81.1	135.2
2001.....	1.2	2.9	5.6	12.4	21.4	34.3	54.7	85.0	143.0
2000.....	1.2	2.8	6.0	11.9	21.1	34.5	53.6	85.9	148.0
1999.....	1.1	3.0	6.3	11.7	20.6	35.2	55.6	83.7	143.0

Non-Hispanic White, female

2007 5/.....	0.6	0.9	1.8	3.1	5.5	9.5	15.9	24.1	39.3
2006 5/.....	0.5	0.9	1.6	3.6	5.6	9.2	14.5	23.2	38.5
2005 5/.....	0.6	1.0	1.8	3.4	5.2	9.6	15.7	22.8	40.1
2004 5/.....	0.5	0.8	1.7	3.5	5.4	9.8	15.8	23.8	38.9
2003 5/.....	0.3	0.8	1.7	3.4	5.8	10.0	14.8	23.6	38.6
2002.....	0.5	1.0	1.7	3.1	6.1	10.7	16.9	24.5	39.0
2001.....	0.3	0.9	1.8	3.2	5.6	9.2	14.3	23.1	41.7
2000.....	0.5	0.8	1.8	3.4	5.8	9.8	14.0	24.1	42.5
1999.....	0.3	1.0	1.6	3.5	5.4	10.3	14.0	23.5	41.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

Non-Hispanic Black, both sexes

2007 5/.....	1.2	2.1	4.5	5.8	10.1	14.7	29.1	38.2	55.2
2006 5/.....	0.8	1.7	3.9	6.4	9.8	18.4	26.2	39.5	55.7
2005 5/.....	1.0	2.1	3.7	6.1	9.9	15.8	25.2	36.5	57.6
2004 5/.....	1.2	2.1	3.4	5.1	8.1	16.7	23.1	34.2	61.3
2003 5/.....	1.2	2.2	3.3	5.1	11.7	12.9	27.0	41.5	53.8
2002.....	*	1.6	4.1	5.1	11.2	19.1	27.0	37.6	64.1
2001.....	0.8	1.8	3.5	7.1	10.3	14.3	24.8	41.7	55.6
2000.....	0.9	1.3	3.0	5.5	9.7	18.0	28.8	44.5	49.6
1999.....	1.0	2.2	4.6	6.8	11.9	19.2	25.4	35.8	60.8

Non-Hispanic Black, male

2007 5/.....	*	2.8	6.3	9.0	14.9	23.0	43.8	49.9	81.1
2006 5/.....	*	2.9	5.7	10.8	12.8	25.3	33.1	57.6	95.3
2005 5/.....	*	2.8	4.8	9.4	13.2	25.0	38.1	57.6	101.8
2004 5/.....	1.6	3.1	4.5	8.1	8.5	23.6	35.4	47.7	83.3
2003 5/.....	1.8	2.8	4.6	6.7	18.7	16.5	41.6	62.6	79.9
2002.....	*	*	4.4	6.9	14.6	24.3	42.3	57.7	91.9
2001.....	*	2.9	4.3	9.4	13.0	21.0	34.1	58.9	78.5
2000.....	*	*	4.7	7.4	12.8	24.3	41.4	73.0	88.1
1999.....	*	3.8	5.6	9.6	15.7	23.9	39.2	60.2	86.8

Non-Hispanic Black, female

2007 5/.....	*	*	3.0	3.2	6.6	9.0	20.3	32.0	44.5
2006 5/.....	*	*	2.4	3.0	7.7	13.7	22.0	30.0	39.6
2005 5/.....	*	*	2.8	3.5	7.4	9.6	17.5	25.4	39.8
2004 5/.....	*	*	2.6	*	7.8	11.9	15.7	27.0	52.6
2003 5/.....	*	1.8	*	3.9	6.6	10.5	18.3	30.4	43.7
2002.....	*	*	3.8	3.7	8.7	15.6	17.9	27.2	53.5
2001.....	*	*	2.8	5.3	8.3	9.9	19.2	32.9	47.1
2000.....	*	*	*	4.0	7.5	13.8	21.2	30.2	35.3
1999.....	*	*	3.8	4.6	9.1	16.1	17.2	23.7	51.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes

2007.....	3.6	5.2	7.9	10.6	14.3	17.0	19.4	20.8	16.1
2006.....	3.5	5.7	8.1	10.8	13.0	16.8	18.5	19.6	16.7
2005.....	3.7	5.6	8.3	10.7	14.0	18.0	20.0	20.4	15.8
2004.....	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
2003.....	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All origins 3/, male

2007.....	4.4	6.3	9.7	13.3	17.5	19.8	23.8	24.6	21.1
2006.....	4.3	6.8	9.9	13.5	16.4	20.4	22.9	25.6	21.2
2005.....	4.3	7.0	10.0	13.1	17.7	21.2	25.3	25.3	19.3
2004.....	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
2003.....	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All origins 3/, female

2007.....	2.8	4.1	6.2	8.1	11.5	14.8	16.1	18.5	13.8
2006.....	2.7	4.6	6.4	8.4	10.0	13.9	15.3	15.8	14.6
2005.....	3.0	4.3	6.8	8.6	10.7	15.4	16.1	17.4	14.3
2004.....	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
2003.....	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Hispanic, both sexes

2007	2.7	3.5	5.3	6.4	7.5	10.8	12.9	16.4	11.1
2006	2.3	3.5	4.5	6.6	6.6	8.7	14.1	11.5	12.1
2005	1.9	4.2	5.3	6.8	9.1	11.6	13.5	13.3	15.4
2004	1.7	3.3	4.9	7.3	8.8	9.2	12.0	13.1	14.3
2003	2.0	3.2	4.7	7.6	8.2	11.1	16.1	13.9	*
2002	2.4	4.6	4.7	7.4	9.1	11.1	14.4	12.5	13.3
2001	2.2	3.0	4.5	7.8	9.5	11.3	13.3	17.4	*
2000	2.3	3.7	3.7	9.3	9.7	9.8	16.2	*	15.9
1999	2.2	4.0	5.2	7.2	8.1	12.5	12.3	17.0	*

Hispanic, male

2007	3.4	3.2	5.8	7.9	7.8	11.7	16.4	20.1	*
2006	2.9	3.9	6.2	5.7	7.6	11.4	17.0	*	*
2005	2.3	5.3	6.3	9.0	10.9	12.3	17.3	19.3	*
2004	1.9	4.3	5.4	9.8	9.9	10.3	16.7	*	*
2003	2.7	3.8	4.8	8.4	10.1	14.1	15.2	*	*
2002	2.9	6.4	5.0	9.0	12.2	12.7	17.3	*	*
2001	3.0	2.9	6.2	7.6	11.3	18.3	16.4	*	*
2000	2.6	*	4.4	12.1	11.9	13.1	19.9	*	*
1999	*	3.8	5.5	10.7	9.7	11.7	*	*	*

Hispanic, female

2007	1.8	3.9	4.9	5.1	7.3	10.0	10.4	14.0	11.0
2006	1.8	3.2	3.0	7.4	5.7	6.5	11.9	*	14.4
2005	*	3.1	4.4	4.8	7.6	11.1	10.9	*	15.0
2004	*	2.3	4.5	5.1	7.9	8.3	8.6	13.1	15.1
2003	*	2.6	4.6	6.9	6.7	8.9	16.8	*	*
2002	1.9	2.9	4.4	6.0	6.6	9.8	12.3	*	*
2001	*	3.2	*	8.0	8.0	*	11.1	*	*
2000	*	4.7	*	6.9	7.9	7.4	13.6	*	*
1999	*	4.1	4.8	*	6.9	13.0	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 4/, both sexes

2007.....	3.7	5.4	8.2	10.9	14.8	17.5	19.8	21.1	16.4
2006.....	3.6	5.9	8.4	11.2	13.5	17.4	18.8	20.0	16.9
2005.....	3.9	5.7	8.6	11.0	14.3	18.4	20.4	20.8	15.9
2004.....	4.0	5.7	8.6	11.2	14.3	18.2	19.4	20.8	16.7
2003.....	4.1	6.0	8.8	11.5	14.2	17.7	19.6	19.8	15.4
2002.....	4.4	6.3	8.8	11.3	15.1	18.2	19.8	18.7	16.5
2001.....	4.2	6.3	8.6	11.8	14.5	17.1	20.6	20.7	16.5
2000.....	4.2	6.5	8.6	11.5	15.1	18.8	19.5	21.3	16.9
1999.....	4.5	6.3	9.5	11.9	16.2	18.5	21.3	19.5	16.1

Non-Hispanic 4/, male

2007.....	4.5	6.7	10.1	13.7	18.3	20.4	24.3	24.8	21.7
2006.....	4.4	7.1	10.2	14.2	17.1	21.0	23.2	26.2	21.9
2005.....	4.5	7.1	10.3	13.4	18.2	21.8	25.8	25.6	19.4
2004.....	5.1	7.3	10.8	14.0	16.4	22.1	23.5	25.1	19.7
2003.....	5.3	7.7	10.1	14.1	16.8	21.8	24.5	25.7	20.1
2002.....	5.6	7.3	11.1	13.3	18.9	22.8	24.7	23.3	20.2
2001.....	5.1	7.9	11.0	14.3	17.7	21.4	25.8	26.7	20.5
2000.....	5.1	8.1	10.6	14.1	18.2	23.3	23.3	26.5	22.3
1999.....	5.5	7.7	10.9	14.9	20.1	22.9	26.5	24.6	21.3

Non-Hispanic 4/, female

2007.....	2.9	4.1	6.3	8.3	11.8	15.2	16.5	18.7	13.9
2006.....	2.8	4.7	6.7	8.5	10.3	14.4	15.5	16.2	14.6
2005.....	3.2	4.4	7.0	8.9	10.9	15.6	16.4	17.8	14.2
2004.....	3.0	4.1	6.6	8.6	12.5	15.1	16.4	18.1	15.4
2003.....	3.0	4.4	7.6	9.2	12.0	14.4	16.1	16.2	13.4
2002.....	3.3	5.3	6.7	9.5	11.8	14.5	16.4	15.9	14.9
2001.....	3.3	4.9	6.3	9.5	11.7	13.6	16.9	17.1	14.9
2000.....	3.4	4.9	6.7	9.1	12.4	15.2	16.8	18.3	14.7
1999.....	3.4	5.0	8.2	9.2	12.8	15.1	17.8	16.5	14.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic White, both sexes

2007 5/.....	4.1	5.9	8.9	12.0	16.1	19.1	21.6	22.6	17.2
2006 5/.....	4.1	6.6	9.1	12.4	14.6	18.8	20.4	21.1	17.7
2005 5/.....	4.3	6.4	9.4	12.0	15.7	19.9	21.6	21.9	16.5
2004 5/.....	4.4	6.2	9.5	12.2	15.7	19.8	20.6	22.1	17.4
2003 5/.....	4.5	6.6	9.5	12.6	15.4	19.0	20.9	20.9	15.9
2002.....	4.9	6.9	9.5	12.3	16.4	19.5	20.9	19.6	17.3
2001.....	4.6	7.0	9.2	12.9	15.6	18.2	22.0	21.8	17.2
2000.....	4.7	7.2	9.2	12.5	16.4	20.3	20.6	22.1	17.5
1999.....	5.0	6.8	10.4	12.9	17.4	20.0	22.6	20.4	16.7

Non-Hispanic White, male

2007 5/.....	5.0	7.3	10.9	15.0	19.7	22.1	26.3	26.8	23.2
2006 5/.....	4.9	7.8	10.9	15.6	18.5	22.6	25.1	27.2	22.9
2005 5/.....	5.0	7.8	11.1	14.4	19.9	23.3	26.9	26.9	20.4
2004 5/.....	5.5	7.9	11.7	15.2	18.0	24.0	24.6	26.4	21.0
2003 5/.....	5.9	8.4	10.8	15.4	17.9	23.2	25.8	26.9	21.0
2002.....	6.2	8.0	11.7	14.4	20.5	24.3	25.9	24.2	21.4
2001.....	5.6	8.6	11.8	15.4	19.1	22.8	27.2	28.0	20.9
2000.....	5.7	8.7	11.4	15.3	19.8	24.9	24.5	27.7	23.5
1999.....	6.0	8.2	11.8	16.0	21.6	24.6	27.9	25.9	22.4

Non-Hispanic White, female

2007 5/.....	3.3	4.5	7.0	9.2	12.8	16.6	18.1	19.9	14.4
2006 5/.....	3.2	5.3	7.4	9.4	11.2	15.7	16.9	17.3	15.3
2005 5/.....	3.6	5.0	7.7	9.8	11.8	17.1	17.7	18.9	14.8
2004 5/.....	3.3	4.6	7.4	9.5	13.6	16.3	17.7	19.4	15.8
2003 5/.....	3.2	5.0	8.2	10.0	13.1	15.6	17.3	17.2	13.7
2002.....	3.7	5.8	7.3	10.5	12.8	15.6	17.4	16.9	15.5
2001.....	3.6	5.4	6.7	10.6	12.4	14.5	18.3	18.0	15.7
2000.....	3.7	5.6	7.2	9.9	13.4	16.6	17.8	18.8	15.1
1999.....	3.9	5.5	9.0	10.1	13.8	16.4	18.8	17.1	14.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic Black, both sexes

2007 5/.....	2.3	3.8	5.1	5.7	9.4	9.9	7.9	10.0	8.9
2006 5/.....	1.9	3.3	5.6	5.1	7.2	9.4	7.4	12.6	9.6
2005 5/.....	2.4	3.2	5.4	5.7	7.3	10.0	13.0	11.2	9.6
2004 5/.....	2.3	3.6	4.2	6.1	7.1	9.0	10.1	10.1	10.6
2003 5/.....	2.7	3.7	5.7	5.6	8.3	9.8	10.6	10.6	10.7
2002.....	2.0	3.7	5.7	5.5	7.8	9.6	12.1	11.5	9.7
2001.....	2.7	3.5	5.5	6.3	9.2	9.6	9.2	11.0	11.9
2000.....	2.2	3.7	5.5	6.2	7.6	9.0	10.5	14.7	10.5
1999.....	2.7	4.0	5.5	6.9	8.7	7.6	12.0	11.2	10.0

Non-Hispanic Black, male

2007 5/.....	2.8	4.5	6.8	7.5	11.1	11.8	9.1	*	*
2006 5/.....	2.5	4.1	7.6	6.6	8.7	12.7	*	19.7	*
2005 5/.....	3.0	4.6	6.8	7.7	8.3	13.7	19.7	15.1	*
2004 5/.....	3.0	5.2	6.1	8.1	8.0	8.3	14.5	14.7	*
2003 5/.....	3.1	5.3	6.8	6.7	11.6	12.1	16.4	*	*
2002.....	2.2	3.3	8.3	7.7	9.0	12.0	15.0	17.9	*
2001.....	3.1	4.6	7.1	9.0	9.4	11.7	11.9	*	*
2000.....	2.2	5.7	6.6	7.9	7.7	12.6	12.0	*	*
1999.....	3.8	5.3	6.1	8.9	11.2	9.0	14.5	*	*

Non-Hispanic Black, female

2007 5/.....	1.8	3.2	3.7	4.3	8.2	8.6	7.1	9.8	9.0
2006 5/.....	1.5	2.6	4.0	4.0	6.1	7.2	7.1	8.9	9.0
2005 5/.....	2.0	2.0	4.2	4.2	6.5	7.6	9.0	9.1	10.0
2004 5/.....	1.6	2.2	2.7	4.5	6.5	9.5	7.5	7.7	11.6
2003 5/.....	2.3	2.3	4.8	4.6	5.8	8.2	7.2	8.7	12.0
2002.....	1.8	4.1	3.5	3.8	7.0	8.0	10.4	8.1	10.5
2001.....	2.3	2.7	4.2	4.3	9.0	8.3	7.6	10.4	9.8
2000.....	2.3	2.0	4.6	4.9	7.5	6.6	9.6	14.3	10.5
1999.....	1.7	2.9	5.0	5.3	6.9	6.7	10.5	12.3	9.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes

2007.....	6.2	10.4	18.1	30.3	50.0	79.6	119.0	160.8	185.5
2006.....	6.3	11.0	18.5	32.4	51.7	81.7	119.0	162.2	186.1
2005.....	6.6	11.0	19.0	33.2	52.9	83.8	121.4	161.5	189.8
2004.....	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
2003.....	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All origins 3/, male

2007.....	7.5	13.1	22.5	38.2	64.1	102.3	156.8	218.0	253.4
2006.....	7.4	14.0	23.4	41.2	65.7	103.4	157.6	219.4	244.7
2005.....	7.8	13.3	23.9	40.7	68.1	109.5	161.0	214.7	262.4
2004.....	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
2003.....	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All origins 3/, female

2007.....	5.0	7.8	14.0	23.1	37.6	61.1	91.2	124.7	153.3
2006.....	5.2	8.2	13.9	24.3	39.4	64.0	90.8	126.3	158.7
2005.....	5.5	8.7	14.4	26.4	39.6	63.0	92.7	128.4	156.4
2004.....	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
2003.....	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Hispanic, both sexes

2007	6.1	8.9	16.9	26.5	38.7	62.1	80.7	109.5	95.1
2006	5.3	9.5	14.9	24.0	41.8	63.6	82.2	110.9	100.9
2005	6.5	10.1	15.5	28.6	42.3	66.1	91.2	107.5	107.8
2004	6.0	10.0	16.5	24.6	39.4	59.7	85.6	110.7	103.2
2003	7.3	9.9	17.1	24.9	38.5	58.0	91.3	111.9	129.6
2002	6.5	9.3	18.1	27.6	47.3	61.7	89.8	119.0	132.5
2001	6.1	10.9	17.0	33.1	43.0	64.5	98.0	112.7	134.4
2000	7.8	10.4	17.9	34.0	45.7	75.2	97.0	114.7	143.3
1999	8.4	12.3	20.0	29.5	49.6	69.7	88.1	124.5	131.9

Hispanic, male

2007	7.3	10.7	17.1	31.1	46.4	75.9	95.7	134.8	133.2
2006	6.6	10.5	18.4	28.5	48.0	77.4	103.0	135.9	129.7
2005	7.8	10.8	18.3	31.7	52.5	81.0	114.0	142.7	144.5
2004	7.1	13.7	18.2	32.3	49.7	75.8	107.7	132.1	119.1
2003	8.3	11.2	19.9	30.8	46.5	71.7	107.6	126.8	149.3
2002	7.3	11.2	21.3	33.1	56.5	72.0	108.1	138.4	180.7
2001	7.5	12.7	21.6	38.8	56.3	81.9	139.8	146.7	180.9
2000	10.7	11.6	23.2	40.6	58.2	88.5	114.4	151.7	197.5
1999	10.0	14.6	23.6	36.9	56.0	85.0	106.9	139.9	135.9

Hispanic, female

2007	4.8	7.1	16.7	22.3	32.1	51.4	69.9	92.6	74.5
2006	3.9	8.6	11.5	20.1	36.7	52.9	67.1	94.3	85.7
2005	5.2	9.4	13.0	25.9	33.7	54.6	74.9	84.2	88.6
2004	5.0	6.4	14.9	17.9	30.9	47.3	70.0	96.6	95.1
2003	6.3	8.7	14.6	19.7	32.0	47.5	79.8	102.2	119.5
2002	5.7	7.5	15.2	22.8	39.8	53.7	76.8	106.6	108.6
2001	4.8	9.1	12.8	28.2	32.1	51.3	68.3	91.7	111.7
2000	4.8	9.3	13.1	28.3	35.6	65.2	84.7	92.3	116.7
1999	6.8	10.2	16.7	23.1	44.4	58.2	74.9	115.3	130.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 4/, both sexes

2007.....	6.2	10.5	18.2	30.6	50.9	80.9	121.6	163.7	190.2
2006.....	6.4	11.2	18.8	33.0	52.4	82.9	121.4	164.8	190.1
2005.....	6.6	11.0	19.3	33.6	53.7	85.0	123.2	164.2	193.6
2004.....	6.7	11.7	20.4	33.7	55.6	85.8	124.8	164.9	193.0
2003.....	7.1	11.8	20.3	36.3	56.4	89.1	129.2	167.4	200.5
2002.....	7.6	12.8	21.7	37.5	59.9	91.7	127.2	170.5	199.8
2001.....	8.2	13.1	23.1	38.3	59.7	93.2	130.3	170.5	202.1
2000.....	8.4	13.6	23.8	38.4	63.2	92.3	132.7	178.1	206.5
1999.....	8.4	13.7	25.9	40.0	63.6	95.1	132.2	175.3	201.4

Non-Hispanic 4/, male

2007.....	7.5	13.3	22.9	38.7	65.5	104.2	160.9	223.0	260.3
2006.....	7.5	14.3	23.8	42.2	67.0	105.2	161.1	224.0	250.7
2005.....	7.8	13.6	24.4	41.4	69.3	111.4	163.9	218.5	268.5
2004.....	8.2	14.4	25.4	42.4	71.5	110.9	161.6	218.6	261.1
2003.....	8.9	14.7	25.5	46.6	73.6	114.9	166.3	220.9	273.6
2002.....	9.2	15.4	27.2	46.6	77.5	119.2	166.8	227.5	272.8
2001.....	10.0	16.2	28.1	48.0	76.3	118.7	166.4	223.6	278.8
2000.....	10.2	16.4	28.7	46.0	79.4	119.1	172.1	233.3	285.0
1999.....	10.2	16.9	31.4	49.4	79.3	120.5	171.8	224.3	284.3

Non-Hispanic 4/, female

2007.....	5.0	7.8	13.8	23.1	38.0	61.8	92.6	126.3	157.1
2006.....	5.4	8.2	14.1	24.6	39.5	64.8	92.3	128.0	162.0
2005.....	5.5	8.6	14.5	26.4	40.0	63.6	93.7	130.6	159.5
2004.....	5.4	9.2	15.7	25.8	41.6	65.6	98.3	132.0	162.6
2003.....	5.4	9.1	15.3	26.8	41.4	68.4	102.8	134.8	168.4
2002.....	6.0	10.3	16.5	29.3	44.6	69.7	99.2	135.9	168.3
2001.....	6.5	10.2	18.5	29.5	45.4	72.9	105.0	138.7	170.2
2000.....	6.6	10.9	19.2	31.4	49.2	71.2	105.3	145.6	174.8
1999.....	6.6	10.6	20.8	31.4	50.1	75.2	104.7	146.7	168.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic White, both sexes

2007 5/.....	5.9	10.0	17.5	30.5	51.2	82.9	125.2	169.2	198.0
2006 5/.....	6.0	11.0	18.4	32.6	53.1	84.9	125.0	171.1	196.6
2005 5/.....	6.2	10.5	18.9	33.6	54.3	86.9	127.5	170.1	201.5
2004 5/.....	6.5	11.3	20.1	33.8	56.3	86.7	127.6	169.7	198.8
2003 5/.....	6.9	11.4	20.3	36.5	57.3	90.7	132.3	172.1	207.2
2002.....	7.3	12.3	21.2	37.6	61.0	93.5	130.4	175.5	206.5
2001.....	8.0	12.9	22.6	38.2	60.0	95.2	133.3	174.9	208.2
2000.....	8.0	13.3	23.6	38.6	64.5	94.1	135.7	183.7	210.7
1999.....	8.2	13.4	25.6	39.8	64.6	96.9	134.7	179.9	206.4

Non-Hispanic White, male

2007 5/.....	7.2	12.8	22.2	39.0	66.3	106.9	165.9	231.0	271.4
2006 5/.....	6.9	14.0	23.2	41.6	67.8	107.5	165.4	232.9	258.6
2005 5/.....	7.3	12.9	23.9	41.7	70.6	113.8	168.8	226.1	281.9
2004 5/.....	7.8	13.8	25.3	42.6	72.0	112.0	165.5	225.5	270.9
2003 5/.....	8.6	14.1	25.4	47.3	75.4	118.0	171.9	228.8	284.6
2002.....	8.9	15.0	26.8	46.6	79.1	121.8	171.7	234.5	283.5
2001.....	9.9	16.0	27.4	47.9	76.7	121.2	170.6	229.3	288.7
2000.....	9.8	15.9	28.4	46.2	81.5	120.4	176.5	240.8	293.8
1999.....	9.9	16.6	31.0	49.4	80.6	122.7	174.6	231.2	291.8

Non-Hispanic White, female

2007 5/.....	4.7	7.3	12.9	22.4	37.7	62.7	94.6	129.8	163.2
2006 5/.....	5.0	8.0	13.7	24.1	40.0	66.0	94.7	132.2	167.7
2005 5/.....	5.2	8.1	14.2	26.0	39.8	64.5	96.9	135.2	164.7
2004 5/.....	5.2	8.9	15.1	25.6	42.2	65.7	99.9	135.1	166.5
2003 5/.....	5.2	8.7	15.3	26.5	41.2	68.2	103.7	137.2	173.1
2002.....	5.6	9.7	15.8	29.2	45.0	70.3	100.9	139.5	173.2
2001.....	6.2	9.8	18.0	29.3	45.2	74.0	106.8	142.1	174.6
2000.....	6.2	10.8	19.0	31.5	49.5	73.0	107.0	149.8	177.2
1999.....	6.5	10.2	20.5	30.9	50.5	76.3	106.6	149.7	172.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic Black, both sexes

2007 5/.....	9.0	15.4	26.1	38.7	59.0	82.4	106.5	133.2	140.4
2006 5/.....	10.3	14.9	26.6	43.8	57.8	85.0	111.2	129.2	145.5
2005 5/.....	9.4	16.5	25.2	39.4	59.6	82.8	104.7	129.2	139.3
2004 5/.....	9.3	17.0	26.9	41.2	61.4	94.4	117.4	136.7	156.3
2003 5/.....	9.8	17.7	24.0	42.2	60.5	91.7	118.4	140.8	152.3
2002.....	10.8	17.5	30.3	44.3	63.1	94.6	113.6	138.5	153.7
2001.....	10.7	16.9	31.0	44.9	68.4	92.2	117.1	144.0	157.3
2000.....	12.6	18.0	29.7	44.8	63.5	91.5	115.8	132.8	177.1
1999.....	10.6	18.5	33.7	46.5	65.8	91.8	120.8	140.0	162.3

Non-Hispanic Black, male

2007 5/.....	10.6	20.0	31.4	46.5	73.7	102.5	138.1	169.2	185.0
2006 5/.....	12.7	19.6	35.3	57.3	77.3	112.7	146.9	169.9	198.8
2005 5/.....	12.2	20.5	33.7	47.6	74.5	110.5	141.4	174.8	168.4
2004 5/.....	11.5	21.5	32.2	51.8	81.1	124.0	140.8	171.8	197.1
2003 5/.....	12.2	23.1	32.7	52.6	74.9	106.8	138.3	169.8	188.6
2002.....	13.5	20.0	36.8	56.4	80.1	125.2	140.4	180.2	203.5
2001.....	12.1	21.4	38.5	57.1	89.4	115.7	142.2	194.6	206.7
2000.....	15.0	22.4	37.6	55.1	74.9	127.1	149.8	170.8	226.6
1999.....	13.4	22.8	42.2	56.2	82.1	116.3	161.0	166.5	221.1

Non-Hispanic Black, female

2007 5/.....	7.7	11.5	21.8	32.7	48.2	68.6	87.5	114.4	121.9
2006 5/.....	8.2	11.0	19.5	33.4	43.5	66.0	89.9	107.8	123.7
2005 5/.....	7.0	13.0	18.2	33.0	48.6	63.8	82.8	105.1	127.5
2004 5/.....	7.4	13.2	22.5	33.0	46.8	74.4	103.5	118.2	140.1
2003 5/.....	7.6	13.2	16.9	34.0	49.7	81.6	106.5	125.6	138.3
2002.....	8.5	15.4	25.0	34.8	50.5	74.1	97.6	116.9	134.6
2001.....	9.5	13.0	24.8	35.3	52.9	76.6	102.1	118.3	139.0
2000.....	10.5	14.3	23.3	36.7	55.0	67.8	95.2	113.6	158.7
1999.....	8.2	14.8	26.9	38.9	53.7	75.6	96.7	126.9	140.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hodgkin's disease (C81)

All origins 3/, both sexes

2007.....	0.3	0.4	0.5	0.6	0.9	1.3	1.8	2.6	2.4
2006.....	0.3	0.4	0.5	0.7	0.9	1.3	2.1	2.3	2.6
2005.....	0.4	0.4	0.5	0.7	1.0	1.2	2.2	2.5	2.1
2004.....	0.4	0.4	0.7	0.8	1.0	1.4	1.8	2.1	2.2
2003.....	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4

All origins 3/, male

2007.....	0.4	0.6	0.5	0.8	1.2	1.5	2.5	2.9	2.8
2006.....	0.3	0.6	0.8	0.9	1.2	1.6	2.3	3.0	3.7
2005.....	0.5	0.4	0.8	0.9	1.2	1.6	2.3	2.7	3.1
2004.....	0.5	0.5	0.9	1.0	1.4	1.9	2.0	3.2	3.2
2003.....	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8

All origins 3/, female

2007.....	*	0.2	0.4	0.5	0.7	1.1	1.4	2.5	2.2
2006.....	0.2	0.2	0.3	0.5	0.7	1.0	2.0	1.9	2.1
2005.....	0.4	0.3	0.3	0.5	0.8	1.0	2.1	2.3	1.6
2004.....	0.3	0.3	0.5	0.5	0.8	1.0	1.6	1.5	1.7
2003.....	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hodgkin's disease (C81)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.4	0.4	0.6	0.9	1.2	1.8	2.6	2.4
2006.....	0.3	0.4	0.5	0.7	0.9	1.2	2.1	2.3	2.6
2005.....	0.4	0.4	0.5	0.7	0.9	1.1	2.1	2.4	2.0
2004.....	0.4	0.5	0.7	0.7	1.0	1.4	1.7	2.1	2.1
2003.....	0.3	0.4	0.6	0.9	0.8	1.5	2.0	2.3	2.3
2002.....	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.2	2.8
2001.....	0.5	0.5	0.4	0.7	1.0	1.2	2.1	2.5	2.2
2000.....	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.4
1999.....	0.4	0.5	0.6	0.8	1.2	1.8	2.0	2.2	2.3

Non-Hispanic 4/, male

2007.....	0.4	0.6	0.5	0.8	1.1	1.4	2.5	2.9	2.7
2006.....	0.4	0.6	0.8	0.9	1.2	1.5	2.3	2.9	3.8
2005.....	0.5	0.4	0.7	0.8	1.1	1.5	2.3	2.5	3.0
2004.....	0.5	0.6	0.9	1.0	1.3	1.9	2.0	3.2	3.1
2003.....	0.4	0.5	0.8	1.1	1.1	1.9	2.3	2.9	2.7
2002.....	0.5	0.5	0.7	1.0	1.4	2.2	2.0	3.1	3.7
2001.....	0.6	0.6	0.5	0.9	1.4	1.6	2.5	2.9	2.5
2000.....	0.4	0.6	0.5	1.0	1.3	1.9	2.1	2.6	3.7
1999.....	0.5	0.5	0.8	1.2	1.6	2.2	2.2	2.5	3.7

Non-Hispanic 4/, female

2007.....	*	0.3	0.4	0.5	0.7	1.1	1.4	2.4	2.2
2006.....	0.3	0.2	0.3	0.5	0.7	1.0	2.0	1.9	2.1
2005.....	0.4	0.3	0.3	0.5	0.7	0.9	2.0	2.4	1.6
2004.....	0.3	0.3	0.5	0.5	0.8	1.0	1.5	1.4	1.6
2003.....	0.2	0.4	0.5	0.6	0.6	1.2	1.9	2.0	2.2
2002.....	0.3	0.5	*	0.7	0.6	1.1	1.5	1.6	2.4
2001.....	0.4	0.4	0.3	0.6	0.7	0.9	1.8	2.2	2.0
2000.....	0.3	0.4	0.5	0.6	0.7	1.3	1.4	2.1	2.0
1999.....	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hodgkin's disease (C81)

Non-Hispanic White, both sexes

2007 5/.....	0.3	0.4	0.4	0.6	0.9	1.4	2.0	2.9	2.5
2006 5/.....	0.3	0.5	0.6	0.7	1.0	1.2	2.2	2.4	2.7
2005 5/.....	0.4	0.4	0.5	0.7	1.0	1.2	2.3	2.7	2.1
2004 5/.....	0.4	0.5	0.7	0.8	1.0	1.5	1.8	2.2	2.1
2003 5/.....	0.4	0.5	0.6	0.9	0.8	1.6	2.1	2.4	2.5
2002.....	0.4	0.5	0.5	0.9	1.0	1.7	1.8	2.3	2.9
2001.....	0.5	0.5	0.4	0.8	1.2	1.2	2.1	2.6	2.3
2000.....	0.4	0.5	0.5	0.8	1.1	1.6	1.9	2.4	2.6
1999.....	0.4	0.5	0.7	0.9	1.2	1.9	2.1	2.3	2.3

Non-Hispanic White, male

2007 5/.....	0.4	0.6	0.4	0.8	1.2	1.5	2.6	3.2	2.9
2006 5/.....	0.3	0.6	0.8	1.0	1.2	1.5	2.4	3.0	3.9
2005 5/.....	0.4	0.4	0.7	0.9	1.2	1.5	2.4	2.8	3.2
2004 5/.....	0.5	0.6	1.0	1.0	1.3	2.0	2.1	3.5	3.3
2003 5/.....	0.5	0.5	0.7	1.1	1.1	2.0	2.3	3.1	2.7
2002.....	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.3	4.0
2001.....	0.6	0.6	0.5	0.9	1.5	1.7	2.6	3.0	2.6
2000.....	0.5	0.6	0.5	1.0	1.4	1.9	2.4	2.7	3.9
1999.....	0.5	0.5	0.9	1.3	1.6	2.4	2.4	2.7	3.8

Non-Hispanic White, female

2007 5/.....	*	*	0.4	0.5	0.7	1.2	1.5	2.7	2.4
2006 5/.....	*	0.3	0.4	0.5	0.8	1.0	2.1	2.1	2.1
2005 5/.....	0.4	0.3	0.4	0.6	0.8	0.9	2.2	2.6	1.5
2004 5/.....	0.4	0.3	0.5	0.5	0.8	1.1	1.7	1.5	1.6
2003 5/.....	0.2	0.4	0.5	0.7	0.6	1.4	2.0	2.0	2.4
2002.....	0.3	0.5	*	0.8	0.6	1.2	1.6	1.7	2.5
2001.....	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.3
2000.....	0.3	0.4	0.5	0.7	0.8	1.4	1.6	2.2	2.1
1999.....	0.4	0.4	0.5	0.5	0.8	1.5	2.0	2.1	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

All origins 3/, both sexes

2007.....	2.4	4.0	7.1	11.2	18.5	28.9	45.0	62.7	72.1
2006.....	2.5	4.2	6.7	12.5	19.0	30.1	46.2	63.4	72.3
2005.....	2.4	4.3	7.6	12.3	19.2	31.5	46.9	64.4	75.2
2004.....	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
2003.....	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7

All origins 3/, male

2007.....	3.1	5.2	9.2	14.1	24.1	36.7	56.0	79.3	91.6
2006.....	3.1	5.6	8.8	16.0	23.8	37.2	57.3	80.7	89.6
2005.....	3.1	5.7	9.7	15.0	24.5	39.9	59.0	79.4	100.5
2004.....	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
2003.....	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4

All origins 3/, female

2007.....	1.8	2.8	5.1	8.6	13.6	22.5	36.8	52.2	62.8
2006.....	1.8	2.8	4.7	9.3	14.8	24.3	38.2	52.6	64.2
2005.....	1.7	3.0	5.7	9.9	14.5	24.7	38.1	55.2	63.5
2004.....	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
2003.....	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

Hispanic, both sexes

2007.....	2.6	3.3	6.4	10.6	13.3	24.5	33.2	50.2	47.6
2006.....	1.9	3.8	6.5	10.6	15.6	23.2	34.8	48.1	40.3
2005.....	2.2	3.7	5.6	10.6	14.4	26.0	40.0	52.0	44.5
2004.....	2.7	4.0	6.8	8.5	15.1	23.4	35.6	48.4	43.8
2003.....	2.8	3.4	7.7	9.4	15.4	23.7	35.8	48.0	54.4
2002.....	2.5	3.5	6.7	13.1	18.6	27.9	41.3	47.7	50.4
2001.....	2.9	4.5	7.9	14.9	14.4	23.2	41.6	46.2	59.4
2000.....	2.9	4.6	8.1	14.9	18.5	34.8	39.2	45.7	59.7
1999.....	3.8	5.2	6.8	12.5	20.2	31.5	37.8	51.7	56.1

Hispanic, male

2007.....	3.3	4.2	6.5	12.1	17.6	31.2	39.2	64.3	52.9
2006.....	2.4	4.2	8.6	12.5	17.5	28.2	40.7	59.0	54.1
2005.....	3.0	3.6	6.9	11.8	17.7	32.2	48.6	59.5	52.3
2004.....	3.8	6.4	7.6	10.5	18.5	31.5	44.3	51.9	50.4
2003.....	3.1	3.8	9.4	12.1	18.8	30.3	40.7	53.0	57.0
2002.....	3.3	4.0	8.4	16.2	23.4	29.4	42.2	51.5	60.2
2001.....	3.6	5.4	10.0	17.7	19.5	30.7	55.9	63.6	68.6
2000.....	3.8	5.7	10.5	18.4	22.4	40.4	44.3	55.9	82.6
1999.....	5.0	6.7	8.8	15.8	21.6	35.1	46.0	51.9	60.4

Hispanic, female

2007.....	1.9	2.3	6.2	9.3	9.8	19.2	28.8	40.9	44.7
2006.....	*	3.4	4.5	8.9	14.0	19.3	30.6	40.9	33.0
2005.....	*	3.9	4.4	9.5	11.6	21.3	33.8	47.0	40.4
2004.....	*	*	6.1	6.7	12.3	17.2	29.5	46.1	40.3
2003.....	2.4	3.0	6.2	7.1	12.5	18.7	32.3	44.7	53.1
2002.....	*	3.1	5.1	10.5	14.7	26.7	40.7	45.3	45.6
2001.....	2.1	3.7	6.0	12.5	10.3	17.4	31.5	35.4	54.9
2000.....	*	3.4	6.0	11.9	15.4	30.6	35.6	39.4	48.5
1999.....	2.6	3.8	5.0	9.8	19.1	28.7	31.9	51.6	54.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic 4/, both sexes

2007.....	2.4	4.0	7.2	11.3	18.9	29.2	45.8	63.3	73.4
2006.....	2.5	4.2	6.7	12.6	19.2	30.6	47.0	64.3	73.9
2005.....	2.5	4.4	7.8	12.5	19.5	31.9	47.3	65.1	76.6
2004.....	2.8	4.8	7.8	12.9	21.5	32.5	47.5	65.5	76.3
2003.....	3.0	4.8	8.1	14.7	22.0	33.3	50.0	66.1	79.6
2002.....	3.0	5.1	9.0	15.0	23.5	35.4	50.5	67.5	78.2
2001.....	3.5	5.6	9.8	15.6	23.5	36.7	51.2	70.6	80.3
2000.....	3.7	5.8	10.8	16.1	25.4	36.3	54.3	72.7	80.9
1999.....	3.5	6.2	11.1	16.5	25.9	38.7	55.2	73.2	78.3

Non-Hispanic 4/, male

2007.....	3.1	5.2	9.4	14.2	24.7	37.0	57.1	80.2	93.9
2006.....	3.2	5.8	8.8	16.2	24.3	37.8	58.3	81.9	91.5
2005.....	3.2	5.9	9.9	15.2	25.0	40.5	59.6	80.4	103.0
2004.....	3.7	6.0	10.0	16.1	27.2	40.2	58.3	83.0	97.8
2003.....	3.9	6.2	10.1	19.1	28.5	41.7	61.8	83.1	105.6
2002.....	3.9	6.4	11.5	18.5	30.4	43.5	62.7	85.5	101.4
2001.....	4.5	7.5	12.1	19.1	29.7	45.5	61.3	88.2	108.1
2000.....	4.8	7.6	13.4	19.1	31.0	45.3	67.2	89.3	106.7
1999.....	4.7	8.0	13.8	19.9	31.3	47.3	68.9	86.5	106.0

Non-Hispanic 4/, female

2007.....	1.7	2.9	5.0	8.6	13.9	22.8	37.4	52.7	63.8
2006.....	1.9	2.7	4.7	9.3	14.8	24.7	38.6	53.3	65.7
2005.....	1.8	2.9	5.8	9.9	14.7	25.0	38.3	55.6	64.6
2004.....	1.9	3.7	5.8	10.1	16.5	26.3	39.8	54.8	66.7
2003.....	2.1	3.5	6.1	10.6	16.4	26.6	41.6	55.8	68.2
2002.....	2.2	3.9	6.5	11.8	17.5	28.8	42.0	56.6	68.2
2001.....	2.5	3.8	7.6	12.4	18.1	29.8	44.1	60.1	68.7
2000.....	2.6	4.2	8.3	13.4	20.7	29.1	45.3	62.9	70.4
1999.....	2.3	4.6	8.6	13.5	21.2	32.0	45.6	65.3	67.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic White, both sexes

2007 5/.....	2.3	4.0	7.2	11.6	19.9	31.0	48.3	67.1	78.2
2006 5/.....	2.3	4.3	6.6	12.9	20.1	32.6	50.0	68.0	78.2
2005 5/.....	2.4	4.2	7.9	12.8	20.6	33.7	50.5	68.6	81.3
2004 5/.....	2.7	4.8	8.0	13.5	22.7	34.2	50.0	69.5	80.2
2003 5/.....	2.9	4.8	8.5	15.2	23.3	34.9	52.6	69.9	84.3
2002.....	3.1	5.0	9.1	15.8	24.7	36.9	53.4	71.1	83.1
2001.....	3.4	5.7	9.9	16.3	24.5	38.7	54.0	74.0	84.1
2000.....	3.7	6.0	11.4	16.9	26.8	38.1	57.1	76.9	85.1
1999.....	3.5	6.5	11.5	17.4	27.5	40.6	57.6	77.3	82.0

Non-Hispanic White, male

2007 5/.....	3.0	5.2	9.4	14.6	25.7	38.8	60.3	84.4	99.6
2006 5/.....	2.9	5.8	8.7	16.6	25.1	40.1	61.8	87.0	96.2
2005 5/.....	3.0	5.7	10.1	15.5	26.4	42.5	63.6	84.8	109.8
2004 5/.....	3.5	6.0	10.1	16.6	28.4	42.0	61.0	87.9	103.5
2003 5/.....	3.8	6.2	10.6	19.8	30.3	43.9	65.2	88.1	112.0
2002.....	4.0	6.4	11.8	19.2	31.9	45.3	66.1	89.8	107.6
2001.....	4.5	7.6	12.1	19.8	30.9	47.5	64.3	92.2	113.1
2000.....	4.7	7.6	13.9	19.9	32.8	46.9	70.8	94.1	113.0
1999.....	4.7	8.2	14.1	20.6	32.9	49.5	71.1	91.8	110.0

Non-Hispanic White, female

2007 5/.....	1.6	2.9	5.0	8.6	14.8	24.5	39.3	56.1	68.0
2006 5/.....	1.8	2.8	4.7	9.5	15.6	26.3	41.2	56.1	69.8
2005 5/.....	1.7	2.8	5.9	10.2	15.5	26.3	40.7	58.6	68.3
2004 5/.....	1.8	3.6	6.0	10.5	17.7	27.8	41.9	58.1	69.8
2003 5/.....	2.1	3.5	6.4	10.9	17.0	27.5	43.6	58.7	72.2
2002.....	2.1	3.7	6.6	12.5	18.4	30.0	44.3	59.7	72.6
2001.....	2.4	3.9	7.8	13.1	18.8	31.6	46.7	63.0	72.0
2000.....	2.6	4.4	9.0	14.1	21.6	31.1	47.6	66.6	73.8
1999.....	2.3	4.8	9.0	14.4	22.7	33.4	48.2	68.8	70.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic Black, both sexes

2007 5/.....	3.4	4.7	8.3	11.0	15.1	18.8	23.8	29.4	31.7
2006 5/.....	4.3	4.8	8.7	12.5	15.4	19.7	25.6	32.0	34.5
2005 5/.....	3.3	5.7	7.6	11.6	14.2	21.0	23.0	32.8	34.4
2004 5/.....	4.0	5.9	7.7	11.0	16.0	21.5	29.3	26.6	39.4
2003 5/.....	4.0	6.1	6.2	12.9	14.9	22.9	27.0	31.1	31.2
2002.....	3.6	6.2	9.4	12.6	18.0	26.2	26.9	35.0	31.2
2001.....	4.3	6.1	9.9	12.1	19.3	24.0	25.9	38.1	40.0
2000.....	4.5	5.6	9.1	12.8	17.0	22.9	26.3	28.5	36.6
1999.....	4.1	5.6	10.8	11.6	17.3	24.4	32.5	31.4	40.4

Non-Hispanic Black, male

2007 5/.....	4.0	6.6	10.7	13.8	22.0	25.8	28.7	34.0	37.5
2006 5/.....	6.1	7.2	12.2	16.6	20.5	24.1	30.8	35.8	43.1
2005 5/.....	4.6	8.5	10.5	15.0	18.1	29.6	26.6	38.1	40.9
2004 5/.....	5.6	7.8	10.2	14.4	20.0	28.3	32.2	35.2	42.7
2003 5/.....	5.8	7.7	8.0	16.5	17.7	24.6	29.0	34.7	35.2
2002.....	4.2	7.6	11.7	16.9	23.3	34.9	31.9	42.1	43.8
2001.....	5.1	8.8	12.4	15.5	25.0	32.4	30.8	48.2	55.4
2000.....	6.2	8.0	13.9	15.9	19.7	32.5	31.3	32.6	43.4
1999.....	5.6	7.9	14.1	15.7	23.9	29.1	47.9	28.3	59.4

Non-Hispanic Black, female

2007 5/.....	2.8	3.1	6.2	8.8	10.1	13.9	20.8	26.9	29.3
2006 5/.....	2.7	2.9	5.8	9.4	11.6	16.7	22.5	30.0	31.0
2005 5/.....	2.2	3.3	5.2	8.9	11.2	15.1	20.8	30.0	31.8
2004 5/.....	2.5	4.4	5.7	8.4	13.1	16.9	27.6	22.0	38.1
2003 5/.....	2.4	4.7	4.8	10.2	12.8	21.8	25.8	29.2	29.7
2002.....	3.1	5.0	7.5	9.2	14.0	20.4	23.8	31.2	26.3
2001.....	3.7	3.7	7.9	9.5	15.0	18.4	22.9	32.9	34.2
2000.....	3.1	3.6	5.1	10.4	14.9	16.6	23.2	26.4	34.0
1999.....	2.9	3.8	8.1	8.4	12.4	21.4	23.3	32.9	33.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

All origins 3/, both sexes

2007.....	2.5	3.7	6.5	11.1	18.6	30.3	45.4	62.0	76.7
2006.....	2.5	4.1	7.0	11.6	19.9	31.3	44.7	62.9	76.9
2005.....	2.7	4.1	6.5	11.8	19.8	31.3	45.7	61.0	78.7
2004.....	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
2003.....	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4

All origins 3/, male

2007.....	2.8	4.5	8.2	14.3	24.6	40.7	63.8	92.4	109.4
2006.....	2.7	4.9	8.7	15.1	26.3	41.6	64.8	91.8	105.1
2005.....	2.9	4.4	8.3	15.4	26.1	43.9	66.0	87.9	112.4
2004.....	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
2003.....	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2

All origins 3/, female

2007.....	2.3	2.9	4.9	8.3	13.4	21.9	31.8	42.9	61.2
2006.....	2.3	3.3	5.3	8.5	14.3	22.8	30.0	44.8	63.7
2005.....	2.5	3.7	4.9	8.5	14.4	21.1	31.0	44.3	63.3
2004.....	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
2003.....	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

Hispanic, both sexes

2007.....	2.3	3.6	5.8	7.8	11.6	17.5	23.8	34.4	30.4
2006.....	2.3	3.3	4.8	7.5	13.3	21.7	24.9	36.3	35.2
2005.....	2.8	3.8	5.6	8.9	15.3	20.7	27.8	29.0	40.6
2004.....	2.2	4.0	5.1	7.9	13.6	18.4	25.1	34.2	36.2
2003.....	3.5	3.8	4.7	8.6	11.9	16.5	29.0	39.3	47.2
2002.....	2.5	2.7	6.5	8.2	13.4	18.1	21.6	37.0	47.1
2001.....	2.6	3.5	4.5	9.2	14.3	21.2	32.0	35.8	42.5
2000.....	3.3	3.8	6.4	9.6	13.5	21.2	29.1	39.0	49.1
1999.....	3.0	3.9	6.8	7.3	14.9	20.3	27.4	37.6	44.2

Hispanic, male

2007.....	2.3	3.9	6.5	9.2	14.9	19.8	29.7	42.6	51.8
2006.....	2.9	3.2	5.5	9.4	16.6	26.7	36.1	47.5	42.9
2005.....	3.1	3.8	7.1	10.3	19.5	27.4	38.3	47.3	57.3
2004.....	2.1	5.0	6.1	11.2	17.9	23.2	34.0	44.4	43.4
2003.....	4.2	3.9	5.0	11.3	14.6	23.1	35.9	48.9	61.6
2002.....	2.1	3.7	6.6	8.0	15.2	22.8	32.0	50.3	65.3
2001.....	3.1	3.6	5.2	12.8	16.3	23.8	49.1	39.0	68.6
2000.....	4.3	3.9	9.0	11.8	16.8	25.8	35.4	57.4	76.6
1999.....	3.6	3.5	7.2	10.4	17.4	31.6	35.1	47.2	49.6

Hispanic, female

2007.....	2.3	3.3	5.1	6.6	8.9	15.8	19.5	29.0	18.8
2006.....	1.8	3.4	4.1	5.8	10.4	17.8	16.8	28.9	31.2
2005.....	2.4	3.8	4.3	7.7	11.8	15.4	20.2	16.8	31.9
2004.....	2.3	2.9	4.2	5.1	10.0	14.7	18.8	27.4	32.4
2003.....	2.9	3.6	4.4	6.2	9.6	11.5	24.1	33.2	39.8
2002.....	2.8	*	6.4	8.5	11.9	14.5	14.2	28.5	38.1
2001.....	*	3.3	3.8	6.1	12.6	19.2	19.8	33.8	29.8
2000.....	2.3	3.7	4.0	7.7	10.9	17.7	24.6	27.8	35.6
1999.....	2.4	4.3	6.5	*	12.8	11.9	22.0	31.9	41.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

Non-Hispanic 4/, both sexes

2007.....	2.6	3.7	6.6	11.4	19.2	31.3	46.8	63.6	79.0
2006.....	2.5	4.1	7.1	11.9	20.4	32.0	46.0	64.3	78.9
2005.....	2.7	4.1	6.6	12.0	20.2	32.1	46.9	62.6	80.6
2004.....	2.5	4.2	7.3	12.3	20.4	32.7	46.8	62.8	80.6
2003.....	2.6	4.0	7.1	12.7	20.1	33.5	48.4	63.9	81.6
2002.....	2.9	4.7	7.5	13.3	21.4	33.2	46.9	64.5	81.7
2001.....	3.1	4.7	7.8	13.4	21.0	33.9	48.4	62.2	82.7
2000.....	3.1	4.5	7.8	12.9	22.2	33.8	48.1	66.0	84.7
1999.....	3.2	4.4	8.6	13.5	22.2	33.7	47.0	63.5	83.7

Non-Hispanic 4/, male

2007.....	2.9	4.6	8.4	14.7	25.4	42.2	66.1	95.3	112.6
2006.....	2.7	5.1	9.0	15.5	27.1	42.7	66.7	94.2	108.3
2005.....	2.9	4.5	8.3	15.8	26.5	45.1	67.7	90.0	115.3
2004.....	2.9	5.0	9.2	16.0	27.4	45.3	65.8	88.7	113.9
2003.....	3.1	4.9	9.2	16.1	27.7	46.2	67.9	90.7	113.7
2002.....	3.3	5.4	9.4	16.7	28.8	47.0	66.5	91.7	116.5
2001.....	3.4	5.3	9.5	17.7	28.3	46.0	66.9	88.7	119.3
2000.....	3.5	5.1	9.6	16.3	30.0	47.3	66.4	94.5	121.5
1999.....	3.6	5.3	10.2	17.2	29.6	46.3	65.8	89.0	122.1

Non-Hispanic 4/, female

2007.....	2.3	2.9	4.9	8.4	13.8	22.3	32.6	43.6	63.2
2006.....	2.3	3.3	5.4	8.7	14.6	23.2	30.8	45.7	65.2
2005.....	2.5	3.7	4.9	8.5	14.6	21.5	31.7	45.7	64.8
2004.....	2.2	3.4	5.5	8.9	14.3	22.6	33.1	46.9	65.7
2003.....	2.2	3.2	5.2	9.6	13.6	23.4	34.5	47.6	67.4
2002.....	2.5	4.0	5.7	10.1	14.9	22.2	33.0	48.0	66.6
2001.....	2.8	4.1	6.2	9.5	14.7	24.3	35.5	46.4	67.5
2000.....	2.6	4.0	6.0	9.8	15.5	23.1	35.4	49.2	69.8
1999.....	2.8	3.5	7.1	10.1	15.8	23.8	34.0	48.5	68.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

Non-Hispanic White, both sexes

2007 5/.....	2.6	3.7	6.6	11.7	19.7	32.4	49.1	66.6	82.9
2006 5/.....	2.5	4.3	7.3	12.1	21.3	33.3	47.4	67.6	82.0
2005 5/.....	2.6	4.0	6.7	12.4	20.9	33.6	48.8	65.6	84.5
2004 5/.....	2.6	4.1	7.4	12.5	21.0	33.5	48.5	65.0	83.6
2003 5/.....	2.7	4.0	7.3	13.0	21.0	34.8	50.4	66.2	84.7
2002.....	2.7	4.7	7.4	13.4	22.4	34.5	48.3	66.9	84.4
2001.....	3.1	4.6	7.9	13.7	21.4	35.2	49.8	64.8	86.1
2000.....	2.9	4.5	7.8	13.1	23.2	35.1	49.9	68.5	87.3
1999.....	3.2	4.3	8.8	13.7	22.6	34.9	48.8	65.0	86.8

Non-Hispanic White, male

2007 5/.....	2.9	4.6	8.5	15.2	26.0	43.9	69.2	100.1	118.1
2006 5/.....	2.7	5.2	9.2	15.8	28.0	44.4	68.4	98.8	112.1
2005 5/.....	2.8	4.5	8.6	16.3	27.6	47.0	69.8	93.7	121.9
2004 5/.....	2.9	4.8	9.5	16.4	27.9	45.9	68.1	92.2	118.8
2003 5/.....	3.3	4.8	9.3	16.8	28.7	48.0	71.1	94.8	119.0
2002.....	3.1	5.5	9.4	16.9	29.9	48.5	68.7	95.1	121.3
2001.....	3.5	5.2	9.7	18.1	28.6	47.8	69.0	92.0	125.3
2000.....	3.4	5.0	9.7	16.5	31.3	48.6	68.9	97.9	125.5
1999.....	3.5	5.2	10.5	17.5	30.2	47.9	68.0	91.7	127.2

Non-Hispanic White, female

2007 5/.....	2.3	2.8	4.8	8.4	14.1	22.8	33.9	45.2	66.2
2006 5/.....	2.3	3.4	5.6	8.7	15.4	24.0	31.7	47.9	68.0
2005 5/.....	2.4	3.6	5.0	8.7	14.9	22.5	33.2	48.0	67.3
2004 5/.....	2.3	3.5	5.4	8.9	14.8	23.2	34.0	48.2	67.8
2003 5/.....	2.2	3.2	5.4	9.6	14.0	23.9	35.4	48.6	69.6
2002.....	2.4	4.0	5.4	10.2	15.6	23.1	33.8	49.6	68.4
2001.....	2.7	3.9	6.2	9.7	15.0	25.0	36.2	48.4	69.8
2000.....	2.5	4.0	6.0	9.9	16.0	24.2	36.6	51.1	72.0
1999.....	2.9	3.4	7.2	10.1	15.9	24.6	35.3	49.2	70.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

Non-Hispanic Black, both sexes

2007 5/.....	2.8	4.7	7.5	11.8	18.1	29.1	35.6	45.6	52.9
2006 5/.....	2.7	4.2	7.5	13.1	18.2	29.2	39.9	42.4	56.8
2005 5/.....	3.2	5.0	6.9	11.0	19.0	24.7	35.6	44.0	53.0
2004 5/.....	2.9	5.3	7.9	14.0	19.5	32.9	37.5	47.8	60.7
2003 5/.....	2.6	5.2	7.2	13.1	18.2	29.9	37.6	48.7	60.4
2002.....	3.8	5.1	9.4	14.2	17.4	29.1	40.0	47.2	65.3
2001.....	3.1	5.7	8.1	12.5	22.4	28.4	41.8	42.5	57.1
2000.....	4.7	5.3	8.1	14.3	18.1	29.0	35.3	44.3	66.1
1999.....	3.6	5.5	8.3	14.0	22.6	28.3	35.6	51.4	55.9

Non-Hispanic Black, male

2007 5/.....	3.4	6.0	8.9	13.6	24.1	38.0	47.9	61.0	78.5
2006 5/.....	2.7	5.3	10.1	17.0	27.3	39.3	59.4	62.5	87.0
2005 5/.....	3.9	5.2	8.5	14.2	24.7	35.1	53.3	68.4	63.8
2004 5/.....	3.0	7.1	9.5	17.5	28.4	49.2	48.5	59.5	81.3
2003 5/.....	2.8	6.9	10.9	14.9	25.1	40.1	50.5	59.6	78.8
2002.....	5.0	5.6	11.4	17.7	23.8	43.4	53.5	65.5	87.5
2001.....	2.9	6.8	10.1	17.6	32.6	36.8	52.6	62.1	75.1
2000.....	5.1	6.3	10.2	18.9	22.9	44.4	50.6	65.2	98.6
1999.....	4.6	7.1	10.6	17.6	29.5	38.4	52.2	62.0	78.4

Non-Hispanic Black, female

2007 5/.....	2.3	3.6	6.3	10.4	13.7	22.9	28.2	37.5	42.3
2006 5/.....	2.7	3.2	5.4	10.1	11.4	22.4	28.3	31.8	44.5
2005 5/.....	2.7	4.9	5.5	8.5	14.7	17.6	25.1	31.1	48.6
2004 5/.....	2.7	3.8	6.5	11.2	12.9	21.9	30.9	41.7	52.6
2003 5/.....	2.5	3.8	4.3	11.7	13.1	22.9	30.0	43.0	53.2
2002.....	2.7	4.7	7.8	11.5	12.7	19.5	32.0	37.7	56.9
2001.....	3.4	4.8	6.5	8.5	14.8	22.7	35.4	32.5	50.5
2000.....	4.3	4.5	6.4	10.6	14.5	18.8	26.1	33.7	54.1
1999.....	2.7	4.2	6.4	11.1	17.5	21.6	25.6	46.1	47.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All origins 3/, both sexes

2007.....	1.0	2.3	4.0	7.3	11.9	19.0	26.7	33.3	34.1
2006.....	1.1	2.4	4.3	7.5	11.8	19.0	25.8	33.3	33.8
2005.....	1.1	2.2	4.3	8.4	12.8	19.6	26.5	33.4	33.6
2004.....	1.0	2.3	4.6	7.7	12.3	18.9	28.3	33.9	33.1
2003.....	1.1	2.5	4.4	7.9	13.1	20.4	28.5	34.5	36.3
2002.....	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001.....	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000.....	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
1999.....	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6

All origins 3/, male

2007.....	1.2	2.8	4.5	9.0	14.2	23.4	34.2	43.4	49.4
2006.....	1.2	2.9	5.1	9.2	14.3	22.9	33.1	43.8	45.8
2005.....	1.3	2.7	5.2	9.4	16.3	23.9	33.5	44.5	46.1
2004.....	1.1	2.7	5.2	9.3	15.3	23.2	35.0	42.8	44.6
2003.....	1.3	3.1	5.4	10.1	16.0	24.3	33.9	43.3	50.4
2002.....	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001.....	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000.....	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
1999.....	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3

All origins 3/, female

2007.....	0.8	1.8	3.5	5.7	9.9	15.5	21.1	27.0	26.8
2006.....	0.9	1.9	3.6	6.0	9.6	15.8	20.5	26.8	28.2
2005.....	0.9	1.7	3.5	7.4	9.9	16.1	21.4	26.5	27.9
2004.....	0.9	1.8	4.0	6.2	9.8	15.5	23.5	28.4	28.0
2003.....	0.9	1.9	3.5	6.0	10.6	17.2	24.7	29.2	30.1
2002.....	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001.....	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000.....	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
1999.....	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Hispanic, both sexes

2007	0.9	1.9	4.0	7.2	12.6	18.0	21.9	22.0	14.3
2006	0.9	2.2	3.1	5.2	12.0	17.0	20.1	23.9	22.7
2005	1.1	2.2	3.7	7.8	10.6	17.0	19.9	23.7	20.1
2004	*	1.9	4.2	7.4	9.5	16.2	22.5	26.3	18.6
2003	*	2.3	4.4	6.4	10.1	15.8	24.2	23.4	24.9
2002	1.2	2.7	4.3	5.9	14.3	14.3	25.1	29.4	33.3
2001	*	2.2	3.9	8.3	12.2	18.2	22.7	28.3	29.4
2000	1.2	1.5	3.2	7.9	13.0	17.4	25.1	23.4	31.2
1999	1.2	2.3	5.6	8.3	12.1	15.1	19.4	30.0	28.1

Hispanic, male

2007	1.5	2.2	3.0	9.1	12.7	22.4	24.3	26.3	24.4
2006	*	2.8	3.7	5.9	12.4	20.6	24.2	25.4	30.4
2005	*	2.7	3.4	7.2	13.0	18.3	23.8	30.6	29.9
2004	*	*	4.1	9.8	11.1	18.2	27.1	34.9	*
2003	*	2.8	5.1	6.6	11.7	15.4	27.3	22.9	*
2002	*	2.8	5.5	8.2	17.3	18.0	32.6	29.7	53.5
2001	*	*	4.8	7.6	17.4	24.2	32.7	41.6	38.1
2000	*	*	*	8.1	17.5	19.4	28.1	33.9	*
1999	*	*	6.7	9.2	14.3	15.3	21.1	31.4	*

Hispanic, female

2007	*	*	4.9	5.6	12.5	14.6	20.2	19.1	*
2006	*	*	2.6	4.5	11.6	14.3	17.1	22.9	18.6
2005	*	*	4.0	8.3	8.6	16.0	17.1	19.2	15.0
2004	*	*	4.3	5.3	8.2	14.7	19.2	20.6	18.7
2003	*	*	3.6	6.2	8.8	16.1	22.0	23.7	22.7
2002	*	2.5	*	*	11.9	11.5	19.7	29.2	23.2
2001	*	*	*	8.9	8.0	13.6	15.5	20.1	25.1
2000	*	*	*	7.7	9.4	15.8	23.0	*	30.7
1999	*	*	4.6	7.4	10.3	14.9	18.2	29.1	31.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 4/, both sexes

2007.....	1.0	2.3	4.0	7.3	11.8	19.1	27.0	34.0	35.1
2006.....	1.1	2.4	4.4	7.7	11.7	19.1	26.2	33.8	34.4
2005.....	1.1	2.2	4.4	8.4	13.0	19.8	26.9	33.8	34.2
2004.....	1.0	2.3	4.6	7.7	12.6	19.1	28.6	34.1	33.7
2003.....	1.1	2.5	4.4	8.0	13.3	20.6	28.7	35.0	36.7
2002.....	1.3	2.5	4.8	8.4	14.0	21.4	27.9	36.1	36.7
2001.....	1.2	2.3	5.2	8.5	14.1	21.2	28.5	34.8	36.5
2000.....	1.2	2.7	4.6	8.5	14.5	20.5	28.4	36.8	38.0
1999.....	1.2	2.5	5.5	9.1	14.2	20.8	27.9	36.3	36.8

Non-Hispanic 4/, male

2007.....	1.2	2.9	4.7	9.0	14.3	23.4	34.9	44.4	50.8
2006.....	1.2	2.9	5.2	9.4	14.4	23.1	33.6	44.7	46.6
2005.....	1.3	2.7	5.4	9.6	16.5	24.3	34.1	45.2	46.9
2004.....	1.2	2.8	5.3	9.2	15.6	23.5	35.4	43.2	45.9
2003.....	1.4	3.1	5.4	10.3	16.3	24.9	34.1	44.2	51.2
2002.....	1.5	3.0	5.5	10.3	16.8	26.3	35.5	46.8	50.6
2001.....	1.6	2.8	6.1	10.2	16.8	25.3	35.6	43.4	48.4
2000.....	1.4	3.0	5.0	9.5	17.0	24.5	36.0	46.5	52.5
1999.....	1.4	3.0	6.5	11.1	16.7	24.6	34.6	46.0	52.2

Non-Hispanic 4/, female

2007.....	0.9	1.8	3.4	5.7	9.6	15.5	21.1	27.4	27.7
2006.....	1.0	2.0	3.6	6.2	9.4	15.9	20.7	26.9	28.6
2005.....	0.9	1.7	3.5	7.3	9.9	16.1	21.7	26.8	28.5
2004.....	0.9	1.8	3.9	6.3	9.9	15.6	23.7	28.6	28.3
2003.....	0.9	1.9	3.5	6.0	10.7	17.2	24.8	29.4	30.3
2002.....	1.1	2.0	4.0	6.7	11.6	17.5	22.5	29.5	30.8
2001.....	0.9	1.9	4.3	6.9	11.8	17.9	23.5	29.7	31.5
2000.....	1.0	2.3	4.2	7.6	12.3	17.4	23.1	31.1	32.1
1999.....	1.1	2.1	4.7	7.3	12.1	17.9	23.2	30.7	30.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic White, both sexes

2007 5/.....	0.8	1.9	3.3	6.5	10.6	18.0	25.6	32.5	34.2
2006 5/.....	0.8	2.0	3.8	6.8	10.7	17.7	25.1	32.8	33.3
2005 5/.....	0.9	1.8	3.7	7.7	11.8	18.4	25.8	33.1	33.5
2004 5/.....	0.9	1.9	3.9	7.0	11.4	17.4	27.2	32.6	32.6
2003 5/.....	0.9	2.0	3.9	7.4	12.2	19.2	27.0	33.5	35.3
2002.....	1.0	2.0	4.1	7.6	12.8	20.2	26.8	35.1	35.7
2001.....	1.0	2.0	4.4	7.4	12.9	19.9	27.2	33.1	35.1
2000.....	1.0	2.3	3.9	7.7	13.3	19.2	26.5	35.6	35.2
1999.....	1.1	2.1	4.6	7.8	13.2	19.4	26.0	35.1	35.0

Non-Hispanic White, male

2007 5/.....	0.9	2.5	3.9	8.3	13.4	22.6	33.4	43.1	50.6
2006 5/.....	1.0	2.5	4.6	8.2	13.4	21.3	32.5	43.9	45.9
2005 5/.....	1.0	2.3	4.5	9.0	15.4	22.8	32.7	44.6	46.5
2004 5/.....	1.0	2.3	4.7	8.5	14.4	22.0	34.1	41.5	44.8
2003 5/.....	1.1	2.5	4.8	9.5	15.3	23.9	33.0	42.8	50.5
2002.....	1.3	2.7	4.9	9.5	15.6	25.3	34.6	46.0	50.1
2001.....	1.4	2.5	5.2	9.1	15.5	23.9	34.6	41.6	47.3
2000.....	1.2	2.6	4.3	8.8	15.8	22.9	34.1	45.7	50.9
1999.....	1.3	2.6	5.5	10.0	15.8	22.9	32.9	44.7	50.6

Non-Hispanic White, female

2007 5/.....	0.6	1.4	2.8	4.8	8.1	14.1	19.8	25.8	26.4
2006 5/.....	0.7	1.5	3.1	5.4	8.3	14.7	19.6	25.9	27.5
2005 5/.....	0.7	1.4	3.0	6.4	8.5	14.7	20.7	26.0	27.5
2004 5/.....	0.8	1.5	3.2	5.7	8.8	13.7	22.1	27.1	27.1
2003 5/.....	0.7	1.6	3.0	5.3	9.5	15.4	22.6	27.8	28.7
2002.....	0.8	1.4	3.4	5.8	10.3	16.0	21.2	28.4	29.5
2001.....	0.7	1.6	3.7	5.8	10.6	16.6	22.0	28.0	30.0
2000.....	0.8	2.0	3.5	6.7	11.1	16.2	21.2	29.7	28.9
1999.....	0.9	1.5	3.8	5.8	10.9	16.6	21.1	29.4	28.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic Black, both sexes

2007 5/.....	2.6	5.1	9.6	15.0	24.9	34.0	46.1	57.3	54.0
2006 5/.....	2.7	5.5	9.8	17.6	23.4	34.8	44.4	53.1	52.0
2005 5/.....	2.4	5.2	10.0	16.5	25.9	35.8	44.7	51.7	50.2
2004 5/.....	2.1	5.2	10.5	15.5	24.7	39.0	49.4	60.5	54.7
2003 5/.....	2.9	6.1	9.6	15.6	26.1	38.1	52.7	60.4	60.4
2002.....	3.0	5.7	10.9	16.6	26.9	38.0	45.5	55.0	55.7
2001.....	2.7	4.8	12.1	19.3	26.2	38.8	47.7	61.8	59.3
2000.....	3.0	6.5	11.9	16.9	27.9	38.0	53.8	58.3	72.8
1999.....	2.5	6.8	13.9	20.5	24.6	38.1	51.6	56.9	63.4

Non-Hispanic Black, male

2007 5/.....	2.7	6.1	10.6	17.7	26.7	38.3	59.7	74.2	65.4
2006 5/.....	3.2	6.4	12.0	23.0	27.8	47.8	55.2	68.8	66.0
2005 5/.....	3.1	6.4	13.2	17.8	30.9	43.6	59.2	68.4	62.8
2004 5/.....	2.2	6.1	11.6	18.7	30.9	45.2	58.8	75.6	72.1
2003 5/.....	3.3	7.8	12.4	20.2	30.1	40.4	57.5	74.7	74.6
2002.....	3.7	6.1	12.6	20.4	32.0	45.2	53.5	71.0	71.1
2001.....	3.5	5.4	15.0	23.0	31.0	45.4	56.9	81.8	72.8
2000.....	3.1	7.4	12.7	18.7	31.7	48.5	67.4	70.4	81.0
1999.....	2.6	6.7	16.4	22.2	26.6	47.4	59.5	75.3	80.8

Non-Hispanic Black, female

2007 5/.....	2.5	4.3	8.7	12.9	23.6	31.1	37.8	48.4	49.2
2006 5/.....	2.3	4.8	8.0	13.3	20.1	25.8	37.9	44.8	46.3
2005 5/.....	1.7	4.2	7.4	15.5	22.3	30.4	36.1	42.9	45.2
2004 5/.....	2.0	4.4	9.6	13.0	20.2	34.7	43.9	52.5	47.8
2003 5/.....	2.5	4.6	7.3	11.9	23.1	36.6	49.9	52.9	54.9
2002.....	2.5	5.4	9.4	13.7	23.2	33.3	40.6	46.7	49.7
2001.....	2.0	4.4	9.8	16.4	22.7	34.4	42.2	51.6	54.3
2000.....	2.9	5.7	11.2	15.4	25.2	31.1	45.6	52.3	69.8
1999.....	2.3	6.8	11.8	19.1	23.2	31.9	46.9	47.9	56.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes

2007.....	9.3	15.7	26.9	41.7	61.3	84.4	122.1	162.2	206.7
2006.....	9.4	16.5	26.9	42.5	62.5	90.2	122.7	162.9	210.1
2005.....	9.9	16.6	27.4	43.5	62.7	90.1	121.9	167.4	212.9
2004.....	9.6	16.2	27.6	43.3	64.6	91.1	126.4	165.2	213.1
2003.....	10.3	17.7	28.1	44.4	64.6	92.9	131.5	167.7	224.0
2002.....	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001.....	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000.....	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
1999.....	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4

All origins 3/, male

2007.....	10.2	18.7	32.2	50.2	73.7	103.5	149.8	208.1	260.9
2006.....	10.5	18.5	31.5	51.7	74.1	107.7	150.6	207.3	259.6
2005.....	11.2	19.1	32.8	52.3	74.7	109.2	148.5	208.5	263.5
2004.....	10.6	18.8	32.5	51.4	78.8	109.3	157.8	207.2	260.6
2003.....	11.9	20.1	33.0	52.9	78.3	111.3	162.7	208.2	275.0
2002.....	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001.....	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000.....	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
1999.....	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9

All origins 3/, female

2007.....	8.5	12.9	22.0	33.8	50.5	68.8	101.6	133.3	180.9
2006.....	8.3	14.5	22.5	34.2	52.4	76.0	102.3	135.1	186.9
2005.....	8.7	14.2	22.4	35.6	52.1	74.7	102.5	141.9	189.6
2004.....	8.6	13.7	23.0	35.9	52.3	76.5	103.8	139.3	191.8
2003.....	8.7	15.5	23.6	36.7	52.8	78.1	109.2	142.9	201.4
2002.....	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001.....	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000.....	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
1999.....	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Hispanic, both sexes

2007	6.8	12.7	18.2	29.0	45.0	53.8	84.9	111.3	136.2
2006	7.5	11.9	18.6	32.7	42.5	69.8	88.9	110.9	145.5
2005	6.1	10.5	17.7	29.8	45.2	69.4	93.5	132.7	139.5
2004	6.4	12.0	17.8	29.5	43.6	64.4	93.5	126.9	140.8
2003	7.1	13.3	18.6	30.7	45.4	75.5	97.3	129.1	168.9
2002	6.9	12.8	20.4	32.3	48.4	75.2	102.6	129.7	164.1
2001	7.4	12.9	19.0	35.4	46.8	78.2	98.3	133.1	193.8
2000	6.6	12.6	23.5	30.5	52.9	78.2	109.3	152.6	192.4
1999	8.6	13.2	23.7	35.2	56.2	75.6	108.5	147.4	202.1

Hispanic, male

2007	7.1	14.8	20.5	32.3	52.7	64.5	104.1	117.0	157.5
2006	7.6	11.5	21.5	35.2	50.5	89.6	103.5	130.2	163.5
2005	5.7	12.2	21.9	36.3	55.2	83.0	105.9	168.1	159.4
2004	6.0	13.8	19.7	33.5	46.9	76.2	115.8	156.7	147.1
2003	9.0	14.2	21.2	36.9	54.9	92.2	117.9	143.5	194.0
2002	7.3	14.5	25.1	35.9	55.5	86.9	127.3	159.0	204.1
2001	8.1	14.3	23.0	39.9	50.0	90.6	130.3	161.0	213.3
2000	6.8	13.2	24.8	35.1	60.8	90.4	138.0	179.6	231.8
1999	9.0	13.9	26.6	42.0	62.2	96.7	124.0	187.1	235.1

Hispanic, female

2007	6.5	10.5	16.0	26.0	38.5	45.3	71.0	107.6	124.7
2006	7.5	12.3	15.8	30.5	35.8	54.3	78.3	98.1	136.0
2005	6.6	8.8	13.8	24.1	36.9	58.9	84.6	109.2	129.0
2004	6.9	10.3	15.9	26.0	40.9	55.3	77.7	107.2	137.6
2003	5.2	12.6	16.2	25.3	37.6	62.6	82.8	119.8	156.3
2002	6.5	11.2	16.0	29.1	42.6	66.2	85.0	111.0	144.2
2001	6.8	11.7	15.4	31.5	44.2	68.8	75.5	115.8	184.3
2000	6.5	12.1	22.4	26.6	46.5	68.9	88.9	136.2	173.1
1999	8.2	12.6	21.1	29.3	51.3	59.8	97.5	123.8	186.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 4/, both sexes

2007.....	9.6	16.0	27.6	42.7	62.6	86.7	124.5	165.2	210.3
2006.....	9.6	16.9	27.6	43.2	64.0	91.6	124.8	165.7	213.1
2005.....	10.4	17.2	28.2	44.6	63.9	91.5	123.5	169.2	216.2
2004.....	9.9	16.6	28.3	44.2	66.2	92.8	128.1	166.8	215.9
2003.....	10.6	18.1	28.8	45.3	65.9	93.8	133.2	169.3	225.8
2002.....	10.7	17.9	29.4	47.5	67.0	98.2	132.1	170.1	232.1
2001.....	10.8	18.4	30.9	47.2	68.5	98.8	134.8	172.0	235.3
2000.....	10.9	18.5	30.8	49.3	72.6	101.4	136.7	177.9	241.0
1999.....	11.7	19.5	33.0	53.6	77.3	109.2	143.5	189.5	251.6

Non-Hispanic 4/, male

2007.....	10.6	19.1	33.2	51.6	75.3	106.2	152.8	213.4	266.7
2006.....	10.8	19.3	32.3	52.9	75.8	108.8	153.4	211.6	264.6
2005.....	11.8	19.7	33.7	53.4	76.1	110.9	151.0	210.5	268.7
2004.....	11.1	19.2	33.4	52.6	81.0	111.2	160.1	209.6	265.7
2003.....	12.1	20.5	33.8	53.9	79.8	112.3	165.0	211.0	278.2
2002.....	12.1	21.3	33.6	57.8	81.3	120.7	163.2	209.4	294.0
2001.....	12.4	22.0	35.7	56.3	81.2	121.3	163.7	212.5	297.9
2000.....	12.5	21.2	35.8	58.6	87.5	123.8	168.2	222.5	303.0
1999.....	13.5	22.7	39.0	63.9	92.2	134.2	175.7	233.9	321.9

Non-Hispanic 4/, female

2007.....	8.7	13.1	22.4	34.5	51.4	70.6	103.7	134.8	183.7
2006.....	8.4	14.7	23.1	34.4	53.7	77.6	103.8	137.1	189.1
2005.....	9.0	14.8	23.1	36.5	53.2	75.8	103.6	143.6	192.2
2004.....	8.8	14.1	23.6	36.6	53.2	77.8	105.2	140.6	193.7
2003.....	9.1	15.7	24.1	37.5	53.9	79.0	110.5	143.8	202.7
2002.....	9.4	14.6	25.5	38.1	54.6	80.2	110.2	146.2	205.4
2001.....	9.3	15.0	26.4	38.8	57.5	81.0	114.6	147.7	209.3
2000.....	9.3	16.0	26.1	40.9	59.7	83.6	114.9	151.7	215.9
1999.....	9.9	16.5	27.5	44.2	64.5	89.6	121.1	163.5	223.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic White, both sexes

2007 5/.....	9.3	15.1	26.8	41.8	62.3	87.7	126.9	168.1	216.2
2006 5/.....	9.2	16.2	26.8	42.5	63.7	93.1	126.4	169.4	217.1
2005 5/.....	9.8	16.4	27.4	43.8	63.6	92.3	125.5	171.8	220.9
2004 5/.....	9.3	15.7	27.3	43.5	65.5	93.9	129.6	168.7	220.6
2003 5/.....	10.0	17.3	28.0	44.9	66.0	94.4	134.5	171.7	229.9
2002.....	10.1	17.1	28.7	46.6	66.5	98.0	132.8	171.4	235.2
2001.....	10.1	17.5	29.7	45.9	67.7	98.1	135.5	172.5	238.9
2000.....	10.2	17.5	30.1	48.3	72.4	100.8	136.2	179.9	244.9
1999.....	11.0	18.6	31.6	52.3	77.5	108.5	143.3	190.3	253.6

Non-Hispanic White, male

2007 5/.....	10.0	18.2	32.2	50.5	74.5	107.9	156.3	218.1	276.0
2006 5/.....	10.3	18.2	31.4	52.5	76.0	110.8	155.3	216.7	271.8
2005 5/.....	11.1	18.9	32.5	52.4	75.1	111.1	154.2	213.6	276.1
2004 5/.....	10.3	17.9	31.8	51.3	79.8	111.9	162.3	213.2	272.4
2003 5/.....	11.2	19.6	32.5	53.2	79.4	112.5	166.5	215.2	284.6
2002.....	11.4	20.4	32.5	56.5	81.6	120.4	163.5	212.7	300.2
2001.....	11.6	20.5	33.8	54.3	79.8	120.8	164.4	213.7	302.9
2000.....	11.5	20.0	34.7	56.1	87.6	123.1	166.3	226.2	309.5
1999.....	12.4	21.3	37.2	62.1	91.7	133.2	176.2	236.1	326.9

Non-Hispanic White, female

2007 5/.....	8.5	12.1	21.7	33.6	51.3	70.8	104.7	136.3	187.9
2006 5/.....	8.1	14.1	22.4	33.1	52.8	78.4	104.8	139.7	191.5
2005 5/.....	8.6	13.9	22.5	35.8	53.3	76.7	104.2	145.6	195.6
2004 5/.....	8.3	13.5	23.0	36.2	52.8	79.0	105.7	141.2	197.4
2003 5/.....	8.7	15.0	23.6	37.2	54.0	79.6	111.3	144.9	205.8
2002.....	8.8	13.9	25.0	37.4	53.2	79.7	110.7	146.2	207.0
2001.....	8.6	14.4	25.8	38.1	57.1	79.6	115.1	147.6	212.3
2000.....	8.8	14.9	25.7	41.0	59.1	82.9	115.1	152.4	218.9
1999.....	9.5	16.0	26.2	43.3	65.0	88.8	120.2	163.2	224.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic Black, both sexes

2007 5/.....	13.9	24.6	38.6	58.7	77.5	91.9	124.6	152.3	173.6
2006 5/.....	13.4	25.2	39.8	59.0	79.0	98.0	130.3	152.5	188.7
2005 5/.....	15.3	25.5	40.3	57.3	77.4	99.5	129.4	171.4	184.6
2004 5/.....	15.0	25.6	41.4	58.4	85.4	100.2	134.7	170.1	188.2
2003 5/.....	16.3	26.9	41.9	58.6	78.5	104.5	141.2	167.7	206.4
2002.....	16.1	25.8	42.4	64.0	83.8	115.8	149.2	175.4	223.2
2001.....	17.2	28.4	45.2	65.3	86.5	120.7	143.0	182.4	218.5
2000.....	17.1	29.5	42.1	65.9	86.8	118.5	156.7	174.4	211.1
1999.....	18.7	29.8	49.2	74.4	86.6	129.5	159.2	194.6	244.8

Non-Hispanic Black, male

2007 5/.....	16.7	29.5	48.3	75.7	98.8	108.4	150.0	192.1	210.3
2006 5/.....	16.2	31.1	48.6	71.5	92.8	118.4	165.3	203.7	216.2
2005 5/.....	18.6	28.9	51.6	73.1	100.9	128.7	146.9	220.2	221.7
2004 5/.....	17.6	33.5	53.4	75.5	109.0	129.7	172.1	207.1	223.5
2003 5/.....	20.2	31.6	54.4	72.8	100.2	131.4	179.0	207.5	239.7
2002.....	18.7	31.6	51.4	82.3	99.1	145.7	190.1	210.6	265.8
2001.....	19.4	36.6	58.6	82.6	111.8	150.5	181.6	218.3	287.6
2000.....	21.4	33.9	52.1	90.4	103.1	150.3	211.9	212.0	253.6
1999.....	23.8	37.4	62.9	95.8	111.0	165.1	193.8	235.6	293.6

Non-Hispanic Black, female

2007 5/.....	11.5	20.4	30.7	45.5	61.9	80.6	109.4	131.5	158.4
2006 5/.....	10.9	20.2	32.6	49.3	68.8	84.0	109.4	125.5	177.5
2005 5/.....	12.4	22.6	31.0	45.1	60.0	79.6	118.9	145.7	169.7
2004 5/.....	12.8	18.9	31.6	45.1	67.9	80.2	112.6	150.7	174.3
2003 5/.....	12.9	22.9	31.6	47.5	62.3	86.3	118.7	146.9	193.6
2002.....	13.9	20.9	35.0	49.6	72.5	95.7	124.7	157.1	206.9
2001.....	15.2	21.5	34.2	51.8	67.9	100.9	120.0	164.1	192.9
2000.....	13.5	25.7	33.8	46.6	74.7	97.3	123.4	155.5	195.4
1999.....	14.4	23.3	38.1	57.7	68.5	105.9	138.3	174.3	226.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes

2007.....	1.3	2.0	3.5	5.4	9.4	16.2	28.2	46.5	76.4
2006.....	1.5	2.2	3.1	5.6	9.0	16.8	28.8	46.3	78.0
2005.....	1.4	2.1	3.3	5.7	9.5	16.3	27.9	44.8	76.7
2004.....	1.4	2.0	3.5	5.8	9.3	16.7	27.3	47.2	77.7
2003.....	1.4	2.0	3.2	6.0	9.8	16.3	28.5	47.4	80.5
2002.....	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001.....	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000.....	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
1999.....	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9

All origins 3/, male

2007.....	1.4	2.2	4.3	6.6	11.6	19.9	37.4	63.9	95.7
2006.....	1.6	2.5	3.8	6.3	11.1	21.5	38.2	60.8	92.3
2005.....	1.5	2.4	3.6	7.2	12.1	20.3	36.3	59.5	92.7
2004.....	1.4	2.3	4.0	7.1	11.7	20.5	34.8	61.3	98.9
2003.....	1.4	2.3	3.6	7.0	11.6	21.2	35.9	61.5	101.7
2002.....	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001.....	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000.....	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
1999.....	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2

All origins 3/, female

2007.....	1.1	1.7	2.7	4.3	7.4	13.2	21.5	35.6	67.2
2006.....	1.4	1.9	2.5	4.9	7.2	13.0	21.8	37.3	71.4
2005.....	1.3	1.8	3.1	4.3	7.3	13.0	21.8	35.6	69.4
2004.....	1.3	1.7	3.0	4.6	7.1	13.7	21.9	38.5	68.1
2003.....	1.3	1.7	2.8	5.1	8.3	12.4	23.2	38.7	71.1
2002.....	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001.....	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000.....	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
1999.....	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, both sexes

2007	0.9	1.6	2.6	3.7	5.4	10.0	19.0	33.2	46.5
2006	1.1	1.7	2.3	5.3	6.8	11.5	19.2	33.7	48.5
2005	*	1.9	2.4	4.2	6.0	10.8	17.2	26.5	34.2
2004	1.3	1.4	2.4	5.8	5.0	13.3	17.9	31.9	48.5
2003	1.1	*	2.2	4.2	7.2	10.6	22.2	39.8	48.2
2002	1.4	1.5	1.8	4.8	6.2	12.2	17.3	37.0	48.8
2001	1.1	1.9	2.4	5.3	7.5	10.5	19.0	30.3	57.5
2000	1.1	1.7	2.4	4.0	6.7	12.2	20.5	31.2	70.3
1999	*	1.9	3.4	4.0	6.7	15.3	18.4	31.7	51.9

Hispanic, male

2007	*	*	2.6	5.1	7.6	9.2	21.3	31.8	45.7
2006	*	2.5	2.7	6.1	7.6	16.0	24.7	39.3	53.0
2005	*	2.4	*	5.7	8.9	13.1	20.5	38.5	37.4
2004	*	*	*	6.3	*	16.6	21.9	37.7	57.4
2003	*	*	*	5.4	8.1	12.4	26.1	46.8	52.3
2002	*	*	*	5.4	7.8	13.6	19.8	52.6	56.9
2001	*	*	*	6.8	7.4	13.3	19.8	49.3	70.5
2000	*	*	*	*	*	14.6	26.6	*	86.7
1999	*	*	*	*	*	17.3	18.7	36.2	49.6

Hispanic, female

2007	*	*	2.6	*	*	10.6	17.3	34.1	46.9
2006	*	*	*	4.5	6.2	8.0	15.3	30.0	46.1
2005	*	*	*	*	*	8.9	14.7	18.6	32.6
2004	*	*	*	5.3	5.4	10.9	15.1	28.0	43.9
2003	*	*	*	*	6.4	9.2	19.4	35.2	46.1
2002	*	*	*	*	*	11.2	15.5	27.0	44.8
2001	*	*	*	*	7.5	8.4	18.4	18.5	51.2
2000	*	*	*	*	7.2	10.3	16.2	34.0	62.3
1999	*	*	*	*	8.1	13.8	18.2	29.1	53.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 4/, both sexes

2007.....	1.3	2.0	3.6	5.5	9.7	16.7	28.8	47.3	77.9
2006.....	1.5	2.3	3.2	5.6	9.2	17.1	29.4	46.9	79.5
2005.....	1.4	2.1	3.4	5.8	9.8	16.6	28.5	45.7	78.6
2004.....	1.4	2.1	3.6	5.8	9.5	16.9	27.8	47.9	78.9
2003.....	1.4	2.1	3.3	6.2	10.0	16.6	28.8	47.7	81.8
2002.....	1.5	2.1	3.9	5.5	9.9	16.6	29.3	46.6	79.8
2001.....	1.5	2.3	4.0	6.3	10.7	18.3	30.0	48.3	83.5
2000.....	1.5	2.3	3.9	6.8	10.8	17.6	29.2	51.2	85.2
1999.....	1.5	2.3	3.5	6.4	11.0	18.8	29.7	48.0	83.8

Non-Hispanic 4/, male

2007.....	1.4	2.3	4.4	6.7	11.9	20.7	38.4	65.8	98.5
2006.....	1.6	2.5	3.8	6.3	11.4	21.8	39.1	61.9	94.5
2005.....	1.6	2.4	3.7	7.3	12.3	20.8	37.3	60.5	95.4
2004.....	1.5	2.4	4.1	7.2	12.2	20.7	35.4	62.5	101.0
2003.....	1.4	2.4	3.7	7.1	11.8	21.6	36.4	62.2	103.9
2002.....	1.6	2.2	4.7	6.4	12.5	20.4	38.0	59.2	102.9
2001.....	1.7	2.4	4.2	7.5	13.1	23.0	40.1	62.7	109.0
2000.....	1.7	2.7	4.2	7.8	13.8	21.6	37.2	66.6	109.4
1999.....	1.5	2.6	4.1	8.1	13.9	24.3	37.9	61.8	111.4

Non-Hispanic 4/, female

2007.....	1.2	1.7	2.7	4.4	7.7	13.4	21.7	35.6	68.2
2006.....	1.5	2.0	2.6	5.0	7.3	13.3	22.3	37.6	72.5
2005.....	1.3	1.8	3.1	4.4	7.6	13.3	22.2	36.5	71.0
2004.....	1.4	1.7	3.1	4.6	7.2	13.8	22.2	38.9	69.0
2003.....	1.3	1.8	2.9	5.2	8.4	12.6	23.4	38.9	72.1
2002.....	1.5	2.0	3.1	4.7	7.7	13.5	23.2	38.9	69.8
2001.....	1.3	2.1	3.8	5.3	8.6	14.6	22.9	39.7	72.9
2000.....	1.3	1.9	3.6	5.8	8.2	14.6	23.7	42.1	75.4
1999.....	1.4	2.0	3.0	4.9	8.5	14.5	24.0	39.9	72.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic White, both sexes

2007 5/.....	1.2	2.0	3.6	5.4	9.7	17.0	29.8	48.8	81.2
2006 5/.....	1.4	2.3	3.2	5.4	9.4	17.7	30.1	48.4	81.6
2005 5/.....	1.3	2.1	3.4	5.6	10.0	17.0	29.4	47.3	81.7
2004 5/.....	1.4	1.9	3.4	5.7	9.7	17.3	28.3	49.3	81.6
2003 5/.....	1.3	2.1	3.2	6.3	10.0	17.1	29.9	49.1	84.4
2002.....	1.6	2.0	3.8	5.6	10.1	17.0	30.2	47.4	81.8
2001.....	1.5	2.3	4.0	6.3	11.0	18.6	30.9	49.4	84.8
2000.....	1.4	2.3	4.0	6.8	11.0	17.6	29.8	52.8	87.6
1999.....	1.3	2.2	3.4	6.4	10.9	19.2	30.1	49.2	86.3

Non-Hispanic White, male

2007 5/.....	1.4	2.3	4.4	6.5	11.8	20.9	39.8	68.0	102.5
2006 5/.....	1.5	2.6	3.9	6.2	11.6	22.7	40.4	63.7	96.0
2005 5/.....	1.4	2.4	3.7	7.2	12.6	21.5	38.6	63.0	100.0
2004 5/.....	1.4	2.3	3.9	7.0	12.6	21.3	36.1	64.5	105.6
2003 5/.....	1.4	2.4	3.5	7.2	12.0	22.3	37.9	64.0	107.9
2002.....	1.7	2.2	4.5	6.5	12.8	21.1	39.2	60.2	106.0
2001.....	1.8	2.5	4.3	7.4	13.5	23.6	41.3	63.9	111.4
2000.....	1.6	2.7	4.4	7.5	13.9	21.3	38.0	69.4	113.1
1999.....	1.5	2.6	3.9	8.2	13.9	24.9	38.6	63.8	116.8

Non-Hispanic White, female

2007 5/.....	1.1	1.7	2.8	4.4	7.9	13.6	22.2	36.5	71.1
2006 5/.....	1.3	1.9	2.6	4.7	7.5	13.5	22.4	38.9	74.9
2005 5/.....	1.2	1.8	3.1	4.2	7.6	13.2	22.6	37.4	73.4
2004 5/.....	1.3	1.6	2.9	4.5	7.1	14.0	22.5	39.9	70.9
2003 5/.....	1.2	1.8	2.9	5.4	8.1	12.9	24.2	39.9	74.0
2002.....	1.4	1.8	3.1	4.8	7.8	13.6	23.8	39.6	71.4
2001.....	1.3	2.1	3.8	5.3	8.7	14.5	23.5	40.6	73.7
2000.....	1.2	1.8	3.6	6.1	8.5	14.6	24.2	43.0	77.3
1999.....	1.2	1.9	3.0	4.7	8.3	14.7	24.2	40.6	74.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic Black, both sexes

2007 5/.....	1.8	2.4	4.2	7.0	10.5	18.3	24.7	38.7	50.6
2006 5/.....	2.4	2.8	3.9	8.1	9.3	15.1	26.7	40.0	63.4
2005 5/.....	2.5	2.5	4.0	7.7	10.2	16.8	24.3	34.3	55.7
2004 5/.....	1.7	3.1	5.7	7.2	10.4	16.4	26.5	40.0	57.9
2003 5/.....	2.3	2.8	4.3	6.7	11.5	15.6	22.5	40.2	58.9
2002.....	2.0	3.1	5.3	6.0	10.8	15.4	24.7	42.4	64.4
2001.....	1.5	2.8	4.6	7.7	11.0	18.9	25.0	42.8	72.7
2000.....	2.4	3.2	4.2	7.3	10.9	20.3	27.0	38.2	64.2
1999.....	2.3	3.1	5.0	8.0	14.0	18.2	29.4	39.0	62.7

Non-Hispanic Black, male

2007 5/.....	1.9	2.7	5.8	9.4	13.9	23.7	29.2	52.7	63.7
2006 5/.....	2.4	2.5	4.9	8.9	12.3	18.7	33.6	55.5	82.5
2005 5/.....	2.7	2.8	4.3	9.4	11.3	18.2	28.5	43.9	56.1
2004 5/.....	1.7	3.6	6.8	9.6	12.7	19.6	35.0	47.7	62.0
2003 5/.....	2.1	3.0	5.6	8.5	12.7	21.6	26.6	52.8	68.2
2002.....	*	2.9	7.7	6.7	12.8	16.4	30.0	54.6	81.0
2001.....	*	2.5	4.5	9.4	13.3	23.4	32.2	59.7	87.8
2000.....	2.2	3.7	4.2	10.4	16.0	23.9	34.7	39.5	76.3
1999.....	*	3.2	6.3	8.5	17.3	21.8	36.3	43.4	64.2

Non-Hispanic Black, female

2007 5/.....	1.7	2.1	2.9	5.1	8.0	14.6	21.9	31.3	45.2
2006 5/.....	2.3	3.1	3.1	7.4	7.1	12.6	22.5	31.8	55.7
2005 5/.....	2.3	2.3	3.8	6.4	9.4	15.8	21.9	29.2	55.5
2004 5/.....	1.6	2.6	4.7	5.2	8.7	14.2	21.5	35.9	56.2
2003 5/.....	2.4	2.6	3.2	5.3	10.7	11.6	20.0	33.6	55.3
2002.....	2.3	3.3	3.4	5.4	9.3	14.7	21.6	36.1	58.1
2001.....	*	3.0	4.6	6.4	9.2	15.8	20.7	34.1	67.2
2000.....	2.5	2.8	4.2	4.9	7.1	17.9	22.4	37.6	59.7
1999.....	2.6	3.1	4.0	7.7	11.6	15.9	25.3	36.9	62.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

All origins 3/, both sexes

2007.....	0.6	0.6	0.8	1.2	2.1	3.9	6.5	13.2	36.1
2006.....	0.5	0.5	0.8	1.2	1.4	3.0	5.1	10.5	30.0
2005.....	0.6	0.6	0.8	1.2	2.0	3.5	6.3	12.8	36.6
2004.....	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
2003.....	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8

All origins 3/, male

2007.....	0.5	0.7	0.8	1.2	2.2	3.9	7.1	14.2	34.3
2006.....	0.5	0.5	0.8	1.1	1.5	2.8	5.4	11.6	29.6
2005.....	0.5	0.5	0.7	1.3	1.9	3.5	6.4	14.1	33.5
2004.....	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
2003.....	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4

All origins 3/, female

2007.....	0.6	0.5	0.8	1.3	1.9	3.9	6.0	12.5	36.9
2006.....	0.6	0.6	0.8	1.2	1.3	3.2	5.0	9.8	30.2
2005.....	0.7	0.7	0.9	1.2	2.0	3.5	6.3	12.1	38.0
2004.....	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
2003.....	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

Non-Hispanic 4/, both sexes

2007.....	0.6	0.6	0.8	1.2	2.1	4.0	6.6	13.1	37.0
2006.....	0.6	0.6	0.8	1.2	1.4	3.1	5.1	10.7	30.3
2005.....	0.6	0.7	0.9	1.3	2.0	3.5	6.4	13.1	37.2
2004.....	0.6	0.6	0.9	1.4	1.8	3.6	6.0	12.3	38.5
2003.....	0.7	0.6	0.7	1.3	1.9	3.5	6.1	12.7	41.2
2002.....	0.6	0.7	0.9	1.2	1.9	3.7	6.5	12.9	40.5
2001.....	0.6	0.7	0.8	1.2	1.9	3.6	7.1	13.7	42.5
2000.....	0.6	0.7	1.0	1.2	2.2	3.4	6.6	13.9	43.2
1999.....	0.7	0.6	0.8	1.2	1.8	3.4	6.8	13.7	44.0

Non-Hispanic 4/, male

2007.....	0.5	0.7	0.8	1.2	2.2	4.1	7.2	14.1	35.0
2006.....	0.5	0.5	0.8	1.2	1.6	2.8	5.3	12.0	30.2
2005.....	0.5	0.6	0.8	1.3	2.0	3.5	6.3	14.4	34.1
2004.....	0.5	0.6	1.0	1.5	2.0	3.8	6.2	13.0	37.1
2003.....	0.7	0.5	0.8	1.6	2.3	3.9	6.6	13.4	38.3
2002.....	0.6	0.6	0.8	1.3	1.7	4.3	6.7	12.5	37.6
2001.....	0.6	0.7	0.8	1.1	2.0	3.9	7.1	14.4	40.8
2000.....	0.6	0.7	1.1	1.3	2.2	3.5	6.6	14.4	41.0
1999.....	0.7	0.6	0.7	1.2	2.1	3.2	7.2	14.7	43.3

Non-Hispanic 4/, female

2007.....	0.7	0.5	0.8	1.3	2.1	4.0	6.2	12.4	38.0
2006.....	0.7	0.6	0.8	1.2	1.3	3.3	5.0	9.9	30.4
2005.....	0.7	0.7	1.0	1.3	2.0	3.4	6.4	12.2	38.6
2004.....	0.6	0.6	0.7	1.3	1.7	3.4	5.8	12.0	39.1
2003.....	0.6	0.6	0.7	1.1	1.5	3.2	5.7	12.3	42.4
2002.....	0.7	0.7	0.9	1.1	2.1	3.2	6.3	13.2	41.8
2001.....	0.5	0.7	0.9	1.3	1.9	3.3	7.0	13.3	43.1
2000.....	0.6	0.7	0.9	1.2	2.3	3.4	6.6	13.6	44.1
1999.....	0.7	0.6	0.9	1.2	1.6	3.6	6.5	13.2	44.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.4	0.5	1.0	1.9	3.7	6.4	12.8	37.4
2006 5/.....	0.2	0.3	0.6	0.9	1.3	2.9	4.9	10.6	30.0
2005 5/.....	0.3	0.3	0.7	1.0	1.6	3.1	6.1	12.6	37.2
2004 5/.....	0.3	0.3	0.5	1.1	1.7	3.3	5.7	12.4	38.1
2003 5/.....	0.3	0.3	0.4	1.0	1.6	3.2	5.8	12.5	41.0
2002.....	0.3	0.3	0.6	0.9	1.7	3.3	5.9	12.6	39.9
2001.....	0.2	0.4	0.5	1.1	1.7	3.3	6.7	13.2	42.1
2000.....	0.2	0.4	0.7	0.9	2.0	3.2	6.3	13.8	42.9
1999.....	0.3	0.3	0.4	0.8	1.6	3.1	6.5	13.7	43.2

Non-Hispanic White, male

2007 5/.....	*	0.5	0.5	1.0	2.0	3.9	6.9	13.9	35.2
2006 5/.....	*	*	0.5	0.9	1.5	2.6	5.0	11.9	30.6
2005 5/.....	0.3	0.3	0.6	1.0	1.5	3.3	5.9	13.8	34.3
2004 5/.....	0.3	0.3	0.5	1.2	1.9	3.6	5.7	12.8	37.3
2003 5/.....	0.3	0.3	0.4	1.1	2.0	3.3	6.4	13.5	38.1
2002.....	0.3	0.4	0.6	1.0	1.5	3.6	6.3	11.9	36.3
2001.....	*	0.4	0.5	1.0	1.8	3.4	6.9	14.2	41.2
2000.....	*	0.4	0.8	0.9	2.1	3.3	6.3	14.1	39.9
1999.....	0.4	0.3	*	0.7	1.9	2.8	7.0	14.7	43.4

Non-Hispanic White, female

2007 5/.....	0.3	0.3	0.5	1.1	1.9	3.5	6.0	12.0	38.5
2006 5/.....	0.3	0.3	0.6	1.0	1.1	3.1	4.8	9.8	29.7
2005 5/.....	0.4	0.3	0.7	1.0	1.7	2.9	6.2	11.9	38.5
2004 5/.....	*	*	0.5	1.0	1.5	3.1	5.6	12.1	38.4
2003 5/.....	*	*	0.3	0.8	1.2	3.1	5.4	11.8	42.2
2002.....	0.3	0.3	0.6	0.8	1.8	3.0	5.6	13.0	41.4
2001.....	*	0.4	0.5	1.1	1.7	3.1	6.5	12.7	42.4
2000.....	*	0.4	0.6	0.9	1.8	3.1	6.3	13.5	44.1
1999.....	0.3	*	0.5	1.0	1.4	3.4	6.1	13.1	43.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

Non-Hispanic Black, both sexes

2007 5/.....	3.1	2.3	3.2	2.9	4.8	7.6	9.6	18.4	36.1
2006 5/.....	2.7	2.5	3.0	3.5	3.1	5.7	8.3	13.6	39.3
2005 5/.....	2.6	3.0	2.6	3.7	5.4	7.5	11.1	20.4	43.4
2004 5/.....	2.5	3.1	3.9	4.2	3.6	6.3	10.8	13.4	48.9
2003 5/.....	3.4	2.7	4.2	4.5	4.3	7.3	11.0	19.7	50.3
2002.....	3.2	3.1	3.2	4.2	4.5	8.1	14.0	19.5	52.9
2001.....	3.3	2.9	3.3	2.7	4.0	7.4	12.6	21.5	49.3
2000.....	3.0	3.2	3.0	4.0	4.9	6.3	10.5	18.1	51.8
1999.....	3.3	3.0	4.3	4.3	3.7	6.6	11.4	16.4	53.0

Non-Hispanic Black, male

2007 5/.....	3.1	2.6	3.5	*	5.4	6.5	10.9	20.1	34.0
2006 5/.....	2.5	2.2	3.4	4.0	*	*	9.7	15.4	30.2
2005 5/.....	2.5	2.7	*	*	6.9	*	13.3	23.7	38.1
2004 5/.....	2.1	2.9	5.3	4.4	*	*	13.5	14.7	43.7
2003 5/.....	4.1	2.6	5.1	6.4	5.6	11.1	13.1	18.9	50.1
2002.....	2.7	2.6	3.1	5.3	*	12.0	13.1	23.4	58.0
2001.....	4.0	3.2	3.4	*	*	10.7	12.3	20.4	42.7
2000.....	3.3	3.5	*	*	*	*	11.1	18.9	56.4
1999.....	3.4	3.3	4.4	6.1	*	8.0	10.6	*	44.0

Non-Hispanic Black, female

2007 5/.....	3.0	2.1	2.9	3.1	4.3	8.4	8.8	17.5	36.9
2006 5/.....	2.8	2.8	2.6	3.1	*	5.9	7.4	12.6	43.0
2005 5/.....	2.7	3.2	3.0	3.9	4.4	8.4	9.8	18.6	45.6
2004 5/.....	2.9	3.2	2.8	4.0	*	6.3	9.1	12.7	51.0
2003 5/.....	2.8	2.8	3.4	3.1	*	4.8	9.7	20.1	50.3
2002.....	3.6	3.5	3.4	3.3	4.7	5.5	14.6	17.5	51.0
2001.....	2.7	2.7	3.3	*	4.0	5.3	12.7	22.1	51.8
2000.....	2.8	3.1	3.0	4.0	5.9	6.4	10.2	17.7	50.2
1999.....	3.2	2.8	4.3	*	*	5.7	11.9	16.2	56.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

All origins 3/, both sexes

2007.....	9.5	17.1	27.9	43.0	64.0	95.7	136.7	196.1	276.2
2006.....	9.6	17.1	28.7	46.4	68.3	98.3	141.9	199.3	285.2
2005.....	9.9	17.3	29.7	47.3	71.7	104.9	151.4	211.2	312.1
2004.....	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
2003.....	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2

All origins 3/, male

2007.....	12.2	20.8	34.4	52.0	78.0	112.9	155.8	228.2	307.4
2006.....	12.3	21.3	34.5	56.8	80.7	115.9	167.5	232.2	321.0
2005.....	12.7	21.0	36.3	56.6	82.4	121.2	174.0	244.1	344.4
2004.....	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
2003.....	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7

All origins 3/, female

2007.....	6.8	13.6	21.8	34.7	51.7	81.7	122.6	175.9	261.4
2006.....	6.9	13.2	23.3	36.9	57.5	84.0	123.1	178.7	268.4
2005.....	7.2	13.8	23.4	38.9	62.3	91.6	134.9	190.8	297.2
2004.....	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
2003.....	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

Hispanic, both sexes

2007	8.7	19.2	32.2	55.8	89.8	130.1	202.8	276.7	337.5
2006	9.4	18.1	34.5	54.1	96.3	146.1	205.6	264.1	369.7
2005	10.8	20.0	36.6	67.1	101.3	158.9	240.2	308.3	404.7
2004	10.3	17.2	36.7	64.7	102.6	153.8	229.6	303.6	365.8
2003	10.1	22.7	40.5	72.1	105.3	165.4	238.3	327.1	428.1
2002	9.2	20.4	38.0	71.3	108.2	168.0	257.0	333.3	437.9
2001	11.9	18.8	38.4	72.8	121.1	166.3	254.5	322.3	489.6
2000	9.8	19.5	45.4	68.5	114.1	190.3	250.1	330.3	465.1
1999	10.1	19.0	40.4	77.1	124.6	181.4	252.4	344.8	493.9

Hispanic, male

2007	11.6	21.6	39.6	67.6	111.8	148.1	218.1	286.7	324.2
2006	11.8	22.0	41.8	62.0	110.3	169.7	225.1	309.5	381.2
2005	13.2	23.4	42.8	81.1	114.2	183.9	266.3	336.2	404.7
2004	11.4	22.1	41.6	79.3	110.2	166.4	249.4	328.4	325.0
2003	11.6	26.8	45.2	87.1	116.7	185.3	254.6	370.1	426.4
2002	10.2	25.4	43.3	80.8	110.7	186.5	282.8	350.0	425.0
2001	14.4	23.8	43.4	79.6	132.5	175.7	262.6	328.5	422.8
2000	11.1	21.4	52.8	80.3	126.0	206.1	267.2	363.7	449.4
1999	12.5	23.5	46.7	82.8	127.9	192.5	262.1	355.2	468.0

Hispanic, female

2007	5.7	16.8	25.1	45.2	71.3	115.9	191.7	270.0	344.8
2006	6.8	14.3	27.6	47.1	84.5	127.7	191.5	233.8	363.7
2005	8.3	16.7	30.8	54.6	90.5	139.4	221.5	289.8	404.7
2004	9.2	12.4	32.1	51.9	96.3	144.1	215.6	287.2	386.8
2003	8.6	18.7	36.1	59.0	96.0	150.1	226.8	299.1	428.9
2002	8.2	15.4	33.1	63.1	106.2	153.8	238.6	322.7	444.3
2001	9.4	14.0	33.8	67.0	111.9	159.1	248.9	318.5	522.2
2000	8.6	17.7	38.7	58.3	104.5	178.2	237.9	310.0	472.8
1999	7.6	14.7	34.6	72.1	121.9	173.1	245.6	338.5	506.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

Non-Hispanic 4/, both sexes

2007.....	9.5	16.8	27.4	41.8	61.8	92.9	131.8	191.1	272.7
2006.....	9.6	17.0	28.1	45.6	65.9	94.5	137.4	195.2	280.5
2005.....	9.8	17.0	29.0	45.6	69.2	100.8	145.4	205.7	307.2
2004.....	10.0	17.1	28.6	45.6	68.8	102.1	144.6	208.1	303.5
2003.....	10.3	17.4	29.5	47.0	72.5	104.8	150.2	210.7	312.0
2002.....	9.9	17.8	29.3	45.4	73.0	104.7	151.8	212.2	315.0
2001.....	9.7	17.6	28.7	46.4	71.9	104.7	150.1	213.9	314.8
2000.....	9.5	17.0	28.3	46.4	71.1	103.6	151.7	209.0	313.4
1999.....	9.8	16.2	28.6	46.7	72.3	105.0	148.7	209.5	310.0

Non-Hispanic 4/, male

2007.....	12.2	20.6	33.8	50.6	75.2	110.0	151.2	224.3	306.4
2006.....	12.3	21.1	33.7	56.1	78.1	111.8	163.5	227.4	317.0
2005.....	12.6	20.7	35.6	54.5	79.8	116.6	167.8	238.6	340.9
2004.....	12.5	21.0	33.7	54.7	80.5	117.8	167.0	242.3	333.5
2003.....	12.7	20.8	34.5	55.4	84.4	120.2	174.2	243.1	335.9
2002.....	12.0	21.1	35.4	53.3	82.0	116.7	169.5	237.7	344.0
2001.....	11.7	21.0	33.6	52.4	81.0	117.1	166.4	238.1	338.9
2000.....	11.4	19.5	31.8	51.9	77.6	114.4	168.6	237.2	339.7
1999.....	12.3	18.9	32.5	51.6	79.4	115.2	167.9	236.6	339.5

Non-Hispanic 4/, female

2007.....	6.9	13.2	21.4	33.7	50.0	78.9	117.5	170.2	256.8
2006.....	6.9	13.0	22.8	35.9	55.2	80.5	118.2	175.1	263.5
2005.....	7.1	13.4	22.6	37.5	59.8	88.0	129.1	185.3	291.9
2004.....	7.5	13.4	23.8	37.4	58.5	89.4	128.5	187.2	290.1
2003.....	8.0	14.2	24.8	39.4	62.2	92.4	133.1	190.9	301.5
2002.....	7.9	14.6	23.7	38.2	65.1	95.2	139.4	196.7	302.5
2001.....	7.7	14.4	24.1	41.0	64.1	94.7	138.7	199.4	304.7
2000.....	7.7	14.6	25.1	41.4	65.5	95.1	139.9	192.4	302.8
1999.....	7.5	13.7	25.0	42.2	66.1	97.0	135.3	193.6	298.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

Non-Hispanic White, both sexes

2007 5/.....	8.1	14.1	23.0	36.1	53.7	81.9	120.0	175.9	257.0
2006 5/.....	8.1	14.1	23.8	39.0	56.6	83.6	124.4	180.3	264.2
2005 5/.....	8.2	14.2	24.5	39.2	59.5	89.0	132.4	191.8	289.8
2004 5/.....	8.6	14.2	24.1	39.0	59.7	89.7	130.7	193.1	284.5
2003 5/.....	8.6	14.2	24.8	39.8	62.6	92.5	136.6	195.4	293.8
2002.....	8.2	14.7	24.7	38.3	61.9	92.7	138.0	197.8	296.4
2001.....	8.1	14.7	23.9	39.4	62.0	92.7	136.3	199.9	297.4
2000.....	7.8	14.0	23.7	39.7	61.2	91.7	137.5	193.0	295.5
1999.....	8.1	13.2	23.0	38.7	62.2	92.0	135.2	194.3	290.2

Non-Hispanic White, male

2007 5/.....	10.6	17.8	29.0	45.0	68.0	100.4	142.2	213.8	298.1
2006 5/.....	10.7	17.9	28.9	49.6	69.7	101.6	152.9	217.1	306.8
2005 5/.....	10.9	17.6	30.6	48.0	71.0	105.3	156.8	228.9	331.5
2004 5/.....	10.9	18.1	28.9	48.3	71.6	106.9	155.5	232.7	321.7
2003 5/.....	11.0	17.7	30.1	48.1	75.3	109.7	164.0	232.9	327.3
2002.....	10.2	17.9	30.7	46.4	72.5	107.2	160.1	228.4	338.4
2001.....	9.9	17.9	29.0	45.9	72.8	108.6	155.6	226.9	332.1
2000.....	9.5	16.7	27.3	45.7	69.6	105.1	157.9	227.9	330.4
1999.....	10.3	16.0	27.2	44.5	70.9	106.0	158.4	227.5	326.0

Non-Hispanic White, female

2007 5/.....	5.7	10.6	17.2	27.7	40.9	66.5	103.2	151.8	237.5
2006 5/.....	5.6	10.3	18.8	29.1	44.9	68.6	103.1	157.1	244.3
2005 5/.....	5.6	10.9	18.6	31.0	49.1	75.4	114.3	168.7	270.7
2004 5/.....	6.4	10.3	19.6	30.4	49.1	75.5	112.5	168.6	267.8
2003 5/.....	6.2	10.9	19.8	32.1	51.4	78.3	116.7	172.3	279.0
2002.....	6.3	11.6	18.8	30.7	52.5	80.8	122.1	179.0	278.3
2001.....	6.3	11.6	18.9	33.3	52.5	79.7	122.6	183.7	283.0
2000.....	6.1	11.3	20.4	34.2	53.7	80.9	123.2	172.2	281.4
1999.....	5.9	10.5	18.9	33.3	54.5	80.8	119.0	174.7	275.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

Non-Hispanic Black, both sexes

2007 5/.....	19.5	37.0	61.8	95.0	132.6	192.8	260.7	360.1	451.8
2006 5/.....	19.6	38.0	63.0	103.6	148.4	197.1	277.5	371.2	475.9
2005 5/.....	21.2	37.3	65.2	104.5	153.5	214.9	286.7	380.6	513.4
2004 5/.....	19.8	39.7	66.1	103.5	151.1	222.3	300.5	388.9	536.5
2003 5/.....	22.7	40.7	71.3	109.1	159.9	225.4	303.4	394.6	532.2
2002.....	22.0	41.8	69.3	107.3	164.9	225.1	303.3	396.4	544.7
2001.....	21.3	41.0	70.6	106.6	159.4	229.9	307.4	385.5	533.8
2000.....	22.4	41.7	67.7	103.7	160.7	226.7	320.5	409.2	530.1
1999.....	23.1	41.1	74.2	113.7	162.6	232.3	297.1	399.2	530.9

Non-Hispanic Black, male

2007 5/.....	24.8	43.0	74.1	107.9	149.6	210.2	274.9	381.4	404.0
2006 5/.....	24.3	46.3	73.6	121.9	165.6	225.1	303.8	380.6	468.2
2005 5/.....	26.4	43.5	80.0	120.3	164.8	240.6	315.5	395.0	471.0
2004 5/.....	24.9	45.2	76.4	117.4	172.6	241.1	312.9	393.6	508.1
2003 5/.....	26.2	45.0	78.9	123.9	174.9	236.2	311.6	393.2	457.1
2002.....	26.1	47.7	78.6	120.6	168.4	226.5	297.7	390.8	423.4
2001.....	25.8	46.4	77.8	114.5	163.9	223.4	310.5	404.7	452.7
2000.....	26.1	44.4	73.5	110.1	155.4	227.2	314.1	384.5	456.8
1999.....	28.2	44.5	79.6	115.2	167.3	218.8	284.2	379.0	506.3

Non-Hispanic Black, female

2007 5/.....	14.9	31.8	51.6	84.9	120.1	180.8	252.2	348.9	471.6
2006 5/.....	15.5	31.0	54.3	89.3	135.8	177.9	261.7	366.2	479.1
2005 5/.....	16.6	32.0	53.1	92.1	145.1	197.3	269.5	373.0	530.4
2004 5/.....	15.3	35.0	57.5	92.7	135.2	209.6	293.2	386.4	547.7
2003 5/.....	19.7	37.2	65.0	97.6	148.9	218.1	298.5	395.3	561.3
2002.....	18.5	36.7	61.8	96.9	162.2	224.1	306.7	399.3	591.1
2001.....	17.4	36.4	64.8	100.4	156.0	234.2	305.5	375.6	563.8
2000.....	19.3	39.4	63.0	98.6	164.6	226.4	324.4	421.7	557.3
1999.....	18.7	38.2	69.8	112.5	159.1	241.2	304.9	409.2	540.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

All origins 3/, both sexes

2007.....	0.1	0.3	0.5	0.8	1.3	2.2	4.3	8.6	24.4
2006.....	0.2	0.3	0.5	0.8	1.1	1.9	4.0	7.5	22.7
2005.....	0.2	0.4	0.5	0.6	1.2	2.7	4.7	10.0	30.8
2004.....	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.0	30.3
2003.....	0.2	0.3	0.5	0.8	1.3	2.5	5.7	10.4	35.5
2002.....	0.2	0.3	0.5	0.8	1.6	2.7	6.1	12.8	41.5
2001.....	0.2	0.3	0.5	0.8	1.4	2.8	5.9	12.8	42.9
2000.....	0.2	0.3	0.5	1.0	2.0	3.0	6.8	15.8	50.0
1999.....	0.2	0.4	0.6	0.9	1.6	3.2	6.4	15.9	53.9

All origins 3/, male

2007.....	*	0.3	0.5	0.8	1.4	2.5	4.2	9.2	22.1
2006.....	*	0.3	0.4	0.7	1.1	2.1	3.9	8.1	18.7
2005.....	0.2	0.3	0.5	0.7	1.2	2.6	4.5	9.3	27.9
2004.....	*	0.4	0.5	0.8	1.3	2.2	4.9	9.6	27.4
2003.....	*	0.3	0.5	0.7	1.4	2.7	5.6	10.9	30.6
2002.....	0.2	0.3	0.6	0.9	1.6	3.0	6.3	12.0	37.0
2001.....	0.3	0.3	0.5	0.9	1.4	2.9	5.6	13.2	35.2
2000.....	0.3	0.2	0.7	1.0	2.0	3.4	6.8	14.4	43.7
1999.....	0.2	0.4	0.7	1.1	1.6	3.8	6.8	16.0	48.4

All origins 3/, female

2007.....	*	0.3	0.4	0.8	1.2	2.0	4.4	8.3	25.5
2006.....	0.2	0.3	0.5	0.8	1.0	1.8	4.0	7.2	24.5
2005.....	0.2	0.4	0.4	0.5	1.2	2.8	5.0	10.4	32.1
2004.....	0.2	0.4	0.4	0.7	1.1	2.0	4.3	10.3	31.7
2003.....	0.2	0.3	0.5	0.8	1.2	2.3	5.8	10.1	37.6
2002.....	*	0.4	0.4	0.7	1.6	2.4	6.0	13.3	43.5
2001.....	0.2	0.4	0.4	0.7	1.3	2.7	6.1	12.6	46.1
2000.....	0.2	0.4	0.4	0.9	2.0	2.7	6.8	16.6	52.6
1999.....	0.2	0.4	0.5	0.7	1.5	2.6	6.2	15.8	56.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Non-Hispanic 4/, both sexes

2007	0.2	0.3	0.5	0.8	1.4	2.3	4.4	8.8	24.8
2006	0.2	0.3	0.5	0.8	1.1	2.0	3.9	7.5	22.8
2005	0.2	0.4	0.5	0.6	1.2	2.7	4.8	9.8	31.1
2004	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.2	30.8
2003	0.2	0.3	0.5	0.8	1.3	2.5	5.8	10.5	35.7
2002	0.2	0.3	0.5	0.8	1.6	2.6	6.0	12.9	41.8
2001	0.2	0.4	0.5	0.8	1.4	2.9	5.9	12.9	43.0
2000	0.2	0.3	0.5	1.0	2.0	3.1	6.8	15.8	50.3
1999	0.2	0.4	0.6	0.9	1.6	3.2	6.5	16.1	53.9

Non-Hispanic 4/, male

2007	*	0.3	0.5	0.8	1.5	2.6	4.4	9.3	22.6
2006	*	0.3	0.5	0.7	1.2	2.1	3.8	8.2	18.9
2005	0.2	0.3	0.6	0.7	1.2	2.5	4.4	9.2	28.0
2004	*	0.4	0.5	0.8	1.3	2.3	5.1	9.7	28.0
2003	*	0.3	0.5	0.7	1.4	2.6	5.8	11.0	30.7
2002	*	0.3	0.6	1.0	1.6	3.0	6.2	12.0	37.1
2001	0.3	0.3	0.5	0.9	1.5	3.0	5.5	13.2	34.8
2000	0.3	*	0.7	1.1	2.0	3.4	6.7	14.3	43.9
1999	0.2	0.3	0.7	1.2	1.7	3.8	6.7	16.4	48.1

Non-Hispanic 4/, female

2007	*	0.3	0.4	0.8	1.2	2.1	4.4	8.5	25.8
2006	0.2	0.3	0.5	0.9	1.1	1.9	4.0	7.1	24.6
2005	0.2	0.4	0.4	0.6	1.2	2.8	5.1	10.2	32.5
2004	0.2	0.5	0.4	0.7	1.2	2.0	4.3	10.5	32.1
2003	*	0.4	0.5	0.8	1.3	2.3	5.8	10.1	37.9
2002	*	0.4	0.5	0.7	1.6	2.4	5.8	13.4	43.9
2001	0.2	0.4	0.4	0.8	1.3	2.7	6.2	12.7	46.4
2000	0.2	0.4	0.4	0.9	2.0	2.8	6.9	16.6	52.8
1999	0.3	0.4	0.5	0.7	1.5	2.6	6.3	16.0	56.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Non-Hispanic White, both sexes

2007 5/.....	0.1	0.3	0.4	0.8	1.2	2.0	4.1	8.4	23.7
2006 5/.....	0.2	0.3	0.4	0.7	1.0	1.9	3.9	7.0	22.2
2005 5/.....	0.2	0.3	0.5	0.5	1.1	2.6	4.4	9.2	30.3
2004 5/.....	0.2	0.4	0.4	0.6	1.0	2.0	4.3	9.9	29.6
2003 5/.....	0.2	0.3	0.5	0.7	1.3	2.3	5.6	9.9	34.4
2002.....	0.2	0.3	0.5	0.7	1.4	2.4	5.7	12.3	40.6
2001.....	0.2	0.3	0.4	0.8	1.2	2.7	5.4	12.0	42.2
2000.....	0.2	0.2	0.5	0.9	1.9	2.7	6.4	15.3	49.0
1999.....	0.2	0.3	0.4	0.8	1.4	2.6	6.0	15.1	52.8

Non-Hispanic White, male

2007 5/.....	*	*	0.4	0.7	1.3	2.2	4.0	8.8	21.7
2006 5/.....	*	*	0.3	0.6	1.0	2.0	3.5	7.3	18.4
2005 5/.....	*	0.3	0.5	0.6	1.1	2.4	4.1	8.4	27.4
2004 5/.....	*	0.3	0.4	0.7	0.9	2.2	4.6	9.3	26.9
2003 5/.....	*	*	0.4	0.6	1.4	2.2	5.3	10.1	29.6
2002.....	*	*	0.5	0.7	1.4	2.7	6.0	11.0	36.5
2001.....	*	*	0.4	0.8	1.2	2.7	5.1	12.3	33.3
2000.....	*	*	0.5	0.9	1.8	2.8	6.1	13.8	43.0
1999.....	*	*	*	1.0	1.4	3.0	5.9	15.0	46.2

Non-Hispanic White, female

2007 5/.....	*	0.3	0.4	0.9	1.2	1.9	4.2	8.2	24.6
2006 5/.....	*	0.3	0.5	0.9	1.0	1.8	4.1	6.9	24.0
2005 5/.....	*	0.3	0.4	0.5	1.1	2.7	4.7	9.7	31.7
2004 5/.....	*	0.4	0.4	0.6	1.2	1.8	4.0	10.2	30.8
2003 5/.....	*	*	0.5	0.8	1.1	2.3	5.9	9.8	36.5
2002.....	*	0.4	0.5	0.7	1.4	2.2	5.5	13.2	42.3
2001.....	*	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.9
2000.....	*	0.3	0.4	0.9	1.9	2.6	6.5	16.1	51.5
1999.....	*	0.4	0.4	0.6	1.5	2.3	6.0	15.1	55.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	*	*	2.8	5.3	8.0	14.3	37.1
2006 5/.....	*	*	*	*	2.7	3.5	5.2	13.8	31.3
2005 5/.....	*	*	*	1.6	2.7	4.2	8.9	18.4	43.1
2004 5/.....	*	0.9	*	1.8	2.8	4.3	8.3	14.2	47.2
2003 5/.....	*	1.0	*	1.7	*	4.7	9.2	19.2	55.9
2002.....	*	*	1.6	2.1	3.6	5.1	10.7	20.8	60.2
2001.....	*	1.1	*	*	3.1	4.5	11.0	24.0	55.2
2000.....	*	*	*	2.2	3.3	6.7	12.1	23.0	67.4
1999.....	*	*	2.0	2.0	3.3	8.4	12.5	31.1	70.8

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	*	6.5	10.0	17.3	34.0
2006 5/.....	*	*	*	*	*	*	*	19.0	29.3
2005 5/.....	*	*	*	*	*	*	*	19.4	37.1
2004 5/.....	*	*	*	*	*	*	10.7	14.7	45.7
2003 5/.....	*	*	*	*	*	7.4	14.0	24.1	49.0
2002.....	*	*	*	*	*	*	9.9	27.3	45.9
2001.....	*	*	*	*	*	6.9	11.9	24.5	58.9
2000.....	*	*	*	*	*	9.9	14.9	23.2	64.6
1999.....	*	*	*	*	*	13.2	16.4	35.4	76.1

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	4.5	6.9	12.7	38.3
2006 5/.....	*	*	*	*	*	*	*	11.1	32.1
2005 5/.....	*	*	*	*	*	4.9	9.3	17.8	45.6
2004 5/.....	*	*	*	*	*	*	6.9	13.9	47.8
2003 5/.....	*	*	*	*	*	*	6.4	16.6	58.6
2002.....	*	*	*	*	*	4.6	11.2	17.5	65.6
2001.....	*	*	*	*	*	*	10.5	23.7	53.9
2000.....	*	*	*	*	*	4.5	10.5	22.9	68.5
1999.....	*	*	*	*	*	5.3	10.2	29.0	68.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

All origins 3/, both sexes

2007.....	0.1	0.3	0.4	0.7	1.2	2.1	4.1	8.0	22.4
2006.....	0.2	0.3	0.4	0.7	1.0	1.8	3.7	7.0	20.9
2005.....	0.2	0.3	0.4	0.6	1.2	2.5	4.6	9.3	28.9
2004.....	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
2003.....	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0

All origins 3/, male

2007.....	*	0.3	0.5	0.7	1.3	2.2	3.9	8.7	20.3
2006.....	*	0.2	0.4	0.6	1.0	1.9	3.7	7.4	17.5
2005.....	0.2	0.3	0.5	0.6	1.2	2.4	4.4	8.8	26.3
2004.....	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
2003.....	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4

All origins 3/, female

2007.....	*	0.3	0.4	0.8	1.2	2.0	4.2	7.6	23.3
2006.....	0.2	0.3	0.5	0.7	1.0	1.8	3.8	6.8	22.4
2005.....	0.2	0.4	0.3	0.5	1.2	2.6	4.7	9.7	30.2
2004.....	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
2003.....	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

Non-Hispanic 4/, both sexes

2007	0.1	0.3	0.4	0.8	1.3	2.2	4.1	8.2	22.7
2006	0.2	0.3	0.5	0.7	1.1	1.9	3.7	7.0	21.0
2005	0.2	0.4	0.5	0.6	1.2	2.5	4.6	9.2	29.2
2004	0.2	0.4	0.4	0.7	1.2	2.1	4.5	9.8	28.6
2003	0.2	0.3	0.5	0.7	1.3	2.3	5.6	10.1	33.7
2002	0.2	0.3	0.5	0.7	1.4	2.5	5.5	12.2	38.7
2001	0.2	0.3	0.5	0.8	1.3	2.6	5.4	12.1	40.2
2000	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.5
1999	0.2	0.3	0.5	0.9	1.5	3.0	6.0	14.9	50.0

Non-Hispanic 4/, male

2007	*	0.3	0.5	0.7	1.4	2.3	4.1	8.8	20.7
2006	*	0.2	0.4	0.6	1.1	2.0	3.6	7.5	17.9
2005	0.2	0.3	0.6	0.6	1.2	2.4	4.4	8.6	26.5
2004	*	0.4	0.5	0.7	1.2	2.3	4.9	9.2	25.8
2003	*	0.3	0.5	0.7	1.3	2.5	5.5	10.5	28.4
2002	*	0.3	0.6	0.9	1.4	2.8	5.7	11.4	35.3
2001	0.2	0.3	0.5	0.8	1.4	2.7	4.9	12.7	33.2
2000	0.2	*	0.6	1.0	1.8	3.2	6.3	13.0	40.9
1999	0.2	0.3	0.6	1.1	1.6	3.6	6.1	15.1	44.3

Non-Hispanic 4/, female

2007	*	0.3	0.4	0.8	1.2	2.1	4.2	7.8	23.6
2006	0.2	0.3	0.5	0.8	1.1	1.8	3.8	6.7	22.5
2005	0.2	0.4	0.4	0.5	1.2	2.6	4.8	9.5	30.5
2004	0.2	0.4	0.3	0.7	1.1	1.9	4.1	10.2	29.9
2003	*	0.3	0.5	0.8	1.2	2.2	5.8	9.8	36.0
2002	*	0.3	0.5	0.6	1.5	2.2	5.4	12.7	40.1
2001	*	0.4	0.4	0.8	1.3	2.6	5.7	11.7	43.2
2000	0.2	0.3	0.3	0.9	1.9	2.5	6.6	15.5	48.7
1999	0.2	0.4	0.4	0.6	1.5	2.5	6.0	14.9	52.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

Non-Hispanic White, both sexes

2007 5/.....	0.1	0.3	0.4	0.7	1.2	1.8	3.9	7.8	21.6
2006 5/.....	0.2	0.2	0.4	0.7	0.9	1.8	3.6	6.5	20.3
2005 5/.....	0.2	0.3	0.4	0.5	1.0	2.4	4.2	8.6	28.5
2004 5/.....	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.5	27.3
2003 5/.....	0.1	0.2	0.4	0.6	1.2	2.1	5.4	9.5	32.3
2002.....	0.2	0.3	0.4	0.6	1.3	2.3	5.2	11.7	37.4
2001.....	0.2	0.2	0.4	0.8	1.2	2.5	5.0	11.2	39.4
2000.....	0.2	0.2	0.4	0.8	1.7	2.4	6.1	14.1	45.3
1999.....	0.2	0.3	0.3	0.8	1.3	2.5	5.6	13.9	48.8

Non-Hispanic White, male

2007 5/.....	*	*	0.4	0.7	1.2	1.9	3.7	8.2	19.7
2006 5/.....	*	*	0.3	0.6	0.9	1.8	3.2	6.7	17.2
2005 5/.....	*	0.3	0.5	0.5	1.0	2.2	4.0	7.9	25.9
2004 5/.....	*	0.3	0.4	0.6	0.9	2.1	4.4	8.8	24.6
2003 5/.....	*	*	0.3	0.6	1.3	2.1	5.0	9.6	27.3
2002.....	*	*	0.4	0.6	1.3	2.6	5.4	10.5	34.5
2001.....	*	*	0.4	0.7	1.1	2.5	4.5	11.8	31.7
2000.....	*	*	0.4	0.8	1.7	2.6	5.8	12.3	39.8
1999.....	*	*	*	1.0	1.3	2.8	5.5	13.7	42.0

Non-Hispanic White, female

2007 5/.....	*	0.3	0.4	0.8	1.1	1.8	4.0	7.5	22.5
2006 5/.....	*	0.3	0.5	0.8	1.0	1.8	3.9	6.4	21.8
2005 5/.....	*	0.3	0.4	0.5	1.0	2.5	4.4	9.0	29.7
2004 5/.....	*	0.4	0.4	0.6	1.1	1.7	3.8	9.9	28.5
2003 5/.....	*	*	0.5	0.7	1.1	2.2	5.8	9.4	34.5
2002.....	*	0.3	0.5	0.6	1.3	2.1	5.1	12.5	38.6
2001.....	*	0.3	0.4	0.8	1.2	2.5	5.3	10.9	42.7
2000.....	*	*	*	0.9	1.8	2.2	6.3	15.1	47.4
1999.....	*	0.4	*	0.5	1.4	2.2	5.6	14.1	51.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	*	*	2.7	5.3	7.5	13.8	35.5
2006 5/.....	*	*	*	*	2.7	3.5	4.5	13.3	30.0
2005 5/.....	*	*	*	*	2.7	4.1	8.9	17.1	41.0
2004 5/.....	*	0.9	*	*	2.5	4.3	8.3	13.7	46.3
2003 5/.....	*	1.0	*	*	*	4.5	9.1	18.9	54.1
2002.....	*	*	1.5	2.0	3.1	4.4	10.4	20.0	57.2
2001.....	*	*	*	*	2.9	4.1	10.3	22.3	52.1
2000.....	*	*	*	2.0	3.2	6.7	11.2	21.3	64.2
1999.....	*	*	1.6	2.0	3.3	8.1	11.3	29.1	66.0

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	*	6.5	9.6	17.3	34.0
2006 5/.....	*	*	*	*	*	*	*	17.6	29.3
2005 5/.....	*	*	*	*	*	*	*	18.0	34.3
2004 5/.....	*	*	*	*	*	*	10.7	*	43.7
2003 5/.....	*	*	*	*	*	7.1	13.5	23.4	46.9
2002.....	*	*	*	*	*	*	9.4	25.0	44.9
2001.....	*	*	*	*	*	*	11.4	23.7	56.6
2000.....	*	*	*	*	*	9.9	14.0	23.2	62.2
1999.....	*	*	*	*	*	12.5	13.5	33.7	74.9

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	4.5	6.3	12.0	36.2
2006 5/.....	*	*	*	*	*	*	*	11.1	30.3
2005 5/.....	*	*	*	*	*	4.7	9.3	16.7	43.7
2004 5/.....	*	*	*	*	*	*	6.9	13.5	47.4
2003 5/.....	*	*	*	*	*	*	6.4	16.6	57.0
2002.....	*	*	*	*	*	*	10.9	17.5	61.9
2001.....	*	*	*	*	*	*	9.6	21.7	50.5
2000.....	*	*	*	*	*	4.5	9.6	20.3	65.0
1999.....	*	*	*	*	*	5.3	9.9	26.8	62.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	*	*	*	0.7	2.1
2006 5/.....	*	*	*	*	*	*	*	0.5	1.9
2005 5/.....	*	*	*	*	*	*	*	0.6	1.8
2004 5/.....	*	*	*	*	*	*	*	*	2.3
2003 5/.....	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	0.5	0.6	3.2
2001.....	*	*	*	*	*	*	0.4	0.8	2.8
2000.....	*	*	*	*	*	0.3	*	1.2	3.8
1999.....	*	*	*	*	*	*	0.4	1.1	3.9

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	*	*	1.9
2006 5/.....	*	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*	1.5
2004 5/.....	*	*	*	*	*	*	*	*	2.3
2003 5/.....	*	*	*	*	*	*	*	*	2.3
2002.....	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	1.5	3.1
1999.....	*	*	*	*	*	*	*	1.3	4.2

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	0.7	2.1
2006 5/.....	*	*	*	*	*	*	*	*	2.2
2005 5/.....	*	*	*	*	*	*	*	0.7	2.0
2004 5/.....	*	*	*	*	*	*	*	*	2.3
2003 5/.....	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	0.7	3.7
2001.....	*	*	*	*	*	*	*	0.9	3.3
2000.....	*	*	*	*	*	*	*	1.0	4.0
1999.....	*	*	*	*	*	*	*	1.0	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

All origins 3/, both sexes

2007.....	0.2	0.3	0.2	0.3	0.4	0.4	0.7	0.7	0.7
2006.....	0.2	0.3	0.2	0.2	0.3	0.5	0.6	0.8	0.9
2005.....	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
2004.....	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	0.9
2003.....	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5

All origins 3/, male

2007.....	0.3	0.3	0.3	0.3	0.4	0.5	0.7	*	*
2006.....	0.3	0.3	0.2	*	*	0.5	*	0.9	*
2005.....	0.3	0.3	0.4	0.3	*	*	*	*	*
2004.....	0.3	0.4	0.3	0.3	*	*	*	*	*
2003.....	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7

All origins 3/, female

2007.....	*	0.3	*	0.3	*	*	0.7	0.6	0.7
2006.....	0.2	0.2	0.3	*	0.4	0.4	0.5	0.8	0.8
2005.....	0.2	0.3	0.4	0.3	*	*	0.5	0.6	0.9
2004.....	*	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7
2003.....	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

Non-Hispanic 4/, both sexes

2007.....	0.2	0.3	0.3	0.3	0.4	0.4	0.6	0.7	0.6
2006.....	0.2	0.3	0.2	0.2	0.3	0.5	0.6	0.8	0.9
2005.....	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
2004.....	0.3	0.3	0.3	0.3	0.4	0.5	0.6	0.7	0.9
2003.....	0.3	0.3	0.3	0.4	0.6	0.5	0.8	0.7	1.1
2002.....	0.3	0.3	0.3	0.4	0.5	0.7	0.6	0.6	1.2
2001.....	0.3	0.3	0.4	0.5	0.5	0.5	0.6	0.9	0.9
2000.....	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.4

Non-Hispanic 4/, male

2007.....	0.3	0.3	0.3	0.3	0.5	*	0.7	*	*
2006.....	0.3	0.3	*	*	*	*	*	1.0	*
2005.....	0.3	0.3	0.4	*	*	*	*	*	*
2004.....	0.3	0.4	0.3	*	*	*	*	*	*
2003.....	0.3	0.4	0.4	0.5	0.6	*	0.9	*	*
2002.....	0.4	0.3	0.4	*	0.6	0.9	*	*	*
2001.....	0.3	0.4	0.4	0.6	0.5	0.7	0.9	*	*
2000.....	0.3	0.3	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.4	0.4	0.3	0.4	0.6	0.8	0.9	1.3	*

Non-Hispanic 4/, female

2007.....	*	0.3	*	0.3	*	*	0.6	0.6	0.6
2006.....	0.2	0.3	0.3	*	0.4	*	0.5	0.7	0.8
2005.....	0.2	0.2	0.4	*	*	*	0.5	0.6	0.8
2004.....	*	0.2	0.3	*	0.5	0.6	0.6	0.6	0.7
2003.....	0.2	*	0.3	*	0.6	0.4	0.7	0.7	1.0
2002.....	*	0.3	*	*	*	0.6	0.5	*	1.2
2001.....	0.2	*	0.4	0.4	0.5	0.4	*	0.9	0.7
2000.....	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.1
1999.....	*	*	*	0.5	*	0.7	0.8	1.1	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.2	0.3	0.2	0.3	0.4	0.6	0.8	0.6
2006 5/.....	0.2	0.3	0.2	0.2	0.3	0.5	0.5	0.8	0.9
2005 5/.....	0.3	0.2	0.4	0.3	0.3	0.4	0.5	0.6	0.8
2004 5/.....	0.2	0.2	0.2	0.3	0.4	0.4	0.6	0.7	0.8
2003 5/.....	0.2	0.2	0.3	0.3	0.5	0.4	0.8	0.6	1.0
2002.....	0.2	0.2	0.2	0.3	0.4	0.6	0.5	0.6	1.2
2001.....	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.8	0.7
2000.....	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.1
1999.....	0.2	0.2	0.3	0.4	0.4	0.6	0.7	1.1	1.3

Non-Hispanic White, male

2007 5/.....	0.3	*	0.3	*	*	*	*	*	*
2006 5/.....	*	0.3	*	*	*	*	*	*	*
2005 5/.....	0.3	0.3	0.4	*	*	*	*	*	*
2004 5/.....	0.3	0.3	*	*	*	*	*	*	*
2003 5/.....	0.3	*	0.3	*	0.6	*	1.0	*	*
2002.....	0.3	*	*	*	0.6	0.7	*	*	*
2001.....	*	0.3	*	0.5	*	0.7	0.8	*	*
2000.....	0.3	*	*	0.5	*	0.9	1.1	*	*
1999.....	*	*	*	*	*	0.7	*	1.3	*

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	0.6	*	*
2006 5/.....	*	0.3	*	*	*	*	*	*	0.8
2005 5/.....	*	*	0.4	*	*	*	*	*	0.8
2004 5/.....	*	*	*	*	*	0.5	0.6	*	*
2003 5/.....	*	*	*	*	0.5	*	0.6	*	0.8
2002.....	*	*	*	*	*	*	*	*	1.3
2001.....	*	*	*	*	*	*	*	0.8	*
2000.....	*	*	*	*	*	*	0.6	1.1	1.0
1999.....	*	*	*	*	*	0.6	0.7	1.0	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

All origins 3/, both sexes

2007.....	*	0.2	0.7	1.9	6.5	18.7	50.5	99.4	143.5
2006.....	*	0.3	0.7	2.1	6.0	19.8	50.1	95.4	144.8
2005.....	0.1	0.2	0.8	2.3	6.8	20.3	50.2	98.7	143.7
2004.....	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
2003.....	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4

All origins 3/, male

2007.....	*	0.3	0.8	2.6	9.4	26.7	74.8	152.8	228.0
2006.....	*	0.4	0.9	2.8	8.5	29.0	76.0	145.6	231.7
2005.....	*	0.3	0.9	3.2	9.6	30.0	75.0	153.6	228.6
2004.....	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
2003.....	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6

All origins 3/, female

2007.....	*	*	0.5	1.3	3.9	12.2	32.6	65.7	103.3
2006.....	*	*	0.4	1.3	3.8	12.2	31.0	64.0	104.1
2005.....	*	*	0.6	1.5	4.3	12.5	32.3	64.6	104.7
2004.....	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
2003.....	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Hispanic, both sexes

2007	*	*	*	*	4.9	12.6	37.8	71.3	76.5
2006	*	*	*	*	3.9	15.4	30.3	58.6	93.9
2005	*	*	*	*	5.0	14.2	36.6	67.0	95.0
2004	*	*	*	*	3.5	11.2	31.6	63.8	85.1
2003	*	*	*	*	3.7	13.2	28.2	59.8	77.7
2002	*	*	*	*	5.9	12.0	27.7	53.0	76.5
2001	*	*	*	*	4.8	13.5	30.6	60.6	86.3
2000	*	*	*	*	5.7	10.5	27.2	56.8	86.3
1999	*	*	*	*	4.1	13.1	31.3	51.1	77.2

Hispanic, male

2007	*	*	*	*	6.5	16.9	54.5	104.6	106.7
2006	*	*	*	*	*	21.4	47.4	90.1	151.1
2005	*	*	*	*	6.2	23.4	52.9	100.7	138.2
2004	*	*	*	*	*	15.7	49.0	95.3	126.1
2003	*	*	*	*	6.5	20.1	38.9	91.5	129.3
2002	*	*	*	*	8.8	19.7	40.3	67.5	113.8
2001	*	*	*	*	*	16.9	43.0	101.3	127.6
2000	*	*	*	*	*	13.1	39.1	86.9	143.1
1999	*	*	*	*	*	16.8	46.8	83.3	105.7

Hispanic, female

2007	*	*	*	*	*	9.2	25.6	49.1	60.1
2006	*	*	*	*	*	10.7	17.9	37.6	63.5
2005	*	*	*	*	*	7.1	24.8	44.7	72.3
2004	*	*	*	*	*	7.7	19.2	43.0	64.1
2003	*	*	*	*	*	7.9	20.7	39.2	51.6
2002	*	*	*	*	*	*	18.7	43.8	58.0
2001	*	*	*	*	*	10.8	21.8	35.4	66.1
2000	*	*	*	*	*	8.5	18.8	38.5	58.4
1999	*	*	*	*	*	10.3	20.4	31.9	63.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Non-Hispanic 4/, both sexes

2007	*	0.2	0.7	2.0	6.6	19.2	51.4	101.0	147.0
2006	*	0.3	0.7	2.1	6.2	20.1	51.4	97.4	147.2
2005	0.1	0.2	0.8	2.3	6.9	20.8	51.1	100.4	145.9
2004	*	0.3	0.6	2.0	6.6	19.0	48.4	95.4	137.9
2003	*	0.3	0.7	2.1	6.6	20.3	50.3	94.6	140.6
2002	*	0.2	0.6	1.9	6.6	18.8	46.2	91.2	137.4
2001	*	0.2	0.7	1.9	5.9	18.4	47.1	91.8	135.6
2000	*	0.2	0.6	1.8	6.6	17.1	45.5	88.5	133.3
1999	*	0.2	0.6	1.6	5.7	16.9	42.2	83.8	125.8

Non-Hispanic 4/, male

2007	*	0.3	0.8	2.7	9.7	27.5	76.2	155.7	235.0
2006	*	0.4	0.9	2.9	8.8	29.6	77.9	148.8	235.9
2005	*	0.3	1.0	3.3	9.9	30.5	76.3	156.5	233.1
2004	*	0.4	0.9	2.6	9.4	28.1	71.8	149.5	224.4
2003	*	0.3	0.9	2.8	9.0	29.7	76.0	142.7	226.5
2002	*	0.3	0.8	2.4	9.8	27.7	68.5	139.5	224.4
2001	*	0.3	1.1	2.8	8.3	27.1	69.9	142.9	227.8
2000	*	*	0.9	2.3	9.1	24.7	68.5	136.7	225.2
1999	*	0.3	0.7	2.4	7.7	25.2	65.1	132.9	209.4

Non-Hispanic 4/, female

2007	*	*	0.6	1.3	4.0	12.4	33.1	66.6	105.4
2006	*	*	0.4	1.4	3.9	12.3	31.9	65.4	106.0
2005	*	*	0.7	1.4	4.3	12.9	32.8	65.6	106.1
2004	*	*	0.4	1.4	4.3	11.7	31.6	62.2	99.2
2003	*	0.2	0.6	1.4	4.5	12.7	32.0	65.3	102.8
2002	*	*	0.5	1.5	3.8	11.6	30.5	61.9	99.8
2001	*	*	0.4	1.0	3.8	11.5	31.1	61.3	97.3
2000	*	*	0.4	1.3	4.4	11.0	29.4	60.2	96.1
1999	*	*	0.4	1.0	4.0	10.4	26.2	55.1	92.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Non-Hispanic White, both sexes

2007 5/.....	*	0.2	0.7	2.0	7.1	20.6	55.2	107.3	155.7
2006 5/.....	*	0.3	0.7	2.2	6.5	21.6	55.1	104.3	157.0
2005 5/.....	0.1	0.2	0.9	2.5	7.3	22.3	54.8	106.3	155.5
2004 5/.....	*	0.3	0.7	2.1	7.1	20.8	51.6	101.5	145.5
2003 5/.....	*	0.3	0.7	2.1	7.0	21.7	53.6	100.6	148.8
2002.....	*	0.3	0.6	2.1	7.0	20.0	49.3	97.4	145.8
2001.....	*	0.2	0.8	2.0	6.3	19.7	50.4	97.5	143.5
2000.....	*	0.2	0.7	1.9	7.1	18.2	48.4	93.7	141.2
1999.....	*	0.2	0.6	1.7	6.1	18.0	44.6	88.6	132.8

Non-Hispanic White, male

2007 5/.....	*	0.3	0.9	2.7	10.1	29.0	81.3	165.0	249.0
2006 5/.....	*	0.4	1.0	3.0	9.1	31.4	82.8	158.0	252.4
2005 5/.....	*	0.3	1.1	3.5	10.2	32.4	81.2	165.6	247.9
2004 5/.....	*	0.4	1.0	2.7	10.0	30.3	76.0	158.9	236.3
2003 5/.....	*	*	0.9	2.8	9.5	31.5	80.4	151.4	237.9
2002.....	*	0.4	0.8	2.7	10.3	29.3	72.7	148.2	237.8
2001.....	*	0.3	1.1	3.1	8.9	28.7	74.7	151.0	241.7
2000.....	*	*	0.9	2.5	9.8	26.2	72.0	143.4	238.6
1999.....	*	0.3	0.8	2.5	8.1	26.9	68.5	140.4	221.0

Non-Hispanic White, female

2007 5/.....	*	*	0.6	1.3	4.4	13.5	35.6	70.5	111.5
2006 5/.....	*	*	0.5	1.4	4.1	13.4	34.5	70.6	112.5
2005 5/.....	*	*	0.7	1.6	4.7	13.8	35.2	69.3	113.2
2004 5/.....	*	*	0.4	1.6	4.5	12.9	33.8	66.0	104.8
2003 5/.....	*	*	0.6	1.5	4.7	13.7	34.3	69.4	109.6
2002.....	*	*	0.5	1.6	4.0	12.4	32.6	66.3	106.0
2001.....	*	*	0.4	0.9	4.0	12.4	33.1	65.2	102.7
2000.....	*	*	0.5	1.4	4.8	11.8	31.8	64.1	101.9
1999.....	*	*	0.4	1.0	4.3	10.8	27.8	58.0	97.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	*	2.3	4.5	10.9	25.5	49.9	66.2
2006 5/.....	*	*	*	1.6	4.2	11.7	25.5	37.8	51.8
2005 5/.....	*	*	*	*	4.3	11.2	23.0	50.9	57.3
2004 5/.....	*	*	*	*	3.9	7.4	22.9	40.0	61.6
2003 5/.....	*	*	*	1.9	4.2	11.0	23.3	38.4	53.5
2002.....	*	*	*	*	4.5	9.6	21.1	32.8	51.7
2001.....	*	*	*	*	2.8	9.1	19.3	35.6	59.0
2000.....	*	*	*	*	3.2	8.9	21.0	37.4	52.5
1999.....	*	*	*	*	3.6	8.7	19.8	32.9	51.7

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	8.3	18.1	38.7	79.1	100.3
2006 5/.....	*	*	*	*	6.8	19.0	43.3	63.9	71.5
2005 5/.....	*	*	*	*	7.6	19.2	36.7	83.5	99.0
2004 5/.....	*	*	*	*	5.2	13.0	35.0	59.5	106.7
2003 5/.....	*	*	*	*	5.6	17.2	37.8	58.1	99.1
2002.....	*	*	*	*	7.4	16.1	31.0	56.2	87.5
2001.....	*	*	*	*	*	15.1	26.1	58.9	99.3
2000.....	*	*	*	*	*	14.0	37.6	65.2	88.1
1999.....	*	*	*	*	5.8	12.5	32.4	49.6	87.9

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	6.0	17.5	34.6	52.1
2006 5/.....	*	*	*	*	*	6.7	14.8	24.1	43.7
2005 5/.....	*	*	*	*	*	5.8	14.8	33.8	40.6
2004 5/.....	*	*	*	*	*	*	15.7	29.7	43.8
2003 5/.....	*	*	*	*	*	6.8	14.7	28.0	35.9
2002.....	*	*	*	*	*	5.3	15.1	20.7	38.0
2001.....	*	*	*	*	*	5.0	15.3	23.7	44.1
2000.....	*	*	*	*	*	5.4	11.0	23.3	39.2
1999.....	*	*	*	*	*	6.2	12.2	24.6	38.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

All origins 3/, both sexes

2007.....	0.1	0.3	1.3	3.4	10.0	33.8	103.4	270.8	849.1
2006.....	0.1	0.4	1.1	3.4	9.6	32.9	101.2	272.6	848.3
2005.....	0.1	0.3	1.1	3.5	9.7	33.3	102.5	275.5	861.6
2004.....	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
2003.....	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3

All origins 3/, male

2007.....	*	0.3	1.2	3.2	9.4	31.6	94.9	240.3	646.0
2006.....	*	0.4	1.0	3.4	10.3	31.6	95.1	244.5	638.6
2005.....	*	0.2	1.0	3.2	9.6	31.4	95.9	246.9	639.9
2004.....	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
2003.....	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7

All origins 3/, female

2007.....	0.2	0.3	1.3	3.6	10.5	35.7	109.7	290.1	945.7
2006.....	*	0.3	1.3	3.3	9.0	34.0	105.7	290.1	946.4
2005.....	*	0.4	1.2	3.7	9.8	34.8	107.3	293.2	963.4
2004.....	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
2003.....	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Hispanic, both sexes

2007	*	*	*	2.7	5.8	21.6	65.7	171.2	490.6
2006	*	*	*	3.4	6.8	19.9	63.1	181.9	514.5
2005	*	*	*	2.7	6.8	21.0	62.8	182.6	507.8
2004	*	*	*	*	6.0	19.8	62.0	158.4	490.9
2003	*	*	*	2.5	6.4	25.0	67.6	162.3	493.4
2002	*	*	*	*	5.9	16.8	56.3	144.4	438.5
2001	*	*	*	*	5.1	15.8	54.9	134.6	399.5
2000	*	*	*	*	7.2	17.8	55.1	123.1	380.2
1999	*	*	*	*	4.3	16.4	53.6	115.1	351.5

Hispanic, male

2007	*	*	*	*	*	24.6	59.0	151.9	390.3
2006	*	*	*	4.1	8.7	19.1	56.1	188.3	404.8
2005	*	*	*	*	5.9	24.6	69.1	183.0	412.2
2004	*	*	*	*	*	20.7	57.6	151.0	374.1
2003	*	*	*	*	*	24.3	69.3	155.9	426.4
2002	*	*	*	*	*	16.2	54.4	127.0	341.3
2001	*	*	*	*	*	12.8	60.7	126.0	306.6
2000	*	*	*	*	*	22.8	55.4	134.0	284.2
1999	*	*	*	*	*	17.3	52.3	95.9	278.2

Hispanic, female

2007	*	*	*	3.4	6.6	19.2	70.6	184.1	545.0
2006	*	*	*	*	5.2	20.5	68.2	177.7	572.8
2005	*	*	*	*	7.6	18.2	58.2	182.4	557.9
2004	*	*	*	*	7.1	19.2	65.0	163.3	551.0
2003	*	*	*	*	6.9	25.6	66.4	166.5	527.3
2002	*	*	*	*	5.8	17.2	57.6	155.5	486.6
2001	*	*	*	*	*	18.1	50.8	139.9	444.9
2000	*	*	*	*	8.2	14.0	54.9	116.5	427.3
1999	*	*	*	*	*	15.7	54.5	126.6	386.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Non-Hispanic 4/, both sexes

2007.....	0.1	0.3	1.3	3.5	10.3	34.7	106.0	276.6	867.5
2006.....	0.1	0.4	1.2	3.4	9.9	33.8	103.6	277.5	864.3
2005.....	*	0.3	1.2	3.5	9.9	34.1	104.9	280.1	877.5
2004.....	0.2	0.3	1.0	3.2	10.4	31.7	101.5	265.9	832.6
2003.....	*	0.3	1.0	3.4	10.3	33.7	100.2	258.9	814.0
2002.....	*	0.2	1.0	3.3	10.0	31.4	97.2	249.6	763.2
2001.....	*	0.3	1.2	3.4	9.3	30.0	90.0	236.3	720.5
2000.....	*	0.3	1.1	3.3	9.2	29.5	86.5	224.5	676.7
1999.....	0.1	0.2	1.0	3.1	9.0	27.3	79.5	209.1	608.9

Non-Hispanic 4/, male

2007.....	*	0.3	1.3	3.3	9.7	32.0	97.3	245.6	660.3
2006.....	*	0.4	1.0	3.4	10.4	32.4	97.5	247.6	651.1
2005.....	*	0.2	1.1	3.3	9.8	31.8	97.4	250.1	650.9
2004.....	*	0.3	0.9	3.2	9.7	30.2	91.1	240.5	631.2
2003.....	*	0.2	1.0	3.0	10.3	32.5	93.4	234.4	613.6
2002.....	*	0.3	0.8	3.2	9.4	30.2	92.6	224.5	581.4
2001.....	*	0.3	1.0	3.3	9.2	28.5	86.6	220.3	565.2
2000.....	*	0.3	1.0	3.0	9.8	29.3	81.0	209.5	531.0
1999.....	*	*	1.2	3.1	9.0	27.4	78.5	201.6	492.8

Non-Hispanic 4/, female

2007.....	*	0.3	1.3	3.7	10.8	37.0	112.4	296.2	965.3
2006.....	*	0.3	1.3	3.4	9.4	34.9	108.1	296.1	963.4
2005.....	*	0.4	1.3	3.7	10.0	36.0	110.3	298.7	980.9
2004.....	0.3	0.4	1.0	3.2	11.0	32.9	109.0	281.5	922.7
2003.....	*	0.3	1.0	3.7	10.2	34.7	105.0	273.8	902.1
2002.....	*	*	1.1	3.4	10.6	32.4	100.5	264.8	841.6
2001.....	*	0.3	1.4	3.5	9.3	31.1	92.3	245.8	785.1
2000.....	*	0.3	1.2	3.6	8.7	29.6	90.3	233.3	735.6
1999.....	*	*	0.9	3.1	9.0	27.2	80.3	213.5	655.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.4	1.5	3.7	11.0	36.0	109.4	287.3	899.8
2006 5/.....	0.1	0.4	1.3	3.6	10.5	35.2	107.5	288.3	896.5
2005 5/.....	*	0.3	1.3	3.8	10.2	35.0	108.0	289.4	909.1
2004 5/.....	0.2	0.4	1.0	3.4	10.8	32.8	104.6	273.8	861.9
2003 5/.....	*	0.3	1.1	3.7	10.5	35.1	103.5	268.7	846.9
2002.....	*	0.3	1.1	3.5	10.5	32.8	101.2	259.6	793.9
2001.....	*	0.4	1.4	3.7	9.6	31.5	92.5	244.5	749.2
2000.....	*	0.3	1.2	3.5	9.7	30.5	89.5	232.7	705.1
1999.....	*	0.3	1.1	3.4	9.4	28.1	82.4	216.8	636.6

Non-Hispanic White, male

2007 5/.....	*	0.4	1.4	3.5	10.5	32.7	99.7	254.6	686.4
2006 5/.....	*	0.4	1.1	3.5	11.0	33.5	101.1	257.2	676.4
2005 5/.....	*	*	1.1	3.6	10.0	32.3	99.2	257.6	673.6
2004 5/.....	*	0.4	1.0	3.4	10.1	30.8	93.5	248.6	653.3
2003 5/.....	*	0.3	1.1	3.1	10.6	33.5	96.2	241.9	638.1
2002.....	*	*	0.9	3.4	9.6	31.4	96.2	232.0	605.4
2001.....	*	0.3	1.0	3.5	9.4	29.7	89.2	226.7	587.2
2000.....	*	*	1.0	3.2	10.3	30.2	83.8	217.3	551.7
1999.....	*	*	1.2	3.4	9.3	28.0	81.3	209.4	515.1

Non-Hispanic White, female

2007 5/.....	*	0.3	1.6	4.0	11.4	38.7	116.7	308.2	1,000.9
2006 5/.....	*	0.3	1.5	3.6	10.1	36.6	112.3	307.9	999.2
2005 5/.....	*	0.4	1.4	3.9	10.4	37.2	114.5	309.3	1,016.8
2004 5/.....	0.3	0.4	1.1	3.4	11.4	34.4	112.8	289.4	955.3
2003 5/.....	*	0.3	1.1	4.2	10.5	36.3	108.8	285.2	938.8
2002.....	*	*	1.2	3.6	11.4	34.0	104.8	276.6	875.3
2001.....	*	0.4	1.7	3.9	9.8	32.9	94.9	255.3	816.6
2000.....	*	*	1.4	3.9	9.2	30.8	93.4	241.9	767.0
1999.....	*	*	0.9	3.4	9.5	28.1	83.2	221.1	685.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	*	2.2	7.6	35.7	105.3	240.0	691.1
2006 5/.....	*	*	*	3.1	7.7	33.4	92.4	229.4	674.5
2005 5/.....	*	*	*	2.8	10.6	38.0	101.6	245.0	698.8
2004 5/.....	*	*	*	2.3	10.4	30.8	97.6	239.7	666.9
2003 5/.....	*	*	*	1.8	10.8	30.7	93.1	210.5	591.1
2002.....	*	*	*	2.5	8.4	27.5	82.1	188.2	548.7
2001.....	*	*	*	1.8	8.9	23.8	88.4	186.8	513.2
2000.....	*	*	*	2.2	7.6	26.5	72.3	173.9	450.9
1999.....	*	*	*	*	7.0	25.8	65.1	154.1	374.7

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	6.4	38.3	103.5	215.0	510.5
2006 5/.....	*	*	*	*	8.5	33.6	90.7	202.3	478.3
2005 5/.....	*	*	*	*	12.0	37.1	104.7	228.1	527.2
2004 5/.....	*	*	*	*	9.5	32.6	91.4	213.0	507.0
2003 5/.....	*	*	*	*	11.4	32.7	94.8	209.8	465.6
2002.....	*	*	*	*	9.2	29.4	82.2	189.5	414.6
2001.....	*	*	*	*	9.9	24.8	86.3	188.1	420.4
2000.....	*	*	*	*	8.0	28.4	65.5	157.9	378.1
1999.....	*	*	*	*	8.2	28.0	62.8	141.7	320.9

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	8.5	33.9	106.4	253.1	766.0
2006 5/.....	*	*	*	2.8	7.1	33.2	93.5	243.7	754.5
2005 5/.....	*	*	*	3.5	9.6	38.7	99.8	253.9	767.9
2004 5/.....	*	*	*	*	11.1	29.5	101.3	253.8	730.0
2003 5/.....	*	*	*	*	10.3	29.3	92.1	210.9	639.7
2002.....	*	*	*	*	7.8	26.2	82.1	187.5	599.9
2001.....	*	*	*	*	8.1	23.2	89.7	186.2	547.5
2000.....	*	*	*	*	7.3	25.2	76.4	181.9	478.0
1999.....	*	*	*	*	6.1	24.4	66.4	160.2	394.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes

2007.....	80.4	132.7	200.6	305.5	470.1	756.7	1,299.7	2,348.8	5,679.0
2006.....	83.2	137.0	205.7	323.5	490.4	798.7	1,368.3	2,468.6	5,928.2
2005.....	84.2	140.0	208.2	336.6	515.2	848.2	1,452.7	2,635.2	6,398.5
2004.....	86.0	139.2	211.1	343.0	537.5	885.1	1,514.3	2,730.4	6,616.6
2003.....	87.7	142.8	224.5	363.7	574.0	946.1	1,627.3	2,918.8	7,138.0
2002.....	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001.....	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000.....	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999.....	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1

All origins 3/, male

2007.....	111.8	188.5	286.1	423.4	629.4	978.4	1,626.6	2,832.4	5,894.7
2006.....	115.5	195.7	291.1	443.8	657.9	1,027.7	1,713.4	2,982.2	6,151.8
2005.....	116.8	199.9	294.8	462.0	679.9	1,093.3	1,820.2	3,159.0	6,615.9
2004.....	120.2	198.5	296.2	472.0	714.7	1,134.9	1,900.8	3,283.3	6,818.1
2003.....	123.1	204.5	314.7	497.9	765.9	1,225.3	2,051.8	3,517.1	7,329.3
2002.....	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001.....	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000.....	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999.....	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2

All origins 3/, female

2007.....	49.7	79.2	119.9	197.3	330.6	575.6	1,059.0	2,043.3	5,576.4
2006.....	51.6	80.8	125.2	213.5	344.0	612.4	1,115.8	2,147.6	5,823.6
2005.....	52.4	82.5	126.6	222.2	371.4	649.7	1,185.8	2,309.9	6,298.7
2004.....	52.6	82.4	130.8	225.5	383.3	684.0	1,236.7	2,389.8	6,526.0
2003.....	53.3	83.7	139.7	241.8	407.6	722.2	1,325.1	2,552.9	7,053.5
2002.....	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001.....	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000.....	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999.....	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

Hispanic, both sexes

2007	50.0	87.7	145.3	237.6	370.5	588.4	1,045.3	1,825.0	3,845.6
2006	52.9	93.3	151.9	241.0	383.6	643.8	1,092.3	1,891.9	4,096.9
2005	52.4	95.7	152.3	263.5	405.1	726.7	1,192.8	2,139.6	4,469.4
2004	53.3	95.0	155.1	268.8	429.5	718.3	1,223.0	2,109.2	4,583.7
2003	54.6	94.7	165.1	277.9	457.5	768.9	1,336.3	2,247.8	5,176.6
2002	56.4	96.2	171.9	299.2	471.4	810.5	1,353.8	2,354.6	5,369.7
2001	54.0	103.2	165.7	283.6	490.6	837.6	1,459.3	2,475.8	6,036.2
2000	56.1	99.6	178.2	310.5	501.0	861.4	1,451.4	2,583.3	6,168.2
1999	59.1	108.5	186.0	313.9	543.7	875.1	1,519.7	2,691.9	6,335.9

Hispanic, male

2007	72.3	124.4	208.7	330.5	502.3	757.9	1,276.0	2,111.4	3,825.8
2006	70.0	134.7	210.2	330.8	511.9	825.5	1,379.3	2,203.6	4,103.7
2005	72.4	138.5	212.3	363.9	537.5	947.4	1,512.9	2,514.6	4,556.5
2004	76.0	134.7	218.3	375.6	580.2	932.3	1,540.6	2,563.2	4,600.7
2003	76.4	138.4	233.3	385.1	608.6	1,004.0	1,642.5	2,633.2	5,079.7
2002	79.0	135.0	247.0	413.5	626.1	1,077.3	1,686.2	2,749.8	5,494.9
2001	77.7	146.6	233.8	383.5	672.6	1,084.7	1,808.6	3,004.8	6,155.6
2000	80.2	144.5	257.4	426.9	658.5	1,145.9	1,849.2	3,116.9	6,304.3
1999	83.2	160.7	259.8	427.7	730.3	1,182.4	1,968.4	3,137.4	6,595.2

Hispanic, female

2007	26.5	50.9	85.0	154.6	259.3	455.8	877.6	1,633.8	3,856.3
2006	34.9	52.0	96.9	161.4	275.9	502.4	884.6	1,684.5	4,093.2
2005	31.6	53.3	96.0	174.8	294.8	555.4	962.9	1,890.9	4,423.7
2004	29.9	56.0	96.1	175.3	304.9	553.3	997.4	1,809.5	4,575.0
2003	32.3	52.3	101.9	184.4	333.4	588.4	1,119.1	1,996.9	5,225.8
2002	33.4	58.5	102.9	199.5	345.0	605.0	1,116.5	2,102.3	5,307.7
2001	30.2	61.7	104.1	197.2	343.2	649.1	1,211.3	2,148.2	5,977.9
2000	32.0	56.8	106.6	210.2	373.5	645.9	1,169.6	2,258.6	6,101.4
1999	35.3	58.9	119.4	215.9	392.8	643.9	1,203.1	2,426.1	6,210.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

Non-Hispanic 4/, both sexes

2007	84.2	137.3	205.0	310.3	477.1	768.1	1,315.5	2,377.5	5,771.9
2006	86.8	141.2	209.8	329.4	497.4	808.3	1,384.1	2,497.2	6,013.2
2005	87.8	144.0	212.4	341.5	522.2	855.4	1,466.1	2,657.2	6,483.0
2004	89.4	143.0	215.1	347.6	544.1	894.0	1,528.3	2,756.4	6,698.1
2003	91.0	146.7	228.4	368.7	580.8	954.6	1,639.3	2,943.4	7,206.9
2002	90.9	147.9	237.6	378.1	609.6	1,001.0	1,713.0	3,056.0	7,451.4
2001	88.3	147.8	239.3	391.7	626.6	1,025.2	1,768.6	3,152.1	7,715.2
2000	91.1	149.9	252.6	412.4	655.5	1,073.4	1,835.5	3,270.0	8,076.4
1999	90.6	151.3	258.7	421.2	684.9	1,119.5	1,898.7	3,370.5	8,239.8

Non-Hispanic 4/, male

2007	117.0	195.0	292.4	429.7	637.7	992.4	1,647.7	2,874.0	6,010.3
2006	121.1	201.5	297.1	451.7	666.9	1,039.3	1,731.8	3,023.0	6,255.9
2005	122.0	205.5	300.8	468.4	688.2	1,101.4	1,834.9	3,189.2	6,716.9
2004	125.0	204.0	301.7	477.5	722.1	1,144.7	1,917.5	3,314.1	6,917.7
2003	127.8	210.0	319.7	503.9	774.5	1,235.1	2,068.8	3,551.2	7,419.6
2002	128.4	211.5	331.6	517.8	810.8	1,295.2	2,157.3	3,663.2	7,713.8
2001	125.4	208.7	333.1	533.8	828.4	1,335.1	2,236.3	3,795.4	7,990.6
2000	130.7	213.2	355.8	563.0	866.4	1,411.9	2,318.5	3,979.0	8,513.1
1999	129.0	215.8	363.7	574.7	916.0	1,469.7	2,402.8	4,101.3	8,657.6

Non-Hispanic 4/, female

2007	52.6	82.1	122.8	200.5	336.1	584.1	1,070.6	2,065.1	5,659.3
2006	53.6	83.6	127.5	217.3	348.7	619.7	1,129.4	2,169.8	5,900.4
2005	54.6	85.1	129.0	225.6	376.9	655.6	1,198.1	2,328.0	6,376.3
2004	54.9	84.7	133.4	229.0	388.6	691.5	1,248.5	2,414.0	6,600.0
2003	55.4	86.3	142.4	245.5	412.2	728.9	1,333.5	2,572.8	7,113.4
2002	54.7	87.1	149.1	251.0	435.1	765.3	1,399.4	2,687.5	7,338.2
2001	52.6	89.5	151.1	262.3	452.1	778.4	1,441.4	2,767.2	7,600.7
2000	53.0	89.2	155.8	275.5	473.9	806.2	1,499.3	2,852.2	7,899.8
1999	53.4	89.4	160.3	281.8	486.4	845.2	1,548.4	2,943.2	8,072.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

Non-Hispanic White, both sexes

2007 5/.....	74.4	120.2	182.5	284.1	442.9	732.2	1,286.8	2,370.4	5,875.7
2006 5/.....	75.9	123.3	187.3	301.9	461.8	771.3	1,354.1	2,484.7	6,119.6
2005 5/.....	76.3	124.7	189.5	312.1	487.5	818.8	1,431.4	2,644.4	6,596.0
2004 5/.....	77.2	124.4	191.6	318.5	507.6	857.0	1,491.4	2,746.7	6,790.0
2003 5/.....	78.3	127.1	205.3	335.9	541.5	912.5	1,603.0	2,932.8	7,304.7
2002.....	78.6	129.6	214.1	346.2	572.1	961.8	1,673.1	3,043.0	7,546.1
2001.....	76.0	129.1	215.0	359.1	584.0	984.9	1,727.4	3,139.5	7,811.4
2000.....	78.1	132.4	228.1	378.9	617.5	1,032.1	1,791.9	3,258.1	8,179.0
1999.....	77.8	133.6	232.2	388.2	648.2	1,074.2	1,856.0	3,358.3	8,342.9

Non-Hispanic White, male

2007 5/.....	106.4	175.1	264.4	397.6	598.3	952.3	1,617.0	2,880.2	6,162.7
2006 5/.....	109.0	180.0	269.0	417.5	624.6	996.4	1,701.5	3,029.3	6,398.5
2005 5/.....	109.3	182.7	272.4	433.7	648.5	1,061.5	1,805.6	3,194.8	6,880.5
2004 5/.....	111.4	182.8	273.4	442.3	682.0	1,108.2	1,883.8	3,329.1	7,045.5
2003 5/.....	113.8	187.0	292.8	467.4	733.0	1,190.2	2,037.8	3,555.0	7,561.6
2002.....	115.3	190.6	304.1	481.2	771.3	1,257.4	2,127.0	3,670.7	7,845.8
2001.....	111.7	188.0	305.3	498.8	785.3	1,292.9	2,203.7	3,797.2	8,116.3
2000.....	117.3	193.1	326.7	527.5	829.2	1,372.2	2,284.2	3,984.9	8,650.5
1999.....	115.3	195.1	333.2	537.4	879.0	1,426.0	2,370.6	4,111.9	8,776.1

Non-Hispanic White, female

2007 5/.....	42.8	66.6	103.3	177.4	303.5	547.7	1,038.2	2,045.4	5,739.7
2006 5/.....	43.2	67.7	108.5	193.6	316.0	583.4	1,094.3	2,141.9	5,989.5
2005 5/.....	43.6	67.8	109.4	198.5	343.5	617.3	1,154.0	2,300.8	6,465.9
2004 5/.....	43.4	67.2	112.7	203.2	352.1	649.7	1,204.0	2,386.2	6,675.5
2003 5/.....	43.1	68.4	121.1	213.6	371.2	684.2	1,288.4	2,550.3	7,191.6
2002.....	42.2	69.9	127.6	220.8	395.7	719.9	1,348.3	2,659.1	7,416.6
2001.....	40.6	71.4	128.4	229.5	406.1	734.6	1,390.3	2,742.5	7,684.6
2000.....	39.4	73.0	133.8	241.2	431.4	758.3	1,445.9	2,826.1	7,988.9
1999.....	40.8	73.5	135.8	249.9	445.8	793.8	1,495.0	2,913.7	8,170.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

Non-Hispanic Black, both sexes

2007 5/.....	161.1	280.0	410.9	598.6	853.6	1,217.9	1,842.8	2,818.4	5,502.3
2006 5/.....	170.1	291.6	420.6	625.5	873.1	1,274.7	1,921.7	3,021.0	5,656.4
2005 5/.....	176.1	302.5	431.2	653.1	896.6	1,322.9	2,069.7	3,202.0	6,088.3
2004 5/.....	183.9	299.5	443.5	661.9	937.9	1,397.2	2,160.8	3,256.7	6,436.5
2003 5/.....	188.3	313.0	460.6	705.3	999.0	1,508.1	2,276.6	3,444.0	6,923.9
2002.....	188.5	308.2	477.1	708.5	1,015.7	1,529.7	2,367.5	3,588.7	7,136.9
2001.....	185.9	313.3	478.6	725.1	1,073.1	1,567.7	2,444.2	3,650.2	7,366.6
2000.....	195.9	310.0	497.1	748.1	1,078.7	1,639.8	2,519.2	3,750.0	7,576.5
1999.....	192.5	317.0	517.6	753.4	1,106.8	1,719.8	2,571.4	3,809.5	7,624.4

Non-Hispanic Black, male

2007 5/.....	208.4	378.1	574.2	828.6	1,144.2	1,578.0	2,327.6	3,364.0	5,259.0
2006 5/.....	222.0	400.0	582.9	867.2	1,177.5	1,669.0	2,419.2	3,552.0	5,493.9
2005 5/.....	229.6	411.6	597.0	890.8	1,176.7	1,696.5	2,541.7	3,703.4	5,778.7
2004 5/.....	240.4	400.9	606.2	906.3	1,228.5	1,753.7	2,659.1	3,720.1	6,280.6
2003 5/.....	245.3	422.8	621.2	935.5	1,291.9	1,922.6	2,820.7	4,073.6	6,625.9
2002.....	242.6	413.5	644.4	949.5	1,310.3	1,933.1	2,831.4	4,127.6	6,982.7
2001.....	242.7	413.4	637.0	947.4	1,361.6	2,006.9	2,966.0	4,316.4	7,413.8
2000.....	252.1	415.3	680.4	967.2	1,346.6	2,057.4	3,018.8	4,434.6	7,755.5
1999.....	251.7	431.3	697.1	1,003.1	1,415.5	2,159.6	3,060.6	4,425.3	7,952.2

Non-Hispanic Black, female

2007 5/.....	120.0	197.0	277.2	418.8	640.4	969.9	1,551.2	2,531.8	5,603.1
2006 5/.....	125.0	200.1	287.1	437.6	648.7	1,004.2	1,624.6	2,741.0	5,722.7
2005 5/.....	129.5	210.4	294.5	468.2	689.2	1,067.5	1,788.7	2,937.5	6,212.9
2004 5/.....	134.7	213.9	309.8	471.7	722.3	1,155.5	1,865.8	3,012.8	6,498.1
2003 5/.....	138.7	220.2	328.7	525.9	781.8	1,228.6	1,953.5	3,114.6	7,039.3
2002.....	141.6	219.1	339.9	520.3	797.4	1,259.2	2,090.4	3,308.3	7,195.9
2001.....	136.8	228.2	348.9	550.3	860.0	1,275.0	2,132.8	3,310.8	7,349.1
2000.....	147.3	220.3	347.1	575.2	881.0	1,362.6	2,218.0	3,405.4	7,510.0
1999.....	141.5	220.2	371.1	557.6	878.0	1,427.9	2,276.7	3,504.3	7,501.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes

2007.....	65.0	107.4	163.4	246.9	369.1	580.2	977.3	1,749.0	4,267.7
2006.....	67.4	111.0	167.7	261.3	386.9	615.9	1,031.7	1,841.0	4,480.8
2005.....	68.0	114.1	171.0	273.2	405.5	654.0	1,090.7	1,946.8	4,778.4
2004.....	69.6	113.7	173.4	278.1	423.7	679.6	1,130.0	2,007.8	4,895.9
2003.....	71.3	116.6	185.0	296.1	456.3	730.9	1,218.3	2,151.6	5,278.4
2002.....	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001.....	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000.....	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999.....	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0

All origins 3/, male

2007.....	94.0	158.2	239.9	351.2	509.8	774.2	1,266.4	2,206.1	4,621.8
2006.....	97.4	164.1	244.4	369.3	534.4	819.7	1,341.1	2,319.5	4,819.9
2005.....	98.7	168.8	249.4	385.1	549.7	869.1	1,414.1	2,428.3	5,143.4
2004.....	101.5	167.5	252.3	393.5	580.0	899.4	1,472.2	2,509.6	5,239.3
2003.....	104.0	173.3	268.1	416.1	624.7	975.4	1,589.0	2,695.8	5,621.5
2002.....	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001.....	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000.....	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999.....	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6

All origins 3/, female

2007.....	36.7	58.6	91.2	151.1	246.0	421.7	764.4	1,460.4	4,099.3
2006.....	38.1	60.0	95.4	162.6	258.0	450.1	805.2	1,541.9	4,322.1
2005.....	38.0	61.6	97.1	171.2	279.6	479.8	856.0	1,647.8	4,610.8
2004.....	38.5	62.2	99.1	173.2	287.7	502.6	884.1	1,698.7	4,741.5
2003.....	39.5	62.4	106.7	187.0	310.3	534.8	954.5	1,818.9	5,126.7
2002.....	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001.....	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000.....	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999.....	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

Hispanic, both sexes

2007	35.9	65.4	108.1	183.1	281.0	444.2	779.3	1,358.2	2,913.8
2006	38.4	69.1	113.0	184.3	290.2	490.1	832.8	1,416.2	3,140.6
2005	37.2	72.1	118.0	203.7	313.7	557.8	902.2	1,601.6	3,407.5
2004	39.0	70.0	116.8	210.6	330.1	545.8	918.3	1,584.5	3,446.8
2003	39.9	71.0	126.8	219.2	353.0	598.1	1,000.2	1,685.0	3,932.9
2002	41.0	73.9	134.3	234.7	376.2	620.0	1,028.8	1,746.8	4,121.9
2001	38.3	77.1	129.3	225.5	391.0	646.2	1,097.2	1,866.0	4,607.5
2000	39.9	76.8	141.1	237.3	388.4	654.8	1,096.6	1,940.5	4,701.8
1999	42.6	84.9	146.0	251.5	425.7	676.2	1,167.9	2,013.5	4,888.6

Hispanic, male

2007	54.9	97.7	160.2	262.2	396.4	586.3	982.1	1,626.3	2,960.8
2006	53.8	104.7	163.8	259.7	401.8	645.8	1,056.9	1,708.2	3,257.9
2005	54.5	109.4	171.8	289.8	427.7	741.7	1,179.1	1,939.4	3,534.2
2004	58.6	103.1	170.7	303.3	452.8	732.3	1,199.7	1,963.0	3,496.8
2003	58.0	108.6	186.7	305.9	487.8	805.4	1,260.2	2,028.2	3,874.5
2002	59.4	108.7	199.5	335.0	511.6	846.9	1,314.4	2,109.2	4,301.8
2001	58.7	115.3	191.0	311.8	547.1	860.1	1,385.1	2,307.5	4,784.3
2000	60.8	115.6	208.7	338.5	525.4	892.6	1,411.5	2,374.9	4,897.5
1999	63.4	130.7	213.6	347.6	588.1	933.4	1,526.8	2,398.7	5,012.2

Hispanic, female

2007	15.7	33.1	58.6	112.4	183.7	332.9	631.9	1,179.2	2,888.2
2006	22.2	33.7	65.0	117.5	196.5	368.8	670.7	1,221.8	3,078.3
2005	19.2	35.1	67.5	127.6	218.7	415.1	703.4	1,377.5	3,341.3
2004	18.7	37.5	66.5	129.5	228.5	401.8	718.4	1,334.7	3,421.2
2003	21.3	34.3	71.3	143.7	242.5	438.9	815.6	1,461.7	3,962.5
2002	22.4	40.2	74.4	147.2	265.7	445.1	824.9	1,515.4	4,032.8
2001	17.8	40.7	73.5	150.9	264.6	483.0	892.8	1,592.5	4,521.1
2000	19.1	39.8	80.0	150.1	277.5	474.6	873.7	1,676.2	4,605.8
1999	21.9	41.5	84.9	168.6	294.4	482.7	914.6	1,783.7	4,828.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic 4/, both sexes

2007	68.7	111.7	167.9	251.5	375.4	589.5	989.6	1,770.4	4,336.0
2006	70.9	115.0	172.0	266.9	393.4	623.6	1,042.7	1,862.1	4,542.5
2005	71.5	117.9	175.0	278.1	411.4	659.7	1,100.3	1,961.8	4,837.9
2004	72.8	117.5	177.6	282.4	429.5	686.6	1,139.9	2,025.0	4,953.4
2003	74.5	120.5	188.9	300.6	462.4	737.2	1,227.0	2,168.1	5,324.5
2002	74.2	122.7	196.7	308.3	484.9	772.1	1,287.0	2,249.3	5,484.9
2001	72.1	122.6	198.4	319.3	499.9	793.7	1,327.9	2,320.3	5,688.6
2000	73.9	123.4	209.4	335.7	522.3	833.0	1,380.1	2,402.3	5,953.6
1999	74.1	126.1	214.7	346.6	552.7	874.4	1,434.0	2,493.7	6,086.3

Non-Hispanic 4/, male

2007	99.1	164.5	246.5	357.4	517.2	786.4	1,283.6	2,239.5	4,714.6
2006	102.8	170.0	250.6	377.2	542.8	829.7	1,357.0	2,351.5	4,899.0
2005	103.9	174.2	255.1	391.5	556.9	876.2	1,425.2	2,451.1	5,222.0
2004	106.2	173.1	258.2	398.7	587.2	907.3	1,484.5	2,532.5	5,317.0
2003	108.7	178.7	273.2	422.3	632.2	982.7	1,602.4	2,720.7	5,690.6
2002	109.3	182.2	283.1	434.2	661.3	1,032.2	1,679.9	2,809.3	5,874.4
2001	106.6	179.5	285.1	448.0	678.1	1,061.9	1,743.4	2,898.2	6,077.8
2000	111.1	182.4	304.2	470.1	710.1	1,127.0	1,798.4	3,028.8	6,474.6
1999	110.6	186.3	311.6	484.3	759.8	1,180.7	1,873.1	3,140.2	6,602.0

Non-Hispanic 4/, female

2007	39.3	61.2	93.9	154.0	250.8	428.0	772.9	1,475.2	4,157.3
2006	40.0	62.5	97.9	166.0	262.5	455.4	812.6	1,557.3	4,376.7
2005	40.1	64.0	99.4	174.3	284.0	483.7	864.3	1,659.0	4,662.6
2004	40.5	64.4	101.5	176.3	291.7	508.4	892.1	1,713.5	4,790.8
2003	41.3	64.8	109.3	189.7	314.5	539.6	959.7	1,831.1	5,163.6
2002	40.4	65.8	115.3	193.8	332.0	563.7	1,009.7	1,909.4	5,316.8
2001	38.7	68.0	117.0	202.2	345.9	580.0	1,037.2	1,974.4	5,526.8
2000	37.9	66.9	120.5	213.4	360.8	600.9	1,089.1	2,033.1	5,742.9
1999	38.8	68.3	124.0	221.4	374.9	634.5	1,129.0	2,115.7	5,879.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic White, both sexes

2007 5/.....	63.0	101.0	153.4	234.2	353.5	567.4	975.0	1,774.7	4,432.5
2006 5/.....	63.8	103.7	157.5	249.2	370.9	600.8	1,027.0	1,862.2	4,642.7
2005 5/.....	64.4	105.3	159.4	258.4	389.5	638.4	1,081.1	1,960.7	4,942.5
2004 5/.....	65.0	105.4	161.8	264.2	406.0	663.9	1,120.6	2,026.9	5,038.3
2003 5/.....	66.2	107.6	173.7	278.8	436.3	710.7	1,206.9	2,168.5	5,412.2
2002.....	66.3	110.8	180.7	286.8	460.5	748.2	1,262.7	2,248.0	5,568.4
2001.....	64.0	110.9	181.6	298.4	471.8	769.0	1,302.6	2,318.8	5,776.0
2000.....	65.8	112.1	192.8	313.1	497.3	805.7	1,353.2	2,402.0	6,047.9
1999.....	65.8	114.4	196.2	324.5	528.0	844.9	1,407.9	2,492.7	6,176.6

Non-Hispanic White, male

2007 5/.....	93.1	151.7	228.4	336.2	492.8	762.9	1,269.8	2,259.9	4,858.5
2006 5/.....	94.8	156.5	232.8	354.7	515.9	804.1	1,341.0	2,368.5	5,032.8
2005 5/.....	95.7	159.5	235.8	368.9	532.8	853.5	1,412.2	2,469.1	5,374.1
2004 5/.....	97.5	159.5	239.2	376.9	561.4	886.3	1,469.9	2,556.1	5,430.9
2003 5/.....	99.5	163.7	255.9	398.9	604.6	954.8	1,588.2	2,735.5	5,821.0
2002.....	100.8	168.9	264.7	409.8	636.5	1,010.8	1,665.4	2,827.0	5,994.1
2001.....	97.9	166.9	265.9	426.1	650.7	1,037.0	1,726.7	2,911.3	6,195.4
2000.....	102.8	169.3	284.7	446.8	686.5	1,102.9	1,780.9	3,046.4	6,606.6
1999.....	101.8	172.6	289.8	459.4	735.6	1,154.2	1,858.9	3,162.2	6,713.7

Non-Hispanic White, female

2007 5/.....	33.2	51.3	81.1	138.4	228.5	403.6	753.0	1,465.3	4,230.8
2006 5/.....	33.2	52.0	84.8	150.5	241.0	431.3	792.2	1,543.5	4,460.8
2005 5/.....	33.3	52.3	85.6	155.3	261.3	459.9	835.6	1,643.3	4,745.1
2004 5/.....	32.9	52.4	87.2	159.1	267.3	480.4	864.7	1,699.3	4,862.4
2003 5/.....	33.2	52.6	94.6	167.1	286.6	510.1	930.9	1,820.0	5,232.2
2002.....	32.2	54.0	99.9	172.5	304.7	533.4	974.5	1,893.8	5,384.5
2001.....	30.5	56.0	100.8	180.0	313.7	551.2	1,002.4	1,961.1	5,601.6
2000.....	29.1	56.0	105.0	189.2	330.9	566.6	1,052.6	2,019.1	5,822.7
1999.....	30.3	57.3	106.9	199.5	346.0	598.4	1,091.5	2,097.7	5,962.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic Black, both sexes

2007 5/.....	120.1	210.2	313.4	457.6	638.3	905.1	1,339.6	2,037.8	3,989.1
2006 5/.....	129.3	218.6	320.2	475.6	655.4	950.3	1,403.3	2,189.6	4,103.9
2005 5/.....	131.2	229.5	333.3	504.0	673.2	974.1	1,515.4	2,304.1	4,373.6
2004 5/.....	138.1	230.5	344.2	500.5	707.1	1,036.2	1,553.5	2,324.0	4,610.3
2003 5/.....	143.0	238.0	355.0	540.9	764.5	1,128.5	1,655.8	2,471.6	4,977.8
2002.....	142.1	236.4	372.5	550.7	774.7	1,140.0	1,731.2	2,577.8	5,151.0
2001.....	140.3	236.6	375.4	554.8	820.9	1,173.2	1,795.1	2,634.8	5,307.7
2000.....	145.2	236.1	386.8	579.6	826.1	1,242.2	1,852.8	2,693.6	5,424.6
1999.....	145.9	244.9	407.2	587.8	861.8	1,297.7	1,889.6	2,762.0	5,514.7

Non-Hispanic Black, male

2007 5/.....	161.3	293.7	446.9	647.2	869.3	1,191.6	1,726.8	2,490.2	3,931.8
2006 5/.....	175.8	306.5	448.6	675.4	903.7	1,271.0	1,836.4	2,664.4	4,129.6
2005 5/.....	179.6	320.4	469.1	692.8	893.1	1,281.0	1,905.3	2,736.4	4,277.2
2004 5/.....	187.0	316.5	480.8	697.0	945.7	1,332.0	1,963.8	2,731.7	4,695.5
2003 5/.....	192.4	329.9	489.3	726.1	1,012.2	1,475.4	2,104.0	3,007.3	4,897.8
2002.....	190.7	324.8	512.2	751.5	1,014.9	1,464.8	2,114.1	3,061.4	5,161.2
2001.....	188.8	320.2	510.4	740.5	1,053.7	1,531.1	2,230.3	3,202.0	5,495.4
2000.....	195.0	325.5	540.6	763.9	1,047.1	1,582.4	2,260.1	3,247.6	5,677.2
1999.....	197.0	341.6	565.4	797.1	1,122.5	1,658.3	2,283.9	3,268.7	5,877.1

Non-Hispanic Black, female

2007 5/.....	84.4	139.6	204.1	309.4	468.6	707.9	1,106.8	1,800.2	4,012.8
2006 5/.....	88.9	144.3	214.7	320.2	472.3	730.4	1,144.6	1,939.3	4,093.4
2005 5/.....	89.1	152.8	221.5	357.1	510.4	764.4	1,283.3	2,076.1	4,412.4
2004 5/.....	95.5	157.9	232.0	347.5	530.1	835.7	1,310.7	2,109.5	4,576.6
2003 5/.....	100.0	160.3	244.6	396.7	580.8	894.7	1,389.7	2,191.3	5,008.9
2002.....	100.0	161.6	258.0	393.8	596.6	922.2	1,502.6	2,326.2	5,147.0
2001.....	98.4	165.5	264.8	408.8	649.0	934.8	1,535.3	2,345.9	5,238.2
2000.....	102.0	159.9	260.9	434.1	663.2	1,016.4	1,607.3	2,414.7	5,330.8
1999.....	101.9	163.0	278.2	423.7	668.6	1,058.5	1,652.1	2,510.8	5,379.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins 3/, both sexes

2007.....	0.4	0.5	0.8	1.3	2.0	3.6	6.0	10.8	17.9
2006.....	0.4	0.4	0.9	1.5	2.5	4.1	6.8	11.0	16.7
2005.....	0.4	0.5	0.9	1.5	2.5	4.2	6.7	10.9	19.0
2004.....	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
2003.....	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0

All origins 3/, male

2007.....	0.3	0.4	0.6	1.1	1.5	2.6	4.9	8.2	14.4
2006.....	0.4	0.4	0.7	1.2	1.9	3.0	5.1	7.7	12.0
2005.....	0.3	0.5	0.9	1.1	1.9	3.1	4.9	8.0	14.0
2004.....	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
2003.....	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3

All origins 3/, female

2007.....	0.4	0.5	0.9	1.5	2.5	4.4	6.9	12.4	19.6
2006.....	0.4	0.5	1.0	1.7	3.0	5.0	8.0	13.0	18.9
2005.....	0.5	0.6	1.0	1.9	3.0	5.1	8.0	12.7	21.3
2004.....	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
2003.....	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.5	0.8	1.3	2.2	3.8	6.1	11.0	18.5
2006.....	0.4	0.5	0.9	1.4	2.6	4.1	6.8	11.1	17.0
2005.....	0.4	0.5	0.9	1.5	2.5	4.3	6.8	11.2	19.5
2004.....	0.4	0.6	0.9	1.3	2.5	4.2	7.4	11.1	17.7
2003.....	0.5	0.6	1.0	1.8	3.0	4.7	8.2	11.8	20.9
2002.....	0.4	0.7	1.3	2.1	2.8	4.9	8.5	11.5	18.1
2001.....	0.5	0.7	1.2	1.7	3.2	5.0	8.2	11.6	18.0
2000.....	0.6	0.8	1.4	2.2	3.4	5.5	8.3	11.8	17.1
1999.....	0.5	0.8	1.4	2.1	3.5	5.9	8.4	12.7	18.3

Non-Hispanic 4/, male

2007.....	0.3	0.4	0.6	1.2	1.6	2.8	5.1	8.2	14.8
2006.....	0.4	0.4	0.7	1.2	1.9	3.1	5.3	7.7	12.3
2005.....	0.3	0.4	0.9	1.1	1.9	3.1	5.1	8.3	14.6
2004.....	0.4	0.5	0.7	0.9	2.0	2.8	5.7	9.6	11.9
2003.....	0.4	0.4	0.8	1.5	2.3	3.7	6.5	9.3	16.6
2002.....	0.4	0.6	1.0	1.4	1.9	4.2	6.0	8.4	12.0
2001.....	0.5	0.5	1.1	1.2	2.3	3.3	6.0	7.8	13.1
2000.....	0.5	0.6	1.1	1.8	2.3	3.5	5.9	8.1	12.5
1999.....	0.4	0.6	1.0	1.3	2.4	4.8	5.6	9.1	10.7

Non-Hispanic 4/, female

2007.....	0.4	0.5	0.9	1.5	2.6	4.6	6.9	12.7	20.2
2006.....	0.4	0.5	1.0	1.7	3.1	5.0	7.9	13.3	19.2
2005.....	0.5	0.5	1.0	1.9	3.0	5.2	8.0	13.0	21.8
2004.....	0.4	0.7	1.1	1.7	3.0	5.4	8.5	11.9	20.2
2003.....	0.6	0.8	1.1	2.1	3.6	5.6	9.4	13.3	22.8
2002.....	0.5	0.9	1.5	2.6	3.6	5.4	10.2	13.4	20.8
2001.....	0.6	0.9	1.3	2.1	4.0	6.3	9.8	13.9	20.0
2000.....	0.7	0.9	1.6	2.5	4.5	7.1	10.0	13.9	19.0
1999.....	0.6	1.0	1.8	2.8	4.4	6.7	10.3	14.8	21.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic White, both sexes

2007 5/.....	0.3	0.4	0.6	1.3	2.1	3.9	6.5	11.6	19.7
2006 5/.....	0.3	0.4	0.8	1.3	2.5	4.2	7.1	11.8	18.4
2005 5/.....	0.3	0.4	0.8	1.4	2.5	4.4	7.1	11.9	21.0
2004 5/.....	0.3	0.6	0.8	1.3	2.5	4.4	7.7	11.7	18.9
2003 5/.....	0.4	0.5	0.8	1.6	3.1	4.8	8.7	12.6	22.3
2002.....	0.4	0.7	1.1	2.0	2.7	5.1	8.7	12.3	19.3
2001.....	0.4	0.6	1.1	1.7	3.1	5.0	8.6	12.4	19.0
2000.....	0.5	0.7	1.3	2.2	3.6	5.6	8.9	12.5	18.0
1999.....	0.5	0.7	1.4	2.0	3.6	6.1	8.8	13.3	19.2

Non-Hispanic White, male

2007 5/.....	*	0.4	0.5	1.1	1.7	2.8	5.3	8.5	16.0
2006 5/.....	0.3	0.3	0.6	1.0	1.8	3.1	5.5	8.5	13.2
2005 5/.....	0.3	0.4	0.8	1.1	1.9	3.4	5.4	8.5	15.4
2004 5/.....	0.3	0.4	0.6	0.9	2.0	3.0	6.1	10.3	12.8
2003 5/.....	0.3	0.3	0.6	1.3	2.3	3.8	7.1	10.1	17.7
2002.....	0.3	0.5	0.9	1.4	1.8	4.4	6.2	9.1	12.9
2001.....	0.4	0.4	1.0	1.2	2.2	3.4	6.3	8.2	13.7
2000.....	0.5	0.6	1.1	1.9	2.3	3.5	6.3	8.7	13.0
1999.....	0.4	0.6	1.0	1.3	2.6	5.0	5.7	9.7	11.6

Non-Hispanic White, female

2007 5/.....	0.4	0.3	0.7	1.4	2.5	4.9	7.4	13.7	21.4
2006 5/.....	0.3	0.4	1.0	1.5	3.2	5.2	8.3	13.9	20.8
2005 5/.....	0.4	0.4	0.9	1.8	3.1	5.3	8.3	14.0	23.6
2004 5/.....	0.3	0.7	0.9	1.8	2.9	5.6	8.9	12.7	21.6
2003 5/.....	0.5	0.6	1.0	1.9	3.8	5.6	9.9	14.1	24.4
2002.....	0.4	0.9	1.4	2.5	3.6	5.7	10.6	14.2	22.0
2001.....	0.4	0.7	1.1	2.1	4.0	6.3	10.2	14.9	21.2
2000.....	0.6	0.8	1.4	2.4	4.7	7.3	10.7	14.8	20.0
1999.....	0.5	0.9	1.8	2.7	4.4	7.0	11.0	15.5	22.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic Black, both sexes

2007 5/.....	*	1.0	1.5	1.9	2.5	*	3.9	5.5	8.7
2006 5/.....	0.8	*	1.2	2.6	2.9	3.2	4.6	6.1	*
2005 5/.....	*	1.3	1.3	1.8	2.5	*	4.3	*	6.3
2004 5/.....	1.0	1.0	2.0	*	2.7	2.7	4.2	*	6.0
2003 5/.....	1.3	1.5	2.2	3.3	3.2	4.9	4.0	*	7.1
2002.....	0.9	1.1	2.1	2.9	3.2	*	4.9	*	*
2001.....	1.1	1.7	2.2	2.0	3.4	5.2	5.0	*	6.9
2000.....	1.0	1.1	2.0	2.1	3.5	3.7	3.8	*	8.6
1999.....	0.9	1.7	2.3	2.1	2.7	4.1	4.2	*	8.4

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*	*
2005 5/.....	*	*	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	*	*	*	9.8
2006 5/.....	*	*	*	*	*	*	*	8.1	*
2005 5/.....	*	*	*	*	*	*	5.5	*	*
2004 5/.....	*	*	*	*	*	*	5.5	*	*
2003 5/.....	1.7	1.8	*	3.4	*	6.4	*	*	*
2002.....	*	*	*	3.7	*	*	5.6	*	*
2001.....	*	2.3	2.8	*	3.8	7.3	6.5	*	*
2000.....	*	*	*	*	4.5	*	*	*	*
1999.....	*	*	2.8	*	*	4.8	*	*	11.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

All origins 3/, both sexes

2007.....	6.8	9.7	12.5	15.3	17.8	24.4	35.5	65.3	192.3
2006.....	6.8	10.0	11.8	14.9	18.2	23.1	35.2	64.2	194.0
2005.....	6.5	9.2	11.7	14.7	17.6	23.8	35.8	65.9	198.6
2004.....	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
2003.....	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9

All origins 3/, male

2007.....	9.5	13.6	17.4	21.3	23.6	27.9	40.2	64.6	146.3
2006.....	9.8	13.8	16.2	20.4	24.0	27.8	40.2	63.1	143.0
2005.....	9.1	13.0	16.5	20.3	22.5	28.5	39.3	62.2	147.5
2004.....	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
2003.....	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6

All origins 3/, female

2007.....	4.2	6.0	7.9	9.7	12.6	21.6	32.1	65.8	214.1
2006.....	3.8	6.3	7.5	9.8	13.2	19.2	31.4	64.9	217.9
2005.....	3.9	5.5	7.2	9.7	13.3	20.0	33.2	68.2	222.0
2004.....	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
2003.....	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

Hispanic, both sexes

2007	3.8	6.9	9.1	12.8	15.3	22.4	31.3	61.1	143.7
2006	3.8	6.4	8.1	11.6	17.3	19.9	35.0	55.3	157.7
2005	3.9	6.2	8.7	13.7	12.6	21.5	35.2	54.5	145.0
2004	3.9	6.0	10.5	11.6	20.3	24.7	37.3	62.3	150.3
2003	5.0	6.6	9.8	14.7	19.8	26.9	42.6	73.0	179.3
2002	4.0	6.1	10.7	14.2	19.8	25.6	43.2	65.1	165.7
2001	3.1	7.2	8.5	11.8	16.8	26.3	37.7	63.1	151.9
2000	3.1	4.6	5.8	11.3	13.3	24.7	35.5	65.2	143.3
1999	2.6	4.4	8.4	9.5	16.4	17.3	33.2	65.8	152.2

Hispanic, male

2007	5.3	9.7	14.1	16.6	19.2	28.6	37.7	62.0	109.8
2006	5.7	8.9	11.4	15.6	23.1	24.8	37.6	55.7	125.2
2005	6.0	8.3	11.0	19.0	15.1	25.4	41.6	52.5	118.3
2004	5.4	7.1	15.1	17.2	24.4	32.3	43.8	74.6	131.7
2003	7.2	9.7	14.4	20.7	25.7	37.6	51.6	84.2	144.7
2002	6.2	9.0	15.4	19.2	24.4	33.8	51.8	77.8	137.2
2001	4.5	10.4	11.4	17.1	22.3	31.6	38.9	66.2	144.7
2000	4.7	6.3	8.1	14.1	16.4	30.1	43.6	64.8	112.9
1999	4.1	5.6	11.8	13.4	20.1	21.9	34.3	59.7	107.8

Hispanic, female

2007	2.2	4.1	4.4	9.4	12.1	17.5	26.7	60.5	162.2
2006	1.7	3.9	4.9	8.2	12.3	16.0	33.2	55.0	174.9
2005	1.7	4.2	6.5	9.1	10.6	18.5	30.6	55.8	159.0
2004	2.2	5.0	6.2	6.7	16.9	18.8	32.7	54.2	159.9
2003	2.8	3.6	5.5	9.4	14.9	18.7	36.2	65.6	196.9
2002	*	3.4	6.4	9.8	16.0	19.3	37.0	56.9	179.9
2001	*	4.1	6.0	7.3	12.3	22.3	36.8	61.1	155.5
2000	*	3.0	*	8.9	10.9	20.6	29.8	65.4	158.3
1999	*	3.2	5.4	6.2	13.4	13.8	32.5	69.4	173.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

Non-Hispanic 4/, both sexes

2007.....	7.2	10.0	12.7	15.4	17.9	24.4	35.7	65.6	194.6
2006.....	7.1	10.3	12.0	15.1	18.2	23.2	35.1	64.5	195.6
2005.....	6.8	9.4	11.9	14.8	17.9	23.9	35.7	66.3	200.7
2004.....	7.1	9.2	12.3	15.0	18.5	24.9	36.6	66.2	194.3
2003.....	6.8	9.6	11.8	14.8	19.3	24.8	39.3	66.3	195.0
2002.....	6.1	8.7	11.0	14.2	18.5	23.8	37.5	65.7	187.1
2001.....	5.2	7.6	9.7	12.4	16.3	23.2	36.4	66.0	190.3
2000.....	4.8	6.7	9.5	13.3	16.5	22.7	36.1	66.0	194.3
1999.....	4.7	6.3	9.6	12.0	16.1	22.2	36.4	64.3	185.3

Non-Hispanic 4/, male

2007.....	10.0	14.0	17.5	21.6	23.9	27.6	40.2	64.8	148.2
2006.....	10.3	14.3	16.5	20.7	23.9	28.0	40.3	63.2	143.8
2005.....	9.5	13.4	16.9	20.3	22.8	28.6	38.9	62.6	148.7
2004.....	9.6	12.8	17.1	20.4	24.6	30.1	38.8	60.0	145.6
2003.....	9.1	13.0	15.9	18.6	25.0	28.6	42.2	62.0	141.7
2002.....	8.2	12.1	15.1	18.9	23.1	26.4	40.1	61.1	133.7
2001.....	7.2	10.2	12.7	16.2	20.4	26.5	38.6	63.4	133.4
2000.....	6.4	9.6	12.3	17.6	20.6	26.7	36.6	61.5	139.6
1999.....	6.2	8.3	12.6	14.8	19.3	24.4	39.4	58.8	134.2

Non-Hispanic 4/, female

2007.....	4.4	6.2	8.2	9.7	12.7	21.8	32.4	66.1	216.6
2006.....	4.1	6.5	7.7	9.9	13.2	19.4	31.2	65.3	219.6
2005.....	4.2	5.6	7.3	9.7	13.5	20.1	33.3	68.6	224.5
2004.....	4.7	5.8	7.8	10.1	13.2	20.7	35.0	69.9	216.0
2003.....	4.5	6.3	7.9	11.3	14.3	21.7	37.2	68.9	218.5
2002.....	4.1	5.6	7.2	9.9	14.5	21.6	35.6	68.5	210.2
2001.....	3.3	5.2	6.8	9.0	12.7	20.5	34.8	67.6	214.0
2000.....	3.3	3.8	6.8	9.3	13.0	19.5	35.8	68.6	216.5
1999.....	3.2	4.3	6.8	9.4	13.4	20.4	34.3	67.6	205.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

Non-Hispanic White, both sexes

2007 5/.....	5.4	7.3	9.5	11.5	14.1	19.4	29.7	58.8	186.6
2006 5/.....	5.2	7.4	9.0	11.4	13.8	17.6	29.3	57.9	188.0
2005 5/.....	4.7	6.5	8.9	11.4	13.4	19.1	29.9	58.1	190.6
2004 5/.....	4.9	6.4	8.8	11.1	14.5	19.9	30.4	58.7	184.6
2003 5/.....	4.2	6.5	8.4	11.1	14.2	19.5	32.4	60.3	184.9
2002.....	4.2	6.0	7.8	10.0	13.7	18.7	31.0	57.5	177.0
2001.....	3.3	5.0	6.7	8.8	11.8	17.9	30.4	58.9	180.4
2000.....	3.1	4.3	6.2	9.1	12.2	17.0	29.7	58.0	182.5
1999.....	2.9	4.0	6.3	8.5	11.7	16.7	29.6	56.8	175.2

Non-Hispanic White, male

2007 5/.....	7.9	10.7	13.5	16.4	18.8	21.9	33.7	56.6	140.0
2006 5/.....	7.9	11.0	12.8	15.7	17.7	21.3	32.9	55.7	134.7
2005 5/.....	7.0	9.7	12.9	16.0	17.7	22.5	33.0	54.8	137.6
2004 5/.....	7.1	9.5	12.5	15.1	19.4	24.3	32.6	52.9	135.9
2003 5/.....	6.2	9.6	11.9	14.4	18.7	22.4	34.9	56.2	130.8
2002.....	6.0	8.8	11.2	13.5	17.3	21.0	33.2	53.3	123.5
2001.....	4.8	7.1	9.4	12.0	15.4	21.1	32.5	55.5	122.0
2000.....	4.5	6.5	8.5	12.4	15.6	20.7	30.2	52.9	129.5
1999.....	4.1	5.7	8.6	10.9	13.9	18.7	32.1	51.2	123.8

Non-Hispanic White, female

2007 5/.....	3.0	4.0	5.7	7.0	9.8	17.2	26.8	60.2	208.6
2006 5/.....	2.5	3.9	5.3	7.3	10.3	14.5	26.7	59.4	212.9
2005 5/.....	2.4	3.5	5.0	7.1	9.6	16.3	27.7	60.2	214.9
2004 5/.....	2.7	3.3	5.2	7.4	10.1	16.2	28.9	62.3	206.5
2003 5/.....	2.3	3.5	5.0	7.9	10.1	17.2	30.6	62.8	208.7
2002.....	2.5	3.3	4.5	6.8	10.5	16.8	29.5	60.0	200.1
2001.....	1.8	2.9	4.2	5.8	8.6	15.3	28.9	60.9	204.7
2000.....	1.7	2.1	4.0	6.0	9.2	14.0	29.4	61.0	203.9
1999.....	1.6	2.4	4.0	6.3	9.7	15.1	27.9	60.1	195.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

Non-Hispanic Black, both sexes

2007 5/.....	19.4	30.5	39.1	51.7	53.9	73.3	102.9	147.8	315.5
2006 5/.....	20.7	32.1	37.9	49.8	57.9	76.5	98.3	148.1	304.2
2005 5/.....	21.3	30.9	38.6	46.8	58.7	70.8	101.6	168.7	341.9
2004 5/.....	22.7	31.6	42.5	50.5	56.3	76.8	104.9	164.3	318.6
2003 5/.....	24.3	33.6	42.5	48.6	64.0	78.2	115.8	147.2	334.9
2002.....	19.8	30.8	40.4	51.2	61.1	77.3	111.6	168.4	322.1
2001.....	18.9	28.5	35.7	44.1	56.4	78.2	106.1	155.9	328.7
2000.....	18.1	25.9	38.4	48.9	55.6	82.1	111.8	165.8	344.1
1999.....	18.2	25.3	38.4	42.2	57.4	79.5	114.9	161.4	308.5

Non-Hispanic Black, male

2007 5/.....	25.6	41.7	53.8	75.2	75.9	90.6	124.0	173.4	268.8
2006 5/.....	28.2	41.5	51.0	72.8	85.3	96.6	133.0	168.5	278.5
2005 5/.....	27.7	43.2	54.3	66.2	75.0	92.7	119.4	169.1	304.5
2004 5/.....	28.5	40.8	60.2	72.8	79.3	98.4	116.6	160.8	271.3
2003 5/.....	30.0	42.4	55.6	60.1	87.6	101.1	132.2	150.2	291.9
2002.....	24.9	40.4	54.2	72.1	78.8	91.0	126.8	166.9	264.7
2001.....	25.5	36.5	45.2	56.4	71.1	91.2	116.6	173.3	288.7
2000.....	22.0	36.5	48.8	65.1	71.2	95.7	117.1	179.4	275.9
1999.....	22.9	32.0	50.0	52.5	72.8	90.7	130.5	166.5	266.2

Non-Hispanic Black, female

2007 5/.....	14.1	21.1	27.1	33.3	37.8	61.3	90.2	134.4	334.9
2006 5/.....	14.2	24.1	27.1	31.9	37.8	62.8	77.5	137.4	314.6
2005 5/.....	15.7	20.5	25.6	31.8	46.6	55.8	91.0	168.5	356.9
2004 5/.....	17.6	23.8	27.9	33.1	39.2	62.2	98.0	166.1	337.3
2003 5/.....	19.2	26.2	31.7	39.6	46.6	62.7	106.0	145.7	351.6
2002.....	15.4	22.6	29.2	34.8	48.0	68.1	102.6	169.2	344.1
2001.....	13.2	21.7	28.0	34.3	45.6	69.5	99.9	147.0	343.5
2000.....	14.8	16.9	29.9	36.1	44.0	73.1	108.6	159.0	369.4
1999.....	14.1	19.7	29.0	34.1	46.1	72.1	105.4	158.9	324.2

Worktable Orig291R. Death rates for I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2007.....	0.3	0.5	0.7	1.1	1.6	2.4	4.5	8.1	22.3
2006.....	0.3	0.5	0.7	1.0	1.4	2.4	4.3	8.6	22.6
2005.....	0.3	0.6	0.6	1.0	1.8	2.9	5.0	9.5	25.3
2004.....	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
2003.....	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1
All origins 3/, male									
2007.....	0.5	0.6	1.0	1.3	1.9	2.5	4.8	8.9	23.0
2006.....	0.3	0.6	0.9	1.3	1.6	2.6	4.7	10.0	23.8
2005.....	0.5	0.7	0.8	1.2	2.1	2.9	5.3	9.8	27.4
2004.....	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
2003.....	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3
All origins 3/, female									
2007.....	0.2	0.3	0.5	0.9	1.4	2.3	4.3	7.6	22.0
2006.....	0.3	0.4	0.4	0.8	1.3	2.3	4.0	7.8	22.1
2005.....	*	0.4	0.5	0.8	1.6	2.8	4.8	9.3	24.3
2004.....	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
2003.....	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

Non-Hispanic 4/, both sexes

2007.....	0.4	0.5	0.7	1.1	1.6	2.3	4.4	8.3	22.4
2006.....	0.3	0.5	0.6	1.0	1.4	2.4	4.3	8.8	23.0
2005.....	0.3	0.6	0.6	0.9	1.7	2.7	5.1	9.4	25.4
2004.....	0.5	0.5	0.7	1.2	1.6	2.6	4.3	9.3	24.9
2003.....	0.4	0.6	0.9	1.1	1.6	2.8	5.1	9.7	25.6
2002.....	0.4	0.5	0.9	1.1	1.6	2.8	4.8	9.4	23.5
2001.....	0.3	0.4	0.8	1.2	1.7	2.7	5.0	9.2	23.1
2000.....	0.3	0.4	0.8	1.0	1.6	3.0	5.0	9.8	24.2
1999.....	0.4	0.5	0.9	1.4	2.0	3.0	6.6	12.3	29.3

Non-Hispanic 4/, male

2007.....	0.5	0.6	1.0	1.3	1.8	2.4	4.9	9.1	23.0
2006.....	0.4	0.7	0.9	1.3	1.5	2.6	4.7	10.2	24.3
2005.....	0.5	0.7	0.8	1.1	1.9	2.9	5.2	9.7	27.7
2004.....	0.6	0.7	0.7	1.6	1.7	2.9	4.9	10.1	24.4
2003.....	0.6	0.8	1.1	1.3	1.7	3.0	5.6	10.1	26.3
2002.....	0.5	0.6	1.1	1.3	1.5	3.2	5.1	9.7	22.9
2001.....	0.5	0.6	0.9	1.1	2.0	2.8	5.3	9.0	23.8
2000.....	0.4	0.5	0.9	1.0	1.7	3.1	4.8	10.7	27.9
1999.....	0.5	0.6	1.0	1.5	2.2	3.1	7.4	13.4	31.4

Non-Hispanic 4/, female

2007.....	0.2	0.3	0.5	0.9	1.4	2.3	4.1	7.8	22.2
2006.....	0.3	0.4	0.4	0.8	1.3	2.3	4.0	8.0	22.5
2005.....	*	0.5	0.5	0.7	1.6	2.6	5.0	9.2	24.4
2004.....	0.4	0.4	0.7	0.8	1.5	2.4	3.9	8.7	25.2
2003.....	0.2	0.5	0.6	0.9	1.5	2.6	4.8	9.4	25.3
2002.....	0.2	0.4	0.7	1.0	1.6	2.4	4.5	9.1	23.7
2001.....	*	0.3	0.7	1.3	1.4	2.7	4.8	9.3	22.8
2000.....	0.2	0.4	0.7	0.9	1.5	2.8	5.2	9.3	22.8
1999.....	0.3	0.3	0.7	1.2	1.9	2.9	6.1	11.7	28.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

Non-Hispanic White, both sexes

2007 5/.....	0.1	0.2	0.3	0.5	0.9	1.5	3.5	7.0	20.9
2006 5/.....	0.1	0.2	0.3	0.5	0.7	1.5	3.0	7.8	21.3
2005 5/.....	0.1	0.2	0.2	0.3	0.9	1.8	3.9	7.7	24.0
2004 5/.....	*	0.2	0.3	0.5	0.9	1.7	3.3	7.9	23.0
2003 5/.....	*	0.2	0.3	0.6	0.8	1.9	4.1	8.4	23.2
2002.....	*	*	0.3	0.5	0.7	1.8	3.6	8.2	20.8
2001.....	*	0.2	0.3	0.6	0.9	1.8	3.8	8.1	20.7
2000.....	*	0.2	0.4	0.4	0.8	2.1	3.9	8.0	21.8
1999.....	*	0.2	0.4	0.6	1.1	1.8	5.1	10.4	26.1

Non-Hispanic White, male

2007 5/.....	*	*	0.4	0.5	1.1	1.5	3.8	8.1	21.5
2006 5/.....	*	*	0.4	0.6	0.7	1.5	3.1	9.2	22.9
2005 5/.....	*	0.3	0.3	0.4	0.9	2.0	4.0	7.9	25.7
2004 5/.....	*	*	*	0.8	0.9	2.1	3.7	8.9	22.8
2003 5/.....	*	*	0.4	0.7	1.0	2.2	4.8	9.2	23.6
2002.....	*	*	0.4	0.5	0.7	2.2	4.1	8.6	20.2
2001.....	*	*	0.4	0.5	1.1	1.9	4.2	8.3	22.2
2000.....	*	*	0.4	*	1.0	2.2	4.2	9.0	24.3
1999.....	*	*	0.4	0.7	1.3	1.8	5.9	12.2	28.7

Non-Hispanic White, female

2007 5/.....	*	*	*	0.5	0.7	1.5	3.2	6.4	20.7
2006 5/.....	*	*	*	0.4	0.8	1.4	2.9	6.9	20.5
2005 5/.....	*	*	*	*	0.9	1.6	3.8	7.6	23.2
2004 5/.....	*	*	*	*	0.8	1.5	3.0	7.2	23.1
2003 5/.....	*	*	*	0.4	0.7	1.6	3.6	7.9	23.0
2002.....	*	*	*	0.5	0.8	1.5	3.3	8.0	21.0
2001.....	*	*	*	0.7	0.8	1.8	3.6	7.9	20.1
2000.....	*	*	*	*	0.7	2.0	3.7	7.4	20.8
1999.....	*	*	*	0.6	0.9	1.7	4.6	9.3	25.1

Worktable Orig291R. Death rates for I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

Non-Hispanic Black, both sexes

2007 5/.....	1.8	2.4	3.9	5.7	7.5	9.8	14.2	22.4	40.1
2006 5/.....	1.7	2.8	3.5	5.9	6.7	11.2	16.9	21.6	42.5
2005 5/.....	1.6	3.5	3.8	5.7	8.9	11.0	16.8	29.1	42.9
2004 5/.....	2.8	3.0	4.0	6.5	7.6	9.9	15.4	26.6	49.2
2003 5/.....	2.4	3.9	5.5	5.6	7.7	11.3	16.9	24.4	53.8
2002.....	2.3	3.3	6.1	6.3	8.1	12.5	17.6	23.2	57.2
2001.....	1.9	2.4	4.4	5.9	8.6	11.8	18.4	22.9	50.6
2000.....	2.0	2.5	4.1	5.4	8.4	11.9	17.0	31.3	50.9
1999.....	2.3	2.3	5.2	7.4	10.3	15.7	22.0	35.8	66.0

Non-Hispanic Black, male

2007 5/.....	2.5	2.8	5.7	8.4	8.5	11.5	17.3	23.6	42.8
2006 5/.....	2.0	3.6	5.2	7.8	8.5	13.6	23.0	23.9	40.3
2005 5/.....	2.8	4.1	4.8	7.9	11.8	10.7	17.0	33.8	50.4
2004 5/.....	3.4	4.0	4.5	8.9	8.2	9.6	21.0	27.9	42.7
2003 5/.....	3.5	4.8	7.8	6.4	6.8	12.1	15.9	24.1	51.1
2002.....	3.1	4.7	7.8	7.5	7.4	14.0	17.4	22.6	62.4
2001.....	3.0	3.6	5.6	6.5	10.9	13.4	18.5	18.8	45.0
2000.....	2.8	3.0	5.1	5.7	9.3	13.3	12.5	33.5	64.6
1999.....	3.1	3.6	6.4	9.6	10.9	17.7	23.7	30.1	60.6

Non-Hispanic Black, female

2007 5/.....	*	1.9	2.5	3.6	6.8	8.6	12.3	21.9	39.1
2006 5/.....	1.4	2.0	2.1	4.4	5.3	9.6	13.2	20.4	43.3
2005 5/.....	*	2.9	3.0	3.9	6.7	11.1	16.7	26.6	39.8
2004 5/.....	2.4	2.2	3.6	4.6	7.2	10.1	12.1	25.9	51.8
2003 5/.....	*	3.2	3.6	4.9	8.3	10.7	17.5	24.5	54.9
2002.....	1.5	2.2	4.7	5.4	8.5	11.5	17.7	23.5	55.2
2001.....	*	*	3.4	5.4	6.9	10.8	18.4	25.0	52.6
2000.....	*	2.1	3.3	5.2	7.7	10.9	19.7	30.2	45.8
1999.....	1.7	*	4.1	5.6	9.8	14.5	21.0	38.6	68.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

All origins 3/, both sexes

2007.....	40.6	71.9	113.7	175.7	261.3	406.2	675.1	1,170.4	2,705.3
2006.....	42.2	75.0	118.1	188.5	279.1	441.7	726.4	1,263.9	2,904.7
2005.....	42.9	78.3	120.9	198.6	295.4	472.3	778.6	1,359.8	3,158.1
2004.....	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
2003.....	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1

All origins 3/, male

2007.....	62.4	111.5	174.6	260.3	376.2	566.1	914.1	1,554.7	3,089.2
2006.....	64.4	116.9	180.0	277.1	400.1	613.0	988.2	1,672.7	3,303.0
2005.....	66.2	121.3	183.8	291.3	415.2	653.7	1,054.3	1,780.2	3,576.8
2004.....	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
2003.....	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5

All origins 3/, female

2007.....	19.3	33.9	56.3	98.1	160.7	275.5	499.1	927.8	2,522.7
2006.....	20.6	34.8	59.7	107.5	173.3	302.4	534.8	1,008.4	2,718.3
2005.....	20.2	37.0	61.5	114.1	190.9	325.4	578.5	1,098.7	2,965.8
2004.....	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
2003.....	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

Hispanic, both sexes

2007	22.3	44.0	77.8	136.9	212.3	328.7	578.8	1,003.0	2,082.1
2006	24.0	47.8	84.0	139.6	220.8	371.8	642.6	1,071.6	2,296.7
2005	23.5	51.8	88.2	154.5	242.5	435.8	697.2	1,240.6	2,528.4
2004	24.6	49.4	87.3	159.9	250.8	424.1	710.4	1,215.2	2,613.4
2003	23.8	50.6	91.5	164.3	267.1	466.2	771.2	1,301.0	2,970.0
2002	26.9	52.6	99.9	184.7	293.4	493.0	805.7	1,353.7	3,182.9
2001	25.8	55.5	98.1	172.8	306.5	518.7	881.4	1,481.0	3,620.2
2000	27.4	55.4	109.9	186.8	312.5	521.9	873.5	1,544.0	3,697.2
1999	29.9	64.4	111.6	199.2	340.5	551.1	948.1	1,596.5	3,876.2

Hispanic, male

2007	36.0	69.0	119.6	205.0	313.2	446.6	749.6	1,247.5	2,143.6
2006	35.2	76.1	125.7	202.6	315.2	500.9	849.8	1,297.9	2,453.9
2005	35.3	81.9	133.8	226.3	342.1	594.3	924.1	1,556.8	2,694.8
2004	39.0	77.0	132.5	237.2	359.9	584.1	961.8	1,525.1	2,695.4
2003	36.0	82.6	141.2	235.0	382.1	642.3	995.3	1,590.5	2,933.9
2002	41.4	80.3	155.3	272.2	413.4	681.9	1,053.3	1,662.0	3,374.9
2001	41.3	85.6	152.6	245.2	441.9	712.3	1,147.8	1,860.8	3,828.2
2000	44.8	86.7	167.7	276.6	435.1	726.3	1,161.2	1,936.1	3,865.6
1999	46.7	104.2	171.1	279.1	483.8	773.0	1,270.9	1,949.1	3,987.7

Hispanic, female

2007	7.8	18.9	38.1	76.1	127.2	236.5	454.6	839.9	2,048.7
2006	12.2	19.6	44.6	83.8	141.6	271.4	492.6	921.0	2,213.2
2005	11.2	22.0	45.4	91.1	159.4	312.7	534.2	1,030.8	2,441.3
2004	9.8	22.5	45.1	92.3	160.6	300.6	531.9	1,010.7	2,571.3
2003	11.4	19.6	45.5	102.7	172.6	331.1	612.2	1,112.5	2,988.3
2002	12.1	25.8	49.1	108.5	195.5	347.5	628.8	1,157.0	3,087.8
2001	10.1	26.6	48.7	110.0	196.9	371.0	692.3	1,245.8	3,518.6
2000	10.0	25.5	57.6	109.4	213.2	367.1	669.8	1,305.3	3,614.6
1999	13.2	26.6	58.0	130.4	224.7	384.2	720.2	1,386.1	3,822.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

Non-Hispanic 4/, both sexes

2007.....	42.9	74.8	116.6	178.4	264.5	411.3	680.6	1,178.9	2,736.0
2006.....	44.4	77.7	120.7	191.9	282.8	445.6	730.5	1,272.7	2,931.2
2005.....	45.1	80.7	123.3	201.6	298.7	474.0	782.0	1,363.6	3,183.8
2004.....	46.1	80.4	126.6	205.7	313.9	499.1	817.5	1,425.0	3,312.1
2003.....	48.1	83.3	135.1	221.8	338.7	540.9	889.9	1,545.2	3,629.3
2002.....	48.4	85.9	142.7	228.2	361.5	572.0	943.8	1,617.7	3,787.0
2001.....	47.4	86.9	145.7	240.2	375.6	594.1	985.9	1,688.1	3,959.4
2000.....	49.7	88.9	156.4	253.1	394.4	631.3	1,036.0	1,764.3	4,187.0
1999.....	50.0	91.8	159.8	262.9	423.4	664.7	1,082.6	1,844.1	4,319.6

Non-Hispanic 4/, male

2007.....	65.9	115.9	179.1	263.9	380.0	573.5	923.5	1,571.7	3,141.5
2006.....	68.0	120.9	184.1	282.2	405.2	619.0	995.2	1,692.0	3,344.5
2005.....	69.9	124.9	187.4	295.6	419.3	656.4	1,059.6	1,789.2	3,618.5
2004.....	71.4	125.0	191.5	301.2	444.3	685.8	1,111.5	1,867.6	3,722.6
2003.....	75.1	130.6	203.9	324.5	480.9	747.7	1,207.3	2,028.4	4,045.5
2002.....	75.4	134.1	214.8	334.2	512.8	794.4	1,281.4	2,107.5	4,228.4
2001.....	74.5	134.5	219.2	349.7	529.1	823.6	1,342.0	2,185.7	4,408.5
2000.....	79.6	137.3	236.5	366.8	555.3	883.8	1,399.7	2,312.5	4,721.1
1999.....	79.5	142.5	241.2	380.4	599.4	927.6	1,464.8	2,409.4	4,854.5

Non-Hispanic 4/, female

2007.....	20.7	35.4	57.8	99.7	163.2	278.2	501.6	931.6	2,544.6
2006.....	21.6	36.3	60.9	109.3	175.5	304.0	536.6	1,011.5	2,739.1
2005.....	21.2	38.4	62.8	115.6	193.0	325.8	580.3	1,100.2	2,985.4
2004.....	21.5	37.9	65.4	118.4	200.1	348.3	606.1	1,153.3	3,128.5
2003.....	22.0	38.0	70.1	128.2	215.0	374.6	664.0	1,250.5	3,446.4
2002.....	22.2	39.7	74.9	131.8	230.2	393.9	705.5	1,320.5	3,596.5
2001.....	21.2	41.4	76.6	140.6	242.9	411.4	736.7	1,390.3	3,772.7
2000.....	20.8	42.4	81.4	149.8	255.9	431.9	782.9	1,441.2	3,971.0
1999.....	21.4	43.1	83.6	156.2	272.3	459.0	817.1	1,513.5	4,105.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

Non-Hispanic White, both sexes

2007 5/.....	41.9	71.0	110.7	170.7	254.2	400.8	674.7	1,185.6	2,794.7
2006 5/.....	42.3	73.6	114.3	183.6	271.4	434.8	723.3	1,275.7	2,992.4
2005 5/.....	43.3	75.7	116.0	192.0	288.6	464.0	773.1	1,367.4	3,253.9
2004 5/.....	44.1	75.7	119.7	197.1	301.1	487.7	810.3	1,430.2	3,369.7
2003 5/.....	45.7	78.2	128.7	210.6	325.4	527.9	879.9	1,548.1	3,691.6
2002.....	46.0	81.2	135.4	217.1	348.4	560.1	931.0	1,622.0	3,848.2
2001.....	44.8	82.1	137.8	228.7	359.7	581.5	973.2	1,692.1	4,022.8
2000.....	46.9	84.3	148.3	240.8	380.8	617.2	1,021.9	1,770.0	4,261.3
1999.....	47.3	86.6	151.0	251.6	410.9	649.3	1,071.3	1,851.7	4,390.2

Non-Hispanic White, male

2007 5/.....	65.3	111.5	171.8	255.0	369.4	563.8	919.5	1,596.0	3,241.0
2006 5/.....	65.9	116.4	176.2	271.8	392.4	608.2	990.1	1,709.8	3,441.2
2005 5/.....	67.8	119.3	178.7	284.9	409.5	648.2	1,057.5	1,808.1	3,732.9
2004 5/.....	69.5	119.7	183.5	291.9	431.4	676.4	1,110.5	1,894.0	3,814.0
2003 5/.....	72.5	124.9	197.4	312.8	468.4	733.9	1,202.8	2,043.6	4,149.5
2002.....	73.5	129.4	206.7	322.1	501.2	786.3	1,278.6	2,128.1	4,324.5
2001.....	72.2	129.4	210.4	338.5	514.8	812.4	1,338.8	2,207.1	4,506.9
2000.....	77.2	132.8	226.8	355.2	545.0	873.4	1,394.6	2,335.6	4,833.4
1999.....	77.1	136.7	231.0	368.1	589.6	915.9	1,466.0	2,439.8	4,954.1

Non-Hispanic White, female

2007 5/.....	18.7	31.3	51.7	91.5	150.9	264.1	490.3	924.0	2,583.4
2006 5/.....	18.8	31.7	54.5	101.0	163.0	290.1	523.8	1,002.4	2,783.1
2005 5/.....	19.0	33.1	55.5	105.3	180.5	311.1	562.3	1,092.3	3,034.8
2004 5/.....	18.8	32.5	58.1	108.7	184.9	331.8	590.3	1,143.1	3,170.6
2003 5/.....	19.2	32.6	62.5	115.5	198.3	358.6	646.2	1,243.6	3,490.0
2002.....	18.8	34.2	66.9	119.6	213.1	375.2	682.3	1,312.3	3,642.5
2001.....	17.7	35.8	68.1	126.7	222.7	393.8	714.3	1,381.2	3,821.4
2000.....	17.0	36.7	73.2	134.8	236.5	410.9	759.9	1,433.9	4,030.7
1999.....	17.9	37.6	74.7	143.8	254.3	436.8	794.3	1,504.7	4,165.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

Non-Hispanic Black, both sexes

2007 5/.....	59.7	117.4	187.0	285.1	403.9	581.7	866.3	1,297.8	2,494.5
2006 5/.....	66.4	124.2	195.6	303.0	429.1	626.5	935.1	1,436.6	2,640.8
2005 5/.....	66.6	132.1	205.7	326.2	437.4	647.6	1,017.6	1,529.7	2,830.2
2004 5/.....	68.9	132.9	211.0	325.1	477.3	701.6	1,040.9	1,568.8	3,038.5
2003 5/.....	73.2	136.6	217.2	357.2	509.6	764.3	1,148.6	1,713.3	3,328.7
2002.....	74.4	139.0	234.9	366.7	530.0	787.4	1,206.0	1,780.5	3,481.1
2001.....	74.3	141.7	239.4	381.7	570.0	820.0	1,257.4	1,844.5	3,628.7
2000.....	79.8	142.9	254.5	398.2	573.6	874.6	1,307.5	1,894.4	3,701.7
1999.....	78.7	152.1	262.9	400.2	602.8	911.2	1,327.7	1,932.5	3,817.9

Non-Hispanic Black, male

2007 5/.....	85.3	172.2	276.3	413.7	564.5	787.3	1,158.8	1,610.2	2,528.7
2006 5/.....	93.6	182.4	286.2	441.3	604.5	864.9	1,257.7	1,811.1	2,685.6
2005 5/.....	97.7	192.2	298.4	462.4	594.5	869.4	1,312.9	1,902.4	2,808.9
2004 5/.....	98.7	193.5	302.3	461.3	648.7	934.0	1,345.9	1,876.2	3,133.7
2003 5/.....	105.8	200.7	306.9	496.8	685.4	1,036.6	1,511.0	2,158.3	3,309.2
2002.....	103.8	199.8	335.4	516.5	713.6	1,034.1	1,508.4	2,183.9	3,572.8
2001.....	104.5	203.8	337.8	522.1	752.5	1,102.1	1,591.8	2,254.3	3,791.8
2000.....	113.9	201.0	371.1	536.0	735.3	1,148.8	1,639.7	2,329.3	3,911.2
1999.....	112.9	221.3	378.6	558.1	795.8	1,191.3	1,633.7	2,312.3	4,082.5

Non-Hispanic Black, female

2007 5/.....	37.6	71.1	113.8	184.5	285.9	440.0	690.4	1,133.7	2,480.4
2006 5/.....	42.8	75.1	121.1	195.4	299.9	463.0	742.5	1,239.0	2,622.6
2005 5/.....	39.5	81.4	129.3	220.2	321.0	496.0	841.9	1,333.1	2,838.7
2004 5/.....	43.0	81.8	135.9	219.2	350.2	544.1	860.4	1,407.1	3,000.9
2003 5/.....	44.9	82.5	143.6	248.4	379.2	580.7	933.3	1,480.5	3,336.2
2002.....	48.9	87.6	152.6	249.7	393.9	621.9	1,025.5	1,570.5	3,446.0
2001.....	48.2	88.8	158.8	271.3	435.3	632.0	1,057.8	1,635.8	3,568.3
2000.....	50.2	93.4	159.1	289.5	454.4	692.6	1,107.3	1,675.5	3,623.9
1999.....	49.3	93.5	168.6	276.3	459.7	725.3	1,143.3	1,744.2	3,719.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

All origins 3/, both sexes

2007.....	15.8	27.9	44.3	67.4	99.6	148.0	228.2	367.1	756.7
2006.....	16.5	29.2	46.7	73.9	107.0	160.4	252.0	406.4	825.8
2005.....	17.3	30.9	47.8	78.9	115.0	176.6	272.1	448.1	923.1
2004.....	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
2003.....	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3

All origins 3/, male

2007.....	24.1	42.8	66.9	98.5	140.2	201.3	298.7	472.4	855.8
2006.....	24.7	44.8	70.0	107.3	151.8	219.9	331.3	527.1	936.3
2005.....	26.5	47.2	72.2	114.0	159.3	240.5	361.8	567.1	1,045.0
2004.....	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
2003.....	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1

All origins 3/, female

2007.....	7.6	13.6	23.1	38.8	64.1	104.4	176.2	300.6	709.5
2006.....	8.4	14.3	24.7	43.4	67.8	112.0	194.0	330.9	774.1
2005.....	8.4	15.2	24.7	47.0	76.4	124.8	206.9	374.1	867.1
2004.....	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
2003.....	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute myocardial infarction (I21-I22)

Hispanic, both sexes

2007	7.9	16.0	28.3	46.1	71.4	113.7	194.0	338.1	636.8
2006	8.4	15.6	30.3	48.4	80.1	127.1	225.5	346.5	692.9
2005	8.7	17.6	32.7	58.4	85.2	163.5	249.3	426.0	787.2
2004	9.0	17.9	33.9	58.7	94.6	153.8	248.3	421.5	816.8
2003	8.3	19.9	35.7	56.7	104.7	179.5	272.6	453.7	952.0
2002	9.5	20.2	40.1	75.6	112.2	188.8	297.8	475.5	1,048.2
2001	10.5	20.5	36.2	67.2	113.4	195.2	328.7	511.1	1,185.5
2000	11.0	21.3	41.7	72.1	118.8	198.8	331.8	537.5	1,177.1
1999	12.9	24.3	38.6	78.2	129.6	207.5	365.1	558.6	1,227.8

Hispanic, male

2007	12.5	25.7	40.8	67.2	100.7	154.0	259.8	425.4	652.5
2006	11.7	23.5	42.9	67.3	112.0	172.4	293.1	411.1	742.0
2005	12.7	27.3	48.0	83.3	115.4	222.1	325.7	536.7	875.4
2004	14.3	27.8	50.0	87.9	134.3	204.1	329.4	524.7	840.6
2003	12.2	32.0	53.1	80.0	145.3	243.0	333.6	538.5	963.6
2002	14.5	29.7	60.9	108.5	155.8	255.8	383.9	595.9	1,107.7
2001	17.2	30.1	53.3	91.3	162.3	264.4	408.5	641.5	1,331.3
2000	17.6	31.2	62.7	102.8	162.9	265.9	435.5	672.9	1,273.8
1999	19.0	37.1	57.3	106.9	181.6	284.1	511.0	655.5	1,276.8

Hispanic, female

2007	3.0	6.2	16.4	27.3	46.7	82.1	146.3	279.8	628.3
2006	5.0	7.7	18.4	31.6	53.3	91.8	176.6	303.6	666.8
2005	4.5	8.0	18.4	36.3	60.0	118.1	194.4	352.5	741.0
2004	3.5	8.3	18.9	33.1	61.8	115.0	190.6	353.3	804.5
2003	4.3	8.1	19.7	36.5	71.5	130.8	229.3	398.6	946.1
2002	4.6	11.1	20.9	47.0	76.6	137.2	236.3	398.6	1,018.8
2001	3.7	11.3	20.6	46.3	73.7	142.4	272.1	430.3	1,114.2
2000	4.5	11.9	22.6	45.7	83.0	148.0	258.3	455.1	1,129.7
1999	6.8	12.2	21.7	53.4	87.5	149.8	262.1	500.8	1,204.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

Non-Hispanic 4/, both sexes

2007.....	16.8	29.1	45.7	69.0	101.7	150.4	230.3	368.3	762.6
2006.....	17.5	30.6	48.0	75.8	108.8	162.5	253.3	409.1	831.0
2005.....	18.3	32.1	49.0	80.4	117.2	177.2	273.2	448.7	928.4
2004.....	18.7	32.9	51.9	84.4	126.0	189.7	293.1	477.8	999.2
2003.....	19.9	34.4	57.0	93.3	136.3	208.7	327.8	535.2	1,123.8
2002.....	20.0	36.2	59.8	97.6	150.8	229.1	354.6	575.7	1,194.0
2001.....	20.6	37.8	63.2	102.1	156.7	237.5	375.3	614.1	1,265.3
2000.....	22.0	39.1	68.7	113.4	169.5	257.1	407.8	652.8	1,340.8
1999.....	22.2	40.1	71.7	117.2	184.5	274.1	424.3	692.4	1,381.0

Non-Hispanic 4/, male

2007.....	25.7	44.6	69.2	100.9	143.0	204.4	300.9	474.6	867.2
2006.....	26.4	46.9	72.2	110.2	154.3	222.7	333.3	533.2	945.2
2005.....	28.1	49.1	74.1	116.1	162.4	241.4	363.5	567.9	1,052.8
2004.....	28.8	50.7	78.4	122.8	175.3	256.8	389.4	612.2	1,132.5
2003.....	31.0	53.9	85.5	135.2	191.2	283.1	434.3	687.7	1,274.5
2002.....	30.8	55.8	88.5	141.0	210.7	314.5	469.4	731.0	1,374.2
2001.....	32.1	57.7	94.0	146.7	217.3	324.5	498.6	779.7	1,463.9
2000.....	35.1	59.5	102.2	162.2	235.0	353.2	538.5	840.4	1,576.2
1999.....	35.0	62.6	108.3	168.4	257.4	377.0	564.9	891.3	1,631.5

Non-Hispanic 4/, female

2007.....	8.2	14.4	23.6	39.7	65.5	106.2	178.2	301.4	713.2
2006.....	8.9	15.0	25.2	44.3	68.9	113.4	194.6	331.9	778.0
2005.....	8.8	15.9	25.3	47.8	77.6	125.0	207.5	374.9	871.7
2004.....	8.8	15.9	26.9	49.3	83.0	135.5	223.8	395.3	939.6
2003.....	9.2	15.8	30.1	55.0	88.6	148.8	251.9	442.2	1,057.5
2002.....	9.6	17.5	32.8	58.1	98.8	160.7	273.6	481.4	1,116.3
2001.....	9.5	18.7	34.2	61.6	104.3	168.3	288.9	515.1	1,182.7
2000.....	9.3	19.4	37.4	69.0	113.2	181.3	316.8	542.3	1,245.6
1999.....	9.9	18.4	37.5	70.6	121.9	193.5	326.6	576.1	1,280.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute myocardial infarction (I21-I22)

Non-Hispanic White, both sexes

2007 5/.....	16.9	28.5	44.9	67.9	99.9	148.4	229.5	368.5	772.5
2006 5/.....	17.2	30.1	46.9	74.8	107.0	160.7	251.4	410.4	841.4
2005 5/.....	18.0	31.6	47.4	79.4	115.8	176.2	270.6	449.7	941.9
2004 5/.....	18.3	31.9	50.6	83.4	123.5	186.7	293.0	480.0	1,010.1
2003 5/.....	19.3	33.2	55.8	91.5	133.9	206.3	326.2	534.8	1,137.4
2002.....	19.4	35.4	58.6	95.4	148.4	226.2	351.1	575.3	1,202.8
2001.....	20.2	36.7	61.6	100.5	153.1	234.9	372.6	614.8	1,276.8
2000.....	21.6	37.9	67.3	111.3	166.6	253.9	403.8	654.9	1,355.3
1999.....	21.8	39.3	70.2	115.6	183.3	270.8	422.0	695.3	1,394.2

Non-Hispanic White, male

2007 5/.....	26.3	44.3	68.9	100.4	142.6	202.7	300.8	480.0	888.1
2006 5/.....	26.4	46.9	71.4	109.3	153.8	222.5	333.6	538.4	964.6
2005 5/.....	28.1	49.2	72.5	116.2	163.1	242.4	364.6	573.5	1,077.9
2004 5/.....	28.8	50.0	77.4	123.4	173.8	255.1	394.1	621.8	1,156.5
2003 5/.....	30.7	52.8	85.2	134.4	190.9	281.3	434.8	690.7	1,300.9
2002.....	30.7	55.5	88.0	140.6	210.8	315.1	469.8	734.8	1,393.0
2001.....	32.4	57.3	93.3	147.1	216.2	324.4	500.3	786.9	1,483.2
2000.....	35.3	59.3	101.2	161.8	234.8	352.6	539.5	847.9	1,600.7
1999.....	35.2	62.6	107.0	168.7	258.1	377.4	569.2	903.1	1,653.8

Non-Hispanic White, female

2007 5/.....	7.6	13.0	21.7	37.4	61.7	102.9	175.8	297.4	717.7
2006 5/.....	8.0	13.6	23.3	42.5	65.0	109.1	190.0	329.8	783.9
2005 5/.....	8.1	14.3	23.2	44.9	73.5	121.2	201.0	372.4	879.7
2004 5/.....	8.0	14.1	24.7	46.1	78.6	130.2	218.9	392.2	944.5
2003 5/.....	8.1	13.9	27.4	51.5	83.3	144.6	247.6	439.0	1,065.3
2002.....	8.3	15.8	30.3	53.5	93.2	153.4	266.1	477.8	1,120.7
2001.....	8.1	16.5	31.1	57.2	97.4	162.3	282.2	510.9	1,190.9
2000.....	8.0	17.0	34.9	64.5	106.7	174.5	308.4	540.2	1,256.3
1999.....	8.6	16.4	35.0	66.5	117.7	185.8	318.7	572.7	1,290.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute myocardial infarction (I21-I22)

Non-Hispanic Black, both sexes

2007 5/.....	20.4	40.2	64.3	96.7	140.2	199.6	282.3	426.2	763.0
2006 5/.....	23.8	42.7	68.5	103.3	147.3	211.4	316.7	455.1	811.7
2005 5/.....	24.9	44.2	71.1	109.7	153.2	220.3	351.4	503.4	901.2
2004 5/.....	25.8	48.8	77.1	114.7	171.2	255.1	353.6	527.6	991.3
2003 5/.....	28.3	51.0	81.0	126.6	183.4	273.0	406.5	605.6	1,103.6
2002.....	28.5	51.8	86.3	138.6	197.2	299.8	440.2	648.9	1,212.9
2001.....	28.6	54.3	90.9	137.8	214.1	308.1	458.0	677.1	1,247.0
2000.....	30.2	56.7	96.2	154.4	222.8	333.5	495.3	705.5	1,281.8
1999.....	30.9	56.3	98.8	153.1	227.4	349.5	499.9	735.3	1,328.6

Non-Hispanic Black, male

2007 5/.....	28.4	56.3	90.4	133.6	183.6	266.0	361.5	500.0	767.9
2006 5/.....	32.4	60.4	97.2	148.2	198.9	277.6	397.7	563.9	824.6
2005 5/.....	35.3	60.5	101.3	148.1	195.0	283.8	433.0	603.0	898.3
2004 5/.....	37.3	69.2	107.5	152.7	225.4	329.2	423.0	600.0	1,020.2
2003 5/.....	39.8	73.1	110.1	171.3	234.1	357.6	522.8	742.6	1,121.9
2002.....	38.0	72.3	117.2	180.6	252.6	375.9	539.9	783.1	1,296.3
2001.....	37.5	72.9	122.5	176.6	267.8	390.8	556.0	802.9	1,389.4
2000.....	41.9	74.6	133.6	200.3	275.0	426.1	596.8	865.1	1,426.6
1999.....	42.3	78.8	141.5	202.7	295.6	436.9	597.4	863.5	1,509.4

Non-Hispanic Black, female

2007 5/.....	13.5	26.6	42.9	67.8	108.3	154.0	234.7	387.5	761.0
2006 5/.....	16.3	27.8	45.0	68.4	109.2	165.9	268.3	397.7	806.4
2005 5/.....	15.7	30.4	46.2	79.8	122.2	176.9	302.9	450.8	902.3
2004 5/.....	15.8	31.6	52.2	85.1	131.0	204.9	312.5	489.5	979.8
2003 5/.....	18.4	32.3	57.2	91.8	145.9	216.1	337.4	533.9	1,096.5
2002.....	20.2	34.5	60.9	105.8	156.2	248.7	380.7	579.1	1,181.0
2001.....	20.8	38.5	65.0	107.3	174.5	253.0	399.5	613.0	1,194.3
2000.....	20.1	41.4	65.7	118.3	184.2	272.0	434.1	625.2	1,228.0
1999.....	21.0	37.3	64.0	114.2	176.8	291.6	441.2	671.7	1,261.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

All origins 3/, both sexes

2007.....	0.6	1.1	1.8	2.5	3.4	4.4	6.4	9.6	20.8
2006.....	0.6	1.1	1.5	2.5	2.9	4.4	6.4	10.0	21.5
2005.....	0.6	1.1	1.4	2.1	2.4	4.2	5.4	9.5	20.8
2004.....	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
2003.....	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9

All origins 3/, male

2007.....	0.9	1.6	2.7	3.6	4.9	6.0	7.7	11.9	18.7
2006.....	0.8	1.6	1.9	3.6	3.6	5.9	8.0	12.1	21.8
2005.....	0.8	1.6	2.1	3.2	3.4	5.3	7.1	10.6	20.2
2004.....	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
2003.....	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3

All origins 3/, female

2007.....	0.4	0.6	0.9	1.5	2.0	3.1	5.5	8.2	21.8
2006.....	0.3	0.6	1.0	1.5	2.2	3.3	5.3	8.8	21.4
2005.....	0.4	0.6	0.7	1.2	1.6	3.3	4.2	8.8	21.1
2004.....	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
2003.....	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

Non-Hispanic 4/, both sexes

2007.....	0.7	1.2	1.8	2.6	3.5	4.5	6.6	10.0	21.4
2006.....	0.6	1.2	1.5	2.6	3.0	4.6	6.6	10.3	22.1
2005.....	0.6	1.2	1.5	2.3	2.5	4.3	5.6	9.7	21.2
2004.....	0.4	0.7	1.2	1.5	1.9	2.7	4.3	6.7	15.3
2003.....	0.5	1.0	1.3	1.8	2.9	3.8	5.4	8.6	20.6
2002.....	0.6	1.3	1.5	2.2	3.2	4.1	5.8	9.3	21.8
2001.....	0.6	1.1	1.5	2.5	3.0	4.3	5.5	9.5	22.0
2000.....	0.5	1.1	1.7	2.6	3.2	4.0	6.1	9.6	22.2
1999.....	0.5	1.0	1.7	2.3	3.2	4.5	5.6	9.5	22.2

Non-Hispanic 4/, male

2007.....	1.0	1.8	2.8	3.7	5.1	6.2	7.9	12.2	19.2
2006.....	0.9	1.8	2.1	3.8	3.8	6.1	8.3	12.4	22.6
2005.....	0.8	1.7	2.2	3.4	3.5	5.5	7.3	10.8	20.9
2004.....	0.5	1.1	1.8	2.1	2.8	3.6	5.1	8.6	17.0
2003.....	0.8	1.4	1.9	2.6	4.0	5.1	7.4	9.8	20.2
2002.....	0.9	1.9	2.2	3.1	4.6	5.2	7.3	11.7	22.1
2001.....	1.0	1.7	2.0	3.7	4.2	5.6	6.9	11.6	20.9
2000.....	0.8	1.7	2.6	3.7	4.6	5.3	7.8	11.9	21.0
1999.....	0.8	1.4	2.6	3.2	4.5	5.9	7.7	11.0	22.9

Non-Hispanic 4/, female

2007.....	0.5	0.6	0.9	1.6	2.1	3.1	5.7	8.5	22.5
2006.....	0.4	0.6	1.0	1.6	2.3	3.4	5.4	9.1	21.9
2005.....	0.4	0.6	0.8	1.2	1.7	3.4	4.4	9.0	21.4
2004.....	0.2	0.4	0.7	0.9	1.1	2.0	3.7	5.6	14.6
2003.....	0.3	0.5	0.8	1.1	1.9	2.7	4.0	7.8	20.7
2002.....	0.3	0.7	0.8	1.4	1.9	3.2	4.8	7.8	21.6
2001.....	0.3	0.5	1.0	1.4	2.1	3.2	4.5	8.2	22.4
2000.....	*	0.6	0.8	1.6	2.1	3.0	4.9	8.2	22.7
1999.....	0.3	0.6	0.8	1.5	2.1	3.3	4.2	8.5	21.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

Non-Hispanic White, both sexes

2007 5/.....	0.7	1.0	1.7	2.4	3.3	4.3	6.4	10.0	21.6
2006 5/.....	0.5	1.0	1.4	2.5	2.9	4.3	6.5	10.1	22.3
2005 5/.....	0.5	1.0	1.3	2.0	2.4	4.0	5.3	9.5	20.9
2004 5/.....	0.3	0.6	1.0	1.3	1.7	2.6	4.1	6.5	15.6
2003 5/.....	0.4	0.9	1.2	1.6	2.7	3.6	5.2	8.4	20.9
2002.....	0.5	1.2	1.4	2.0	3.1	3.9	5.8	9.2	22.0
2001.....	0.6	0.9	1.3	2.2	2.7	4.2	5.3	9.4	22.0
2000.....	0.4	1.0	1.4	2.4	3.0	3.8	5.7	9.5	22.5
1999.....	0.4	0.9	1.5	2.2	3.1	4.2	5.4	9.2	22.7

Non-Hispanic White, male

2007 5/.....	1.0	1.6	2.7	3.4	4.9	5.9	7.4	12.4	19.4
2006 5/.....	0.8	1.6	1.9	3.6	3.7	5.7	8.0	11.9	22.7
2005 5/.....	0.8	1.5	1.9	3.0	3.3	5.1	6.8	10.5	20.3
2004 5/.....	0.4	0.9	1.5	1.9	2.5	3.2	4.7	8.6	16.9
2003 5/.....	0.6	1.4	1.7	2.3	3.8	5.0	6.9	9.3	20.4
2002.....	0.8	1.7	2.2	2.9	4.4	4.8	7.3	11.5	22.7
2001.....	0.9	1.4	1.9	3.3	3.6	5.5	6.7	11.9	20.9
2000.....	0.8	1.6	2.3	3.4	4.2	5.0	7.4	11.8	21.3
1999.....	0.6	1.3	2.4	2.9	4.3	5.5	7.7	10.5	23.6

Non-Hispanic White, female

2007 5/.....	0.4	0.5	0.9	1.5	1.9	3.0	5.6	8.5	22.7
2006 5/.....	0.3	0.5	1.0	1.4	2.1	3.2	5.4	9.0	22.2
2005 5/.....	*	0.4	0.7	1.1	1.7	3.1	4.2	8.8	21.2
2004 5/.....	*	0.4	0.6	0.8	1.0	2.0	3.6	5.2	15.0
2003 5/.....	0.2	0.5	0.7	1.0	1.7	2.5	3.9	7.9	21.1
2002.....	0.3	0.7	0.8	1.2	1.9	3.1	4.7	7.8	21.8
2001.....	*	0.5	0.8	1.2	2.0	3.1	4.4	7.9	22.4
2000.....	*	0.5	0.6	1.4	2.0	2.9	4.6	8.2	23.0
1999.....	*	0.5	0.7	1.5	2.0	3.1	3.9	8.4	22.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

Non-Hispanic Black, both sexes

2007 5/.....	0.8	2.2	3.1	4.0	5.4	7.2	11.0	11.7	22.0
2006 5/.....	1.2	2.4	2.7	4.6	4.8	7.7	9.0	14.5	21.0
2005 5/.....	1.3	2.8	3.3	4.7	4.3	7.5	9.4	15.2	27.9
2004 5/.....	1.0	1.9	3.0	3.0	3.9	4.4	8.3	10.1	15.5
2003 5/.....	1.2	1.5	2.7	3.4	5.0	6.1	8.9	11.9	20.2
2002.....	1.1	2.5	2.1	4.2	5.1	7.3	7.7	12.0	21.5
2001.....	1.1	2.2	2.9	5.3	5.4	6.3	6.9	11.0	24.3
2000.....	1.0	2.1	3.9	4.9	5.2	6.1	10.1	11.8	21.9
1999.....	1.5	2.3	3.3	3.9	4.4	7.5	8.9	14.1	17.5

Non-Hispanic Black, male

2007 5/.....	*	3.5	4.7	6.5	7.6	10.9	16.0	*	20.9
2006 5/.....	*	3.3	4.0	7.1	5.8	11.1	12.9	19.7	23.8
2005 5/.....	*	3.5	5.2	7.1	6.6	10.1	14.2	18.0	28.5
2004 5/.....	*	2.7	4.6	4.8	7.0	7.3	12.1	*	23.4
2003 5/.....	1.8	2.3	4.6	5.2	7.3	8.1	14.5	19.6	24.5
2002.....	2.0	3.8	3.2	5.9	8.2	10.9	*	16.4	21.9
2001.....	*	3.7	3.5	8.4	9.4	7.9	11.4	*	24.3
2000.....	*	3.5	5.9	6.8	8.3	9.2	13.5	*	*
1999.....	2.5	2.9	5.7	6.5	6.9	10.4	10.2	17.7	*

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	3.8	4.7	8.0	11.3	22.4
2006 5/.....	*	*	*	*	4.1	5.4	6.6	11.8	19.8
2005 5/.....	*	2.2	*	*	*	5.8	6.6	13.7	27.6
2004 5/.....	*	*	*	*	*	*	6.1	9.3	12.4
2003 5/.....	*	*	*	*	*	4.8	5.5	7.9	18.6
2002.....	*	*	*	*	*	4.8	7.0	9.7	21.3
2001.....	*	*	*	*	*	5.3	*	12.1	24.4
2000.....	*	*	*	3.4	*	*	8.1	9.9	22.2
1999.....	*	*	*	*	*	5.5	8.2	12.3	16.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes

2007.....	24.2	42.9	67.6	105.8	158.3	253.8	440.5	793.8	1,927.9
2006.....	25.2	44.7	70.0	112.1	169.3	276.9	468.0	847.5	2,057.4
2005.....	25.1	46.3	71.7	117.5	178.0	291.6	501.1	902.3	2,214.2
2004.....	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
2003.....	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0

All origins 3/, male

2007.....	37.4	67.1	105.0	158.2	231.1	358.7	607.8	1,070.5	2,214.7
2006.....	38.8	70.5	108.1	166.2	244.7	387.2	649.0	1,133.5	2,344.9
2005.....	39.0	72.5	109.5	174.1	252.6	407.9	685.4	1,202.5	2,511.6
2004.....	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
2003.....	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1

All origins 3/, female

2007.....	11.2	19.8	32.4	57.8	94.6	168.0	317.4	619.0	1,791.4
2006.....	11.8	19.9	34.1	62.7	103.3	187.1	335.6	668.6	1,922.9
2005.....	11.5	21.2	36.1	65.9	112.9	197.3	367.3	715.8	2,077.6
2004.....	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
2003.....	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

Hispanic, both sexes

2007	14.4	27.9	49.0	89.8	139.7	211.9	381.2	660.9	1,437.0
2006	15.5	32.0	53.2	90.7	139.6	242.9	414.2	720.2	1,595.2
2005	14.6	34.0	55.3	95.7	156.3	270.1	445.3	810.7	1,730.5
2004	15.6	31.4	53.2	100.8	155.7	269.9	460.7	791.9	1,795.7
2003	15.4	30.6	55.6	107.1	161.9	285.4	496.6	845.6	2,012.8
2002	17.1	32.1	59.5	108.1	180.7	302.3	505.5	875.6	2,129.7
2001	15.3	34.8	61.3	104.8	191.7	321.5	549.9	968.0	2,426.6
2000	16.1	33.9	67.6	114.2	193.0	320.8	539.0	1,002.0	2,508.8
1999	17.0	39.6	72.6	119.8	210.1	343.0	579.7	1,036.7	2,640.7

Hispanic, male

2007	23.5	43.0	77.8	136.3	210.9	288.2	485.8	815.9	1,481.9
2006	23.4	52.3	82.3	134.5	201.8	325.8	555.2	880.3	1,705.1
2005	22.3	54.3	85.4	142.4	225.2	370.6	594.1	1,013.9	1,811.9
2004	24.6	48.9	82.2	148.9	225.3	379.2	630.1	998.5	1,853.5
2003	23.7	50.4	88.1	154.3	236.2	398.0	659.9	1,051.0	1,965.7
2002	26.6	50.1	93.8	162.1	257.3	422.2	666.8	1,062.6	2,262.2
2001	24.1	55.3	98.3	152.9	277.5	444.2	735.9	1,216.7	2,479.8
2000	27.0	55.4	104.3	173.3	270.7	457.0	722.0	1,257.4	2,577.7
1999	27.6	66.6	113.3	171.0	301.0	488.3	754.4	1,292.1	2,706.7

Hispanic, female

2007	4.7	12.6	21.5	48.3	79.5	152.1	305.1	557.5	1,412.7
2006	7.2	11.8	25.7	51.9	87.4	178.5	312.3	613.6	1,536.8
2005	6.5	13.8	27.1	54.5	98.9	192.1	338.3	676.0	1,687.9
2004	6.3	14.2	26.2	58.8	98.0	185.6	340.4	655.5	1,766.0
2003	6.9	11.4	25.5	66.1	100.8	199.0	380.7	711.9	2,036.7
2002	7.4	14.6	28.0	61.1	118.1	209.9	390.3	756.3	2,064.1
2001	6.4	15.2	27.9	63.2	122.2	227.9	417.8	813.9	2,400.7
2000	5.3	13.5	34.3	63.3	130.1	217.6	409.4	846.6	2,475.0
1999	6.4	14.0	35.9	75.7	136.6	233.7	456.5	884.3	2,608.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 4/, both sexes

2007.....	25.4	44.5	69.1	106.8	159.3	256.4	443.7	800.6	1,952.0
2006.....	26.3	45.9	71.2	113.5	171.0	278.5	470.6	853.2	2,078.0
2005.....	26.2	47.4	72.8	118.9	179.0	292.4	503.2	905.2	2,234.1
2004.....	27.0	46.8	73.5	119.8	186.1	306.7	520.1	940.5	2,297.5
2003.....	27.7	47.9	76.7	126.8	199.5	328.5	556.8	1,001.4	2,484.9
2002.....	27.8	48.4	81.4	128.4	207.5	338.9	583.4	1,032.8	2,571.2
2001.....	26.1	48.0	81.1	135.6	215.9	352.3	605.2	1,064.5	2,672.2
2000.....	27.2	48.7	86.0	137.1	221.6	370.1	622.1	1,101.9	2,824.0
1999.....	27.2	50.7	86.5	143.5	235.7	386.2	652.7	1,142.2	2,916.3

Non-Hispanic 4/, male

2007.....	39.2	69.6	107.1	159.3	231.8	363.0	614.7	1,084.8	2,255.1
2006.....	40.7	72.2	109.8	168.2	247.1	390.2	653.5	1,146.4	2,376.7
2005.....	41.0	74.1	111.1	176.1	253.4	409.5	688.8	1,210.4	2,544.8
2004.....	42.0	73.3	111.3	176.4	266.2	425.4	717.0	1,246.9	2,573.1
2003.....	43.3	75.3	116.5	186.7	285.7	459.5	765.6	1,330.9	2,750.7
2002.....	43.7	76.5	124.1	190.0	297.4	474.7	804.7	1,364.8	2,832.2
2001.....	41.4	75.1	123.2	199.4	307.7	493.5	836.5	1,394.4	2,923.7
2000.....	43.7	76.1	131.7	200.9	315.7	525.3	853.4	1,460.2	3,123.9
1999.....	43.7	78.5	130.3	208.8	337.5	544.7	892.2	1,507.1	3,200.1

Non-Hispanic 4/, female

2007.....	12.1	20.5	33.3	58.4	95.6	169.0	317.7	621.7	1,808.9
2006.....	12.4	20.7	34.7	63.4	104.3	187.3	336.6	670.6	1,939.2
2005.....	12.0	21.9	36.7	66.6	113.7	197.3	368.4	716.3	2,092.3
2004.....	12.5	21.5	37.8	68.2	116.1	210.7	378.6	752.4	2,174.3
2003.....	12.6	21.7	39.2	72.1	124.5	223.1	408.1	800.5	2,368.1
2002.....	12.3	21.5	41.2	72.3	129.5	230.0	427.1	831.3	2,458.6
2001.....	11.4	22.1	41.5	77.6	136.6	239.9	443.3	867.0	2,567.6
2000.....	11.3	22.4	43.2	79.1	140.7	247.6	461.2	890.7	2,702.7
1999.....	11.3	24.1	45.4	84.1	148.3	262.1	486.4	928.9	2,802.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic White, both sexes

2007 5/.....	24.2	41.4	64.0	100.4	150.9	248.1	438.8	807.1	2,000.6
2006 5/.....	24.6	42.5	65.9	106.3	161.6	269.7	465.4	855.2	2,128.7
2005 5/.....	24.8	43.2	67.3	110.7	170.3	283.8	497.1	908.3	2,291.1
2004 5/.....	25.4	43.2	68.1	112.4	176.0	298.4	513.2	943.7	2,343.9
2003 5/.....	25.9	44.2	71.7	117.5	188.8	318.0	548.5	1,004.8	2,533.4
2002.....	26.0	44.7	75.4	119.7	196.9	330.1	574.2	1,037.5	2,623.4
2001.....	24.0	44.5	74.9	126.0	203.8	342.4	595.3	1,067.8	2,724.1
2000.....	24.9	45.3	79.5	127.1	211.2	359.4	612.4	1,105.6	2,883.6
1999.....	25.2	46.5	79.3	133.8	224.5	374.3	643.9	1,147.2	2,973.2

Non-Hispanic White, male

2007 5/.....	38.0	65.6	100.2	151.2	221.8	355.3	611.3	1,103.6	2,333.5
2006 5/.....	38.7	67.9	102.9	158.9	235.0	379.9	648.5	1,159.5	2,453.9
2005 5/.....	39.0	68.6	104.3	165.7	243.0	400.6	686.1	1,224.0	2,634.7
2004 5/.....	40.3	68.8	104.6	166.6	255.2	418.1	711.8	1,263.5	2,640.6
2003 5/.....	41.1	70.7	110.5	176.0	273.7	447.6	761.1	1,343.5	2,828.2
2002.....	42.1	72.2	116.6	178.6	285.9	466.3	801.5	1,381.9	2,908.8
2001.....	39.0	70.6	115.2	188.2	295.0	482.6	831.9	1,408.3	3,002.8
2000.....	41.1	72.0	123.2	189.9	306.1	515.8	847.8	1,475.9	3,211.3
1999.....	41.4	72.8	121.6	196.5	327.2	533.0	889.2	1,526.1	3,276.7

Non-Hispanic White, female

2007 5/.....	10.6	17.8	29.1	52.6	87.2	158.2	308.9	618.0	1,843.0
2006 5/.....	10.6	17.6	30.3	57.1	95.9	177.8	328.4	663.6	1,977.0
2005 5/.....	10.7	18.4	31.6	59.3	105.3	186.8	357.1	711.1	2,133.9
2004 5/.....	10.6	18.1	32.9	61.8	105.3	199.6	367.8	745.7	2,211.1
2003 5/.....	10.9	18.2	34.3	63.0	113.3	211.5	394.7	796.6	2,403.6
2002.....	10.2	17.7	35.8	64.9	118.0	218.6	411.5	826.8	2,500.1
2001.....	9.3	18.8	36.2	68.2	123.3	228.4	427.7	862.3	2,608.1
2000.....	8.9	19.2	37.8	68.9	127.8	233.5	447.0	885.5	2,751.4
1999.....	9.2	20.7	38.9	75.8	134.6	247.9	471.7	923.6	2,852.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic Black, both sexes

2007 5/.....	38.5	75.0	119.6	184.4	258.3	374.8	573.0	859.9	1,709.6
2006 5/.....	41.5	79.1	124.4	195.0	277.0	407.4	609.5	966.9	1,808.1
2005 5/.....	40.4	85.1	131.3	211.8	279.9	419.8	656.8	1,011.2	1,901.2
2004 5/.....	42.2	82.2	130.8	207.4	302.1	442.1	679.0	1,031.1	2,031.7
2003 5/.....	43.7	84.1	133.4	227.2	321.2	485.2	733.2	1,095.8	2,204.8
2002.....	44.8	84.7	146.5	223.9	327.7	480.3	758.1	1,119.5	2,246.7
2001.....	44.6	85.2	145.6	238.5	350.5	505.5	792.5	1,156.4	2,357.3
2000.....	48.5	84.1	154.4	238.9	345.6	535.0	802.0	1,177.1	2,398.0
1999.....	46.4	93.5	160.9	243.2	371.0	554.2	818.9	1,183.2	2,471.9

Non-Hispanic Black, male

2007 5/.....	55.8	112.4	181.2	273.6	373.4	510.4	781.3	1,097.8	1,739.9
2006 5/.....	59.7	118.6	184.9	286.0	399.8	576.2	847.1	1,227.6	1,837.1
2005 5/.....	60.9	128.2	191.9	307.2	392.9	575.5	865.6	1,281.5	1,882.1
2004 5/.....	60.2	121.6	190.2	303.8	416.2	597.5	910.8	1,264.5	2,090.1
2003 5/.....	64.3	125.3	192.1	320.4	444.0	671.0	973.7	1,396.1	2,162.8
2002.....	63.8	123.8	215.0	329.9	452.8	647.2	959.6	1,384.5	2,254.6
2001.....	65.5	127.3	211.7	337.1	475.2	703.4	1,024.4	1,442.4	2,378.1
2000.....	70.5	122.9	231.6	328.9	452.0	713.5	1,029.4	1,448.7	2,463.5
1999.....	68.1	139.6	231.4	348.9	493.3	744.0	1,026.2	1,431.1	2,552.9

Non-Hispanic Black, female

2007 5/.....	23.6	43.3	69.2	114.6	173.8	281.4	447.7	734.9	1,697.0
2006 5/.....	25.6	45.8	74.5	124.3	186.5	291.7	467.6	829.5	1,796.3
2005 5/.....	22.6	48.8	81.3	137.6	196.2	313.3	532.5	868.6	1,908.8
2004 5/.....	26.5	49.0	82.1	132.4	217.5	336.7	541.9	908.3	2,008.6
2003 5/.....	25.8	49.3	85.2	154.6	230.1	359.9	590.4	938.7	2,221.1
2002.....	28.4	51.7	90.4	141.0	234.9	368.4	637.8	981.7	2,243.7
2001.....	26.6	49.4	91.4	161.0	258.3	373.7	654.1	1,010.7	2,349.6
2000.....	29.5	51.1	91.2	167.9	267.2	416.5	665.0	1,040.4	2,373.7
1999.....	27.6	54.5	103.3	160.3	280.3	428.3	694.0	1,060.2	2,441.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins 3/, both sexes

2007.....	9.6	17.7	25.9	36.8	44.5	59.3	86.0	139.0	314.7
2006.....	9.4	17.9	26.1	38.5	47.4	64.4	89.1	147.1	345.5
2005.....	9.6	18.6	26.9	37.7	46.8	64.6	96.5	161.9	376.4
2004.....	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
2003.....	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4

All origins 3/, male

2007.....	15.2	28.2	41.3	56.7	65.8	83.9	115.5	179.7	326.1
2006.....	14.9	29.3	41.5	58.9	69.9	90.6	117.9	188.1	365.3
2005.....	15.3	30.0	42.5	57.7	68.3	91.5	127.0	204.6	390.7
2004.....	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
2003.....	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6

All origins 3/, female

2007.....	4.1	7.6	11.3	18.7	25.8	39.3	64.2	113.4	309.3
2006.....	4.1	7.1	11.5	20.0	27.7	43.1	67.9	121.4	336.1
2005.....	4.0	7.7	12.2	19.4	28.1	42.9	74.3	135.4	369.9
2004.....	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
2003.....	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

Hispanic, both sexes

2007.....	5.7	12.1	20.8	34.9	44.0	50.6	74.1	109.8	210.6
2006.....	6.1	13.0	23.0	35.8	45.2	65.3	81.7	124.7	260.2
2005.....	7.2	15.7	23.8	36.6	48.2	64.2	94.4	143.9	274.7
2004.....	5.9	14.8	22.2	39.4	45.3	69.4	99.0	150.1	279.7
2003.....	7.1	13.3	22.1	39.5	46.6	70.8	105.2	156.2	357.1
2002.....	6.8	11.8	22.7	39.2	49.8	75.4	105.3	155.5	358.6
2001.....	6.2	15.1	23.5	37.9	54.9	79.6	112.1	168.4	417.0
2000.....	6.3	13.8	24.0	38.9	54.1	80.2	101.3	177.1	452.5
1999.....	6.5	15.4	27.1	40.4	62.9	83.7	124.0	190.3	469.4

Hispanic, male

2007.....	9.0	18.8	34.0	55.5	73.2	76.3	95.7	130.9	226.7
2006.....	9.2	23.5	39.2	57.1	70.3	95.7	105.1	161.3	277.4
2005.....	11.6	26.4	40.2	59.0	75.0	93.4	135.6	189.1	308.8
2004.....	9.8	25.7	38.2	60.0	67.6	104.3	147.4	205.7	297.0
2003.....	12.0	24.0	38.2	63.7	74.1	107.6	148.9	204.8	397.1
2002.....	10.7	20.3	40.4	61.8	74.8	111.5	140.1	178.4	415.0
2001.....	10.0	26.6	39.2	59.7	87.2	117.1	159.6	222.0	440.0
2000.....	10.9	24.1	39.9	64.5	74.9	115.2	158.7	225.3	493.8
1999.....	10.6	27.4	42.5	59.6	103.6	131.9	171.6	238.9	485.3

Hispanic, female

2007.....	2.2	5.5	8.2	16.5	19.4	30.4	58.4	95.7	201.9
2006.....	2.9	2.5	7.7	16.9	24.2	41.6	64.8	100.3	251.0
2005.....	2.6	5.2	8.4	16.8	25.8	41.6	64.8	113.8	256.8
2004.....	1.8	4.1	7.3	21.4	26.8	42.5	64.6	113.4	270.8
2003.....	2.1	2.9	7.3	18.4	24.0	42.6	74.1	124.5	336.7
2002.....	2.8	3.5	6.4	19.5	29.3	47.7	80.4	140.9	330.7
2001.....	2.5	4.1	9.2	19.0	28.7	51.0	78.4	135.1	405.8
2000.....	*	4.0	9.5	16.9	37.1	53.8	60.6	147.8	432.3
1999.....	2.5	4.1	13.1	23.9	30.0	47.5	90.3	161.3	461.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 4/, both sexes

2007.....	10.1	18.2	26.2	36.7	44.2	59.6	86.2	140.4	319.6
2006.....	9.8	18.4	26.1	38.5	47.2	63.9	89.1	147.9	349.3
2005.....	9.8	18.9	27.0	37.5	46.3	64.3	96.1	162.2	380.6
2004.....	9.9	18.5	26.7	38.8	49.7	68.4	100.3	166.6	404.7
2003.....	10.1	18.9	26.7	40.1	53.5	72.9	110.2	186.8	445.3
2002.....	10.0	17.8	28.1	39.2	53.7	75.9	116.2	188.3	477.8
2001.....	9.1	17.5	27.7	40.5	54.8	75.9	121.3	197.5	494.8
2000.....	9.6	17.7	28.6	40.3	55.3	81.3	125.0	207.9	535.9
1999.....	9.3	17.4	28.5	43.0	58.9	86.0	130.8	217.9	564.9

Non-Hispanic 4/, male

2007.....	15.9	29.1	41.8	56.3	64.7	83.7	116.0	182.1	330.8
2006.....	15.6	29.7	41.4	58.6	69.3	89.4	118.0	188.9	369.6
2005.....	15.7	30.3	42.4	57.2	67.1	90.8	125.7	204.7	394.0
2004.....	15.9	29.8	42.0	57.9	73.0	94.7	132.1	209.5	418.3
2003.....	16.3	30.9	41.6	59.6	76.8	101.5	145.1	233.3	463.3
2002.....	16.0	28.7	44.5	58.7	78.4	105.3	153.9	233.6	493.3
2001.....	14.9	28.2	43.2	61.4	77.6	106.0	158.8	247.2	503.0
2000.....	15.8	28.4	45.7	60.2	80.4	114.3	165.7	252.5	551.0
1999.....	15.2	27.7	44.8	63.8	85.4	118.6	170.8	263.4	580.2

Non-Hispanic 4/, female

2007.....	4.4	7.8	11.5	18.8	26.1	39.8	64.3	114.1	314.2
2006.....	4.2	7.5	11.8	20.2	27.8	43.0	67.9	122.3	339.9
2005.....	4.2	7.9	12.4	19.5	28.2	42.8	74.6	136.0	374.5
2004.....	4.1	7.7	12.3	21.4	29.4	47.1	77.5	140.3	398.6
2003.....	4.1	7.5	12.6	22.4	33.2	49.8	85.3	158.4	437.4
2002.....	4.2	7.3	12.6	21.4	32.3	52.3	89.7	160.7	471.1
2001.....	3.4	7.2	13.1	21.5	35.1	51.9	95.0	167.7	491.4
2000.....	3.7	7.4	12.5	22.3	33.8	55.2	96.7	181.6	529.8
1999.....	3.6	7.5	13.3	24.0	36.1	60.5	103.0	191.3	558.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic White, both sexes

2007 5/.....	9.4	16.5	23.6	33.3	39.4	54.4	80.4	135.3	317.2
2006 5/.....	8.9	16.4	23.2	34.4	41.9	57.6	82.7	141.3	346.8
2005 5/.....	9.2	16.5	23.6	32.7	41.6	57.8	88.9	155.0	377.8
2004 5/.....	9.0	16.5	23.7	34.4	43.9	62.2	92.4	159.8	399.9
2003 5/.....	9.0	16.9	24.0	34.6	47.1	65.6	101.9	179.7	441.5
2002.....	9.3	15.8	24.6	34.4	46.6	68.5	107.0	180.7	473.5
2001.....	8.1	15.4	23.6	34.4	46.7	68.0	111.2	190.8	490.7
2000.....	8.5	15.7	24.5	34.3	48.6	72.3	115.5	198.2	530.2
1999.....	8.0	14.9	24.2	36.6	51.0	76.8	120.3	209.3	558.8

Non-Hispanic White, male

2007 5/.....	15.1	26.8	37.8	51.2	58.4	76.5	108.2	175.4	330.0
2006 5/.....	14.4	26.9	37.0	52.2	61.4	80.3	109.5	180.6	366.7
2005 5/.....	14.9	27.0	37.7	50.1	60.3	81.2	116.7	194.4	396.1
2004 5/.....	14.9	27.3	37.5	51.2	65.3	87.0	121.5	201.6	412.6
2003 5/.....	14.9	27.9	37.9	51.8	68.1	90.5	134.7	224.0	459.5
2002.....	15.1	26.0	39.3	52.0	68.2	96.0	142.1	222.9	489.2
2001.....	13.6	25.3	37.7	52.9	66.9	95.2	146.4	239.6	499.4
2000.....	14.2	25.6	39.5	52.4	71.8	102.1	152.6	240.8	544.7
1999.....	13.4	24.2	38.7	55.0	75.0	106.5	157.5	252.7	572.8

Non-Hispanic White, female

2007 5/.....	3.8	6.5	9.8	16.4	22.3	35.8	59.5	109.7	311.1
2006 5/.....	3.5	6.1	9.8	17.8	24.5	38.7	62.6	116.5	337.5
2005 5/.....	3.5	6.3	10.0	16.4	24.9	38.4	68.4	130.4	369.4
2004 5/.....	3.1	5.9	10.4	18.8	24.9	41.7	71.1	134.0	394.2
2003 5/.....	3.3	6.2	10.7	18.6	28.5	45.1	78.1	152.5	433.5
2002.....	3.5	5.8	10.5	18.0	27.5	46.0	81.9	154.9	466.8
2001.....	2.6	5.7	10.1	17.3	28.8	45.9	86.3	161.3	487.0
2000.....	2.8	5.9	10.2	17.4	28.2	48.3	89.5	172.9	524.4
1999.....	2.8	5.9	10.3	19.7	30.0	53.1	94.2	183.6	553.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic Black, both sexes

2007 5/.....	16.5	33.2	50.9	74.3	93.2	116.3	157.5	216.6	402.7
2006 5/.....	17.0	35.6	53.0	79.7	98.6	128.0	168.3	247.5	433.7
2005 5/.....	15.9	38.4	57.6	84.4	93.2	131.9	185.1	266.3	475.2
2004 5/.....	17.8	36.2	56.3	82.3	105.5	140.1	200.1	271.4	509.2
2003 5/.....	17.7	36.3	53.6	92.0	113.3	150.1	215.7	286.5	546.7
2002.....	17.3	35.9	60.8	86.1	120.6	158.1	233.1	303.8	580.4
2001.....	17.1	36.3	64.5	95.5	129.4	164.1	244.8	300.8	595.9
2000.....	19.7	36.9	66.3	94.4	120.8	178.9	240.8	342.8	652.2
1999.....	19.5	39.1	68.6	98.4	135.0	187.7	260.8	339.9	670.3

Non-Hispanic Black, male

2007 5/.....	25.2	50.9	81.2	116.2	139.9	173.5	225.2	305.8	415.3
2006 5/.....	25.2	55.4	83.0	128.3	154.0	193.1	238.9	332.2	471.0
2005 5/.....	23.5	59.8	88.7	131.8	141.8	202.9	253.5	364.8	452.0
2004 5/.....	25.5	54.6	88.7	129.6	154.1	198.5	280.3	353.9	540.6
2003 5/.....	26.8	58.7	81.8	139.3	163.8	229.2	302.8	384.9	588.1
2002.....	26.0	54.5	97.1	131.4	182.2	220.6	321.1	420.4	603.9
2001.....	26.3	56.7	97.3	145.3	185.0	241.7	332.8	385.9	609.8
2000.....	30.5	57.2	107.3	137.8	170.0	259.7	339.1	429.1	711.6
1999.....	30.5	60.8	105.2	146.7	198.2	267.3	346.6	436.6	714.3

Non-Hispanic Black, female

2007 5/.....	8.9	18.2	26.1	41.5	59.0	77.0	116.8	169.7	397.5
2006 5/.....	9.8	18.8	28.3	41.9	57.7	83.4	126.2	202.9	418.5
2005 5/.....	9.2	20.3	31.9	47.5	57.1	83.3	144.3	214.4	484.5
2004 5/.....	11.1	20.6	29.7	45.5	69.4	100.5	152.7	227.9	496.7
2003 5/.....	9.8	17.4	30.4	55.2	75.8	96.8	164.0	235.0	530.7
2002.....	9.7	20.1	31.1	50.8	75.0	116.1	180.5	243.1	571.5
2001.....	9.2	18.9	37.6	56.4	88.3	112.4	192.4	257.4	590.7
2000.....	10.4	19.5	32.7	60.2	84.5	125.2	181.5	299.4	630.2
1999.....	10.1	20.7	38.7	60.4	88.2	134.9	209.1	291.9	653.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes

2007.....	14.6	25.2	41.7	69.0	113.8	194.5	354.6	654.7	1,613.2
2006.....	15.7	26.8	43.9	73.6	121.9	212.5	379.0	700.4	1,712.0
2005.....	15.5	27.7	44.9	79.9	131.2	226.9	404.6	740.4	1,837.8
2004.....	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
2003.....	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6

All origins 3/, male

2007.....	22.2	38.9	63.6	101.5	165.2	274.9	492.2	890.8	1,888.6
2006.....	23.9	41.2	66.6	107.4	174.8	296.7	531.1	945.4	1,979.5
2005.....	23.7	42.4	67.0	116.4	184.3	316.4	558.3	997.9	2,120.9
2004.....	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
2003.....	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4

All origins 3/, female

2007.....	7.1	12.2	21.1	39.1	68.8	128.8	253.1	505.6	1,482.2
2006.....	7.7	12.9	22.6	42.7	75.6	144.0	267.7	547.2	1,586.8
2005.....	7.4	13.5	23.9	46.5	84.8	154.5	293.0	580.4	1,707.7
2004.....	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
2003.....	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Hispanic, both sexes

2007.....	8.7	15.7	28.2	54.9	95.7	161.3	307.1	551.1	1,226.4
2006.....	9.4	19.0	30.2	54.9	94.4	177.7	332.5	595.5	1,335.0
2005.....	7.4	18.3	31.5	59.1	108.2	205.9	350.9	666.9	1,455.8
2004.....	9.7	16.6	31.0	61.4	110.4	200.5	361.7	641.8	1,516.0
2003.....	8.2	17.3	33.5	67.6	115.3	214.6	391.4	689.4	1,655.7
2002.....	10.3	20.3	36.8	68.9	130.9	226.8	400.2	720.1	1,771.1
2001.....	9.0	19.7	37.9	66.9	136.8	241.9	437.7	799.6	2,009.6
2000.....	9.9	20.1	43.6	75.3	139.0	240.5	437.7	824.9	2,056.3
1999.....	10.4	24.2	45.5	79.4	147.2	259.3	455.8	846.4	2,171.4

Hispanic, male

2007.....	14.5	24.3	43.8	80.8	137.7	211.9	390.2	684.9	1,255.3
2006.....	14.2	28.7	43.1	77.4	131.5	230.0	450.1	719.0	1,427.7
2005.....	10.8	28.0	45.2	83.3	150.2	277.3	458.6	824.8	1,503.1
2004.....	14.7	23.2	44.0	88.8	157.7	274.9	482.6	792.7	1,556.5
2003.....	11.7	26.5	50.0	90.6	162.2	290.4	511.0	846.2	1,568.6
2002.....	15.9	29.8	53.4	100.3	182.5	310.7	526.6	884.2	1,847.2
2001.....	14.1	28.7	59.1	93.2	190.3	327.1	576.3	994.7	2,039.8
2000.....	16.0	31.3	64.5	108.8	195.8	341.8	563.3	1,032.1	2,084.0
1999.....	17.0	39.2	70.7	111.4	197.4	356.4	582.8	1,053.1	2,221.4

Hispanic, female

2007.....	2.5	7.1	13.3	31.7	60.2	121.7	246.8	461.8	1,210.8
2006.....	4.3	9.3	18.0	35.0	63.2	136.9	247.4	513.4	1,285.7
2005.....	3.9	8.6	18.6	37.7	73.1	150.5	273.5	562.2	1,431.1
2004.....	4.5	10.1	18.9	37.4	71.2	143.1	275.7	542.1	1,495.2
2003.....	4.8	8.5	18.2	47.6	76.8	156.4	306.5	587.4	1,700.0
2002.....	4.6	11.1	21.6	41.6	88.7	162.2	309.9	615.4	1,733.3
2001.....	3.9	11.1	18.6	44.2	93.5	176.9	339.4	678.8	1,994.8
2000.....	3.7	9.5	24.8	46.4	93.0	163.9	348.7	698.8	2,042.7
1999.....	3.9	9.9	22.7	51.8	106.6	186.2	366.2	723.0	2,147.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 4/, both sexes

2007.....	15.3	26.2	42.8	70.0	115.1	196.8	357.5	660.2	1,632.5
2006.....	16.5	27.5	45.0	75.0	123.8	214.6	381.5	705.4	1,728.7
2005.....	16.4	28.5	45.9	81.4	132.6	228.2	407.1	742.9	1,853.4
2004.....	17.1	28.3	46.7	81.0	136.3	238.3	419.8	773.9	1,892.8
2003.....	17.6	28.9	50.0	86.6	146.0	255.6	446.6	814.6	2,039.6
2002.....	17.7	30.6	53.3	89.2	153.8	263.0	467.1	844.5	2,093.4
2001.....	17.1	30.6	53.4	95.1	161.1	276.4	483.9	867.0	2,177.4
2000.....	17.6	31.0	57.4	96.8	166.3	288.8	497.1	894.0	2,288.1
1999.....	17.9	33.3	57.9	100.5	176.8	300.2	522.0	924.3	2,351.4

Non-Hispanic 4/, male

2007.....	23.3	40.4	65.3	103.0	167.1	279.2	498.7	902.7	1,924.3
2006.....	25.1	42.5	68.5	109.6	177.8	300.8	535.6	957.6	2,007.1
2005.....	25.3	43.7	68.7	118.9	186.3	318.7	563.2	1,005.7	2,150.8
2004.....	26.1	43.4	69.3	118.5	193.2	330.7	584.9	1,037.4	2,154.8
2003.....	27.1	44.4	74.8	127.1	208.9	358.0	620.5	1,097.6	2,287.4
2002.....	27.7	47.8	79.6	131.3	219.0	369.4	650.8	1,131.1	2,338.9
2001.....	26.5	46.9	79.9	138.0	230.1	387.4	677.7	1,147.3	2,420.7
2000.....	27.9	47.7	86.0	140.8	235.3	411.0	687.7	1,207.7	2,572.9
1999.....	28.6	50.8	85.6	145.0	252.0	426.1	721.4	1,243.7	2,619.9

Non-Hispanic 4/, female

2007.....	7.7	12.7	21.7	39.7	69.4	129.1	253.4	507.6	1,494.7
2006.....	8.2	13.2	22.9	43.2	76.5	144.2	268.7	548.3	1,599.3
2005.....	7.8	14.0	24.3	47.1	85.6	154.6	293.8	580.3	1,717.8
2004.....	8.4	13.8	25.5	46.9	86.6	163.7	301.1	612.1	1,775.7
2003.....	8.5	14.2	26.6	49.6	91.3	173.2	322.7	642.1	1,930.7
2002.....	8.1	14.1	28.7	51.0	97.2	177.8	337.4	670.6	1,987.5
2001.....	8.0	14.9	28.4	56.1	101.5	188.0	348.3	699.3	2,076.2
2000.....	7.6	15.0	30.7	56.9	106.9	192.4	364.5	709.1	2,172.9
1999.....	7.7	16.6	32.1	60.1	112.2	201.6	383.4	737.6	2,243.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic White, both sexes

2007 5/.....	14.8	24.9	40.5	67.1	111.5	193.7	358.4	671.8	1,683.4
2006 5/.....	15.7	26.2	42.7	71.9	119.7	212.2	382.7	713.9	1,781.9
2005 5/.....	15.6	26.7	43.7	78.0	128.7	226.0	408.2	753.3	1,913.3
2004 5/.....	16.4	26.7	44.4	78.0	132.0	236.2	420.8	783.9	1,944.1
2003 5/.....	16.9	27.2	47.7	82.9	141.7	252.4	446.6	825.1	2,091.9
2002.....	16.7	28.9	50.8	85.3	150.3	261.5	467.2	856.7	2,149.8
2001.....	16.0	29.1	51.3	91.6	157.1	274.4	484.0	877.0	2,233.4
2000.....	16.4	29.6	55.0	92.8	162.6	287.1	496.9	907.4	2,353.3
1999.....	17.1	31.5	55.1	97.1	173.5	297.5	523.6	937.9	2,414.4

Non-Hispanic White, male

2007 5/.....	22.9	38.8	62.4	100.0	163.5	278.8	503.1	928.2	2,003.4
2006 5/.....	24.3	41.0	65.9	106.7	173.5	299.7	539.0	978.9	2,087.2
2005 5/.....	24.1	41.6	66.6	115.6	182.7	319.4	569.3	1,029.6	2,238.6
2004 5/.....	25.4	41.5	67.1	115.5	189.9	331.1	590.3	1,062.0	2,228.0
2003 5/.....	26.3	42.8	72.7	124.2	205.6	357.1	626.4	1,119.6	2,368.6
2002.....	27.0	46.2	77.2	126.6	217.7	370.3	659.5	1,158.9	2,419.7
2001.....	25.4	45.4	77.5	135.4	228.0	387.4	685.5	1,168.6	2,503.4
2000.....	26.9	46.3	83.7	137.5	234.2	413.6	695.2	1,235.1	2,666.6
1999.....	28.0	48.6	82.9	141.5	252.2	426.5	731.7	1,273.5	2,703.9

Non-Hispanic White, female

2007 5/.....	6.8	11.3	19.3	36.2	64.9	122.4	249.4	508.3	1,531.9
2006 5/.....	7.1	11.6	20.4	39.3	71.4	139.1	265.8	547.1	1,639.5
2005 5/.....	7.1	12.1	21.6	42.8	80.4	148.4	288.7	580.7	1,764.6
2004 5/.....	7.5	12.1	22.5	43.0	80.5	157.9	296.7	611.7	1,816.9
2003 5/.....	7.6	12.0	23.6	44.4	84.8	166.4	316.6	644.1	1,970.1
2002.....	6.7	12.0	25.3	46.9	90.6	172.6	329.6	671.9	2,033.3
2001.....	6.7	13.1	26.2	51.0	94.5	182.5	341.4	701.0	2,121.0
2000.....	6.1	13.2	27.6	51.5	99.6	185.3	357.5	712.6	2,227.0
1999.....	6.4	14.8	28.6	56.1	104.6	194.7	377.5	739.9	2,299.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic Black, both sexes

2007 5/.....	22.1	41.8	68.7	110.1	165.1	258.4	415.5	643.3	1,306.9
2006 5/.....	24.5	43.5	71.4	115.4	178.5	279.4	441.1	719.4	1,374.4
2005 5/.....	24.6	46.7	73.7	127.4	186.7	287.9	471.7	744.8	1,426.0
2004 5/.....	24.4	46.0	74.5	125.1	196.7	302.0	478.9	759.7	1,522.6
2003 5/.....	26.0	47.8	79.9	135.1	207.9	335.1	517.5	809.4	1,658.1
2002.....	27.6	48.9	85.7	137.7	207.0	322.3	525.0	815.8	1,666.3
2001.....	27.5	48.9	81.1	143.0	221.1	341.4	547.6	855.6	1,761.4
2000.....	28.8	47.3	88.1	144.5	224.8	356.1	561.3	834.3	1,745.8
1999.....	26.8	54.4	92.3	144.8	236.0	366.5	558.1	843.3	1,801.6

Non-Hispanic Black, male

2007 5/.....	30.6	61.5	100.0	157.3	233.5	337.0	556.1	791.9	1,324.6
2006 5/.....	34.5	63.2	101.9	157.7	245.8	383.0	608.1	895.4	1,366.1
2005 5/.....	37.4	68.4	103.2	175.5	251.1	372.6	612.1	916.7	1,430.2
2004 5/.....	34.7	67.0	101.5	174.2	262.1	399.0	630.5	910.6	1,549.6
2003 5/.....	37.5	66.6	110.4	181.0	280.2	441.8	671.0	1,011.2	1,574.7
2002.....	37.8	69.4	117.8	198.6	270.6	426.6	638.5	964.1	1,650.8
2001.....	39.2	70.5	114.5	191.9	290.3	461.7	691.6	1,056.4	1,768.3
2000.....	40.0	65.6	124.3	191.1	282.0	453.8	690.3	1,019.6	1,751.9
1999.....	37.7	78.8	126.1	202.2	295.1	476.7	679.6	994.5	1,838.6

Non-Hispanic Black, female

2007 5/.....	14.7	25.1	43.0	73.2	114.9	204.4	330.9	565.2	1,299.5
2006 5/.....	15.8	27.0	46.2	82.5	128.8	208.3	341.4	626.6	1,377.8
2005 5/.....	13.4	28.5	49.4	90.1	139.1	230.0	388.1	654.2	1,424.3
2004 5/.....	15.4	28.4	52.4	86.9	148.2	236.2	389.2	680.4	1,511.9
2003 5/.....	15.9	31.9	54.8	99.4	154.3	263.1	426.4	703.7	1,690.4
2002.....	18.7	31.6	59.4	90.2	160.0	252.3	457.2	738.6	1,672.2
2001.....	17.4	30.5	53.8	104.7	170.0	261.3	461.7	753.4	1,758.9
2000.....	19.2	31.6	58.5	107.7	182.6	291.3	483.5	741.0	1,743.5
1999.....	17.5	33.7	64.7	99.9	192.1	293.4	484.9	768.3	1,787.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

All origins 3/, both sexes

2007.....	16.9	24.8	35.7	53.5	86.5	143.6	256.2	494.3	1,329.8
2006.....	17.7	25.0	36.3	55.5	85.7	144.6	259.0	493.3	1,342.8
2005.....	17.9	25.5	36.8	57.4	88.2	150.8	264.6	500.7	1,377.4
2004.....	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
2003.....	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9

All origins 3/, male

2007.....	21.4	32.1	46.4	67.2	106.6	175.1	302.4	569.6	1,348.9
2006.....	22.6	32.4	46.6	69.5	106.9	173.2	302.8	566.0	1,338.1
2005.....	22.6	33.3	47.4	71.3	108.0	180.8	310.3	568.1	1,377.8
2004.....	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
2003.....	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9

All origins 3/, female

2007.....	12.6	17.9	25.6	40.9	68.9	117.8	222.1	446.8	1,320.8
2006.....	12.9	18.0	26.6	42.7	67.2	121.3	226.9	447.9	1,345.0
2005.....	13.3	18.0	26.9	44.8	70.9	126.4	231.4	458.9	1,377.3
2004.....	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
2003.....	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

Hispanic, both sexes

2007	9.1	13.4	19.7	30.2	51.0	88.2	159.2	281.3	659.7
2006	10.0	14.3	19.3	30.5	48.5	92.0	144.6	276.2	663.2
2005	9.3	12.9	19.5	32.3	53.9	92.2	159.6	289.8	703.3
2004	9.9	13.4	17.8	34.5	54.1	91.2	158.8	286.4	659.3
2003	10.2	12.8	24.0	37.7	61.4	97.5	175.0	296.3	749.9
2002	9.2	14.2	21.4	31.4	57.4	91.6	167.4	307.5	740.0
2001	8.9	13.5	21.3	36.8	61.7	92.5	168.2	304.0	798.5
2000	8.5	15.6	23.3	35.8	58.2	98.1	178.7	314.7	829.4
1999	9.1	15.0	23.6	39.4	61.5	100.3	173.0	332.5	827.9

Hispanic, male

2007	12.7	17.3	25.1	37.9	61.3	106.3	188.9	303.7	679.0
2006	12.2	19.0	25.0	38.9	59.3	114.8	162.2	341.5	658.6
2005	12.6	18.0	25.2	41.6	65.8	115.6	205.2	316.1	697.4
2004	13.6	18.0	22.5	44.9	64.2	109.7	181.4	340.7	641.6
2003	13.9	15.7	29.3	48.2	75.4	120.9	202.3	341.0	752.7
2002	11.2	18.4	26.6	38.7	69.8	122.0	202.2	354.6	758.0
2001	12.5	18.3	25.4	45.9	75.5	111.2	190.3	359.7	775.2
2000	10.7	20.9	31.3	44.9	70.8	124.5	194.9	362.2	890.8
1999	11.6	20.5	29.1	51.2	79.6	132.4	213.0	377.2	888.6

Hispanic, female

2007	5.3	9.5	14.6	23.3	42.4	74.0	137.6	266.3	649.2
2006	7.7	9.6	14.0	23.0	39.5	74.2	131.9	232.7	665.6
2005	5.9	7.9	14.1	24.1	44.0	74.0	126.9	272.4	706.4
2004	6.0	9.0	13.4	25.4	45.7	77.0	142.8	250.5	668.4
2003	6.4	9.9	19.2	28.5	49.9	79.7	155.6	267.3	748.4
2002	7.2	10.2	16.7	25.0	47.3	68.3	142.6	277.4	731.1
2001	5.3	8.9	17.6	28.9	50.5	78.2	152.5	269.4	809.8
2000	6.3	10.5	16.1	28.0	48.0	78.1	167.3	285.8	799.3
1999	6.6	9.7	18.6	29.3	46.9	76.2	144.8	305.7	798.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

Non-Hispanic 4/, both sexes

2007.....	17.9	26.0	37.1	55.3	89.2	147.7	262.7	506.8	1,364.4
2006.....	18.7	26.1	37.7	57.5	88.4	148.2	266.1	505.0	1,375.6
2005.....	18.8	26.7	38.2	59.3	90.7	154.8	270.8	511.4	1,408.5
2004.....	18.8	26.7	37.1	59.2	92.9	155.8	274.1	513.5	1,404.5
2003.....	18.7	26.4	40.2	61.2	99.8	164.0	284.5	535.2	1,453.7
2002.....	19.0	26.8	40.8	62.7	100.6	168.7	292.4	545.0	1,469.2
2001.....	18.7	26.9	41.1	63.8	103.1	168.7	292.4	545.3	1,497.8
2000.....	18.4	26.7	41.3	66.1	106.4	170.6	294.7	550.4	1,530.9
1999.....	18.6	26.7	43.0	68.2	107.7	178.5	300.0	560.2	1,533.9

Non-Hispanic 4/, male

2007.....	22.5	33.6	48.2	69.5	109.9	180.0	310.0	585.8	1,387.0
2006.....	23.9	33.7	48.3	71.9	110.2	177.0	311.5	578.4	1,374.1
2005.....	23.7	34.8	49.1	73.4	110.9	185.2	316.3	581.4	1,412.5
2004.....	24.2	34.2	48.3	74.6	114.6	185.8	323.6	585.1	1,412.5
2003.....	23.6	33.8	51.5	76.5	122.4	199.8	340.8	610.9	1,460.5
2002.....	24.8	34.8	51.1	78.4	122.0	204.1	347.2	622.6	1,477.3
2001.....	24.0	33.8	51.2	79.7	124.2	205.6	351.5	632.2	1,499.0
2000.....	24.1	34.3	53.3	83.0	130.2	209.8	351.4	636.0	1,573.6
1999.....	24.0	34.2	55.7	86.2	136.5	220.7	356.0	649.4	1,571.2

Non-Hispanic 4/, female

2007.....	13.5	18.7	26.5	42.3	71.0	121.1	227.9	457.0	1,353.8
2006.....	13.6	18.8	27.7	44.3	69.4	124.7	232.9	459.2	1,376.4
2005.....	14.1	19.0	27.9	46.4	73.0	130.1	237.8	468.0	1,406.6
2004.....	13.5	19.5	26.7	45.2	73.9	131.7	238.6	469.6	1,400.9
2003.....	13.9	19.3	29.6	47.2	80.1	135.2	244.3	489.0	1,450.7
2002.....	13.3	19.2	31.1	48.4	82.1	140.3	253.8	497.9	1,465.7
2001.....	13.4	20.3	31.5	49.3	84.9	139.2	251.1	493.4	1,497.3
2000.....	12.8	19.3	30.1	50.8	86.0	139.6	255.3	500.0	1,513.7
1999.....	13.3	19.6	31.0	51.9	83.0	145.5	261.1	508.0	1,519.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

Non-Hispanic White, both sexes

2007 5/.....	15.3	22.1	32.3	50.2	82.3	141.9	260.7	511.6	1,410.6
2006 5/.....	15.9	22.1	33.1	52.6	82.4	142.7	264.3	509.0	1,422.6
2005 5/.....	15.9	22.5	33.4	53.2	84.0	149.1	267.1	515.6	1,453.0
2004 5/.....	15.7	22.6	32.4	54.1	87.0	150.3	268.9	518.4	1,442.1
2003 5/.....	15.8	22.2	35.6	55.0	92.8	156.6	281.8	539.1	1,490.1
2002.....	15.6	22.8	36.0	57.1	94.9	162.5	288.2	548.1	1,503.1
2001.....	15.4	23.0	35.7	58.7	96.3	162.8	286.7	547.4	1,533.2
2000.....	15.2	22.6	36.7	60.6	99.8	163.9	288.8	553.5	1,564.2
1999.....	15.1	22.8	37.2	61.7	100.8	171.0	293.1	560.5	1,565.9

Non-Hispanic White, male

2007 5/.....	19.5	28.9	42.1	63.1	101.8	172.8	307.6	590.8	1,440.1
2006 5/.....	20.6	28.6	42.8	65.6	103.2	169.9	309.4	585.4	1,420.8
2005 5/.....	20.4	29.8	43.1	66.4	102.8	177.4	312.3	589.8	1,462.5
2004 5/.....	20.3	29.6	42.4	68.3	107.7	180.5	317.0	590.1	1,445.4
2003 5/.....	20.3	28.7	45.6	69.7	114.2	192.5	338.7	616.3	1,499.3
2002.....	20.8	30.1	45.5	72.3	115.5	197.0	343.3	627.9	1,512.9
2001.....	20.3	29.7	44.8	73.9	117.3	198.2	344.9	632.2	1,530.7
2000.....	20.5	29.2	47.9	76.8	122.7	203.1	345.6	640.2	1,606.4
1999.....	20.1	29.4	48.8	78.5	128.2	212.7	349.2	649.4	1,595.5

Non-Hispanic White, female

2007 5/.....	11.1	15.6	22.8	38.0	64.7	116.0	225.3	461.1	1,396.7
2006 5/.....	11.4	15.8	23.7	40.3	63.8	120.1	230.6	460.8	1,423.5
2005 5/.....	11.4	15.3	24.1	40.8	67.2	125.5	233.5	469.3	1,448.6
2004 5/.....	11.0	15.7	22.7	40.9	68.6	125.3	233.7	474.0	1,440.6
2003 5/.....	11.3	15.8	25.9	41.4	73.7	127.1	240.5	491.6	1,486.1
2002.....	10.5	15.6	27.0	43.1	76.7	134.2	248.8	499.2	1,498.8
2001.....	10.6	16.5	27.1	44.6	77.7	134.0	245.4	496.2	1,534.2
2000.....	9.9	16.2	26.1	45.6	79.8	132.3	248.8	502.0	1,547.2
1999.....	10.1	16.3	26.2	46.2	76.7	137.8	253.7	508.0	1,554.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

Non-Hispanic Black, both sexes

2007 5/.....	38.5	58.9	81.9	113.3	170.5	238.1	352.3	564.2	1,130.2
2006 5/.....	39.6	58.8	82.0	114.3	158.8	232.9	348.4	577.3	1,111.9
2005 5/.....	41.0	61.8	84.0	123.5	165.8	242.6	375.1	571.9	1,152.4
2004 5/.....	42.6	61.9	84.7	117.3	163.2	245.2	388.1	560.2	1,197.9
2003 5/.....	41.9	62.4	87.6	126.3	180.0	269.9	370.6	582.3	1,253.2
2002.....	44.8	62.1	89.0	123.6	172.3	260.2	391.1	601.9	1,285.5
2001.....	44.0	62.3	93.6	121.1	182.4	258.0	408.2	608.1	1,292.9
2000.....	44.3	63.6	87.8	125.0	185.1	270.0	412.7	597.8	1,319.4
1999.....	45.8	63.5	98.4	135.9	188.6	287.1	420.9	627.5	1,314.0

Non-Hispanic Black, male

2007 5/.....	47.2	76.3	109.8	148.1	219.0	300.5	422.1	676.8	1,085.5
2006 5/.....	51.3	78.4	105.0	151.1	203.0	293.1	418.0	658.7	1,122.4
2005 5/.....	50.8	79.8	110.3	155.0	209.4	307.6	453.7	623.8	1,107.6
2004 5/.....	55.7	77.1	111.8	153.1	207.0	289.3	478.5	663.8	1,244.7
2003 5/.....	52.2	80.9	116.6	159.6	229.1	322.9	442.0	673.2	1,239.1
2002.....	58.3	78.8	112.9	153.6	211.9	322.9	457.7	685.6	1,258.0
2001.....	54.7	75.2	120.1	154.0	216.3	322.3	501.0	753.1	1,362.9
2000.....	55.6	84.4	114.4	156.1	229.2	321.9	487.5	704.6	1,416.1
1999.....	57.3	83.5	128.8	175.5	241.5	355.6	491.1	756.3	1,466.6

Non-Hispanic Black, female

2007 5/.....	30.9	44.1	59.0	86.1	134.8	195.1	310.4	505.1	1,148.7
2006 5/.....	29.6	42.3	63.1	85.7	126.1	191.5	306.8	534.4	1,107.6
2005 5/.....	32.5	46.5	62.2	99.0	133.5	198.2	328.3	544.6	1,170.4
2004 5/.....	31.3	49.1	62.5	89.4	130.8	215.2	334.6	505.7	1,179.4
2003 5/.....	32.9	46.7	63.7	100.3	143.6	234.2	328.2	534.7	1,258.7
2002.....	33.1	48.1	69.3	100.2	142.9	218.2	351.3	558.4	1,296.0
2001.....	34.8	51.3	71.9	95.3	157.3	215.2	352.8	534.3	1,267.0
2000.....	34.5	45.9	66.0	100.5	152.6	235.5	367.6	544.0	1,283.4
1999.....	35.8	46.5	73.6	105.0	149.4	241.7	378.6	563.7	1,257.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and subacute endocarditis (I33)

All origins 3/, both sexes

2007.....	0.4	0.6	0.6	0.7	1.1	1.4	2.1	2.8	2.4
2006.....	0.4	0.5	0.7	0.8	1.0	1.4	2.2	2.7	2.2
2005.....	0.4	0.5	0.6	0.7	1.1	1.4	2.4	2.7	3.0
2004.....	0.4	0.5	0.5	0.7	1.0	1.5	2.1	2.9	3.2
2003.....	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9

All origins 3/, male

2007.....	0.4	0.8	0.8	0.9	1.3	1.7	2.9	3.8	3.4
2006.....	0.6	0.7	0.9	0.9	1.2	1.6	2.3	3.3	2.3
2005.....	0.4	0.7	0.8	0.9	1.2	1.8	3.1	3.5	5.1
2004.....	0.5	0.7	0.6	0.8	1.4	1.4	2.3	4.2	4.8
2003.....	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7

All origins 3/, female

2007.....	0.3	0.4	0.4	0.5	0.9	1.2	1.5	2.1	2.0
2006.....	0.3	0.3	0.4	0.7	0.7	1.2	2.2	2.3	2.1
2005.....	0.3	0.3	0.4	0.6	1.0	1.1	1.9	2.2	2.1
2004.....	0.3	0.4	0.3	0.6	0.7	1.7	2.0	2.1	2.5
2003.....	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and subacute endocarditis (I33)

Non-Hispanic 4/, both sexes

2007.....	0.4	0.5	0.6	0.7	1.1	1.5	2.1	2.8	2.5
2006.....	0.5	0.5	0.7	0.8	1.0	1.4	2.2	2.8	2.1
2005.....	0.4	0.5	0.6	0.7	1.1	1.4	2.4	2.7	3.1
2004.....	0.4	0.5	0.5	0.6	1.0	1.6	2.1	2.9	3.2
2003.....	0.4	0.5	0.6	1.0	1.0	1.6	2.3	2.7	3.0
2002.....	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.7
2001.....	0.5	0.5	0.5	0.8	1.1	1.7	2.2	3.1	2.6
2000.....	0.5	0.5	0.7	0.7	1.1	1.3	2.2	2.8	2.9
1999.....	0.4	0.5	0.7	0.7	1.1	1.5	2.1	2.8	2.9

Non-Hispanic 4/, male

2007.....	0.4	0.8	0.8	1.0	1.3	1.8	2.9	3.9	3.6
2006.....	0.6	0.7	1.0	1.0	1.2	1.6	2.3	3.5	2.3
2005.....	0.5	0.7	0.8	0.9	1.1	1.7	3.1	3.6	5.3
2004.....	0.5	0.7	0.6	0.7	1.5	1.4	2.4	4.2	4.8
2003.....	0.6	0.7	0.8	1.2	1.1	1.7	3.0	3.9	4.1
2002.....	0.5	0.7	0.6	0.9	1.5	1.7	2.6	3.7	3.6
2001.....	0.7	0.7	0.6	1.1	1.2	2.2	3.1	4.5	3.9
2000.....	0.6	0.7	0.8	0.9	1.3	1.5	3.2	4.1	4.2
1999.....	0.5	0.7	1.0	0.8	1.2	1.6	2.7	4.0	4.7

Non-Hispanic 4/, female

2007.....	0.3	0.4	0.4	0.5	1.0	1.2	1.5	2.1	2.0
2006.....	0.3	0.3	0.4	0.7	0.8	1.3	2.1	2.3	2.1
2005.....	0.3	0.3	0.4	0.6	1.1	1.2	1.9	2.2	2.2
2004.....	0.3	0.4	0.3	0.6	0.6	1.7	1.9	2.1	2.6
2003.....	0.3	0.3	0.4	0.8	0.8	1.5	1.9	2.0	2.5
2002.....	0.3	0.3	0.4	0.8	0.8	1.4	1.9	2.1	2.3
2001.....	0.3	0.3	0.4	0.6	1.1	1.3	1.5	2.2	2.0
2000.....	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	0.3	0.5	0.7	1.0	1.4	1.7	2.1	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and subacute endocarditis (I33)

Non-Hispanic White, both sexes

2007 5/.....	0.3	0.4	0.5	0.7	1.0	1.3	2.0	3.0	2.5
2006 5/.....	0.4	0.4	0.6	0.7	1.0	1.4	2.0	2.9	2.1
2005 5/.....	0.3	0.4	0.4	0.6	1.0	1.5	2.3	2.7	3.2
2004 5/.....	0.3	0.4	0.4	0.5	1.0	1.5	2.0	3.0	3.3
2003 5/.....	0.3	0.4	0.5	0.8	0.9	1.4	2.3	2.8	3.2
2002.....	0.2	0.4	0.3	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.1	2.7
2000.....	0.4	0.4	0.5	0.7	1.1	1.2	2.2	2.8	2.8
1999.....	0.3	0.3	0.5	0.7	1.0	1.4	2.1	2.8	3.0

Non-Hispanic White, male

2007 5/.....	*	0.6	0.6	0.9	1.1	1.6	2.8	4.0	3.6
2006 5/.....	0.6	0.5	0.8	0.8	1.1	1.7	2.2	3.5	2.5
2005 5/.....	0.4	0.5	0.5	0.7	1.1	1.7	3.2	3.6	5.6
2004 5/.....	0.3	0.5	0.6	0.6	1.3	1.4	2.3	4.3	4.9
2003 5/.....	0.4	0.5	0.6	1.1	1.1	1.6	2.9	4.0	4.5
2002.....	0.3	0.6	0.4	0.9	1.4	1.6	2.7	3.9	3.7
2001.....	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.5	4.0
2000.....	0.5	0.5	0.7	0.9	1.2	1.6	3.3	4.0	4.3
1999.....	0.4	0.4	0.7	0.8	1.2	1.5	2.9	4.0	5.1

Non-Hispanic White, female

2007 5/.....	0.3	0.3	0.4	0.4	0.9	1.1	1.4	2.3	2.0
2006 5/.....	0.3	*	0.3	0.6	0.8	1.2	1.9	2.5	1.9
2005 5/.....	0.3	*	*	0.5	0.9	1.3	1.7	2.0	2.1
2004 5/.....	*	*	*	0.5	0.6	1.6	1.8	2.1	2.5
2003 5/.....	*	*	0.4	0.6	0.8	1.3	1.8	2.0	2.7
2002.....	*	*	*	0.7	0.7	1.3	1.8	2.3	2.3
2001.....	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000.....	*	*	0.4	0.5	1.0	1.0	1.5	2.0	2.2
1999.....	*	*	*	0.6	0.8	1.3	1.6	2.0	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All origins 3/, both sexes

2007.....	0.3	0.3	0.4	0.4	0.6	0.7	1.0	1.1	2.1
2006.....	0.2	0.3	0.3	0.5	0.4	0.8	1.1	1.3	1.8
2005.....	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.9
2004.....	0.2	0.3	0.3	0.5	0.6	0.6	1.1	1.3	1.8
2003.....	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0

All origins 3/, male

2007.....	0.3	0.3	0.4	0.5	0.7	0.9	1.3	*	2.3
2006.....	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.7	1.8
2005.....	0.4	0.4	0.3	0.5	0.7	1.0	1.4	1.5	1.9
2004.....	0.2	0.4	0.4	0.6	0.8	0.7	1.2	1.4	2.3
2003.....	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5

All origins 3/, female

2007.....	0.2	0.3	0.3	0.3	0.5	0.5	0.8	1.3	2.0
2006.....	*	0.2	0.2	0.4	*	0.7	1.1	1.0	1.8
2005.....	0.3	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.9
2004.....	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.3	0.4	0.4	0.6	0.7	1.0	1.1	2.2
2006.....	0.2	0.3	0.3	0.5	0.5	0.8	1.1	1.3	1.8
2005.....	0.3	0.3	0.3	0.5	0.6	0.8	1.1	1.3	1.9
2004.....	0.2	0.3	0.3	0.5	0.6	0.7	1.2	1.3	1.8
2003.....	0.3	0.3	0.4	0.4	0.8	0.9	1.0	1.5	1.7
2002.....	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.2	1.8
1999.....	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.4	2.1

Non-Hispanic 4/, male

2007.....	0.3	0.3	0.4	0.5	0.7	0.9	1.2	*	2.4
2006.....	0.3	0.3	0.4	0.6	0.6	0.9	1.2	1.8	1.8
2005.....	0.3	0.4	0.3	0.5	0.7	1.0	1.5	1.6	2.0
2004.....	0.2	0.4	0.4	0.6	0.8	0.7	1.3	1.3	2.4
2003.....	0.4	0.3	0.4	0.4	0.9	1.1	1.1	1.6	1.6
2002.....	0.4	0.3	0.4	0.6	0.7	1.1	1.2	1.3	2.3
2001.....	0.3	0.3	0.4	0.6	0.7	0.8	1.3	1.2	1.8
2000.....	0.3	*	0.5	*	0.6	1.2	1.4	1.5	2.1
1999.....	0.3	0.3	0.7	0.6	0.8	0.9	1.3	1.5	2.6

Non-Hispanic 4/, female

2007.....	0.3	0.3	0.3	0.3	0.5	0.5	0.8	1.3	2.1
2006.....	*	0.2	0.2	0.4	*	0.8	1.1	1.1	1.9
2005.....	0.3	0.2	0.3	0.4	0.6	0.6	0.8	1.2	1.9
2004.....	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	0.2	0.4	*	0.5	0.9	1.1	1.5	2.0
2001.....	0.3	0.3	0.3	*	*	0.9	1.1	1.4	1.9
2000.....	*	0.3	*	*	0.5	0.7	0.9	1.1	1.7
1999.....	0.2	0.3	0.4	0.6	0.7	0.8	1.1	1.4	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic White, both sexes

2007 5/.....	0.3	0.3	0.4	0.3	0.5	0.7	1.0	1.2	2.2
2006 5/.....	0.2	0.3	0.3	0.5	0.5	0.8	1.2	1.4	1.9
2005 5/.....	0.3	0.3	0.3	0.4	0.7	0.7	1.0	1.3	1.9
2004 5/.....	0.2	0.3	0.3	0.4	0.6	0.7	1.1	1.3	1.8
2003 5/.....	0.3	0.2	0.4	0.4	0.8	0.9	1.0	1.5	1.7
2002.....	0.2	0.2	0.3	0.4	0.6	1.0	1.0	1.4	2.1
2001.....	0.3	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.3	1.8
1999.....	0.3	0.3	0.5	0.6	0.7	0.8	1.2	1.4	2.0

Non-Hispanic White, male

2007 5/.....	0.3	*	0.4	0.4	0.6	0.9	1.2	*	2.4
2006 5/.....	0.3	0.3	0.4	0.6	0.6	0.9	1.3	1.9	1.9
2005 5/.....	0.3	0.4	0.3	0.4	0.7	1.0	1.3	1.6	2.0
2004 5/.....	*	0.3	0.4	0.5	0.7	0.7	1.2	1.3	2.3
2003 5/.....	0.3	0.3	0.4	0.4	1.0	1.1	1.1	1.7	1.8
2002.....	0.3	*	0.3	0.6	0.6	1.2	1.1	1.2	2.3
2001.....	0.3	0.3	0.4	0.5	*	0.9	1.3	1.2	2.0
2000.....	*	*	0.5	*	*	1.1	1.2	1.7	2.1
1999.....	0.3	*	0.7	0.7	0.7	0.9	1.3	1.5	2.6

Non-Hispanic White, female

2007 5/.....	0.3	0.3	0.3	*	0.5	*	0.8	1.4	2.0
2006 5/.....	*	*	*	*	*	0.7	1.1	1.1	1.9
2005 5/.....	*	*	*	0.4	0.6	0.6	0.8	1.1	1.8
2004 5/.....	*	*	*	*	0.5	0.6	0.9	1.3	1.6
2003 5/.....	*	*	0.4	*	0.7	0.7	0.9	1.4	1.7
2002.....	*	*	0.4	*	0.5	0.9	0.9	1.4	2.1
2001.....	0.3	*	*	*	*	0.9	1.2	1.4	1.8
2000.....	*	*	*	*	0.6	0.7	0.9	1.1	1.7
1999.....	*	*	*	0.6	0.7	0.8	1.1	1.4	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

All origins 3/, both sexes

2007.....	1.7	3.3	6.2	11.2	21.3	40.2	81.7	174.6	554.4
2006.....	2.0	3.7	6.6	12.0	21.9	43.1	88.7	190.0	614.4
2005.....	1.7	3.5	6.2	11.8	21.5	43.6	88.8	190.5	618.6
2004.....	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
2003.....	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9

All origins 3/, male

2007.....	2.2	4.0	8.1	13.4	26.1	48.8	94.2	197.7	552.8
2006.....	2.6	4.9	8.2	14.5	26.6	50.0	103.3	213.5	599.0
2005.....	2.1	4.3	7.6	14.0	24.6	50.1	102.1	213.7	603.1
2004.....	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
2003.....	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0

All origins 3/, female

2007.....	1.2	2.6	4.5	9.2	17.1	33.2	72.5	160.0	555.1
2006.....	1.5	2.6	5.1	9.7	17.9	37.6	78.0	175.2	621.6
2005.....	1.4	2.7	4.9	9.8	18.7	38.4	79.1	176.1	625.6
2004.....	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
2003.....	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cause of death	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)									
Hispanic, both sexes									
2007.....	0.9	1.8	3.1	7.8	12.7	28.5	53.0	103.3	280.0
2006.....	1.0	2.1	3.2	6.3	12.6	26.9	48.6	113.5	304.8
2005.....	*	1.7	3.6	5.7	14.0	24.3	53.1	110.3	316.6
2004.....	1.0	1.8	3.2	6.7	13.7	26.3	53.6	103.2	287.3
2003.....	1.0	1.6	4.1	7.8	16.8	29.7	55.7	111.9	330.1
2002.....	*	1.8	3.1	6.7	14.2	27.1	53.3	111.8	316.0
2001.....	*	1.7	2.9	9.1	14.3	24.4	50.4	112.2	349.5
2000.....	*	*	4.0	8.3	12.2	27.2	52.3	108.1	369.6
1999.....	*	*	3.5	7.6	12.3	28.6	53.9	109.3	377.4
Hispanic, male									
2007.....	*	2.4	3.8	9.8	15.1	36.7	55.5	107.7	288.7
2006.....	*	3.0	3.7	6.8	14.4	34.3	56.7	142.5	276.3
2005.....	*	2.3	3.4	5.5	15.1	30.2	70.2	122.6	295.1
2004.....	*	*	4.6	8.1	16.0	30.2	59.3	124.6	263.4
2003.....	*	*	5.3	11.3	18.2	36.3	68.7	134.1	312.5
2002.....	*	*	4.0	8.2	17.3	30.7	64.0	124.7	296.2
2001.....	*	*	*	9.5	19.5	24.7	47.1	129.9	333.3
2000.....	*	*	5.3	10.9	16.4	31.6	52.4	117.8	380.9
1999.....	*	*	*	8.3	15.5	36.7	67.9	113.2	358.0
Hispanic, female									
2007.....	*	*	*	5.9	10.7	22.1	51.2	100.3	275.3
2006.....	*	*	2.7	5.8	11.1	21.1	42.9	94.3	319.9
2005.....	*	*	3.7	6.0	13.0	19.7	40.7	102.2	327.8
2004.....	*	*	*	5.5	11.7	23.3	49.5	89.1	299.6
2003.....	*	*	*	4.7	15.7	24.6	46.6	97.4	339.1
2002.....	*	*	*	5.4	11.6	24.3	45.7	103.7	325.8
2001.....	*	*	*	8.7	10.0	24.1	52.8	101.3	357.4
2000.....	*	*	*	6.0	8.8	23.9	52.3	102.1	364.0
1999.....	*	*	*	6.9	9.7	22.6	44.0	106.9	386.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

Non-Hispanic 4/, both sexes

2007.....	1.8	3.4	6.5	11.5	22.0	41.1	83.7	178.8	568.6
2006.....	2.2	3.9	6.9	12.4	22.6	44.3	91.2	194.0	629.3
2005.....	1.8	3.7	6.4	12.2	22.0	45.0	90.9	194.6	632.5
2004.....	1.8	3.5	6.3	11.8	23.0	44.9	90.6	191.7	635.3
2003.....	1.7	3.4	6.6	12.4	24.1	45.5	91.7	196.1	653.2
2002.....	1.8	3.5	6.3	12.8	23.5	46.7	90.7	197.0	656.8
2001.....	1.8	3.4	6.8	13.3	24.4	46.8	94.5	202.2	685.6
2000.....	1.5	3.2	6.3	13.1	24.7	47.5	95.2	202.8	694.8
1999.....	1.6	3.0	6.7	11.9	24.1	49.2	95.8	204.5	691.0

Non-Hispanic 4/, male

2007.....	2.3	4.1	8.5	13.6	26.9	49.7	96.8	203.2	567.9
2006.....	2.8	5.0	8.6	15.1	27.4	51.0	106.2	217.3	616.3
2005.....	2.2	4.5	8.0	14.6	25.3	51.4	103.9	218.6	618.8
2004.....	2.3	4.1	7.7	14.1	27.1	52.4	105.6	215.4	621.3
2003.....	2.2	4.2	8.1	14.9	29.0	54.0	107.2	217.1	643.2
2002.....	2.2	4.6	7.5	15.3	27.7	54.4	105.9	221.4	634.7
2001.....	2.3	4.2	8.4	15.3	28.5	55.8	110.7	230.4	658.0
2000.....	2.1	3.8	7.5	15.5	28.9	56.8	110.3	233.4	683.3
1999.....	2.0	3.8	8.2	14.5	28.7	60.0	109.9	228.4	677.1

Non-Hispanic 4/, female

2007.....	1.2	2.8	4.7	9.4	17.6	34.0	74.0	163.5	568.9
2006.....	1.6	2.8	5.3	10.0	18.4	38.7	80.2	179.6	635.3
2005.....	1.4	2.9	5.0	10.1	19.2	39.7	81.5	179.7	638.7
2004.....	1.4	2.9	5.0	9.7	19.4	38.9	79.9	177.2	641.6
2003.....	1.3	2.7	5.1	10.1	19.8	38.6	80.6	183.3	657.6
2002.....	1.4	2.5	5.1	10.5	19.8	40.6	79.9	182.2	666.4
2001.....	1.3	2.7	5.2	11.5	20.8	39.6	83.2	185.4	697.0
2000.....	0.9	2.6	5.2	10.9	21.1	40.1	84.7	184.8	699.5
1999.....	1.2	2.2	5.3	9.6	20.2	40.8	86.1	190.5	696.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

Non-Hispanic White, both sexes

2007 5/.....	1.3	2.6	5.3	10.2	20.3	40.1	83.4	181.9	590.2
2006 5/.....	1.5	2.9	5.8	11.4	21.1	43.3	91.6	197.7	653.7
2005 5/.....	1.3	2.9	5.4	10.9	20.3	43.6	89.8	196.7	655.9
2004 5/.....	1.4	2.6	5.2	10.6	22.0	44.1	89.9	195.3	657.1
2003 5/.....	1.3	2.7	5.7	10.9	22.7	44.0	91.3	198.9	673.5
2002.....	1.2	2.8	5.5	11.7	22.2	45.1	89.8	198.8	677.3
2001.....	1.4	2.8	5.9	12.2	22.9	45.5	93.9	204.3	706.8
2000.....	1.1	2.7	5.5	12.0	23.3	45.8	94.5	204.9	714.3
1999.....	1.2	2.3	5.7	11.1	22.6	47.7	94.0	205.5	709.1

Non-Hispanic White, male

2007 5/.....	1.7	3.2	6.9	12.1	25.3	48.6	96.2	207.3	592.5
2006 5/.....	2.0	3.7	7.6	13.9	26.0	49.3	107.3	222.0	641.8
2005 5/.....	1.6	3.5	6.7	13.0	23.2	49.7	102.3	223.2	646.8
2004 5/.....	1.6	3.2	6.4	12.4	25.8	52.1	104.3	220.1	643.0
2003 5/.....	1.7	3.2	7.1	13.1	27.5	52.9	107.8	220.6	665.3
2002.....	1.6	3.7	6.6	14.0	26.1	52.5	105.5	224.9	656.2
2001.....	1.9	3.5	7.5	14.4	26.9	54.4	110.5	232.1	679.1
2000.....	1.7	3.2	6.9	14.2	27.8	55.5	110.1	235.8	701.6
1999.....	1.5	3.0	7.3	13.5	26.8	59.1	108.8	229.9	690.8

Non-Hispanic White, female

2007 5/.....	0.9	2.0	3.8	8.4	15.8	33.0	73.7	165.7	589.1
2006 5/.....	1.0	2.1	4.2	9.1	16.7	38.4	79.9	182.4	659.3
2005 5/.....	1.0	2.2	4.2	8.8	17.7	38.6	80.5	180.2	660.0
2004 5/.....	1.1	2.0	4.1	8.9	18.6	37.4	79.3	179.9	663.4
2003 5/.....	0.9	2.1	4.4	8.8	18.4	36.6	79.3	185.5	677.2
2002.....	0.9	1.9	4.4	9.6	18.9	39.0	78.6	182.8	686.4
2001.....	0.8	2.1	4.3	10.2	19.4	38.2	82.2	187.4	718.4
2000.....	0.6	2.1	4.2	9.9	19.3	38.1	83.6	186.6	719.4
1999.....	0.8	1.7	4.2	8.8	18.9	38.7	83.6	191.1	716.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

Non-Hispanic Black, both sexes

2007 5/.....	5.1	9.9	17.3	25.7	41.3	63.6	113.8	192.0	467.9
2006 5/.....	6.6	11.1	16.6	25.1	41.9	65.7	111.4	204.1	492.6
2005 5/.....	5.2	10.2	16.1	27.5	42.4	70.5	125.8	219.1	497.8
2004 5/.....	5.2	10.8	16.7	24.4	36.6	65.0	122.2	198.0	513.2
2003 5/.....	4.9	9.7	15.6	28.2	42.2	71.8	116.8	206.1	538.7
2002.....	5.6	9.4	14.1	25.3	39.3	72.8	119.4	217.6	534.5
2001.....	4.7	8.9	16.0	25.1	42.1	72.3	122.2	216.6	551.2
2000.....	4.3	7.9	14.4	24.8	42.0	73.4	121.4	211.8	564.8
1999.....	4.9	8.6	16.6	22.4	43.0	74.9	131.6	218.4	557.4

Non-Hispanic Black, male

2007 5/.....	6.9	11.9	23.8	33.1	49.4	79.7	136.3	216.4	442.4
2006 5/.....	8.3	15.5	18.8	32.3	50.3	83.3	128.4	229.6	473.7
2005 5/.....	6.4	13.0	20.7	33.7	50.0	84.9	154.3	227.4	431.0
2004 5/.....	7.1	12.6	21.4	31.6	46.6	69.8	149.7	224.0	510.1
2003 5/.....	5.7	12.0	20.0	36.1	52.4	80.2	127.1	228.7	521.0
2002.....	6.7	12.3	16.6	32.2	48.6	88.9	138.0	240.2	489.0
2001.....	5.2	10.7	19.3	26.8	50.5	87.1	141.7	260.8	547.5
2000.....	5.7	9.3	15.2	30.0	46.4	84.1	137.3	246.3	589.4
1999.....	6.5	11.3	19.3	27.6	52.6	84.5	146.9	241.8	609.7

Non-Hispanic Black, female

2007 5/.....	3.5	8.2	12.0	20.0	35.4	52.5	100.3	179.2	478.5
2006 5/.....	5.2	7.4	14.7	19.4	35.8	53.6	101.2	190.7	500.4
2005 5/.....	4.1	7.9	12.2	22.6	36.8	60.7	108.8	214.8	524.7
2004 5/.....	3.5	9.2	12.8	18.9	29.2	61.8	106.0	184.3	514.4
2003 5/.....	4.1	7.8	12.0	22.0	34.7	66.1	110.7	194.3	545.6
2002.....	4.7	7.0	12.1	19.9	32.4	61.9	108.2	205.8	551.8
2001.....	4.2	7.3	13.3	23.8	35.8	62.4	110.6	194.1	552.6
2000.....	3.1	6.6	13.7	20.6	38.7	66.2	111.8	194.4	555.6
1999.....	3.5	6.3	14.4	18.3	35.8	68.5	122.3	206.8	537.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes

2007.....	14.6	20.7	28.5	41.2	63.5	101.3	171.4	315.8	770.9
2006.....	15.0	20.5	28.7	42.2	62.4	99.2	167.0	299.4	724.4
2005.....	15.5	21.2	29.7	44.5	65.0	104.9	172.3	306.2	754.0
2004.....	15.5	21.4	28.8	45.0	66.3	105.9	176.1	311.3	747.7
2003.....	15.5	21.3	31.7	46.3	71.8	113.2	186.0	328.5	781.5
2002.....	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001.....	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000.....	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
1999.....	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1

All origins 3/, male

2007.....	18.5	27.1	37.1	52.4	78.5	123.7	204.1	367.3	790.5
2006.....	19.1	26.5	37.0	53.5	78.5	120.7	196.1	347.5	735.0
2005.....	19.8	27.9	38.7	55.8	81.5	128.0	203.6	349.3	767.7
2004.....	20.2	27.8	37.9	57.6	82.7	128.0	209.5	357.0	765.2
2003.....	19.7	27.5	40.8	58.6	88.8	139.6	224.8	380.7	795.9
2002.....	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001.....	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000.....	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
1999.....	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7

All origins 3/, female

2007.....	10.9	14.6	20.3	30.9	50.3	83.0	147.3	283.3	761.7
2006.....	11.1	14.8	20.9	31.9	48.3	81.7	145.7	269.3	719.5
2005.....	11.4	14.8	21.3	34.1	50.5	86.2	149.6	279.5	747.6
2004.....	11.0	15.3	20.3	33.4	52.0	88.0	152.2	283.2	739.9
2003.....	11.5	15.4	23.1	35.1	57.0	92.0	158.4	296.6	775.1
2002.....	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001.....	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000.....	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
1999.....	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Hispanic, both sexes

2007.....	7.8	10.8	16.2	21.6	37.2	59.1	103.3	175.2	378.7
2006.....	8.5	11.5	15.4	23.3	35.0	63.4	93.2	161.0	354.5
2005.....	8.1	10.6	14.9	25.6	38.4	65.6	103.4	177.4	384.2
2004.....	8.2	11.1	13.7	25.8	38.8	63.5	102.4	178.3	369.1
2003.....	8.5	10.7	18.9	29.1	41.8	67.1	117.0	180.8	418.2
2002.....	7.8	11.9	17.2	23.2	41.7	62.4	111.4	192.1	419.6
2001.....	7.5	11.2	17.8	26.6	46.5	66.3	114.7	188.7	444.6
2000.....	7.8	13.4	18.4	26.5	44.2	68.3	123.7	203.9	457.2
1999.....	7.9	13.1	19.3	31.2	47.7	69.3	114.9	219.1	447.6

Hispanic, male

2007.....	10.8	14.1	20.8	26.6	45.4	68.9	128.4	193.7	389.3
2006.....	10.1	15.0	20.3	30.9	43.2	78.2	103.0	198.2	378.9
2005.....	11.2	14.8	20.6	34.8	47.8	81.8	130.2	191.8	401.0
2004.....	11.2	14.8	16.4	34.0	46.6	77.8	120.4	208.6	374.1
2003.....	11.7	14.0	22.9	35.9	52.6	84.1	131.9	205.8	437.2
2002.....	9.8	15.4	20.9	28.7	50.5	88.6	136.3	225.3	455.1
2001.....	10.4	15.5	22.0	34.5	55.3	83.7	139.1	229.8	438.1
2000.....	10.0	18.7	25.0	33.4	52.6	89.9	140.3	237.0	505.9
1999.....	10.5	18.1	23.6	42.6	63.0	92.2	139.7	259.4	526.2

Hispanic, female

2007.....	4.7	7.6	11.8	17.0	30.3	51.4	85.0	162.9	372.9
2006.....	6.9	7.9	10.8	16.5	28.2	52.0	86.1	136.2	341.5
2005.....	4.9	6.5	9.5	17.6	30.5	53.0	84.2	167.8	375.4
2004.....	5.2	7.4	11.2	18.7	32.4	52.4	89.6	158.3	366.6
2003.....	5.3	7.5	15.2	23.2	32.8	54.1	106.5	164.4	408.6
2002.....	5.8	8.5	13.8	18.3	34.6	42.3	93.7	170.8	402.0
2001.....	4.7	7.2	13.9	19.7	39.3	53.0	97.3	163.3	447.7
2000.....	5.5	8.5	12.5	20.6	37.4	51.9	111.9	183.7	433.3
1999.....	5.2	8.4	15.4	21.3	35.3	52.1	97.5	195.1	409.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 4/, both sexes

2007.....	15.5	21.7	29.6	42.8	65.6	104.5	176.0	324.0	791.1
2006.....	15.8	21.4	29.8	43.8	64.4	101.7	171.6	306.8	742.4
2005.....	16.3	22.2	30.9	45.9	66.9	107.6	176.4	312.7	770.9
2004.....	16.3	22.4	30.0	46.4	68.3	108.7	180.2	317.6	764.1
2003.....	16.2	22.2	32.7	47.5	73.9	116.0	189.5	334.8	795.8
2002.....	16.6	22.6	33.6	48.7	75.4	119.4	198.4	343.8	807.6
2001.....	16.2	22.7	33.4	49.2	77.1	119.3	194.6	338.8	807.8
2000.....	16.2	22.7	34.0	52.0	80.1	120.9	196.2	343.5	831.4
1999.....	16.3	23.0	35.0	54.9	81.7	126.9	200.9	351.5	837.9

Non-Hispanic 4/, male

2007.....	19.5	28.5	38.6	54.4	81.0	127.7	209.1	377.8	813.2
2006.....	20.2	27.6	38.4	55.3	81.0	123.5	201.8	355.8	753.7
2005.....	20.7	29.2	40.1	57.4	83.8	131.1	207.9	357.6	786.5
2004.....	21.2	29.0	39.6	59.3	85.2	131.3	214.4	364.2	784.1
2003.....	20.5	28.6	42.1	60.0	91.3	142.9	229.5	388.3	811.6
2002.....	21.8	29.2	42.6	61.6	92.2	146.8	237.6	396.1	836.7
2001.....	20.8	28.6	41.7	62.8	93.9	146.8	236.4	396.2	835.3
2000.....	21.1	29.7	44.5	66.2	99.4	150.3	236.4	396.9	883.9
1999.....	21.1	29.5	45.8	70.3	105.8	158.2	242.1	415.5	886.8

Non-Hispanic 4/, female

2007.....	11.7	15.3	21.1	32.0	52.0	85.4	151.6	290.1	780.7
2006.....	11.6	15.5	21.7	33.2	49.8	83.9	149.4	276.3	737.2
2005.....	12.1	15.5	22.3	35.4	52.0	88.6	153.6	284.9	763.9
2004.....	11.6	16.0	21.0	34.6	53.4	90.5	155.7	289.0	755.1
2003.....	12.1	16.1	23.7	36.0	58.7	94.4	160.9	302.1	788.8
2002.....	11.5	16.2	25.1	36.9	60.9	97.4	170.8	312.1	795.0
2001.....	11.7	17.1	25.6	36.9	62.6	97.4	165.3	304.4	796.3
2000.....	11.4	16.1	24.2	39.0	63.5	97.7	168.2	312.1	810.2
1999.....	11.7	16.9	24.9	41.0	61.1	102.4	172.3	314.0	818.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic White, both sexes

2007 5/.....	13.4	18.8	26.2	39.0	60.4	99.7	174.3	325.5	815.7
2006 5/.....	13.8	18.6	26.4	40.0	59.8	97.2	169.5	307.0	765.0
2005 5/.....	14.0	18.9	27.4	41.3	62.0	103.2	173.9	315.0	792.0
2004 5/.....	13.8	19.4	26.4	42.5	63.5	104.0	176.0	318.9	779.9
2003 5/.....	13.9	18.9	29.0	42.9	68.3	110.3	187.2	335.9	811.6
2002.....	13.9	19.4	29.9	44.2	71.1	114.9	195.2	345.0	820.9
2001.....	13.4	19.6	29.1	45.3	72.0	114.9	189.5	338.8	821.8
2000.....	13.5	19.3	30.4	47.6	75.0	115.9	191.0	344.5	845.3
1999.....	13.4	19.9	30.5	49.4	76.5	121.0	195.8	350.7	851.7

Non-Hispanic White, male

2007 5/.....	17.3	24.8	34.2	49.7	74.9	121.6	207.4	378.5	841.6
2006 5/.....	17.7	24.0	34.1	50.3	75.4	118.0	198.6	358.0	774.7
2005 5/.....	18.1	25.4	35.6	52.3	77.8	125.0	205.5	361.5	808.1
2004 5/.....	18.2	25.7	35.1	54.6	79.8	126.3	209.2	364.4	795.1
2003 5/.....	17.9	24.6	37.5	55.0	84.6	136.9	226.8	390.1	827.7
2002.....	18.5	25.6	38.1	56.8	87.4	141.7	234.0	397.9	850.7
2001.....	17.6	25.4	36.4	58.0	88.9	141.1	230.1	394.4	845.6
2000.....	18.2	25.2	39.8	61.4	93.3	144.9	231.0	398.7	898.3
1999.....	17.9	25.7	40.1	63.6	99.5	151.1	236.2	413.9	897.0

Non-Hispanic White, female

2007 5/.....	9.6	12.9	18.4	28.9	47.5	81.4	149.3	291.7	803.5
2006 5/.....	9.9	13.2	19.0	30.3	45.9	79.8	147.7	274.8	760.4
2005 5/.....	10.0	12.6	19.4	31.0	47.9	85.1	150.5	286.0	784.6
2004 5/.....	9.5	13.2	18.0	31.2	48.8	85.6	151.7	290.7	773.1
2003 5/.....	10.0	13.4	20.9	31.7	53.8	88.5	158.6	302.7	804.5
2002.....	9.3	13.3	21.9	32.6	56.6	93.1	167.4	312.7	808.1
2001.....	9.2	13.9	22.1	33.5	57.0	93.6	160.7	305.2	811.9
2000.....	8.9	13.6	21.4	34.9	58.9	92.6	162.9	312.3	824.0
1999.....	9.0	14.2	21.3	36.2	56.3	97.1	167.5	313.5	833.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic Black, both sexes

2007 5/.....	32.1	47.0	62.4	85.0	125.4	171.1	234.2	370.6	656.9
2006 5/.....	31.8	45.9	63.0	86.8	115.0	164.4	231.5	369.7	614.7
2005 5/.....	34.4	49.7	65.0	93.3	120.6	169.7	244.0	346.3	649.7
2004 5/.....	35.7	48.9	66.6	90.6	123.5	177.0	260.3	358.0	678.7
2003 5/.....	35.3	50.8	69.8	95.4	135.3	194.1	249.6	371.5	711.8
2002.....	37.5	50.6	72.2	96.2	129.8	183.6	266.8	381.5	747.1
2001.....	37.7	51.0	75.4	93.3	137.0	182.1	280.5	386.8	737.9
2000.....	38.1	53.9	70.9	98.3	140.3	193.5	287.0	382.5	748.9
1999.....	39.3	52.9	78.7	111.2	142.1	208.0	285.7	405.3	752.7

Non-Hispanic Black, male

2007 5/.....	38.3	61.8	82.9	111.5	164.0	217.1	280.4	457.0	637.0
2006 5/.....	41.3	60.2	82.7	116.2	149.4	208.0	285.4	423.5	647.8
2005 5/.....	42.7	64.4	86.2	117.2	157.0	220.1	293.4	390.0	672.7
2004 5/.....	46.2	61.5	88.6	119.2	155.4	216.5	323.2	434.7	728.6
2003 5/.....	44.1	66.1	93.3	121.3	174.4	238.3	310.2	440.0	718.1
2002.....	49.1	64.1	92.7	119.1	159.5	230.2	316.4	440.7	763.6
2001.....	46.8	61.6	98.2	123.3	161.0	229.6	352.7	484.9	813.1
2000.....	47.4	72.9	96.8	124.0	177.8	235.1	344.9	454.0	821.9
1999.....	48.9	68.9	105.2	145.4	185.7	266.2	342.7	509.2	855.7

Non-Hispanic Black, female

2007 5/.....	26.6	34.5	45.7	64.2	97.0	139.4	206.4	325.2	665.1
2006 5/.....	23.5	33.8	46.7	63.9	89.6	134.4	199.3	341.4	601.3
2005 5/.....	27.1	37.2	47.6	74.8	93.6	135.3	214.6	323.3	640.4
2004 5/.....	26.7	38.3	48.6	68.3	99.9	150.3	223.0	317.6	659.0
2003 5/.....	27.7	37.8	50.4	75.3	106.2	164.3	213.6	335.7	709.4
2002.....	27.5	39.3	55.3	78.3	107.8	152.3	237.2	350.6	740.8
2001.....	29.7	42.1	56.7	69.7	119.3	150.5	237.4	336.9	710.1
2000.....	30.0	37.6	49.6	78.0	112.7	165.8	252.1	346.5	721.7
1999.....	31.1	39.3	57.1	84.3	109.9	169.3	251.3	353.9	714.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All origins 3/, both sexes

2007.....	1.9	3.8	5.2	8.1	12.8	20.4	35.1	67.9	191.1
2006.....	2.1	4.0	5.5	8.8	13.3	21.0	36.6	69.7	189.4
2005.....	2.2	3.3	5.0	8.3	13.9	22.2	40.5	75.4	210.0
2004.....	2.2	3.4	4.9	8.1	13.0	21.9	38.5	71.3	198.5
2003.....	1.8	3.3	5.0	8.0	13.1	21.3	38.4	69.9	188.9
2002.....	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001.....	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000.....	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
1999.....	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1

All origins 3/, male

2007.....	2.4	4.8	6.5	10.0	14.6	22.4	38.4	66.8	157.9
2006.....	2.6	5.1	6.7	10.6	15.1	22.8	36.6	72.2	159.2
2005.....	2.8	4.4	6.4	9.9	15.9	23.3	41.9	74.5	169.3
2004.....	2.6	4.5	5.6	9.8	14.9	22.2	40.0	72.7	166.0
2003.....	2.5	4.2	6.0	9.2	14.9	22.5	37.5	66.1	153.5
2002.....	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001.....	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000.....	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
1999.....	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4

All origins 3/, female

2007.....	1.4	2.9	4.0	6.3	11.2	18.8	32.7	68.6	207.0
2006.....	1.6	2.8	4.4	7.1	11.6	19.6	36.7	68.1	203.5
2005.....	1.7	2.3	3.8	6.8	12.2	21.4	39.4	76.0	228.7
2004.....	1.7	2.3	4.3	6.6	11.3	21.6	37.4	70.4	213.2
2003.....	1.2	2.4	4.0	6.9	11.4	20.4	39.1	72.2	204.6
2002.....	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001.....	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000.....	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
1999.....	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Hispanic, both sexes

2007.....	1.6	3.4	5.0	8.2	14.6	19.5	33.4	69.8	157.0
2006.....	1.6	3.4	6.1	8.6	13.0	22.0	27.7	58.9	147.1
2005.....	1.3	2.7	4.6	8.8	14.6	23.8	39.1	74.4	169.4
2004.....	1.4	3.3	4.3	7.1	12.3	21.1	34.9	58.9	172.2
2003.....	1.7	2.3	4.3	7.9	12.6	21.9	34.3	61.1	171.0
2002.....	*	1.4	4.4	8.7	10.0	22.3	34.7	68.2	148.6
2001.....	1.5	2.2	2.9	6.6	11.4	22.8	38.2	67.5	160.1
2000.....	*	2.7	3.5	10.0	10.8	20.3	36.1	63.5	148.0
1999.....	1.7	2.7	3.2	7.3	12.3	19.0	35.5	71.7	129.8

Hispanic, male

2007.....	1.7	4.1	5.8	10.8	17.0	24.6	31.7	74.4	151.4
2006.....	1.7	4.2	6.6	11.1	16.1	22.9	30.4	52.4	131.9
2005.....	1.6	2.8	4.3	8.7	18.3	23.4	38.3	79.7	140.7
2004.....	*	4.6	5.9	7.2	16.0	21.5	38.0	66.1	147.1
2003.....	2.4	3.3	4.5	11.6	11.0	21.4	33.4	63.4	133.9
2002.....	*	*	5.5	9.2	8.8	23.7	39.0	59.5	132.2
2001.....	2.2	*	*	5.7	10.6	24.7	39.6	72.7	129.5
2000.....	*	4.2	4.4	13.0	9.7	22.4	42.1	73.6	141.1
1999.....	*	*	*	8.6	16.2	21.4	43.7	67.6	131.6

Hispanic, female

2007.....	*	2.6	4.2	5.9	12.5	15.5	34.6	66.7	160.0
2006.....	*	2.5	5.7	6.3	10.4	21.4	25.7	63.2	155.2
2005.....	*	2.6	4.8	8.9	11.6	24.1	39.6	70.9	184.4
2004.....	*	*	*	6.9	9.2	20.8	32.7	54.2	185.1
2003.....	*	*	4.1	4.7	13.9	22.3	34.9	59.5	189.8
2002.....	*	*	3.4	8.3	11.1	21.3	31.5	73.7	156.7
2001.....	*	*	*	7.3	12.1	21.3	37.3	64.3	175.0
2000.....	*	*	*	7.4	11.8	18.8	31.9	57.3	151.3
1999.....	*	*	*	6.2	9.1	17.2	29.7	74.1	128.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic 4/, both sexes

2007.....	1.9	3.8	5.2	8.1	12.6	20.4	35.2	67.6	192.8
2006.....	2.2	4.0	5.4	8.8	13.3	20.9	37.2	70.2	191.2
2005.....	2.4	3.4	5.0	8.2	13.8	22.1	40.5	75.3	211.8
2004.....	2.3	3.4	5.0	8.2	13.0	21.8	38.7	71.8	199.4
2003.....	1.8	3.4	5.0	8.0	13.0	21.1	38.6	70.1	189.4
2002.....	1.9	3.0	4.5	7.2	11.8	20.5	34.9	66.5	181.1
2001.....	1.9	3.0	4.5	7.5	12.3	18.7	35.2	65.0	172.0
2000.....	1.8	3.0	4.5	7.6	11.5	18.9	33.5	62.9	162.9
1999.....	1.7	2.8	4.2	7.3	11.9	18.6	32.5	59.1	152.3

Non-Hispanic 4/, male

2007.....	2.5	4.8	6.6	10.0	14.5	22.2	38.8	66.2	158.2
2006.....	2.8	5.2	6.7	10.6	15.1	22.7	36.9	73.2	160.5
2005.....	2.9	4.5	6.5	9.9	15.6	23.3	42.1	74.1	170.7
2004.....	2.8	4.4	5.5	10.0	14.7	22.1	40.1	73.0	166.7
2003.....	2.5	4.3	6.1	9.0	15.1	22.4	37.7	66.2	154.4
2002.....	2.4	3.7	5.4	8.1	12.8	22.0	35.6	64.8	153.9
2001.....	2.3	3.8	5.3	8.3	13.8	19.6	35.6	60.6	151.1
2000.....	2.3	3.8	5.4	8.7	11.5	20.4	33.9	62.4	142.0
1999.....	2.0	3.5	5.1	8.1	13.2	20.0	32.6	58.5	131.9

Non-Hispanic 4/, female

2007.....	1.4	2.9	4.0	6.3	11.0	19.0	32.6	68.6	209.2
2006.....	1.7	2.9	4.2	7.1	11.7	19.5	37.4	68.3	205.5
2005.....	1.8	2.3	3.7	6.7	12.3	21.1	39.3	76.1	230.6
2004.....	1.8	2.4	4.5	6.5	11.4	21.5	37.6	71.1	214.0
2003.....	1.2	2.5	4.0	7.1	11.2	20.1	39.2	72.6	204.7
2002.....	1.4	2.3	3.7	6.3	10.9	19.3	34.3	67.5	192.8
2001.....	1.4	2.3	3.8	6.8	10.9	18.0	35.0	67.6	180.7
2000.....	1.3	2.2	3.5	6.6	11.5	17.8	33.3	63.2	171.3
1999.....	1.3	2.1	3.2	6.5	10.9	17.4	32.5	59.5	160.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic White, both sexes

2007 5/.....	1.1	2.4	3.5	5.8	9.6	16.8	30.2	61.4	184.1
2006 5/.....	1.3	2.5	3.7	6.4	9.8	16.9	31.8	63.6	180.8
2005 5/.....	1.4	2.0	3.4	5.9	10.2	17.1	34.5	68.9	201.5
2004 5/.....	1.3	2.1	3.1	5.6	9.6	17.2	32.7	65.0	188.3
2003 5/.....	1.1	2.2	3.1	5.6	9.5	16.1	32.9	64.1	181.0
2002.....	1.1	2.0	2.9	5.1	8.9	16.3	29.3	59.7	170.6
2001.....	1.0	1.8	2.9	5.2	8.9	14.7	30.3	58.1	162.8
2000.....	1.0	1.7	2.9	5.0	8.4	14.9	28.3	57.0	152.7
1999.....	1.0	1.5	2.7	4.8	8.8	14.9	27.1	52.2	141.7

Non-Hispanic White, male

2007 5/.....	1.5	3.2	4.9	7.2	11.0	18.1	33.7	60.1	149.9
2006 5/.....	1.8	3.3	4.5	8.3	11.3	17.8	31.9	65.7	152.0
2005 5/.....	1.9	2.8	4.5	7.3	11.5	17.6	35.0	68.0	160.8
2004 5/.....	1.7	2.8	3.6	7.2	11.2	17.8	33.4	66.4	159.6
2003 5/.....	1.5	2.8	4.0	6.1	11.6	17.6	32.7	60.0	145.4
2002.....	1.4	2.6	3.6	6.1	9.8	17.7	30.2	57.3	145.4
2001.....	1.3	2.2	3.5	5.9	10.2	15.6	30.8	53.4	142.1
2000.....	1.3	2.3	3.7	5.9	8.7	16.3	29.1	56.5	131.8
1999.....	1.3	1.9	3.6	5.6	10.0	16.2	27.5	51.2	119.8

Non-Hispanic White, female

2007 5/.....	0.7	1.8	2.2	4.5	8.3	15.7	27.6	62.2	200.4
2006 5/.....	0.9	1.6	2.8	4.7	8.5	16.2	31.7	62.2	194.2
2005 5/.....	0.9	1.3	2.3	4.6	8.9	16.6	34.2	69.5	220.1
2004 5/.....	0.9	1.4	2.6	4.2	8.2	16.7	32.1	64.2	201.2
2003 5/.....	0.7	1.7	2.3	5.0	7.6	14.9	33.0	66.6	196.6
2002.....	0.7	1.4	2.2	4.2	8.0	15.1	28.7	61.1	181.4
2001.....	0.7	1.3	2.3	4.5	7.8	14.0	29.9	60.9	171.5
2000.....	0.6	1.1	2.2	4.2	8.0	13.8	27.8	57.2	161.1
1999.....	0.7	1.1	1.8	4.1	7.8	13.7	26.9	52.8	150.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Non-Hispanic Black, both sexes

2007 5/.....	7.3	13.9	18.8	28.6	39.8	53.8	88.9	135.2	301.7
2006 5/.....	7.9	15.3	19.5	29.2	43.6	58.4	95.7	151.5	322.8
2005 5/.....	8.8	13.1	18.7	29.2	45.3	67.7	102.8	155.5	345.2
2004 5/.....	8.6	12.6	20.0	30.2	42.4	66.0	105.9	152.7	339.6
2003 5/.....	7.2	12.3	21.1	29.6	43.7	69.6	103.4	147.2	304.3
2002.....	7.6	11.1	18.8	24.8	36.9	62.6	98.1	153.2	312.4
2001.....	7.9	12.9	18.3	27.6	41.2	57.4	90.7	150.1	291.5
2000.....	7.3	13.1	18.3	29.5	41.0	59.9	93.8	132.8	285.9
1999.....	6.5	13.2	17.3	27.3	38.7	58.3	96.0	140.3	272.2

Non-Hispanic Black, male

2007 5/.....	9.0	17.5	21.7	36.5	48.7	62.9	103.9	140.8	251.3
2006 5/.....	9.5	19.9	25.4	33.8	50.5	73.8	99.4	174.2	274.9
2005 5/.....	10.2	17.7	24.9	37.1	53.4	80.3	127.2	157.6	294.0
2004 5/.....	10.1	17.0	22.9	37.2	50.1	67.5	125.4	160.8	268.3
2003 5/.....	9.9	16.8	25.0	36.9	50.6	75.5	103.3	149.4	280.2
2002.....	9.5	13.4	22.2	26.3	40.7	68.4	107.0	166.9	276.8
2001.....	10.2	16.7	21.9	30.9	49.5	63.4	95.8	161.1	277.2
2000.....	9.0	15.9	22.9	35.3	40.8	65.3	93.9	139.9	268.9
1999.....	8.1	17.1	19.9	30.9	43.8	63.4	98.6	156.8	268.6

Non-Hispanic Black, female

2007 5/.....	5.9	10.9	16.4	22.5	33.3	47.6	79.8	132.2	322.6
2006 5/.....	6.5	11.4	14.6	25.7	38.5	47.8	93.5	139.6	342.3
2005 5/.....	7.6	9.2	13.7	23.0	39.4	59.1	88.3	154.4	365.8
2004 5/.....	7.3	8.9	17.7	24.9	36.6	64.9	94.4	148.4	367.8
2003 5/.....	4.8	8.5	17.8	24.0	38.5	65.7	103.5	146.1	313.6
2002.....	6.0	9.1	16.0	23.7	34.2	58.7	92.8	146.1	326.1
2001.....	5.9	9.6	15.4	25.0	35.0	53.4	87.7	144.5	296.9
2000.....	5.9	10.6	14.5	25.0	41.1	56.3	93.8	129.2	292.2
1999.....	5.1	9.8	15.2	24.4	34.8	54.9	94.4	132.1	273.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes

2007.....	11.4	17.9	26.0	39.8	69.4	122.6	231.7	438.8	1,015.5
2006.....	11.2	18.5	26.4	42.7	70.7	127.5	240.3	458.6	1,039.6
2005.....	11.7	18.8	25.8	42.7	73.8	133.6	253.6	497.3	1,141.8
2004.....	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
2003.....	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8

All origins 3/, male

2007.....	12.5	20.4	31.2	46.7	79.3	138.9	253.2	445.8	895.7
2006.....	12.0	21.2	31.2	49.2	81.9	140.9	259.6	468.5	932.4
2005.....	12.1	21.5	30.1	50.0	84.6	149.5	275.8	513.0	1,023.3
2004.....	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
2003.....	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8

All origins 3/, female

2007.....	10.4	15.6	21.2	33.4	60.7	109.3	215.9	434.4	1,072.4
2006.....	10.4	15.8	21.9	36.8	60.9	116.5	226.2	452.3	1,089.8
2005.....	11.3	16.2	21.8	36.0	64.3	120.7	237.6	487.5	1,196.1
2004.....	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
2003.....	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Hispanic, both sexes

2007	11.2	17.2	28.8	40.9	65.0	104.0	193.8	339.0	652.9
2006	11.5	18.2	29.2	41.4	67.2	113.4	196.5	361.2	687.0
2005	12.5	18.2	26.4	43.5	64.5	121.5	205.5	383.7	738.4
2004	11.4	18.9	28.8	43.5	73.2	127.7	229.9	392.2	813.0
2003	11.6	19.3	29.1	43.9	77.5	124.8	252.4	421.4	880.5
2002	13.0	18.4	28.2	46.2	70.3	143.8	240.2	453.2	898.0
2001	12.4	20.3	28.6	44.5	74.7	145.5	271.5	463.4	1,042.9
2000	13.8	17.0	28.2	54.6	83.1	151.1	262.9	491.8	1,082.9
1999	13.6	18.3	31.4	45.6	90.5	150.2	265.7	498.7	1,077.6

Hispanic, male

2007	13.7	20.2	37.9	50.2	75.3	120.6	213.7	341.7	594.6
2006	12.8	22.6	34.6	51.0	76.2	132.4	244.6	369.3	581.9
2005	14.3	22.5	30.9	54.2	75.0	148.2	228.5	400.1	692.4
2004	13.7	22.6	34.9	54.7	93.8	142.0	251.7	440.7	778.9
2003	14.2	23.3	34.7	56.1	91.0	143.9	275.9	441.8	837.4
2002	16.2	22.0	36.0	57.7	86.7	173.8	269.4	462.1	856.7
2001	14.6	24.1	34.2	56.2	95.3	168.4	311.0	529.8	1,005.6
2000	16.2	20.5	37.7	63.3	99.6	183.8	310.8	556.5	1,029.9
1999	16.2	23.8	36.3	58.7	105.5	187.4	330.0	526.6	1,136.6

Hispanic, female

2007	8.5	14.1	20.2	32.6	56.3	91.0	179.4	337.2	684.5
2006	10.2	13.8	24.1	33.0	59.7	98.6	161.7	355.9	742.9
2005	10.6	13.9	22.2	34.0	55.9	100.8	188.9	372.8	762.5
2004	9.0	15.3	23.2	33.8	56.2	116.6	214.4	360.1	830.4
2003	8.9	15.4	23.8	33.2	66.4	110.1	235.8	408.1	902.3
2002	9.7	14.9	21.1	36.2	57.0	120.7	219.4	447.5	918.5
2001	10.2	16.7	23.5	34.3	58.0	128.1	243.5	422.2	1,061.2
2000	11.4	13.8	19.6	47.1	69.8	126.3	229.0	452.4	1,108.9
1999	11.0	13.1	26.9	34.4	78.4	122.2	220.2	482.0	1,049.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Non-Hispanic 4/, both sexes

2007.....	11.5	18.0	25.7	39.6	69.6	123.8	234.0	444.4	1,034.0
2006.....	11.1	18.4	26.1	42.7	70.7	128.3	242.9	463.3	1,056.5
2005.....	11.6	18.8	25.7	42.5	74.3	134.2	256.2	502.6	1,159.8
2004.....	11.9	18.1	26.2	44.4	78.6	142.9	279.0	538.4	1,263.8
2003.....	11.8	18.5	27.6	45.9	79.7	152.0	296.0	577.2	1,388.4
2002.....	12.2	18.3	28.6	48.3	84.6	161.4	311.7	606.9	1,465.0
2001.....	12.0	18.2	28.8	49.8	86.3	165.1	321.7	627.7	1,514.0
2000.....	12.9	19.4	30.9	53.5	91.3	169.6	335.1	658.3	1,604.5
1999.....	12.3	18.3	31.6	51.6	90.9	174.2	343.0	668.9	1,630.4

Non-Hispanic 4/, male

2007.....	12.3	20.3	30.4	46.3	79.4	139.9	255.4	451.8	912.5
2006.....	11.9	21.0	30.8	48.9	82.1	141.3	260.0	473.7	950.4
2005.....	11.8	21.3	30.0	49.6	85.2	149.3	278.2	518.4	1,039.8
2004.....	12.6	20.9	29.4	51.8	89.7	159.7	303.6	556.6	1,133.7
2003.....	12.7	20.8	30.9	53.0	91.2	172.1	324.6	602.2	1,252.9
2002.....	13.0	20.2	32.3	55.4	97.5	178.7	337.3	621.1	1,336.0
2001.....	13.1	20.2	32.5	56.9	98.5	188.7	350.0	658.1	1,392.9
2000.....	13.9	21.3	34.9	62.0	102.6	192.7	369.1	698.8	1,499.9
1999.....	12.9	20.4	35.5	59.7	102.3	197.6	377.8	712.2	1,524.8

Non-Hispanic 4/, female

2007.....	10.7	15.7	21.2	33.4	61.0	110.6	218.3	439.7	1,091.4
2006.....	10.4	16.0	21.7	36.9	60.7	117.8	230.3	456.8	1,105.8
2005.....	11.4	16.4	21.7	36.1	64.8	121.9	240.3	492.9	1,214.5
2004.....	11.1	15.5	23.2	37.7	68.9	129.4	261.4	527.2	1,321.9
2003.....	10.9	16.3	24.5	39.4	69.6	135.8	275.7	562.0	1,448.0
2002.....	11.3	16.4	25.1	41.7	73.4	147.5	293.5	598.3	1,520.7
2001.....	10.9	16.4	25.2	43.2	75.8	146.3	302.0	609.6	1,564.4
2000.....	12.0	17.6	27.1	45.7	81.7	151.4	311.4	634.5	1,646.8
1999.....	11.7	16.3	28.0	44.2	81.1	155.9	318.9	643.6	1,672.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Non-Hispanic White, both sexes

2007 5/.....	8.6	13.6	19.9	33.4	60.8	113.2	224.4	438.4	1,045.4
2006 5/.....	8.5	14.0	20.4	35.5	61.6	117.7	233.4	457.3	1,068.3
2005 5/.....	8.5	13.9	20.5	35.4	65.5	123.6	246.1	496.2	1,171.8
2004 5/.....	8.7	13.4	20.9	36.8	69.3	132.3	266.4	532.1	1,276.4
2003 5/.....	8.6	13.4	21.8	37.6	69.9	140.7	284.2	571.5	1,401.1
2002.....	9.0	13.4	23.3	40.4	74.2	149.9	300.9	600.4	1,479.2
2001.....	8.8	13.0	23.4	41.1	74.9	153.2	310.4	621.9	1,525.0
2000.....	9.2	15.1	25.0	45.6	81.4	158.8	323.1	651.1	1,616.8
1999.....	8.8	14.0	25.6	43.0	81.8	161.2	331.1	662.9	1,648.0

Non-Hispanic White, male

2007 5/.....	9.2	15.6	23.1	38.6	68.4	126.6	243.0	441.6	922.4
2006 5/.....	9.0	15.4	23.3	39.8	71.0	128.2	249.2	469.2	960.2
2005 5/.....	8.8	15.7	23.4	40.5	74.1	136.7	268.1	510.2	1,050.5
2004 5/.....	9.1	15.4	22.8	42.0	79.2	146.7	289.6	551.4	1,145.3
2003 5/.....	9.4	15.0	23.7	43.3	80.7	158.7	311.1	596.6	1,264.2
2002.....	9.8	14.6	25.9	45.8	84.7	166.1	325.1	615.2	1,345.9
2001.....	9.6	14.4	26.0	46.9	85.6	174.7	337.7	651.5	1,399.0
2000.....	10.0	16.4	27.5	52.8	90.8	179.5	355.4	689.2	1,505.9
1999.....	9.0	15.3	28.7	49.6	91.7	182.4	363.1	705.6	1,535.1

Non-Hispanic White, female

2007 5/.....	7.9	11.6	16.8	28.4	54.0	102.1	210.4	436.3	1,103.7
2006 5/.....	7.9	12.5	17.5	31.5	53.2	109.0	221.5	449.8	1,118.7
2005 5/.....	8.3	12.2	17.6	30.6	57.8	112.7	229.8	487.5	1,227.3
2004 5/.....	8.3	11.4	19.0	31.9	60.6	120.4	249.4	520.1	1,335.1
2003 5/.....	7.7	11.9	20.1	32.4	60.4	126.0	264.8	556.1	1,461.3
2002.....	8.1	12.2	20.7	35.3	64.9	136.6	283.5	591.4	1,536.7
2001.....	8.0	11.6	20.8	35.7	65.4	135.7	291.1	604.0	1,577.4
2000.....	8.3	13.8	22.5	38.8	73.2	142.2	300.4	628.4	1,661.6
1999.....	8.6	12.7	22.6	37.0	73.2	144.3	308.7	637.7	1,693.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Non-Hispanic Black, both sexes

2007 5/.....	28.8	47.8	67.2	95.6	149.1	220.2	351.8	547.1	1,012.8
2006 5/.....	28.0	49.7	69.2	103.7	145.9	225.8	355.9	564.9	1,023.0
2005 5/.....	31.3	51.6	67.3	101.9	147.1	235.2	377.1	618.6	1,122.3
2004 5/.....	32.4	49.1	68.1	110.5	158.1	251.6	425.2	656.2	1,226.1
2003 5/.....	32.2	53.9	73.9	114.5	159.8	261.6	440.4	686.0	1,366.0
2002.....	32.7	52.4	72.4	112.1	172.6	276.8	452.2	721.0	1,398.0
2001.....	33.1	56.2	71.9	119.3	181.5	281.4	465.4	725.7	1,465.5
2000.....	38.0	52.4	78.6	115.6	178.0	284.5	479.0	775.9	1,545.8
1999.....	35.1	51.3	79.6	119.8	172.9	312.5	493.4	762.8	1,501.5

Non-Hispanic Black, male

2007 5/.....	31.4	54.6	88.9	121.9	190.7	275.6	417.6	629.0	883.0
2006 5/.....	29.8	62.5	91.0	135.6	182.7	275.1	407.0	589.9	894.3
2005 5/.....	32.8	62.0	86.2	136.8	188.1	282.2	412.8	663.4	998.2
2004 5/.....	36.4	57.8	87.1	142.9	192.0	304.2	478.5	685.1	1,077.1
2003 5/.....	35.0	64.3	92.5	143.7	187.0	314.8	511.2	737.3	1,200.7
2002.....	34.8	63.0	90.8	141.7	217.1	334.6	508.9	738.6	1,266.8
2001.....	37.3	65.4	89.2	145.9	222.5	340.9	530.4	776.0	1,359.4
2000.....	41.5	61.7	99.4	138.8	216.4	343.4	548.2	872.8	1,479.5
1999.....	39.8	62.3	94.3	151.1	207.7	372.2	565.0	824.5	1,455.9

Non-Hispanic Black, female

2007 5/.....	26.6	42.1	49.4	75.0	118.5	182.1	312.3	504.0	1,066.6
2006 5/.....	26.4	38.8	51.2	78.9	118.8	192.0	325.5	551.8	1,075.5
2005 5/.....	30.1	42.9	51.8	74.7	116.8	203.1	355.9	595.0	1,172.3
2004 5/.....	28.9	41.9	52.5	85.4	133.0	215.9	393.6	641.0	1,285.0
2003 5/.....	29.7	45.2	58.6	91.7	139.7	225.8	398.4	659.1	1,430.0
2002.....	30.9	43.4	57.4	88.9	139.7	238.1	418.3	711.8	1,448.1
2001.....	29.6	48.4	57.8	98.4	151.2	241.8	426.6	700.1	1,504.8
2000.....	34.9	44.5	61.7	97.3	149.6	245.5	437.3	727.1	1,570.4
1999.....	31.1	42.0	67.7	95.2	147.1	272.9	450.2	732.3	1,518.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

All origins 3/, both sexes

2007.....	0.2	0.4	0.8	1.4	2.8	6.1	12.2	24.6	83.3
2006.....	0.2	0.4	0.9	1.7	3.2	6.1	12.6	27.2	89.9
2005.....	0.3	0.7	1.0	2.1	3.9	7.2	16.9	37.8	132.9
2004.....	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
2003.....	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0

All origins 3/, male

2007.....	0.3	0.6	1.1	2.0	3.7	7.4	13.7	27.2	76.5
2006.....	0.3	0.6	1.2	2.2	3.8	7.3	14.2	29.5	85.3
2005.....	0.4	1.0	1.2	2.7	4.8	8.7	20.0	42.0	121.7
2004.....	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
2003.....	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0

All origins 3/, female

2007.....	*	0.3	0.6	0.7	2.0	5.1	11.1	23.0	86.5
2006.....	*	0.2	0.5	1.3	2.6	5.1	11.4	25.7	92.1
2005.....	0.2	0.4	0.8	1.5	3.1	5.9	14.7	35.2	138.1
2004.....	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
2003.....	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Hispanic, both sexes

2007	*	*	*	*	*	5.0	11.5	20.2	50.4
2006	*	*	*	*	*	*	8.0	16.7	50.5
2005	*	*	*	*	*	5.9	11.5	27.2	79.6
2004	*	*	*	*	3.2	5.8	14.4	25.1	74.7
2003	*	*	*	*	3.1	6.5	11.9	30.3	86.0
2002	*	*	*	*	3.7	5.0	15.5	34.8	108.6
2001	*	*	*	*	*	6.3	17.0	33.8	130.7
2000	*	*	*	*	4.0	8.2	16.2	39.5	132.7
1999	*	*	*	*	*	8.5	19.4	41.1	129.1

Hispanic, male

2007	*	*	*	*	*	*	13.9	22.5	42.7
2006	*	*	*	*	*	*	*	18.8	45.1
2005	*	*	*	*	*	*	14.6	29.8	83.4
2004	*	*	*	*	*	*	17.9	31.1	88.3
2003	*	*	*	*	*	9.4	18.8	33.3	95.4
2002	*	*	*	*	*	*	18.6	41.2	102.1
2001	*	*	*	*	*	*	21.1	39.0	127.6
2000	*	*	*	*	*	12.6	25.1	45.6	104.8
1999	*	*	*	*	*	11.2	23.4	47.2	151.0

Hispanic, female

2007	*	*	*	*	*	*	9.7	18.6	54.6
2006	*	*	*	*	*	*	7.8	15.3	53.3
2005	*	*	*	*	*	*	9.3	25.6	77.6
2004	*	*	*	*	*	*	11.9	21.2	67.7
2003	*	*	*	*	*	*	*	28.4	81.3
2002	*	*	*	*	*	*	13.3	30.7	111.9
2001	*	*	*	*	*	*	14.0	30.6	132.2
2000	*	*	*	*	*	*	*	35.8	146.4
1999	*	*	*	*	*	*	16.5	37.5	118.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Non-Hispanic 4/, both sexes

2007.....	0.2	0.4	0.9	1.4	2.9	6.2	12.3	24.9	84.9
2006.....	0.2	0.4	0.9	1.8	3.2	6.3	12.9	27.7	91.8
2005.....	0.3	0.7	1.0	2.2	4.0	7.3	17.2	38.3	135.3
2004.....	0.3	0.5	1.1	2.4	4.2	8.7	17.6	38.6	139.7
2003.....	0.4	0.6	1.1	2.5	4.6	9.2	20.2	43.9	157.1
2002.....	0.3	0.7	1.4	2.5	5.3	9.5	20.7	48.3	170.2
2001.....	0.3	0.6	1.4	2.6	5.1	10.0	22.0	49.6	180.5
2000.....	0.4	0.6	1.3	2.7	5.2	10.6	23.0	52.8	190.3
1999.....	0.3	0.8	1.3	3.0	5.2	11.3	24.3	53.8	203.9

Non-Hispanic 4/, male

2007.....	0.3	0.6	1.2	2.2	3.9	7.4	13.7	27.5	78.4
2006.....	0.4	0.6	1.2	2.3	3.9	7.5	14.5	30.0	87.5
2005.....	0.4	1.0	1.2	2.8	4.9	8.8	20.3	42.6	123.7
2004.....	0.4	0.7	1.2	3.1	5.0	10.5	20.3	44.1	130.1
2003.....	0.6	0.9	1.5	3.3	5.5	10.6	24.1	48.8	140.3
2002.....	0.4	0.9	1.8	3.4	6.5	11.8	24.6	52.9	158.7
2001.....	0.5	0.8	1.7	3.4	6.4	12.4	25.0	53.6	168.2
2000.....	0.5	0.8	1.8	3.7	6.7	13.1	27.5	60.2	178.7
1999.....	0.4	1.0	1.8	3.8	6.4	13.4	28.2	60.2	189.9

Non-Hispanic 4/, female

2007.....	*	0.3	0.6	0.8	2.1	5.2	11.2	23.2	88.0
2006.....	*	0.3	0.6	1.4	2.7	5.4	11.6	26.3	93.9
2005.....	0.2	0.3	0.8	1.6	3.3	6.0	15.0	35.7	140.6
2004.....	*	0.3	0.9	1.7	3.4	7.2	15.7	35.2	144.0
2003.....	*	0.4	0.7	1.8	3.7	8.0	17.5	40.9	164.5
2002.....	*	0.5	1.1	1.7	4.2	7.7	17.9	45.4	175.1
2001.....	0.2	0.5	1.1	1.9	4.0	8.2	19.9	47.3	185.5
2000.....	0.2	0.4	0.9	1.7	4.0	8.6	19.8	48.5	195.1
1999.....	*	0.5	0.9	2.2	4.1	9.7	21.5	50.1	209.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.4	0.9	1.3	2.9	6.1	12.4	25.5	87.9
2006 5/.....	0.2	0.4	0.8	1.7	3.2	6.3	13.0	28.0	95.2
2005 5/.....	0.3	0.6	1.0	2.1	4.0	7.1	17.5	39.4	141.3
2004 5/.....	0.3	0.5	1.0	2.4	4.1	8.7	18.0	40.0	145.0
2003 5/.....	0.4	0.6	1.0	2.5	4.4	9.1	20.6	44.5	162.2
2002.....	0.3	0.6	1.4	2.3	5.4	9.4	20.5	48.8	177.0
2001.....	0.3	0.6	1.4	2.4	5.1	10.0	22.4	50.8	187.4
2000.....	0.3	0.5	1.3	2.5	5.3	10.5	23.3	53.4	196.8
1999.....	0.3	0.7	1.2	2.9	5.1	11.4	24.3	54.9	209.9

Non-Hispanic White, male

2007 5/.....	0.2	0.6	1.2	1.9	3.9	7.3	14.0	28.4	81.3
2006 5/.....	0.3	0.6	1.1	2.2	3.7	7.3	14.6	29.8	90.3
2005 5/.....	0.4	0.9	1.2	2.7	4.9	8.5	21.0	43.5	130.6
2004 5/.....	0.5	0.7	1.2	3.1	5.0	10.7	20.6	45.5	136.8
2003 5/.....	0.5	0.8	1.4	3.2	5.4	10.3	24.0	49.0	145.2
2002.....	0.4	0.7	1.7	3.1	6.7	11.6	24.2	53.5	165.7
2001.....	0.4	0.7	1.7	3.2	6.4	12.4	25.1	54.4	175.2
2000.....	0.4	0.7	1.8	3.6	6.6	13.3	28.3	60.8	183.6
1999.....	0.4	1.0	1.6	3.7	6.3	13.3	27.9	61.0	194.6

Non-Hispanic White, female

2007 5/.....	*	0.3	0.6	0.8	2.1	5.1	11.2	23.6	91.0
2006 5/.....	*	*	0.6	1.2	2.7	5.4	11.8	26.8	97.6
2005 5/.....	*	*	0.8	1.5	3.2	6.0	15.0	36.8	146.2
2004 5/.....	*	*	0.8	1.8	3.2	7.1	16.1	36.6	148.7
2003 5/.....	*	0.4	0.7	1.8	3.6	8.1	18.1	41.7	169.6
2002.....	*	0.4	1.0	1.6	4.3	7.6	17.9	45.9	181.9
2001.....	*	0.5	1.1	1.6	4.0	8.1	20.4	48.6	192.6
2000.....	*	0.3	0.8	1.5	4.1	8.3	19.8	49.1	202.1
1999.....	*	0.4	0.9	2.1	4.0	9.8	21.9	51.4	215.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	1.1	2.5	3.3	8.4	13.4	23.6	70.8
2006 5/.....	*	*	1.4	3.2	4.6	9.0	14.6	29.8	72.7
2005 5/.....	*	1.3	1.4	3.1	5.8	9.8	16.8	34.5	99.7
2004 5/.....	*	*	1.7	3.0	5.5	9.9	18.6	32.7	103.6
2003 5/.....	*	1.3	2.0	3.7	6.6	12.3	20.9	45.6	122.9
2002.....	*	1.5	2.0	4.3	5.3	11.7	27.0	52.1	118.0
2001.....	*	1.0	2.0	5.2	6.3	12.1	23.2	42.5	126.1
2000.....	*	1.5	2.1	4.8	5.8	13.1	23.6	54.6	140.2
1999.....	*	1.3	2.0	4.8	7.0	12.2	27.6	47.0	155.8

Non-Hispanic Black, male

2007 5/.....	*	*	*	4.6	4.7	10.6	13.7	22.9	64.6
2006 5/.....	*	*	2.6	3.8	6.8	11.7	17.0	40.0	76.0
2005 5/.....	*	*	*	4.3	7.6	12.7	17.4	41.0	77.1
2004 5/.....	*	*	*	4.8	6.2	11.6	22.9	41.9	78.2
2003 5/.....	*	*	*	5.0	8.1	17.2	29.0	58.9	105.5
2002.....	*	2.4	*	6.5	6.4	15.4	35.7	60.1	109.4
2001.....	*	*	*	6.1	7.8	15.1	30.8	48.2	113.2
2000.....	*	*	*	6.2	8.8	14.0	25.0	67.8	150.3
1999.....	*	*	*	5.9	9.0	15.9	37.7	56.7	166.4

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	6.9	13.2	24.0	73.4
2006 5/.....	*	*	*	*	*	7.2	13.2	24.4	71.4
2005 5/.....	*	*	*	*	4.5	7.8	16.4	31.1	108.8
2004 5/.....	*	*	*	*	5.0	8.8	16.0	27.8	113.6
2003 5/.....	*	*	*	*	5.4	9.1	16.1	38.7	129.6
2002.....	*	*	*	*	4.6	9.2	21.9	47.9	121.2
2001.....	*	*	*	4.4	5.2	10.1	18.7	39.6	130.9
2000.....	*	*	*	3.7	*	12.5	22.7	48.0	136.5
1999.....	*	*	*	3.9	5.5	9.6	21.5	42.1	151.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

All origins 3/, both sexes

2007.....	1.9	3.2	5.1	9.4	16.0	27.3	43.3	68.5	121.4
2006.....	2.2	3.2	5.3	9.0	16.4	28.2	47.1	72.2	128.5
2005.....	2.0	3.2	5.4	10.2	18.0	31.2	50.9	77.9	135.5
2004.....	2.1	3.4	5.2	9.9	18.3	33.0	51.7	81.1	139.3
2003.....	2.4	3.5	5.7	11.3	20.5	34.2	56.6	82.7	146.0
2002.....	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001.....	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000.....	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
1999.....	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2

All origins 3/, male

2007.....	2.7	4.5	7.4	13.4	22.0	35.4	54.8	86.6	142.8
2006.....	3.1	4.5	7.6	12.4	22.7	37.0	61.9	92.5	155.0
2005.....	2.9	4.4	7.8	14.2	24.8	42.7	68.4	101.2	158.2
2004.....	3.0	4.8	7.2	13.6	24.7	44.1	67.1	105.8	166.5
2003.....	3.2	5.1	7.9	15.9	29.4	46.1	78.9	111.3	179.7
2002.....	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001.....	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000.....	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
1999.....	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5

All origins 3/, female

2007.....	1.1	1.9	3.0	5.7	10.7	20.7	34.8	57.0	111.3
2006.....	1.4	1.9	3.0	5.8	10.8	21.0	36.3	59.5	116.1
2005.....	1.1	2.0	3.2	6.6	12.1	21.9	38.2	63.4	125.0
2004.....	1.2	2.1	3.3	6.6	12.8	24.0	40.6	65.9	127.0
2003.....	1.6	2.1	3.7	7.2	12.8	24.6	40.7	65.2	131.1
2002.....	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001.....	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000.....	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
1999.....	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

Hispanic, both sexes

2007	1.3	1.3	3.0	4.9	8.7	15.8	27.3	37.8	71.5
2006	1.3	2.2	3.2	5.7	10.8	15.4	27.2	38.9	71.6
2005	1.3	2.4	2.7	6.1	9.9	17.7	34.5	52.7	74.4
2004	1.6	2.5	4.5	6.2	10.8	18.0	25.6	48.4	77.1
2003	1.4	1.8	3.8	5.7	11.3	17.6	37.6	50.0	106.2
2002	1.4	2.1	4.2	8.2	11.1	19.5	34.7	51.7	92.6
2001	1.6	2.8	3.7	6.2	11.3	16.8	35.4	45.2	95.0
2000	*	2.7	4.7	7.2	14.7	27.0	39.5	47.9	102.8
1999	*	1.8	4.3	7.6	12.6	21.2	31.3	67.0	110.8

Hispanic, male

2007	1.8	*	4.3	6.6	12.2	19.4	34.7	46.5	76.2
2006	1.6	2.7	4.7	8.0	15.0	20.2	39.1	54.9	86.8
2005	1.9	3.5	4.7	9.2	13.0	26.6	52.4	65.7	105.9
2004	2.2	4.1	5.7	9.5	13.9	29.4	33.4	62.3	89.7
2003	*	2.8	5.8	9.6	15.3	23.9	54.1	66.5	138.5
2002	*	2.8	4.8	10.3	14.9	26.3	44.8	77.8	102.1
2001	2.1	3.6	4.4	9.0	16.7	24.2	51.8	55.8	108.6
2000	*	3.6	6.1	10.4	19.0	34.5	59.8	66.3	131.0
1999	*	*	5.8	10.1	17.0	29.0	44.5	97.5	163.9

Hispanic, female

2007	*	*	*	3.4	5.7	12.9	22.0	32.1	69.0
2006	*	*	*	3.6	7.3	11.6	18.6	28.3	63.5
2005	*	*	*	*	7.4	10.8	21.7	44.1	58.0
2004	*	*	3.3	*	8.2	9.3	20.0	39.3	70.6
2003	*	*	*	*	8.0	12.8	25.9	39.2	89.8
2002	*	*	3.7	6.5	8.0	14.2	27.4	35.0	87.9
2001	*	*	*	*	6.9	11.2	23.7	38.6	88.4
2000	*	*	*	*	11.2	21.4	25.1	36.7	89.0
1999	*	*	*	5.4	9.1	15.3	22.0	48.8	85.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

Non-Hispanic 4/, both sexes

2007.....	1.9	3.3	5.3	9.8	16.5	28.2	44.4	70.2	124.0
2006.....	2.3	3.3	5.4	9.2	16.7	29.1	48.4	73.9	131.2
2005.....	2.1	3.3	5.6	10.5	18.6	32.2	51.8	79.1	138.2
2004.....	2.1	3.5	5.3	10.2	18.9	34.0	53.1	82.6	141.8
2003.....	2.5	3.7	5.9	11.7	21.2	35.1	57.5	84.1	147.5
2002.....	2.3	3.2	6.3	11.9	23.0	37.5	58.8	85.1	150.3
2001.....	2.1	3.3	6.2	12.5	23.0	37.7	61.7	89.6	160.1
2000.....	2.2	3.5	6.5	13.0	25.0	41.3	63.8	93.7	165.0
1999.....	2.2	3.4	6.9	12.8	24.1	41.1	64.8	95.0	166.8

Non-Hispanic 4/, male

2007.....	2.8	4.8	7.7	13.9	22.7	36.6	56.1	89.0	146.6
2006.....	3.3	4.7	7.8	12.7	23.1	38.1	63.3	94.6	158.5
2005.....	3.0	4.5	8.0	14.5	25.6	43.8	69.1	102.9	160.6
2004.....	3.0	4.8	7.3	13.9	25.4	45.0	69.0	108.0	170.1
2003.....	3.4	5.3	8.0	16.3	30.4	47.3	80.1	113.3	181.5
2002.....	3.3	4.5	8.9	16.8	32.8	50.5	79.9	115.1	190.7
2001.....	2.9	4.4	8.5	17.3	31.6	52.6	82.2	124.9	200.5
2000.....	2.9	5.0	9.5	18.5	35.6	58.7	89.6	128.8	217.9
1999.....	3.1	4.6	9.7	18.7	34.3	58.0	91.1	130.2	209.0

Non-Hispanic 4/, female

2007.....	1.1	2.0	3.1	6.0	11.1	21.3	35.7	58.4	113.4
2006.....	1.4	1.9	3.1	6.0	11.1	21.7	37.5	61.1	118.5
2005.....	1.2	2.1	3.4	6.9	12.5	22.8	39.2	64.4	128.0
2004.....	1.3	2.1	3.3	6.8	13.2	25.0	41.8	67.1	129.2
2003.....	1.7	2.2	3.8	7.6	13.1	25.3	41.4	66.2	132.6
2002.....	1.4	2.0	4.0	7.5	14.6	27.1	44.0	66.9	132.8
2001.....	1.2	2.2	3.9	8.1	15.6	25.9	47.3	68.4	143.3
2000.....	1.5	2.1	3.7	8.1	16.0	27.5	45.8	73.0	143.6
1999.....	1.4	2.2	4.2	7.5	15.3	27.8	46.6	74.5	149.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

Non-Hispanic White, both sexes

2007 5/.....	1.6	2.8	4.8	9.4	16.1	28.7	44.8	70.5	125.7
2006 5/.....	2.1	2.8	5.0	9.0	16.4	29.6	48.9	73.7	132.6
2005 5/.....	1.7	2.8	5.2	10.3	18.3	32.6	52.1	79.2	138.9
2004 5/.....	1.9	3.1	4.8	9.6	18.6	34.8	53.7	82.7	141.9
2003 5/.....	2.1	3.2	5.6	11.4	21.4	35.8	58.5	84.3	148.3
2002.....	1.9	2.8	5.9	11.6	23.2	38.1	59.8	86.1	150.9
2001.....	1.8	2.9	5.7	12.1	23.3	38.0	61.8	90.0	160.1
2000.....	1.9	3.0	6.1	12.7	25.2	42.1	64.0	94.6	164.8
1999.....	1.9	3.0	6.5	12.9	24.4	41.8	65.5	95.6	166.7

Non-Hispanic White, male

2007 5/.....	2.4	4.0	6.9	13.7	22.3	37.5	56.6	90.2	150.7
2006 5/.....	3.0	4.1	7.2	12.5	22.8	39.1	64.8	96.2	163.3
2005 5/.....	2.5	3.8	7.5	14.4	25.1	45.2	69.3	104.0	164.5
2004 5/.....	2.6	4.4	6.6	13.1	25.2	46.7	70.2	109.7	173.0
2003 5/.....	2.9	4.7	7.8	15.9	30.8	48.9	81.8	114.0	185.8
2002.....	2.8	3.9	8.2	16.5	33.7	51.2	82.2	117.6	194.6
2001.....	2.5	3.8	8.2	16.9	32.5	53.3	83.4	126.6	204.7
2000.....	2.7	4.4	9.0	18.4	36.6	60.3	90.5	132.0	222.7
1999.....	2.8	4.1	9.5	19.1	35.4	59.8	93.2	131.9	212.8

Non-Hispanic White, female

2007 5/.....	0.9	1.6	2.6	5.3	10.5	21.3	36.0	58.0	113.9
2006 5/.....	1.1	1.5	2.8	5.6	10.6	21.6	37.0	59.6	118.3
2005 5/.....	0.9	1.8	3.1	6.5	12.2	22.2	39.3	63.7	127.2
2004 5/.....	1.1	1.8	3.0	6.3	12.8	25.0	41.6	66.0	128.0
2003 5/.....	1.3	1.8	3.5	7.3	13.0	25.1	41.7	66.0	131.8
2002.....	1.0	1.7	3.7	7.2	13.9	27.3	43.7	66.9	132.1
2001.....	1.1	1.9	3.4	7.7	15.3	25.5	46.5	67.9	141.6
2000.....	1.1	1.7	3.3	7.5	15.2	27.5	45.3	72.3	141.5
1999.....	1.1	2.0	3.6	7.2	14.8	27.5	46.1	74.1	148.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

Non-Hispanic Black, both sexes

2007 5/.....	4.2	7.4	10.5	14.3	23.2	30.4	49.1	74.7	127.8
2006 5/.....	4.5	7.3	10.3	13.8	23.6	31.2	52.2	85.1	134.0
2005 5/.....	4.4	6.9	10.4	14.9	25.1	36.0	57.6	89.2	147.5
2004 5/.....	4.4	6.4	9.5	17.6	24.7	33.4	57.6	91.1	156.9
2003 5/.....	5.5	7.5	8.7	16.6	24.5	36.0	56.1	93.6	152.9
2002.....	5.4	6.8	11.3	16.6	26.2	38.6	59.0	84.6	157.6
2001.....	4.1	6.6	10.9	18.2	23.2	43.5	69.8	97.1	175.7
2000.....	4.6	6.9	11.3	18.6	27.8	40.1	69.9	93.1	180.0
1999.....	4.7	6.3	11.4	13.8	26.5	39.1	64.9	97.4	180.1

Non-Hispanic Black, male

2007 5/.....	5.9	11.3	15.0	18.4	30.7	37.4	65.6	81.1	128.3
2006 5/.....	6.2	10.2	15.3	18.6	33.8	37.4	59.4	83.6	119.1
2005 5/.....	6.7	9.6	15.1	19.7	34.5	40.3	79.0	105.1	132.3
2004 5/.....	6.4	8.8	13.9	24.5	34.4	38.2	68.6	100.6	161.6
2003 5/.....	7.1	10.1	11.6	23.8	33.9	39.8	73.4	120.7	141.7
2002.....	6.9	9.9	16.6	23.4	31.2	49.9	65.7	100.6	168.5
2001.....	5.7	9.6	13.2	24.0	28.1	56.5	78.7	129.2	168.6
2000.....	5.6	10.2	15.2	23.0	33.6	52.3	91.5	106.4	179.7
1999.....	6.3	8.9	14.3	18.1	32.4	49.9	75.4	118.7	184.2

Non-Hispanic Black, female

2007 5/.....	2.8	4.2	6.8	11.1	17.7	25.5	39.2	71.4	127.7
2006 5/.....	3.0	4.9	6.1	10.1	16.0	26.9	47.8	85.9	140.1
2005 5/.....	2.4	4.6	6.6	11.2	18.1	33.1	44.8	80.8	153.6
2004 5/.....	2.7	4.4	5.9	12.3	17.6	30.2	51.1	86.2	155.0
2003 5/.....	4.1	5.2	6.4	11.0	17.5	33.4	45.8	79.4	157.2
2002.....	4.2	4.2	7.0	11.3	22.4	31.0	54.9	76.3	153.4
2001.....	2.7	4.1	9.0	13.6	19.6	34.9	64.5	80.8	178.4
2000.....	3.8	4.2	8.0	15.1	23.6	32.0	56.9	86.4	180.1
1999.....	3.3	4.1	9.1	10.4	22.0	31.9	58.5	86.9	178.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

All origins 3/, both sexes

2007.....	1.3	2.1	3.3	6.0	10.2	17.7	27.2	41.3	59.5
2006.....	1.6	2.1	3.4	5.6	10.6	18.0	29.5	43.2	62.7
2005.....	1.4	2.2	3.3	6.8	11.7	19.8	31.6	46.6	64.9
2004.....	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
2003.....	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3

All origins 3/, male

2007.....	2.0	3.3	5.1	9.2	15.0	24.5	36.0	55.4	81.8
2006.....	2.4	3.3	5.3	8.4	15.2	24.8	41.1	59.0	87.6
2005.....	2.3	3.3	5.3	10.0	17.1	29.0	43.8	65.4	91.0
2004.....	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
2003.....	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3

All origins 3/, female

2007.....	0.7	1.0	1.7	3.1	6.0	12.2	20.7	32.4	48.9
2006.....	0.8	1.0	1.6	3.1	6.5	12.5	21.0	33.3	51.0
2005.....	0.6	1.1	1.5	3.8	7.0	12.4	22.7	35.0	52.9
2004.....	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
2003.....	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Hispanic, both sexes

2007.....	0.9	1.0	1.5	2.7	4.3	9.2	13.8	19.8	34.3
2006.....	0.8	1.4	1.8	3.4	5.9	8.7	14.3	19.3	31.7
2005.....	*	1.7	1.8	3.9	5.9	9.0	17.6	23.4	29.1
2004.....	1.0	1.7	3.0	3.8	6.0	10.3	13.9	24.4	30.4
2003.....	*	*	2.5	3.9	5.7	11.7	25.0	29.5	49.2
2002.....	*	*	2.7	5.3	6.4	9.0	19.2	24.5	31.0
2001.....	*	1.6	1.9	3.4	7.1	9.5	21.2	25.8	31.9
2000.....	*	2.1	2.7	4.3	7.7	15.7	25.4	30.1	41.8
1999.....	*	*	2.6	4.3	7.3	11.6	18.4	34.1	40.7

Hispanic, male

2007.....	*	*	*	4.2	7.0	11.4	18.3	29.4	45.7
2006.....	*	*	3.0	5.1	9.3	13.0	21.1	28.7	47.4
2005.....	*	2.7	3.5	6.6	8.9	13.1	30.8	30.6	43.6
2004.....	*	3.1	4.3	7.2	8.0	18.6	19.0	35.9	40.6
2003.....	*	*	4.1	6.6	8.4	17.9	37.1	40.5	80.0
2002.....	*	*	3.9	7.7	9.5	15.4	25.6	43.5	43.5
2001.....	*	*	*	5.7	12.4	16.5	35.5	33.8	53.3
2000.....	*	3.0	*	6.6	12.3	20.9	42.1	42.7	64.5
1999.....	*	*	*	7.1	11.6	18.8	24.2	53.4	79.8

Hispanic, female

2007.....	*	*	*	*	*	7.5	10.4	13.4	28.1
2006.....	*	*	*	*	*	*	9.3	13.1	23.4
2005.....	*	*	*	*	*	*	8.1	18.6	21.5
2004.....	*	*	*	*	*	*	10.2	16.8	25.2
2003.....	*	*	*	*	*	6.9	16.4	22.3	33.6
2002.....	*	*	*	*	*	*	14.6	*	24.9
2001.....	*	*	*	*	*	*	11.1	20.9	21.4
2000.....	*	*	*	*	*	11.8	13.6	22.4	30.7
1999.....	*	*	*	*	*	*	14.3	22.5	21.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Non-Hispanic 4/, both sexes

2007.....	1.4	2.3	3.5	6.3	10.6	18.4	28.1	42.5	60.8
2006.....	1.6	2.2	3.5	5.8	10.9	18.7	30.5	44.4	64.2
2005.....	1.5	2.2	3.5	7.0	12.1	20.6	32.4	47.8	66.5
2004.....	1.4	2.2	3.1	6.5	11.8	21.8	33.7	50.0	66.2
2003.....	1.8	2.5	3.8	7.6	14.1	23.4	37.1	51.6	70.7
2002.....	1.6	2.1	4.1	7.6	15.2	24.8	38.3	52.6	70.8
2001.....	1.4	2.1	3.9	8.1	15.5	24.5	39.8	56.4	79.0
2000.....	1.4	2.2	4.5	8.7	16.8	28.4	41.6	57.3	80.3
1999.....	1.5	2.2	4.5	8.4	17.0	28.1	43.2	58.9	81.7

Non-Hispanic 4/, male

2007.....	2.1	3.5	5.4	9.6	15.6	25.5	37.2	57.0	83.9
2006.....	2.5	3.4	5.5	8.6	15.6	25.7	42.3	60.7	89.8
2005.....	2.4	3.3	5.4	10.2	17.7	30.1	44.5	67.3	93.3
2004.....	2.2	3.4	4.9	9.8	17.6	31.3	45.7	72.0	97.5
2003.....	2.7	3.9	6.0	11.8	21.0	33.8	54.6	75.8	106.6
2002.....	2.5	3.3	6.5	11.7	23.8	36.1	56.4	75.7	109.1
2001.....	2.2	3.2	6.0	12.4	22.8	37.0	57.2	84.2	117.0
2000.....	2.0	3.5	6.9	13.5	25.7	42.9	62.6	86.5	129.6
1999.....	2.3	3.4	7.0	13.3	26.1	42.1	65.0	89.7	127.0

Non-Hispanic 4/, female

2007.....	0.7	1.0	1.8	3.2	6.3	12.5	21.4	33.5	49.9
2006.....	0.8	1.1	1.7	3.2	6.8	13.0	21.8	34.3	52.3
2005.....	0.6	1.2	1.6	4.0	7.3	12.8	23.6	35.8	54.2
2004.....	0.7	1.1	1.5	3.5	6.8	14.1	25.1	36.5	52.3
2003.....	0.9	1.1	1.8	3.8	8.1	15.0	24.7	36.8	55.0
2002.....	0.8	1.0	1.8	3.8	7.7	15.8	25.5	38.6	54.3
2001.....	0.7	0.9	1.9	4.1	9.2	14.6	27.7	39.8	63.2
2000.....	0.8	1.0	2.2	4.3	9.2	16.9	27.0	40.2	60.4
1999.....	0.7	1.0	2.1	3.9	9.3	17.1	28.0	40.9	63.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Non-Hispanic White, both sexes

2007 5/.....	1.1	1.9	3.2	6.2	10.6	19.3	29.2	43.7	63.1
2006 5/.....	1.5	1.8	3.3	5.8	10.9	19.3	31.5	45.6	66.0
2005 5/.....	1.2	1.8	3.3	6.9	12.2	21.5	33.4	48.9	67.9
2004 5/.....	1.2	2.0	2.8	6.3	12.2	22.9	34.9	51.0	68.4
2003 5/.....	1.5	2.2	3.7	7.6	14.7	24.4	38.7	53.1	73.2
2002.....	1.3	1.8	3.8	7.7	16.0	25.8	39.9	54.3	72.8
2001.....	1.2	1.7	3.7	8.1	16.1	25.5	40.7	58.1	81.3
2000.....	1.2	2.0	4.2	8.8	17.3	29.7	42.6	59.1	82.0
1999.....	1.2	1.9	4.3	8.8	17.7	29.2	44.7	60.7	83.7

Non-Hispanic White, male

2007 5/.....	1.7	3.0	5.0	9.7	15.7	26.8	38.4	59.1	88.2
2006 5/.....	2.3	3.0	5.2	8.7	15.7	26.7	44.2	63.0	92.8
2005 5/.....	2.0	2.8	5.2	10.2	17.8	31.9	45.9	69.3	96.7
2004 5/.....	1.8	3.0	4.5	9.4	18.1	32.9	47.5	73.9	101.4
2003 5/.....	2.2	3.5	5.9	11.8	22.0	35.6	57.1	78.1	111.2
2002.....	2.0	2.8	6.1	11.9	25.3	37.4	59.1	79.1	114.7
2001.....	1.9	2.8	5.9	12.3	23.9	38.5	58.7	87.4	122.9
2000.....	1.9	3.2	6.7	13.8	27.0	45.0	64.3	90.3	135.3
1999.....	2.0	3.0	7.0	14.1	27.4	44.2	68.3	92.7	131.8

Non-Hispanic White, female

2007 5/.....	0.5	0.8	1.5	2.8	5.9	13.0	22.2	34.0	51.2
2006 5/.....	0.6	0.7	1.4	3.0	6.6	13.1	22.1	34.6	53.4
2005 5/.....	0.5	0.9	1.5	3.7	7.2	12.9	24.2	36.2	54.8
2004 5/.....	0.5	0.9	1.2	3.5	6.9	14.6	25.7	36.8	53.6
2003 5/.....	0.7	0.9	1.5	3.7	8.2	15.1	25.4	37.8	56.5
2002.....	0.6	0.8	1.7	3.8	7.7	16.3	26.2	39.2	54.7
2001.....	0.5	0.7	1.6	4.1	9.2	14.9	27.8	40.5	64.0
2000.....	0.5	0.8	1.9	4.1	8.8	17.4	27.4	40.5	60.5
1999.....	0.5	0.9	1.8	3.8	9.2	17.3	28.1	41.8	64.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Non-Hispanic Black, both sexes

2007 5/.....	3.2	5.0	6.3	7.9	12.7	13.8	23.3	31.3	40.9
2006 5/.....	3.1	5.2	6.2	7.6	13.2	16.2	24.1	34.2	47.8
2005 5/.....	3.4	5.0	5.6	8.9	13.7	16.2	26.0	38.5	51.3
2004 5/.....	3.3	4.0	5.8	10.1	11.8	15.2	26.4	41.3	44.6
2003 5/.....	4.0	4.8	5.9	9.2	12.0	17.8	26.3	36.6	47.3
2002.....	3.9	4.8	6.8	8.3	11.7	18.8	25.1	37.9	50.5
2001.....	2.8	4.5	6.2	9.5	11.6	18.6	34.5	43.3	55.9
2000.....	2.9	3.7	7.2	9.5	14.6	20.5	34.1	41.1	63.0
1999.....	3.6	4.1	6.4	6.7	14.2	20.0	29.8	40.8	61.1

Non-Hispanic Black, male

2007 5/.....	4.7	8.2	9.7	10.2	16.5	17.8	32.8	30.5	42.8
2006 5/.....	4.6	7.5	9.4	10.4	18.6	20.3	27.2	35.1	52.2
2005 5/.....	5.4	7.2	8.7	12.2	19.4	17.2	34.4	46.8	52.3
2004 5/.....	5.3	5.9	8.9	16.0	18.5	18.6	29.8	49.2	57.9
2003 5/.....	6.0	7.2	8.2	14.9	16.5	19.9	33.2	50.6	52.2
2002.....	5.8	7.7	11.2	13.2	14.8	25.3	31.0	41.3	49.2
2001.....	4.1	7.0	8.5	14.8	14.9	26.5	42.7	55.6	49.7
2000.....	3.6	6.1	10.3	13.2	18.1	28.4	47.7	46.3	70.5
1999.....	5.5	6.6	9.1	9.6	19.4	25.6	30.0	54.9	77.3

Non-Hispanic Black, female

2007 5/.....	1.8	2.4	3.6	6.0	9.9	11.2	17.5	31.7	40.1
2006 5/.....	1.9	3.1	3.5	5.4	9.3	13.5	22.3	33.7	46.0
2005 5/.....	1.6	3.2	3.0	6.4	9.4	15.6	21.0	34.2	50.9
2004 5/.....	1.6	2.5	3.4	5.5	6.8	12.8	24.3	37.1	39.4
2003 5/.....	2.2	2.8	3.9	4.8	8.6	16.4	22.2	29.2	45.4
2002.....	2.2	2.3	3.3	4.5	9.3	14.5	21.6	36.1	51.0
2001.....	1.7	2.3	4.4	5.3	9.2	13.3	29.7	37.1	58.2
2000.....	2.4	*	4.7	6.5	12.0	15.2	25.8	38.5	60.2
1999.....	2.0	*	4.3	4.4	10.4	16.3	29.7	33.8	55.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes

2007.....	0.5	1.0	1.8	3.4	5.8	9.6	16.1	27.2	62.0
2006.....	0.7	1.0	1.9	3.3	5.8	10.2	17.6	29.1	65.8
2005.....	0.5	1.0	2.1	3.5	6.3	11.4	19.3	31.3	70.6
2004.....	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
2003.....	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9

All origins 3/, male

2007.....	0.7	1.2	2.3	4.1	7.0	10.9	18.8	31.2	61.0
2006.....	0.7	1.2	2.3	4.0	7.5	12.1	20.8	33.5	67.3
2005.....	0.6	1.1	2.5	4.2	7.7	13.7	24.6	35.8	67.2
2004.....	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
2003.....	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2

All origins 3/, female

2007.....	0.4	0.9	1.3	2.7	4.7	8.5	14.1	24.6	62.4
2006.....	0.6	0.9	1.5	2.7	4.3	8.6	15.3	26.2	65.0
2005.....	0.5	0.9	1.7	2.8	5.2	9.6	15.5	28.4	72.2
2004.....	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
2003.....	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Hispanic, both sexes

2007	*	*	1.5	2.2	4.3	6.6	13.6	18.0	37.2
2006	*	*	1.4	2.3	4.9	6.7	13.0	19.6	39.9
2005	*	*	*	2.2	4.0	8.7	16.9	29.3	45.3
2004	*	*	*	2.4	4.8	7.8	11.7	24.0	46.6
2003	*	*	*	*	5.6	5.9	12.6	20.5	57.0
2002	*	*	*	3.0	4.7	10.5	15.5	27.2	61.5
2001	*	*	*	2.8	4.1	7.3	14.2	19.4	63.2
2000	*	*	*	2.9	7.0	11.3	14.1	17.8	61.0
1999	*	*	*	3.3	5.4	9.6	12.9	32.9	70.2

Hispanic, male

2007	*	*	*	*	*	8.1	16.4	17.0	30.5
2006	*	*	*	*	5.6	*	18.0	26.2	39.5
2005	*	*	*	*	*	13.5	21.6	35.0	62.3
2004	*	*	*	*	*	10.8	14.4	26.4	49.0
2003	*	*	*	*	6.8	*	17.0	26.0	58.5
2002	*	*	*	*	*	11.0	19.2	34.3	58.6
2001	*	*	*	*	*	*	16.4	*	55.2
2000	*	*	*	*	*	13.6	17.7	*	66.5
1999	*	*	*	*	*	10.2	20.3	44.0	84.1

Hispanic, female

2007	*	*	*	*	*	*	11.5	18.6	40.8
2006	*	*	*	*	*	6.2	9.3	15.3	40.1
2005	*	*	*	*	*	*	13.6	25.6	36.5
2004	*	*	*	*	*	*	9.8	22.4	45.4
2003	*	*	*	*	*	*	9.5	16.9	56.3
2002	*	*	*	*	*	10.1	12.8	22.6	63.0
2001	*	*	*	*	*	7.0	12.6	17.7	67.0
2000	*	*	*	*	7.2	9.6	11.5	*	58.4
1999	*	*	*	*	*	9.2	*	26.3	63.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 4/, both sexes

2007.....	0.6	1.1	1.8	3.5	5.9	9.8	16.2	27.7	63.2
2006.....	0.7	1.1	1.9	3.4	5.8	10.4	17.9	29.5	67.0
2005.....	0.5	1.1	2.2	3.6	6.5	11.6	19.4	31.3	71.7
2004.....	0.7	1.3	2.1	3.6	7.1	12.2	19.4	32.6	75.6
2003.....	0.7	1.2	2.0	4.1	7.1	11.7	20.4	32.5	76.7
2002.....	0.7	1.1	2.3	4.3	7.8	12.7	20.6	32.5	79.5
2001.....	0.6	1.2	2.3	4.4	7.5	13.2	21.9	33.1	81.1
2000.....	0.8	1.3	2.1	4.4	8.2	12.9	22.2	36.3	84.7
1999.....	0.7	1.2	2.4	4.4	7.1	13.0	21.7	36.1	85.1

Non-Hispanic 4/, male

2007.....	0.7	1.3	2.3	4.3	7.1	11.2	18.9	32.0	62.7
2006.....	0.8	1.3	2.4	4.1	7.6	12.4	21.0	33.9	68.7
2005.....	0.6	1.1	2.6	4.3	7.9	13.7	24.7	35.7	67.3
2004.....	0.8	1.5	2.4	4.0	7.8	13.8	23.3	36.0	72.6
2003.....	0.7	1.4	2.0	4.5	9.4	13.5	25.5	37.6	74.9
2002.....	0.8	1.2	2.4	5.0	9.0	14.4	23.5	39.3	81.7
2001.....	0.7	1.2	2.5	4.9	8.8	15.6	25.0	40.7	83.5
2000.....	0.9	1.5	2.6	4.9	9.9	15.8	27.0	42.3	88.2
1999.....	0.8	1.3	2.7	5.3	8.3	15.9	26.1	40.5	82.0

Non-Hispanic 4/, female

2007.....	0.4	0.9	1.4	2.7	4.8	8.8	14.3	24.9	63.5
2006.....	0.6	0.9	1.5	2.8	4.3	8.7	15.7	26.8	66.2
2005.....	0.5	1.0	1.8	2.9	5.3	9.9	15.6	28.6	73.7
2004.....	0.6	1.0	1.9	3.3	6.4	10.9	16.7	30.6	76.9
2003.....	0.8	1.0	2.0	3.8	5.1	10.3	16.7	29.4	77.6
2002.....	0.6	1.0	2.1	3.7	6.9	11.3	18.5	28.3	78.5
2001.....	0.6	1.2	2.1	4.0	6.4	11.3	19.6	28.6	80.1
2000.....	0.7	1.1	1.6	3.8	6.8	10.6	18.8	32.8	83.3
1999.....	0.6	1.2	2.1	3.6	6.0	10.7	18.6	33.6	86.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic White, both sexes

2007 5/.....	0.5	0.9	1.6	3.2	5.5	9.4	15.7	26.8	62.6
2006 5/.....	0.6	0.9	1.7	3.2	5.5	10.3	17.4	28.2	66.6
2005 5/.....	0.5	1.0	1.9	3.4	6.1	11.1	18.7	30.2	70.9
2004 5/.....	0.7	1.1	2.0	3.2	6.5	11.9	18.8	31.7	73.6
2003 5/.....	0.6	1.1	2.0	3.8	6.6	11.4	19.8	31.1	75.1
2002.....	0.6	1.0	2.1	3.9	7.2	12.3	19.8	31.8	78.2
2001.....	0.6	1.1	2.1	4.0	7.3	12.5	21.1	31.9	78.8
2000.....	0.7	1.0	1.9	3.9	7.8	12.4	21.3	35.5	82.8
1999.....	0.7	1.1	2.1	4.2	6.7	12.6	20.9	34.8	83.0

Non-Hispanic White, male

2007 5/.....	0.6	1.0	2.0	4.0	6.6	10.6	18.2	31.1	62.5
2006 5/.....	0.7	1.1	2.0	3.8	7.1	12.4	20.6	33.2	70.5
2005 5/.....	0.4	1.0	2.2	4.1	7.4	13.2	23.5	34.7	67.8
2004 5/.....	0.8	1.3	2.1	3.7	7.2	13.8	22.7	35.8	71.6
2003 5/.....	0.7	1.2	2.0	4.1	8.8	13.2	24.7	35.9	74.6
2002.....	0.8	1.1	2.1	4.5	8.4	13.8	23.1	38.5	79.9
2001.....	0.6	1.0	2.3	4.5	8.6	14.8	24.6	39.2	81.8
2000.....	0.8	1.2	2.4	4.5	9.5	15.3	26.2	41.7	87.4
1999.....	0.9	1.2	2.5	5.1	8.0	15.6	25.0	39.2	81.0

Non-Hispanic White, female

2007 5/.....	0.4	0.8	1.2	2.5	4.6	8.3	13.8	24.0	62.6
2006 5/.....	0.5	0.8	1.4	2.6	4.0	8.5	14.9	25.0	64.9
2005 5/.....	0.5	0.9	1.6	2.7	5.0	9.3	15.1	27.5	72.4
2004 5/.....	0.5	0.9	1.8	2.8	5.8	10.4	15.9	29.2	74.4
2003 5/.....	0.6	0.9	1.9	3.6	4.8	9.9	16.3	28.2	75.4
2002.....	0.5	1.0	2.0	3.4	6.2	11.0	17.5	27.7	77.4
2001.....	0.5	1.2	1.8	3.6	6.1	10.6	18.7	27.5	77.6
2000.....	0.6	0.9	1.3	3.4	6.4	10.1	17.9	31.8	81.0
1999.....	0.6	1.1	1.8	3.4	5.5	10.2	18.0	32.2	83.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic Black, both sexes

2007 5/.....	1.1	2.4	4.2	6.4	10.5	16.5	25.9	43.5	86.9
2006 5/.....	1.4	2.2	4.1	6.2	10.4	14.9	28.1	50.9	86.3
2005 5/.....	1.0	1.9	4.8	6.0	11.5	19.8	31.5	50.7	96.1
2004 5/.....	1.2	2.4	3.7	7.5	13.0	18.3	31.2	49.9	112.2
2003 5/.....	1.5	2.6	2.9	7.4	12.5	18.2	29.8	57.0	105.6
2002.....	1.6	2.0	4.5	8.3	14.5	19.8	33.9	46.7	107.1
2001.....	1.3	2.2	4.6	8.7	11.6	24.9	35.3	53.8	119.9
2000.....	1.7	3.2	4.0	9.1	13.2	19.6	35.9	52.0	117.0
1999.....	1.0	2.2	5.0	7.1	12.2	19.1	35.1	56.6	119.0

Non-Hispanic Black, male

2007 5/.....	*	3.1	5.3	8.2	14.2	19.6	32.8	50.6	85.5
2006 5/.....	1.7	2.7	5.9	8.2	15.2	17.1	32.2	48.5	66.9
2005 5/.....	*	2.5	6.3	7.5	15.2	23.1	44.5	58.3	79.9
2004 5/.....	*	2.9	5.0	8.5	16.0	19.6	38.7	51.4	103.6
2003 5/.....	*	2.9	3.3	8.9	17.5	19.9	40.2	70.2	89.5
2002.....	*	2.1	5.4	10.2	16.4	24.6	34.7	59.3	119.2
2001.....	*	2.6	4.7	9.2	13.3	30.0	36.0	73.6	119.0
2000.....	2.0	4.1	4.9	9.8	15.5	23.9	43.8	60.1	109.2
1999.....	*	*	5.2	8.5	13.0	24.2	45.4	63.8	107.0

Non-Hispanic Black, female

2007 5/.....	*	1.8	3.2	5.1	7.8	14.4	21.7	39.7	87.5
2006 5/.....	*	1.8	2.6	4.7	6.8	13.5	25.6	52.2	94.2
2005 5/.....	*	*	3.5	4.8	8.7	17.6	23.8	46.7	102.6
2004 5/.....	*	1.9	2.6	6.7	10.7	17.4	26.8	49.1	115.7
2003 5/.....	1.9	2.4	2.5	6.2	8.8	17.0	23.6	50.2	111.8
2002.....	2.0	1.9	3.8	6.8	13.1	16.5	33.4	40.2	102.4
2001.....	*	*	4.6	8.4	10.4	21.6	34.8	43.7	120.2
2000.....	*	2.4	3.3	8.6	11.6	16.8	31.1	48.0	119.9
1999.....	*	*	4.8	6.0	11.6	15.6	28.8	53.1	123.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

All origins 3/, both sexes

2007.....	1.2	1.4	1.6	2.0	2.3	3.5	5.7	8.2	17.4
2006.....	1.1	1.4	1.6	1.9	2.4	3.5	5.6	8.7	18.4
2005.....	1.3	1.6	1.9	2.4	3.3	4.4	7.0	10.8	23.0
2004.....	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
2003.....	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4

All origins 3/, male

2007.....	1.4	1.7	1.9	2.3	2.6	3.8	5.8	8.3	15.8
2006.....	1.3	1.8	1.9	2.3	2.4	3.5	6.1	8.3	16.2
2005.....	1.6	2.1	2.3	3.0	4.0	5.0	6.6	11.1	19.2
2004.....	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
2003.....	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8

All origins 3/, female

2007.....	1.0	1.1	1.3	1.7	2.1	3.3	5.7	8.2	18.2
2006.....	1.0	0.9	1.2	1.6	2.5	3.6	5.3	9.0	19.5
2005.....	1.0	1.2	1.5	1.8	2.7	4.0	7.2	10.7	24.8
2004.....	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
2003.....	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

Non-Hispanic 4/, both sexes

2007.....	1.3	1.4	1.6	2.0	2.3	3.6	5.8	8.2	17.8
2006.....	1.2	1.3	1.6	2.0	2.4	3.6	5.8	8.8	18.8
2005.....	1.3	1.7	2.0	2.4	3.3	4.5	7.0	10.8	23.4
2004.....	1.3	1.5	1.9	2.5	3.2	4.4	7.3	11.7	23.5
2003.....	1.3	1.5	1.7	2.5	3.3	5.1	7.0	12.1	24.3
2002.....	1.2	1.5	2.1	2.6	3.5	4.3	7.6	13.3	24.7
2001.....	1.1	1.3	2.0	2.7	3.8	5.4	7.8	11.9	25.8
2000.....	1.3	1.5	1.8	2.7	3.4	5.4	7.7	12.0	25.6
1999.....	1.0	1.0	1.6	2.2	3.3	4.2	6.9	12.7	23.3

Non-Hispanic 4/, male

2007.....	1.5	1.8	2.0	2.3	2.6	3.9	5.9	8.5	16.3
2006.....	1.3	1.8	2.0	2.3	2.4	3.6	6.2	8.5	16.5
2005.....	1.6	2.2	2.4	3.0	4.0	5.0	6.8	11.4	19.3
2004.....	1.4	1.9	2.2	2.9	3.4	4.7	8.1	12.6	20.2
2003.....	1.4	1.7	2.2	2.7	3.6	5.0	6.9	11.0	19.3
2002.....	1.4	1.6	2.4	2.6	3.6	4.3	7.1	13.5	21.1
2001.....	1.2	1.5	2.3	3.0	3.9	6.1	8.3	12.0	21.6
2000.....	1.5	1.7	2.0	2.7	3.9	5.7	8.0	12.3	22.1
1999.....	1.0	1.2	1.7	2.3	3.8	4.7	7.6	13.3	20.5

Non-Hispanic 4/, female

2007.....	1.1	1.1	1.3	1.8	2.1	3.4	5.8	8.1	18.5
2006.....	1.1	0.9	1.3	1.7	2.4	3.7	5.4	9.1	19.8
2005.....	1.1	1.2	1.6	1.9	2.7	4.0	7.2	10.4	25.3
2004.....	1.1	1.1	1.7	2.1	3.0	4.2	6.7	11.2	25.0
2003.....	1.2	1.3	1.3	2.3	3.1	5.2	7.0	12.7	26.5
2002.....	1.1	1.4	1.8	2.6	3.3	4.3	7.9	13.2	26.2
2001.....	0.9	1.0	1.8	2.4	3.8	4.9	7.4	11.8	27.6
2000.....	1.1	1.3	1.6	2.7	3.0	5.1	7.5	11.8	27.0
1999.....	1.0	0.9	1.5	2.1	2.9	3.8	6.5	12.4	24.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

Non-Hispanic White, both sexes

2007 5/.....	1.0	1.4	1.4	1.8	2.3	3.5	5.8	8.2	17.8
2006 5/.....	1.0	1.2	1.5	1.7	2.3	3.5	5.7	8.5	18.5
2005 5/.....	1.2	1.5	1.8	2.1	3.2	4.2	6.7	10.4	23.9
2004 5/.....	1.1	1.2	1.8	2.3	3.0	4.0	7.0	11.6	23.4
2003 5/.....	1.1	1.4	1.6	2.3	3.1	4.8	6.5	11.8	23.8
2002.....	1.1	1.4	1.9	2.2	3.2	4.1	7.3	12.8	24.5
2001.....	0.9	1.1	1.8	2.4	3.5	5.1	7.3	11.8	25.7
2000.....	1.2	1.4	1.6	2.4	2.9	5.1	7.3	11.9	25.5
1999.....	0.8	0.9	1.6	2.1	2.9	4.0	6.6	12.4	23.0

Non-Hispanic White, male

2007 5/.....	1.2	1.6	1.6	2.1	2.5	3.9	5.9	8.4	16.4
2006 5/.....	1.1	1.5	1.9	1.9	2.3	3.5	6.1	7.7	16.3
2005 5/.....	1.4	2.0	2.2	2.6	3.8	4.6	6.5	10.9	19.4
2004 5/.....	1.3	1.5	2.1	2.7	3.2	4.4	7.8	12.2	20.3
2003 5/.....	1.2	1.5	2.1	2.3	3.2	4.8	6.4	10.3	18.7
2002.....	1.2	1.5	2.1	2.3	3.3	4.0	6.8	12.5	20.5
2001.....	1.2	1.3	2.1	2.5	3.6	5.7	7.8	11.9	21.2
2000.....	1.4	1.6	1.7	2.4	3.4	5.6	7.8	12.1	21.8
1999.....	0.9	1.0	1.7	2.3	3.3	4.5	7.1	12.8	20.3

Non-Hispanic White, female

2007 5/.....	0.8	1.1	1.3	1.6	2.1	3.1	5.8	8.1	18.4
2006 5/.....	0.9	0.8	1.1	1.5	2.2	3.5	5.4	9.0	19.5
2005 5/.....	1.0	1.0	1.4	1.7	2.6	3.9	6.8	10.2	26.0
2004 5/.....	0.9	0.9	1.4	1.9	2.9	3.7	6.3	11.2	24.8
2003 5/.....	1.1	1.2	1.0	2.3	2.9	4.9	6.5	12.7	26.1
2002.....	1.0	1.3	1.7	2.2	3.2	4.1	7.6	12.9	26.3
2001.....	0.7	1.0	1.6	2.3	3.5	4.7	6.9	11.8	27.6
2000.....	1.0	1.1	1.4	2.4	2.5	4.7	7.0	11.8	27.0
1999.....	0.8	0.8	1.5	1.9	2.6	3.6	6.3	12.2	24.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

Non-Hispanic Black, both sexes

2007 5/.....	3.1	2.4	3.4	4.4	3.8	5.6	8.0	10.7	23.0
2006 5/.....	2.8	3.0	2.8	4.5	4.1	5.9	8.3	15.0	26.0
2005 5/.....	2.6	3.6	4.1	5.3	5.6	7.7	11.5	17.6	24.3
2004 5/.....	2.5	3.7	3.5	5.1	5.0	9.1	13.2	14.9	29.1
2003 5/.....	2.7	2.6	3.7	5.3	6.3	9.0	13.9	17.9	33.6
2002.....	2.4	2.9	4.3	5.9	6.4	7.4	13.0	22.7	29.9
2001.....	2.2	2.4	4.0	6.2	7.1	8.9	14.0	14.9	29.7
2000.....	2.5	3.2	4.0	5.3	8.7	9.3	14.5	14.4	29.9
1999.....	2.4	2.4	2.1	4.0	7.0	6.5	11.6	17.9	28.5

Non-Hispanic Black, male

2007 5/.....	3.6	3.1	5.1	5.5	5.0	*	*	*	20.1
2006 5/.....	2.7	4.6	3.0	6.2	*	*	*	17.6	22.9
2005 5/.....	3.5	4.2	4.8	7.1	8.1	10.1	11.5	23.0	23.8
2004 5/.....	2.2	5.2	3.3	6.4	6.0	9.0	14.5	16.9	25.4
2003 5/.....	3.5	3.5	3.8	7.5	7.8	9.1	15.9	21.1	33.0
2002.....	3.1	2.9	5.5	5.7	7.7	7.9	12.2	28.9	31.7
2001.....	1.8	3.1	4.8	8.8	8.1	10.7	16.1	*	28.9
2000.....	3.2	3.8	4.9	5.7	9.3	8.9	13.0	*	30.5
1999.....	2.7	3.0	*	*	9.6	6.9	15.0	21.3	*

Non-Hispanic Black, female

2007 5/.....	2.7	1.8	2.0	3.6	*	6.0	7.7	10.2	24.2
2006 5/.....	2.9	1.6	2.7	3.1	3.7	6.1	8.2	13.7	27.3
2005 5/.....	1.8	3.0	3.4	3.9	3.8	6.0	11.5	14.8	24.5
2004 5/.....	2.7	2.5	3.7	4.2	4.3	9.2	12.4	13.9	30.5
2003 5/.....	2.0	1.8	3.7	3.6	5.1	8.9	12.8	16.2	33.8
2002.....	1.8	3.0	3.3	6.0	5.5	7.1	13.5	19.5	29.3
2001.....	2.5	*	3.3	4.1	6.3	7.8	12.7	14.6	29.9
2000.....	2.0	2.7	3.3	4.9	8.3	9.5	15.4	14.7	29.7
1999.....	2.2	*	*	4.3	5.1	6.2	9.6	16.2	30.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J09-J18)

All origins 3/, both sexes

2007.....	3.5	5.3	7.7	12.1	20.4	39.0	78.4	160.0	463.2
2006.....	3.6	5.7	7.6	13.2	23.1	42.9	86.2	181.8	502.5
2005.....	4.2	6.1	8.8	14.6	24.5	48.6	96.1	202.7	593.9
2004.....	3.6	5.6	8.3	14.2	23.8	47.2	94.7	198.7	582.6
2003.....	4.2	6.4	8.2	15.2	26.6	49.4	103.3	216.8	666.1
2002.....	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001.....	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000.....	3.9	5.6	8.9	15.6	26.6	52.6	106.7	240.8	744.1
1999.....	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8

All origins 3/, male

2007.....	4.1	6.4	9.4	15.1	24.5	46.8	95.1	202.1	542.7
2006.....	4.2	7.1	9.4	16.4	28.1	52.5	108.1	226.8	586.4
2005.....	5.1	7.5	10.7	17.5	29.6	59.0	119.3	253.7	673.5
2004.....	4.3	7.1	9.9	17.2	29.8	58.6	118.3	252.3	677.2
2003.....	4.9	8.1	9.6	17.9	33.3	62.1	130.1	272.3	757.8
2002.....	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001.....	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000.....	4.7	7.0	10.9	18.9	32.9	65.7	134.1	308.8	892.0
1999.....	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7

All origins 3/, female

2007.....	2.9	4.2	6.0	9.4	16.8	32.5	66.0	133.5	425.5
2006.....	3.1	4.4	6.0	10.3	18.8	35.0	70.2	153.7	463.3
2005.....	3.4	4.9	6.9	12.0	20.1	40.2	79.3	171.1	557.4
2004.....	3.0	4.2	6.8	11.4	18.6	37.9	77.7	165.7	540.0
2003.....	3.5	4.8	6.8	12.7	20.7	39.2	84.2	182.9	625.6
2002.....	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001.....	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000.....	3.1	4.3	7.0	12.5	21.2	42.2	87.6	200.7	683.9
1999.....	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J09-J18)

Hispanic, both sexes

2007	1.7	4.0	6.3	11.3	18.8	38.8	70.1	145.8	340.4
2006	3.3	4.1	6.9	14.0	19.8	44.9	87.6	154.4	390.9
2005	3.4	4.9	8.1	13.6	24.4	47.4	89.0	173.9	456.9
2004	2.8	5.0	7.6	12.5	22.8	46.7	96.2	170.8	482.3
2003	2.9	5.3	7.9	14.7	24.6	48.2	93.6	200.4	516.2
2002	2.9	4.0	8.3	13.4	22.8	47.5	102.6	209.4	566.0
2001	2.9	4.4	6.8	12.1	22.2	51.9	98.0	206.1	660.3
2000	3.0	4.8	8.5	14.5	23.0	55.9	95.8	210.0	643.6
1999	2.9	4.4	6.4	13.4	24.7	40.4	81.3	186.2	614.6

Hispanic, male

2007	2.0	5.0	8.8	14.3	24.3	46.2	87.3	188.3	370.0
2006	3.7	5.5	9.3	17.4	24.8	52.6	104.6	174.4	440.9
2005	4.2	6.3	11.3	15.3	28.6	60.8	108.6	216.3	496.9
2004	3.7	6.6	9.1	15.1	26.9	55.1	119.8	204.8	528.2
2003	3.5	7.6	9.1	15.3	30.9	59.8	111.8	257.8	577.2
2002	3.9	5.8	11.0	18.0	29.8	61.9	120.9	266.5	655.9
2001	4.9	5.3	10.0	13.3	25.2	60.8	122.8	270.1	765.6
2000	4.6	6.2	11.0	19.9	28.0	69.5	122.5	295.9	763.9
1999	3.8	6.2	7.4	14.3	31.7	53.5	100.6	243.6	720.3

Hispanic, female

2007	*	3.1	4.0	8.6	14.1	33.0	57.6	117.4	324.3
2006	3.0	2.6	4.5	11.1	15.6	38.9	75.3	141.1	364.3
2005	2.5	3.4	5.1	12.2	20.9	37.0	74.9	145.8	436.0
2004	2.0	3.5	6.2	10.2	19.4	40.2	79.4	148.3	458.8
2003	2.3	3.1	6.8	14.2	19.5	39.3	80.6	163.1	485.2
2002	1.9	*	5.9	9.4	17.1	36.5	89.6	173.0	521.4
2001	*	3.6	4.0	11.0	19.8	45.0	80.4	166.5	608.8
2000	*	3.6	6.4	9.9	19.0	45.7	76.9	157.7	584.6
1999	*	*	5.4	12.6	19.1	30.6	67.7	151.9	563.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J09-J18)

Non-Hispanic 4/, both sexes

2007	3.7	5.4	7.7	12.2	20.5	38.9	78.9	160.7	469.2
2006	3.7	5.9	7.7	13.1	23.3	42.5	86.0	183.1	507.5
2005	4.3	6.3	8.8	14.7	24.5	48.6	96.3	203.9	599.9
2004	3.7	5.6	8.3	14.2	23.7	47.0	94.3	199.7	586.0
2003	4.3	6.5	8.2	15.1	26.5	49.3	103.6	217.2	671.1
2002	4.3	5.6	8.5	14.9	25.3	51.0	106.3	227.3	700.2
2001	4.0	5.4	8.2	14.0	24.4	48.8	101.0	216.7	684.8
2000	3.9	5.6	8.8	15.6	26.6	52.2	106.7	241.0	745.7
1999	4.1	5.3	8.5	14.2	24.8	50.8	106.6	234.7	754.0

Non-Hispanic 4/, male

2007	4.4	6.5	9.4	15.1	24.5	46.8	95.5	202.6	552.2
2006	4.2	7.3	9.4	16.3	28.2	52.2	108.2	229.4	593.8
2005	5.2	7.6	10.6	17.6	29.6	58.7	119.6	255.2	682.2
2004	4.3	7.1	9.8	17.2	29.7	58.6	118.0	254.1	683.2
2003	5.0	8.0	9.6	17.9	33.2	62.0	130.8	272.6	764.6
2002	5.0	6.8	10.1	18.0	31.1	63.7	134.2	287.4	810.7
2001	4.8	6.7	9.9	17.2	30.5	61.1	130.6	274.2	808.2
2000	4.7	7.0	10.8	18.7	32.9	65.2	134.0	307.9	894.9
1999	5.2	6.7	9.7	16.7	31.5	64.5	134.4	302.7	897.9

Non-Hispanic 4/, female

2007	3.1	4.3	6.1	9.4	17.1	32.4	66.6	134.3	430.0
2006	3.1	4.6	6.1	10.2	19.0	34.6	69.7	154.2	467.4
2005	3.5	5.0	7.1	12.0	19.9	40.4	79.4	172.1	562.3
2004	3.1	4.3	6.8	11.5	18.5	37.7	77.3	166.2	542.5
2003	3.7	5.0	6.8	12.5	20.7	39.1	84.2	183.4	630.0
2002	3.5	4.5	6.9	12.0	20.4	40.8	86.7	190.7	652.6
2001	3.2	4.1	6.5	11.1	19.1	39.0	80.2	182.3	633.5
2000	3.3	4.4	7.0	12.7	21.2	41.9	87.7	201.6	685.4
1999	3.1	4.0	7.3	11.9	19.1	40.1	87.2	194.9	696.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J09-J18)

Non-Hispanic White, both sexes

2007 5/.....	3.2	4.9	7.0	11.3	19.4	37.0	77.3	160.4	475.8
2006 5/.....	3.2	5.3	7.2	12.5	21.7	42.0	85.0	182.7	514.4
2005 5/.....	3.9	5.5	7.9	13.7	23.0	46.6	95.1	204.9	612.6
2004 5/.....	3.2	4.9	7.6	13.3	22.0	45.9	92.4	199.1	593.8
2003 5/.....	3.8	5.7	7.2	13.8	25.0	47.6	102.5	218.5	684.5
2002.....	3.9	5.0	7.6	13.8	24.5	49.8	105.0	227.7	712.7
2001.....	3.3	4.8	7.5	12.9	23.2	47.4	98.2	216.2	693.9
2000.....	3.3	4.9	7.9	14.2	24.4	49.7	105.5	240.7	759.8
1999.....	3.4	4.6	7.7	12.9	23.8	48.9	106.1	236.5	766.9

Non-Hispanic White, male

2007 5/.....	3.5	5.8	8.6	13.8	22.5	43.8	92.1	200.9	557.8
2006 5/.....	3.7	6.3	8.8	15.2	26.1	50.7	106.2	228.4	598.8
2005 5/.....	4.5	6.6	9.1	16.2	27.3	55.6	117.8	255.1	693.4
2004 5/.....	3.7	6.1	9.0	15.9	26.7	56.5	113.8	252.6	691.1
2003 5/.....	4.3	7.1	8.3	15.5	30.5	58.8	127.6	271.3	774.1
2002.....	4.4	6.0	8.8	16.3	29.6	62.0	131.3	286.1	820.6
2001.....	4.0	5.7	8.9	15.3	28.3	58.5	125.1	271.6	813.9
2000.....	3.7	5.7	9.2	16.9	29.5	61.4	131.6	305.1	908.1
1999.....	4.2	5.5	8.6	15.0	29.7	61.0	132.9	302.9	909.3

Non-Hispanic White, female

2007 5/.....	2.8	4.0	5.5	9.0	16.6	31.4	66.2	134.5	436.9
2006 5/.....	2.7	4.4	5.6	10.0	17.8	34.7	69.1	153.9	475.0
2005 5/.....	3.2	4.5	6.8	11.3	19.2	39.2	78.3	173.6	575.7
2004 5/.....	2.8	3.8	6.3	10.9	17.8	37.2	76.8	165.9	550.2
2003 5/.....	3.2	4.4	6.2	12.1	20.0	38.4	84.4	186.0	645.1
2002.....	3.4	4.1	6.3	11.5	20.0	39.8	86.1	192.0	666.1
2001.....	2.6	3.9	6.1	10.6	18.7	38.3	79.2	182.8	644.0
2000.....	2.9	4.0	6.7	11.7	20.0	40.2	87.2	202.5	700.0
1999.....	2.6	3.7	6.8	11.0	18.6	39.3	87.3	197.2	710.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J09-J18)

Non-Hispanic Black, both sexes

2007 5/.....	7.5	9.6	14.9	20.9	32.8	58.8	98.3	172.9	406.0
2006 5/.....	7.3	11.0	13.2	20.8	40.5	51.7	101.4	197.4	439.6
2005 5/.....	8.1	12.5	17.2	24.8	40.0	70.0	117.2	202.0	475.4
2004 5/.....	7.2	11.8	15.5	23.9	40.6	62.9	118.6	216.5	511.7
2003 5/.....	8.7	12.7	16.1	27.6	42.1	71.1	123.9	207.9	528.3
2002.....	7.7	10.3	17.6	26.8	35.6	70.0	124.3	233.8	579.8
2001.....	9.2	10.8	15.4	23.7	34.9	66.2	133.9	224.9	583.1
2000.....	8.6	12.4	18.1	27.6	46.2	78.5	124.8	247.4	604.5
1999.....	9.9	11.9	16.3	26.6	37.0	76.4	119.8	220.4	634.7

Non-Hispanic Black, male

2007 5/.....	10.1	12.7	18.1	29.1	45.4	80.7	136.8	231.6	480.8
2006 5/.....	8.4	15.7	16.4	30.3	51.5	70.0	141.8	243.7	512.2
2005 5/.....	10.6	16.4	24.4	33.6	54.9	93.0	146.0	262.6	537.6
2004 5/.....	8.8	16.1	19.1	32.9	62.1	88.1	172.1	298.1	607.6
2003 5/.....	10.5	16.8	22.0	40.5	62.0	101.8	179.4	287.5	642.5
2002.....	10.1	13.2	24.2	36.7	48.1	91.0	175.1	324.5	695.8
2001.....	11.6	15.9	20.8	34.3	50.8	95.4	203.4	309.9	718.4
2000.....	11.7	17.6	26.2	35.5	65.0	108.7	173.4	345.9	763.2
1999.....	13.6	17.4	21.8	34.2	52.6	110.8	162.4	311.7	803.4

Non-Hispanic Black, female

2007 5/.....	5.2	7.1	12.3	14.5	23.6	43.7	75.1	142.0	375.1
2006 5/.....	6.3	7.0	10.5	13.5	32.4	39.1	77.2	172.9	409.9
2005 5/.....	5.9	9.2	11.3	17.9	29.0	54.2	100.0	170.0	450.4
2004 5/.....	5.9	8.1	12.6	16.9	24.6	45.8	87.0	173.5	473.8
2003 5/.....	7.1	9.2	11.3	17.6	27.4	50.4	91.0	166.3	484.1
2002.....	5.5	7.9	12.1	19.1	26.4	56.0	93.9	186.7	535.5
2001.....	7.1	6.6	11.1	15.4	23.1	46.8	92.5	181.6	533.0
2000.....	6.0	8.1	11.4	21.3	32.2	58.5	95.5	197.9	545.6
1999.....	6.7	7.3	11.8	20.7	25.4	53.5	94.1	175.2	571.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J09-J11)

All origins 3/, both sexes

2007.....	*	*	*	0.1	*	0.3	0.4	0.8	2.5
2006.....	*	*	0.1	0.2	0.3	0.6	1.3	2.5	7.0
2005.....	*	0.1	0.1	0.3	0.4	1.1	2.4	5.4	19.3
2004.....	*	*	0.1	0.3	0.3	0.8	1.6	3.2	11.4
2003.....	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	0.7	2.2
2000.....	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2

All origins 3/, male

2007.....	*	*	*	*	*	*	*	1.0	3.0
2006.....	*	*	*	*	*	0.7	1.3	3.0	8.1
2005.....	*	*	*	0.4	*	1.1	2.5	5.3	18.2
2004.....	*	*	*	*	*	1.1	1.8	3.3	11.1
2003.....	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9

All origins 3/, female

2007.....	*	*	*	*	*	*	*	0.7	2.3
2006.....	*	*	*	*	*	0.5	1.3	2.2	6.5
2005.....	*	*	*	*	0.5	1.1	2.3	5.5	19.8
2004.....	*	*	*	*	*	0.6	1.4	3.1	11.5
2003.....	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	0.6	1.1	1.9	5.4	21.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J09-J11)

Non-Hispanic 4/, both sexes

2007	*	*	*	*	*	0.3	0.5	0.9	2.6
2006	*	*	0.1	*	0.3	0.6	1.3	2.6	7.3
2005	*	0.1	0.2	0.3	0.5	1.1	2.4	5.6	19.8
2004	*	*	0.2	0.3	0.3	0.8	1.6	3.3	11.8
2003	0.1	0.2	0.3	0.3	0.8	1.4	2.8	4.8	17.5
2002	*	*	*	*	*	0.4	0.8	2.3	9.0
2001	*	*	*	*	*	*	*	0.7	2.3
2000	0.1	0.1	0.3	0.5	0.8	1.5	2.6	6.2	20.5
1999	*	*	0.2	0.4	0.6	1.2	2.1	6.2	21.6

Non-Hispanic 4/, male

2007	*	*	*	*	*	*	*	1.1	3.1
2006	*	*	*	*	*	0.8	1.3	3.2	8.4
2005	*	*	*	0.4	*	1.1	2.7	5.5	18.7
2004	*	*	*	*	*	1.1	1.9	3.4	11.6
2003	*	*	0.4	*	0.8	1.4	3.0	4.7	17.5
2002	*	*	*	*	*	*	0.9	2.7	9.2
2001	*	*	*	*	*	*	*	*	2.3
2000	*	*	*	0.5	1.0	1.7	2.8	6.2	21.1
1999	*	*	*	0.5	0.6	1.2	2.4	7.5	20.7

Non-Hispanic 4/, female

2007	*	*	*	*	*	*	*	0.7	2.3
2006	*	*	*	*	*	0.5	1.3	2.3	6.7
2005	*	*	*	*	0.5	1.1	2.3	5.7	20.3
2004	*	*	*	*	*	0.6	1.5	3.2	11.8
2003	*	*	*	0.3	0.7	1.3	2.6	4.8	17.6
2002	*	*	*	*	*	*	0.7	2.1	8.9
2001	*	*	*	*	*	*	*	0.8	2.2
2000	*	*	*	0.6	0.7	1.3	2.5	6.2	20.3
1999	*	*	*	*	0.7	1.2	1.9	5.4	22.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J09-J11)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	*	0.3	0.5	0.9	2.8
2006 5/.....	*	*	0.2	*	0.3	0.7	1.5	2.8	7.8
2005 5/.....	*	*	0.2	0.3	0.5	1.2	2.6	6.0	21.5
2004 5/.....	*.1	*	0.2	0.3	0.3	0.9	1.6	3.6	12.7
2003 5/.....	0.1	0.2	0.3	0.3	0.8	1.5	3.0	5.2	18.8
2002.....	*	*	*	*	*	0.5	0.9	2.5	9.6
2001.....	*	*	*	*	*	*	*	0.7	2.4
2000.....	*	0.2	0.3	0.6	0.9	1.5	2.8	6.5	21.7
1999.....	*	*	0.2	0.4	0.7	1.2	2.2	6.5	23.3

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	*	1.1	3.4
2006 5/.....	*	*	*	*	*	0.8	1.4	3.3	8.9
2005 5/.....	*	*	*	*	*	1.2	2.8	5.8	19.9
2004 5/.....	*	*	*	*	*	1.1	1.7	3.7	12.7
2003 5/.....	*	*	0.3	*	0.9	1.6	3.2	5.2	18.9
2002.....	*	*	*	*	*	*	1.0	2.9	10.0
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	0.5	1.1	1.8	2.8	6.7	22.0
1999.....	*	*	*	0.5	0.6	1.2	2.7	7.8	22.5

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	0.8	2.6
2006 5/.....	*	*	*	*	*	0.6	1.5	2.5	7.3
2005 5/.....	*	*	*	*	0.5	1.2	2.5	6.1	22.2
2004 5/.....	*	*	*	*	*	0.7	1.5	3.5	12.8
2003 5/.....	*	*	*	*	0.7	1.5	2.8	5.1	18.7
2002.....	*	*	*	*	*	*	0.8	2.3	9.5
2001.....	*	*	*	*	*	*	*	0.7	2.4
2000.....	*	*	*	0.6	0.7	1.2	2.8	6.4	21.6
1999.....	*	*	*	*	0.8	1.2	1.9	5.7	23.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

All origins 3/, both sexes

2007.....	3.5	5.2	7.6	12.0	20.3	38.7	77.9	159.2	460.7
2006.....	3.6	5.6	7.5	13.0	22.8	42.3	85.0	179.3	495.5
2005.....	4.2	6.0	8.6	14.3	24.1	47.5	93.7	197.3	574.7
2004.....	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
2003.....	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6

All origins 3/, male

2007.....	4.1	6.3	9.3	15.0	24.4	46.5	94.6	201.1	539.7
2006.....	4.2	7.0	9.3	16.3	27.8	51.8	106.9	223.8	578.3
2005.....	5.0	7.4	10.5	17.1	29.2	57.9	116.8	248.4	655.3
2004.....	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
2003.....	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7

All origins 3/, female

2007.....	2.9	4.1	5.9	9.3	16.7	32.2	65.6	132.8	423.2
2006.....	3.1	4.3	5.9	10.1	18.5	34.5	69.0	151.5	456.7
2005.....	3.4	4.7	6.8	11.7	19.6	39.1	77.0	165.6	537.6
2004.....	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
2003.....	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

Hispanic, both sexes

2007.....	1.7	4.0	6.3	11.0	18.8	38.5	70.1	145.5	339.3
2006.....	3.3	4.0	6.9	13.8	19.7	44.9	87.3	153.8	388.9
2005.....	3.3	4.7	8.1	13.5	24.4	46.7	87.8	172.1	450.5
2004.....	2.8	5.0	7.6	12.5	22.6	46.2	95.9	170.4	479.5
2003.....	2.9	5.3	7.7	14.3	24.2	48.2	92.6	197.2	511.0
2002.....	2.9	4.0	8.3	13.4	22.7	47.4	102.1	209.0	563.2
2001.....	2.9	4.4	6.8	12.1	22.2	51.7	97.1	205.1	660.3
2000.....	2.9	4.8	8.4	14.4	22.5	55.7	94.9	206.1	633.0
1999.....	2.9	4.2	6.1	13.0	24.0	39.6	79.7	183.8	607.6

Hispanic, male

2007.....	1.9	4.8	8.8	14.0	24.3	45.8	87.3	188.3	370.0
2006.....	3.6	5.4	9.3	17.2	24.5	52.6	104.6	174.4	438.7
2005.....	4.0	6.1	11.3	15.1	28.6	60.0	108.6	214.5	488.2
2004.....	3.7	6.6	9.1	15.1	26.9	54.2	119.2	203.8	526.8
2003.....	3.5	7.6	8.9	15.0	29.9	59.8	111.2	251.6	571.1
2002.....	3.8	5.8	11.0	18.0	29.5	61.9	120.9	265.4	654.2
2001.....	4.8	5.3	10.0	13.3	25.2	60.8	122.8	267.5	765.6
2000.....	4.5	6.2	10.7	19.6	27.2	69.5	121.8	290.1	751.8
1999.....	3.8	5.9	6.9	13.7	30.1	51.9	99.9	242.1	718.2

Hispanic, female

2007.....	*	3.1	4.0	8.4	14.1	32.7	57.6	116.9	322.7
2006.....	3.0	2.5	4.5	10.7	15.6	38.9	74.9	140.1	362.5
2005.....	2.5	3.3	5.0	12.2	20.9	36.4	72.9	144.0	430.8
2004.....	1.8	3.4	6.2	10.2	19.2	39.9	79.4	148.3	455.2
2003.....	2.2	3.0	6.5	13.7	19.5	39.3	79.3	161.7	480.5
2002.....	1.9	*	5.9	9.4	17.1	36.2	88.7	173.0	518.1
2001.....	*	3.6	4.0	11.0	19.8	44.7	78.9	166.5	608.8
2000.....	*	3.6	6.4	9.9	18.7	45.3	75.8	155.0	574.7
1999.....	*	*	5.4	12.3	19.1	30.3	65.5	149.1	554.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

Non-Hispanic 4/, both sexes

2007	3.7	5.3	7.6	12.0	20.4	38.6	78.4	159.8	466.6
2006	3.6	5.8	7.6	13.0	23.0	41.9	84.6	180.4	500.2
2005	4.2	6.1	8.6	14.3	24.0	47.5	93.9	198.2	580.1
2004	3.6	5.6	8.1	14.0	23.4	46.2	92.7	196.4	574.2
2003	4.2	6.3	7.9	14.8	25.8	47.9	100.8	212.4	653.6
2002	4.2	5.5	8.4	14.7	25.1	50.5	105.5	225.0	691.2
2001	4.0	5.4	8.1	13.9	24.3	48.7	100.8	216.0	682.5
2000	3.8	5.5	8.5	15.1	25.8	50.7	104.1	234.8	725.2
1999	4.1	5.3	8.2	13.8	24.2	49.6	104.4	228.5	732.4

Non-Hispanic 4/, male

2007	4.4	6.4	9.3	15.0	24.4	46.5	95.0	201.5	549.1
2006	4.2	7.2	9.3	16.2	27.9	51.5	106.8	226.2	585.5
2005	5.0	7.5	10.4	17.3	29.2	57.6	117.0	249.7	663.5
2004	4.3	7.0	9.7	17.0	29.2	57.5	116.1	250.7	671.6
2003	4.9	7.8	9.3	17.7	32.3	60.6	127.8	267.8	747.1
2002	4.9	6.7	10.0	17.7	30.8	63.1	133.3	284.7	801.5
2001	4.8	6.7	9.8	17.1	30.4	60.9	130.4	273.6	805.9
2000	4.5	6.8	10.5	18.2	31.9	63.5	131.3	301.6	873.8
1999	5.1	6.6	9.5	16.3	30.9	63.3	132.0	295.2	877.2

Non-Hispanic 4/, female

2007	3.0	4.2	6.0	9.3	16.9	32.1	66.2	133.5	427.7
2006	3.1	4.5	5.9	10.0	18.7	34.1	68.4	151.9	460.6
2005	3.5	4.8	7.0	11.7	19.4	39.3	77.2	166.4	542.0
2004	3.0	4.2	6.7	11.2	18.2	37.1	75.9	163.0	530.7
2003	3.5	4.8	6.5	12.1	20.0	37.7	81.6	178.6	612.4
2002	3.5	4.4	6.8	11.9	20.2	40.4	86.0	188.7	643.7
2001	3.1	4.1	6.5	10.9	19.0	38.9	80.0	181.6	631.3
2000	3.1	4.3	6.8	12.2	20.5	40.6	85.2	195.4	665.2
1999	3.0	4.0	7.0	11.6	18.4	38.9	85.3	189.6	674.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

Non-Hispanic White, both sexes

2007 5/.....	3.1	4.8	6.9	11.2	19.2	36.7	76.9	159.4	472.9
2006 5/.....	3.2	5.2	7.0	12.3	21.4	41.3	83.5	179.9	506.5
2005 5/.....	3.8	5.4	7.8	13.3	22.5	45.4	92.5	198.9	591.2
2004 5/.....	3.2	4.9	7.4	13.0	21.7	45.0	90.8	195.5	581.1
2003 5/.....	3.6	5.6	6.9	13.4	24.2	46.1	99.5	213.4	665.7
2002.....	3.8	5.0	7.5	13.7	24.3	49.3	104.1	225.2	703.1
2001.....	3.3	4.7	7.4	12.8	23.1	47.3	98.0	215.5	691.5
2000.....	3.2	4.7	7.6	13.7	23.6	48.2	102.7	234.2	738.1
1999.....	3.3	4.5	7.5	12.5	23.1	47.7	103.9	230.0	743.7

Non-Hispanic White, male

2007 5/.....	3.5	5.7	8.5	13.6	22.3	43.5	91.6	199.8	554.4
2006 5/.....	3.6	6.2	8.7	15.0	25.7	49.9	104.8	225.2	589.9
2005 5/.....	4.4	6.5	8.9	15.8	26.8	54.4	115.0	249.3	673.6
2004 5/.....	3.7	6.1	8.9	15.6	26.3	55.4	112.1	248.9	678.4
2003 5/.....	4.2	6.8	8.0	15.3	29.6	57.3	124.4	266.2	755.2
2002.....	4.3	5.9	8.7	16.1	29.4	61.4	130.3	283.2	810.6
2001.....	4.0	5.6	8.8	15.3	28.3	58.4	124.9	270.9	811.5
2000.....	3.7	5.6	8.9	16.4	28.5	59.6	128.7	298.4	886.1
1999.....	4.1	5.5	8.4	14.5	29.1	59.8	130.3	295.2	886.8

Non-Hispanic White, female

2007 5/.....	2.7	4.0	5.4	8.9	16.5	31.1	65.8	133.7	434.3
2006 5/.....	2.7	4.2	5.5	9.8	17.5	34.1	67.7	151.4	467.7
2005 5/.....	3.2	4.3	6.6	11.0	18.6	37.9	75.8	167.5	553.5
2004 5/.....	2.7	3.7	6.1	10.6	17.6	36.5	75.3	162.5	537.5
2003 5/.....	3.1	4.3	6.0	11.7	19.3	36.9	81.5	180.9	626.4
2002.....	3.4	4.0	6.2	11.4	19.9	39.4	85.4	189.8	656.6
2001.....	2.6	3.9	6.0	10.5	18.6	38.2	78.9	182.1	641.7
2000.....	2.8	3.9	6.3	11.1	19.3	39.0	84.5	196.0	678.4
1999.....	2.5	3.6	6.6	10.6	17.8	38.1	85.3	191.5	686.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

Non-Hispanic Black, both sexes

2007 5/.....	7.5	9.5	14.7	20.9	32.7	58.7	97.9	172.9	405.8
2006 5/.....	7.2	10.9	13.2	20.7	40.3	51.5	101.0	195.4	437.7
2005 5/.....	8.0	12.3	17.1	24.4	39.8	69.6	116.3	199.8	471.3
2004 5/.....	7.2	11.7	15.5	23.6	40.3	62.3	116.5	215.4	509.2
2003 5/.....	8.4	12.5	15.8	27.5	41.5	70.6	123.1	206.1	522.4
2002.....	7.7	10.3	17.5	26.5	35.2	69.8	124.1	233.0	577.4
2001.....	9.2	10.8	15.4	23.5	34.6	66.2	133.9	223.8	581.5
2000.....	8.4	12.3	17.9	27.2	45.6	77.0	123.9	245.1	595.0
1999.....	9.9	11.8	16.0	26.4	36.7	75.0	118.8	217.5	628.9

Non-Hispanic Black, male

2007 5/.....	10.1	12.7	17.9	29.1	45.4	80.4	136.3	231.6	480.8
2006 5/.....	8.4	15.6	16.4	30.1	51.2	70.0	141.3	240.2	509.4
2005 5/.....	10.4	16.2	24.3	32.8	54.9	92.3	145.6	261.2	530.0
2004 5/.....	8.7	16.1	19.1	32.7	61.8	87.1	167.9	297.4	606.6
2003 5/.....	10.1	16.6	21.4	40.3	61.3	101.4	178.5	286.8	639.3
2002.....	10.1	13.2	24.1	36.3	47.4	91.0	175.1	322.9	693.6
2001.....	11.6	15.9	20.6	33.9	50.3	95.4	203.4	309.9	716.1
2000.....	11.4	17.4	25.9	35.3	64.5	107.6	171.5	345.0	750.3
1999.....	13.5	17.3	21.8	34.0	52.1	109.4	161.9	307.3	799.9

Non-Hispanic Black, female

2007 5/.....	5.2	6.9	12.2	14.4	23.4	43.7	74.8	142.0	374.7
2006 5/.....	6.3	7.0	10.5	13.3	32.2	38.9	77.0	171.8	408.4
2005 5/.....	5.9	9.0	11.3	17.8	28.7	54.0	98.9	167.3	447.7
2004 5/.....	5.9	8.0	12.6	16.5	24.2	45.5	86.1	172.3	470.6
2003 5/.....	6.9	9.0	11.2	17.5	26.8	49.8	90.2	163.9	477.1
2002.....	5.5	7.9	12.1	18.9	26.2	55.5	93.6	186.3	533.0
2001.....	7.1	6.6	11.1	15.4	23.1	46.8	92.5	179.9	531.7
2000.....	5.8	8.1	11.4	20.8	31.7	56.7	95.2	194.9	537.3
1999.....	6.7	7.2	11.4	20.5	25.4	52.2	92.9	173.0	565.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22,U04)

All origins 3/, both sexes

2007	*	*	*	*	*	*	*	0.4	1.5
2006	*	*	*	*	*	*	0.3	0.6	2.0
2005	*	*	*	*	*	*	0.4	0.9	3.3
2004	*	*	*	*	*	*	0.3	0.6	2.6
2003	*	*	*	*	*	*	0.5	0.8	3.5
2002	*	*	*	*	*	0.3	0.3	0.8	3.3
2001	*	*	*	*	*	*	0.3	0.7	3.3
2000	*	*	*	*	*	0.3	0.6	0.8	4.5
1999	*	*	*	*	*	0.2	0.6	1.3	6.0

All origins 3/, male

2007	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	1.8
2005	*	*	*	*	*	*	*	1.3	3.1
2004	*	*	*	*	*	*	*	*	2.5
2003	*	*	*	*	*	*	*	*	2.4
2002	*	*	*	*	*	*	*	*	2.8
2001	*	*	*	*	*	*	*	*	2.3
2000	*	*	*	*	*	*	0.7	*	3.8
1999	*	*	*	*	*	*	0.7	1.6	4.0

All origins 3/, female

2007	*	*	*	*	*	*	*	*	1.7
2006	*	*	*	*	*	*	*	0.7	2.1
2005	*	*	*	*	*	*	*	0.6	3.5
2004	*	*	*	*	*	*	*	0.7	2.7
2003	*	*	*	*	*	*	*	0.8	3.9
2002	*	*	*	*	*	*	*	0.9	3.5
2001	*	*	*	*	*	*	*	0.7	3.7
2000	*	*	*	*	*	*	0.5	0.9	4.8
1999	*	*	*	*	*	*	0.5	1.1	6.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22,U04)

Non-Hispanic 4/, both sexes

2007.....	*	*	*	*	*	*	*	0.4	1.5
2006.....	*	*	*	*	*	*	0.3	0.7	2.1
2005.....	*	*	*	*	*	*	0.4	0.9	3.4
2004.....	*	*	*	*	*	*	0.3	0.6	2.6
2003.....	*	*	*	*	*	*	0.5	0.9	3.6
2002.....	*	*	*	*	*	0.3	0.3	0.9	3.4
2001.....	*	*	*	*	*	*	0.3	0.6	3.3
2000.....	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	0.2	0.6	1.3	6.1

Non-Hispanic 4/, male

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	1.9
2005.....	*	*	*	*	*	*	*	1.3	3.2
2004.....	*	*	*	*	*	*	*	*	2.5
2003.....	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	0.7	*	3.7
1999.....	*	*	*	*	*	*	0.8	1.7	4.1

Non-Hispanic 4/, female

2007.....	*	*	*	*	*	*	*	*	1.8
2006.....	*	*	*	*	*	*	*	0.8	2.2
2005.....	*	*	*	*	*	*	*	0.6	3.6
2004.....	*	*	*	*	*	*	*	0.7	2.7
2003.....	*	*	*	*	*	*	*	0.8	4.1
2002.....	*	*	*	*	*	*	*	0.9	3.6
2001.....	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	0.5	1.1	6.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22,U04)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	*	*	*	*	0.5	1.6
2006 5/.....	*	*	*	*	*	*	*	0.8	2.3
2005 5/.....	*	*	*	*	*	*	0.4	0.9	3.6
2004 5/.....	*	*	*	*	*	*	*	0.7	2.8
2003 5/.....	*	*	*	*	*	*	0.5	0.9	3.8
2002.....	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	0.7	1.4	6.4

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	*	*	*
2006 5/.....	*	*	*	*	*	*	*	*	2.0
2005 5/.....	*	*	*	*	*	*	*	1.4	3.4
2004 5/.....	*	*	*	*	*	*	*	*	2.6
2003 5/.....	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	3.0
2001.....	*	*	*	*	*	*	*	*	2.5
2000.....	*	*	*	*	*	*	0.8	*	3.7
1999.....	*	*	*	*	*	*	0.8	1.7	4.1

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	*	1.9
2006 5/.....	*	*	*	*	*	*	*	0.9	2.4
2005 5/.....	*	*	*	*	*	*	*	*	3.7
2004 5/.....	*	*	*	*	*	*	*	0.8	2.9
2003 5/.....	*	*	*	*	*	*	*	0.8	4.3
2002.....	*	*	*	*	*	*	*	0.9	3.7
2001.....	*	*	*	*	*	*	*	0.8	3.9
2000.....	*	*	*	*	*	*	*	1.0	5.1
1999.....	*	*	*	*	*	*	0.5	1.2	7.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes

2007.....	6.1	13.1	25.4	56.2	111.4	194.1	310.7	443.7	596.1
2006.....	5.8	12.7	25.8	57.4	111.8	194.8	308.5	434.8	589.1
2005.....	6.5	12.6	26.9	62.1	118.6	210.3	330.6	457.8	637.2
2004.....	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
2003.....	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0

All origins 3/, male

2007.....	6.2	13.8	28.0	61.4	123.7	219.0	363.8	550.2	758.7
2006.....	5.5	13.4	27.4	62.0	119.8	218.0	364.6	541.3	760.8
2005.....	6.4	13.6	28.5	67.6	132.9	240.0	393.6	564.7	819.9
2004.....	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
2003.....	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7

All origins 3/, female

2007.....	6.0	12.4	23.0	51.4	100.6	173.7	271.6	376.4	518.7
2006.....	6.1	12.1	24.2	53.3	104.8	175.8	267.5	368.3	508.7
2005.....	6.6	11.7	25.5	57.1	106.1	186.2	284.8	391.5	553.3
2004.....	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
2003.....	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

Hispanic, both sexes

2007.....	2.3	3.9	7.8	15.6	33.5	60.2	125.6	223.0	357.2
2006.....	1.8	3.7	8.1	17.7	32.1	62.9	120.6	203.2	374.4
2005.....	2.4	5.0	8.6	18.3	33.3	72.9	136.6	226.6	414.6
2004.....	2.1	4.4	9.4	16.3	35.8	72.3	124.4	212.4	391.5
2003.....	2.8	4.4	9.4	17.4	36.9	74.0	129.9	237.7	460.2
2002.....	2.4	4.9	9.0	20.6	38.4	73.1	136.0	242.9	457.3
2001.....	2.4	5.2	9.5	19.9	42.0	73.1	145.3	212.1	480.2
2000.....	2.5	4.4	9.1	20.9	40.0	69.6	143.2	252.3	481.1
1999.....	3.1	4.9	10.4	19.9	45.3	84.6	155.9	265.5	536.0

Hispanic, male

2007.....	2.6	4.6	9.3	19.3	39.7	88.0	169.0	292.1	461.5
2006.....	1.8	4.0	8.9	21.7	39.5	79.7	155.0	262.0	484.9
2005.....	1.9	5.8	9.4	25.4	42.2	94.5	184.2	315.2	526.8
2004.....	1.7	4.2	9.6	19.8	45.4	99.4	167.0	270.9	518.4
2003.....	2.8	3.7	11.8	23.4	40.9	103.8	170.7	334.7	646.5
2002.....	2.7	5.4	11.7	25.7	51.5	92.2	199.7	321.4	602.4
2001.....	2.3	4.4	10.6	23.6	57.1	99.3	195.1	279.2	676.1
2000.....	2.4	4.8	8.8	30.5	50.7	100.6	191.9	366.6	671.1
1999.....	2.5	5.0	11.6	25.3	52.5	120.7	216.9	385.1	836.8

Hispanic, female

2007.....	1.9	3.1	6.3	12.3	28.3	38.5	94.0	176.9	300.6
2006.....	1.8	3.4	7.3	14.1	25.8	49.9	95.8	164.0	315.7
2005.....	2.9	4.3	7.7	12.0	25.8	56.1	102.4	167.8	355.8
2004.....	2.4	4.7	9.2	13.2	27.8	51.4	94.1	173.8	326.3
2003.....	2.8	5.1	7.1	12.2	33.6	51.1	100.9	174.6	365.6
2002.....	2.1	4.4	6.4	16.1	27.6	58.5	90.5	192.7	385.5
2001.....	2.6	6.0	8.5	16.7	29.8	53.0	109.9	170.5	384.4
2000.....	2.7	4.0	9.3	12.7	31.4	46.0	108.8	182.8	387.8
1999.....	3.7	4.9	9.4	15.1	39.4	57.5	112.9	194.1	391.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 4/, both sexes

2007	6.6	14.1	27.0	59.5	117.5	204.3	323.2	456.2	608.3
2006	6.3	13.6	27.3	60.5	118.0	204.3	320.5	447.3	599.3
2005	6.9	13.3	28.4	65.5	125.1	219.9	342.3	469.5	647.1
2004	5.9	12.4	26.8	63.3	119.4	209.0	326.8	445.1	610.3
2003	6.2	12.5	28.9	67.2	127.1	219.5	340.6	465.1	641.2
2002	6.1	12.6	29.6	63.8	124.6	219.7	339.4	474.4	643.5
2001	5.7	12.4	30.9	65.9	129.7	223.3	337.5	461.7	649.4
2000	6.1	12.3	31.0	65.3	129.3	226.0	335.4	480.7	653.2
1999	5.7	12.5	33.5	69.7	135.5	234.6	347.3	491.4	648.0

Non-Hispanic 4/, male

2007	6.7	14.8	29.6	64.7	130.1	228.7	376.8	565.2	775.3
2006	5.9	14.3	29.0	65.0	125.9	227.6	377.8	557.0	775.2
2005	6.9	14.3	30.0	70.7	139.5	249.8	406.0	577.7	833.9
2004	6.0	13.3	28.3	67.9	131.7	236.2	390.8	556.6	824.6
2003	6.7	13.0	30.7	72.3	138.8	252.6	407.6	579.0	874.6
2002	6.2	13.5	30.9	68.8	136.7	254.8	415.0	606.7	906.0
2001	6.1	12.9	32.9	69.9	142.2	261.1	416.4	594.6	924.8
2000	6.5	12.6	33.4	70.3	144.8	265.6	412.8	637.1	978.0
1999	5.9	12.6	35.9	77.7	156.5	285.5	440.8	656.5	1,000.2

Non-Hispanic 4/, female

2007	6.5	13.4	24.5	54.7	106.5	184.2	283.8	387.7	529.5
2006	6.6	13.0	25.7	56.5	111.0	185.4	278.6	379.0	517.5
2005	7.0	12.4	27.0	60.7	112.4	195.7	296.1	402.5	561.8
2004	5.8	11.5	25.4	59.1	108.6	187.1	280.7	376.7	514.5
2003	5.7	12.0	27.2	62.5	116.9	192.9	293.0	395.7	538.6
2002	6.0	11.7	28.5	59.2	114.1	191.6	286.1	394.1	530.3
2001	5.3	12.0	29.0	62.3	118.9	193.3	282.4	382.2	534.8
2000	5.7	12.1	28.8	60.8	115.9	194.7	281.5	388.6	521.9
1999	5.5	12.4	31.2	62.4	117.4	194.8	282.3	394.8	507.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic White, both sexes

2007 5/.....	6.4	14.3	28.3	63.4	126.2	220.9	344.3	483.1	640.6
2006 5/.....	6.3	13.8	28.4	64.1	126.9	221.1	340.5	472.0	629.8
2005 5/.....	6.7	13.2	29.4	69.7	134.0	236.0	363.2	494.9	679.0
2004 5/.....	5.5	12.3	27.6	67.1	128.6	224.4	345.9	469.6	640.0
2003 5/.....	6.0	12.4	29.6	71.3	136.6	233.4	359.4	489.6	671.6
2002.....	5.8	12.4	30.6	67.2	133.4	233.0	357.7	496.4	673.1
2001.....	5.4	12.3	31.8	69.9	137.9	236.4	355.6	483.3	676.9
2000.....	5.5	12.0	31.7	68.8	136.5	238.6	351.8	503.7	680.8
1999.....	5.0	12.3	34.3	73.2	143.2	246.1	361.3	512.6	675.6

Non-Hispanic White, male

2007 5/.....	6.6	15.1	30.7	67.0	136.9	242.1	391.8	588.5	806.5
2006 5/.....	5.8	14.2	29.6	66.7	132.0	241.0	393.5	577.2	804.2
2005 5/.....	6.6	14.0	30.2	73.8	146.3	262.5	420.7	597.9	861.9
2004 5/.....	5.7	13.1	28.1	70.1	138.9	247.7	406.1	578.0	852.5
2003 5/.....	6.5	12.8	30.9	75.1	146.5	262.0	419.8	596.6	903.6
2002.....	5.9	13.1	31.1	70.6	143.1	264.3	428.0	624.9	935.4
2001.....	5.9	12.5	33.4	72.5	147.9	270.1	429.6	611.8	951.5
2000.....	5.9	12.1	33.6	72.5	148.9	274.6	425.1	658.0	1,008.6
1999.....	5.2	12.1	36.1	79.7	162.0	293.9	450.4	673.4	1,033.5

Non-Hispanic White, female

2007 5/.....	6.2	13.5	25.9	59.9	116.5	203.0	308.5	415.9	561.9
2006 5/.....	6.8	13.3	27.2	61.6	122.3	204.5	300.9	405.8	548.5
2005 5/.....	6.8	12.5	28.7	65.8	123.0	214.1	320.6	430.5	595.3
2004 5/.....	5.4	11.6	27.1	64.2	119.4	205.1	301.8	402.5	544.8
2003 5/.....	5.4	11.9	28.4	67.7	127.9	209.9	315.7	423.8	569.5
2002.....	5.8	11.7	30.2	64.0	124.8	207.4	307.4	417.7	559.8
2001.....	4.8	12.1	30.3	67.4	129.1	209.0	303.2	405.8	562.7
2000.....	5.1	11.9	29.9	65.3	125.7	209.7	300.2	412.0	548.7
1999.....	4.8	12.6	32.5	67.2	126.8	208.0	298.8	417.7	532.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic Black, both sexes

2007 5/.....	9.5	17.2	26.9	48.4	86.4	129.8	206.3	265.1	342.9
2006 5/.....	7.8	17.5	28.2	52.9	85.2	125.1	203.8	266.7	337.4
2005 5/.....	10.8	18.4	31.2	53.3	93.6	145.2	223.6	275.5	369.2
2004 5/.....	9.9	16.8	28.6	53.6	83.1	130.2	208.4	251.4	335.9
2003 5/.....	9.4	17.1	32.0	53.7	88.4	149.3	214.6	274.0	354.9
2002.....	9.3	17.6	31.4	54.8	87.4	149.7	217.7	300.6	370.8
2001.....	9.3	16.8	32.3	52.7	95.9	150.7	208.7	279.2	379.3
2000.....	11.3	18.5	33.8	58.1	98.7	152.8	213.1	273.6	376.2
1999.....	11.5	17.5	36.9	58.6	101.6	164.5	245.5	297.3	377.0

Non-Hispanic Black, male

2007 5/.....	9.0	17.3	31.2	66.8	114.6	177.5	310.9	416.8	522.7
2006 5/.....	8.0	20.3	34.0	71.5	114.6	172.5	294.2	422.8	522.3
2005 5/.....	11.3	20.6	38.3	66.9	128.6	201.9	349.5	431.0	596.6
2004 5/.....	10.2	19.6	38.6	69.5	110.7	184.9	306.9	398.0	569.0
2003 5/.....	10.0	18.2	38.9	68.7	114.4	216.0	331.7	475.4	608.4
2002.....	10.0	20.3	38.4	72.7	118.0	216.2	345.1	479.7	653.1
2001.....	9.1	19.0	37.8	66.7	129.2	224.1	333.2	475.9	682.6
2000.....	11.8	19.6	40.3	74.2	140.2	228.6	328.0	451.4	696.3
1999.....	12.3	20.0	44.3	80.5	141.6	244.1	391.5	524.3	707.2

Non-Hispanic Black, female

2007 5/.....	10.0	17.1	23.4	34.0	65.8	96.9	143.4	185.4	268.4
2006 5/.....	7.7	15.1	23.4	38.5	63.6	92.5	149.8	184.4	262.0
2005 5/.....	10.3	16.4	25.2	42.7	67.7	106.4	148.7	193.5	277.7
2004 5/.....	9.6	14.5	20.4	41.2	62.5	93.1	150.2	174.2	243.8
2003 5/.....	8.8	16.2	26.4	42.1	69.1	104.3	145.1	168.6	256.7
2002.....	8.7	15.4	25.7	40.7	64.7	105.1	141.6	207.4	263.0
2001.....	9.5	14.9	27.8	41.7	71.4	101.9	134.4	179.1	266.9
2000.....	10.8	17.4	28.4	45.5	68.0	102.5	143.7	184.1	257.3
1999.....	10.9	15.3	31.0	41.5	72.1	111.7	157.5	184.8	253.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

All origins 3/, both sexes

2007.....	*	*	0.1	0.2	0.4	0.6	0.8	2.1	4.8
2006.....	*	*	0.1	0.2	0.3	0.6	1.1	2.1	6.0
2005.....	*	*	0.2	0.3	0.5	0.9	1.1	2.2	7.4
2004.....	*	0.1	0.2	0.3	0.4	0.7	1.2	2.0	6.5
2003.....	*	*	0.1	0.2	0.4	0.7	1.4	2.6	7.3
2002.....	*	0.1	0.1	0.3	0.5	1.0	1.8	2.9	8.1
2001.....	0.1	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000.....	*	*	0.1	0.4	0.7	1.0	2.1	3.6	11.8
1999.....	*	0.1	0.3	0.6	0.7	1.2	1.9	4.1	10.7

All origins 3/, male

2007.....	*	*	*	*	*	*	1.1	2.3	5.0
2006.....	*	*	*	*	*	0.6	1.1	2.2	5.9
2005.....	*	*	*	0.4	*	0.9	1.4	2.5	7.3
2004.....	*	*	*	*	0.5	0.8	1.2	2.2	7.0
2003.....	*	*	*	*	*	0.8	1.9	2.6	7.0
2002.....	*	*	*	*	0.7	1.1	2.2	2.9	8.8
2001.....	*	*	*	0.4	0.8	1.1	2.2	2.5	9.2
2000.....	*	*	*	0.4	0.9	1.0	2.8	4.9	12.9
1999.....	*	*	0.4	0.5	0.8	1.5	2.4	4.7	11.9

All origins 3/, female

2007.....	*	*	*	*	0.4	0.7	0.6	2.0	4.8
2006.....	*	*	*	*	*	0.6	1.1	2.0	6.0
2005.....	*	*	0.2	*	0.5	0.8	0.8	2.0	7.4
2004.....	*	*	*	*	*	0.6	1.2	1.9	6.3
2003.....	*	*	*	*	0.4	0.7	1.1	2.5	7.5
2002.....	*	*	*	*	*	0.9	1.4	2.9	7.8
2001.....	*	*	*	*	0.4	0.7	1.4	2.6	9.1
2000.....	*	*	*	0.4	0.5	1.1	1.6	2.8	11.4
1999.....	*	*	*	0.6	0.5	1.0	1.5	3.8	10.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic 4/, both sexes

2007	*	*	0.1	0.2	0.4	0.6	0.9	2.1	4.8
2006	*	*	0.1	0.2	0.4	0.7	1.1	2.1	6.0
2005	*	*	0.2	0.3	0.5	0.9	1.1	2.2	7.3
2004	*	0.1	0.2	0.3	0.4	0.7	1.2	1.9	6.6
2003	*	*	0.1	0.2	0.4	0.8	1.5	2.5	7.3
2002	*	*	0.2	0.3	0.5	1.0	1.7	2.9	8.0
2001	*	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000	*	*	*	0.5	0.7	1.0	2.1	3.5	11.8
1999	*	0.2	0.3	0.6	0.7	1.3	1.9	4.1	10.7

Non-Hispanic 4/, male

2007	*	*	*	*	*	*	1.2	2.1	4.9
2006	*	*	*	*	*	0.7	1.1	2.2	6.1
2005	*	*	*	0.4	*	1.0	1.5	2.5	7.4
2004	*	*	*	*	0.6	0.8	1.2	2.1	7.0
2003	*	*	*	*	*	0.8	1.9	2.6	6.9
2002	*	*	*	*	0.7	1.2	2.3	2.9	8.8
2001	*	*	*	0.4	0.8	1.1	2.1	2.6	9.3
2000	*	*	*	0.5	1.0	0.9	2.9	4.8	12.8
1999	*	*	0.4	0.6	0.8	1.5	2.5	4.7	11.9

Non-Hispanic 4/, female

2007	*	*	*	*	0.4	0.7	0.6	2.1	4.8
2006	*	*	*	*	*	0.7	1.2	2.0	6.0
2005	*	*	0.2	*	0.6	0.8	0.8	2.0	7.2
2004	*	*	*	*	*	0.5	1.2	1.9	6.4
2003	*	*	*	*	0.5	0.8	1.1	2.5	7.5
2002	*	*	*	*	*	0.9	1.4	2.9	7.6
2001	*	*	*	*	0.4	0.8	1.4	2.6	9.1
2000	*	*	*	0.5	0.5	1.1	1.7	2.8	11.4
1999	*	*	*	0.6	0.6	1.1	1.5	3.7	10.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic White, both sexes

2007 5/.....	*	*	0.2	*	0.4	0.6	1.0	2.2	4.9
2006 5/.....	*	*	*	0.2	0.4	0.7	1.2	2.1	6.3
2005 5/.....	*	*	0.2	0.3	0.5	0.9	1.2	2.2	7.8
2004 5/.....	*	*	0.2	0.3	0.5	0.7	1.2	2.1	6.7
2003 5/.....	*	*	*	0.2	0.5	0.8	1.6	2.5	7.5
2002.....	*	*	*	0.3	0.6	1.1	1.9	2.9	8.4
2001.....	*	0.1	0.2	0.4	0.6	0.9	1.7	2.7	9.7
2000.....	*	*	*	0.5	0.7	1.0	2.2	3.6	12.6
1999.....	*	*	0.3	0.6	0.7	1.3	1.9	4.1	11.2

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	1.3	2.2	5.1
2006 5/.....	*	*	*	*	*	0.7	1.0	2.2	6.5
2005 5/.....	*	*	*	0.4	*	1.0	1.5	2.4	7.5
2004 5/.....	*	*	*	*	0.6	0.9	1.2	2.3	7.2
2003 5/.....	*	*	*	*	*	0.7	2.0	2.5	7.1
2002.....	*	*	*	*	0.8	1.2	2.4	2.9	9.3
2001.....	*	*	*	*	0.9	1.1	2.1	2.6	9.6
2000.....	*	*	*	0.5	1.0	0.9	2.9	5.1	13.8
1999.....	*	*	0.4	0.6	0.9	1.5	2.5	4.5	12.4

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	0.7	0.7	2.2	4.9
2006 5/.....	*	*	*	*	*	0.7	1.2	2.0	6.2
2005 5/.....	*	*	*	*	0.6	0.9	0.9	2.0	7.9
2004 5/.....	*	*	*	*	*	0.6	1.2	1.9	6.5
2003 5/.....	*	*	*	*	0.5	0.8	1.2	2.5	7.6
2002.....	*	*	*	*	*	1.0	1.5	3.0	8.0
2001.....	*	*	*	*	*	0.7	1.5	2.8	9.7
2000.....	*	*	*	0.5	0.5	1.2	1.7	2.8	12.2
1999.....	*	*	*	0.7	0.6	1.1	1.5	3.8	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

All origins 3/, both sexes

2007.....	0.7	1.5	3.3	6.9	13.0	22.1	33.1	42.4	45.5
2006.....	0.7	1.4	3.2	7.1	13.2	22.3	33.1	41.6	45.3
2005.....	0.8	1.4	3.7	8.0	15.3	25.8	37.8	46.0	51.9
2004.....	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
2003.....	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1

All origins 3/, male

2007.....	0.9	1.9	3.9	8.1	15.3	26.4	40.9	53.8	61.6
2006.....	0.7	1.9	3.5	7.9	15.1	25.2	40.3	52.8	63.7
2005.....	1.1	1.6	4.4	9.7	18.4	30.7	47.5	59.5	71.5
2004.....	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
2003.....	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4

All origins 3/, female

2007.....	0.5	1.1	2.7	5.7	11.0	18.5	27.3	35.1	37.9
2006.....	0.7	1.0	2.8	6.4	11.5	19.9	27.8	34.6	36.7
2005.....	0.5	1.1	3.0	6.5	12.6	21.9	30.8	37.6	42.8
2004.....	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
2003.....	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

Hispanic, both sexes

2007	*	*	*	*	3.3	7.4	8.1	22.0	21.5
2006	*	*	*	2.2	3.5	4.7	13.2	19.3	20.7
2005	*	*	*	*	3.9	7.5	14.5	19.9	24.8
2004	*	*	*	*	4.2	6.0	13.6	19.9	30.0
2003	*	*	*	*	3.5	9.1	13.6	30.3	34.7
2002	*	*	*	2.4	3.8	10.3	10.9	32.1	36.6
2001	*	*	*	*	5.2	8.3	17.3	25.8	39.4
2000	*	*	*	*	4.3	8.2	18.4	29.5	31.2
1999	*	*	*	*	4.8	10.1	18.1	33.5	44.2

Hispanic, male

2007	*	*	*	*	*	12.1	11.4	33.3	32.5
2006	*	*	*	*	*	*	21.1	22.9	31.6
2005	*	*	*	*	6.8	11.1	20.5	26.3	38.6
2004	*	*	*	*	*	9.1	15.6	28.3	47.6
2003	*	*	*	*	*	13.2	21.9	41.6	53.9
2002	*	*	*	*	6.8	12.3	17.3	43.5	53.5
2001	*	*	*	*	*	12.4	28.0	45.4	60.9
2000	*	*	*	*	*	12.6	25.8	45.6	58.4
1999	*	*	*	*	*	16.8	26.5	55.0	77.6

Hispanic, female

2007	*	*	*	*	*	*	*	14.5	15.4
2006	*	*	*	*	*	*	7.5	16.9	15.0
2005	*	*	*	*	*	*	10.1	15.7	17.6
2004	*	*	*	*	*	*	12.3	14.3	20.9
2003	*	*	*	*	*	*	*	23.0	25.0
2002	*	*	*	*	*	8.8	*	24.8	28.2
2001	*	*	*	*	*	*	9.7	*	28.9
2000	*	*	*	*	*	*	13.1	19.7	*
1999	*	*	*	*	*	*	12.1	20.6	28.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

Non-Hispanic 4/, both sexes

2007.....	0.8	1.6	3.5	7.3	13.7	23.2	34.8	43.5	46.7
2006.....	0.8	1.5	3.3	7.5	13.9	23.6	34.4	42.8	46.5
2005.....	0.8	1.4	3.9	8.5	16.2	27.1	39.3	47.3	53.1
2004.....	0.6	1.6	3.6	8.6	16.6	27.5	39.0	46.9	50.7
2003.....	0.8	1.6	4.2	9.8	17.5	29.7	43.4	52.6	55.4
2002.....	0.8	1.7	4.6	9.6	18.3	32.6	44.8	55.6	60.1
2001.....	0.7	1.8	5.3	10.7	20.8	34.6	48.2	56.6	62.8
2000.....	0.8	1.9	5.4	11.0	21.8	36.9	48.7	61.5	64.3
1999.....	0.7	2.0	5.9	12.8	24.7	39.9	52.6	66.3	62.5

Non-Hispanic 4/, male

2007.....	1.0	2.1	4.1	8.5	16.2	27.5	42.9	55.0	63.2
2006.....	0.8	2.0	3.6	8.2	15.9	26.5	41.6	54.5	65.3
2005.....	1.2	1.7	4.6	10.3	19.3	31.9	49.1	61.2	73.1
2004.....	0.9	1.8	4.0	10.1	19.3	33.1	48.4	62.1	75.5
2003.....	1.0	2.0	4.7	10.8	19.3	35.7	53.4	68.6	80.4
2002.....	1.0	2.0	4.9	11.3	20.6	39.3	58.0	76.3	91.3
2001.....	0.9	2.1	6.0	11.4	23.4	41.8	60.6	76.0	102.7
2000.....	1.1	2.2	6.0	11.7	25.6	44.0	61.9	85.1	105.9
1999.....	0.9	2.3	6.6	14.2	29.2	50.5	68.9	92.7	106.1

Non-Hispanic 4/, female

2007.....	0.6	1.2	2.9	6.1	11.6	19.7	28.8	36.3	39.0
2006.....	0.7	1.1	3.0	6.8	12.3	21.2	29.2	35.5	37.8
2005.....	0.5	1.2	3.1	6.9	13.5	23.1	32.1	38.7	44.0
2004.....	0.4	1.4	3.3	7.3	14.3	23.0	32.2	37.6	39.6
2003.....	0.5	1.3	3.8	8.9	16.0	24.9	36.3	42.8	44.5
2002.....	0.6	1.4	4.3	8.2	16.4	27.2	35.5	43.1	46.6
2001.....	0.5	1.5	4.5	10.1	18.6	28.9	39.6	44.9	46.3
2000.....	0.6	1.6	4.8	10.3	18.5	31.3	39.5	47.7	47.5
1999.....	0.5	1.8	5.2	11.5	20.8	31.6	41.3	50.9	45.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

Non-Hispanic White, both sexes

2007 5/.....	0.8	1.7	3.8	7.8	14.9	25.3	37.0	46.3	49.5
2006 5/.....	0.8	1.7	3.6	8.0	15.1	25.9	36.7	45.3	49.2
2005 5/.....	0.9	1.4	4.1	9.2	17.4	29.5	41.6	49.8	55.9
2004 5/.....	0.7	1.6	3.8	9.2	18.2	29.9	41.7	49.8	53.4
2003 5/.....	0.8	1.8	4.5	10.5	19.2	31.9	46.0	55.2	57.9
2002.....	0.8	1.8	4.9	10.4	19.8	34.8	47.7	58.4	63.1
2001.....	0.8	1.8	5.5	11.6	22.7	36.9	51.4	59.9	65.9
2000.....	0.8	1.9	5.6	11.9	23.2	39.5	51.6	64.9	66.9
1999.....	0.7	2.1	6.3	13.8	26.8	42.6	55.3	69.9	65.2

Non-Hispanic White, male

2007 5/.....	1.0	2.1	4.3	8.7	17.0	29.5	44.3	57.2	65.6
2006 5/.....	0.8	2.0	3.8	8.3	16.6	28.6	43.1	56.5	68.4
2005 5/.....	1.3	1.6	4.7	10.7	20.2	33.9	51.1	62.5	75.5
2004 5/.....	0.8	1.7	4.0	10.5	20.6	35.1	50.6	65.0	78.2
2003 5/.....	1.0	2.0	4.8	11.2	20.7	37.4	54.9	70.8	81.6
2002.....	1.0	2.0	5.1	11.7	21.2	41.1	60.3	78.9	94.3
2001.....	1.0	2.0	6.1	12.0	25.1	43.3	63.4	79.2	106.6
2000.....	0.9	2.1	6.2	12.3	26.3	46.3	64.8	88.6	107.3
1999.....	0.8	2.1	6.8	14.9	30.8	52.4	71.4	95.7	110.0

Non-Hispanic White, female

2007 5/.....	0.6	1.3	3.3	6.9	12.9	21.8	31.4	39.3	41.9
2006 5/.....	0.8	1.3	3.4	7.6	13.6	23.6	31.9	38.2	40.3
2005 5/.....	0.5	1.2	3.5	7.8	14.9	25.8	34.5	41.8	47.0
2004 5/.....	0.5	1.4	3.7	8.1	16.1	25.6	35.3	40.4	42.3
2003 5/.....	0.5	1.5	4.2	9.8	17.8	27.4	39.6	45.7	47.4
2002.....	0.6	1.6	4.7	9.2	18.6	29.6	38.8	45.9	49.7
2001.....	0.5	1.5	4.9	11.3	20.6	31.7	42.9	48.3	48.9
2000.....	0.6	1.6	5.1	11.5	20.5	34.1	42.3	50.8	50.6
1999.....	0.5	2.0	5.7	12.9	23.2	34.7	43.9	54.8	47.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

Non-Hispanic Black, both sexes

2007 5/.....	1.1	1.6	2.7	5.8	9.6	12.2	22.6	23.9	24.3
2006 5/.....	*	1.3	2.6	6.7	9.1	11.9	21.0	25.0	24.2
2005 5/.....	1.0	1.9	3.6	6.1	12.0	14.0	26.7	26.6	27.3
2004 5/.....	*	2.2	3.2	6.7	8.5	14.1	21.2	24.6	27.9
2003 5/.....	1.1	1.3	3.7	7.4	10.5	16.7	25.4	31.1	31.2
2002.....	1.1	1.5	3.4	6.1	12.1	21.4	23.7	35.0	32.1
2001.....	*	2.4	5.0	6.3	11.7	20.4	21.8	26.5	35.9
2000.....	1.6	2.6	4.8	7.8	15.3	19.2	24.5	31.0	37.5
1999.....	1.4	2.5	4.7	8.2	14.2	21.1	31.8	31.7	37.8

Non-Hispanic Black, male

2007 5/.....	*	2.3	3.8	9.4	14.9	18.7	37.4	38.8	47.1
2006 5/.....	*	2.4	3.9	11.1	13.8	14.9	33.1	42.1	40.3
2005 5/.....	*	2.9	5.8	9.9	18.4	20.8	40.4	46.8	50.4
2004 5/.....	*	3.1	5.0	10.0	14.0	21.9	35.4	39.7	57.9
2003 5/.....	*	*	5.4	10.5	13.7	25.3	43.9	51.3	71.4
2002.....	*	2.4	4.6	9.6	20.7	31.5	42.3	61.6	65.6
2001.....	*	3.0	6.9	9.2	15.9	32.4	34.6	49.9	69.3
2000.....	2.5	3.6	6.1	10.8	25.9	29.4	34.7	53.2	89.2
1999.....	2.0	4.2	6.6	12.2	23.1	37.4	49.8	66.4	73.7

Non-Hispanic Black, female

2007 5/.....	*	*	*	2.9	5.7	7.7	13.7	16.0	14.8
2006 5/.....	*	*	*	3.3	5.7	9.8	13.7	15.9	17.6
2005 5/.....	*	*	*	3.1	7.3	9.3	18.6	15.9	18.0
2004 5/.....	*	*	*	4.0	4.4	8.8	12.7	16.6	16.1
2003 5/.....	*	*	2.4	4.9	8.1	10.9	14.4	20.5	15.7
2002.....	*	*	*	3.3	5.7	14.7	12.6	21.1	19.2
2001.....	*	*	3.4	4.1	8.7	12.4	14.1	14.6	23.5
2000.....	*	*	3.7	5.4	7.5	12.5	18.3	19.9	18.3
1999.....	*	*	3.1	5.1	7.7	10.3	21.0	14.5	24.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

All origins 3/, both sexes

2007.....	1.1	1.3	1.3	1.5	2.0	2.3	3.5	5.5	12.0
2006.....	1.2	1.4	1.4	1.8	2.0	2.7	3.6	6.4	12.3
2005.....	1.4	1.4	1.5	1.9	2.3	2.9	4.1	7.1	14.5
2004.....	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
2003.....	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3

All origins 3/, male

2007.....	0.9	1.0	1.2	1.3	1.5	1.6	2.4	4.4	7.1
2006.....	0.9	1.1	1.0	1.2	1.4	2.1	2.9	4.4	9.1
2005.....	0.9	1.2	1.1	1.2	1.7	1.9	3.7	4.8	8.7
2004.....	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
2003.....	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7

All origins 3/, female

2007.....	1.4	1.6	1.5	1.7	2.4	3.0	4.2	6.3	14.2
2006.....	1.6	1.7	1.7	2.3	2.6	3.2	4.2	7.6	13.9
2005.....	1.8	1.7	1.9	2.6	2.8	3.7	4.4	8.6	17.2
2004.....	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
2003.....	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

Hispanic, both sexes

2007	*	*	*	*	*	*	5.0	*	9.7
2006	*	*	*	2.2	*	*	*	*	16.4
2005	*	*	1.4	*	*	3.6	*	*	13.7
2004	0.9	1.5	1.8	*	2.8	4.0	5.0	*	11.4
2003	1.6	1.6	2.4	*	4.1	*	*	9.0	14.5
2002	1.2	*	*	2.9	3.8	3.8	5.9	8.9	12.2
2001	1.0	2.0	2.5	*	*	4.8	7.4	*	13.8
2000	*	1.6	*	4.4	5.7	5.4	7.7	*	16.6
1999	1.5	1.8	2.5	*	5.5	6.1	8.7	13.5	17.5

Hispanic, male

2007	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*

Hispanic, female

2007	*	*	*	*	*	*	*	*	11.0
2006	*	*	*	*	*	*	*	*	19.8
2005	*	*	*	*	*	*	*	*	13.0
2004	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	5.6	*	*	*	17.2
2002	*	*	*	*	5.8	*	*	*	*
2001	*	2.8	*	*	*	*	*	*	18.6
2000	*	*	*	*	6.0	*	*	*	*
1999	*	*	*	*	6.9	*	11.0	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

Non-Hispanic 4/, both sexes

2007.....	1.2	1.4	1.4	1.5	2.0	2.4	3.3	5.6	12.1
2006.....	1.3	1.5	1.4	1.8	2.0	2.7	3.6	6.4	12.1
2005.....	1.5	1.5	1.5	2.0	2.2	2.8	4.1	7.2	14.5
2004.....	1.4	1.4	1.7	1.9	2.5	2.9	4.1	6.6	14.2
2003.....	1.4	1.6	1.8	2.2	2.6	3.2	4.9	7.1	14.1
2002.....	1.4	1.7	2.0	2.1	2.6	3.7	5.4	7.7	14.7
2001.....	1.4	1.7	1.8	2.6	3.0	3.9	5.0	8.1	16.1
2000.....	1.6	1.7	2.0	2.3	3.3	4.5	5.5	8.7	16.6
1999.....	1.6	1.7	2.6	2.2	3.6	4.6	6.1	9.5	17.3

Non-Hispanic 4/, male

2007.....	0.9	1.1	1.2	1.3	1.5	1.5	2.3	4.5	7.1
2006.....	0.9	1.2	1.0	1.2	1.4	2.1	2.9	4.4	9.0
2005.....	0.9	1.2	1.1	1.2	1.6	1.9	3.6	4.8	8.4
2004.....	1.2	0.9	1.4	1.3	1.9	2.3	2.8	4.5	11.1
2003.....	1.2	1.1	1.4	1.7	1.7	2.4	3.8	5.5	10.7
2002.....	1.0	1.3	1.5	1.6	2.0	2.8	4.1	6.2	11.1
2001.....	1.0	1.3	1.1	1.9	2.3	2.9	3.7	5.6	12.2
2000.....	1.3	1.2	1.6	1.6	2.6	3.2	3.8	6.7	12.7
1999.....	1.0	1.2	1.9	1.8	2.5	3.8	4.8	7.1	12.6

Non-Hispanic 4/, female

2007.....	1.5	1.7	1.5	1.7	2.4	3.1	4.1	6.3	14.4
2006.....	1.7	1.7	1.8	2.3	2.5	3.2	4.1	7.6	13.6
2005.....	2.0	1.7	1.9	2.6	2.8	3.7	4.5	8.6	17.3
2004.....	1.6	1.8	1.9	2.5	2.9	3.5	5.0	7.8	15.5
2003.....	1.7	2.1	2.2	2.7	3.3	3.8	5.7	8.1	15.6
2002.....	1.8	2.0	2.4	2.5	3.1	4.5	6.2	8.5	16.3
2001.....	1.7	2.1	2.5	3.2	3.6	4.7	6.0	9.6	17.7
2000.....	1.9	2.1	2.4	3.0	3.9	5.5	6.7	9.8	18.2
1999.....	2.2	2.3	3.3	2.7	4.5	5.3	7.0	10.9	19.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

Non-Hispanic White, both sexes

2007 5/.....	0.8	1.0	0.9	1.2	1.6	2.0	3.1	5.1	11.9
2006 5/.....	0.9	1.1	1.0	1.3	1.6	2.3	3.4	6.0	11.8
2005 5/.....	1.0	1.1	1.1	1.6	1.8	2.1	3.8	6.4	14.5
2004 5/.....	0.8	1.0	1.2	1.4	2.0	2.6	3.5	6.2	13.8
2003 5/.....	1.0	1.1	1.4	1.7	2.1	2.6	4.4	6.6	14.1
2002.....	1.0	1.1	1.4	1.7	2.0	3.1	5.1	7.4	14.7
2001.....	0.9	1.3	1.3	1.8	2.6	3.3	4.5	7.7	15.7
2000.....	1.0	1.2	1.4	1.8	2.5	3.9	5.0	8.4	15.8
1999.....	1.1	1.2	2.0	1.7	2.9	3.9	5.6	9.2	16.5

Non-Hispanic White, male

2007 5/.....	0.6	0.8	0.8	0.9	1.1	1.2	1.9	3.8	6.6
2006 5/.....	0.6	0.7	0.6	0.8	1.0	1.6	2.8	3.9	9.0
2005 5/.....	0.6	1.0	0.8	1.0	1.2	1.3	3.1	4.1	8.2
2004 5/.....	0.6	0.7	1.0	0.8	1.5	1.9	2.1	3.9	10.4
2003 5/.....	0.8	0.7	1.0	1.2	1.5	1.9	3.0	5.1	10.6
2002.....	0.7	0.9	1.0	1.2	1.5	2.3	3.6	5.8	10.8
2001.....	0.7	0.9	0.8	1.2	1.9	2.3	3.2	5.0	11.4
2000.....	0.9	0.9	1.0	1.1	1.8	2.9	3.5	6.3	11.1
1999.....	0.6	0.8	1.4	1.3	1.9	2.9	4.2	6.9	10.6

Non-Hispanic White, female

2007 5/.....	1.0	1.1	1.1	1.4	2.1	2.7	4.0	5.8	14.4
2006 5/.....	1.3	1.4	1.4	1.8	2.1	2.8	3.8	7.4	13.2
2005 5/.....	1.4	1.2	1.4	2.1	2.2	2.8	4.3	7.8	17.4
2004 5/.....	1.0	1.3	1.5	2.0	2.5	3.3	4.5	7.6	15.3
2003 5/.....	1.3	1.4	1.7	2.3	2.8	3.1	5.4	7.5	15.7
2002.....	1.4	1.3	1.7	2.0	2.5	3.8	6.1	8.4	16.4
2001.....	1.2	1.7	1.8	2.4	3.2	4.1	5.5	9.3	17.5
2000.....	1.2	1.4	1.7	2.3	3.2	4.8	6.1	9.7	17.7
1999.....	1.6	1.6	2.5	2.1	3.8	4.6	6.7	10.5	18.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

Non-Hispanic Black, both sexes

2007 5/.....	3.8	4.2	4.5	4.3	4.5	6.0	5.1	8.4	11.8
2006 5/.....	3.7	4.4	4.5	6.2	4.9	6.3	5.2	7.5	12.7
2005 5/.....	4.5	4.0	5.5	5.3	5.8	9.0	7.9	12.9	12.0
2004 5/.....	4.9	4.3	5.3	6.4	5.8	5.8	9.2	9.6	15.3
2003 5/.....	4.2	5.8	5.3	5.1	6.8	7.7	8.4	10.6	11.9
2002.....	4.2	5.8	7.3	5.7	6.2	10.3	7.2	9.1	11.8
2001.....	4.4	5.0	6.1	7.9	6.3	9.5	10.3	11.3	16.5
2000.....	5.6	5.3	6.7	7.2	9.1	9.8	10.9	9.8	22.3
1999.....	5.0	5.9	7.5	6.3	8.6	11.2	10.0	11.4	20.7

Non-Hispanic Black, male

2007 5/.....	2.8	3.3	4.2	4.9	5.0	*	*	*	*
2006 5/.....	2.9	4.3	4.1	5.7	*	*	*	*	*
2005 5/.....	3.4	2.6	4.4	*	*	6.5	9.6	*	*
2004 5/.....	4.9	3.0	4.9	5.8	5.0	6.7	*	*	*
2003 5/.....	3.5	4.6	3.9	5.0	5.1	*	9.3	*	*
2002.....	3.4	4.7	6.1	5.5	6.4	8.9	*	*	*
2001.....	3.8	4.6	3.7	6.1	*	9.6	*	*	*
2000.....	4.1	3.2	6.3	6.0	8.8	6.8	*	*	24.7
1999.....	3.7	4.3	5.2	5.9	6.9	12.1	9.7	*	*

Non-Hispanic Black, female

2007 5/.....	4.7	5.0	4.7	3.9	4.2	6.9	*	8.4	14.1
2006 5/.....	4.4	4.5	4.8	6.7	5.5	6.7	6.0	8.5	15.3
2005 5/.....	5.5	5.2	6.5	6.7	6.9	10.7	6.8	13.3	13.8
2004 5/.....	5.0	5.3	5.6	6.9	6.5	5.2	9.9	9.3	16.9
2003 5/.....	4.8	6.9	6.4	5.3	8.1	8.9	7.8	11.8	13.6
2002.....	4.8	6.8	8.2	5.9	6.1	11.2	7.8	9.7	12.5
2001.....	5.0	5.3	8.0	9.4	7.5	9.4	11.6	12.1	17.1
2000.....	6.9	7.1	7.1	8.2	9.2	11.8	14.5	9.9	21.4
1999.....	6.0	7.2	9.4	6.7	9.8	10.6	10.2	14.0	21.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

All origins 3/, both sexes

2007.....	4.2	10.2	20.7	47.7	96.1	169.1	273.4	393.6	533.8
2006.....	3.8	9.8	21.1	48.3	96.3	169.1	270.7	384.8	525.4
2005.....	4.3	9.7	21.5	51.8	100.6	180.7	287.5	402.5	563.5
2004.....	3.6	8.8	20.2	49.7	95.1	170.8	272.9	380.4	531.1
2003.....	3.6	8.6	21.5	52.2	101.4	178.5	281.8	394.6	558.9
2002.....	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001.....	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000.....	3.4	8.2	22.4	49.1	99.2	177.1	272.5	400.7	556.7
1999.....	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8

All origins 3/, male

2007.....	4.3	10.8	22.8	51.9	106.5	190.7	319.4	489.7	684.9
2006.....	3.9	10.3	22.8	52.7	103.1	190.1	320.3	481.9	682.2
2005.....	4.3	10.7	22.8	56.3	112.4	206.5	341.0	497.9	732.4
2004.....	3.6	9.8	21.5	53.5	105.2	193.5	328.2	476.3	718.6
2003.....	4.1	9.1	23.3	57.0	112.2	206.5	338.3	493.3	769.0
2002.....	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001.....	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000.....	3.8	8.7	24.4	54.5	111.6	210.9	337.1	533.3	838.0
1999.....	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7

All origins 3/, female

2007.....	4.1	9.6	18.7	43.8	86.9	151.4	239.5	333.0	461.9
2006.....	3.8	9.3	19.6	44.3	90.3	152.0	234.3	324.2	452.1
2005.....	4.3	8.8	20.4	47.7	90.2	159.8	248.7	343.2	485.9
2004.....	3.6	7.8	18.9	46.2	86.2	152.5	233.1	321.3	446.8
2003.....	3.2	8.0	19.8	47.8	92.1	156.0	241.5	334.2	466.0
2002.....	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001.....	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000.....	2.9	7.8	20.5	44.3	88.7	150.4	227.6	322.5	442.2
1999.....	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

Hispanic, both sexes

2007	1.5	2.7	5.7	12.6	28.2	50.6	112.2	194.5	321.1
2006	1.1	2.4	5.9	13.3	26.0	55.8	103.6	175.4	331.8
2005	1.5	3.7	6.0	15.0	27.0	61.5	117.4	197.6	367.9
2004	1.0	2.7	6.6	13.1	28.8	61.8	104.8	183.5	345.4
2003	1.0	2.6	6.0	13.4	29.3	61.9	111.2	195.9	403.7
2002	1.1	3.3	6.9	15.1	30.4	59.0	117.3	198.7	397.4
2001	1.1	3.0	5.9	15.5	33.6	59.6	118.4	176.8	418.9
2000	1.2	2.2	6.6	14.1	29.7	54.5	116.0	209.4	422.7
1999	1.3	2.6	7.2	15.3	34.6	68.0	127.8	213.2	462.3

Hispanic, male

2007	1.7	3.5	6.5	15.5	34.0	73.7	153.2	251.8	416.7
2006	*	2.7	6.9	16.4	32.7	71.0	131.9	230.9	439.8
2005	*	4.5	6.9	22.1	33.1	81.0	158.8	281.9	468.2
2004	*	2.8	7.2	17.2	38.0	87.8	146.9	236.9	455.3
2003	*	*	8.0	18.2	35.4	87.1	145.2	278.6	574.2
2002	*	3.7	9.9	19.8	43.0	77.2	175.3	268.8	535.4
2001	*	2.9	6.6	19.8	47.1	82.3	157.5	227.2	605.7
2000	*	*	7.0	22.2	38.8	79.7	159.5	307.7	584.5
1999	*	*	8.8	20.6	41.0	98.8	184.1	319.1	735.4

Hispanic, female

2007	*	1.9	5.0	10.0	23.2	32.4	82.5	156.2	269.2
2006	*	2.1	4.9	10.5	20.4	43.9	83.1	138.4	274.4
2005	1.8	2.9	5.1	8.7	21.9	46.3	87.7	141.7	315.4
2004	*	2.6	6.1	9.6	21.2	41.8	74.9	148.3	288.8
2003	*	3.2	4.1	9.2	24.3	42.6	87.1	142.1	317.2
2002	*	2.9	4.2	11.0	20.2	45.0	75.9	154.0	329.1
2001	*	3.2	5.2	11.8	22.7	42.2	90.5	145.6	327.7
2000	*	*	6.2	7.2	22.3	35.3	85.2	149.6	343.3
1999	*	*	5.8	10.8	29.4	44.8	88.1	150.0	330.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 4/, both sexes

2007.....	4.6	11.0	22.0	50.5	101.4	178.1	284.3	405.0	544.7
2006.....	4.2	10.5	22.4	51.1	101.7	177.3	281.4	396.1	534.6
2005.....	4.6	10.3	22.8	54.7	106.2	189.1	297.9	412.8	572.1
2004.....	3.9	9.3	21.3	52.5	99.9	177.9	282.5	389.7	538.8
2003.....	3.9	9.1	22.7	55.0	106.5	185.9	290.8	403.0	564.3
2002.....	3.8	9.1	22.9	51.8	103.1	182.4	287.5	408.2	560.8
2001.....	3.5	8.8	23.6	52.3	105.2	184.0	282.5	394.4	561.3
2000.....	3.5	8.7	23.5	51.5	103.5	183.6	279.1	407.0	560.5
1999.....	3.3	8.6	24.7	54.0	106.6	188.8	286.7	411.5	557.6

Non-Hispanic 4/, male

2007.....	4.7	11.5	24.2	54.8	112.1	199.3	330.4	503.6	700.0
2006.....	4.2	11.1	24.1	55.4	108.3	198.3	332.2	495.9	694.8
2005.....	4.7	11.3	24.1	58.8	118.3	215.0	351.8	509.2	745.0
2004.....	3.9	10.4	22.7	56.2	109.9	200.0	338.5	487.8	730.9
2003.....	4.4	9.8	24.4	59.6	117.3	213.8	348.5	502.4	776.7
2002.....	4.0	10.0	24.3	55.6	113.4	211.5	350.7	521.4	794.9
2001.....	4.0	9.4	25.5	56.2	115.6	215.3	350.0	510.4	800.6
2000.....	4.1	9.2	25.7	56.4	115.7	217.5	344.3	540.5	846.6
1999.....	3.9	9.0	27.0	61.1	124.1	229.6	364.7	552.0	869.6

Non-Hispanic 4/, female

2007.....	4.4	10.5	19.9	46.6	92.1	160.7	250.3	342.9	471.4
2006.....	4.2	10.0	20.8	47.1	95.9	160.2	244.1	333.9	460.1
2005.....	4.5	9.4	21.7	50.9	95.6	168.1	258.7	353.1	493.2
2004.....	3.8	8.3	19.9	49.1	91.1	160.2	242.3	329.4	453.0
2003.....	3.4	8.4	21.0	50.8	97.1	163.4	249.8	342.3	471.0
2002.....	3.5	8.3	21.7	48.3	94.3	159.0	243.0	339.5	459.8
2001.....	3.0	8.2	21.9	48.7	96.2	159.0	235.3	325.1	461.7
2000.....	3.0	8.2	21.5	47.0	93.0	156.8	233.7	328.4	444.8
1999.....	2.7	8.2	22.5	47.5	91.6	156.9	232.5	329.4	432.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic White, both sexes

2007 5/.....	4.7	11.5	23.4	54.2	109.3	192.9	303.3	429.6	574.2
2006 5/.....	4.5	10.9	23.7	54.6	109.8	192.2	299.3	418.6	562.5
2005 5/.....	4.8	10.6	24.0	58.6	114.3	203.5	316.7	436.5	600.8
2004 5/.....	4.0	9.7	22.3	56.2	107.8	191.2	299.4	411.6	566.1
2003 5/.....	4.1	9.4	23.6	58.8	114.8	198.1	307.5	425.2	592.2
2002.....	3.9	9.4	24.2	54.9	111.0	194.0	303.0	427.6	586.9
2001.....	3.6	9.1	24.8	56.0	112.0	195.4	298.0	413.0	585.7
2000.....	3.6	8.9	24.5	54.6	110.1	194.1	292.9	426.8	585.5
1999.....	3.2	8.9	25.8	57.0	112.9	198.4	298.5	429.4	582.8

Non-Hispanic White, male

2007 5/.....	4.8	12.1	25.4	57.2	118.4	211.0	344.3	525.3	729.2
2006 5/.....	4.4	11.4	25.1	57.5	113.9	210.0	346.5	514.6	720.3
2005 5/.....	4.7	11.2	24.5	61.7	124.5	226.3	365.0	528.8	770.7
2004 5/.....	4.1	10.6	22.9	58.6	116.2	209.9	352.3	506.8	756.7
2003 5/.....	4.6	10.0	24.9	62.4	123.9	222.0	360.0	518.3	804.2
2002.....	4.1	10.1	24.9	57.4	119.7	219.6	361.6	537.3	821.0
2001.....	4.1	9.5	26.2	58.9	120.0	223.5	361.0	525.0	823.9
2000.....	4.0	9.0	26.3	58.6	119.8	224.5	353.8	558.0	876.5
1999.....	3.8	9.1	27.5	63.0	128.5	237.0	372.4	566.3	900.5

Non-Hispanic White, female

2007 5/.....	4.7	10.9	21.4	51.4	101.1	177.7	272.4	368.6	500.8
2006 5/.....	4.6	10.5	22.4	51.8	106.2	177.4	263.9	358.2	488.9
2005 5/.....	4.9	9.9	23.6	55.6	105.2	184.6	280.9	378.8	523.1
2004 5/.....	3.9	8.8	21.8	53.9	100.3	175.7	260.7	352.6	480.7
2003 5/.....	3.6	9.0	22.4	55.4	106.8	178.5	269.5	368.0	498.8
2002.....	3.8	8.8	23.6	52.6	103.3	173.0	261.1	360.4	485.7
2001.....	3.0	8.7	23.5	53.4	104.9	172.5	253.3	345.3	486.6
2000.....	3.2	8.7	22.9	51.0	101.5	169.7	250.1	348.8	468.2
1999.....	2.6	8.8	24.1	51.5	99.2	167.5	246.7	348.6	456.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic Black, both sexes

2007 5/.....	4.6	11.3	19.8	38.0	71.6	111.3	178.4	230.9	303.3
2006 5/.....	3.3	11.7	20.8	39.9	70.9	106.1	176.2	232.3	296.8
2005 5/.....	5.3	12.3	21.7	41.5	75.2	121.7	188.3	234.5	328.2
2004 5/.....	4.2	10.1	19.8	40.1	68.6	109.7	177.2	216.5	288.1
2003 5/.....	3.9	9.7	22.6	41.0	70.9	124.3	180.3	229.9	306.1
2002.....	3.9	10.2	20.5	42.8	68.7	117.7	186.1	254.1	322.7
2001.....	4.2	9.4	21.2	38.2	77.1	119.8	175.2	240.6	323.4
2000.....	4.0	10.3	22.2	42.9	73.5	122.6	175.9	229.9	312.6
1999.....	5.0	8.6	24.5	43.7	78.4	130.6	201.5	250.1	314.0

Non-Hispanic Black, male

2007 5/.....	4.9	11.7	23.3	52.5	94.0	153.5	267.6	366.8	466.0
2006 5/.....	3.9	13.6	25.5	54.6	96.4	151.0	255.0	372.9	473.7
2005 5/.....	6.5	15.0	27.8	53.0	105.1	173.6	297.6	369.8	534.8
2004 5/.....	4.0	12.8	28.6	53.1	91.5	155.6	262.5	348.1	495.9
2003 5/.....	4.6	11.4	29.4	53.0	95.7	184.0	277.1	412.8	526.3
2002.....	4.9	12.9	27.5	57.4	90.1	175.5	295.3	408.7	572.1
2001.....	4.5	11.3	27.2	51.0	107.6	180.8	288.7	414.6	592.5
2000.....	5.2	12.7	27.9	56.8	105.0	191.0	286.6	386.2	578.9
1999.....	6.2	11.0	32.1	62.0	111.3	192.5	329.2	442.8	610.9

Non-Hispanic Black, female

2007 5/.....	4.4	11.0	16.9	26.7	55.2	82.1	124.7	159.5	235.8
2006 5/.....	2.9	10.1	16.9	28.5	52.0	75.4	129.2	158.1	224.6
2005 5/.....	4.2	10.0	16.6	32.5	53.1	86.2	123.3	163.2	245.1
2004 5/.....	4.4	7.8	12.6	30.0	51.6	78.7	126.7	147.2	206.0
2003 5/.....	3.2	8.3	17.1	31.7	52.6	84.1	122.9	134.3	220.8
2002.....	3.1	7.9	14.8	31.3	52.8	78.9	120.8	173.7	227.4
2001.....	3.9	7.8	16.2	28.1	54.6	79.1	107.5	152.0	223.7
2000.....	2.9	8.3	17.5	31.9	50.3	77.1	109.2	151.2	213.7
1999.....	3.9	6.6	18.3	29.4	53.9	89.6	124.6	154.5	203.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

All origins 3/, both sexes

2007.....	0.8	1.2	2.2	3.4	5.8	12.7	25.6	58.1	153.3
2006.....	0.8	1.2	2.0	3.7	6.3	12.8	26.5	58.7	155.8
2005.....	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
2004.....	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6

All origins 3/, male

2007.....	1.0	1.6	2.9	4.2	7.5	17.2	36.2	81.7	222.0
2006.....	1.0	1.6	2.5	4.9	8.7	17.5	37.3	85.7	224.8
2005.....	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
2004.....	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
2003.....	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4

All origins 3/, female

2007.....	0.6	0.9	1.4	2.7	4.4	9.1	17.8	43.2	120.7
2006.....	0.7	0.9	1.5	2.6	4.3	8.9	18.6	41.9	123.5
2005.....	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
2004.....	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Hispanic, both sexes

2007	*	*	1.9	2.1	3.3	7.7	14.0	27.3	75.1
2006	*	*	*	*	3.5	8.5	18.6	37.0	77.5
2005	*	*	1.4	*	2.8	8.5	14.5	39.5	109.9
2004	*	*	*	2.2	4.5	6.0	18.2	39.0	101.3
2003	*	*	1.6	*	4.2	9.5	15.9	34.0	113.0
2002	*	*	*	2.7	3.7	7.3	19.5	45.0	131.4
2001	*	*	*	*	4.3	8.3	19.0	42.2	161.3
2000	*	*	*	*	3.5	9.0	20.8	40.7	137.4
1999	*	*	*	*	3.6	7.9	18.4	35.8	130.5

Hispanic, male

2007	*	*	*	*	*	10.3	21.8	33.3	100.6
2006	*	*	*	*	*	11.1	24.2	44.2	86.8
2005	*	*	*	*	*	9.1	17.8	55.2	154.4
2004	*	*	*	*	6.5	8.7	21.9	48.1	121.9
2003	*	*	*	*	*	12.0	20.7	52.0	123.1
2002	*	*	*	*	*	*	26.9	72.1	167.3
2001	*	*	*	*	*	11.4	25.2	57.1	220.9
2000	*	*	*	*	*	12.2	32.5	63.3	169.3
1999	*	*	*	*	*	*	26.5	48.7	159.6

Hispanic, female

2007	*	*	*	*	*	5.7	8.3	23.3	61.2
2006	*	*	*	*	*	6.5	14.5	32.2	72.5
2005	*	*	*	*	*	8.0	12.0	29.0	86.7
2004	*	*	*	*	*	*	15.5	33.0	90.8
2003	*	*	*	*	*	7.5	12.5	22.3	107.8
2002	*	*	*	*	*	6.8	14.2	27.7	113.6
2001	*	*	*	*	*	*	14.5	33.0	132.2
2000	*	*	*	*	*	*	12.5	26.9	121.7
1999	*	*	*	*	*	*	12.7	28.1	116.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Non-Hispanic 4/, both sexes

2007.....	0.9	1.3	2.2	3.5	6.0	13.1	26.4	59.9	157.4
2006.....	0.9	1.3	2.1	3.8	6.5	13.1	27.0	60.0	159.6
2005.....	0.8	1.5	2.1	3.9	6.6	13.1	29.3	62.1	166.3
2004.....	0.8	1.2	1.9	3.5	6.1	14.2	29.0	63.3	167.4
2003.....	0.9	1.1	2.0	3.9	6.4	13.5	30.8	66.8	178.5
2002.....	0.7	1.2	1.9	3.5	6.7	13.7	31.9	68.1	187.8
2001.....	0.7	1.3	1.8	3.7	6.3	14.4	30.9	68.8	190.0
2000.....	0.8	1.2	1.9	3.2	7.0	14.5	30.8	66.4	189.0
1999.....	0.7	1.0	2.1	3.1	6.2	13.4	28.5	61.4	176.8

Non-Hispanic 4/, male

2007.....	1.1	1.6	3.0	4.4	7.7	17.7	37.1	84.5	228.8
2006.....	1.0	1.7	2.6	5.1	9.0	18.0	38.1	88.1	232.3
2005.....	1.0	1.9	2.4	5.1	8.7	18.0	39.5	87.7	232.8
2004.....	1.0	1.6	2.4	4.5	8.2	19.3	39.9	89.3	242.7
2003.....	1.1	1.4	2.6	4.9	8.7	18.1	42.5	94.0	251.5
2002.....	0.9	1.6	2.2	4.2	8.8	18.7	44.2	95.8	272.1
2001.....	0.8	1.7	2.3	4.4	8.4	19.5	41.7	99.6	271.6
2000.....	1.0	1.5	2.3	4.0	9.6	20.6	42.6	96.0	279.7
1999.....	0.9	1.0	2.5	4.3	8.5	18.7	42.1	91.4	261.6

Non-Hispanic 4/, female

2007.....	0.6	0.9	1.4	2.7	4.6	9.3	18.5	44.4	123.6
2006.....	0.7	0.9	1.6	2.7	4.4	9.1	18.9	42.4	125.8
2005.....	0.7	1.2	1.7	2.8	4.8	9.2	21.9	46.3	136.0
2004.....	0.6	0.9	1.6	2.6	4.3	10.2	21.1	47.3	133.8
2003.....	0.7	0.8	1.5	3.0	4.4	9.8	22.5	50.2	146.5
2002.....	0.5	0.8	1.5	2.9	4.9	9.7	23.2	51.4	151.4
2001.....	0.6	0.9	1.4	3.1	4.5	10.4	23.4	50.4	156.0
2000.....	0.6	1.0	1.5	2.5	4.7	9.6	22.6	48.9	152.3
1999.....	0.5	0.9	1.7	2.0	4.3	9.3	19.0	43.8	142.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Non-Hispanic White, both sexes

2007 5/.....	0.9	1.2	2.0	3.4	6.0	13.3	26.6	60.8	162.0
2006 5/.....	0.8	1.3	2.0	3.9	6.4	12.7	26.8	61.4	164.1
2005 5/.....	0.8	1.5	2.0	3.7	6.3	12.9	28.8	62.4	169.0
2004 5/.....	0.8	1.2	1.9	3.1	6.0	13.9	28.8	64.0	170.4
2003 5/.....	0.9	1.2	2.0	3.5	6.0	13.3	30.4	66.3	181.0
2002.....	0.7	1.1	1.8	3.3	6.3	13.4	31.5	69.0	191.3
2001.....	0.7	1.2	1.8	3.6	6.1	13.9	30.7	69.0	191.2
2000.....	0.7	1.1	1.8	3.0	6.6	13.6	30.5	66.3	192.5
1999.....	0.6	0.9	1.9	3.0	5.9	12.7	27.9	60.9	179.3

Non-Hispanic White, male

2007 5/.....	1.2	1.6	2.7	4.2	7.6	17.7	36.9	85.4	236.9
2006 5/.....	1.0	1.7	2.5	5.1	8.6	17.4	37.2	90.1	239.9
2005 5/.....	0.9	1.7	2.4	4.7	8.2	17.6	38.3	87.6	237.8
2004 5/.....	1.0	1.6	2.2	3.8	7.9	18.5	39.3	90.4	247.4
2003 5/.....	1.0	1.5	2.5	4.3	8.4	17.5	41.6	93.4	256.5
2002.....	0.9	1.4	2.1	4.1	8.2	18.0	43.3	97.3	278.4
2001.....	0.9	1.6	2.3	4.1	8.1	18.8	41.3	100.4	276.9
2000.....	0.9	1.3	2.3	3.6	8.8	19.0	41.7	96.9	288.1
1999.....	0.8	0.9	2.2	4.0	7.9	17.4	40.9	90.2	267.4

Non-Hispanic White, female

2007 5/.....	0.6	0.9	1.4	2.6	4.5	9.6	18.8	45.1	126.5
2006 5/.....	0.6	0.9	1.5	2.8	4.4	8.8	19.0	43.3	128.7
2005 5/.....	0.8	1.2	1.6	2.8	4.7	9.1	21.7	46.7	137.5
2004 5/.....	0.5	0.8	1.6	2.5	4.3	10.1	21.1	47.6	135.9
2003 5/.....	0.7	0.8	1.5	2.7	4.0	9.8	22.3	49.7	147.8
2002.....	0.5	0.7	1.5	2.6	4.6	9.6	23.0	51.8	153.7
2001.....	0.5	0.9	1.4	3.1	4.3	9.8	23.2	50.1	155.5
2000.....	0.5	0.9	1.4	2.3	4.6	9.3	22.6	48.2	153.9
1999.....	0.4	0.8	1.5	2.0	4.2	8.9	18.8	43.7	144.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Non-Hispanic Black, both sexes

2007 5/.....	1.0	1.9	4.0	5.8	8.3	15.5	30.5	61.6	127.6
2006 5/.....	1.2	1.7	3.0	5.0	9.3	20.1	33.6	56.0	130.9
2005 5/.....	1.2	2.4	3.2	6.6	10.5	17.3	41.8	69.8	155.4
2004 5/.....	1.1	1.7	3.1	7.5	9.0	20.7	35.0	66.1	147.4
2003 5/.....	1.3	1.3	2.7	8.6	11.1	17.4	40.7	77.5	165.7
2002.....	0.8	2.2	2.4	5.6	12.0	19.2	42.7	68.3	167.3
2001.....	1.0	2.5	2.5	6.1	9.5	22.7	39.2	76.2	187.3
2000.....	1.7	2.3	3.0	5.9	11.5	25.5	39.1	72.4	160.6
1999.....	1.3	1.8	4.0	5.2	10.2	24.0	40.9	71.6	168.5

Non-Hispanic Black, male

2007 5/.....	*	2.3	6.1	7.5	12.1	24.3	48.3	94.3	163.2
2006 5/.....	*	2.4	3.8	7.7	15.5	28.8	56.2	87.1	177.8
2005 5/.....	1.6	3.5	3.7	10.1	15.4	26.0	63.8	107.9	205.5
2004 5/.....	*	2.2	4.5	11.4	14.0	31.6	52.2	94.7	207.3
2003 5/.....	1.8	*	3.9	11.7	14.7	26.6	60.7	110.2	215.2
2002.....	*	3.2	3.7	6.5	17.1	29.1	63.4	92.0	237.4
2001.....	*	3.5	3.2	8.4	12.8	31.3	56.9	107.9	243.7
2000.....	2.4	3.2	3.4	7.4	17.9	42.0	62.1	97.8	210.2
1999.....	*	*	5.2	8.3	15.7	37.7	65.3	111.6	229.4

Non-Hispanic Black, female

2007 5/.....	*	*	2.4	4.4	5.6	9.4	19.7	44.4	112.8
2006 5/.....	*	*	2.3	2.8	4.8	14.1	20.1	39.6	111.7
2005 5/.....	*	*	2.8	3.8	6.9	11.3	28.7	49.7	135.2
2004 5/.....	*	*	*	4.5	5.4	13.3	24.8	51.0	123.7
2003 5/.....	*	*	*	6.2	8.4	11.1	28.9	60.4	146.5
2002.....	*	*	*	4.9	8.2	12.6	30.3	56.0	140.5
2001.....	*	*	*	4.3	7.1	17.0	28.6	60.0	166.4
2000.....	*	*	*	4.7	6.9	14.5	25.3	59.6	142.2
1999.....	*	*	3.0	*	6.1	14.9	26.2	51.8	145.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes

2007.....	2.3	4.0	7.0	12.2	21.9	37.3	59.4	90.9	138.9
2006.....	2.3	4.0	6.7	11.9	21.4	36.6	59.1	91.9	136.6
2005.....	2.4	4.1	6.6	11.8	21.5	37.5	59.4	89.7	136.5
2004.....	2.6	4.0	7.3	12.4	21.7	39.5	61.4	91.3	139.4
2003.....	2.6	4.0	6.6	11.6	21.0	36.2	57.3	83.8	138.1
2002.....	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001.....	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000.....	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
1999.....	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6

All origins 3/, male

2007.....	2.7	4.7	8.0	14.4	26.0	47.0	76.0	116.2	169.2
2006.....	2.4	4.3	7.7	14.0	26.0	46.3	75.7	118.0	163.7
2005.....	2.7	4.4	7.3	14.1	25.9	46.0	74.5	113.9	157.7
2004.....	2.7	4.5	8.1	14.3	26.6	49.4	77.3	118.1	168.8
2003.....	2.8	4.3	7.5	13.8	25.9	43.8	71.8	105.5	165.0
2002.....	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001.....	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000.....	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
1999.....	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3

All origins 3/, female

2007.....	2.0	3.3	6.0	10.3	18.3	29.3	47.3	74.9	124.5
2006.....	2.3	3.8	5.7	10.0	17.3	28.7	47.0	75.6	124.0
2005.....	2.1	3.8	5.9	9.8	17.7	30.6	48.5	74.6	126.7
2004.....	2.6	3.4	6.6	10.7	17.4	31.6	50.0	74.8	126.1
2003.....	2.5	3.7	5.7	9.6	16.8	30.0	47.0	70.5	126.2
2002.....	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001.....	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000.....	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
1999.....	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, both sexes

2007	1.8	3.4	5.9	11.7	20.4	35.4	51.3	82.5	105.1
2006	2.0	3.4	5.6	10.5	20.3	32.1	46.9	78.5	106.8
2005	1.9	3.5	6.1	12.7	20.8	34.9	53.3	90.1	104.0
2004	1.8	3.3	6.3	9.9	19.7	37.0	60.5	87.4	112.7
2003	2.0	3.0	7.5	10.8	18.5	34.3	48.9	73.4	118.7
2002	2.2	3.4	6.2	11.1	18.7	35.3	49.3	72.6	112.0
2001	2.2	4.3	7.9	10.3	19.8	32.1	52.7	87.9	134.4
2000	2.2	3.1	8.6	11.2	20.5	31.4	53.3	84.1	135.4
1999	2.3	4.3	5.8	14.3	22.6	30.2	56.2	78.1	134.0

Hispanic, male

2007	2.4	4.7	6.8	14.2	27.0	38.5	57.5	93.8	128.1
2006	2.1	3.6	7.7	13.7	23.4	36.2	55.6	86.0	112.8
2005	2.3	3.9	6.5	14.9	24.2	35.8	65.4	108.6	114.6
2004	1.8	4.3	6.8	12.3	25.3	38.9	70.8	101.0	112.1
2003	2.3	4.2	8.9	14.8	21.8	42.7	56.5	74.8	127.8
2002	2.2	3.2	7.9	12.6	24.4	47.8	49.9	89.2	108.8
2001	2.5	4.7	8.6	13.3	23.4	38.9	64.8	103.9	167.6
2000	2.5	3.6	9.6	15.0	22.7	37.9	67.9	107.5	133.0
1999	2.6	3.7	7.2	17.3	26.7	35.6	71.0	105.3	161.8

Hispanic, female

2007	*	2.1	5.1	9.4	14.8	33.0	46.8	75.0	92.7
2006	2.0	3.3	3.6	7.6	17.8	28.8	40.6	73.6	103.6
2005	*	3.1	5.7	10.8	18.0	34.2	44.6	77.8	98.4
2004	*	2.3	5.8	7.7	15.1	35.5	53.2	78.5	113.1
2003	*	*	6.2	7.3	15.7	27.9	43.5	72.4	114.1
2002	2.1	3.6	4.7	9.8	14.1	25.7	48.9	62.0	113.6
2001	*	4.0	7.2	7.8	16.9	26.9	44.1	78.0	118.2
2000	*	*	7.7	7.9	18.7	26.5	42.9	69.9	136.5
1999	*	4.9	4.6	11.8	19.4	26.0	45.7	61.9	120.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 4/, both sexes

2007.....	2.4	4.1	7.0	12.3	22.0	37.4	59.9	91.4	140.6
2006.....	2.4	4.1	6.7	12.0	21.4	36.9	59.9	92.6	138.1
2005.....	2.5	4.1	6.6	11.7	21.6	37.6	59.7	89.6	138.0
2004.....	2.7	4.0	7.4	12.6	21.8	39.6	61.4	91.4	140.5
2003.....	2.7	4.1	6.5	11.6	21.2	36.2	57.7	84.2	138.7
2002.....	2.5	4.1	7.1	11.9	20.6	37.7	57.1	84.0	134.1
2001.....	2.6	4.4	7.2	12.8	23.0	38.2	62.1	89.4	142.7
2000.....	2.7	4.7	7.6	12.7	23.0	37.0	60.7	89.1	141.6
1999.....	2.7	4.1	6.9	12.5	22.4	37.0	59.5	87.7	147.7

Non-Hispanic 4/, male

2007.....	2.7	4.7	8.1	14.4	25.9	47.7	77.1	117.6	171.6
2006.....	2.5	4.3	7.7	14.0	26.2	47.0	77.0	119.9	166.4
2005.....	2.8	4.5	7.3	14.0	26.0	46.6	74.9	114.1	159.9
2004.....	2.8	4.6	8.2	14.5	26.6	50.0	77.6	118.7	171.6
2003.....	2.9	4.3	7.3	13.6	26.2	43.8	72.5	106.9	166.5
2002.....	2.5	4.3	7.8	13.9	24.3	47.7	71.6	104.7	158.5
2001.....	2.9	4.9	7.6	15.1	27.3	47.2	76.5	110.0	169.1
2000.....	2.9	5.1	8.7	14.6	27.0	46.0	77.5	112.3	181.3
1999.....	2.8	4.5	7.3	14.5	27.2	46.8	74.8	114.8	182.7

Non-Hispanic 4/, female

2007.....	2.1	3.4	6.0	10.3	18.5	29.0	47.3	74.9	126.0
2006.....	2.3	3.8	5.8	10.2	17.2	28.7	47.3	75.5	124.9
2005.....	2.2	3.8	5.9	9.7	17.7	30.3	48.6	74.4	128.0
2004.....	2.7	3.5	6.6	10.9	17.6	31.3	49.8	74.6	126.6
2003.....	2.5	3.8	5.7	9.8	16.8	30.1	47.1	70.4	126.4
2002.....	2.4	3.9	6.4	10.1	17.4	29.6	46.9	71.4	123.6
2001.....	2.4	3.9	6.8	10.7	19.2	31.0	52.0	77.1	131.8
2000.....	2.4	4.3	6.6	11.1	19.6	30.0	49.0	75.5	125.5
1999.....	2.6	3.7	6.6	10.8	18.2	29.4	48.8	71.8	133.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic White, both sexes

2007 5/.....	2.1	3.7	6.6	12.1	22.2	38.0	62.1	94.9	145.7
2006 5/.....	2.0	3.7	6.3	11.7	21.7	38.3	62.1	96.0	142.9
2005 5/.....	2.1	3.6	6.2	11.4	21.6	38.5	60.9	92.7	141.8
2004 5/.....	2.3	3.7	7.1	12.4	22.1	41.0	63.6	94.2	145.4
2003 5/.....	2.3	3.7	6.1	11.5	21.6	37.3	58.9	87.9	143.7
2002.....	2.0	3.8	6.7	11.6	21.0	38.2	59.0	85.8	138.6
2001.....	2.2	3.8	6.9	12.5	23.2	39.0	63.8	92.2	145.9
2000.....	2.2	4.4	7.4	12.5	22.9	37.9	62.1	91.0	145.7
1999.....	2.3	3.7	6.4	12.4	22.5	37.2	60.6	89.6	150.7

Non-Hispanic White, male

2007 5/.....	2.3	4.3	7.6	14.0	26.2	48.3	80.6	121.3	178.3
2006 5/.....	2.2	4.0	7.5	13.6	26.7	48.4	79.8	124.1	171.5
2005 5/.....	2.4	4.0	6.8	13.4	26.4	47.9	75.7	117.2	162.2
2004 5/.....	2.3	4.3	7.9	14.3	27.0	51.4	80.0	122.1	177.3
2003 5/.....	2.5	4.0	6.9	13.5	26.6	45.1	73.5	111.4	172.0
2002.....	2.1	4.0	7.5	13.4	25.2	48.1	73.6	107.1	164.1
2001.....	2.5	4.4	7.5	14.5	27.7	48.6	78.9	113.3	173.4
2000.....	2.4	4.7	8.4	14.2	27.0	46.6	79.0	114.9	185.9
1999.....	2.3	4.2	6.5	14.4	27.6	47.4	76.5	117.4	186.1

Non-Hispanic White, female

2007 5/.....	1.9	3.1	5.6	10.3	18.5	29.4	48.2	78.1	130.3
2006 5/.....	1.9	3.3	5.1	10.0	17.2	29.9	48.8	78.3	129.6
2005 5/.....	1.9	3.3	5.7	9.5	17.3	30.7	49.9	77.3	132.5
2004 5/.....	2.3	3.1	6.4	10.6	17.7	32.4	51.5	76.9	131.0
2003 5/.....	2.0	3.5	5.4	9.7	17.2	31.0	48.3	73.4	131.2
2002.....	2.0	3.6	5.9	9.9	17.2	30.2	48.5	72.8	127.6
2001.....	2.0	3.3	6.3	10.6	19.1	31.2	53.1	79.4	134.5
2000.....	2.1	4.0	6.4	10.8	19.3	30.8	50.3	76.9	129.5
1999.....	2.2	3.3	6.3	10.6	18.0	29.1	49.4	73.2	136.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic Black, both sexes

2007 5/.....	4.6	7.4	11.6	16.6	23.9	37.1	47.1	63.8	96.1
2006 5/.....	5.0	7.2	11.3	16.4	20.5	32.8	44.8	62.1	96.6
2005 5/.....	5.0	8.0	10.1	15.5	25.1	35.2	54.0	61.6	104.0
2004 5/.....	5.7	7.3	11.2	16.9	23.5	33.4	46.6	64.6	94.4
2003 5/.....	5.8	7.5	10.6	14.5	20.6	29.7	51.7	50.8	91.0
2002.....	5.6	6.9	11.1	17.2	21.2	34.5	41.4	68.9	90.7
2001.....	5.9	9.5	11.2	17.4	24.3	36.1	50.3	67.0	110.8
2000.....	6.1	7.9	11.8	15.9	28.2	33.5	50.2	71.8	99.8
1999.....	6.0	7.3	12.1	15.6	25.2	38.7	53.6	66.9	112.8

Non-Hispanic Black, male

2007 5/.....	5.7	8.7	13.6	21.0	26.7	46.1	54.7	86.0	116.1
2006 5/.....	5.0	6.8	10.6	20.3	24.9	44.6	58.5	81.5	119.1
2005 5/.....	5.5	8.6	12.3	20.1	28.9	41.6	72.1	83.5	134.2
2004 5/.....	6.5	7.3	12.9	19.1	29.2	42.2	61.6	83.0	117.9
2003 5/.....	5.7	7.4	12.9	18.7	25.8	38.8	70.6	66.4	116.1
2002.....	5.8	7.5	12.3	21.8	22.8	45.5	54.5	88.1	102.8
2001.....	6.4	9.6	10.8	22.2	27.9	39.6	59.7	85.0	131.7
2000.....	7.4	8.7	13.7	18.1	32.2	41.3	68.9	83.3	135.0
1999.....	6.8	7.9	14.1	18.1	28.7	44.7	67.2	85.0	146.2

Non-Hispanic Black, female

2007 5/.....	3.8	6.4	9.9	13.2	21.9	30.9	42.5	52.1	87.9
2006 5/.....	5.1	7.5	11.9	13.5	17.3	24.8	36.6	51.8	87.4
2005 5/.....	4.5	7.5	8.3	12.0	22.3	30.9	43.2	50.1	91.9
2004 5/.....	5.0	7.3	9.9	15.3	19.2	27.5	37.8	54.9	85.1
2003 5/.....	5.9	7.5	8.7	11.3	16.7	23.6	40.5	42.6	81.3
2002.....	5.4	6.3	10.2	13.5	20.1	27.1	33.6	58.8	86.1
2001.....	5.5	9.4	11.5	13.6	21.7	33.7	44.7	57.9	103.1
2000.....	5.1	7.2	10.3	14.3	25.2	28.4	38.9	66.1	86.8
1999.....	5.3	6.8	10.4	13.7	22.6	34.7	45.4	58.0	100.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

All origins 3/, both sexes

2007.....	0.5	0.9	1.0	1.3	2.1	3.0	4.9	8.2	17.3
2006.....	0.5	0.8	1.0	1.5	2.1	3.5	5.3	9.6	20.7
2005.....	0.5	0.8	0.9	1.6	2.2	3.8	5.8	10.8	21.4
2004.....	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
2003.....	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6

All origins 3/, male

2007.....	0.8	1.1	1.3	1.7	2.7	3.4	5.5	9.5	18.0
2006.....	0.8	1.0	1.4	1.9	2.6	4.0	6.3	11.0	22.0
2005.....	0.8	1.2	1.3	2.2	2.6	4.8	7.3	13.1	22.4
2004.....	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
2003.....	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5

All origins 3/, female

2007.....	0.2	0.6	0.8	1.0	1.6	2.7	4.5	7.4	16.9
2006.....	0.3	0.5	0.6	1.1	1.7	3.1	4.6	8.8	20.1
2005.....	0.3	0.5	0.6	1.1	1.8	3.0	4.7	9.4	21.0
2004.....	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
2003.....	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

Non-Hispanic 4/, both sexes

2007.....	0.5	0.9	1.0	1.3	2.1	3.1	5.1	8.4	17.7
2006.....	0.6	0.8	1.0	1.6	2.1	3.6	5.3	9.8	20.7
2005.....	0.5	0.8	1.0	1.6	2.2	3.8	5.8	10.9	21.6
2004.....	0.6	0.8	1.0	1.7	2.6	3.9	6.9	11.7	24.6
2003.....	0.6	0.8	1.1	1.7	2.3	4.1	7.3	13.5	26.7
2002.....	0.6	0.8	1.4	1.7	2.6	4.8	8.0	13.2	28.6
2001.....	0.7	0.8	1.3	2.2	3.1	4.8	8.3	16.0	33.6
2000.....	0.7	1.0	1.5	2.2	3.2	5.1	8.8	15.5	35.7
1999.....	0.6	0.9	1.4	2.2	3.3	5.9	9.1	17.2	34.8

Non-Hispanic 4/, male

2007.....	0.8	1.2	1.3	1.7	2.7	3.4	5.6	9.6	18.4
2006.....	0.8	1.1	1.5	1.9	2.5	4.0	6.2	11.3	22.2
2005.....	0.8	1.2	1.3	2.2	2.7	4.8	7.2	13.3	23.0
2004.....	0.7	1.0	1.2	2.1	2.9	4.6	8.3	13.0	24.8
2003.....	0.7	1.1	1.4	2.3	3.0	5.2	8.8	16.1	29.6
2002.....	0.6	1.2	1.7	2.0	3.0	6.1	9.3	16.8	31.7
2001.....	0.9	1.1	1.8	2.8	4.1	5.6	10.2	16.8	35.5
2000.....	0.9	1.6	2.1	2.8	3.9	6.9	11.3	18.1	41.0
1999.....	0.7	1.2	1.7	3.0	4.5	8.0	10.9	21.1	37.7

Non-Hispanic 4/, female

2007.....	0.3	0.7	0.8	1.0	1.6	2.8	4.6	7.5	17.3
2006.....	0.3	0.6	0.6	1.2	1.7	3.2	4.6	8.9	20.0
2005.....	0.3	0.5	0.6	1.1	1.8	3.0	4.8	9.4	21.0
2004.....	0.4	0.5	0.7	1.3	2.3	3.4	5.9	10.9	24.6
2003.....	0.4	0.5	0.9	1.1	1.7	3.3	6.3	11.9	25.4
2002.....	0.5	0.5	1.0	1.4	2.2	3.7	7.1	11.0	27.3
2001.....	0.5	0.6	0.8	1.6	2.3	4.3	7.0	15.5	32.8
2000.....	0.5	0.5	1.0	1.7	2.5	3.6	7.0	14.0	33.5
1999.....	0.5	0.5	1.1	1.5	2.3	4.3	7.9	14.9	33.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

Non-Hispanic White, both sexes

2007 5/.....	0.4	0.8	1.0	1.2	2.0	3.1	5.0	8.4	18.1
2006 5/.....	0.5	0.7	1.0	1.4	2.0	3.4	5.2	10.1	21.2
2005 5/.....	0.5	0.7	1.0	1.6	2.2	3.7	5.7	11.0	21.8
2004 5/.....	0.5	0.7	0.9	1.6	2.5	3.8	6.9	11.8	25.4
2003 5/.....	0.5	0.7	1.1	1.6	2.3	4.2	7.2	13.3	27.3
2002.....	0.5	0.8	1.2	1.6	2.5	4.8	7.9	13.7	29.4
2001.....	0.6	0.7	1.1	2.1	3.0	4.8	8.3	16.3	34.7
2000.....	0.6	0.9	1.4	2.2	3.1	5.1	8.7	15.9	36.3
1999.....	0.5	0.7	1.3	2.2	3.1	5.8	9.2	17.2	35.6

Non-Hispanic White, male

2007 5/.....	0.6	1.1	1.2	1.5	2.5	3.3	5.4	9.6	18.5
2006 5/.....	0.7	0.8	1.4	1.8	2.3	3.6	5.9	11.6	22.2
2005 5/.....	0.6	1.0	1.3	2.1	2.6	4.6	6.9	13.7	23.1
2004 5/.....	0.6	1.0	1.2	2.0	2.9	4.4	8.1	12.7	25.7
2003 5/.....	0.6	0.9	1.2	2.2	2.9	5.1	8.5	15.5	29.8
2002.....	0.6	1.0	1.5	1.9	2.9	6.0	9.2	17.3	32.6
2001.....	0.8	0.9	1.5	2.7	3.9	5.3	9.9	16.9	36.3
2000.....	0.8	1.3	2.0	2.8	3.8	6.7	11.0	18.1	41.0
1999.....	0.6	1.1	1.5	2.9	4.1	7.9	11.0	21.1	38.6

Non-Hispanic White, female

2007 5/.....	0.3	0.6	0.8	0.9	1.6	3.0	4.7	7.7	18.0
2006 5/.....	0.3	0.5	0.5	1.1	1.6	3.3	4.6	9.1	20.7
2005 5/.....	0.3	0.5	0.7	1.1	1.9	2.9	4.8	9.3	21.2
2004 5/.....	0.3	0.5	0.7	1.3	2.2	3.3	6.0	11.2	25.3
2003 5/.....	0.3	0.5	0.9	1.1	1.8	3.4	6.2	12.0	26.2
2002.....	0.5	0.5	1.0	1.3	2.2	3.9	7.1	11.5	27.9
2001.....	0.5	0.6	0.8	1.5	2.3	4.5	7.1	15.9	34.0
2000.....	0.4	0.5	0.9	1.7	2.4	3.7	7.0	14.5	34.4
1999.....	0.4	0.4	1.1	1.6	2.2	4.3	7.9	14.9	34.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

Non-Hispanic Black, both sexes

2007 5/.....	1.1	1.5	1.8	2.5	3.0	2.7	5.1	6.4	11.5
2006 5/.....	1.1	1.7	1.8	2.6	3.0	4.8	7.2	7.5	14.9
2005 5/.....	1.1	1.4	1.3	2.2	2.8	5.5	5.5	9.9	17.5
2004 5/.....	1.3	1.3	1.7	2.7	3.6	5.2	6.8	9.4	16.1
2003 5/.....	1.2	1.8	2.0	2.4	2.2	4.5	8.7	14.8	17.3
2002.....	0.9	1.6	2.4	2.9	3.3	5.4	8.8	6.7	18.8
2001.....	1.1	1.6	2.5	3.1	3.5	5.6	8.0	10.5	20.9
2000.....	1.4	2.1	2.6	2.7	4.2	5.7	9.4	11.2	26.7
1999.....	1.3	1.9	2.5	3.2	5.1	6.5	7.6	16.4	25.5

Non-Hispanic Black, male

2007 5/.....	2.0	2.0	2.8	3.4	*	*	*	*	*
2006 5/.....	1.8	2.9	2.6	*	*	8.2	10.1	*	*
2005 5/.....	1.9	2.6	*	*	*	8.1	*	*	20.0
2004 5/.....	2.1	*	*	4.0	*	*	10.3	*	*
2003 5/.....	1.7	3.0	3.0	4.0	*	7.1	11.7	24.1	*
2002.....	*	2.9	3.7	4.3	*	8.2	11.7	*	*
2001.....	1.9	2.8	4.0	4.6	5.2	9.3	10.9	*	24.3
2000.....	*	3.4	3.6	*	*	10.3	12.5	*	31.7
1999.....	1.9	2.9	3.8	5.2	7.7	9.0	*	18.6	28.5

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	*	*	*	11.2
2006 5/.....	*	*	*	*	*	*	5.5	*	14.2
2005 5/.....	*	*	*	*	*	*	*	10.6	16.5
2004 5/.....	*	*	*	*	3.7	4.5	*	7.7	16.9
2003 5/.....	*	*	*	*	*	*	6.9	9.9	16.1
2002.....	*	*	*	*	*	*	7.0	*	18.4
2001.....	*	*	*	*	*	*	6.2	9.2	19.7
2000.....	*	*	*	*	*	*	7.6	8.6	24.9
1999.....	*	*	*	*	*	4.8	7.0	15.4	24.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

All origins 3/, both sexes

2007.....	0.1	0.1	0.2	0.2	0.3	0.6	0.7	1.0	1.5
2006.....	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.7
2005.....	*	0.1	0.2	0.2	0.4	0.4	0.8	0.9	2.1
2004.....	*	0.2	0.1	0.2	0.3	0.6	0.8	1.4	1.7
2003.....	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

All origins 3/, male

2007.....	*	*	0.2	0.3	*	0.8	0.9	1.7	2.1
2006.....	*	*	0.2	*	0.5	0.8	1.1	1.5	2.3
2005.....	*	*	0.3	*	0.6	0.7	0.9	1.3	2.8
2004.....	*	0.2	*	*	*	0.7	1.1	1.7	2.5
2003.....	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	0.5	0.9	*	2.9

All origins 3/, female

2007.....	*	*	*	*	*	0.5	0.6	0.6	1.2
2006.....	*	*	*	*	*	*	*	0.7	1.5
2005.....	*	*	*	*	*	*	0.7	0.7	1.8
2004.....	*	*	*	*	*	0.5	0.6	1.2	1.4
2003.....	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	1.1	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

Non-Hispanic 4/, both sexes

2007.....	*	*	0.1	0.2	0.3	0.6	0.7	1.1	1.5
2006.....	*	*	0.2	0.2	0.4	0.6	0.5	1.0	1.7
2005.....	*	*	0.2	0.2	0.4	0.5	0.8	1.0	2.2
2004.....	*	0.1	0.1	0.2	0.3	0.6	0.8	1.5	1.8
2003.....	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.6
2002.....	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.1
2001.....	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.2
2000.....	*	0.2	0.2	0.3	0.3	0.5	0.7	1.4	1.7
1999.....	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

Non-Hispanic 4/, male

2007.....	*	*	*	0.3	*	0.8	0.9	1.8	2.2
2006.....	*	*	0.2	*	0.6	0.9	1.0	1.5	2.3
2005.....	*	*	0.3	*	0.6	0.7	0.9	1.4	3.0
2004.....	*	*	*	*	*	0.7	1.1	1.8	2.5
2003.....	*	*	*	*	*	0.7	0.9	2.3	2.2
2002.....	*	*	*	0.4	0.6	0.9	1.1	1.7	2.3
2001.....	*	*	*	*	0.5	0.8	0.7	1.5	3.0
2000.....	*	*	*	0.5	*	0.7	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.9	*	3.0

Non-Hispanic 4/, female

2007.....	*	*	*	*	*	0.5	0.6	0.6	1.2
2006.....	*	*	*	*	*	*	*	0.7	1.5
2005.....	*	*	*	*	*	*	0.7	0.7	1.8
2004.....	*	*	*	*	*	0.5	0.6	1.3	1.5
2003.....	*	*	*	*	*	*	0.7	0.9	1.4
2002.....	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	0.5	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.6	1.2	1.5
1999.....	*	*	*	*	*	*	*	1.1	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

Non-Hispanic White, both sexes

2007 5/.....	*	*	*	0.2	0.3	0.6	0.7	1.0	1.6
2006 5/.....	*	*	0.2	*	0.4	0.6	0.5	1.0	1.8
2005 5/.....	*	*	0.2	*	0.4	0.5	0.7	1.0	2.3
2004 5/.....	*	*	0.2	*	0.3	0.6	0.8	1.4	1.8
2003 5/.....	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	0.2	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.3	0.3	0.4	0.6	1.3	1.8
1999.....	*	0.2	*	*	0.3	0.4	0.6	1.1	1.6

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	0.8	0.9	1.7	2.3
2006 5/.....	*	*	*	*	0.6	0.8	0.9	1.5	2.4
2005 5/.....	*	*	*	*	0.6	0.7	0.8	1.5	3.1
2004 5/.....	*	*	*	*	*	0.7	1.1	1.8	2.3
2003 5/.....	*	*	*	*	*	0.6	0.8	2.4	2.4
2002.....	*	*	*	0.5	*	0.8	1.1	1.7	2.1
2001.....	*	*	*	*	*	0.9	*	1.4	2.6
2000.....	*	*	*	0.5	*	*	1.0	1.5	2.1
1999.....	*	*	*	*	*	*	0.9	*	3.1

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	0.6	*	1.3
2006 5/.....	*	*	*	*	*	*	*	0.7	1.5
2005 5/.....	*	*	*	*	*	*	0.7	0.7	1.9
2004 5/.....	*	*	*	*	*	*	0.6	1.1	1.6
2003 5/.....	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.6	0.9	2.1
2001.....	*	*	*	*	*	*	*	*	2.0
2000.....	*	*	*	*	*	*	*	1.2	1.6
1999.....	*	*	*	*	*	*	*	1.1	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

All origins 3/, both sexes

2007.....	0.2	0.3	0.4	0.7	1.0	1.6	2.7	4.8	11.3
2006.....	0.1	0.3	0.5	0.7	0.9	1.7	2.8	5.4	12.1
2005.....	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.8	12.1
2004.....	0.2	0.2	0.4	0.5	0.9	1.8	2.9	5.3	11.0
2003.....	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7

All origins 3/, male

2007.....	0.2	0.3	0.5	0.7	0.7	1.9	2.7	4.8	12.0
2006.....	*	0.3	0.4	0.8	0.8	1.9	2.9	5.7	13.5
2005.....	*	0.3	0.4	0.9	0.7	1.3	2.4	4.6	12.7
2004.....	0.2	0.3	0.4	0.5	0.8	2.2	3.2	4.7	11.3
2003.....	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2

All origins 3/, female

2007.....	*	0.2	0.3	0.6	1.2	1.4	2.7	4.9	11.0
2006.....	*	0.3	0.5	0.7	1.0	1.5	2.7	5.1	11.4
2005.....	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
2004.....	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	10.8
2003.....	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

Non-Hispanic 4/, both sexes

2007	0.2	0.3	0.4	0.7	0.9	1.7	2.8	4.9	11.4
2006	0.1	0.3	0.4	0.8	0.9	1.7	2.8	5.4	12.3
2005	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.9	12.1
2004	0.2	0.2	0.4	0.5	0.9	1.7	2.9	5.3	11.2
2003	0.2	0.3	0.4	0.7	1.0	1.6	2.7	4.8	12.1
2002	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.2	13.0
2001	0.1	0.3	0.4	0.6	0.9	1.5	2.4	5.0	12.5
2000	0.1	0.3	0.4	0.6	0.9	1.4	2.9	4.9	13.1
1999	*	0.2	0.3	0.7	0.8	1.5	2.5	5.4	12.7

Non-Hispanic 4/, male

2007	0.2	0.3	0.5	0.8	0.7	2.0	2.8	4.9	12.3
2006	*	0.3	0.4	0.8	0.8	1.9	3.0	5.5	13.9
2005	*	0.3	0.4	0.9	0.7	1.4	2.5	4.8	12.8
2004	*	0.2	0.4	0.5	0.8	2.1	3.3	4.7	11.6
2003	0.2	0.4	0.4	0.8	1.0	1.7	3.0	5.2	12.5
2002	*	*	0.4	0.5	0.9	1.7	2.7	5.1	11.9
2001	*	0.3	0.4	0.8	0.9	1.4	2.4	5.2	12.1
2000	*	0.4	*	0.6	0.9	1.3	2.9	5.6	12.8
1999	*	*	0.4	0.7	0.8	1.4	2.5	6.5	13.3

Non-Hispanic 4/, female

2007	*	0.2	0.3	0.7	1.1	1.4	2.7	4.9	11.0
2006	*	0.3	0.5	0.7	0.9	1.5	2.7	5.3	11.6
2005	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
2004	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	11.0
2003	*	0.2	0.5	0.7	1.0	1.6	2.4	4.5	12.0
2002	*	0.3	0.4	0.7	0.9	1.5	2.9	5.3	13.4
2001	*	0.3	0.5	0.6	0.8	1.5	2.5	4.8	12.7
2000	*	0.3	0.5	0.5	0.8	1.5	3.0	4.5	13.2
1999	*	*	*	0.6	0.8	1.6	2.5	4.7	12.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.2	0.4	0.7	0.9	1.7	2.9	4.8	12.2
2006 5/.....	0.1	0.3	0.4	0.8	0.9	1.7	2.8	5.3	12.8
2005 5/.....	0.1	0.3	0.4	0.9	0.9	1.4	2.5	5.1	12.5
2004 5/.....	0.2	0.2	0.4	0.5	0.9	1.7	3.1	5.4	11.4
2003 5/.....	0.2	0.3	0.4	0.7	1.0	1.6	2.8	4.7	12.4
2002.....	0.1	0.2	0.4	0.5	0.8	1.6	3.0	5.3	13.6
2001.....	*	0.3	0.4	0.6	0.9	1.5	2.4	5.0	12.8
2000.....	*	0.3	0.4	0.6	0.8	1.5	2.9	5.0	13.3
1999.....	*	0.2	0.3	0.7	0.7	1.5	2.5	5.5	13.0

Non-Hispanic White, male

2007 5/.....	*	*	0.4	0.7	0.7	2.0	2.7	4.6	12.8
2006 5/.....	*	0.3	0.3	0.9	0.8	1.8	3.0	5.3	14.5
2005 5/.....	*	0.3	0.4	0.9	0.7	1.4	2.4	4.9	12.9
2004 5/.....	*	*	0.4	0.5	0.8	2.2	3.4	4.7	11.7
2003 5/.....	*	0.4	0.4	0.7	1.0	1.6	3.2	5.0	12.6
2002.....	*	*	0.4	*	0.8	1.6	2.9	5.2	12.2
2001.....	*	0.3	*	0.7	0.9	1.4	2.4	5.2	11.9
2000.....	*	0.3	*	0.6	0.9	1.4	3.0	5.7	13.0
1999.....	*	*	0.4	0.8	0.8	1.5	2.5	6.6	13.0

Non-Hispanic White, female

2007 5/.....	*	*	0.4	0.7	1.1	1.4	3.0	5.0	11.8
2006 5/.....	*	*	0.5	0.8	0.9	1.5	2.6	5.4	12.0
2005 5/.....	*	*	0.4	0.8	1.0	1.4	2.6	5.2	12.3
2004 5/.....	*	*	0.4	0.4	1.0	1.4	2.8	5.8	11.3
2003 5/.....	*	*	0.4	0.7	1.0	1.7	2.5	4.5	12.4
2002.....	*	*	0.3	0.6	0.9	1.5	3.1	5.3	14.2
2001.....	*	0.3	0.5	0.5	0.8	1.5	2.4	4.9	13.1
2000.....	*	0.3	0.5	0.5	0.7	1.5	2.8	4.6	13.4
1999.....	*	*	*	0.6	0.6	1.5	2.4	4.8	12.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes

2007.....	15.5	22.2	24.6	24.3	25.5	28.2	29.3	27.2	19.8
2006.....	14.8	21.2	22.8	22.9	24.8	27.4	29.6	28.2	19.4
2005.....	15.0	20.7	23.0	24.1	25.4	29.3	29.8	27.9	19.7
2004.....	15.1	21.3	21.7	23.7	26.2	29.5	30.1	27.0	19.7
2003.....	15.9	21.1	22.0	24.4	28.0	31.2	31.0	28.5	20.1
2002.....	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001.....	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000.....	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
1999.....	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2

All origins 3/, male

2007.....	21.5	32.6	37.1	35.4	36.0	36.5	38.2	34.6	25.8
2006.....	20.7	31.8	33.7	33.1	34.1	35.5	36.9	34.4	26.1
2005.....	21.3	31.3	34.5	35.3	33.9	38.1	37.2	35.1	27.6
2004.....	21.5	31.9	32.3	33.4	35.7	38.7	38.6	34.0	25.5
2003.....	23.4	32.3	32.4	35.6	38.4	40.0	38.1	35.0	27.5
2002.....	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001.....	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000.....	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
1999.....	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1

All origins 3/, female

2007.....	9.6	12.2	12.9	14.1	16.3	21.4	22.7	22.6	17.0
2006.....	9.0	11.1	12.5	13.6	16.7	20.9	24.2	24.3	16.2
2005.....	8.9	10.6	12.2	13.9	17.9	22.2	24.4	23.4	16.0
2004.....	9.0	11.1	11.8	14.9	18.0	22.1	24.0	22.7	17.1
2003.....	8.6	10.3	12.2	14.2	19.0	24.1	26.0	24.5	16.8
2002.....	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001.....	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000.....	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
1999.....	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

Hispanic, both sexes

2007	19.4	31.2	31.4	41.1	44.9	54.3	56.5	46.2	35.8
2006	19.4	29.4	33.0	36.1	41.2	45.2	52.3	57.6	40.3
2005	19.5	28.7	36.4	37.5	47.0	51.2	58.0	47.8	34.7
2004	20.3	30.7	31.7	38.7	43.6	52.1	62.7	56.3	26.2
2003	21.4	28.1	35.6	42.3	47.2	56.0	66.8	54.1	31.1
2002	23.1	31.9	37.1	40.1	50.5	63.2	52.8	54.8	45.5
2001	25.0	31.9	35.9	43.2	51.1	58.6	62.0	57.6	36.9
2000	24.8	30.4	39.6	43.7	55.6	61.8	65.8	60.2	48.4
1999	25.2	28.4	35.6	43.2	53.6	59.7	60.4	57.6	51.2

Hispanic, male

2007	30.5	49.0	50.6	63.4	64.3	74.4	69.4	42.6	43.7
2006	30.7	46.3	52.7	56.3	55.6	58.7	62.3	59.0	51.9
2005	31.7	46.0	56.5	61.2	66.1	73.1	69.7	67.4	36.1
2004	32.7	51.7	49.4	56.7	54.9	65.4	76.0	62.3	29.4
2003	35.4	46.6	56.7	70.1	59.8	63.2	79.6	57.2	35.4
2002	38.0	52.4	61.1	58.5	76.5	79.4	60.8	70.9	45.2
2001	41.9	52.5	58.9	70.6	78.0	75.9	81.8	64.9	41.9
2000	41.2	51.5	64.7	67.9	86.9	78.8	89.3	61.8	60.5
1999	42.3	46.9	59.4	66.7	76.5	82.5	72.6	67.6	66.9

Hispanic, female

2007	7.6	13.4	13.2	21.1	28.5	38.5	47.2	48.6	31.4
2006	7.5	12.6	14.4	18.1	29.1	34.7	45.1	56.7	34.2
2005	6.8	11.6	17.5	16.6	31.0	34.2	49.7	34.8	33.9
2004	7.5	10.1	15.2	23.0	34.2	41.8	53.2	52.3	24.5
2003	7.0	10.0	16.2	18.0	36.8	50.5	57.8	52.1	28.9
2002	7.9	12.1	15.0	24.2	29.3	50.7	47.1	44.5	45.6
2001	8.0	12.2	15.0	19.5	29.3	45.4	47.9	53.1	34.4
2000	8.4	10.3	16.9	22.8	30.2	49.0	49.1	59.1	42.5
1999	8.2	10.8	14.2	22.8	35.0	42.5	51.8	51.6	43.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 4/, both sexes

2007.....	14.9	21.2	23.9	22.8	23.9	26.1	27.3	26.0	19.0
2006.....	14.1	20.3	21.8	21.7	23.4	26.1	28.0	26.4	18.3
2005.....	14.4	19.8	21.8	22.9	23.6	27.7	27.9	26.8	18.9
2004.....	14.5	20.2	20.8	22.5	24.8	27.9	28.1	25.5	19.4
2003.....	15.2	20.3	20.8	22.8	26.5	29.5	29.0	27.2	19.6
2002.....	15.1	19.1	20.3	23.1	26.1	29.0	31.3	28.7	20.4
2001.....	16.3	19.0	19.8	23.2	27.2	29.4	29.7	27.4	21.6
2000.....	15.9	17.5	20.4	24.7	26.9	29.0	30.3	28.3	22.1
1999.....	15.5	17.6	20.3	25.2	28.0	30.0	31.3	29.8	22.1

Non-Hispanic 4/, male

2007.....	20.2	30.6	35.7	33.0	33.7	33.5	36.0	34.1	24.7
2006.....	19.3	30.2	31.9	31.1	32.2	33.7	35.1	32.8	24.6
2005.....	19.8	29.6	32.4	33.1	31.3	35.5	35.1	33.3	27.1
2004.....	20.0	29.7	30.7	31.5	34.1	36.8	36.3	32.5	25.3
2003.....	21.9	30.7	30.3	32.7	36.7	38.4	35.7	33.9	27.1
2002.....	21.5	28.8	29.4	32.4	34.3	37.4	38.1	35.3	27.7
2001.....	23.4	28.0	29.1	32.4	36.8	38.6	35.4	34.1	31.4
2000.....	23.7	26.4	29.1	35.3	36.0	38.5	38.4	33.4	33.0
1999.....	23.6	27.1	30.3	35.0	39.4	39.2	38.5	37.6	31.6

Non-Hispanic 4/, female

2007.....	9.9	12.1	12.8	13.5	15.3	20.0	20.9	21.0	16.3
2006.....	9.1	10.9	12.3	13.2	15.7	19.8	22.8	22.5	15.3
2005.....	9.1	10.4	11.7	13.6	16.8	21.3	22.7	22.8	15.2
2004.....	9.1	11.1	11.5	14.2	16.6	20.7	22.3	21.2	16.7
2003.....	8.7	10.3	11.8	13.9	17.6	22.3	24.2	23.2	16.3
2002.....	9.0	9.9	11.8	14.6	19.0	22.3	26.6	24.7	17.3
2001.....	9.3	10.3	11.1	14.8	18.8	22.0	25.7	23.4	17.5
2000.....	8.4	9.0	12.3	15.1	19.1	21.5	24.7	25.3	17.7
1999.....	7.8	8.5	10.9	16.3	18.3	22.9	26.3	25.3	18.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic White, both sexes

2007 5/.....	15.5	21.6	23.8	23.2	24.7	27.0	28.5	27.4	19.7
2006 5/.....	14.7	20.7	22.1	22.2	24.1	27.6	29.0	27.7	19.0
2005 5/.....	14.6	19.8	22.0	22.7	24.4	29.0	28.8	27.5	19.5
2004 5/.....	14.7	20.3	20.4	22.4	25.3	29.2	29.7	26.4	19.9
2003 5/.....	15.3	20.0	20.5	22.8	27.1	30.8	30.7	28.5	20.3
2002.....	15.0	18.7	20.1	23.1	26.9	30.2	32.9	29.6	21.2
2001.....	16.0	18.1	19.1	23.5	27.7	30.2	30.9	28.6	22.5
2000.....	15.7	16.7	19.9	24.7	27.2	29.7	31.6	29.1	22.8
1999.....	14.6	17.2	19.4	24.7	28.3	30.8	32.2	30.9	22.6

Non-Hispanic White, male

2007 5/.....	21.1	31.1	35.4	33.4	34.2	34.3	37.0	35.9	26.1
2006 5/.....	20.1	30.6	32.1	31.6	32.8	35.4	36.0	34.7	25.6
2005 5/.....	19.9	29.7	32.5	32.5	32.0	36.8	35.7	33.8	28.1
2004 5/.....	20.1	29.8	30.0	31.4	34.4	38.1	38.1	33.7	26.4
2003 5/.....	22.2	30.1	29.5	32.5	37.5	40.0	37.6	34.9	27.9
2002.....	21.6	28.2	29.1	32.3	34.8	38.7	40.2	36.3	28.7
2001.....	23.2	26.7	27.7	32.3	36.8	39.6	36.6	35.8	32.9
2000.....	23.8	25.1	28.5	35.4	36.4	39.2	39.7	34.7	34.1
1999.....	22.3	26.4	29.0	34.7	39.5	40.0	39.3	39.1	32.7

Non-Hispanic White, female

2007 5/.....	10.0	12.3	12.6	13.6	16.1	21.0	22.2	22.0	16.7
2006 5/.....	9.4	11.0	12.5	13.4	16.3	21.0	23.8	23.4	15.9
2005 5/.....	9.3	10.1	11.8	13.5	17.7	22.4	23.6	23.6	15.6
2004 5/.....	9.4	11.0	11.2	14.1	17.2	21.9	23.5	21.9	17.0
2003 5/.....	8.6	10.0	11.7	13.8	18.0	23.3	25.7	24.5	17.0
2002.....	8.5	9.5	11.5	14.7	19.8	23.3	27.7	25.5	18.0
2001.....	9.0	9.7	10.8	15.3	19.6	22.6	26.8	24.3	18.2
2000.....	7.7	8.4	11.8	14.8	19.0	22.1	26.0	25.9	18.2
1999.....	7.0	8.1	10.3	15.5	18.5	23.5	27.2	26.1	18.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic Black, both sexes

2007 5/.....	12.7	20.9	27.9	23.9	22.0	22.7	18.3	11.5	10.0
2006 5/.....	11.2	21.3	23.5	20.8	21.1	16.4	19.6	11.6	10.9
2005 5/.....	14.6	22.1	24.9	25.8	20.3	20.7	19.9	18.4	12.0
2004 5/.....	14.7	23.5	26.3	25.3	23.3	20.4	13.9	15.2	12.1
2003 5/.....	15.3	26.1	26.7	25.3	25.8	22.0	13.8	14.5	12.5
2002.....	17.8	25.3	24.9	25.8	23.2	19.9	19.7	16.3	12.1
2001.....	19.4	27.2	28.3	23.8	27.1	23.0	20.6	14.6	10.6
2000.....	19.8	26.3	26.0	26.3	27.8	25.0	18.3	17.8	14.0
1999.....	22.8	23.2	28.4	31.4	28.3	25.1	23.3	19.1	16.8

Non-Hispanic Black, male

2007 5/.....	16.0	31.9	43.8	37.2	37.6	34.9	31.9	13.9	*
2006 5/.....	15.3	32.2	36.4	30.1	34.6	22.5	32.2	*	*
2005 5/.....	21.5	32.5	39.0	41.2	31.1	31.9	33.1	23.7	*
2004 5/.....	21.6	35.1	41.8	37.4	36.2	32.3	19.6	22.0	*
2003 5/.....	22.0	42.0	42.5	39.3	37.2	29.0	18.7	22.6	22.4
2002.....	23.9	38.9	37.3	39.3	35.1	28.7	22.1	21.8	*
2001.....	28.9	42.8	44.8	38.5	43.8	34.4	27.0	*	*
2000.....	27.2	41.2	38.9	37.8	38.4	37.9	27.9	19.7	*
1999.....	34.8	37.1	44.3	43.3	44.1	37.0	32.4	25.7	*

Non-Hispanic Black, female

2007 5/.....	9.7	11.6	14.9	13.5	10.6	14.4	10.1	10.2	10.1
2006 5/.....	7.6	12.2	12.9	13.6	11.2	12.2	12.1	11.1	9.0
2005 5/.....	8.7	13.4	13.3	13.9	12.3	13.1	12.0	15.6	10.7
2004 5/.....	8.6	13.8	13.5	15.9	13.7	12.4	10.5	11.6	11.2
2003 5/.....	9.5	12.6	13.7	14.4	17.3	17.3	10.8	10.3	8.7
2002.....	12.6	13.9	14.7	15.3	14.4	14.0	18.2	13.4	10.5
2001.....	11.3	14.0	14.8	12.3	14.8	15.4	16.7	14.6	9.8
2000.....	13.3	13.6	15.4	17.3	20.1	16.3	12.5	16.9	11.8
1999.....	12.6	11.4	15.5	22.0	16.5	17.2	17.8	15.8	15.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

All origins 3/, both sexes

2007.....	10.1	13.4	13.7	12.5	10.8	9.2	6.9	4.3	2.1
2006.....	9.2	12.0	12.5	11.6	10.1	8.6	7.0	4.5	2.2
2005.....	9.3	12.0	12.4	12.0	10.3	9.4	6.4	4.8	1.8
2004.....	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
2003.....	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8

All origins 3/, male

2007.....	14.2	20.2	21.5	20.1	17.9	15.7	12.2	7.9	4.7
2006.....	12.9	18.4	19.4	18.1	16.9	14.1	12.0	7.5	4.7
2005.....	13.2	18.5	19.5	19.4	16.5	15.0	11.2	8.7	4.2
2004.....	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
2003.....	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1

All origins 3/, female

2007.....	6.1	6.9	6.3	5.6	4.7	3.9	3.0	1.9	0.9
2006.....	5.6	5.9	6.0	5.6	4.2	4.1	3.4	2.6	1.1
2005.....	5.4	5.8	5.7	5.3	4.9	4.8	3.0	2.4	0.7
2004.....	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
2003.....	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

Hispanic, both sexes

2007	13.4	19.6	18.9	22.1	20.4	19.2	17.7	9.0	7.2
2006	12.7	18.3	18.9	20.0	17.5	16.2	16.2	11.8	*
2005	12.0	17.5	19.2	18.6	20.5	19.8	13.1	8.7	*
2004	13.1	17.9	17.5	16.8	16.3	15.9	17.0	10.9	*
2003	13.2	16.0	21.0	22.0	20.1	14.7	16.1	10.7	*
2002	13.6	19.3	20.1	19.5	21.2	20.4	11.7	*	*
2001	16.0	18.6	20.9	22.2	22.2	19.8	19.3	12.9	*
2000	15.4	18.6	22.7	24.3	25.5	19.7	19.0	*	*
1999	16.1	17.8	21.5	22.1	21.4	19.5	16.5	*	*

Hispanic, male

2007	21.5	32.3	33.2	39.8	37.5	37.4	33.2	*	*
2006	20.5	30.4	32.5	35.8	31.9	32.4	30.9	21.3	*
2005	19.8	29.3	33.3	33.7	37.5	38.9	24.8	21.9	*
2004	22.6	32.1	29.5	30.0	28.1	29.4	34.0	18.9	*
2003	23.0	28.8	36.8	40.6	31.8	25.2	30.4	*	*
2002	22.6	33.0	36.7	35.7	41.3	39.1	19.2	*	*
2001	27.5	31.7	37.8	40.5	43.9	39.3	34.1	*	*
2000	26.7	32.8	41.9	44.0	51.8	39.9	41.3	*	*
1999	27.7	31.0	40.2	39.6	39.8	39.2	29.6	*	*

Hispanic, female

2007	4.8	6.8	5.2	6.2	5.9	*	*	*	*
2006	4.5	6.3	6.0	6.0	5.4	*	*	*	*
2005	3.9	5.9	5.9	5.2	6.4	*	*	*	*
2004	3.3	4.0	6.4	5.3	6.6	*	*	*	*
2003	3.2	3.6	6.3	5.8	10.4	6.6	*	*	*
2002	4.5	6.0	4.9	5.4	*	*	*	*	*
2001	4.5	6.0	5.6	6.3	*	*	*	*	*
2000	4.1	5.0	5.4	7.2	*	*	*	*	*
1999	4.5	5.3	4.6	6.9	6.6	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

Non-Hispanic 4/, both sexes

2007.....	9.6	12.7	13.2	11.7	10.0	8.5	6.1	4.0	1.9
2006.....	8.7	11.4	11.9	10.8	9.5	8.0	6.4	4.0	1.9
2005.....	8.9	11.4	11.8	11.5	9.5	8.5	6.0	4.6	1.7
2004.....	8.6	11.5	11.2	11.1	10.4	8.5	5.8	3.8	2.1
2003.....	9.0	11.3	10.8	10.8	10.3	7.8	6.0	4.1	1.7
2002.....	8.9	10.6	10.6	11.2	9.8	8.8	6.4	4.2	1.9
2001.....	9.6	10.6	10.3	11.1	10.3	8.2	6.3	3.6	2.2
2000.....	9.4	10.0	10.8	12.0	10.4	8.5	6.6	4.2	2.2
1999.....	9.4	10.2	10.8	11.9	10.9	8.8	6.8	5.1	2.5

Non-Hispanic 4/, male

2007.....	13.1	18.7	20.3	18.4	16.3	14.1	10.7	7.5	3.9
2006.....	11.8	17.1	18.1	16.6	15.6	12.8	10.7	6.7	3.9
2005.....	12.3	17.2	18.2	18.2	14.8	13.2	10.4	7.9	3.8
2004.....	11.6	17.1	17.1	17.3	16.9	13.9	10.2	7.4	4.5
2003.....	13.1	17.2	16.5	16.8	16.6	13.4	10.2	7.7	3.4
2002.....	12.7	16.2	15.9	17.8	15.5	13.9	10.3	7.6	4.3
2001.....	14.0	15.7	16.3	17.3	15.9	13.6	11.0	6.3	5.1
2000.....	14.0	15.5	16.1	18.4	16.4	14.3	12.4	7.3	5.0
1999.....	14.3	16.0	17.0	18.5	18.3	14.8	11.2	9.2	5.5

Non-Hispanic 4/, female

2007.....	6.3	6.9	6.4	5.5	4.6	3.9	2.7	1.8	0.9
2006.....	5.7	5.9	6.0	5.5	4.1	4.1	3.3	2.4	1.0
2005.....	5.6	5.8	5.7	5.3	4.8	4.8	2.9	2.5	0.8
2004.....	5.7	6.2	5.7	5.4	4.6	4.0	2.8	1.5	1.0
2003.....	4.9	5.6	5.5	5.3	4.8	3.4	3.1	1.9	1.0
2002.....	5.1	5.3	5.6	5.1	4.9	4.7	3.6	2.2	0.9
2001.....	5.3	5.8	4.6	5.5	5.4	4.0	3.0	1.9	1.0
2000.....	4.8	4.8	5.8	6.1	5.2	3.9	2.6	2.4	1.0
1999.....	4.6	4.7	4.9	5.8	4.5	4.0	3.8	2.6	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

Non-Hispanic White, both sexes

2007 5/.....	10.1	13.2	13.5	11.9	10.2	8.5	6.3	4.1	1.9
2006 5/.....	9.2	11.8	12.3	11.2	9.6	8.3	6.6	4.2	2.0
2005 5/.....	9.2	11.7	12.1	11.4	9.7	8.7	6.2	4.7	1.8
2004 5/.....	8.9	11.8	11.1	11.1	10.5	8.8	6.1	3.8	2.2
2003 5/.....	9.1	11.2	10.8	10.6	10.4	8.1	6.4	4.2	1.7
2002.....	9.0	10.5	10.5	11.2	10.0	9.0	6.5	4.3	1.9
2001.....	9.7	10.1	9.9	11.3	10.1	8.4	6.4	3.6	2.1
2000.....	9.2	9.7	10.4	11.9	10.4	8.6	6.7	4.1	2.2
1999.....	9.0	10.0	10.3	11.5	10.7	8.8	6.7	5.0	2.4

Non-Hispanic White, male

2007 5/.....	13.8	19.3	20.9	18.6	16.3	13.9	10.9	7.9	4.1
2006 5/.....	12.3	17.7	18.6	17.2	15.6	13.2	11.0	7.1	4.0
2005 5/.....	12.5	17.6	18.7	18.0	15.0	13.3	10.4	8.1	3.9
2004 5/.....	11.8	17.6	16.8	17.2	17.0	14.2	10.5	7.4	4.7
2003 5/.....	13.2	17.1	16.4	16.5	16.7	13.7	10.6	7.6	3.5
2002.....	13.0	16.0	15.9	17.8	15.6	14.1	10.5	7.7	4.1
2001.....	14.3	15.1	15.6	17.4	15.4	13.6	11.2	6.4	4.8
2000.....	14.1	15.0	15.8	18.4	16.5	14.3	12.6	7.3	5.1
1999.....	13.9	15.7	16.2	18.3	17.8	14.7	10.9	9.3	5.1

Non-Hispanic White, female

2007 5/.....	6.4	7.2	6.4	5.6	4.8	4.0	2.9	1.8	0.9
2006 5/.....	6.1	6.0	6.2	5.6	4.3	4.2	3.4	2.5	1.1
2005 5/.....	5.8	5.9	5.8	5.2	4.9	4.9	3.0	2.5	0.8
2004 5/.....	6.0	6.1	5.6	5.4	4.7	4.3	2.9	1.5	1.0
2003 5/.....	5.1	5.5	5.4	5.0	4.8	3.4	3.3	2.1	1.0
2002.....	5.0	5.1	5.3	5.0	5.0	4.8	3.6	2.1	1.0
2001.....	5.2	5.3	4.5	5.6	5.4	4.1	3.0	1.9	1.0
2000.....	4.4	4.4	5.3	5.8	5.1	4.0	2.5	2.3	1.0
1999.....	4.2	4.4	4.6	5.3	4.5	4.0	3.7	2.5	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

Non-Hispanic Black, both sexes

2007 5/.....	7.1	9.9	12.5	11.5	9.9	9.5	5.1	*	*
2006 5/.....	6.0	9.7	10.0	8.8	8.8	6.4	5.3	*	*
2005 5/.....	8.0	10.4	11.1	12.1	9.1	8.4	5.3	*	*
2004 5/.....	7.6	11.1	13.2	12.2	11.2	6.6	*	*	*
2003 5/.....	7.7	12.8	12.3	13.1	11.3	7.1	*	*	*
2002.....	8.5	12.4	12.4	12.1	9.6	7.6	5.8	*	*
2001.....	9.1	14.9	14.4	11.6	13.1	8.0	5.7	*	*
2000.....	11.3	13.6	14.0	12.8	12.0	8.6	6.3	*	*
1999.....	12.8	12.9	15.0	15.9	13.2	10.2	8.2	*	*

Non-Hispanic Black, male

2007 5/.....	9.4	16.0	19.8	19.9	18.7	18.7	11.4	*	*
2006 5/.....	8.8	14.3	16.4	13.0	16.7	9.8	10.1	*	*
2005 5/.....	12.1	15.5	17.0	21.0	15.4	15.3	11.9	*	*
2004 5/.....	10.9	16.0	21.4	19.8	20.0	13.6	*	*	*
2003 5/.....	12.2	20.8	19.7	21.0	18.7	12.8	*	*	*
2002.....	11.9	19.1	17.2	20.4	16.4	13.3	9.9	*	*
2001.....	13.3	22.5	24.0	18.8	23.4	15.5	*	*	*
2000.....	15.5	21.0	20.5	18.5	19.5	17.1	11.6	*	*
1999.....	19.2	20.4	25.3	23.7	24.7	18.3	14.5	*	*

Non-Hispanic Black, female

2007 5/.....	5.1	4.7	6.4	4.9	3.5	*	*	*	*
2006 5/.....	3.6	5.7	4.8	5.5	*	*	*	*	*
2005 5/.....	4.4	6.1	6.3	5.2	4.4	*	*	*	*
2004 5/.....	4.7	6.9	6.5	6.3	4.6	*	*	*	*
2003 5/.....	3.7	6.2	6.3	7.0	5.8	*	*	*	*
2002.....	5.5	6.7	8.6	5.6	4.6	*	*	*	*
2001.....	5.6	8.4	6.5	5.9	5.4	*	*	*	*
2000.....	7.7	7.3	8.7	8.4	6.5	*	*	*	*
1999.....	7.4	6.6	6.7	9.7	4.7	4.8	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/, both sexes

2007.....	5.4	8.8	10.9	11.8	14.7	18.9	22.4	23.0	17.7
2006.....	5.6	9.2	10.3	11.4	14.7	18.9	22.5	23.7	17.2
2005.....	5.7	8.7	10.6	12.0	15.0	20.0	23.3	23.1	17.8
2004.....	6.0	9.1	9.9	12.2	15.4	20.5	23.6	22.9	17.6
2003.....	6.5	9.3	10.4	12.7	17.0	22.9	24.5	24.1	18.2
2002.....	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001.....	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000.....	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
1999.....	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4

All origins 3/, male

2007.....	7.3	12.5	15.6	15.3	18.2	20.7	26.0	26.7	21.0
2006.....	7.8	13.4	14.4	15.0	17.2	21.3	24.9	26.9	21.4
2005.....	8.1	12.8	14.9	15.9	17.4	23.1	26.0	26.5	23.3
2004.....	8.6	13.3	14.1	15.2	17.9	23.8	27.1	26.0	21.0
2003.....	9.2	14.0	14.3	17.0	20.7	25.8	26.8	26.8	23.9
2002.....	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001.....	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000.....	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
1999.....	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0

All origins 3/, female

2007.....	3.5	5.3	6.6	8.5	11.7	17.5	19.7	20.6	16.1
2006.....	3.4	5.1	6.5	8.1	12.5	16.8	20.7	21.7	15.2
2005.....	3.5	4.7	6.5	8.5	13.0	17.4	21.4	21.0	15.3
2004.....	3.5	5.1	6.0	9.5	13.2	18.0	21.1	21.0	16.1
2003.....	3.8	4.8	6.6	8.9	13.8	20.5	22.8	22.5	15.8
2002.....	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001.....	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000.....	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
1999.....	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

Hispanic, both sexes

2007	6.0	11.7	12.6	19.0	24.5	35.1	38.8	37.2	28.6
2006	6.7	11.1	14.1	16.1	23.7	29.1	36.1	45.8	32.9
2005	7.5	11.2	17.2	19.0	26.4	31.4	44.9	39.1	30.4
2004	7.2	12.8	14.1	21.9	27.2	36.2	45.7	45.4	24.3
2003	8.2	12.1	14.6	20.3	27.1	41.4	50.7	43.4	26.9
2002	9.5	12.6	17.0	20.7	29.4	42.8	41.1	46.3	38.8
2001	9.0	13.3	15.0	21.0	28.9	38.8	42.8	44.7	30.6
2000	9.4	11.8	16.9	19.5	30.0	42.1	46.8	51.2	44.5
1999	9.1	10.5	14.1	21.1	32.1	40.2	43.9	47.0	40.0

Hispanic, male

2007	9.1	16.8	17.4	23.6	26.7	37.0	36.2	27.9	25.4
2006	10.2	16.0	20.2	20.5	23.7	26.3	31.4	37.7	33.8
2005	11.9	16.7	23.3	27.6	28.6	34.2	44.8	45.5	*
2004	10.1	19.6	19.9	26.7	26.9	36.0	42.0	43.4	*
2003	12.4	17.8	19.9	29.5	27.9	38.0	49.2	37.4	*
2002	15.4	19.4	24.4	22.8	35.2	40.4	41.6	54.9	*
2001	14.3	20.8	21.2	30.1	34.0	36.6	47.7	41.6	*
2000	14.4	18.7	22.8	23.9	35.1	38.9	48.0	42.7	50.4
1999	14.6	15.8	19.2	27.1	36.7	43.3	42.9	45.6	45.3

Hispanic, female

2007	2.8	6.6	8.0	14.8	22.6	33.6	40.7	43.4	30.3
2006	3.0	6.3	8.4	12.2	23.7	31.2	39.5	51.2	32.4
2005	2.9	5.7	11.6	11.4	24.6	29.3	45.0	34.8	33.9
2004	4.2	6.1	8.8	17.7	27.6	36.4	48.3	46.7	24.5
2003	3.8	6.5	9.8	12.2	26.4	43.9	51.7	47.4	26.6
2002	3.4	6.1	10.1	18.8	24.6	44.6	40.7	40.9	42.3
2001	3.5	6.2	9.4	13.2	24.7	40.5	39.2	46.6	33.5
2000	4.4	5.3	11.5	15.6	26.0	44.6	46.0	56.4	41.5
1999	3.7	5.5	9.6	15.9	28.4	37.9	44.6	47.8	37.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 4/, both sexes

2007.....	5.3	8.5	10.7	11.1	13.9	17.7	21.2	22.1	17.1
2006.....	5.4	8.9	9.9	10.9	13.9	18.1	21.6	22.4	16.4
2005.....	5.5	8.4	10.0	11.4	14.1	19.1	21.9	22.2	17.2
2004.....	5.8	8.7	9.6	11.4	14.4	19.4	22.3	21.7	17.3
2003.....	6.3	9.0	10.0	12.1	16.2	21.6	22.9	23.1	17.9
2002.....	6.3	8.5	9.7	11.9	16.2	20.2	25.0	24.5	18.5
2001.....	6.7	8.4	9.6	12.1	16.9	21.1	23.4	23.8	19.4
2000.....	6.6	7.5	9.6	12.7	16.5	20.5	23.7	24.1	19.9
1999.....	6.1	7.4	9.5	13.3	17.2	21.3	24.5	24.8	19.6

Non-Hispanic 4/, male

2007.....	7.0	11.9	15.3	14.6	17.5	19.5	25.2	26.6	20.7
2006.....	7.5	13.1	13.8	14.5	16.7	21.0	24.4	26.1	20.8
2005.....	7.5	12.3	14.2	14.9	16.5	22.3	24.8	25.3	23.3
2004.....	8.4	12.6	13.5	14.3	17.2	22.8	26.2	25.1	20.7
2003.....	8.8	13.5	13.8	15.9	20.1	25.0	25.6	26.2	23.7
2002.....	8.7	12.7	13.6	14.6	18.7	23.4	27.8	27.7	23.4
2001.....	9.5	12.4	12.9	15.1	20.8	25.0	24.4	27.7	26.4
2000.....	9.7	11.0	13.0	16.9	19.6	24.1	26.0	26.1	28.0
1999.....	9.3	11.1	13.3	16.5	21.1	24.4	27.3	28.4	26.2

Non-Hispanic 4/, female

2007.....	3.6	5.2	6.4	8.0	10.7	16.2	18.2	19.2	15.4
2006.....	3.4	5.0	6.3	7.7	11.5	15.7	19.5	20.1	14.3
2005.....	3.5	4.6	6.0	8.3	12.0	16.6	19.8	20.3	14.4
2004.....	3.4	5.0	5.8	8.8	12.0	16.6	19.5	19.7	15.7
2003.....	3.8	4.7	6.4	8.6	12.7	18.9	21.1	21.3	15.3
2002.....	3.9	4.6	6.2	9.5	14.1	17.7	23.0	22.5	16.4
2001.....	4.0	4.6	6.5	9.3	13.5	18.0	22.7	21.5	16.5
2000.....	3.5	4.3	6.5	9.0	14.0	17.6	22.1	22.9	16.7
1999.....	3.1	3.9	5.9	10.5	13.8	18.9	22.5	22.6	17.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic White, both sexes

2007 5/.....	5.4	8.4	10.3	11.3	14.5	18.5	22.2	23.3	17.8
2006 5/.....	5.6	8.9	9.8	11.0	14.4	19.3	22.4	23.5	16.9
2005 5/.....	5.5	8.1	9.8	11.3	14.7	20.3	22.6	22.9	17.7
2004 5/.....	5.9	8.5	9.3	11.4	14.8	20.4	23.6	22.7	17.7
2003 5/.....	6.2	8.7	9.7	12.3	16.7	22.8	24.3	24.3	18.6
2002.....	6.0	8.2	9.7	11.9	16.9	21.2	26.5	25.3	19.3
2001.....	6.3	8.0	9.2	12.3	17.5	21.9	24.5	25.0	20.4
2000.....	6.5	7.0	9.5	12.8	16.7	21.1	25.0	25.0	20.6
1999.....	5.6	7.2	9.1	13.2	17.6	22.1	25.5	25.9	20.2

Non-Hispanic White, male

2007 5/.....	7.3	11.8	14.5	14.8	17.9	20.4	26.1	28.0	22.0
2006 5/.....	7.8	12.9	13.5	14.3	17.2	22.2	25.0	27.6	21.5
2005 5/.....	7.4	12.1	13.7	14.5	17.0	23.6	25.3	25.7	24.2
2004 5/.....	8.3	12.2	13.1	14.2	17.4	23.9	27.6	26.3	21.7
2003 5/.....	8.9	13.0	13.2	16.0	20.8	26.3	26.9	27.3	24.5
2002.....	8.6	12.2	13.2	14.4	19.2	24.6	29.7	28.6	24.7
2001.....	8.9	11.6	12.1	14.9	21.4	25.9	25.4	29.3	28.1
2000.....	9.7	10.1	12.7	16.9	20.0	24.8	27.1	27.4	29.0
1999.....	8.5	10.7	12.8	16.4	21.6	25.2	28.4	29.8	27.5

Non-Hispanic White, female

2007 5/.....	3.6	5.0	6.2	8.0	11.4	16.9	19.3	20.3	15.7
2006 5/.....	3.3	5.1	6.2	7.8	12.0	16.8	20.4	20.9	14.8
2005 5/.....	3.5	4.2	6.1	8.3	12.7	17.5	20.6	21.1	14.8
2004 5/.....	3.4	4.8	5.5	8.7	12.5	17.6	20.6	20.4	16.0
2003 5/.....	3.5	4.5	6.3	8.8	13.1	19.9	22.4	22.4	16.0
2002.....	3.5	4.4	6.3	9.6	14.8	18.5	24.1	23.4	17.0
2001.....	3.8	4.4	6.3	9.8	14.2	18.5	23.8	22.4	17.2
2000.....	3.3	4.0	6.5	9.0	13.8	18.1	23.5	23.6	17.2
1999.....	2.8	3.7	5.7	10.3	14.0	19.5	23.5	23.5	17.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic Black, both sexes

2007 5/.....	5.5	11.0	15.5	12.3	12.1	13.2	13.2	9.3	8.4
2006 5/.....	5.1	11.7	13.5	12.1	12.3	9.9	14.3	9.2	10.1
2005 5/.....	6.7	11.7	13.8	13.7	11.3	12.3	14.6	14.9	10.7
2004 5/.....	7.1	12.5	13.1	13.1	12.1	13.8	11.1	11.4	11.2
2003 5/.....	7.6	13.2	14.4	12.2	14.4	14.9	10.6	10.6	11.0
2002.....	9.4	13.0	12.4	13.8	13.6	12.4	13.9	12.3	9.7
2001.....	10.3	12.3	13.9	12.2	14.1	15.0	14.9	11.6	7.8
2000.....	8.5	12.7	12.0	13.5	15.8	16.4	12.0	13.5	12.4
1999.....	10.0	10.2	13.4	15.5	15.0	14.9	15.1	13.5	12.3

Non-Hispanic Black, male

2007 5/.....	6.6	16.0	24.0	17.2	18.9	16.2	20.5	*	*
2006 5/.....	6.5	17.8	20.0	17.2	17.9	12.7	22.1	*	*
2005 5/.....	9.4	17.0	22.0	20.2	15.7	16.6	21.1	19.4	*
2004 5/.....	10.7	19.0	20.4	17.5	16.2	18.6	15.4	14.7	*
2003 5/.....	9.8	21.3	22.9	18.3	18.5	16.2	13.5	*	*
2002.....	12.1	19.8	20.1	18.9	18.7	15.4	12.2	*	*
2001.....	15.6	20.2	20.8	19.6	20.3	18.9	18.0	*	*
2000.....	11.8	20.2	18.5	19.3	18.9	20.8	16.4	*	*
1999.....	15.6	16.6	19.0	19.6	19.4	18.7	17.9	*	*

Non-Hispanic Black, female

2007 5/.....	4.6	6.8	8.5	8.5	7.1	11.2	8.8	8.7	9.0
2006 5/.....	4.0	6.4	8.1	8.1	8.2	8.0	9.6	9.3	9.0
2005 5/.....	4.3	7.3	7.1	8.6	8.0	9.3	10.7	12.5	10.0
2004 5/.....	3.9	6.9	7.1	9.6	9.1	10.6	8.6	9.7	10.8
2003 5/.....	5.7	6.4	7.4	7.4	11.5	14.1	8.9	9.5	8.3
2002.....	7.0	7.2	6.2	9.7	9.9	10.3	14.9	11.0	9.6
2001.....	5.7	5.6	8.3	6.4	9.4	12.4	13.0	12.9	*
2000.....	5.6	6.2	6.7	8.9	13.6	13.4	9.3	14.3	11.3
1999.....	5.2	4.8	8.8	12.3	11.8	12.4	13.4	12.3	12.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins 3/, both sexes

2007.....	0.2	0.4	0.6	0.8	1.9	3.2	5.7	10.4	23.3
2006.....	0.2	0.4	0.6	1.2	1.8	3.3	6.2	9.9	21.8
2005.....	0.2	0.4	0.5	0.9	1.7	3.3	5.8	11.1	22.9
2004.....	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
2003.....	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9

All origins 3/, male

2007.....	0.3	0.4	0.8	0.9	2.3	4.0	7.1	13.0	23.8
2006.....	0.2	0.5	0.6	1.2	2.3	4.0	7.6	12.5	21.7
2005.....	*	0.5	0.5	1.2	2.3	4.3	7.1	12.7	24.3
2004.....	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
2003.....	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7

All origins 3/, female

2007.....	0.2	0.4	0.5	0.8	1.6	2.5	4.7	8.8	23.1
2006.....	0.2	0.3	0.5	1.2	1.3	2.7	5.2	8.3	21.9
2005.....	0.2	0.3	0.6	0.7	1.2	2.4	4.8	10.1	22.3
2004.....	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
2003.....	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 4/, both sexes

2007.....	0.2	0.4	0.6	0.8	1.8	3.2	5.6	10.4	23.2
2006.....	0.2	0.4	0.6	1.2	1.8	3.3	6.1	9.9	21.9
2005.....	0.2	0.4	0.5	0.9	1.6	3.2	5.8	11.1	22.6
2004.....	0.3	0.4	0.6	1.1	1.8	3.2	5.6	10.4	23.6
2003.....	0.2	0.4	0.5	0.9	1.9	3.3	5.8	11.2	21.9
2002.....	0.2	0.4	0.6	1.0	2.0	3.5	5.5	10.7	22.8
2001.....	0.3	0.4	0.6	1.3	2.0	3.6	5.8	9.8	23.4
2000.....	0.2	0.3	0.6	1.0	1.8	2.9	5.5	10.1	24.6
1999.....	0.3	0.4	0.7	1.0	1.8	3.3	5.5	10.7	23.6

Non-Hispanic 4/, male

2007.....	0.2	0.4	0.7	0.9	2.1	4.0	7.0	13.0	23.6
2006.....	0.2	0.5	0.6	1.2	2.4	4.0	7.5	12.6	22.0
2005.....	*	0.5	0.5	1.2	2.2	4.2	7.1	12.5	24.0
2004.....	0.3	0.4	0.7	1.2	2.2	3.9	6.7	12.6	25.8
2003.....	*	0.4	0.6	1.1	2.1	4.2	7.4	13.5	23.0
2002.....	0.3	0.5	0.8	1.1	2.3	3.9	7.3	13.6	23.9
2001.....	0.4	0.4	0.8	1.5	2.1	4.2	7.5	11.9	26.6
2000.....	*	0.3	0.7	1.1	2.1	3.5	7.0	12.6	27.7
1999.....	0.3	0.3	0.9	1.2	2.0	4.2	6.6	13.1	27.5

Non-Hispanic 4/, female

2007.....	0.2	0.4	0.5	0.8	1.6	2.5	4.6	8.7	23.0
2006.....	0.2	0.3	0.5	1.2	1.3	2.7	5.1	8.2	21.9
2005.....	0.2	0.2	0.5	0.7	1.1	2.3	4.9	10.2	21.9
2004.....	0.2	0.4	0.5	1.0	1.4	2.7	4.8	9.0	22.6
2003.....	0.2	0.4	0.4	0.7	1.8	2.6	4.6	9.8	21.4
2002.....	*	0.3	0.5	0.9	1.7	3.2	4.3	8.9	22.4
2001.....	0.3	0.3	0.4	1.2	1.8	3.1	4.6	8.6	22.1
2000.....	*	0.4	0.5	0.9	1.5	2.5	4.5	8.6	23.4
1999.....	0.2	0.5	0.6	0.8	1.6	2.7	4.7	9.2	22.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.3	0.6	0.7	1.8	3.2	5.4	10.5	23.9
2006 5/.....	0.2	0.3	0.5	1.1	1.5	3.0	5.9	9.9	22.4
2005 5/.....	*	0.3	0.5	0.9	1.4	3.0	5.7	11.1	22.7
2004 5/.....	0.2	0.3	0.5	1.0	1.7	3.0	5.4	10.3	24.0
2003 5/.....	0.2	0.3	0.4	0.8	1.8	3.4	5.7	11.2	22.0
2002.....	0.2	0.3	0.6	0.9	2.0	3.4	5.3	10.6	23.4
2001.....	0.3	0.3	0.5	1.2	1.7	3.4	5.7	9.6	23.6
2000.....	0.1	0.3	0.5	0.9	1.7	2.6	5.5	9.9	25.3
1999.....	0.2	0.4	0.6	0.8	1.5	3.2	5.5	10.7	24.0

Non-Hispanic White, male

2007 5/.....	*	0.3	0.6	0.9	2.2	3.9	7.2	13.1	24.2
2006 5/.....	*	0.4	0.5	1.1	2.1	3.7	7.4	12.8	23.2
2005 5/.....	*	0.4	0.5	1.1	1.9	4.0	7.1	12.9	24.5
2004 5/.....	0.2	0.3	0.7	1.2	2.1	3.8	6.6	12.7	27.0
2003 5/.....	*	0.3	0.5	1.0	2.0	4.3	7.4	13.4	23.3
2002.....	0.3	0.4	0.7	1.0	2.4	3.8	7.3	13.5	24.5
2001.....	0.3	0.4	0.7	1.4	1.8	4.1	7.5	11.8	26.4
2000.....	*	*	0.7	1.0	2.1	3.2	7.1	12.8	28.9
1999.....	*	0.3	0.7	1.1	1.8	4.1	6.7	13.6	27.8

Non-Hispanic White, female

2007 5/.....	*	0.3	0.5	0.6	1.4	2.5	4.1	8.8	23.8
2006 5/.....	*	*	0.5	1.1	1.0	2.3	4.9	8.1	22.0
2005 5/.....	*	*	0.4	0.7	0.8	2.1	4.7	9.9	21.8
2004 5/.....	*	0.3	0.4	0.8	1.3	2.4	4.6	8.8	22.7
2003 5/.....	*	0.3	0.3	0.6	1.6	2.7	4.6	9.8	21.4
2002.....	*	0.3	0.4	0.8	1.6	3.0	3.9	8.8	22.9
2001.....	*	*	0.4	1.1	1.6	2.9	4.4	8.3	22.4
2000.....	*	0.3	0.4	0.8	1.3	2.2	4.3	8.2	23.9
1999.....	*	0.4	0.5	0.6	1.3	2.5	4.7	9.0	22.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic Black, both sexes

2007 5/.....	*	*	1.1	1.5	2.5	3.3	8.2	9.1	18.2
2006 5/.....	*	1.0	1.1	1.9	3.5	4.8	9.3	9.2	17.5
2005 5/.....	*	*	*	*	3.8	4.8	7.2	12.7	21.0
2004 5/.....	*	1.0	1.3	2.0	2.3	5.2	7.1	12.4	19.6
2003 5/.....	*	1.0	*	*	3.1	2.8	6.1	11.1	21.1
2002.....	*	*	*	*	2.2	4.7	7.7	12.5	15.1
2001.....	*	*	1.5	2.0	3.5	6.3	6.2	11.6	20.0
2000.....	*	*	*	*	2.8	4.9	6.5	11.2	18.1
1999.....	*	*	*	2.0	3.3	4.1	4.7	11.4	15.5

Non-Hispanic Black, male

2007 5/.....	*	*	*	*	*	*	*	*	20.1
2006 5/.....	*	*	*	*	*	*	10.6	*	*
2005 5/.....	*	*	*	*	*	6.5	10.1	*	20.0
2004 5/.....	*	*	*	*	*	*	*	*	*
2003 5/.....	*	*	*	*	*	*	*	*	21.3
2002.....	*	*	*	*	*	*	*	16.4	*
2001.....	*	*	*	*	*	*	*	*	27.7
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	*	8.2	*	17.4
2006 5/.....	*	*	*	*	*	5.4	8.5	9.6	20.2
2005 5/.....	*	*	*	*	*	*	5.5	13.7	21.4
2004 5/.....	*	*	*	*	*	5.6	6.6	13.1	21.7
2003 5/.....	*	*	*	*	*	*	*	9.5	21.0
2002.....	*	*	*	*	*	5.5	7.8	10.6	16.3
2001.....	*	*	*	*	*	6.4	7.1	12.9	17.1
2000.....	*	*	*	*	*	4.8	6.4	11.7	19.6
1999.....	*	*	*	*	*	*	*	12.3	15.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes

2007.....	3.9	6.4	10.8	17.0	30.2	52.4	87.2	146.1	290.6
2006.....	3.8	6.7	10.7	18.1	30.5	50.3	86.4	144.0	290.5
2005.....	3.8	5.8	10.6	17.7	29.2	51.3	85.6	142.8	288.3
2004.....	3.7	6.4	10.6	17.6	29.6	49.0	85.4	139.0	286.6
2003.....	3.8	6.1	10.4	17.9	30.7	50.6	84.5	143.9	293.1
2002.....	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001.....	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000.....	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
1999.....	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9

All origins 3/, male

2007.....	4.3	7.5	12.5	19.7	33.7	60.8	104.7	189.6	376.3
2006.....	4.7	8.0	12.4	21.0	33.8	59.4	105.1	182.1	377.1
2005.....	4.8	7.2	12.4	21.1	31.4	59.1	102.7	183.8	376.5
2004.....	4.4	7.7	11.8	19.6	33.3	56.2	102.2	181.2	372.2
2003.....	4.5	7.6	12.2	19.6	34.6	58.1	102.9	187.9	387.6
2002.....	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001.....	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000.....	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
1999.....	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1

All origins 3/, female

2007.....	3.4	5.4	9.2	14.5	27.1	45.5	74.3	118.6	249.9
2006.....	2.9	5.5	9.0	15.5	27.6	42.8	72.7	120.2	250.0
2005.....	2.9	4.5	8.8	14.7	27.3	44.9	73.2	117.2	247.8
2004.....	3.1	5.2	9.4	15.9	26.4	43.2	73.3	113.0	248.1
2003.....	3.2	4.7	8.6	16.4	27.4	44.7	71.4	117.1	251.4
2002.....	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001.....	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000.....	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
1999.....	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Hispanic, both sexes

2007.....	3.3	6.8	11.0	17.7	33.4	54.9	84.7	118.2	190.2
2006.....	3.5	6.7	11.3	18.9	34.6	47.6	78.9	130.6	206.2
2005.....	3.0	5.0	10.8	20.5	29.4	49.8	82.2	117.0	193.4
2004.....	2.6	7.1	11.2	18.9	34.5	49.8	74.2	119.7	183.6
2003.....	3.2	6.3	12.0	19.7	33.0	48.0	80.7	105.7	233.2
2002.....	2.7	6.3	10.5	17.7	32.9	40.3	74.9	111.8	201.2
2001.....	2.7	5.5	11.1	15.1	34.6	49.9	83.8	113.2	222.0
2000.....	2.5	4.7	9.1	19.3	32.5	43.8	71.9	108.1	229.6
1999.....	3.0	4.4	10.5	18.5	29.7	43.1	77.1	109.8	232.2

Hispanic, male

2007.....	3.5	9.2	12.0	21.7	37.5	63.1	98.2	149.5	223.6
2006.....	4.7	8.2	12.2	21.1	41.5	57.2	89.1	153.1	241.3
2005.....	4.1	6.6	12.8	24.9	29.5	57.2	98.8	135.7	259.0
2004.....	3.2	8.1	13.5	22.1	34.6	57.5	82.9	152.9	231.2
2003.....	3.6	8.4	12.8	20.9	36.1	50.8	91.8	134.1	297.1
2002.....	2.8	7.6	11.7	19.8	38.9	43.4	92.1	147.6	235.9
2001.....	2.8	5.1	12.6	18.7	35.4	53.5	102.3	131.2	281.9
2000.....	3.6	5.6	9.9	20.1	33.6	50.6	85.6	132.5	292.2
1999.....	3.6	5.4	13.4	23.5	32.5	47.9	103.8	124.2	336.4

Hispanic, female

2007.....	3.1	4.4	10.0	14.2	29.9	48.5	74.9	97.2	172.1
2006.....	2.3	5.3	10.4	17.0	28.9	40.1	71.5	115.5	187.5
2005.....	1.9	3.3	8.8	16.6	29.3	44.1	70.2	104.5	159.0
2004.....	2.0	6.0	8.9	16.1	34.5	43.8	67.9	97.8	159.2
2003.....	2.9	4.3	11.3	18.7	30.4	45.9	72.9	87.3	200.8
2002.....	2.7	5.1	9.3	15.9	27.9	37.9	62.6	89.1	184.0
2001.....	2.5	5.8	9.8	12.0	33.9	47.1	70.7	102.1	192.7
2000.....	*	3.9	8.3	18.6	31.7	38.7	62.2	93.2	198.8
1999.....	2.4	3.5	7.9	14.1	27.5	39.5	58.4	101.3	182.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 4/, both sexes

2007.....	3.9	6.4	10.8	16.9	29.9	52.1	87.3	147.7	295.6
2006.....	3.8	6.7	10.6	18.0	30.1	50.3	86.7	144.6	294.4
2005.....	3.9	5.9	10.5	17.5	29.1	51.3	85.8	144.0	292.5
2004.....	3.8	6.3	10.5	17.5	29.2	48.8	85.9	139.5	290.9
2003.....	3.9	6.1	10.2	17.7	30.5	50.7	84.5	145.4	294.9
2002.....	3.8	5.8	9.9	16.7	29.1	50.3	85.7	143.2	291.2
2001.....	3.7	5.8	9.8	17.0	31.5	49.1	82.0	136.3	289.6
2000.....	3.5	5.5	9.4	16.6	29.2	47.2	79.6	133.4	279.0
1999.....	3.1	5.2	8.8	15.5	28.0	46.8	77.6	127.7	269.2

Non-Hispanic 4/, male

2007.....	4.4	7.3	12.5	19.5	33.3	60.4	105.1	192.0	384.6
2006.....	4.7	8.0	12.4	21.0	33.1	59.5	105.9	183.5	384.0
2005.....	4.9	7.2	12.4	20.7	31.5	59.1	102.8	186.4	382.0
2004.....	4.5	7.6	11.6	19.3	33.2	55.9	103.1	182.2	378.6
2003.....	4.6	7.5	12.1	19.4	34.3	58.4	103.2	190.1	390.8
2002.....	4.5	6.8	10.9	18.4	33.1	60.5	106.0	183.7	391.0
2001.....	4.5	6.8	10.8	18.1	36.5	57.0	100.3	178.2	397.6
2000.....	4.5	6.4	10.8	18.3	32.7	54.3	100.3	174.1	384.0
1999.....	3.8	5.8	9.6	17.3	30.8	55.1	98.8	170.2	370.9

Non-Hispanic 4/, female

2007.....	3.5	5.4	9.1	14.5	26.9	45.3	74.2	119.8	253.6
2006.....	3.0	5.5	8.9	15.3	27.4	42.9	72.6	120.4	252.7
2005.....	3.0	4.6	8.8	14.5	27.1	44.9	73.4	117.7	251.6
2004.....	3.2	5.1	9.4	15.9	25.7	43.1	73.5	113.4	251.6
2003.....	3.2	4.7	8.4	16.1	27.2	44.5	71.1	118.1	252.7
2002.....	3.1	4.9	9.1	15.0	25.6	42.2	71.3	118.6	248.2
2001.....	2.9	4.8	8.8	16.0	27.1	42.8	69.1	111.2	244.6
2000.....	2.6	4.7	8.2	15.1	26.2	41.6	65.2	109.4	236.6
1999.....	2.4	4.6	8.1	14.0	25.7	40.3	62.9	102.8	228.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic White, both sexes

2007 5/.....	2.6	4.5	7.9	12.6	24.1	44.2	79.8	141.2	287.1
2006 5/.....	2.7	4.4	7.7	13.7	23.5	42.4	78.9	136.4	284.8
2005 5/.....	2.6	4.2	7.3	13.1	23.3	43.8	78.1	136.1	284.0
2004 5/.....	2.5	4.1	7.4	13.0	23.3	42.3	78.1	132.6	282.1
2003 5/.....	2.5	4.0	7.4	13.5	24.4	42.9	77.0	137.2	285.9
2002.....	2.6	3.6	7.0	12.6	23.5	43.1	78.0	136.0	281.1
2001.....	2.3	3.9	6.8	12.5	25.0	42.4	74.4	128.0	279.7
2000.....	2.2	3.6	6.6	12.5	23.0	40.2	72.7	125.6	269.4
1999.....	1.8	3.4	6.2	11.6	22.4	40.1	70.5	118.6	255.3

Non-Hispanic White, male

2007 5/.....	3.1	5.2	9.5	14.6	26.9	52.8	98.8	187.1	382.0
2006 5/.....	3.4	5.3	9.1	16.7	26.4	51.8	98.2	176.0	380.7
2005 5/.....	3.3	5.3	8.4	16.0	26.7	52.2	95.3	180.3	379.1
2004 5/.....	2.9	4.9	8.6	14.5	26.8	49.6	96.1	177.2	373.0
2003 5/.....	3.0	4.7	8.7	14.9	28.2	50.3	96.8	181.7	386.3
2002.....	3.0	4.2	7.6	13.8	27.5	53.2	99.0	178.5	382.6
2001.....	2.7	4.8	7.8	13.9	29.9	50.9	94.0	170.0	392.6
2000.....	2.8	4.2	7.7	14.3	27.3	48.4	93.5	167.6	378.9
1999.....	2.2	3.7	7.2	13.2	25.5	48.8	90.8	160.8	359.6

Non-Hispanic White, female

2007 5/.....	2.2	3.8	6.4	10.7	21.6	37.0	65.4	111.9	242.2
2006 5/.....	2.0	3.5	6.5	10.9	20.9	34.6	64.5	111.5	240.1
2005 5/.....	2.0	3.2	6.2	10.4	20.2	36.7	65.4	108.5	240.5
2004 5/.....	2.1	3.4	6.3	11.6	20.2	36.2	64.9	104.9	241.4
2003 5/.....	2.0	3.3	6.0	12.2	21.0	36.9	62.7	109.8	241.7
2002.....	2.1	3.1	6.4	11.6	20.1	34.8	63.0	110.0	237.3
2001.....	1.8	3.1	5.9	11.2	20.6	35.5	60.5	102.6	232.7
2000.....	1.7	3.1	5.6	10.9	19.3	33.6	58.2	100.6	225.2
1999.....	1.5	3.0	5.3	10.3	19.8	33.1	56.2	93.7	213.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic Black, both sexes

2007 5/.....	12.4	20.1	32.9	55.6	82.6	126.7	178.2	240.0	443.9
2006 5/.....	11.3	23.1	33.4	56.7	88.2	125.1	181.2	259.7	449.4
2005 5/.....	12.4	18.5	36.8	57.1	81.5	127.0	179.9	258.4	429.8
2004 5/.....	12.9	22.1	36.4	58.1	80.8	116.6	178.3	244.3	437.2
2003 5/.....	13.0	21.4	35.1	55.1	85.7	130.7	173.7	263.6	446.5
2002.....	12.0	22.1	34.4	51.4	80.3	128.1	179.7	244.8	444.0
2001.....	12.9	20.3	33.3	54.2	90.4	121.3	174.0	250.8	432.9
2000.....	13.1	20.1	33.0	51.3	84.6	121.2	167.2	245.1	410.2
1999.....	12.1	20.1	29.6	47.9	78.9	118.2	165.6	246.3	436.8

Non-Hispanic Black, male

2007 5/.....	13.5	23.8	37.8	67.3	97.8	140.5	204.2	283.6	500.9
2006 5/.....	14.1	28.5	41.3	66.1	101.3	138.7	215.0	318.8	496.6
2005 5/.....	16.3	23.1	47.2	69.9	81.3	135.9	213.1	307.2	487.2
2004 5/.....	15.8	28.6	40.6	65.8	94.5	130.3	202.9	287.9	520.3
2003 5/.....	15.3	29.8	44.3	63.7	96.4	159.1	194.8	335.8	526.3
2002.....	15.0	27.4	41.0	62.3	90.4	150.9	208.4	284.7	568.9
2001.....	17.0	23.6	38.6	56.2	103.4	132.6	190.6	312.3	517.4
2000.....	18.0	24.3	38.3	54.9	87.9	128.5	201.4	292.7	496.7
1999.....	15.9	24.1	29.8	55.5	82.3	131.9	210.3	325.9	532.4

Non-Hispanic Black, female

2007 5/.....	11.4	16.9	28.8	46.5	71.5	117.3	162.6	217.1	420.3
2006 5/.....	8.9	18.4	26.9	49.4	78.6	115.8	161.1	228.5	430.1
2005 5/.....	9.1	14.7	28.3	47.1	81.6	120.9	160.2	232.6	406.7
2004 5/.....	10.4	16.6	33.0	52.1	70.7	107.3	163.7	221.4	404.4
2003 5/.....	11.1	14.3	27.6	48.4	77.7	111.6	161.2	225.9	415.6
2002.....	9.4	17.5	28.9	42.8	72.9	112.9	162.6	224.0	396.3
2001.....	9.3	17.5	28.9	52.6	80.8	113.8	164.1	219.5	401.7
2000.....	8.9	16.4	28.7	48.5	82.2	116.4	146.6	221.2	378.1
1999.....	8.9	16.8	29.4	42.0	76.4	109.1	138.6	206.8	401.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All origins 3/, both sexes

2007.....	*	*	*	*	*	*	0.3	0.7	0.9
2006.....	*	*	*	*	*	*	*	0.4	0.6
2005.....	*	*	*	*	*	*	0.3	*	0.8
2004.....	*	*	*	*	*	*	0.3	0.4	0.8
2003.....	*	*	*	*	*	*	*	0.6	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	0.4	0.4	0.9
2000.....	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	0.3	0.5	0.5	0.8

All origins 3/, male

2007.....	*	*	*	*	*	*	*	1.0	1.2
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	*

All origins 3/, female

2007.....	*	*	*	*	*	*	*	*	0.7
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	0.7
2004.....	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	0.5	*	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Non-Hispanic 4/, both sexes

2007.....	*	*	*	*	*	*	0.3	0.6	0.9
2006.....	*	*	*	*	*	*	*	0.4	0.7
2005.....	*	*	*	*	*	*	0.3	*	0.7
2004.....	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	*	*	*	*	*	*	*	0.6	0.5
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	0.4	*	0.8
2000.....	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	0.3	0.5	0.5	0.8

Non-Hispanic 4/, male

2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2007.....	*	*	*	*	*	*	*	*	0.8
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	0.7
2004.....	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	0.5	*	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins 3/, both sexes

2007.....	0.3	0.3	0.4	0.9	1.4	2.7	5.1	9.8	22.3
2006.....	0.1	0.3	0.4	0.6	1.0	1.8	3.2	6.0	13.4
2005.....	*	0.2	0.3	0.4	0.5	0.9	1.6	2.4	5.4
2004.....	*	0.1	0.2	0.3	0.4	0.6	1.0	1.6	2.8
2003.....	*	0.2	0.1	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	0.1	0.1	0.2	*	0.3	0.6	1.2	1.7	2.8
2001.....	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.4	2.0	3.2

All origins 3/, male

2007.....	0.3	0.4	0.6	1.1	1.7	3.4	6.1	12.3	29.9
2006.....	*	0.3	0.4	0.8	1.3	1.8	4.3	7.7	17.2
2005.....	*	0.2	0.3	0.4	0.6	1.1	1.9	3.2	7.5
2004.....	*	*	0.2	0.3	0.4	0.7	1.3	2.0	4.0
2003.....	*	0.2	*	*	0.5	0.6	1.3	2.0	4.1
2002.....	*	*	0.3	*	*	0.9	1.6	2.1	3.8
2001.....	*	*	*	*	0.5	1.0	1.3	2.3	3.7
2000.....	*	*	*	*	0.5	0.8	1.5	2.1	4.5
1999.....	*	*	*	*	0.5	0.6	1.9	2.7	5.0

All origins 3/, female

2007.....	0.3	0.2	0.3	0.7	1.2	2.1	4.3	8.1	18.7
2006.....	*	0.3	0.4	0.4	0.8	1.7	2.3	5.0	11.6
2005.....	*	*	*	0.3	0.5	0.8	1.4	1.9	4.4
2004.....	*	*	*	*	*	0.5	0.8	1.4	2.2
2003.....	*	*	*	*	*	0.8	0.9	1.4	2.2
2002.....	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	0.5	0.5	0.7	1.0	2.4
2000.....	*	*	*	*	*	0.6	0.8	1.8	2.1
1999.....	*	*	*	*	0.4	0.7	1.1	1.5	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic 4/, both sexes

2007.....	0.3	0.3	0.4	0.9	1.4	2.7	5.0	9.7	22.8
2006.....	0.1	0.3	0.4	0.6	1.0	1.8	3.2	6.0	13.6
2005.....	*	0.2	0.3	0.4	0.6	0.9	1.6	2.4	5.5
2004.....	*	0.1	0.2	0.3	0.4	0.6	1.1	1.7	2.7
2003.....	*	0.2	*	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	0.1	0.1	0.2	*	0.3	0.7	1.1	1.8	2.8
2001.....	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	*	*	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.5	2.0	3.2

Non-Hispanic 4/, male

2007.....	0.3	0.4	0.6	1.0	1.6	3.4	6.1	12.2	31.0
2006.....	*	0.3	0.4	0.8	1.3	1.8	4.3	7.7	17.4
2005.....	*	0.2	0.3	0.4	0.6	1.0	2.0	3.3	7.5
2004.....	*	*	*	*	*	0.7	1.4	2.1	3.8
2003.....	*	0.2	*	*	0.5	0.6	1.3	1.9	4.3
2002.....	*	*	*	*	*	1.0	1.5	2.1	3.8
2001.....	*	*	*	*	0.5	1.1	1.3	2.3	3.7
2000.....	*	*	*	*	0.6	0.8	1.5	2.1	4.2
1999.....	*	*	*	*	*	0.7	2.0	2.8	5.1

Non-Hispanic 4/, female

2007.....	0.3	*	0.3	0.8	1.2	2.2	4.2	8.2	18.9
2006.....	*	0.3	0.3	0.4	0.8	1.8	2.4	4.9	11.9
2005.....	*	*	*	0.3	0.5	0.8	1.3	1.8	4.5
2004.....	*	*	*	*	*	0.5	0.9	1.4	2.2
2003.....	*	*	*	*	*	0.8	0.8	1.4	2.2
2002.....	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	0.5	0.5	0.7	1.1	2.4
2000.....	*	*	*	*	*	0.6	0.8	1.8	2.2
1999.....	*	*	*	*	*	0.7	1.1	1.5	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic White, both sexes

2007 5/.....	0.2	0.2	0.3	0.7	1.1	2.3	4.6	8.8	21.4
2006 5/.....	*	0.2	0.3	0.4	0.7	1.6	2.9	5.4	12.7
2005 5/.....	*	*	0.2	0.3	0.5	0.7	1.5	2.3	5.2
2004 5/.....	*	*	*	0.2	0.3	0.6	1.0	1.7	2.7
2003 5/.....	*	*	*	*	0.4	0.6	0.9	1.5	2.7
2002.....	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001.....	*	*	*	*	0.4	0.7	0.9	1.5	2.8
2000.....	*	*	0.2	*	0.4	0.7	1.0	1.8	2.4
1999.....	*	*	*	0.2	0.3	0.6	1.4	1.9	2.9

Non-Hispanic White, male

2007 5/.....	*	0.3	0.5	0.8	1.3	3.0	5.7	11.4	29.7
2006 5/.....	*	*	*	0.6	0.9	1.7	3.9	7.2	16.2
2005 5/.....	*	*	*	*	0.6	0.9	1.8	3.2	7.3
2004 5/.....	*	*	*	*	*	*	1.3	2.2	3.6
2003 5/.....	*	*	*	*	*	*	1.1	1.8	4.0
2002.....	*	*	*	*	*	1.0	1.3	2.1	3.8
2001.....	*	*	*	*	*	1.1	1.2	2.3	4.0
2000.....	*	*	*	*	*	0.8	1.4	2.2	3.7
1999.....	*	*	*	*	*	0.6	2.0	2.7	4.6

Non-Hispanic White, female

2007 5/.....	*	*	*	0.6	0.9	1.7	3.8	7.1	17.5
2006 5/.....	*	*	0.3	*	0.5	1.5	2.2	4.2	11.1
2005 5/.....	*	*	*	*	*	0.6	1.2	1.8	4.3
2004 5/.....	*	*	*	*	*	*	0.8	1.4	2.3
2003 5/.....	*	*	*	*	*	0.7	0.8	1.3	2.1
2002.....	*	*	*	*	*	*	0.7	1.5	2.0
2001.....	*	*	*	*	*	*	0.7	1.0	2.4
2000.....	*	*	*	*	*	0.6	0.7	1.6	1.9
1999.....	*	*	*	*	*	0.7	1.0	1.4	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

All origins 3/, both sexes

2007.....	3.6	6.1	10.3	16.0	28.6	49.4	81.7	135.6	267.3
2006.....	3.7	6.4	10.2	17.5	29.3	48.4	83.0	137.5	276.4
2005.....	3.8	5.6	10.3	17.3	28.6	50.2	83.7	140.0	282.1
2004.....	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
2003.....	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8

All origins 3/, male

2007.....	4.0	7.1	11.9	18.5	31.9	57.1	98.1	176.1	345.0
2006.....	4.5	7.7	12.0	20.2	32.4	57.4	100.4	174.0	358.8
2005.....	4.7	6.9	12.0	20.5	30.7	57.8	100.4	180.3	368.0
2004.....	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
2003.....	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6

All origins 3/, female

2007.....	3.2	5.1	8.9	13.7	25.8	43.2	69.7	110.1	230.4
2006.....	2.8	5.2	8.6	15.1	26.6	41.0	70.2	114.8	237.8
2005.....	2.8	4.4	8.6	14.3	26.7	44.0	71.6	115.0	242.6
2004.....	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
2003.....	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

Hispanic, both sexes

2007.....	3.1	6.3	10.6	16.7	31.6	52.0	78.4	106.4	174.8
2006.....	3.4	6.4	10.9	18.1	33.2	45.9	76.1	123.7	197.2
2005.....	2.9	4.8	10.5	20.1	29.1	48.3	79.9	114.5	187.4
2004.....	2.6	6.8	10.7	18.7	33.8	49.0	73.2	119.0	179.3
2003.....	3.1	6.1	11.6	19.4	32.4	47.3	79.4	103.3	230.6
2002.....	2.7	6.3	10.1	17.3	32.3	40.1	73.0	110.1	198.4
2001.....	2.4	5.3	10.8	14.6	33.9	48.7	82.7	111.7	217.6
2000.....	2.4	4.4	8.9	18.9	32.0	43.0	70.7	105.3	225.6
1999.....	2.8	4.4	10.2	17.8	28.2	42.2	75.5	107.5	228.0

Hispanic, male

2007.....	3.3	8.4	11.5	20.0	35.1	58.7	91.2	131.7	208.4
2006.....	4.5	8.0	11.9	19.9	40.4	54.9	85.0	146.6	227.8
2005.....	4.0	6.4	12.4	24.5	29.5	54.8	97.8	134.0	249.1
2004.....	3.2	8.0	12.7	21.6	33.6	56.7	81.8	151.9	224.2
2003.....	3.3	8.0	12.3	20.9	35.4	50.4	90.5	129.9	295.5
2002.....	2.8	7.6	11.0	19.2	38.6	43.4	88.9	145.3	232.6
2001.....	2.8	4.8	12.2	17.9	34.7	52.6	100.9	128.6	278.1
2000.....	3.4	5.3	9.6	19.3	33.6	49.6	84.2	129.6	280.1
1999.....	3.5	5.4	12.9	22.6	30.9	47.9	102.2	122.6	330.0

Hispanic, female

2007.....	2.8	4.1	9.7	13.8	28.7	46.8	69.2	89.5	156.7
2006.....	2.3	4.9	9.9	16.5	27.2	38.9	69.7	108.5	180.9
2005.....	1.9	3.2	8.8	16.2	28.8	43.2	67.1	101.6	155.1
2004.....	2.0	5.7	8.8	16.1	34.0	43.1	67.1	97.2	156.3
2003.....	2.9	4.3	10.9	18.0	29.9	44.9	71.6	85.9	197.7
2002.....	2.5	5.0	9.3	15.7	27.1	37.5	61.7	87.6	181.5
2001.....	2.1	5.7	9.6	11.8	33.3	45.7	69.7	101.3	188.0
2000.....	*	3.6	8.1	18.6	30.8	37.9	61.2	90.5	198.8
1999.....	*	3.5	7.7	13.6	25.9	37.9	56.7	98.5	178.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

Non-Hispanic 4/, both sexes

2007.....	3.6	6.0	10.3	15.9	28.3	49.2	82.0	137.3	271.9
2006.....	3.7	6.4	10.1	17.4	28.9	48.4	83.2	138.2	280.0
2005.....	3.9	5.7	10.2	17.0	28.5	50.2	83.9	141.2	286.2
2004.....	3.7	6.1	10.3	17.2	28.6	48.0	84.4	137.3	287.2
2003.....	3.8	5.9	10.0	17.4	29.9	49.8	83.1	143.1	291.4
2002.....	3.6	5.6	9.7	16.4	28.6	49.5	84.2	140.7	287.4
2001.....	3.6	5.6	9.6	16.7	30.8	48.2	80.6	134.3	285.9
2000.....	3.4	5.3	9.2	16.3	28.6	46.2	78.3	130.6	274.9
1999.....	3.0	5.0	8.6	15.1	27.6	45.7	75.6	125.0	265.0

Non-Hispanic 4/, male

2007.....	4.0	6.9	11.9	18.3	31.5	56.8	98.5	178.8	352.4
2006.....	4.5	7.6	11.9	20.2	31.6	57.5	101.3	175.2	365.5
2005.....	4.8	7.0	12.0	20.2	30.8	57.8	100.4	182.7	373.6
2004.....	4.4	7.4	11.4	18.9	32.6	55.0	101.3	179.3	374.0
2003.....	4.4	7.3	12.0	19.1	33.6	57.6	101.7	187.5	386.2
2002.....	4.3	6.6	10.6	18.2	32.6	59.4	104.1	181.0	386.2
2001.....	4.4	6.6	10.6	17.8	35.9	55.8	98.7	175.3	392.7
2000.....	4.3	6.1	10.5	17.9	31.9	53.2	98.4	171.3	377.7
1999.....	3.6	5.5	9.4	16.9	30.3	54.0	96.3	166.3	364.4

Non-Hispanic 4/, female

2007.....	3.2	5.2	8.7	13.7	25.5	42.9	69.7	111.2	233.9
2006.....	2.9	5.2	8.5	14.9	26.5	41.0	70.0	115.0	240.2
2005.....	3.0	4.5	8.5	14.1	26.5	44.0	71.8	115.5	246.4
2004.....	3.1	4.9	9.2	15.6	25.1	42.4	72.2	111.5	248.5
2003.....	3.2	4.6	8.2	15.8	26.7	43.5	69.9	116.1	249.7
2002.....	3.0	4.7	8.8	14.7	25.2	41.6	70.1	116.3	244.8
2001.....	2.8	4.7	8.6	15.8	26.5	42.1	68.0	109.7	241.5
2000.....	2.5	4.6	8.0	14.8	25.8	40.7	64.3	106.7	233.4
1999.....	2.4	4.4	7.8	13.5	25.2	39.2	61.2	100.9	225.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

Non-Hispanic White, both sexes

2007 5/.....	2.4	4.2	7.5	11.8	22.9	41.7	74.8	131.8	264.8
2006 5/.....	2.6	4.2	7.4	13.2	22.7	40.8	75.7	130.6	271.3
2005 5/.....	2.6	4.1	7.0	12.6	22.7	42.8	76.2	133.4	278.1
2004 5/.....	2.4	4.0	7.2	12.7	22.9	41.5	76.7	130.4	278.6
2003 5/.....	2.4	3.9	7.2	13.2	23.9	42.1	75.8	135.1	282.6
2002.....	2.4	3.5	6.8	12.4	23.1	42.2	76.7	133.7	277.6
2001.....	2.2	3.8	6.7	12.3	24.4	41.4	73.1	126.1	276.1
2000.....	2.1	3.6	6.4	12.2	22.6	39.2	71.5	122.8	265.7
1999.....	1.7	3.2	6.0	11.3	22.0	39.1	68.5	116.2	251.6

Non-Hispanic White, male

2007 5/.....	2.8	4.9	9.0	13.6	25.5	49.6	92.6	174.8	351.1
2006 5/.....	3.3	5.1	8.7	16.1	25.4	50.0	93.9	168.4	363.4
2005 5/.....	3.2	5.1	8.1	15.5	26.1	51.0	93.0	176.7	370.9
2004 5/.....	2.8	4.7	8.4	14.2	26.3	48.7	94.5	174.4	368.8
2003 5/.....	2.9	4.5	8.6	14.7	27.6	49.5	95.5	179.5	381.9
2002.....	2.9	4.1	7.3	13.5	26.9	52.1	97.4	175.9	377.8
2001.....	2.7	4.6	7.6	13.7	29.4	49.7	92.4	167.2	387.6
2000.....	2.6	4.1	7.4	14.0	26.6	47.3	91.7	164.6	373.4
1999.....	2.1	3.5	7.1	12.8	25.0	47.9	88.3	157.2	353.8

Non-Hispanic White, female

2007 5/.....	2.0	3.6	6.1	10.0	20.6	35.1	61.5	104.4	223.9
2006 5/.....	1.9	3.3	6.1	10.6	20.3	33.1	62.1	106.8	228.4
2005 5/.....	1.9	3.1	6.0	9.9	19.7	36.1	63.8	106.4	235.7
2004 5/.....	2.1	3.2	6.1	11.3	19.7	35.5	63.7	103.1	238.2
2003 5/.....	1.9	3.2	5.9	11.8	20.5	36.0	61.5	107.8	238.9
2002.....	2.0	2.9	6.2	11.3	19.6	34.0	61.8	107.9	234.3
2001.....	1.8	3.0	5.8	11.1	20.0	34.7	59.4	101.3	229.7
2000.....	1.6	3.0	5.4	10.6	19.0	32.7	57.3	98.0	222.2
1999.....	1.4	2.9	5.1	10.0	19.4	32.1	54.6	92.0	210.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

Non-Hispanic Black, both sexes

2007 5/.....	11.4	19.1	31.6	52.8	78.1	119.6	166.9	219.7	402.7
2006 5/.....	10.9	21.9	32.0	54.8	84.2	120.7	175.6	246.8	425.2
2005 5/.....	12.3	17.9	35.9	56.2	80.4	124.2	176.7	253.4	418.9
2004 5/.....	12.5	21.8	35.9	57.5	79.7	115.4	176.2	241.0	432.6
2003 5/.....	12.9	21.0	35.0	54.5	84.7	128.9	171.5	260.3	442.3
2002.....	11.6	21.5	33.9	51.1	79.9	127.3	177.1	241.6	436.5
2001.....	12.6	19.8	33.0	53.6	88.9	120.5	172.2	248.1	428.9
2000.....	12.7	19.2	32.6	50.6	83.2	120.1	164.9	241.7	404.2
1999.....	12.0	19.4	29.0	46.9	77.7	116.8	162.8	241.9	428.7

Non-Hispanic Black, male

2007 5/.....	12.6	22.4	36.1	63.6	92.2	131.7	191.5	260.7	452.0
2006 5/.....	13.5	27.2	39.1	63.3	96.0	135.2	207.2	302.7	462.7
2005 5/.....	16.1	22.4	46.1	68.6	80.6	133.6	207.6	300.8	473.9
2004 5/.....	15.3	28.4	39.9	64.7	93.5	128.7	200.1	284.9	514.2
2003 5/.....	14.9	29.2	44.3	62.9	94.9	157.0	191.1	330.5	521.0
2002.....	14.8	26.7	40.4	62.3	89.8	149.5	204.2	281.6	562.3
2001.....	16.6	22.8	38.2	55.4	101.4	131.5	189.1	309.9	514.0
2000.....	17.6	22.9	37.9	53.8	85.8	127.8	198.9	290.1	487.3
1999.....	15.7	23.3	29.3	54.4	81.3	130.2	206.9	319.7	519.4

Non-Hispanic Black, female

2007 5/.....	10.5	16.4	27.8	44.4	67.8	111.3	152.2	198.1	382.3
2006 5/.....	8.6	17.5	26.1	48.1	75.5	110.8	156.7	217.4	409.9
2005 5/.....	8.9	14.1	27.5	46.5	80.3	117.8	158.3	228.4	396.8
2004 5/.....	10.1	16.3	32.7	51.8	69.4	106.4	162.0	217.9	400.4
2003 5/.....	11.1	14.0	27.3	47.9	77.2	110.0	159.8	223.5	411.9
2002.....	8.8	17.0	28.7	42.3	72.5	112.4	160.9	220.8	388.4
2001.....	9.2	17.2	28.8	52.2	79.6	113.1	162.1	216.6	397.4
2000.....	8.5	16.0	28.3	48.1	81.2	115.0	144.3	217.3	373.3
1999.....	8.8	16.1	28.7	41.0	75.0	108.0	136.3	203.3	394.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

All origins 3/, both sexes

2007.....	0.1	0.1	0.2	0.2	0.3	0.6	0.9	1.8	3.5
2006.....	0.1	0.1	0.1	0.2	0.4	0.6	1.0	1.6	4.7
2005.....	0.1	0.1	0.1	0.2	0.4	0.7	1.3	2.5	5.2
2004.....	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.6	5.2
2003.....	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.8	5.9
2002.....	*	0.1	0.2	0.3	0.5	0.7	1.6	2.3	6.3
2001.....	0.1	0.1	0.2	0.2	0.4	0.6	1.7	2.4	6.0
2000.....	0.1	0.1	0.2	0.3	0.4	0.7	1.5	2.7	7.5
1999.....	0.1	0.2	0.2	0.3	0.5	0.9	1.5	2.7	7.5

All origins 3/, male

2007.....	*	*	*	*	*	0.5	0.8	1.4	3.3
2006.....	*	*	*	*	*	0.7	*	1.2	3.7
2005.....	*	*	*	*	*	0.7	0.9	2.5	3.6
2004.....	*	*	0.3	*	*	*	1.0	2.2	4.5
2003.....	*	*	*	*	*	*	0.8	2.4	5.4
2002.....	*	*	*	*	*	0.7	1.4	1.4	5.3
2001.....	*	*	*	*	*	0.5	1.2	2.1	4.3
2000.....	*	*	*	*	*	*	1.1	2.2	5.6
1999.....	*	*	*	*	*	0.6	1.2	2.6	6.8

All origins 3/, female

2007.....	0.2	*	0.2	*	0.4	0.7	1.0	2.1	3.6
2006.....	0.2	*	*	*	0.4	0.6	1.3	1.9	5.2
2005.....	*	*	*	*	0.5	0.8	1.6	2.6	5.9
2004.....	*	*	*	*	0.5	0.9	1.5	2.8	5.5
2003.....	*	*	0.3	*	0.8	0.7	1.6	3.1	6.1
2002.....	*	*	*	0.4	0.6	0.7	1.7	2.8	6.7
2001.....	*	0.2	*	*	*	0.6	2.1	2.6	6.7
2000.....	*	*	0.3	0.4	0.5	1.1	1.8	3.0	8.3
1999.....	*	0.3	*	*	0.6	1.1	1.7	2.8	7.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic 4/, both sexes

2007.....	0.1	0.1	0.2	0.2	0.3	0.7	0.9	1.9	3.5
2006.....	0.1	0.1	0.1	0.2	0.3	0.6	1.0	1.6	4.8
2005.....	0.1	0.1	*	0.2	0.4	0.7	1.3	2.5	5.2
2004.....	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.5	5.3
2003.....	0.2	0.2	0.2	0.2	0.6	0.5	1.2	2.8	6.0
2002.....	*	*	0.2	0.3	0.5	0.6	1.5	2.3	6.4
2001.....	*	0.1	0.2	0.2	0.3	0.6	1.7	2.4	6.1
2000.....	*	0.1	0.2	0.3	0.4	0.8	1.5	2.7	7.6
1999.....	*	0.2	0.2	0.3	0.5	0.9	1.4	2.6	7.5

Non-Hispanic 4/, male

2007.....	*	*	*	*	*	0.6	0.8	1.4	3.5
2006.....	*	*	*	*	*	0.7	*	1.2	3.9
2005.....	*	*	*	*	*	0.6	0.9	2.3	3.5
2004.....	*	*	*	*	*	*	0.9	2.2	4.6
2003.....	*	*	*	*	*	*	0.7	2.5	5.4
2002.....	*	*	*	*	*	0.7	1.3	1.5	5.3
2001.....	*	*	*	*	*	*	1.2	2.2	4.3
2000.....	*	*	*	*	*	*	1.2	2.1	5.8
1999.....	*	*	*	*	*	0.6	1.2	2.4	6.9

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	*	0.8	1.0	2.2	3.5
2006.....	*	*	*	*	0.4	0.6	1.3	1.9	5.2
2005.....	*	*	*	*	0.5	0.8	1.6	2.6	5.9
2004.....	*	*	*	*	0.4	0.9	1.5	2.7	5.6
2003.....	*	*	*	*	0.7	0.6	1.6	3.1	6.2
2002.....	*	*	*	0.4	0.6	0.6	1.7	2.7	6.8
2001.....	*	*	*	*	*	0.6	2.1	2.5	6.9
2000.....	*	*	*	0.4	0.5	1.1	1.8	3.0	8.3
1999.....	*	0.3	*	*	0.7	1.1	1.6	2.8	7.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic White, both sexes

2007 5/.....	*	0.1	0.2	0.2	0.3	0.6	0.9	1.9	3.7
2006 5/.....	*	0.2	*	0.2	0.4	0.6	0.9	1.7	5.0
2005 5/.....	*	*	*	0.2	0.4	0.7	1.2	2.6	5.2
2004 5/.....	0.1	*	0.2	*	0.4	0.7	1.2	2.5	5.3
2003 5/.....	0.1	0.2	*	*	0.5	0.6	1.2	2.7	6.0
2002.....	*	*	0.2	0.2	0.5	0.7	1.5	2.3	6.5
2001.....	*	*	*	*	0.3	0.6	1.8	2.2	6.2
2000.....	*	*	0.2	0.3	0.3	0.7	1.4	2.5	7.6
1999.....	*	0.2	0.2	*	0.5	0.8	1.4	2.7	7.7

Non-Hispanic White, male

2007 5/.....	*	*	*	*	*	*	0.9	1.3	3.7
2006 5/.....	*	*	*	*	*	0.7	*	1.2	4.0
2005 5/.....	*	*	*	*	*	*	0.9	2.4	3.4
2004 5/.....	*	*	*	*	*	*	0.8	2.3	4.8
2003 5/.....	*	*	*	*	*	*	*	2.2	5.8
2002.....	*	*	*	*	*	0.7	1.3	1.4	5.4
2001.....	*	*	*	*	*	*	1.1	2.1	4.2
2000.....	*	*	*	*	*	*	1.1	2.0	6.0
1999.....	*	*	*	*	*	*	1.2	2.4	7.3

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	0.8	1.0	2.3	3.7
2006 5/.....	*	*	*	*	0.5	*	1.2	2.0	5.5
2005 5/.....	*	*	*	*	0.5	0.7	1.4	2.7	6.0
2004 5/.....	*	*	*	*	0.5	1.0	1.4	2.6	5.6
2003 5/.....	*	*	*	*	0.7	0.7	1.5	3.0	6.1
2002.....	*	*	*	*	0.6	0.7	1.7	2.9	6.9
2001.....	*	*	*	*	*	0.7	2.2	2.3	7.1
2000.....	*	*	*	*	0.5	1.0	1.6	2.8	8.2
1999.....	*	*	*	*	0.6	1.1	1.6	2.9	7.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/, both sexes

2007.....	1.4	1.7	2.1	2.2	1.8	1.7	2.6	3.8	4.8
2006.....	1.5	1.6	2.2	2.4	1.9	2.0	2.7	3.5	5.6
2005.....	1.5	1.8	2.3	2.4	1.7	2.0	2.4	4.0	6.4
2004.....	1.5	1.8	2.2	2.2	2.3	2.3	3.1	4.3	6.1
2003.....	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8

All origins 3/, male

2007.....	1.6	1.7	2.4	2.4	2.0	1.8	3.0	3.8	5.0
2006.....	1.7	1.9	2.5	2.7	1.9	2.3	3.0	4.5	6.8
2005.....	1.5	1.9	2.5	2.6	1.6	2.0	2.8	4.9	6.9
2004.....	1.7	2.0	2.3	2.2	2.3	2.7	3.7	3.8	6.8
2003.....	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5

All origins 3/, female

2007.....	1.2	1.8	1.8	1.9	1.7	1.7	2.2	3.9	4.7
2006.....	1.3	1.3	1.9	2.1	1.8	1.8	2.4	2.8	5.1
2005.....	1.4	1.8	2.1	2.3	1.7	2.0	2.2	3.5	6.2
2004.....	1.3	1.7	2.0	2.1	2.3	2.0	2.7	4.6	5.8
2003.....	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 4/, both sexes

2007.....	1.5	1.8	2.2	2.2	1.9	1.7	2.6	4.0	4.9
2006.....	1.5	1.6	2.3	2.4	1.9	2.0	2.7	3.5	5.8
2005.....	1.5	1.9	2.4	2.5	1.7	2.0	2.5	4.0	6.6
2004.....	1.6	1.9	2.2	2.2	2.4	2.4	3.1	4.4	6.2
2003.....	1.6	1.9	2.4	2.4	2.4	2.2	3.0	4.6	6.7
2002.....	1.7	2.0	2.4	2.5	2.4	2.6	3.1	4.8	6.7
2001.....	1.5	1.9	2.2	2.6	2.3	2.7	3.2	4.3	7.2
2000.....	1.5	1.9	2.2	2.5	2.6	2.5	3.2	4.9	7.5
1999.....	1.7	1.8	2.4	2.4	2.6	2.4	3.4	4.9	7.9

Non-Hispanic 4/, male

2007.....	1.7	1.7	2.5	2.4	2.0	1.7	3.1	4.0	5.0
2006.....	1.8	2.0	2.6	2.8	1.9	2.3	3.0	4.5	6.8
2005.....	1.6	2.0	2.6	2.6	1.7	2.1	2.7	4.8	7.2
2004.....	1.8	2.0	2.4	2.2	2.4	2.7	3.6	4.0	7.0
2003.....	1.7	1.9	2.5	2.6	2.5	2.3	2.8	4.6	6.7
2002.....	1.9	2.1	2.4	2.8	2.5	2.9	3.4	4.6	7.0
2001.....	1.5	1.9	2.2	2.4	2.5	3.0	3.5	4.8	7.6
2000.....	1.6	2.2	2.0	2.5	2.8	2.6	3.1	5.2	10.4
1999.....	1.9	2.0	2.5	2.6	2.4	2.5	3.4	5.7	8.5

Non-Hispanic 4/, female

2007.....	1.3	1.8	1.8	2.0	1.7	1.8	2.3	4.0	4.8
2006.....	1.3	1.3	2.0	2.1	1.9	1.8	2.5	2.9	5.3
2005.....	1.4	1.8	2.2	2.4	1.8	2.0	2.3	3.4	6.3
2004.....	1.3	1.8	2.1	2.2	2.3	2.1	2.7	4.7	5.9
2003.....	1.4	1.8	2.4	2.3	2.3	2.2	3.2	4.5	6.8
2002.....	1.5	1.9	2.4	2.3	2.3	2.3	2.9	4.9	6.6
2001.....	1.5	1.8	2.2	2.8	2.1	2.4	3.0	4.1	7.0
2000.....	1.3	1.6	2.4	2.5	2.5	2.4	3.3	4.8	6.4
1999.....	1.6	1.7	2.2	2.2	2.8	2.3	3.5	4.5	7.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic White, both sexes

2007 5/.....	1.5	1.8	2.3	2.4	1.9	1.7	2.8	4.0	5.0
2006 5/.....	1.6	1.6	2.4	2.6	1.9	2.1	2.9	3.5	5.9
2005 5/.....	1.5	2.0	2.5	2.6	1.9	2.2	2.5	4.2	6.7
2004 5/.....	1.6	2.0	2.4	2.4	2.5	2.5	3.0	4.5	6.5
2003 5/.....	1.5	2.0	2.6	2.5	2.4	2.3	3.1	4.7	6.9
2002.....	1.7	2.0	2.5	2.6	2.5	2.6	3.2	4.8	6.9
2001.....	1.4	2.0	2.3	2.8	2.4	2.7	3.3	4.4	7.4
2000.....	1.5	2.0	2.2	2.6	2.7	2.5	3.3	5.1	7.8
1999.....	1.8	1.9	2.5	2.5	2.7	2.4	3.4	5.0	8.0

Non-Hispanic White, male

2007 5/.....	1.7	1.8	2.6	2.6	2.2	1.8	3.2	4.0	5.2
2006 5/.....	1.7	2.0	2.7	2.9	1.9	2.3	3.0	4.7	6.8
2005 5/.....	1.7	2.1	2.7	2.7	1.9	2.3	2.8	5.2	7.2
2004 5/.....	1.9	2.1	2.5	2.4	2.5	2.9	3.4	4.1	7.4
2003 5/.....	1.6	2.0	2.6	2.6	2.4	2.2	2.8	4.4	6.8
2002.....	1.9	2.1	2.4	2.8	2.6	2.8	3.4	4.4	7.3
2001.....	1.4	2.0	2.3	2.6	2.7	3.1	3.8	4.5	7.4
2000.....	1.6	2.3	2.0	2.7	2.6	2.5	3.2	5.5	10.8
1999.....	1.9	2.0	2.7	2.7	2.4	2.5	3.0	5.9	8.6

Non-Hispanic White, female

2007 5/.....	1.3	1.8	2.0	2.1	1.7	1.7	2.4	3.9	4.9
2006 5/.....	1.4	1.3	2.2	2.3	1.9	1.9	2.7	2.8	5.5
2005 5/.....	1.4	2.0	2.3	2.6	1.9	2.2	2.3	3.6	6.5
2004 5/.....	1.3	2.0	2.3	2.4	2.5	2.3	2.8	4.7	6.1
2003 5/.....	1.5	1.9	2.5	2.5	2.3	2.3	3.3	4.8	6.9
2002.....	1.5	2.0	2.5	2.4	2.4	2.4	3.0	5.1	6.7
2001.....	1.5	1.9	2.3	3.0	2.1	2.4	3.0	4.3	7.4
2000.....	1.4	1.8	2.4	2.5	2.7	2.5	3.4	4.9	6.6
1999.....	1.6	1.8	2.4	2.2	2.9	2.3	3.6	4.5	7.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes

2007.....	4.6	5.5	6.4	7.4	10.0	16.6	30.4	63.2	251.4
2006.....	4.9	5.6	5.9	7.2	9.5	15.5	28.7	58.6	236.1
2005.....	6.4	7.0	6.9	7.7	10.4	15.7	27.6	54.0	224.4
2004.....	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
2003.....	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All origins 3/, male

2007.....	5.8	7.0	8.3	9.6	12.0	20.0	33.2	63.6	197.6
2006.....	6.3	7.2	7.7	9.0	11.9	18.5	32.3	61.3	191.1
2005.....	8.2	9.3	9.2	9.6	12.8	18.8	31.0	54.8	183.9
2004.....	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
2003.....	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All origins 3/, female

2007.....	3.4	4.0	4.6	5.4	8.2	13.8	28.4	63.0	277.0
2006.....	3.5	4.1	4.2	5.6	7.3	13.1	26.0	56.9	257.2
2005.....	4.6	4.8	4.7	6.0	8.3	13.2	25.1	53.5	243.0
2004.....	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
2003.....	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, both sexes

2007	3.4	3.8	4.2	4.5	5.1	10.5	18.4	32.3	108.7
2006	3.1	2.9	3.4	4.4	5.5	8.7	13.2	27.2	117.0
2005	3.3	4.0	3.1	4.3	6.7	9.2	13.8	26.2	105.2
2004	4.2	2.9	3.0	5.2	4.3	8.7	12.0	23.3	99.4
2003	3.5	4.0	3.5	3.6	5.3	7.2	14.6	24.6	110.9
2002	3.1	5.0	3.9	6.1	5.5	9.0	11.7	30.3	91.5
2001	9.0	7.4	6.7	7.1	8.7	11.3	14.4	27.8	108.2
2000	8.1	7.9	7.5	5.6	8.2	12.2	16.5	27.8	102.2
1999	3.8	2.3	3.2	4.7	6.2	8.1	13.9	29.4	87.0

Hispanic, male

2007	4.8	5.2	4.6	7.4	6.5	13.6	21.8	33.3	91.5
2006	4.3	4.1	4.5	6.8	7.9	12.2	12.4	25.4	90.2
2005	5.1	6.1	4.3	6.3	8.3	12.7	16.7	28.0	99.6
2004	5.6	3.9	4.3	7.9	*	11.2	12.1	21.7	86.9
2003	5.7	5.5	5.6	5.4	6.5	10.2	15.2	24.9	95.4
2002	3.8	5.8	4.8	8.5	7.8	11.8	15.4	37.7	88.7
2001	14.0	10.7	10.0	11.4	12.8	16.0	18.4	37.7	87.6
2000	12.4	12.3	9.0	8.9	13.1	16.0	19.9	36.8	106.8
1999	6.2	*	*	6.9	8.5	*	21.8	*	77.6

Hispanic, female

2007	2.0	2.3	3.9	*	*	8.0	15.9	31.5	118.0
2006	1.8	*	*	*	*	5.9	13.8	28.3	131.2
2005	*	*	*	*	5.4	6.5	11.6	25.0	108.2
2004	2.8	*	*	*	*	6.7	11.9	24.3	105.9
2003	*	2.5	*	*	*	*	14.2	24.4	118.8
2002	2.4	4.2	*	*	*	6.8	9.1	25.5	92.8
2001	4.0	4.4	3.8	*	*	7.7	11.6	21.7	118.2
2000	3.8	3.6	6.2	*	*	9.2	14.1	22.4	99.9
1999	*	*	*	*	*	8.0	*	30.0	91.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 4/, both sexes

2007.....	4.7	5.6	6.5	7.6	10.3	17.0	31.2	65.1	258.9
2006.....	5.0	5.9	6.1	7.4	9.7	16.0	29.7	60.3	241.9
2005.....	6.6	7.2	7.1	7.9	10.7	16.0	28.2	55.2	229.6
2004.....	5.9	6.0	6.6	8.0	10.6	15.8	28.8	56.9	226.7
2003.....	7.6	7.0	7.0	8.5	11.0	17.9	27.3	57.1	228.4
2002.....	6.7	7.0	7.2	8.3	10.7	16.6	29.1	54.1	211.9
2001.....	9.0	8.8	8.5	10.0	12.9	18.2	30.4	55.3	209.6
2000.....	8.6	7.9	8.5	10.5	14.5	19.3	30.6	57.4	211.6
1999.....	6.0	6.0	7.1	9.1	12.2	17.9	27.0	51.2	190.9

Non-Hispanic 4/, male

2007.....	5.9	7.1	8.6	9.8	12.4	20.5	33.9	65.5	203.6
2006.....	6.5	7.5	7.9	9.1	12.2	18.8	33.5	63.4	196.3
2005.....	8.5	9.5	9.5	9.7	13.1	19.1	31.7	55.8	187.5
2004.....	7.6	7.9	8.5	10.1	13.2	18.2	31.6	59.9	188.0
2003.....	9.7	9.1	9.0	10.9	14.1	21.8	32.0	59.4	181.1
2002.....	9.0	9.2	9.2	11.2	12.9	20.1	33.7	61.1	178.6
2001.....	11.6	11.8	11.1	13.2	15.7	21.1	33.8	61.6	178.2
2000.....	11.5	10.9	11.3	13.5	19.0	22.6	36.6	63.0	185.4
1999.....	8.3	8.1	9.3	12.1	15.4	22.3	31.8	55.2	169.5

Non-Hispanic 4/, female

2007.....	3.6	4.2	4.6	5.6	8.5	14.2	29.3	64.8	285.0
2006.....	3.6	4.4	4.3	5.9	7.6	13.6	26.8	58.4	263.1
2005.....	4.9	5.0	4.9	6.2	8.5	13.5	25.7	54.8	248.9
2004.....	4.2	4.2	4.7	6.1	8.3	13.8	26.8	55.1	244.0
2003.....	5.6	5.0	5.1	6.2	8.2	14.8	23.9	55.6	249.2
2002.....	4.5	5.0	5.2	5.7	8.8	13.7	25.9	49.8	226.3
2001.....	6.4	6.0	6.1	7.1	10.5	15.8	28.0	51.6	222.7
2000.....	5.7	5.0	5.8	7.7	10.5	16.7	26.5	54.1	222.3
1999.....	3.9	4.0	5.1	6.5	9.4	14.5	23.6	48.9	199.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic White, both sexes

2007 5/.....	4.2	5.1	6.1	7.2	9.7	16.7	31.0	65.1	264.8
2006 5/.....	4.9	5.5	5.7	7.2	9.4	15.4	29.2	60.1	247.4
2005 5/.....	6.2	6.8	6.5	7.4	10.0	15.4	27.9	55.0	234.9
2004 5/.....	5.7	5.5	6.0	7.5	10.2	15.0	28.5	56.7	230.0
2003 5/.....	7.0	6.4	6.5	7.9	10.4	17.2	27.1	57.6	232.9
2002.....	6.1	6.1	6.6	7.9	10.1	16.3	28.7	54.1	215.8
2001.....	8.7	8.3	7.9	9.5	12.1	17.5	30.1	55.0	213.3
2000.....	8.0	7.1	7.7	9.7	13.2	18.7	30.0	57.0	215.8
1999.....	5.2	5.2	6.3	8.4	11.3	17.3	25.9	50.1	193.8

Non-Hispanic White, male

2007 5/.....	5.3	6.3	7.8	9.2	11.6	20.0	33.3	65.7	207.6
2006 5/.....	6.1	6.9	7.4	8.9	11.7	17.4	32.7	62.6	199.4
2005 5/.....	8.1	8.9	8.6	9.0	12.1	18.2	31.2	54.9	189.8
2004 5/.....	7.3	7.1	7.7	9.3	12.6	17.6	31.1	59.6	191.0
2003 5/.....	8.9	8.3	8.2	10.0	13.2	21.0	31.8	59.3	183.3
2002.....	8.1	7.8	8.3	10.6	12.1	19.7	32.3	60.6	181.5
2001.....	11.2	11.1	10.0	12.1	14.4	20.5	33.9	61.6	180.4
2000.....	10.9	9.7	10.0	12.5	17.1	21.6	35.9	62.6	189.6
1999.....	6.9	7.0	8.2	11.2	14.2	21.5	30.5	53.3	171.1

Non-Hispanic White, female

2007 5/.....	3.2	3.9	4.4	5.3	8.0	13.9	29.2	64.7	291.9
2006 5/.....	3.6	4.1	4.0	5.5	7.3	13.7	26.7	58.5	269.7
2005 5/.....	4.4	4.8	4.5	6.0	8.0	13.1	25.5	55.1	255.5
2004 5/.....	4.1	4.0	4.4	5.8	8.2	12.9	26.6	54.9	247.5
2003 5/.....	5.0	4.4	4.9	6.0	8.0	14.2	23.7	56.5	254.7
2002.....	4.1	4.4	4.9	5.4	8.3	13.6	26.1	50.2	230.6
2001.....	6.2	5.5	5.9	7.0	10.0	15.1	27.4	51.0	227.0
2000.....	5.2	4.5	5.6	7.1	9.7	16.3	25.9	53.6	226.4
1999.....	3.5	3.5	4.6	5.9	8.8	14.0	22.7	48.2	202.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic Black, both sexes

2007 5/.....	8.7	10.5	11.2	13.3	18.1	25.0	43.0	82.1	245.2
2006 5/.....	7.7	10.4	10.6	11.3	15.3	25.0	42.2	77.6	226.9
2005 5/.....	11.1	12.4	13.9	13.9	19.4	25.6	39.1	70.3	213.3
2004 5/.....	8.6	11.2	12.4	14.7	16.4	26.5	40.1	72.2	223.3
2003 5/.....	13.7	13.5	13.2	15.9	18.4	31.1	35.3	61.2	214.5
2002.....	12.2	15.9	13.9	13.8	18.0	23.3	39.7	65.4	198.1
2001.....	13.2	14.9	14.6	15.7	21.8	28.1	39.2	67.6	194.2
2000.....	13.9	16.0	15.8	19.1	28.3	30.3	44.7	74.1	189.2
1999.....	13.2	13.2	15.0	17.1	21.9	27.8	43.3	73.4	182.0

Non-Hispanic Black, male

2007 5/.....	11.2	14.3	16.6	18.2	24.3	33.3	52.0	84.6	187.6
2006 5/.....	11.0	14.4	14.5	13.3	21.0	37.0	54.3	94.8	196.1
2005 5/.....	13.6	17.2	19.5	19.9	26.5	33.5	49.1	83.5	196.0
2004 5/.....	11.6	16.5	18.1	20.0	22.7	28.9	44.8	82.2	182.9
2003 5/.....	17.8	17.9	18.8	23.2	26.6	39.8	42.1	70.2	197.1
2002.....	17.3	22.4	19.8	20.4	22.8	31.1	57.3	87.4	182.7
2001.....	17.4	19.2	22.2	25.9	29.7	32.0	41.2	71.1	186.0
2000.....	18.3	23.5	24.7	25.5	41.0	38.6	55.4	85.0	163.2
1999.....	20.0	18.7	22.3	23.9	30.0	36.4	54.6	89.4	180.7

Non-Hispanic Black, female

2007 5/.....	6.6	7.3	6.8	9.5	13.5	19.3	37.6	80.8	269.1
2006 5/.....	4.7	7.0	7.4	9.8	11.0	16.7	34.9	68.5	239.5
2005 5/.....	8.9	8.3	9.4	9.3	14.1	20.2	33.1	63.4	220.2
2004 5/.....	5.9	6.8	7.7	10.5	11.7	24.8	37.3	66.8	239.3
2003 5/.....	10.1	9.8	8.6	10.2	12.4	25.2	31.4	56.5	221.2
2002.....	7.9	10.4	9.1	8.8	14.4	18.1	29.2	54.0	204.0
2001.....	9.6	11.2	8.4	7.7	16.0	25.5	37.9	65.8	197.2
2000.....	10.1	9.7	8.5	14.1	18.9	24.7	38.3	68.7	198.9
1999.....	7.5	8.5	9.1	11.8	15.9	22.1	36.4	65.4	182.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/, both sexes

2007.....	30.8	45.0	60.5	82.0	118.8	186.8	337.6	639.2	1,701.9
2006.....	33.3	47.6	61.8	85.0	118.6	191.7	335.2	646.8	1,724.2
2005.....	33.6	47.8	60.3	81.9	115.6	182.4	313.2	593.5	1,559.1
2004.....	33.6	45.0	58.1	78.7	111.5	176.4	300.6	554.8	1,461.6
2003.....	33.3	45.3	56.6	80.9	114.5	178.3	303.7	559.8	1,508.4
2002.....	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001.....	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000.....	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
1999.....	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7

All origins 3/, male

2007.....	36.8	54.7	71.3	92.6	131.3	201.0	351.0	649.0	1,449.1
2006.....	39.7	57.9	73.0	96.3	131.0	207.6	349.9	652.5	1,455.0
2005.....	40.7	58.5	71.9	93.1	125.3	194.2	327.1	595.6	1,305.3
2004.....	40.6	56.0	68.3	88.8	123.1	187.2	315.5	549.4	1,236.6
2003.....	40.3	55.2	65.6	91.0	125.7	191.0	316.5	563.7	1,258.1
2002.....	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001.....	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000.....	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
1999.....	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4

All origins 3/, female

2007.....	25.0	35.7	50.3	72.2	107.8	175.3	327.7	633.0	1,822.1
2006.....	27.0	37.8	51.3	74.7	107.8	178.8	324.4	643.2	1,850.2
2005.....	26.7	37.6	49.4	71.7	107.2	172.9	303.1	592.2	1,675.6
2004.....	26.7	34.6	48.4	69.5	101.4	167.6	289.9	558.0	1,562.8
2003.....	26.5	36.0	48.1	71.7	104.8	168.0	294.7	557.4	1,619.1
2002.....	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001.....	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000.....	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
1999.....	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Hispanic, both sexes

2007.....	21.7	33.4	43.3	62.5	83.5	136.4	242.0	432.7	1,049.8
2006.....	23.9	35.5	47.1	61.2	89.9	146.3	249.1	443.0	1,067.7
2005.....	23.4	34.8	44.7	63.7	86.3	134.9	212.0	412.4	890.7
2004.....	24.3	32.3	43.6	56.7	79.6	128.6	225.1	357.7	880.5
2003.....	25.3	36.9	46.3	63.5	87.0	132.0	219.9	389.4	887.2
2002.....	25.6	30.1	43.3	55.6	90.9	136.7	212.2	372.5	870.8
2001.....	22.8	30.7	42.2	59.7	83.9	136.4	209.8	396.3	991.0
2000.....	24.8	31.0	45.2	50.5	84.3	127.4	217.6	404.9	913.0
1999.....	26.0	31.2	43.3	61.6	88.6	135.5	202.4	374.7	1,039.7

Hispanic, male

2007.....	27.6	42.2	53.5	69.5	94.0	151.8	260.8	410.7	913.8
2006.....	30.2	46.7	59.4	71.4	104.1	159.4	254.4	461.0	897.6
2005.....	31.3	43.9	55.8	78.5	94.2	144.2	230.6	412.4	789.5
2004.....	32.8	43.8	53.5	63.5	89.2	137.0	228.6	323.7	770.5
2003.....	32.5	46.4	59.4	72.1	97.5	141.4	229.1	426.2	757.3
2002.....	34.0	37.5	51.9	62.6	109.0	147.9	233.6	391.2	808.2
2001.....	30.6	40.0	51.7	69.0	96.0	144.6	205.3	403.8	939.0
2000.....	33.3	38.7	59.2	55.8	105.9	138.1	221.5	378.4	870.7
1999.....	35.4	39.2	53.2	74.8	96.6	159.9	220.0	389.8	923.1

Hispanic, female

2007.....	15.5	24.7	33.7	56.4	74.7	124.3	228.4	447.4	1,123.6
2006.....	17.3	24.2	35.5	52.0	77.9	136.0	245.2	431.1	1,158.1
2005.....	15.2	25.8	34.4	50.6	79.7	127.7	198.6	412.3	943.7
2004.....	15.5	21.0	34.4	50.8	71.7	122.0	222.6	380.1	937.0
2003.....	17.8	27.6	34.2	56.0	78.4	124.9	213.4	365.4	953.1
2002.....	17.0	22.9	35.4	49.4	76.0	128.1	197.0	360.6	901.9
2001.....	14.9	21.9	33.5	51.7	74.0	130.2	213.0	391.7	1,016.5
2000.....	16.3	23.7	32.5	45.9	66.7	119.3	214.9	421.1	933.8
1999.....	16.7	23.5	34.4	50.3	82.2	117.2	190.0	365.7	1,096.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 4/, both sexes

2007.....	31.9	46.1	61.9	83.3	121.4	190.5	343.9	651.0	1,735.2
2006.....	34.4	48.7	63.0	86.8	120.7	194.8	340.4	657.7	1,755.2
2005.....	34.8	49.0	61.4	83.2	117.7	185.6	319.1	602.4	1,589.3
2004.....	34.5	46.1	59.1	80.2	113.7	179.3	304.6	563.8	1,485.2
2003.....	34.0	46.0	57.3	82.0	116.2	181.0	307.7	567.1	1,531.7
2002.....	34.3	43.4	57.0	78.4	111.5	175.8	301.5	561.8	1,514.9
2001.....	32.9	41.5	53.8	77.5	113.6	174.0	299.1	543.9	1,491.3
2000.....	31.3	40.9	53.4	75.6	107.1	169.8	291.3	541.5	1,462.0
1999.....	30.0	38.8	51.3	72.6	108.5	169.1	284.1	514.7	1,420.1

Non-Hispanic 4/, male

2007.....	37.9	55.9	72.6	94.2	134.0	204.4	356.7	663.0	1,479.5
2006.....	40.8	58.8	73.9	98.1	132.7	210.5	355.5	663.2	1,484.0
2005.....	41.8	59.8	72.9	94.0	127.4	197.3	332.5	605.0	1,330.8
2004.....	41.3	56.8	69.3	90.4	125.2	190.2	320.1	560.5	1,257.5
2003.....	41.0	55.7	65.9	92.0	127.3	193.8	320.6	569.7	1,278.3
2002.....	41.5	53.4	65.0	89.3	123.3	188.7	312.9	561.0	1,280.6
2001.....	39.4	49.5	60.0	86.7	125.7	188.5	311.8	544.1	1,282.3
2000.....	38.7	48.0	60.8	84.6	117.8	184.8	306.1	542.6	1,279.8
1999.....	36.1	45.6	59.6	80.8	117.9	185.0	300.8	532.1	1,234.0

Non-Hispanic 4/, female

2007.....	26.1	36.7	51.8	73.3	110.4	179.0	334.4	643.4	1,856.0
2006.....	28.1	39.1	52.6	76.5	110.2	181.9	329.4	654.3	1,881.3
2005.....	28.0	38.7	50.6	73.3	109.3	176.0	309.4	600.9	1,707.2
2004.....	27.9	35.8	49.5	70.9	103.6	170.5	293.5	565.8	1,587.0
2003.....	27.3	36.6	49.1	72.8	106.6	170.7	298.6	565.5	1,643.1
2002.....	27.3	33.9	49.5	68.5	101.3	165.4	293.4	562.3	1,615.9
2001.....	26.6	33.9	47.9	69.1	103.1	162.4	290.1	543.7	1,578.3
2000.....	24.2	34.1	46.5	67.3	97.8	158.0	281.0	540.9	1,535.7
1999.....	24.2	32.3	43.6	65.1	100.4	156.6	272.4	504.6	1,494.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic White, both sexes

2007 5/.....	30.1	43.1	59.0	80.2	118.2	188.3	343.1	658.5	1,771.7
2006 5/.....	32.0	45.5	59.0	84.3	117.9	192.2	341.4	661.8	1,790.9
2005 5/.....	32.0	45.6	58.3	79.9	114.5	182.6	317.1	605.4	1,618.2
2004 5/.....	31.6	42.8	55.5	76.8	109.9	176.3	302.7	567.5	1,511.9
2003 5/.....	31.2	41.8	53.8	77.8	113.4	178.0	305.7	569.1	1,558.9
2002.....	31.2	39.8	53.6	74.5	108.5	171.7	299.2	562.5	1,540.5
2001.....	29.2	37.5	49.8	73.3	110.2	170.7	296.0	545.5	1,512.9
2000.....	27.3	37.3	49.3	71.2	103.2	165.9	287.8	540.7	1,481.3
1999.....	26.0	34.8	47.4	68.6	103.6	164.6	279.7	516.0	1,443.8

Non-Hispanic White, male

2007 5/.....	36.5	52.5	69.0	90.3	129.3	200.2	352.6	669.0	1,512.0
2006 5/.....	38.9	55.6	69.2	94.6	130.1	205.7	353.3	663.9	1,512.3
2005 5/.....	39.4	56.3	69.2	89.9	122.5	193.7	327.8	604.2	1,354.9
2004 5/.....	38.8	53.1	65.0	86.4	119.8	186.0	313.7	561.7	1,273.4
2003 5/.....	38.4	50.8	61.8	86.8	122.7	188.5	315.2	566.4	1,300.3
2002.....	38.3	49.3	60.8	84.6	118.5	182.6	307.9	558.6	1,293.4
2001.....	35.5	45.0	55.1	82.2	121.0	183.6	306.0	544.6	1,293.1
2000.....	34.7	44.1	56.1	78.5	112.4	178.8	300.6	538.2	1,288.3
1999.....	31.8	41.2	55.0	76.3	112.0	179.6	293.3	529.5	1,239.8

Non-Hispanic White, female

2007 5/.....	23.7	33.9	49.3	70.8	108.3	178.3	335.9	651.8	1,894.7
2006 5/.....	25.2	35.6	49.1	74.7	107.0	181.0	332.5	660.5	1,920.7
2005 5/.....	24.6	35.0	47.8	70.4	107.4	173.3	309.2	606.2	1,738.7
2004 5/.....	24.5	32.7	46.3	67.8	101.2	168.3	294.7	571.1	1,618.7
2003 5/.....	24.1	32.9	46.1	69.4	105.1	169.4	298.8	570.8	1,672.7
2002.....	24.2	30.4	46.7	65.1	99.6	162.8	293.0	564.9	1,647.2
2001.....	23.0	30.2	44.7	65.1	100.7	160.1	288.9	546.0	1,604.4
2000.....	19.9	30.7	42.7	64.4	95.2	155.5	278.8	542.2	1,559.1
1999.....	20.3	28.5	40.1	61.5	96.2	152.6	270.1	508.1	1,525.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic Black, both sexes

2007 5/.....	50.7	77.0	99.9	130.3	176.3	250.8	431.4	712.3	1,656.9
2006 5/.....	56.3	82.8	108.4	129.4	169.1	261.0	406.5	743.9	1,660.0
2005 5/.....	59.9	81.9	101.1	131.2	167.4	251.6	408.1	691.7	1,525.4
2004 5/.....	60.9	79.6	102.3	128.1	169.5	248.1	391.4	633.2	1,425.6
2003 5/.....	59.7	85.3	98.5	133.8	164.7	245.2	387.7	636.4	1,451.6
2002.....	61.9	79.5	97.9	126.0	158.0	246.8	373.5	648.9	1,443.4
2001.....	64.1	80.9	97.2	124.2	165.8	239.3	384.5	619.4	1,431.5
2000.....	65.4	77.2	97.4	124.6	159.8	237.4	375.9	624.5	1,403.3
1999.....	63.0	77.3	93.5	118.8	169.4	242.1	373.1	560.6	1,295.3

Non-Hispanic Black, male

2007 5/.....	56.6	93.3	121.7	157.0	213.9	292.8	491.9	767.7	1,389.1
2006 5/.....	63.0	97.5	129.9	157.9	192.9	309.3	464.0	816.7	1,437.6
2005 5/.....	67.8	96.3	122.0	154.3	197.7	279.0	456.5	755.5	1,261.7
2004 5/.....	68.3	97.8	124.5	150.6	204.7	284.0	473.8	666.8	1,316.9
2003 5/.....	67.5	104.5	115.9	160.6	194.9	294.2	447.6	722.2	1,243.4
2002.....	72.8	96.4	118.3	148.7	188.1	292.8	421.6	699.6	1,331.3
2001.....	75.8	97.7	116.3	144.0	199.8	276.5	449.4	669.7	1,329.4
2000.....	75.8	90.8	116.7	153.5	191.1	284.0	432.1	698.6	1,340.9
1999.....	74.5	91.6	112.6	135.8	192.6	274.5	439.4	638.5	1,303.8

Non-Hispanic Black, female

2007 5/.....	45.6	63.3	82.1	109.4	148.7	221.9	395.1	683.2	1,767.9
2006 5/.....	50.4	70.4	90.6	107.3	151.6	227.8	372.2	705.4	1,750.7
2005 5/.....	53.1	69.8	83.8	113.3	144.9	232.9	379.4	658.0	1,631.5
2004 5/.....	54.4	64.3	84.1	110.6	143.4	223.8	342.6	615.5	1,468.5
2003 5/.....	52.9	69.1	84.1	112.9	142.3	212.2	352.1	591.6	1,532.3
2002.....	52.5	65.3	81.1	108.4	135.7	215.9	344.8	622.5	1,486.2
2001.....	54.1	66.6	81.6	108.6	140.6	214.5	345.7	593.9	1,469.3
2000.....	56.4	65.7	81.6	101.8	136.6	206.4	342.1	587.2	1,426.5
1999.....	53.2	65.2	78.0	105.5	152.2	220.6	333.1	522.0	1,292.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes

2007.....	46.9	45.7	38.9	35.2	38.9	53.1	80.8	137.1	286.7
2006.....	47.0	43.7	37.1	35.0	39.6	50.5	81.3	136.0	274.9
2005.....	44.6	41.6	35.5	36.0	40.1	53.7	82.7	137.0	279.5
2004.....	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
2003.....	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4

All origins 3/, male

2007.....	65.2	64.6	55.9	49.3	54.8	73.5	106.4	179.7	349.0
2006.....	66.8	63.0	52.9	49.6	55.1	66.4	107.3	176.5	337.1
2005.....	63.5	60.7	51.3	51.4	54.6	72.6	110.0	180.8	344.2
2004.....	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
2003.....	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4

All origins 3/, female

2007.....	29.0	27.5	23.0	22.3	25.1	36.4	62.0	110.3	257.0
2006.....	27.8	25.2	22.2	21.7	26.0	37.5	62.3	110.7	245.8
2005.....	26.0	23.3	20.7	22.0	27.4	38.5	62.8	109.7	249.8
2004.....	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
2003.....	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, both sexes

2007	34.2	35.9	30.5	28.2	31.1	44.3	61.8	99.2	152.0
2006	36.2	34.9	34.1	31.9	35.0	40.9	63.8	91.6	142.8
2005	33.8	34.5	29.5	31.4	35.2	44.8	63.4	104.7	153.6
2004	31.9	29.8	28.8	33.0	33.3	42.4	59.6	97.2	160.8
2003	33.9	31.5	27.8	34.5	35.0	42.1	59.0	90.2	151.8
2002	33.0	30.5	27.3	30.5	34.7	45.4	69.9	90.0	153.0
2001	32.6	30.5	29.7	31.6	32.5	49.3	62.9	92.9	166.3
2000	30.5	30.2	28.3	29.3	31.2	48.4	59.1	95.8	184.5
1999	32.5	29.6	27.4	29.8	42.3	42.6	59.1	88.7	182.4

Hispanic, male

2007	52.4	57.2	44.6	40.6	47.3	64.9	84.3	128.6	186.0
2006	53.6	54.6	52.4	49.3	51.9	57.2	84.5	113.8	174.8
2005	52.3	53.8	45.5	48.3	52.5	66.3	87.0	137.5	168.1
2004	49.4	45.8	41.4	49.8	48.8	56.3	82.4	127.4	193.3
2003	53.3	46.9	42.1	50.0	49.7	58.1	88.7	125.8	195.5
2002	50.0	47.0	43.1	44.1	56.2	61.9	97.3	121.2	224.2
2001	50.0	47.3	42.6	48.3	46.1	66.3	92.8	128.6	198.1
2000	49.1	47.4	42.3	44.3	47.0	70.5	86.4	109.0	251.9
1999	52.9	46.6	42.3	45.0	62.6	64.2	78.0	128.9	258.8

Hispanic, female

2007	14.9	14.5	17.0	17.2	17.5	28.1	45.4	79.6	133.5
2006	17.9	15.2	16.9	16.5	20.8	28.2	48.8	76.8	125.8
2005	14.4	15.4	14.5	16.4	20.7	28.1	46.6	83.0	146.0
2004	13.9	14.2	17.0	18.3	20.4	31.6	43.4	77.3	144.0
2003	14.0	16.5	14.6	21.0	22.9	29.8	37.9	67.0	129.7
2002	15.8	14.6	12.8	18.6	17.1	32.8	50.3	70.1	117.7
2001	15.2	14.6	18.1	17.2	21.5	36.3	41.6	70.8	150.8
2000	12.0	13.8	15.7	16.4	18.4	31.7	39.7	87.8	151.3
1999	12.2	13.5	14.0	16.7	25.9	26.4	45.7	64.7	145.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 4/, both sexes

2007.....	48.4	46.5	39.5	35.6	39.4	53.7	81.9	139.3	293.6
2006.....	48.2	44.4	37.3	35.1	39.9	51.1	82.3	138.1	281.3
2005.....	45.7	42.1	35.9	36.3	40.3	54.3	83.7	138.5	285.1
2004.....	43.6	39.2	33.0	33.6	38.6	51.0	80.7	137.6	281.5
2003.....	41.3	36.8	33.1	32.6	38.1	51.2	79.7	135.3	283.7
2002.....	39.1	34.4	30.4	32.6	38.4	50.9	80.3	132.4	279.8
2001.....	36.0	31.8	29.0	31.2	36.9	49.1	81.1	131.5	279.8
2000.....	34.3	30.5	29.8	32.3	35.5	48.6	76.0	125.4	276.1
1999.....	33.9	28.9	30.0	31.1	38.3	51.2	79.2	134.3	285.0

Non-Hispanic 4/, male

2007.....	66.7	65.0	56.6	49.8	55.2	74.0	107.5	182.6	358.0
2006.....	68.2	63.6	52.8	49.4	55.2	66.9	108.5	179.5	345.9
2005.....	64.6	61.1	51.5	51.4	54.5	72.9	111.2	182.9	352.9
2004.....	61.4	57.1	47.1	46.4	52.8	67.5	105.0	182.1	352.5
2003.....	59.2	52.4	47.3	46.6	52.1	67.1	105.4	177.2	358.9
2002.....	56.9	49.5	44.5	45.5	52.8	68.9	106.5	171.3	351.2
2001.....	54.0	46.5	41.4	43.9	50.0	65.2	108.8	174.4	364.5
2000.....	51.4	45.4	42.8	45.0	47.9	64.0	100.1	166.5	363.1
1999.....	51.1	42.9	43.1	44.1	52.2	68.2	103.5	182.5	378.7

Non-Hispanic 4/, female

2007.....	30.6	28.8	23.4	22.6	25.6	37.0	63.0	112.0	263.2
2006.....	28.9	26.1	22.6	22.1	26.4	38.2	63.1	112.4	251.3
2005.....	27.3	23.9	21.2	22.4	27.9	39.1	63.8	111.0	254.1
2004.....	26.4	22.0	19.7	22.0	26.2	37.6	63.2	110.3	249.8
2003.....	23.8	22.0	19.7	19.9	25.9	38.5	61.4	109.7	250.6
2002.....	21.8	20.0	17.2	20.9	25.9	36.4	61.7	108.8	249.0
2001.....	18.7	17.8	17.4	19.7	25.6	36.2	61.8	105.8	244.6
2000.....	17.7	16.2	17.5	20.6	24.9	36.5	59.2	101.1	240.9
1999.....	17.4	15.6	17.7	19.4	26.3	37.9	62.2	106.1	247.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic White, both sexes

2007 5/.....	50.0	46.8	38.9	35.4	39.0	54.2	83.8	145.5	309.9
2006 5/.....	48.7	44.0	36.2	34.3	39.7	51.5	84.1	143.4	295.9
2005 5/.....	46.6	41.0	35.0	35.4	39.9	54.2	84.3	141.6	297.2
2004 5/.....	44.3	38.5	32.1	33.1	38.4	50.9	81.7	142.0	295.2
2003 5/.....	41.4	35.7	32.3	32.1	37.2	51.3	81.1	138.4	296.5
2002.....	38.8	33.5	29.4	32.0	37.3	51.0	81.2	136.1	290.9
2001.....	34.7	30.8	28.1	30.1	36.5	48.5	81.9	134.4	289.3
2000.....	33.0	28.7	28.6	31.7	34.5	47.8	76.2	128.1	285.6
1999.....	32.3	27.4	28.6	29.2	37.1	49.7	78.5	136.1	293.3

Non-Hispanic White, male

2007 5/.....	68.0	64.3	54.7	49.0	53.5	74.2	108.6	189.9	376.3
2006 5/.....	67.9	61.5	50.3	47.4	54.6	66.6	109.3	186.3	362.3
2005 5/.....	65.0	58.6	49.3	48.9	53.3	71.7	111.3	185.2	366.3
2004 5/.....	61.9	55.1	44.8	44.7	52.4	67.0	105.8	186.4	366.8
2003 5/.....	58.6	50.4	45.5	45.3	50.4	67.1	106.1	179.6	372.9
2002.....	56.6	47.3	42.2	44.1	50.1	68.3	106.5	174.1	361.6
2001.....	51.3	44.3	39.2	41.8	48.8	64.2	108.7	176.3	374.7
2000.....	49.1	42.2	40.2	43.3	45.4	62.5	98.8	168.3	373.8
1999.....	47.9	40.1	40.3	41.3	49.9	65.4	101.3	184.2	387.1

Non-Hispanic White, female

2007 5/.....	32.3	29.7	23.7	22.7	26.0	37.5	65.2	117.2	278.4
2006 5/.....	29.7	26.9	22.7	22.1	26.4	39.0	65.3	116.4	264.9
2005 5/.....	28.3	23.7	21.3	22.8	28.0	39.6	64.2	114.4	265.6
2004 5/.....	26.9	22.2	19.8	22.3	26.0	37.7	64.1	114.4	263.1
2003 5/.....	24.3	21.4	19.5	19.7	25.4	38.3	62.9	113.1	262.8
2002.....	21.1	20.1	17.2	20.8	25.9	36.8	63.2	112.8	260.3
2001.....	18.3	17.5	17.4	19.4	25.6	35.9	63.0	109.1	253.8
2000.....	17.1	15.5	17.6	20.9	24.9	35.9	60.3	104.3	250.0
1999.....	16.9	14.9	17.5	18.0	25.9	37.1	62.5	107.7	255.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic Black, both sexes

2007 5/.....	50.2	56.8	52.0	43.6	46.9	54.7	69.0	84.3	156.2
2006 5/.....	55.1	57.6	52.4	45.0	46.1	52.0	72.5	95.5	153.2
2005 5/.....	50.4	58.4	50.1	46.3	48.5	60.2	87.6	112.5	176.7
2004 5/.....	50.1	51.6	45.9	42.5	44.4	53.7	78.6	102.3	155.1
2003 5/.....	49.0	49.8	44.6	40.8	48.6	54.1	68.4	105.0	161.8
2002.....	49.1	45.9	42.7	40.8	47.4	52.6	76.0	97.7	175.4
2001.....	50.8	45.7	41.2	43.9	45.9	58.0	79.2	105.7	183.9
2000.....	47.9	47.6	42.7	39.6	46.8	60.4	75.9	100.3	183.2
1999.....	51.3	44.5	44.0	46.4	51.3	67.5	90.7	125.0	208.9

Non-Hispanic Black, male

2007 5/.....	76.2	87.2	83.7	70.2	78.5	81.6	104.8	112.3	200.7
2006 5/.....	83.8	93.3	83.8	73.5	70.6	78.2	110.9	127.1	202.5
2005 5/.....	76.6	92.7	81.2	80.4	74.7	92.3	130.0	169.1	227.4
2004 5/.....	73.4	83.5	74.5	68.2	68.1	82.1	116.1	160.1	218.5
2003 5/.....	75.6	76.0	71.3	65.7	75.4	74.1	106.1	152.4	228.0
2002.....	73.0	73.6	72.1	65.4	75.5	79.4	119.2	151.3	262.5
2001.....	83.0	73.2	67.0	69.8	69.8	81.6	121.4	167.6	256.4
2000.....	76.2	77.4	72.6	65.3	75.2	87.1	120.4	156.2	277.1
1999.....	84.1	73.2	72.8	70.5	78.9	103.5	136.3	178.0	318.5

Non-Hispanic Black, female

2007 5/.....	27.6	31.1	26.0	22.8	23.8	36.2	47.4	69.6	137.8
2006 5/.....	30.1	27.5	26.6	22.9	28.0	34.1	49.5	78.9	133.0
2005 5/.....	27.6	29.4	24.5	19.7	29.0	38.2	62.3	82.7	156.3
2004 5/.....	29.8	24.7	22.4	22.5	26.8	34.5	56.3	71.9	130.1
2003 5/.....	25.8	27.6	22.6	21.3	28.7	40.7	46.1	80.2	136.2
2002.....	28.4	22.4	18.6	21.6	26.6	34.6	50.2	69.8	142.1
2001.....	23.0	22.3	20.0	23.5	28.3	42.2	54.0	74.1	157.0
2000.....	23.5	22.3	18.3	19.3	26.0	42.7	49.1	72.2	148.3
1999.....	23.0	20.2	20.5	27.5	30.9	43.7	63.2	98.8	167.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes

2007.....	16.7	16.1	15.4	14.1	15.2	18.1	21.3	25.0	24.1
2006.....	17.2	16.4	15.6	16.1	16.2	17.5	21.6	26.3	24.4
2005.....	16.8	16.6	16.2	16.3	17.2	18.9	23.0	26.0	26.1
2004.....	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
2003.....	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0

All origins 3/, male

2007.....	25.1	24.4	23.1	20.5	22.3	26.3	31.5	37.1	41.7
2006.....	25.7	24.5	22.9	22.9	22.9	23.9	30.1	37.5	42.9
2005.....	24.9	25.0	23.7	23.2	23.9	27.0	32.5	38.0	46.0
2004.....	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
2003.....	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5

All origins 3/, female

2007.....	8.4	8.2	8.1	8.2	9.0	11.4	13.7	17.3	15.7
2006.....	8.9	8.6	8.6	9.9	10.2	12.3	15.3	19.3	15.7
2005.....	8.9	8.6	9.0	10.0	11.3	12.4	16.1	18.5	17.0
2004.....	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
2003.....	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Hispanic, both sexes

2007	14.3	14.8	13.9	11.5	13.7	18.4	18.2	25.1	17.5
2006	15.7	15.6	15.2	15.8	17.6	17.0	22.9	17.3	11.0
2005	14.3	15.7	14.3	14.4	16.2	16.3	19.9	30.7	15.4
2004	14.4	14.2	14.2	18.2	15.9	18.4	19.4	25.9	20.0
2003	15.2	15.0	14.6	17.4	17.7	19.7	20.9	27.9	24.9
2002	15.3	15.0	13.6	17.3	16.6	20.6	27.5	24.1	25.5
2001	16.3	16.5	17.1	19.0	18.1	22.8	28.0	33.8	23.8
2000	15.8	16.7	15.9	15.3	16.4	25.6	23.3	25.1	25.2
1999	14.6	15.5	16.2	14.5	22.8	21.0	24.5	25.8	25.3

Hispanic, male

2007	21.5	22.9	20.7	15.7	22.1	27.9	26.8	36.4	29.5
2006	22.7	23.4	22.1	23.5	24.0	24.4	35.0	20.5	*
2005	21.4	23.3	21.1	20.8	24.8	27.0	27.0	49.0	*
2004	21.6	21.5	18.8	25.8	23.1	24.8	31.1	40.6	28.0
2003	22.9	21.7	20.4	22.9	25.3	27.3	30.4	43.7	46.2
2002	22.1	20.9	20.9	23.9	27.8	29.4	40.3	32.0	45.2
2001	22.4	24.1	21.4	29.1	25.9	31.1	45.7	61.0	43.8
2000	23.0	25.4	23.5	23.6	23.5	37.4	39.1	30.9	44.3
1999	22.0	23.4	21.7	20.0	32.8	30.0	35.9	48.7	*

Hispanic, female

2007	6.7	6.8	7.5	7.8	6.6	10.9	11.9	17.6	11.0
2006	8.4	7.7	8.7	8.9	12.3	11.3	14.2	15.3	*
2005	6.9	8.2	8.0	8.7	9.1	8.0	14.7	18.6	*
2004	6.9	7.0	10.0	11.6	10.0	13.4	11.0	16.2	15.8
2003	7.4	8.5	9.2	12.7	11.5	13.8	14.2	17.6	*
2002	8.4	9.2	6.9	11.6	7.5	13.9	18.3	19.0	*
2001	10.1	9.3	13.2	10.3	11.8	16.4	15.5	16.9	*
2000	8.7	8.3	9.1	8.2	10.6	16.6	12.0	21.5	*
1999	7.3	8.1	11.3	9.8	14.7	14.2	16.5	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	16.9	16.2	15.4	14.2	15.3	18.0	21.4	25.0	24.4
2006.....	17.3	16.4	15.5	16.1	16.0	17.5	21.4	26.7	25.0
2005.....	17.0	16.7	16.3	16.4	17.2	19.1	23.1	25.6	26.5
2004.....	17.3	16.4	15.3	15.7	17.0	18.0	23.8	28.1	27.5
2003.....	16.9	16.3	15.9	15.8	16.0	18.8	24.0	29.4	29.8
2002.....	16.9	15.8	14.9	16.5	16.6	20.1	24.8	29.7	28.9
2001.....	16.2	15.4	15.0	15.0	16.3	19.2	26.0	30.1	29.0
2000.....	16.1	15.2	15.6	16.3	16.7	18.9	25.1	30.0	31.7
1999.....	15.9	14.8	16.2	15.7	17.7	20.8	25.9	32.0	32.2

Non-Hispanic 4/, male

2007.....	25.5	24.5	23.2	20.8	22.3	26.1	31.7	37.1	42.4
2006.....	25.9	24.5	22.9	22.7	22.8	23.8	29.7	38.3	44.2
2005.....	25.2	25.1	23.9	23.3	23.8	27.0	32.8	37.3	47.1
2004.....	25.1	24.5	22.4	22.0	23.1	24.2	32.2	42.6	47.3
2003.....	25.0	23.6	23.3	22.5	22.1	24.7	33.0	42.0	52.0
2002.....	25.3	22.8	22.2	22.6	23.0	27.2	34.1	42.7	53.2
2001.....	24.0	22.3	21.3	21.3	21.8	26.4	35.7	44.6	52.9
2000.....	23.6	21.9	22.2	22.3	22.5	25.5	34.3	44.3	60.4
1999.....	23.1	21.3	22.9	21.8	24.2	27.2	36.1	47.7	64.3

Non-Hispanic 4/, female

2007.....	8.7	8.3	8.1	8.2	9.2	11.4	13.8	17.3	15.9
2006.....	8.9	8.6	8.6	10.0	10.0	12.3	15.3	19.5	16.1
2005.....	9.2	8.6	9.1	10.1	11.5	12.7	16.2	18.4	17.1
2004.....	9.7	8.7	8.6	9.9	11.6	13.1	17.7	19.2	18.6
2003.....	9.1	9.3	8.9	9.6	10.6	14.1	17.7	21.8	20.1
2002.....	8.9	9.1	8.0	11.0	11.1	14.3	18.3	21.9	18.4
2001.....	8.6	8.9	9.0	9.3	11.5	13.5	19.3	21.4	19.1
2000.....	8.9	8.9	9.5	10.9	11.8	13.8	18.8	21.7	20.1
1999.....	9.0	8.6	9.9	10.2	12.2	15.7	18.8	22.9	19.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Non-Hispanic White, both sexes

2007 5/.....	17.3	16.4	15.5	14.3	15.4	18.1	21.7	25.4	24.9
2006 5/.....	17.3	16.7	15.3	15.8	16.2	17.7	21.5	27.2	26.0
2005 5/.....	17.5	16.6	16.5	16.3	17.4	19.1	23.1	26.4	27.5
2004 5/.....	17.4	16.1	15.2	15.6	17.1	18.1	23.8	29.0	28.8
2003 5/.....	17.0	16.1	15.6	15.5	15.7	19.0	24.0	30.0	30.6
2002.....	16.9	15.8	14.8	16.5	16.3	20.3	25.3	30.4	29.7
2001.....	15.7	15.4	14.8	14.7	16.2	19.2	26.5	30.8	29.9
2000.....	15.9	14.8	15.4	16.4	16.5	18.7	25.3	30.9	32.4
1999.....	15.6	14.5	15.8	15.3	17.6	20.7	26.0	32.4	32.5

Non-Hispanic White, male

2007 5/.....	25.8	24.5	23.1	20.7	22.2	26.1	31.6	37.1	42.7
2006 5/.....	25.7	24.7	22.5	22.2	22.8	23.7	29.4	39.2	45.4
2005 5/.....	25.7	24.8	24.0	23.0	23.5	26.7	32.4	38.2	48.7
2004 5/.....	25.2	23.8	22.1	21.5	23.1	23.9	32.1	43.1	49.6
2003 5/.....	24.8	23.3	22.8	22.3	21.7	25.1	32.6	42.5	52.8
2002.....	25.4	22.5	21.7	22.4	22.1	27.0	33.9	42.9	54.0
2001.....	23.1	22.2	20.8	20.7	21.4	26.2	35.7	45.7	54.3
2000.....	23.2	21.1	21.4	21.9	21.6	24.9	34.0	44.8	62.3
1999.....	22.3	20.7	22.3	21.0	23.8	26.7	35.7	47.8	64.6

Non-Hispanic White, female

2007 5/.....	8.8	8.5	8.2	8.3	9.3	11.4	14.3	17.9	16.5
2006 5/.....	8.9	8.9	8.4	9.9	10.2	12.7	15.6	19.7	17.0
2005 5/.....	9.4	8.5	9.3	10.0	11.9	12.8	16.2	19.1	17.8
2004 5/.....	9.7	8.6	8.6	10.1	11.7	13.3	17.7	20.3	19.5
2003 5/.....	9.2	9.1	8.7	9.2	10.3	14.0	17.8	22.3	20.8
2002.....	8.6	9.2	8.1	10.9	11.2	14.8	19.1	22.8	19.2
2001.....	8.5	8.8	9.0	9.1	11.6	13.5	20.0	21.9	19.8
2000.....	8.7	8.5	9.6	11.3	12.0	13.7	19.2	22.6	20.4
1999.....	8.9	8.4	9.7	10.1	12.1	15.8	19.1	23.3	19.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Non-Hispanic Black, both sexes

2007 5/.....	18.1	17.9	17.2	15.8	14.9	18.2	18.7	19.1	17.9
2006 5/.....	19.2	17.1	19.4	18.8	15.7	15.6	20.0	20.4	15.9
2005 5/.....	17.2	19.3	17.1	17.6	17.6	19.3	22.3	16.6	16.9
2004 5/.....	19.6	20.2	17.6	17.7	16.6	16.1	23.1	19.7	15.0
2003 5/.....	18.5	18.6	19.2	17.9	18.8	17.5	23.2	21.8	21.4
2002.....	19.0	16.8	17.2	17.4	17.2	17.3	20.5	23.8	18.8
2001.....	20.7	17.2	18.1	18.9	18.9	18.7	22.3	19.9	17.8
2000.....	19.8	19.6	18.4	16.4	18.6	21.5	19.4	20.7	21.9
1999.....	20.0	18.3	19.1	17.8	18.2	21.1	24.7	28.2	27.8

Non-Hispanic Black, male

2007 5/.....	29.5	29.5	27.8	26.4	25.0	29.3	34.6	35.4	39.3
2006 5/.....	30.4	28.1	30.0	30.1	25.6	25.6	33.6	28.8	37.6
2005 5/.....	25.9	31.6	27.8	28.3	29.4	31.2	38.1	28.1	34.3
2004 5/.....	30.2	33.0	27.5	29.1	26.7	25.9	36.8	41.1	29.5
2003 5/.....	30.0	29.4	30.8	28.0	28.9	23.3	38.3	37.7	43.7
2002.....	29.2	26.6	29.6	27.3	28.4	29.1	39.0	43.7	47.0
2001.....	34.5	26.1	29.3	31.3	28.9	25.5	39.8	31.9	35.8
2000.....	30.0	30.3	31.5	27.2	31.7	33.8	34.7	39.5	45.8
1999.....	31.8	29.2	29.8	28.7	27.4	32.2	40.6	48.7	68.9

Non-Hispanic Black, female

2007 5/.....	8.1	8.1	8.5	7.5	7.5	10.5	9.0	10.6	9.0
2006 5/.....	9.5	7.8	10.6	9.9	8.4	8.7	11.8	15.9	*
2005 5/.....	9.7	9.0	8.3	9.3	8.9	11.1	12.8	10.6	10.0
2004 5/.....	10.3	9.3	9.4	8.8	9.1	9.5	14.9	8.5	9.2
2003 5/.....	8.6	9.5	9.7	10.0	11.3	13.6	14.1	13.4	12.8
2002.....	10.2	8.4	7.0	9.7	8.9	9.4	9.5	13.4	*
2001.....	8.8	9.6	8.8	9.2	11.5	14.2	11.9	13.7	11.1
2000.....	11.0	10.5	7.8	7.9	8.8	13.4	10.2	11.2	13.1
1999.....	9.7	9.1	10.4	9.2	11.4	13.8	15.1	18.0	12.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes

2007.....	15.2	14.6	13.8	12.7	14.0	16.7	20.3	23.8	23.2
2006.....	15.7	14.8	14.0	14.7	14.9	16.1	20.2	25.0	23.4
2005.....	15.3	14.8	14.6	14.8	15.8	17.8	21.6	24.6	25.1
2004.....	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
2003.....	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All origins 3/, male

2007.....	22.5	21.8	20.5	18.0	20.1	23.6	29.7	35.0	39.8
2006.....	23.2	22.0	20.5	20.4	20.8	21.5	27.8	35.0	41.0
2005.....	22.2	22.1	21.0	20.7	21.5	25.2	30.1	35.7	44.1
2004.....	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
2003.....	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All origins 3/, female

2007.....	8.0	7.6	7.6	7.8	8.7	11.0	13.4	16.8	15.2
2006.....	8.4	8.0	7.9	9.4	9.6	11.7	14.6	18.7	15.2
2005.....	8.5	7.9	8.5	9.4	10.7	11.8	15.5	17.8	16.4
2004.....	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
2003.....	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Hispanic, both sexes

2007.....	13.0	13.6	12.6	10.3	12.6	17.2	16.5	24.2	15.4
2006.....	14.7	14.6	14.5	15.0	16.6	16.4	21.2	16.7	10.6
2005.....	13.2	14.6	13.3	13.9	15.0	16.1	18.3	29.0	15.0
2004.....	13.3	13.1	13.5	16.8	15.0	17.7	18.4	25.9	19.5
2003.....	14.1	14.2	13.4	16.2	16.4	18.7	20.4	27.5	24.4
2002.....	14.2	13.6	12.7	16.6	16.1	20.0	25.6	22.3	24.9
2001.....	13.8	13.8	14.4	16.0	15.2	20.8	23.5	31.3	20.0
2000.....	14.8	15.4	14.7	14.0	14.2	23.7	21.7	23.4	22.6
1999.....	13.3	14.1	15.5	13.4	20.6	18.6	23.2	24.1	23.2

Hispanic, male

2007.....	19.1	20.8	18.0	14.0	19.7	25.3	23.3	35.6	27.4
2006.....	20.9	21.8	21.1	22.1	22.3	23.7	31.9	18.8	*
2005.....	19.4	21.8	19.0	19.9	23.0	26.6	23.8	45.5	*
2004.....	19.6	19.6	17.4	23.0	21.3	23.6	28.8	40.6	28.0
2003.....	21.1	20.3	18.8	21.2	23.4	26.5	29.2	42.6	46.2
2002.....	20.4	18.6	19.3	22.3	26.7	28.5	37.8	28.6	43.5
2001.....	19.9	20.5	18.6	24.7	21.6	28.4	38.2	55.8	*
2000.....	21.0	23.3	21.5	20.7	20.1	33.5	36.9	*	*
1999.....	19.7	20.6	20.1	17.6	29.0	25.0	32.8	45.6	*

Hispanic, female

2007.....	6.5	6.5	7.5	7.1	6.6	10.9	11.5	16.5	*
2006.....	8.1	7.4	8.3	8.7	11.8	10.7	13.4	15.3	*
2005.....	6.7	7.5	7.9	8.7	8.4	8.0	14.4	18.0	*
2004.....	6.7	6.8	9.8	11.4	9.7	13.1	11.0	16.2	15.1
2003.....	7.0	8.3	8.4	11.8	10.7	12.8	14.2	17.6	*
2002.....	7.9	8.7	6.6	11.6	7.5	13.5	16.9	18.2	*
2001.....	7.7	7.4	10.7	8.5	10.0	15.0	13.1	16.1	*
2000.....	8.6	7.9	8.5	8.2	9.4	16.2	11.0	21.5	*
1999.....	7.0	7.9	11.3	9.8	13.8	13.8	16.5	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic 4/, both sexes

2007.....	15.4	14.6	13.9	12.8	14.1	16.6	20.5	23.8	23.6
2006.....	15.8	14.8	14.0	14.6	14.7	16.0	20.1	25.3	24.1
2005.....	15.5	14.8	14.7	14.8	15.8	17.9	21.8	24.3	25.5
2004.....	15.7	14.7	13.9	14.1	15.5	16.7	22.9	26.9	26.3
2003.....	15.1	14.6	14.1	14.0	14.6	17.5	22.6	28.2	28.9
2002.....	15.2	14.4	13.4	15.0	15.2	18.8	23.6	28.5	28.1
2001.....	14.5	13.8	13.6	13.5	14.7	17.7	24.6	28.7	28.0
2000.....	14.5	13.6	14.1	14.7	15.1	17.7	23.9	28.6	30.6
1999.....	14.0	13.0	14.2	13.9	16.1	19.2	24.1	30.1	30.5

Non-Hispanic 4/, male

2007.....	22.9	21.8	20.7	18.3	20.1	23.4	29.9	35.0	40.6
2006.....	23.4	21.9	20.4	20.2	20.6	21.3	27.5	35.7	42.3
2005.....	22.5	22.1	21.1	20.7	21.4	25.1	30.4	35.1	45.2
2004.....	22.5	21.7	20.1	19.3	20.7	21.8	30.8	40.3	45.0
2003.....	22.1	20.7	20.3	19.4	19.7	22.3	30.5	39.8	49.9
2002.....	22.2	20.3	19.5	19.9	20.3	24.6	31.9	40.0	51.9
2001.....	21.2	19.5	18.9	18.6	19.0	23.8	33.3	42.4	51.6
2000.....	21.1	19.2	19.6	19.5	19.7	23.2	32.0	41.7	58.3
1999.....	19.9	18.3	19.5	18.7	21.5	24.6	33.5	44.7	60.7

Non-Hispanic 4/, female

2007.....	8.2	7.7	7.5	7.8	8.8	11.0	13.5	16.8	15.6
2006.....	8.4	8.0	7.9	9.4	9.4	11.8	14.7	18.9	15.6
2005.....	8.6	7.9	8.6	9.4	10.9	12.0	15.5	17.7	16.5
2004.....	9.1	8.1	8.1	9.3	11.0	12.5	17.3	18.7	18.0
2003.....	8.4	8.8	8.3	9.2	10.2	13.7	17.0	21.1	19.6
2002.....	8.5	8.7	7.6	10.4	10.8	14.1	17.8	21.5	17.8
2001.....	8.0	8.3	8.6	8.8	11.0	12.9	18.4	20.4	18.2
2000.....	8.2	8.2	9.0	10.3	11.1	13.3	18.2	20.9	19.4
1999.....	8.2	7.9	9.2	9.5	11.4	15.1	17.5	21.6	18.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic White, both sexes

2007 5/.....	15.8	14.7	14.0	12.8	14.2	16.6	20.8	24.2	24.1
2006 5/.....	15.7	15.1	13.7	14.2	14.7	16.2	20.2	25.8	25.1
2005 5/.....	15.9	14.8	14.8	14.6	15.9	17.8	21.8	25.1	26.6
2004 5/.....	15.8	14.4	13.8	14.0	15.6	16.7	23.0	27.7	27.6
2003 5/.....	15.2	14.3	13.7	13.7	14.3	17.6	22.6	28.7	29.6
2002.....	15.2	14.3	13.3	14.8	15.0	19.0	24.1	29.2	28.9
2001.....	14.1	13.8	13.4	13.1	14.7	17.6	25.0	29.4	29.0
2000.....	14.3	13.0	13.9	14.7	14.9	17.4	24.1	29.3	31.3
1999.....	13.6	12.7	13.8	13.5	15.9	19.1	24.2	30.6	30.9

Non-Hispanic White, male

2007 5/.....	23.2	21.8	20.7	18.0	20.1	23.4	29.9	35.0	40.8
2006 5/.....	23.1	22.1	20.0	19.6	20.6	21.1	27.0	36.4	43.6
2005 5/.....	23.0	21.9	21.0	20.3	21.2	24.7	30.1	35.9	46.7
2004 5/.....	22.5	20.9	19.7	18.7	20.5	21.5	30.7	40.6	47.1
2003 5/.....	21.9	20.3	19.7	19.0	19.4	22.6	30.2	40.4	50.7
2002.....	22.3	20.0	19.1	19.7	19.5	24.5	31.9	40.2	52.6
2001.....	20.3	19.4	18.5	17.8	18.7	23.5	33.3	43.5	53.1
2000.....	20.7	18.3	18.9	18.9	18.8	22.7	31.7	42.0	60.0
1999.....	19.1	17.7	18.8	18.0	21.1	24.2	33.2	44.9	61.2

Non-Hispanic White, female

2007 5/.....	8.4	7.8	7.6	7.9	8.9	10.9	13.9	17.4	16.2
2006 5/.....	8.3	8.2	7.6	9.2	9.5	12.0	15.0	19.1	16.5
2005 5/.....	8.9	7.8	8.7	9.3	11.2	12.2	15.6	18.4	17.3
2004 5/.....	9.1	8.0	8.1	9.6	11.1	12.8	17.3	19.7	18.9
2003 5/.....	8.6	8.5	8.0	8.8	9.9	13.5	17.2	21.6	20.3
2002.....	8.2	8.7	7.7	10.3	11.0	14.5	18.5	22.4	18.6
2001.....	8.0	8.2	8.6	8.8	11.1	12.8	19.2	21.0	19.0
2000.....	8.0	7.8	9.1	10.7	11.4	13.2	18.7	21.8	19.8
1999.....	8.1	7.7	9.0	9.3	11.4	15.1	17.9	22.1	18.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic Black, both sexes

2007 5/.....	16.3	16.4	15.5	14.4	13.9	16.6	17.8	18.1	17.6
2006 5/.....	17.6	15.6	17.8	18.0	14.8	14.9	18.8	19.4	14.3
2005 5/.....	15.3	17.0	16.1	16.6	16.3	18.5	21.2	15.4	16.1
2004 5/.....	17.8	18.6	16.3	15.8	15.6	15.2	21.0	19.5	15.0
2003 5/.....	16.6	17.2	17.9	16.4	17.2	17.0	21.6	20.5	21.1
2002.....	17.0	15.5	15.3	16.3	15.8	16.2	18.8	22.4	18.5
2001.....	18.5	15.7	16.2	17.2	16.8	18.0	20.4	19.0	16.9
2000.....	18.0	18.6	17.0	15.5	16.9	20.3	18.1	19.8	20.7
1999.....	18.2	16.4	17.4	15.8	16.7	19.7	22.9	25.8	25.5

Non-Hispanic Black, male

2007 5/.....	26.2	26.8	24.6	23.9	22.9	25.8	32.8	33.3	39.3
2006 5/.....	27.4	25.2	27.1	28.3	23.9	24.1	31.8	28.1	33.9
2005 5/.....	22.8	27.4	25.8	26.4	26.2	30.2	35.8	26.6	33.3
2004 5/.....	27.3	30.0	25.1	25.6	25.2	23.9	33.6	40.4	29.5
2003 5/.....	26.6	26.5	28.1	24.8	25.6	22.2	36.0	34.7	42.6
2002.....	25.5	24.0	25.8	24.8	25.1	26.3	34.3	40.6	45.9
2001.....	30.6	23.7	25.3	28.8	24.5	23.8	37.0	31.1	35.8
2000.....	27.0	28.5	28.6	25.5	28.2	31.8	32.3	38.6	44.6
1999.....	28.8	25.8	26.7	24.8	25.0	29.1	39.2	45.2	61.8

Non-Hispanic Black, female

2007 5/.....	7.8	7.7	8.1	7.1	7.3	10.3	8.8	10.2	8.7
2006 5/.....	9.1	7.6	10.0	9.9	8.0	8.7	11.0	14.8	*
2005 5/.....	8.7	8.3	8.0	8.9	8.9	10.4	12.6	9.5	9.2
2004 5/.....	9.5	9.0	9.1	8.1	8.5	9.2	13.5	8.5	9.2
2003 5/.....	8.0	9.3	9.5	9.9	11.1	13.4	13.0	13.0	12.8
2002.....	9.6	8.3	6.8	9.5	8.9	9.4	9.5	13.0	*
2001.....	8.1	8.9	8.7	8.1	11.2	14.2	10.5	12.9	9.8
2000.....	10.3	10.2	7.5	7.5	8.5	12.7	9.6	10.4	11.8
1999.....	9.0	8.5	9.8	8.7	10.6	13.6	13.1	16.2	12.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins 3/, both sexes

2007.....	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.6	*
2006.....	0.5	0.6	0.5	0.4	0.5	0.5	0.6	0.6	0.5
2005.....	0.5	0.7	0.5	0.5	0.4	0.4	0.6	0.6	0.5
2004.....	0.6	0.5	0.4	0.5	0.4	0.6	0.4	0.6	0.6
2003.....	0.6	0.5	0.6	0.5	0.4	0.5	0.6	0.7	0.5
2002.....	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	*
2001.....	0.7	0.6	0.5	0.6	0.7	0.6	0.8	0.8	0.7
2000.....	0.6	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.7	0.8	0.8	0.8	0.7	1.3	1.4	1.3

All origins 3/, male

2007.....	0.9	0.9	0.7	0.6	0.7	0.8	0.6	0.9	*
2006.....	0.9	1.0	0.8	0.7	0.7	1.0	1.0	1.1	*
2005.....	0.9	1.2	0.7	0.8	0.7	0.6	1.1	1.1	*
2004.....	0.9	0.8	0.7	0.7	0.6	0.8	0.6	1.2	*
2003.....	1.1	0.8	0.9	0.8	0.8	0.7	0.8	1.3	*
2002.....	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.2	*
2001.....	1.0	0.9	0.8	0.9	1.1	0.9	1.2	*	*
2000.....	0.9	0.9	0.7	1.0	0.9	0.9	1.1	1.3	*
1999.....	1.4	1.2	1.2	1.1	1.2	1.2	1.7	2.1	2.9

All origins 3/, female

2007.....	0.2	*	*	*	*	*	*	*	*
2006.....	0.2	0.2	0.3	*	*	*	*	*	*
2005.....	*	0.4	*	*	*	*	*	*	*
2004.....	0.2	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.4	0.2	*	*	*	*	0.5	0.7	*
2000.....	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.3	0.4	0.4	0.5	0.4	0.9	1.0	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic 4/, both sexes

2007.....	0.5	0.5	0.4	0.3	0.3	0.4	0.3	0.6	*
2006.....	0.5	0.6	0.5	0.4	0.5	0.5	0.6	0.6	0.5
2005.....	0.5	0.8	0.4	0.5	0.4	0.5	0.6	0.6	0.5
2004.....	0.5	0.5	0.5	0.4	0.4	0.6	0.4	0.6	0.6
2003.....	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.6	0.4	0.4	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.6	0.5	0.4	0.6	0.7	0.6	0.7	0.7	0.6
2000.....	0.6	0.6	0.4	0.5	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.7	0.8	0.7	0.7	0.7	1.3	1.3	1.3

Non-Hispanic 4/, male

2007.....	0.9	0.9	0.6	0.6	0.6	0.8	*	1.0	*
2006.....	0.8	1.0	0.8	0.7	0.7	1.0	1.0	1.1	*
2005.....	0.9	1.2	0.7	0.8	0.7	0.6	1.1	1.0	*
2004.....	0.9	0.8	0.7	0.7	0.5	0.8	*	1.3	*
2003.....	1.1	0.9	0.9	0.8	0.7	0.8	0.9	1.3	*
2002.....	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.1	*
2001.....	1.0	0.9	0.8	0.8	1.2	0.9	1.0	*	*
2000.....	0.9	0.8	0.7	0.9	0.8	0.9	1.1	1.2	*
1999.....	1.4	1.1	1.3	1.1	1.0	1.0	1.6	2.0	3.0

Non-Hispanic 4/, female

2007.....	0.2	*	*	*	*	*	*	*	*
2006.....	*	0.2	0.3	*	*	*	*	*	*
2005.....	*	0.4	*	*	*	*	*	*	*
2004.....	0.2	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	0.6	*
2000.....	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.3	0.4	0.4	0.5	0.4	1.0	1.0	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic White, both sexes

2007 5/.....	0.5	0.5	0.3	0.3	0.3	0.4	0.4	0.5	*
2006 5/.....	0.5	0.5	0.5	0.4	0.5	0.6	0.5	0.6	0.5
2005 5/.....	0.5	0.6	0.5	0.4	0.4	0.4	0.6	0.6	0.5
2004 5/.....	0.5	0.4	0.5	0.4	0.4	0.6	*	0.7	0.6
2003 5/.....	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.5	0.4	0.3	0.4	0.3	0.4	0.4	0.5	*
2001.....	0.6	0.5	0.4	0.5	0.6	0.6	0.7	0.7	0.5
2000.....	0.5	0.6	0.4	0.5	0.4	0.5	0.6	0.8	0.8
1999.....	0.9	0.7	0.8	0.7	0.7	0.7	1.2	1.3	1.3

Non-Hispanic White, male

2007 5/.....	0.9	0.9	0.5	0.6	0.6	0.7	*	*	*
2006 5/.....	0.8	0.8	0.8	0.6	0.7	1.0	1.0	1.2	*
2005 5/.....	0.8	1.0	0.7	0.6	*	*	1.0	*	*
2004 5/.....	0.7	0.6	0.7	0.6	0.5	0.8	*	1.4	*
2003 5/.....	0.9	0.8	0.8	0.8	0.7	0.7	0.8	1.2	*
2002.....	0.9	0.8	0.6	0.7	*	0.7	*	1.2	*
2001.....	0.9	0.9	0.7	0.8	1.0	0.9	1.0	*	*
2000.....	0.8	0.8	0.5	1.0	0.6	0.9	1.1	1.3	*
1999.....	1.3	1.0	1.3	1.1	1.0	0.9	1.6	1.9	2.8

Non-Hispanic White, female

2007 5/.....	*	*	*	*	*	*	*	*	*
2006 5/.....	*	*	0.3	*	*	*	*	*	*
2005 5/.....	*	0.3	*	*	*	*	*	*	*
2004 5/.....	*	*	*	*	*	*	*	*	*
2003 5/.....	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.5	0.3	0.4	0.5	*	*	0.9	1.0	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins 3/, both sexes

2007.....	0.9	1.1	1.1	1.1	0.8	1.0	0.6	0.6	0.6
2006.....	0.9	0.9	1.0	1.0	0.8	0.9	0.8	0.7	0.4
2005.....	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.7	0.5
2004.....	1.0	1.2	0.9	1.1	1.0	0.8	0.5	0.5	0.5
2003.....	1.1	1.1	1.2	1.2	0.9	0.8	0.8	0.5	*
2002.....	1.1	1.0	1.1	1.1	1.0	0.8	0.8	0.7	0.4
2001.....	1.1	1.2	1.0	1.1	1.0	0.9	0.8	0.7	0.5
2000.....	0.9	1.0	1.0	1.1	1.2	0.7	0.6	0.7	*
1999.....	1.0	1.0	1.1	1.1	0.9	0.8	0.5	0.6	*

All origins 3/, male

2007.....	1.6	1.7	1.9	1.8	1.5	1.9	1.2	1.2	1.1
2006.....	1.6	1.5	1.6	1.7	1.4	1.4	1.3	1.4	*
2005.....	1.7	1.8	1.9	1.7	1.7	1.3	1.3	1.2	*
2004.....	1.7	2.0	1.5	2.1	1.8	1.5	0.9	1.0	*
2003.....	1.7	2.0	2.0	2.3	1.7	1.5	1.6	*	*
2002.....	1.9	1.6	1.9	1.9	2.0	1.7	1.4	1.5	*
2001.....	1.8	1.9	1.7	1.9	1.8	1.7	1.3	1.4	*
2000.....	1.6	1.8	1.8	1.9	2.0	1.4	1.2	1.4	*
1999.....	1.7	1.8	2.0	1.9	1.6	1.6	0.9	*	*

All origins 3/, female

2007.....	0.2	0.4	0.4	0.4	*	*	*	*	*
2006.....	0.3	0.3	0.4	0.4	*	*	*	*	*
2005.....	0.3	0.3	0.3	0.4	0.4	*	*	*	*
2004.....	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.4	0.3	0.4	*	*	*	*	*	*
2002.....	0.3	0.3	0.3	0.4	*	*	*	*	*
2001.....	0.4	0.4	0.4	*	*	*	0.5	*	*
2000.....	0.3	0.3	*	0.4	0.4	*	*	*	*
1999.....	0.3	0.3	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic 4/, both sexes

2007.....	0.9	1.1	1.2	1.1	0.9	1.0	0.6	0.6	0.5
2006.....	1.0	1.0	1.0	1.1	0.9	0.9	0.8	0.8	0.5
2005.....	1.1	1.1	1.2	1.1	1.0	0.8	0.7	0.7	0.5
2004.....	1.1	1.2	0.9	1.2	1.1	0.8	0.5	0.6	0.6
2003.....	1.1	1.2	1.2	1.3	1.0	0.8	0.8	0.5	*
2002.....	1.1	1.0	1.1	1.1	1.0	0.9	0.8	0.7	*
2001.....	1.1	1.1	1.0	1.0	0.9	0.9	0.8	0.7	*
2000.....	1.0	1.1	1.1	1.1	1.2	0.7	0.6	0.7	*
1999.....	1.0	1.1	1.2	1.1	0.9	0.9	0.5	0.6	*

Non-Hispanic 4/, male

2007.....	1.7	1.8	1.9	1.9	1.6	1.9	1.2	1.2	*
2006.....	1.7	1.7	1.7	1.8	1.4	1.5	1.3	1.5	*
2005.....	1.8	1.9	2.0	1.8	1.7	1.3	1.3	1.3	*
2004.....	1.8	2.1	1.5	2.1	1.9	1.5	0.9	1.0	*
2003.....	1.9	2.1	2.2	2.4	1.7	1.6	1.6	*	*
2002.....	2.1	1.7	2.0	2.0	2.1	1.8	1.4	1.6	*
2001.....	1.9	1.9	1.7	1.9	1.6	1.7	1.3	1.3	*
2000.....	1.7	1.9	1.9	1.9	2.1	1.4	1.2	1.3	*
1999.....	1.8	1.9	2.1	2.0	1.7	1.6	0.9	*	*

Non-Hispanic 4/, female

2007.....	0.2	0.4	0.4	0.3	*	*	*	*	*
2006.....	0.3	0.3	0.4	0.4	*	*	*	*	*
2005.....	0.4	0.3	0.3	0.4	*	*	*	*	*
2004.....	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.4	0.3	0.3	*	*	*	*	*	*
2002.....	0.3	0.4	0.3	0.4	*	*	*	*	*
2001.....	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	0.4	0.5	*	*	*	*
1999.....	0.3	0.4	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic White, both sexes

2007 5/.....	1.0	1.2	1.2	1.2	0.9	1.1	0.6	0.6	0.5
2006 5/.....	1.1	1.1	1.1	1.2	0.9	1.0	0.8	0.8	*
2005 5/.....	1.1	1.1	1.3	1.2	1.1	0.9	0.8	0.7	0.5
2004 5/.....	1.2	1.3	1.0	1.2	1.1	0.8	0.5	0.6	0.6
2003 5/.....	1.2	1.3	1.3	1.3	0.9	0.9	0.8	0.5	*
2002.....	1.2	1.1	1.2	1.2	1.0	0.9	0.8	0.7	*
2001.....	1.1	1.1	0.9	1.1	0.9	0.9	0.8	0.7	*
2000.....	1.1	1.2	1.1	1.2	1.2	0.7	0.6	0.7	*
1999.....	1.1	1.1	1.2	1.1	1.0	0.9	0.5	0.5	*

Non-Hispanic White, male

2007 5/.....	1.7	1.8	2.0	2.0	1.6	1.9	1.2	1.2	*
2006 5/.....	1.8	1.8	1.8	2.0	1.5	1.6	1.3	1.6	*
2005 5/.....	1.8	1.9	2.2	2.0	1.9	1.5	1.4	1.2	*
2004 5/.....	2.0	2.3	1.6	2.2	2.1	1.6	1.0	1.1	*
2003 5/.....	2.0	2.2	2.3	2.6	1.7	1.8	1.6	*	*
2002.....	2.3	1.7	2.0	2.1	2.1	1.8	1.3	1.6	*
2001.....	1.9	2.0	1.6	2.0	1.6	1.8	1.3	1.4	*
2000.....	1.7	2.0	2.0	2.0	2.1	1.4	1.2	1.4	*
1999.....	1.9	1.9	2.1	2.0	1.7	1.6	1.0	*	*

Non-Hispanic White, female

2007 5/.....	0.3	0.5	0.4	0.3	*	*	*	*	*
2006 5/.....	0.4	0.4	0.5	0.4	*	*	*	*	*
2005 5/.....	0.3	0.4	0.4	0.5	*	*	*	*	*
2004 5/.....	0.4	0.4	0.3	*	*	*	*	*	*
2003 5/.....	0.4	0.4	0.4	*	*	*	*	*	*
2002.....	*	0.4	0.4	0.4	*	*	*	*	*
2001.....	*	0.3	*	*	*	*	*	*	*
2000.....	0.4	*	*	0.5	*	*	*	*	*
1999.....	0.4	0.3	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes

2007.....	30.2	29.5	23.6	21.1	23.7	35.0	59.6	112.1	262.6
2006.....	29.9	27.4	21.6	18.9	23.4	33.0	59.8	109.7	250.5
2005.....	27.8	25.0	19.4	19.8	22.9	34.8	59.7	111.0	253.4
2004.....	25.5	22.3	17.6	17.8	21.4	32.4	56.0	107.8	249.5
2003.....	23.9	20.3	17.0	16.9	21.9	31.9	54.9	104.1	249.2
2002.....	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001.....	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000.....	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
1999.....	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3

All origins 3/, male

2007.....	40.1	40.2	32.8	28.8	32.4	47.2	74.8	142.6	307.3
2006.....	41.1	38.5	30.0	26.8	32.2	42.5	77.2	139.0	294.3
2005.....	38.7	35.7	27.5	28.3	30.7	45.6	77.5	142.8	298.3
2004.....	35.6	32.1	24.7	24.5	29.6	42.8	71.7	137.0	299.0
2003.....	34.1	28.7	24.0	24.3	29.7	41.8	72.0	133.0	300.5
2002.....	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001.....	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000.....	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
1999.....	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9

All origins 3/, female

2007.....	20.5	19.4	14.9	14.0	16.0	25.0	48.3	92.9	241.4
2006.....	18.9	16.7	13.6	11.8	15.8	25.3	47.0	91.4	230.1
2005.....	17.1	14.7	11.7	12.0	16.0	26.1	46.7	91.3	232.8
2004.....	15.8	12.8	10.8	11.7	14.3	24.1	44.8	89.9	227.3
2003.....	14.1	12.3	10.4	10.2	15.1	23.9	42.7	86.4	226.5
2002.....	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001.....	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000.....	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
1999.....	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Hispanic, both sexes

2007	19.9	21.0	16.6	16.7	17.4	25.9	43.6	74.1	134.4
2006	20.5	19.3	18.9	16.2	17.4	23.9	40.9	74.3	131.9
2005	19.4	18.8	15.2	17.0	18.9	28.5	43.6	74.0	138.2
2004	17.6	15.7	14.5	14.8	17.3	24.0	40.2	71.3	140.8
2003	18.7	16.5	13.3	17.1	17.3	22.4	38.1	62.3	127.0
2002	17.7	15.6	13.7	13.1	18.1	24.8	42.4	65.9	127.5
2001	16.3	14.0	12.6	12.6	14.4	26.5	34.8	59.1	142.6
2000	14.7	13.5	12.4	14.0	14.8	22.8	35.8	70.7	159.2
1999	17.8	14.1	11.2	15.3	19.5	21.6	34.5	62.8	157.2

Hispanic, male

2007	31.0	34.3	23.9	24.9	25.1	37.0	57.5	92.2	156.5
2006	30.9	31.2	30.3	25.8	27.9	32.8	49.4	93.4	155.6
2005	30.9	30.5	24.4	27.6	27.7	39.3	60.0	88.4	146.9
2004	27.7	24.4	22.6	24.0	25.6	31.5	51.3	86.8	165.3
2003	30.4	25.2	21.7	27.1	24.4	30.7	58.3	82.1	149.3
2002	27.9	26.1	22.2	20.3	28.4	32.5	57.0	89.2	179.0
2001	27.5	23.1	21.2	19.3	20.2	35.2	47.1	67.5	154.3
2000	26.1	22.0	18.9	20.7	23.5	33.1	47.2	78.0	207.6
1999	30.9	23.2	20.6	25.0	29.8	34.1	42.1	80.2	217.8

Hispanic, female

2007	8.2	7.7	9.6	9.4	10.9	17.2	33.5	62.1	122.5
2006	9.5	7.4	8.2	7.6	8.5	16.9	34.7	61.6	119.2
2005	7.5	7.2	6.5	7.7	11.6	20.0	31.8	64.5	133.6
2004	7.0	7.1	7.0	6.7	10.5	18.2	32.3	61.1	128.2
2003	6.6	8.0	5.4	8.4	11.5	16.1	23.7	49.4	115.6
2002	7.5	5.4	5.9	6.9	9.7	18.9	32.0	51.1	102.0
2001	5.1	5.3	4.9	6.8	9.8	19.9	26.1	53.9	136.8
2000	3.4	5.5	6.5	8.2	7.9	15.1	27.7	66.3	135.5
1999	4.9	5.5	*	6.9	11.3	12.3	29.2	52.5	127.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	31.4	30.3	24.1	21.4	24.1	35.6	60.5	114.3	269.2
2006.....	30.9	28.0	21.7	19.0	23.9	33.6	60.9	111.4	256.3
2005.....	28.6	25.4	19.6	19.9	23.1	35.2	60.6	112.8	258.6
2004.....	26.3	22.7	17.7	18.0	21.6	32.9	56.9	109.5	254.1
2003.....	24.3	20.5	17.2	16.8	22.1	32.4	55.7	105.9	253.9
2002.....	22.1	18.6	15.5	16.1	21.8	30.8	55.4	102.7	250.9
2001.....	19.8	16.4	14.1	16.2	20.6	29.8	55.1	101.4	250.8
2000.....	18.1	15.2	14.2	15.9	18.8	29.7	50.8	95.3	244.4
1999.....	18.0	14.1	13.8	15.4	20.5	30.4	53.3	102.3	252.8

Non-Hispanic 4/, male

2007.....	41.2	40.6	33.4	29.0	32.9	47.9	75.8	145.5	315.6
2006.....	42.2	39.1	29.9	26.6	32.4	43.1	78.8	141.2	301.7
2005.....	39.5	36.0	27.6	28.1	30.8	45.9	78.4	145.6	305.9
2004.....	36.3	32.6	24.8	24.4	29.7	43.4	72.8	139.4	305.1
2003.....	34.2	28.7	24.0	24.0	29.9	42.4	72.4	135.3	306.9
2002.....	31.7	26.6	22.3	22.9	29.8	41.7	72.4	128.6	298.0
2001.....	29.9	24.2	20.1	22.6	28.1	38.8	73.1	129.8	311.6
2000.....	27.8	23.5	20.6	22.7	25.4	38.5	65.9	122.2	302.7
1999.....	28.0	21.6	20.2	22.3	28.0	40.9	67.4	134.9	314.3

Non-Hispanic 4/, female

2007.....	22.0	20.6	15.3	14.3	16.5	25.6	49.2	94.7	247.3
2006.....	20.0	17.5	14.0	12.1	16.4	25.9	47.8	92.9	235.2
2005.....	18.1	15.4	12.1	12.4	16.4	26.4	47.6	92.5	237.0
2004.....	16.7	13.3	11.1	12.1	14.5	24.5	45.4	91.1	231.2
2003.....	14.8	12.7	10.8	10.3	15.3	24.4	43.7	87.9	230.6
2002.....	12.9	10.8	9.1	9.9	14.8	22.1	43.5	86.9	230.6
2001.....	10.1	8.9	8.4	10.5	14.1	22.7	42.5	84.4	225.5
2000.....	8.8	7.3	8.1	9.7	13.1	22.7	40.4	79.5	220.8
1999.....	8.4	6.9	7.9	9.1	14.1	22.1	43.4	83.2	228.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Non-Hispanic White, both sexes

2007 5/.....	32.8	30.4	23.4	21.1	23.6	36.2	62.1	120.2	285.0
2006 5/.....	31.5	27.3	20.9	18.5	23.5	33.8	62.6	116.2	269.8
2005 5/.....	29.1	24.4	18.5	19.1	22.6	35.1	61.2	115.2	269.7
2004 5/.....	26.9	22.4	16.9	17.5	21.4	32.8	57.9	113.0	266.4
2003 5/.....	24.4	19.6	16.6	16.5	21.5	32.3	57.0	108.4	265.9
2002.....	21.8	17.7	14.6	15.6	20.9	30.7	56.0	105.6	261.2
2001.....	19.0	15.3	13.3	15.5	20.3	29.4	55.4	103.6	259.4
2000.....	17.2	13.9	13.2	15.3	18.0	29.1	50.9	97.3	253.2
1999.....	16.7	12.9	12.8	13.9	19.5	29.0	52.5	103.7	260.8

Non-Hispanic White, male

2007 5/.....	42.2	39.8	31.6	28.3	31.3	48.2	77.0	152.8	333.6
2006 5/.....	42.2	36.8	27.7	25.3	31.8	42.8	79.9	147.1	316.9
2005 5/.....	39.3	33.8	25.3	25.9	29.8	45.0	78.9	147.0	317.6
2004 5/.....	36.7	31.3	22.7	23.1	29.3	43.0	73.7	143.3	317.3
2003 5/.....	33.8	27.1	22.7	23.0	28.7	42.0	73.5	137.2	320.0
2002.....	31.1	24.8	20.4	21.7	28.0	41.3	72.6	131.2	307.6
2001.....	28.2	22.0	18.4	21.1	27.5	38.0	73.0	130.7	320.4
2000.....	25.9	21.1	18.8	21.4	23.8	37.6	64.9	123.5	311.5
1999.....	25.6	19.4	18.0	20.3	26.0	38.7	65.5	136.4	322.6

Non-Hispanic White, female

2007 5/.....	23.4	21.2	15.5	14.3	16.7	26.1	50.9	99.3	261.9
2006 5/.....	20.9	18.0	14.3	12.2	16.2	26.3	49.7	96.7	247.9
2005 5/.....	19.0	15.3	12.0	12.8	16.1	26.8	48.0	95.4	247.8
2004 5/.....	17.1	13.6	11.2	12.2	14.3	24.4	46.4	94.2	243.7
2003 5/.....	15.1	12.3	10.8	10.5	15.1	24.4	45.1	90.8	242.0
2002.....	12.6	10.9	9.1	9.9	14.6	22.0	44.1	90.0	241.1
2001.....	9.8	8.7	8.4	10.3	14.0	22.3	43.0	87.2	234.0
2000.....	8.5	6.9	7.9	9.6	12.9	22.2	41.0	81.7	229.7
1999.....	8.0	6.5	7.8	8.0	13.9	21.3	43.4	84.3	236.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Non-Hispanic Black, both sexes

2007 5/.....	32.1	38.9	34.8	27.8	32.0	36.6	50.3	65.2	138.3
2006 5/.....	35.9	40.6	33.1	26.3	30.4	36.5	52.5	75.2	137.2
2005 5/.....	33.2	39.0	33.0	28.6	30.8	40.9	65.3	95.9	159.8
2004 5/.....	30.5	31.4	28.3	24.8	27.8	37.6	55.5	82.5	140.2
2003 5/.....	30.4	31.1	25.3	22.9	29.9	36.6	45.3	83.2	140.4
2002.....	30.1	29.1	25.5	23.4	30.2	35.3	55.5	73.9	156.7
2001.....	30.1	28.5	23.1	24.9	27.0	39.2	56.9	85.8	166.1
2000.....	28.1	28.0	24.3	23.2	28.3	38.9	56.5	79.6	161.2
1999.....	31.3	26.2	24.9	28.6	33.1	46.4	66.0	96.9	181.1

Non-Hispanic Black, male

2007 5/.....	46.7	57.6	55.9	43.8	53.4	52.3	70.2	77.0	161.4
2006 5/.....	53.4	65.3	53.8	43.4	45.0	52.6	77.3	98.3	164.9
2005 5/.....	50.7	61.1	53.4	52.1	45.3	61.1	91.8	141.0	193.2
2004 5/.....	43.1	50.5	47.0	39.1	41.4	56.2	79.3	119.0	189.0
2003 5/.....	45.7	46.6	40.5	37.7	46.6	50.9	67.7	114.7	184.3
2002.....	43.8	47.1	42.5	38.1	47.1	50.3	80.3	107.6	215.5
2001.....	48.6	47.1	37.7	38.5	40.9	56.1	81.5	135.7	220.6
2000.....	46.2	47.1	41.1	38.1	43.4	53.3	85.7	116.7	231.3
1999.....	52.3	44.0	43.0	41.8	51.5	71.3	95.7	129.3	249.6

Non-Hispanic Black, female

2007 5/.....	19.5	23.1	17.5	15.3	16.3	25.7	38.4	59.0	128.8
2006 5/.....	20.6	19.7	16.0	12.9	19.6	25.4	37.7	63.0	125.9
2005 5/.....	17.9	20.4	16.2	10.4	20.1	27.1	49.5	72.1	146.3
2004 5/.....	19.4	15.4	13.0	13.6	17.8	25.0	41.4	63.4	120.9
2003 5/.....	17.2	18.1	12.8	11.3	17.5	27.0	31.9	66.7	123.4
2002.....	18.2	14.0	11.6	11.9	17.6	25.2	40.6	56.4	134.2
2001.....	14.2	12.7	11.2	14.3	16.7	28.0	42.2	60.4	145.9
2000.....	12.5	11.8	10.5	11.4	17.1	29.3	38.9	60.9	135.2
1999.....	13.2	11.1	10.1	18.3	19.5	29.9	48.1	80.8	155.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

All origins 3/, both sexes

2007.....	2.7	3.3	4.6	6.3	9.8	17.8	34.7	70.4	166.7
2006.....	2.5	3.5	4.4	6.0	9.5	16.5	33.5	66.2	151.6
2005.....	2.4	3.2	3.9	5.9	8.8	16.7	31.9	63.6	147.7
2004.....	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
2003.....	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2

All origins 3/, male

2007.....	4.3	4.8	6.6	8.6	13.5	23.4	44.3	88.6	196.7
2006.....	4.0	5.5	6.4	8.5	13.3	20.6	42.4	83.7	181.1
2005.....	3.7	5.0	5.9	8.5	12.1	21.4	41.0	82.1	174.8
2004.....	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
2003.....	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5

All origins 3/, female

2007.....	1.1	1.8	2.6	4.1	6.6	13.3	27.7	58.8	152.4
2006.....	1.1	1.5	2.4	3.7	6.2	13.1	27.0	55.3	137.8
2005.....	1.2	1.4	2.1	3.6	6.0	13.0	25.4	52.2	135.2
2004.....	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
2003.....	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

Hispanic, both sexes

2007	2.7	3.1	3.8	6.5	6.6	14.2	25.2	49.6	92.6
2006	2.5	4.0	4.3	5.0	8.9	13.0	25.9	47.4	90.4
2005	2.4	3.0	3.4	6.6	9.5	14.8	27.1	46.8	91.6
2004	2.3	2.4	4.3	5.8	8.7	12.3	23.7	42.0	87.1
2003	2.3	3.0	3.4	7.3	6.2	13.2	19.9	34.4	80.8
2002	2.1	2.9	4.0	5.0	7.0	13.7	21.3	42.3	71.5
2001	2.7	2.4	3.8	4.8	5.5	13.5	19.0	31.8	70.0
2000	2.2	2.8	3.6	5.2	6.7	9.6	18.4	41.2	81.0
1999	2.0	2.4	3.3	4.5	8.1	8.1	17.8	35.8	86.3

Hispanic, male

2007	4.4	5.3	5.0	9.1	6.8	19.1	33.2	56.6	106.7
2006	4.4	7.0	7.1	8.6	14.1	16.4	30.4	59.0	102.6
2005	4.1	5.7	5.4	10.5	14.8	18.7	38.9	59.5	87.2
2004	4.2	4.1	7.0	9.8	13.0	15.3	29.4	52.8	106.5
2003	3.9	4.9	5.1	10.8	8.1	16.7	31.0	44.7	90.8
2002	3.4	4.5	6.6	7.4	11.2	17.1	25.6	57.2	97.0
2001	5.1	3.6	6.4	8.1	7.4	16.5	22.5	37.7	72.4
2000	3.9	5.1	5.9	7.5	10.4	15.6	25.1	41.2	104.8
1999	3.3	4.5	6.5	8.0	10.8	12.2	22.6	48.7	120.8

Hispanic, female

2007	*	*	2.7	4.2	6.4	10.3	19.5	45.0	84.9
2006	*	*	*	*	*	10.4	22.7	39.8	83.9
2005	*	*	*	*	5.2	11.7	18.6	38.3	93.8
2004	*	*	*	*	5.1	9.9	19.6	34.9	77.1
2003	*	*	*	4.3	*	10.5	12.1	27.7	75.8
2002	*	*	*	*	*	11.2	18.3	32.8	58.9
2001	*	*	*	*	*	11.2	16.5	28.1	68.9
2000	*	*	*	*	*	*	13.6	41.2	69.2
1999	*	*	*	*	*	*	14.3	28.1	69.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

Non-Hispanic 4/, both sexes

2007.....	2.7	3.3	4.6	6.2	10.1	18.1	35.2	71.5	170.4
2006.....	2.5	3.4	4.4	6.0	9.6	16.7	33.9	67.2	154.6
2005.....	2.4	3.2	3.9	5.9	8.7	16.9	32.2	64.4	150.2
2004.....	2.6	3.2	4.0	5.7	8.9	16.3	30.7	62.7	145.5
2003.....	2.3	2.8	3.7	5.1	8.2	14.7	29.1	58.4	137.9
2002.....	2.3	2.8	3.4	4.9	8.0	13.9	27.3	54.3	132.6
2001.....	2.4	2.8	3.2	4.8	7.9	12.3	25.5	50.4	123.1
2000.....	2.0	2.6	3.3	4.6	6.2	12.1	22.2	44.5	113.5
1999.....	2.3	2.2	3.0	4.5	6.8	11.4	22.3	45.6	110.7

Non-Hispanic 4/, male

2007.....	4.2	4.8	6.8	8.5	14.0	23.6	44.9	90.4	201.5
2006.....	3.9	5.3	6.3	8.4	13.2	20.9	43.1	85.0	185.4
2005.....	3.6	4.9	5.9	8.4	11.8	21.6	41.0	83.2	179.2
2004.....	4.0	4.9	6.1	8.0	12.3	22.0	38.9	80.0	177.2
2003.....	3.6	4.3	5.3	7.0	10.9	19.0	37.1	75.1	168.3
2002.....	3.6	4.2	5.1	7.0	10.6	18.9	35.5	68.0	161.3
2001.....	4.0	4.5	4.9	6.9	10.7	15.8	34.6	64.9	159.0
2000.....	3.3	4.0	4.9	6.8	8.4	15.7	29.5	57.9	146.5
1999.....	3.8	3.5	4.5	6.8	9.7	15.4	27.9	61.7	142.9

Non-Hispanic 4/, female

2007.....	1.1	1.9	2.6	4.1	6.7	13.5	28.1	59.6	155.7
2006.....	1.2	1.6	2.5	3.8	6.4	13.3	27.2	56.1	140.2
2005.....	1.3	1.5	2.1	3.6	6.1	13.0	25.7	52.8	136.9
2004.....	1.2	1.6	2.0	3.7	5.9	11.8	24.8	52.1	131.3
2003.....	1.1	1.3	2.1	3.4	5.9	11.3	23.3	48.3	124.6
2002.....	1.0	1.4	1.8	3.0	5.7	9.9	21.5	46.0	120.2
2001.....	0.9	1.1	1.6	2.9	5.5	9.4	19.2	41.7	108.1
2000.....	0.8	1.2	1.8	2.5	4.3	9.3	17.2	36.7	100.1
1999.....	0.8	1.0	1.6	2.4	4.4	8.3	18.5	36.2	97.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

Non-Hispanic White, both sexes

2007 5/.....	2.8	3.4	4.8	6.4	10.3	19.1	37.4	76.2	182.9
2006 5/.....	2.6	3.4	4.4	6.1	9.8	17.6	36.0	71.0	164.5
2005 5/.....	2.5	3.2	4.0	5.9	9.0	17.7	33.7	67.5	159.6
2004 5/.....	2.6	3.2	4.0	5.8	9.2	17.0	32.4	66.3	155.1
2003 5/.....	2.3	2.7	3.6	5.2	8.3	15.2	30.6	61.1	146.8
2002.....	2.3	2.8	3.3	5.0	8.1	14.3	28.5	57.5	140.4
2001.....	2.4	2.8	3.1	4.9	8.1	12.7	26.6	53.1	129.9
2000.....	2.1	2.5	3.2	4.5	6.1	12.5	23.2	46.7	120.2
1999.....	2.2	2.1	2.8	4.3	6.8	11.5	23.1	47.9	116.6

Non-Hispanic White, male

2007 5/.....	4.3	4.9	6.9	8.7	14.0	24.7	47.0	96.0	215.6
2006 5/.....	4.0	5.2	6.2	8.4	13.4	21.7	44.7	89.5	196.1
2005 5/.....	3.5	4.9	5.9	8.2	12.0	22.6	43.2	85.9	188.7
2004 5/.....	4.1	4.8	5.9	7.7	12.9	22.5	41.0	84.0	186.9
2003 5/.....	3.5	4.0	5.2	7.0	10.7	19.6	38.8	77.9	178.7
2002.....	3.6	4.1	4.9	7.1	10.6	19.3	36.7	71.3	169.2
2001.....	3.8	4.4	4.7	6.9	10.9	16.3	35.7	67.5	166.0
2000.....	3.4	3.7	4.6	6.8	8.2	16.2	30.4	60.3	154.1
1999.....	3.6	3.2	4.1	6.4	9.4	15.4	28.4	64.4	149.7

Non-Hispanic White, female

2007 5/.....	1.2	2.0	2.8	4.2	7.0	14.5	30.2	63.7	167.4
2006 5/.....	1.2	1.7	2.6	3.9	6.5	14.2	29.4	59.3	149.8
2005 5/.....	1.4	1.6	2.1	3.7	6.3	13.6	26.7	56.0	146.3
2004 5/.....	1.2	1.6	2.1	4.0	5.8	12.4	26.2	55.4	140.9
2003 5/.....	1.0	1.3	2.1	3.5	6.2	11.7	24.6	50.8	132.8
2002.....	1.0	1.5	1.8	3.2	5.9	10.2	22.6	49.1	128.0
2001.....	1.0	1.2	1.6	3.0	5.6	9.7	20.1	44.4	114.8
2000.....	0.8	1.2	1.8	2.5	4.3	9.5	18.2	38.6	106.5
1999.....	0.8	0.9	1.5	2.3	4.5	8.4	19.4	38.1	103.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

Non-Hispanic Black, both sexes

2007 5/.....	2.6	3.2	4.3	5.1	8.5	10.7	16.6	26.0	56.3
2006 5/.....	2.6	3.7	4.6	5.2	8.9	10.8	16.9	33.7	54.9
2005 5/.....	2.5	3.9	3.9	5.7	7.3	11.1	20.6	33.0	53.0
2004 5/.....	2.5	3.5	5.0	6.4	8.1	12.8	17.2	26.1	47.5
2003 5/.....	2.9	3.6	4.4	5.5	7.7	10.7	13.2	28.3	44.6
2002.....	2.6	3.2	3.9	4.8	6.1	10.8	14.9	18.4	49.3
2001.....	2.6	3.4	4.0	4.9	8.0	9.8	16.7	21.8	44.0
2000.....	2.0	3.6	4.6	5.1	7.1	9.7	11.8	20.7	39.1
1999.....	3.3	3.7	4.5	6.7	9.0	9.7	13.3	23.5	45.3

Non-Hispanic Black, male

2007 5/.....	4.6	5.0	7.0	8.2	14.4	15.9	22.8	32.6	65.4
2006 5/.....	4.4	6.5	7.7	8.4	12.8	16.8	26.2	47.1	68.7
2005 5/.....	4.5	6.3	6.3	9.7	11.0	14.6	24.3	53.2	63.8
2004 5/.....	3.8	6.0	8.9	12.1	10.0	20.6	24.2	38.9	73.2
2003 5/.....	4.7	5.8	6.7	7.9	12.9	12.1	17.8	35.5	56.5
2002.....	4.1	5.5	6.3	7.5	9.2	15.4	22.5	28.9	76.6
2001.....	4.9	6.1	6.4	7.5	12.0	13.4	24.6	36.8	68.1
2000.....	3.3	6.3	8.0	8.5	10.7	12.3	18.8	26.6	61.1
1999.....	5.7	6.1	7.5	10.2	15.1	14.9	19.8	31.9	58.2

Non-Hispanic Black, female

2007 5/.....	*	1.6	2.0	2.7	4.2	7.1	12.9	22.6	52.4
2006 5/.....	*	*	2.0	*	6.1	6.7	11.3	26.7	49.3
2005 5/.....	*	1.8	*	*	4.5	8.7	18.3	22.4	48.6
2004 5/.....	*	*	*	*	6.7	7.4	13.0	19.3	37.3
2003 5/.....	*	*	2.6	3.6	3.8	9.8	10.5	24.5	40.0
2002.....	*	*	*	*	3.8	7.8	10.4	13.0	38.9
2001.....	*	*	*	*	5.0	7.3	11.9	14.2	35.1
2000.....	*	*	*	*	4.5	7.9	7.6	17.7	31.0
1999.....	*	*	*	3.9	4.5	6.2	9.3	19.3	40.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental drowning and submersion (W65-W74)

All origins 3/, both sexes

2007.....	1.1	1.1	0.9	1.1	0.9	1.1	1.4	1.4	1.6
2006.....	1.2	1.0	1.0	0.9	1.1	1.2	1.6	1.4	1.4
2005.....	1.2	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
2004.....	1.1	1.0	0.9	0.9	1.1	1.0	1.0	1.5	1.6
2003.....	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6

All origins 3/, male

2007.....	1.7	1.8	1.4	1.9	1.5	1.9	2.1	2.4	3.0
2006.....	2.0	1.6	1.6	1.4	1.6	1.9	2.5	2.3	2.4
2005.....	2.0	1.7	1.5	1.5	1.4	1.8	2.1	2.5	3.9
2004.....	1.8	1.7	1.5	1.5	1.7	1.6	1.7	2.8	2.9
2003.....	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3

All origins 3/, female

2007.....	0.4	0.4	0.4	0.4	0.5	0.4	0.9	0.8	0.9
2006.....	0.5	0.4	0.5	0.4	0.7	0.6	0.9	0.8	0.9
2005.....	0.5	0.4	0.3	0.3	0.5	0.7	0.8	0.7	0.9
2004.....	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.0
2003.....	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental drowning and submersion (W65-W74)

Non-Hispanic 4/, both sexes

2007.....	1.1	1.1	0.9	1.2	0.9	1.1	1.4	1.4	1.6
2006.....	1.3	1.1	1.1	0.9	1.1	1.2	1.6	1.4	1.4
2005.....	1.3	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
2004.....	1.1	1.1	0.9	1.0	1.1	1.0	1.1	1.5	1.6
2003.....	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.0	0.9	0.8	0.8	0.9	1.0	1.1	1.5	1.7
2001.....	1.0	0.8	0.8	0.9	0.8	1.0	1.2	1.4	1.7
2000.....	0.9	1.0	0.9	0.9	0.9	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	1.2	1.2	1.5	1.7

Non-Hispanic 4/, male

2007.....	1.8	1.8	1.4	1.9	1.5	1.9	2.1	2.4	3.0
2006.....	2.1	1.7	1.7	1.4	1.6	1.9	2.6	2.3	2.4
2005.....	2.0	1.8	1.5	1.5	1.4	1.9	2.3	2.6	3.8
2004.....	1.8	1.7	1.5	1.5	1.7	1.5	1.8	2.8	2.7
2003.....	1.6	1.6	1.6	1.5	1.6	2.0	1.9	2.3	3.4
2002.....	1.6	1.5	1.3	1.2	1.4	1.7	1.7	2.4	3.6
2001.....	1.8	1.4	1.3	1.4	1.4	1.7	1.8	2.5	3.4
2000.....	1.6	1.7	1.6	1.6	1.5	1.7	1.3	2.7	2.8
1999.....	1.7	1.7	1.6	1.5	1.5	1.6	2.0	2.8	2.4

Non-Hispanic 4/, female

2007.....	0.4	0.5	0.4	0.4	0.5	*	0.9	0.8	0.9
2006.....	0.6	0.4	0.5	0.5	0.7	0.6	0.9	0.8	0.9
2005.....	0.5	0.5	0.3	0.3	0.5	0.7	0.9	0.7	0.9
2004.....	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.1
2003.....	0.4	0.4	0.4	0.4	0.5	0.6	0.9	*	1.1
2002.....	0.4	0.4	0.3	0.4	*	*	0.7	0.9	0.9
2001.....	0.3	0.3	0.4	0.5	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	*	*	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.4	0.4	0.4	0.8	0.6	0.7	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental drowning and submersion (W65-W74)

Non-Hispanic White, both sexes

2007 5/.....	1.2	1.2	0.9	1.2	0.9	1.1	1.4	1.5	1.6
2006 5/.....	1.4	1.0	1.1	0.9	1.2	1.1	1.6	1.5	1.4
2005 5/.....	1.3	1.1	0.9	0.9	0.9	1.2	1.4	1.3	1.8
2004 5/.....	1.1	1.0	0.9	0.9	1.1	0.9	1.0	1.4	1.5
2003 5/.....	1.0	1.0	1.0	0.9	1.1	1.2	1.2	1.3	1.7
2002.....	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	0.9	0.8	0.7	0.9	0.8	1.0	1.2	1.4	1.7
2000.....	0.9	0.9	0.9	0.9	0.9	1.0	0.9	1.8	1.4
1999.....	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.5	1.6

Non-Hispanic White, male

2007 5/.....	1.9	1.8	1.4	1.9	1.5	1.8	2.0	2.6	3.1
2006 5/.....	2.2	1.6	1.6	1.3	1.7	1.9	2.6	2.5	2.6
2005 5/.....	2.0	1.7	1.4	1.4	1.4	1.7	2.2	2.4	3.9
2004 5/.....	1.8	1.6	1.4	1.4	1.8	1.4	1.7	2.9	2.6
2003 5/.....	1.5	1.6	1.6	1.4	1.6	1.9	1.9	2.5	3.4
2002.....	1.4	1.4	1.3	1.2	1.4	1.5	1.7	2.4	3.7
2001.....	1.6	1.3	1.1	1.3	1.4	1.7	1.8	2.4	3.5
2000.....	1.5	1.5	1.6	1.7	1.4	1.6	1.1	3.0	2.7
1999.....	1.8	1.7	1.5	1.4	1.4	1.5	1.9	2.8	2.4

Non-Hispanic White, female

2007 5/.....	0.5	0.5	0.5	0.4	*	*	0.9	0.8	0.9
2006 5/.....	0.6	0.5	0.5	0.5	0.7	*	0.8	0.8	0.9
2005 5/.....	0.5	0.5	*	0.4	*	0.7	0.8	*	0.8
2004 5/.....	0.4	0.4	0.4	0.5	*	*	*	*	1.0
2003 5/.....	0.4	0.4	0.4	0.5	0.5	0.5	0.8	*	0.9
2002.....	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001.....	0.3	0.4	0.4	0.5	*	*	0.7	0.7	1.0
2000.....	0.3	0.3	*	*	*	0.6	0.7	1.1	0.9
1999.....	0.4	0.3	0.4	*	*	0.7	0.5	*	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes

2007.....	1.0	1.2	1.3	1.7	2.0	2.4	2.7	4.1	4.9
2006.....	1.1	1.1	1.4	1.5	2.1	2.2	3.2	3.8	4.3
2005.....	1.2	1.1	1.1	1.6	1.9	2.3	3.1	4.2	5.0
2004.....	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
2003.....	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9

All origins 3/, male

2007.....	1.3	1.6	1.9	2.4	2.7	3.0	3.2	5.1	6.5
2006.....	1.4	1.5	1.9	1.9	2.7	2.5	4.0	5.1	6.3
2005.....	1.5	1.6	1.5	2.0	2.5	2.8	3.9	5.6	7.0
2004.....	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
2003.....	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3

All origins 3/, female

2007.....	0.8	0.9	0.8	1.1	1.4	1.8	2.4	3.5	4.2
2006.....	0.8	0.8	1.0	1.1	1.5	2.0	2.6	2.9	3.4
2005.....	0.9	0.7	0.8	1.2	1.4	1.9	2.5	3.4	4.0
2004.....	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
2003.....	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 4/, both sexes

2007.....	1.1	1.3	1.4	1.8	2.1	2.4	2.7	4.2	5.0
2006.....	1.1	1.2	1.5	1.5	2.1	2.3	3.3	3.9	4.5
2005.....	1.3	1.2	1.2	1.6	2.0	2.3	3.1	4.3	5.1
2004.....	1.2	1.3	1.3	1.7	1.8	2.0	2.9	4.4	5.6
2003.....	1.3	1.3	1.3	1.6	2.1	2.7	3.0	4.4	5.4
2002.....	1.0	1.3	1.2	1.4	1.8	2.5	2.9	4.2	4.8
2001.....	1.1	1.2	1.6	1.5	2.1	2.6	3.1	3.9	5.7
2000.....	1.2	1.3	1.4	1.7	2.0	2.4	3.4	4.3	6.3
1999.....	1.0	1.0	1.2	1.5	2.1	2.6	3.5	4.7	6.0

Non-Hispanic 4/, male

2007.....	1.4	1.7	2.0	2.5	2.8	3.1	3.1	5.0	6.6
2006.....	1.4	1.5	2.0	2.0	2.7	2.6	4.1	5.3	6.5
2005.....	1.6	1.7	1.5	2.0	2.5	2.8	4.0	5.7	7.2
2004.....	1.6	1.8	1.7	2.2	2.6	2.3	3.5	6.0	7.3
2003.....	1.8	1.6	1.8	2.1	2.5	3.5	4.0	5.1	7.3
2002.....	1.5	1.6	1.4	2.0	2.6	3.3	3.6	6.2	7.0
2001.....	1.5	1.6	2.1	1.8	2.6	3.1	3.5	5.2	8.3
2000.....	1.6	1.9	2.0	2.3	2.5	3.2	4.4	5.9	9.3
1999.....	1.5	1.4	1.6	1.8	2.5	3.5	4.2	6.6	9.2

Non-Hispanic 4/, female

2007.....	0.9	0.9	0.9	1.2	1.5	1.9	2.4	3.7	4.2
2006.....	0.9	0.8	1.1	1.0	1.6	2.1	2.7	3.0	3.5
2005.....	0.9	0.7	0.8	1.3	1.5	1.9	2.4	3.4	4.1
2004.....	0.9	0.7	1.0	1.3	1.1	1.7	2.5	3.4	4.9
2003.....	0.9	1.1	0.9	1.2	1.7	2.0	2.3	4.0	4.5
2002.....	0.6	1.0	0.9	0.9	1.1	1.8	2.4	2.9	3.8
2001.....	0.6	0.7	1.1	1.2	1.6	2.2	2.8	3.1	4.6
2000.....	0.7	0.8	0.9	1.2	1.7	1.7	2.8	3.4	5.0
1999.....	0.6	0.7	0.9	1.2	1.7	1.9	3.1	3.6	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic White, both sexes

2007 5/.....	1.0	1.2	1.3	1.7	1.9	2.1	2.3	3.8	4.1
2006 5/.....	1.1	1.1	1.4	1.4	2.0	2.2	2.9	3.4	3.9
2005 5/.....	1.3	1.1	0.9	1.4	1.7	2.2	2.6	3.8	4.3
2004 5/.....	1.0	1.2	1.2	1.5	1.6	1.6	2.4	3.8	4.9
2003 5/.....	1.2	1.2	1.2	1.5	1.8	2.3	2.6	3.8	4.7
2002.....	1.0	1.1	1.0	1.1	1.6	2.1	2.5	3.7	3.9
2001.....	0.9	1.0	1.3	1.3	1.7	2.2	2.6	3.3	4.7
2000.....	1.0	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	1.1	1.2	1.7	2.1	2.8	3.9	4.7

Non-Hispanic White, male

2007 5/.....	1.2	1.6	1.7	2.3	2.4	2.7	2.7	4.5	5.3
2006 5/.....	1.3	1.5	1.8	1.8	2.4	2.5	3.4	4.9	6.2
2005 5/.....	1.6	1.5	1.1	1.7	2.1	2.6	3.4	5.1	6.1
2004 5/.....	1.3	1.6	1.4	2.0	2.2	1.9	2.9	5.4	6.5
2003 5/.....	1.6	1.4	1.6	1.9	2.2	3.1	3.2	4.2	6.5
2002.....	1.4	1.4	1.3	1.7	2.2	2.7	3.1	5.8	5.7
2001.....	1.3	1.3	1.8	1.5	2.2	2.5	2.8	4.5	6.9
2000.....	1.4	1.7	1.6	1.8	2.0	2.6	3.6	4.9	8.1
1999.....	1.3	1.2	1.4	1.6	2.0	2.7	3.2	5.7	7.8

Non-Hispanic White, female

2007 5/.....	0.8	0.9	0.8	1.2	1.5	1.7	2.0	3.4	3.5
2006 5/.....	0.9	0.8	1.0	1.0	1.6	1.9	2.4	2.5	2.8
2005 5/.....	0.9	0.7	0.7	1.2	1.3	1.8	2.0	3.0	3.5
2004 5/.....	0.8	0.7	0.9	1.1	1.0	1.4	2.1	2.8	4.2
2003 5/.....	0.8	0.9	0.8	1.0	1.5	1.7	2.2	3.5	3.9
2002.....	0.6	0.9	0.8	0.7	1.0	1.6	2.0	2.4	3.2
2001.....	0.6	0.7	0.9	1.1	1.4	1.9	2.5	2.6	3.8
2000.....	0.6	0.7	0.7	1.0	1.5	1.4	2.3	2.6	3.9
1999.....	0.5	0.6	0.9	0.8	1.4	1.5	2.6	2.8	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic Black, both sexes

2007 5/.....	1.9	2.1	2.8	2.8	4.5	5.7	7.4	8.8	15.6
2006 5/.....	2.0	1.9	2.8	2.7	4.3	4.0	9.1	10.2	12.5
2005 5/.....	1.8	2.2	3.6	3.2	4.4	4.4	8.1	11.4	14.5
2004 5/.....	2.7	2.2	2.9	4.3	4.1	6.0	8.0	12.2	15.3
2003 5/.....	2.4	2.9	2.9	3.0	4.5	6.5	8.7	12.7	14.9
2002.....	1.6	2.8	2.6	4.1	3.9	6.5	7.4	10.7	16.0
2001.....	2.3	2.6	3.8	3.3	5.1	7.3	8.7	11.9	17.8
2000.....	2.2	2.5	3.9	4.6	5.1	7.2	11.1	16.1	20.7
1999.....	2.1	2.3	2.7	4.4	5.9	8.4	12.2	15.6	22.0

Non-Hispanic Black, male

2007 5/.....	2.6	2.8	4.5	4.4	8.3	7.8	9.1	*	23.6
2006 5/.....	2.6	2.7	3.9	4.2	6.8	*	14.7	*	*
2005 5/.....	2.4	3.9	5.2	5.2	6.9	*	11.9	15.8	21.9
2004 5/.....	4.0	3.6	4.4	4.8	7.5	8.0	12.1	15.4	*
2003 5/.....	3.5	3.2	4.8	4.0	6.6	9.1	15.4	17.4	*
2002.....	2.2	4.2	3.4	6.1	6.4	9.6	9.4	*	26.3
2001.....	3.6	4.4	5.2	5.0	7.3	11.4	12.3	*	26.6
2000.....	3.2	4.0	5.6	7.2	7.7	10.6	15.9	21.5	28.2
1999.....	3.1	3.4	4.2	4.4	8.0	13.5	17.4	20.4	29.7

Non-Hispanic Black, female

2007 5/.....	*	*	*	*	*	4.3	6.3	7.3	12.3
2006 5/.....	1.4	*	*	*	*	*	5.8	8.9	12.0
2005 5/.....	*	*	2.3	*	*	*	5.7	9.1	11.5
2004 5/.....	1.5	*	*	3.9	*	4.7	5.5	10.4	13.7
2003 5/.....	1.5	2.6	*	*	*	4.8	*	10.3	13.2
2002.....	*	*	*	*	*	*	6.2	8.9	12.1
2001.....	*	*	2.8	*	*	4.6	6.5	10.4	14.5
2000.....	*	*	*	*	*	5.0	8.1	13.4	17.9
1999.....	*	*	*	4.4	4.3	5.1	9.0	13.2	19.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes

2007.....	21.6	19.4	12.1	6.4	3.5	2.7	2.6	2.9	3.5
2006.....	20.7	17.1	9.8	4.6	2.8	2.2	2.4	2.9	3.8
2005.....	18.2	14.4	8.3	4.4	2.6	2.0	2.3	2.6	3.5
2004.....	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
2003.....	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4

All origins 3/, male

2007.....	26.7	24.5	15.6	7.6	4.4	3.4	2.4	3.5	3.7
2006.....	26.9	22.5	12.7	6.1	3.0	2.7	3.0	2.7	4.6
2005.....	24.0	19.1	10.9	5.7	3.0	2.3	3.2	2.8	4.6
2004.....	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
2003.....	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4

All origins 3/, female

2007.....	16.6	14.5	8.7	5.3	2.6	2.1	2.7	2.5	3.4
2006.....	14.7	11.9	7.1	3.4	2.6	1.9	1.9	3.0	3.4
2005.....	12.6	10.0	5.8	3.2	2.3	1.7	1.7	2.4	3.0
2004.....	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
2003.....	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

Hispanic, both sexes

2007.....	13.5	12.8	7.8	4.6	2.6	*	*	*	*
2006.....	13.4	11.0	8.7	4.7	*	*	*	*	*
2005.....	11.6	10.1	6.3	4.3	2.7	*	*	*	*
2004.....	10.5	8.7	6.8	3.4	*	*	*	*	*
2003.....	11.9	8.6	5.2	3.2	*	*	*	*	*
2002.....	11.7	7.9	4.7	3.1	*	*	*	*	*
2001.....	8.6	6.6	4.7	*	*	*	*	*	*
2000.....	8.9	6.8	4.0	*	*	*	*	*	*
1999.....	11.4	7.6	4.1	2.9	*	*	*	*	*

Hispanic, male

2007.....	21.2	19.8	10.9	7.0	*	*	*	*	*
2006.....	19.1	16.8	13.9	8.4	*	*	*	*	*
2005.....	17.4	15.9	10.7	7.2	*	*	*	*	*
2004.....	15.3	12.8	10.4	5.6	*	*	*	*	*
2003.....	18.8	13.2	8.6	5.7	*	*	*	*	*
2002.....	17.6	13.4	7.2	5.4	*	*	*	*	*
2001.....	14.0	11.5	8.0	*	*	*	*	*	*
2000.....	15.7	10.8	5.3	*	*	*	*	*	*
1999.....	19.7	12.0	7.6	*	*	*	*	*	*

Hispanic, female

2007.....	5.5	5.8	5.0	*	*	*	*	*	*
2006.....	7.4	5.2	3.8	*	*	*	*	*	*
2005.....	5.6	4.3	*	*	*	*	*	*	*
2004.....	5.5	4.7	3.4	*	*	*	*	*	*
2003.....	4.9	4.2	*	*	*	*	*	*	*
2002.....	5.7	2.6	*	*	*	*	*	*	*
2001.....	3.1	*	*	*	*	*	*	*	*
2000.....	*	3.0	*	*	*	*	*	*	*
1999.....	3.1	3.5	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 4/, both sexes

2007.....	22.5	20.0	12.3	6.5	3.5	2.6	2.7	3.0	3.6
2006.....	21.6	17.7	9.9	4.6	2.9	2.3	2.4	2.9	3.9
2005.....	18.9	14.7	8.4	4.4	2.6	2.0	2.4	2.6	3.5
2004.....	17.4	12.2	6.7	3.7	2.4	2.1	2.3	2.6	3.4
2003.....	15.7	11.0	6.1	3.4	2.4	2.0	2.0	2.6	3.5
2002.....	14.0	9.3	5.0	2.9	2.2	1.9	2.0	2.7	4.1
2001.....	11.0	6.9	3.6	2.4	1.7	1.5	2.1	2.4	3.6
2000.....	9.9	5.9	3.2	2.2	1.5	1.5	1.6	2.6	4.4
1999.....	9.5	5.1	3.3	1.9	1.6	1.6	1.9	2.6	3.5

Non-Hispanic 4/, male

2007.....	27.3	24.9	15.9	7.6	4.4	3.3	2.4	3.7	3.7
2006.....	27.8	23.0	12.6	5.7	3.1	2.7	3.0	2.7	4.7
2005.....	24.7	19.2	10.8	5.5	2.9	2.3	3.3	2.9	4.7
2004.....	22.2	16.5	8.2	4.1	2.7	2.4	2.5	3.0	4.2
2003.....	20.9	14.6	7.4	4.3	2.9	2.2	2.3	3.6	4.0
2002.....	18.7	12.7	6.6	3.4	2.7	2.0	2.2	3.2	5.3
2001.....	15.7	9.6	4.6	2.9	1.9	2.0	2.7	2.9	3.9
2000.....	14.4	8.5	3.8	2.7	1.7	1.7	1.7	2.9	6.2
1999.....	14.4	7.3	4.4	2.7	2.1	1.9	2.1	3.0	4.4

Non-Hispanic 4/, female

2007.....	17.9	15.4	9.0	5.4	2.7	2.1	2.8	2.5	3.5
2006.....	15.6	12.6	7.4	3.5	2.8	2.0	2.0	3.0	3.5
2005.....	13.3	10.5	6.1	3.3	2.4	1.8	1.8	2.4	3.0
2004.....	12.6	8.2	5.3	3.3	2.1	1.8	2.2	2.3	3.1
2003.....	10.6	7.6	4.8	2.6	1.9	1.9	1.7	2.0	3.3
2002.....	9.4	6.1	3.5	2.5	1.8	1.7	2.0	2.3	3.5
2001.....	6.5	4.3	2.6	1.9	1.5	1.0	1.7	2.2	3.4
2000.....	5.5	3.3	2.6	1.8	1.4	1.3	1.6	2.4	3.6
1999.....	4.8	3.0	2.3	1.3	1.2	1.4	1.8	2.4	3.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic White, both sexes

2007 5/.....	23.9	20.1	11.8	6.3	3.3	2.6	2.7	3.0	3.7
2006 5/.....	22.1	17.1	9.2	4.1	2.6	2.2	2.4	2.9	3.8
2005 5/.....	19.6	14.1	7.7	4.1	2.3	1.9	2.4	2.5	3.6
2004 5/.....	17.8	12.0	6.1	3.5	2.3	2.0	2.3	2.5	3.5
2003 5/.....	15.9	10.6	5.9	3.4	2.0	2.0	2.0	2.8	3.5
2002.....	13.7	8.7	4.5	2.7	2.2	1.9	2.0	2.7	4.1
2001.....	10.5	6.0	3.2	2.1	1.6	1.4	2.0	2.3	3.4
2000.....	9.2	5.0	2.9	2.1	1.5	1.4	1.7	2.7	4.1
1999.....	8.7	4.4	3.0	1.6	1.5	1.5	2.0	2.7	3.3

Non-Hispanic White, male

2007 5/.....	28.5	24.3	14.4	7.0	3.8	3.2	2.3	3.8	3.9
2006 5/.....	27.9	21.4	10.9	4.6	2.5	2.3	2.9	2.8	4.8
2005 5/.....	25.0	17.8	9.3	4.6	2.2	1.9	3.2	2.6	4.9
2004 5/.....	22.7	15.5	7.0	3.4	2.3	2.3	2.3	2.8	4.3
2003 5/.....	20.9	13.7	6.8	4.1	2.1	2.1	2.3	3.7	3.6
2002.....	18.2	11.4	5.5	2.7	2.5	1.9	2.2	2.9	5.4
2001.....	14.7	8.0	3.8	2.3	1.8	1.9	2.4	2.7	3.8
2000.....	13.2	7.0	3.0	2.3	1.4	1.6	1.7	3.0	5.8
1999.....	12.9	5.9	3.6	2.0	1.9	1.7	2.2	3.1	4.6

Non-Hispanic White, female

2007 5/.....	19.4	16.0	9.2	5.6	2.9	2.1	2.9	2.6	3.6
2006 5/.....	16.4	12.9	7.7	3.6	2.7	2.1	2.0	3.0	3.3
2005 5/.....	14.2	10.4	6.2	3.5	2.4	1.8	1.8	2.4	2.9
2004 5/.....	13.0	8.5	5.3	3.5	2.3	1.8	2.3	2.3	3.1
2003 5/.....	11.0	7.6	5.0	2.7	1.8	1.9	1.8	2.2	3.5
2002.....	9.4	6.1	3.5	2.7	1.9	1.8	1.9	2.5	3.5
2001.....	6.3	4.0	2.7	1.9	1.5	0.9	1.8	2.0	3.2
2000.....	5.3	3.0	2.7	1.9	1.5	1.3	1.7	2.5	3.4
1999.....	4.5	2.9	2.4	1.2	1.2	1.2	1.8	2.4	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic Black, both sexes

2007 5/.....	21.8	26.5	20.0	10.5	6.3	3.9	3.6	*	*
2006 5/.....	24.9	27.5	18.1	9.4	5.5	3.6	*	*	5.6
2005 5/.....	21.1	23.6	16.6	8.2	5.0	3.8	*	*	*
2004 5/.....	20.0	18.3	12.8	6.2	4.2	3.2	*	*	*
2003 5/.....	19.2	17.1	9.7	4.9	5.5	*	*	*	*
2002.....	19.8	16.5	10.6	6.0	2.8	*	*	*	*
2001.....	18.1	15.6	7.3	5.2	2.5	*	*	*	6.6
2000.....	16.8	14.3	6.9	3.8	2.6	*	*	*	8.3
1999.....	18.2	12.2	6.5	5.4	2.8	3.3	*	*	*

Non-Hispanic Black, male

2007 5/.....	29.5	38.0	32.2	16.7	11.3	*	*	*	*
2006 5/.....	35.6	42.7	30.1	17.2	8.7	6.6	*	*	*
2005 5/.....	30.5	35.3	27.3	15.2	9.3	6.8	*	*	*
2004 5/.....	26.1	29.3	20.8	10.4	7.5	*	*	*	*
2003 5/.....	27.3	25.8	15.8	8.9	10.4	*	*	*	*
2002.....	27.9	26.2	18.3	11.2	5.1	*	*	*	*
2001.....	28.4	25.2	12.6	9.0	*	*	*	*	*
2000.....	27.0	23.2	11.7	6.8	*	*	*	*	*
1999.....	30.0	20.7	12.2	9.6	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	15.1	16.8	10.1	5.7	*	*	*	*	*
2006 5/.....	15.7	14.7	8.2	3.4	*	*	*	*	*
2005 5/.....	12.9	13.8	7.7	*	*	*	*	*	*
2004 5/.....	14.6	9.0	6.2	*	*	*	*	*	*
2003 5/.....	12.3	9.8	4.6	*	*	*	*	*	*
2002.....	12.8	8.3	4.4	*	*	*	*	*	*
2001.....	9.2	7.5	3.0	*	*	*	*	*	*
2000.....	8.1	6.7	3.0	*	*	*	*	*	*
1999.....	8.0	4.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes

2007.....	3.7	4.3	4.6	5.5	7.2	10.9	17.9	33.1	85.8
2006.....	4.0	4.5	4.7	5.8	7.7	10.7	18.8	35.2	89.3
2005.....	4.4	4.9	5.0	6.8	8.4	12.3	20.7	38.9	95.1
2004.....	3.9	4.5	4.5	5.7	7.0	11.0	19.2	37.4	95.5
2003.....	3.8	4.0	4.7	5.5	8.1	11.2	19.8	38.1	102.5
2002.....	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001.....	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000.....	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
1999.....	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9

All origins 3/, male

2007.....	5.9	7.1	7.0	8.1	10.0	15.3	22.3	42.3	97.1
2006.....	6.5	7.1	7.1	8.6	11.2	14.5	24.5	44.8	99.4
2005.....	7.1	7.8	7.4	10.2	11.5	16.6	26.8	49.0	107.1
2004.....	6.3	7.0	6.7	8.4	9.8	14.3	25.1	46.1	109.8
2003.....	6.0	6.1	7.1	8.7	11.4	15.1	26.5	47.6	120.2
2002.....	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001.....	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000.....	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
1999.....	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7

All origins 3/, female

2007.....	1.6	1.7	2.3	3.1	4.8	7.3	14.6	27.3	80.4
2006.....	1.6	2.0	2.5	3.2	4.7	7.7	14.6	29.2	84.5
2005.....	1.8	2.1	2.7	3.6	5.7	8.7	16.3	32.6	89.6
2004.....	1.5	2.1	2.3	3.2	4.6	8.4	15.0	32.1	89.1
2003.....	1.6	2.1	2.5	2.7	5.3	8.2	15.0	32.3	94.7
2002.....	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001.....	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000.....	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
1999.....	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Hispanic, both sexes

2007	2.7	3.5	3.6	4.2	6.7	6.8	12.5	19.2	35.0
2006	3.1	3.2	4.3	4.5	5.4	7.7	10.2	21.6	37.2
2005	3.5	4.2	4.1	4.1	4.8	9.4	11.7	20.3	38.1
2004	3.6	2.9	2.7	4.6	3.5	7.4	10.0	21.8	44.2
2003	2.9	3.1	3.2	4.9	6.7	5.7	12.9	23.4	37.3
2002	2.4	3.3	3.0	3.9	6.4	5.9	14.9	16.0	48.2
2001	3.2	4.1	3.3	4.4	5.9	10.1	12.7	21.9	63.8
2000	2.8	2.6	3.2	4.9	5.3	7.3	13.5	22.3	70.3
1999	2.9	2.6	3.0	5.8	7.1	10.7	11.3	21.7	65.9

Hispanic, male

2007	4.0	6.4	6.0	7.0	11.3	10.3	15.9	26.3	38.6
2006	5.3	5.5	6.6	6.5	8.7	11.1	12.4	27.8	45.1
2005	6.3	6.4	6.0	7.0	6.5	13.9	16.2	21.0	46.1
2004	6.2	4.9	4.0	6.7	*	8.7	13.2	24.5	43.4
2003	4.9	4.4	5.3	7.9	9.1	9.0	21.3	31.2	44.6
2002	4.3	5.4	5.3	5.9	9.1	*	21.1	*	70.3
2001	5.3	7.1	5.4	6.0	8.5	13.7	17.7	26.0	68.6
2000	4.8	4.1	5.3	8.3	7.8	9.7	16.2	*	88.7
1999	5.3	4.2	5.3	7.4	10.4	16.8	*	*	84.1

Hispanic, female

2007	*	*	*	*	*	*	10.1	14.5	33.1
2006	*	*	*	*	*	*	8.6	17.4	33.0
2005	*	*	*	*	*	*	8.5	19.7	33.9
2004	*	*	*	*	*	6.4	*	19.9	44.7
2003	*	*	*	*	*	*	*	18.3	33.6
2002	*	*	*	*	*	*	10.5	*	37.3
2001	*	*	*	*	*	7.3	*	19.3	61.4
2000	*	*	*	*	*	*	11.5	19.7	61.3
1999	*	*	*	*	*	*	11.6	19.7	57.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 4/, both sexes

2007.....	3.8	4.4	4.6	5.5	7.3	11.2	18.2	33.9	88.5
2006.....	4.1	4.6	4.7	5.9	7.9	10.9	19.3	35.9	91.8
2005.....	4.5	4.9	5.0	7.0	8.7	12.4	21.3	39.9	97.7
2004.....	3.9	4.6	4.6	5.8	7.2	11.2	19.7	38.1	97.7
2003.....	3.8	4.1	4.9	5.6	8.2	11.6	20.1	38.7	105.1
2002.....	3.6	4.0	4.9	5.8	8.8	11.4	21.9	39.8	107.6
2001.....	4.0	4.5	4.6	6.4	7.9	12.4	22.9	42.9	116.6
2000.....	4.0	4.2	5.1	6.4	7.9	12.4	22.3	42.0	118.5
1999.....	3.9	4.5	5.1	6.3	9.0	13.4	24.1	47.7	130.7

Non-Hispanic 4/, male

2007.....	6.1	7.1	7.0	8.1	9.9	15.7	22.8	43.3	100.5
2006.....	6.6	7.2	7.1	8.8	11.4	14.7	25.3	45.5	102.2
2005.....	7.2	7.9	7.5	10.4	11.8	16.8	27.4	50.6	110.2
2004.....	6.2	7.1	7.0	8.5	10.0	14.6	25.7	47.2	113.1
2003.....	6.0	6.2	7.2	8.7	11.6	15.5	26.6	48.2	123.5
2002.....	5.9	6.2	7.4	8.9	12.2	15.4	29.0	48.2	120.4
2001.....	6.4	6.8	6.7	9.1	11.0	15.9	29.8	53.6	136.5
2000.....	6.6	6.8	7.9	9.1	10.9	15.8	28.5	52.4	136.9
1999.....	6.1	7.3	7.8	9.1	12.0	18.0	30.7	60.3	154.8

Non-Hispanic 4/, female

2007.....	1.6	1.8	2.4	3.2	5.0	7.6	14.8	28.0	82.8
2006.....	1.7	2.1	2.6	3.2	4.9	7.9	15.0	29.9	87.0
2005.....	2.0	2.1	2.7	3.8	5.9	8.9	16.8	33.2	92.0
2004.....	1.6	2.2	2.4	3.2	4.8	8.5	15.4	32.6	90.9
2003.....	1.7	2.1	2.6	2.7	5.3	8.5	15.5	32.9	97.0
2002.....	1.3	1.9	2.6	3.1	5.8	8.3	16.9	34.7	102.2
2001.....	1.6	2.3	2.6	3.9	5.2	9.6	18.1	36.5	108.4
2000.....	1.4	1.7	2.4	3.9	5.4	9.8	18.0	35.8	111.1
1999.....	1.8	1.8	2.7	3.7	6.4	9.8	19.5	40.3	121.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic White, both sexes

2007 5/.....	3.8	4.3	4.5	5.3	7.0	11.1	18.0	35.2	92.5
2006 5/.....	4.1	4.5	4.7	5.8	7.8	10.6	19.5	37.2	96.0
2005 5/.....	4.3	4.6	4.7	6.7	8.4	11.9	20.8	39.8	100.2
2004 5/.....	4.0	4.7	4.5	5.7	7.0	11.0	19.6	38.7	101.3
2003 5/.....	3.8	3.9	4.7	5.4	8.0	11.4	20.3	39.1	109.1
2002.....	3.6	4.0	4.7	5.7	8.0	11.3	21.6	40.1	110.9
2001.....	3.8	4.4	4.5	6.0	7.8	12.0	22.7	43.2	119.5
2000.....	3.8	4.1	4.9	6.2	7.5	12.0	21.9	42.5	122.0
1999.....	3.6	4.3	4.7	5.7	8.6	12.7	23.3	47.6	134.3

Non-Hispanic White, male

2007 5/.....	6.0	6.8	6.9	8.0	9.3	15.6	22.3	45.3	105.2
2006 5/.....	6.5	6.8	6.9	8.7	11.2	14.3	25.5	47.1	106.7
2005 5/.....	6.8	7.4	7.0	9.6	11.7	15.6	26.5	50.2	113.3
2004 5/.....	6.4	7.1	6.7	8.5	9.7	14.4	25.4	47.7	116.4
2003 5/.....	5.8	6.0	7.0	8.2	11.4	15.0	26.7	48.1	127.5
2002.....	6.1	6.0	7.0	8.7	10.9	15.2	28.4	48.2	123.1
2001.....	6.2	6.5	6.4	8.7	10.8	15.3	29.6	52.7	139.7
2000.....	6.1	6.5	7.4	8.6	10.2	15.3	27.5	52.0	139.7
1999.....	5.5	7.0	7.1	8.5	11.2	16.9	29.4	60.1	157.3

Non-Hispanic White, female

2007 5/.....	1.5	1.8	2.2	2.9	4.8	7.4	14.8	28.8	86.5
2006 5/.....	1.7	2.1	2.5	3.1	4.7	7.6	14.9	31.0	91.0
2005 5/.....	1.9	1.9	2.6	3.9	5.4	8.8	16.6	33.3	94.2
2004 5/.....	1.7	2.3	2.4	3.1	4.6	8.3	15.2	33.1	94.5
2003 5/.....	1.7	2.0	2.5	2.7	5.0	8.5	15.8	33.5	101.0
2002.....	1.2	2.0	2.6	2.9	5.3	8.0	16.8	35.1	105.7
2001.....	1.5	2.4	2.7	3.6	5.1	9.2	17.9	37.4	111.2
2000.....	1.5	1.7	2.4	4.0	5.2	9.4	18.0	36.9	114.9
1999.....	1.7	1.7	2.5	3.2	6.3	9.4	19.0	40.3	125.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic Black, both sexes

2007 5/.....	4.8	6.0	6.6	8.2	11.5	14.9	21.7	27.5	61.6
2006 5/.....	5.1	6.1	6.3	7.7	10.3	16.2	22.4	28.4	62.6
2005 5/.....	6.5	7.8	7.8	10.3	12.8	19.7	32.2	47.5	87.4
2004 5/.....	4.1	5.7	6.4	6.9	10.4	13.7	26.5	38.2	71.4
2003 5/.....	4.8	6.4	6.9	8.3	11.5	15.1	20.0	39.4	74.1
2002.....	4.4	5.2	7.3	7.4	16.3	14.7	28.4	39.8	84.4
2001.....	5.6	6.0	6.1	10.5	10.5	18.9	27.5	44.7	96.5
2000.....	5.8	5.8	8.1	8.8	12.6	18.1	30.8	39.4	90.9
1999.....	6.7	6.6	9.5	11.0	14.1	23.2	36.3	54.0	106.7

Non-Hispanic Black, male

2007 5/.....	7.9	9.8	10.0	12.3	17.5	19.9	31.9	29.8	68.1
2006 5/.....	8.5	10.8	9.6	11.3	14.7	22.8	29.5	35.8	76.0
2005 5/.....	11.0	12.5	12.5	19.1	15.9	29.6	48.2	64.0	99.0
2004 5/.....	6.8	9.0	10.8	10.0	15.0	19.0	35.9	55.1	89.4
2003 5/.....	8.0	9.5	10.7	14.3	15.7	22.9	29.9	55.8	92.7
2002.....	6.6	8.8	12.6	11.4	24.3	19.2	41.8	54.6	107.2
2001.....	8.9	9.7	10.1	14.8	15.9	25.1	35.6	73.6	115.5
2000.....	10.4	10.3	14.2	14.2	18.7	23.6	47.7	64.4	124.5
1999.....	11.3	11.0	15.9	16.1	21.3	35.3	51.2	70.0	155.7

Non-Hispanic Black, female

2007 5/.....	2.0	2.7	3.9	5.1	7.1	11.4	15.6	26.2	59.0
2006 5/.....	2.1	2.1	3.5	4.8	6.9	11.7	18.1	24.4	57.2
2005 5/.....	2.7	3.9	4.0	3.4	10.5	12.9	22.7	38.7	82.7
2004 5/.....	1.7	2.8	2.8	4.5	7.0	10.1	21.0	29.4	64.3
2003 5/.....	2.0	3.7	3.8	3.6	8.4	9.8	14.1	30.8	66.9
2002.....	2.6	2.1	3.0	4.3	10.4	11.7	20.5	32.1	75.7
2001.....	2.7	2.8	2.8	7.1	6.5	14.7	22.6	30.0	89.4
2000.....	1.8	*	3.0	4.5	8.1	14.5	20.6	26.8	78.5
1999.....	2.8	2.9	4.3	7.0	8.9	15.2	27.4	46.1	88.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes

2007.....	17.7	17.7	16.2	14.6	12.9	12.3	16.4	16.1	15.6
2006.....	17.1	17.3	15.3	13.4	12.6	12.6	15.6	16.3	15.9
2005.....	16.7	16.1	14.3	13.3	11.8	13.5	15.4	18.7	16.9
2004.....	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
2003.....	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 6/.....	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All origins 3/, male

2007.....	26.9	27.0	25.0	23.3	22.4	22.7	33.2	35.9	41.8
2006.....	26.0	26.4	23.9	21.1	22.0	23.6	31.6	35.7	43.2
2005.....	25.4	25.0	22.4	22.0	20.6	25.4	31.9	41.4	45.0
2004.....	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
2003.....	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 6/.....	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All origins 3/, female

2007.....	8.7	8.8	8.0	6.5	4.5	3.9	4.1	3.6	3.1
2006.....	8.4	8.4	7.2	6.3	4.4	3.7	3.8	4.2	3.1
2005.....	8.2	7.7	6.6	5.5	4.2	3.8	3.5	4.7	4.0
2004.....	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
2003.....	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 6/.....	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Hispanic, both sexes

2007.....	7.6	9.0	7.8	6.1	6.7	7.2	8.8	8.1	7.9
2006.....	7.2	5.7	5.4	6.9	4.8	5.7	6.7	8.8	*
2005.....	7.2	5.8	6.1	5.8	4.7	7.5	9.7	7.3	9.0
2004.....	7.7	7.3	7.5	6.3	5.6	8.5	6.0	13.9	*
2003.....	7.4	6.9	7.2	6.5	6.4	6.7	7.3	9.0	10.9
2002.....	7.2	7.0	7.9	6.2	6.8	8.4	8.0	11.1	12.7
2001 6/.....	6.8	7.4	6.2	6.2	6.7	8.7	6.8	12.4	13.8
2000.....	7.9	7.6	6.5	5.3	7.2	8.6	9.8	15.0	*
1999.....	7.2	6.9	7.1	6.5	7.9	5.9	11.3	16.4	15.4

Hispanic, male

2007.....	12.0	14.6	13.5	11.1	13.2	14.3	18.8	20.1	*
2006.....	11.1	8.9	9.5	12.1	8.2	10.3	14.4	18.8	*
2005.....	11.9	9.8	10.3	10.1	7.4	15.1	20.5	*	*
2004.....	12.0	11.9	11.6	10.9	11.4	16.6	13.2	30.2	*
2003.....	11.8	11.1	13.4	12.6	12.3	14.1	15.8	21.8	*
2002.....	11.0	11.7	14.5	11.3	13.2	16.7	16.0	25.2	35.1
2001 6/.....	11.0	11.9	11.2	11.9	13.8	17.4	15.7	28.6	41.9
2000.....	12.6	12.3	11.8	10.1	14.5	17.0	19.9	38.3	*
1999.....	12.4	10.4	12.7	11.6	15.8	11.2	23.4	37.7	*

Hispanic, female

2007.....	3.0	3.4	2.5	*	*	*	*	*	*
2006.....	3.0	2.4	*	*	*	*	*	*	*
2005.....	2.4	*	*	*	*	*	*	*	*
2004.....	3.3	2.8	3.7	*	*	*	*	*	*
2003.....	2.9	2.9	*	*	*	*	*	*	*
2002.....	3.3	2.5	*	*	*	*	*	*	*
2001 6/.....	2.5	3.0	*	*	*	*	*	*	*
2000.....	3.3	3.0	*	*	*	*	*	*	*
1999.....	*	3.6	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007.....	19.0	18.7	17.0	15.2	13.3	12.7	16.9	16.5	15.9
2006.....	18.3	18.4	16.2	13.9	13.1	13.1	16.1	16.7	16.3
2005.....	17.8	17.1	14.9	13.9	12.4	13.9	15.7	19.3	17.2
2004.....	17.9	17.0	14.7	13.7	11.8	13.8	16.6	16.9	16.8
2003.....	16.9	16.5	15.2	13.1	12.3	14.0	16.0	18.1	17.1
2002.....	17.2	15.7	15.1	12.7	12.8	14.9	17.0	19.7	18.1
2001 6/.....	16.5	15.1	14.5	12.3	13.1	14.2	16.9	19.1	17.5
2000.....	15.4	14.4	13.1	11.8	11.5	14.0	16.8	19.4	19.9
1999.....	14.9	13.8	13.2	11.7	12.6	14.9	17.6	19.2	19.3

Non-Hispanic 4/, male

2007.....	28.9	28.3	26.0	24.2	23.0	23.3	34.1	36.8	43.1
2006.....	27.8	28.2	25.1	21.8	22.9	24.6	32.6	36.7	44.4
2005.....	27.0	26.4	23.3	22.8	21.6	26.1	32.4	42.8	45.9
2004.....	26.6	25.6	23.1	22.5	20.8	25.9	33.5	38.7	46.2
2003.....	25.7	25.6	23.8	21.7	21.7	26.8	32.6	41.0	48.6
2002.....	26.3	24.8	24.1	20.9	22.3	28.5	35.2	44.8	50.9
2001 6/.....	25.6	23.2	23.1	19.9	23.1	27.2	34.6	44.6	51.3
2000.....	24.2	22.0	20.6	19.1	20.0	26.5	35.5	44.9	58.4
1999.....	23.1	21.2	20.8	19.5	22.3	28.5	36.0	44.2	57.6

Non-Hispanic 4/, female

2007.....	9.4	9.4	8.5	7.0	4.7	4.0	4.2	3.7	3.1
2006.....	9.0	9.0	7.7	6.7	4.5	3.8	4.0	4.3	3.2
2005.....	8.9	8.2	6.9	5.7	4.3	4.0	3.5	4.8	4.1
2004.....	9.4	8.9	6.8	5.7	3.9	4.0	4.4	3.4	3.6
2003.....	8.4	7.9	7.1	5.2	4.1	3.8	4.1	4.1	3.3
2002.....	8.5	7.1	6.5	5.2	4.5	4.0	4.2	4.4	3.9
2001 6/.....	7.8	7.4	6.5	5.5	4.3	3.8	4.4	3.8	3.5
2000.....	7.0	7.1	6.1	5.2	4.1	4.1	3.8	4.4	4.3
1999.....	7.1	6.7	6.1	4.6	4.4	4.2	4.8	4.6	4.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic White, both sexes

2007 5/.....	22.0	21.3	19.4	16.9	14.8	14.0	18.3	17.9	17.3
2006 5/.....	21.0	21.0	18.3	15.5	14.5	14.5	17.6	18.1	17.4
2005 5/.....	20.4	19.4	16.8	15.5	13.7	15.3	17.1	20.7	18.3
2004 5/.....	20.3	19.4	16.3	15.1	13.0	14.8	17.9	17.9	17.9
2003 5/.....	19.3	18.8	16.9	14.4	13.5	15.2	17.3	19.4	18.4
2002.....	19.7	17.7	16.7	14.1	14.1	16.2	18.1	21.3	19.2
2001 6/.....	18.8	17.0	16.0	13.6	14.2	15.2	18.2	20.3	18.7
2000.....	17.3	15.9	14.4	13.1	12.6	15.0	18.0	20.6	21.3
1999.....	16.6	15.3	14.5	12.7	13.6	16.1	18.8	20.4	20.8

Non-Hispanic White, male

2007 5/.....	33.2	31.9	29.2	26.6	25.3	25.6	36.8	39.8	46.9
2006 5/.....	31.6	31.8	27.9	24.0	25.1	27.0	35.3	39.8	47.6
2005 5/.....	30.6	29.6	26.0	25.2	23.9	28.5	34.9	45.9	49.4
2004 5/.....	29.8	28.8	25.1	24.6	22.6	27.7	36.0	40.9	49.8
2003 5/.....	29.1	28.8	26.2	23.7	23.6	28.8	35.0	44.0	52.5
2002.....	29.4	27.7	26.5	23.2	24.3	31.0	37.3	48.2	54.3
2001 6/.....	28.8	25.8	25.2	21.5	25.1	28.8	37.0	47.4	55.2
2000.....	26.8	24.0	22.4	21.1	21.6	28.0	37.9	47.9	62.7
1999.....	25.5	23.2	22.5	21.0	23.8	30.5	38.4	46.9	62.2

Non-Hispanic White, female

2007 5/.....	11.0	10.9	9.8	7.7	5.4	4.4	4.4	3.9	3.3
2006 5/.....	10.5	10.5	8.9	7.5	5.0	4.0	4.3	4.4	3.4
2005 5/.....	10.3	9.4	7.9	6.3	4.6	4.3	3.8	5.0	4.0
2004 5/.....	11.0	10.1	7.9	6.2	4.4	4.2	4.7	3.6	3.6
2003 5/.....	9.7	8.9	7.9	5.8	4.5	4.1	4.4	4.3	3.4
2002.....	10.0	7.8	7.4	5.6	5.0	4.0	4.3	4.8	4.1
2001 6/.....	8.8	8.3	7.2	6.2	4.6	4.1	4.8	3.9	3.5
2000.....	8.0	7.9	6.8	5.8	4.6	4.5	4.0	4.4	4.6
1999.....	7.9	7.6	6.8	5.0	4.6	4.6	5.1	4.8	4.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes

2007.....	7.7	8.4	8.5	8.5	8.8	8.8	12.5	11.8	11.0
2006.....	7.8	8.3	8.1	8.1	8.5	9.4	11.8	12.2	10.9
2005.....	8.0	8.3	8.1	8.2	8.2	9.9	11.9	13.6	11.2
2004.....	8.1	8.0	7.8	8.3	8.0	9.8	12.2	12.3	11.1
2003.....	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6

All origins 3/, male

2007.....	12.9	14.7	15.0	15.4	16.8	18.0	27.5	28.5	32.9
2006.....	13.1	14.3	14.3	14.4	16.3	19.2	26.4	29.7	32.7
2005.....	13.7	14.6	14.1	15.3	15.5	20.2	26.6	33.3	33.8
2004.....	13.4	13.6	13.8	14.9	15.5	20.0	26.7	30.7	34.4
2003.....	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6

All origins 3/, female

2007.....	2.6	2.4	2.3	2.3	1.7	1.4	1.5	1.2	0.6
2006.....	2.7	2.6	2.3	2.4	1.8	1.4	1.1	1.3	0.7
2005.....	2.5	2.4	2.3	1.8	1.7	1.6	1.3	1.3	0.8
2004.....	2.8	2.7	2.1	2.3	1.4	1.7	1.7	1.0	*
2003.....	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic 4/, both sexes

2007.....	8.4	9.0	9.0	9.0	9.2	9.2	13.0	12.2	11.3
2006.....	8.5	8.9	8.7	8.5	9.0	10.0	12.3	12.6	11.2
2005.....	8.6	9.0	8.5	8.7	8.7	10.3	12.3	14.1	11.4
2004.....	8.6	8.5	8.2	8.7	8.4	10.1	12.7	12.5	11.4
2003.....	8.3	8.8	8.9	8.6	9.0	10.5	12.3	13.5	12.1
2002.....	8.2	8.4	9.0	8.4	9.1	11.1	12.8	14.7	12.1
2001.....	7.9	8.4	8.8	8.3	9.3	10.6	12.8	14.4	12.4
2000.....	7.7	8.1	7.8	8.0	8.6	10.5	12.7	14.3	13.4
1999.....	7.8	7.5	8.4	7.8	8.9	11.3	13.1	13.7	12.7

Non-Hispanic 4/, male

2007.....	14.1	15.7	15.9	16.2	17.5	18.7	28.5	29.5	34.1
2006.....	14.4	15.3	15.3	15.1	17.1	20.3	27.5	30.6	33.8
2005.....	14.7	15.7	14.9	16.2	16.5	21.0	27.3	34.6	34.7
2004.....	14.4	14.4	14.5	15.6	16.2	20.6	27.9	31.3	35.6
2003.....	14.1	15.3	15.4	15.9	17.2	21.5	27.2	33.5	37.5
2002.....	14.1	14.8	16.0	15.3	17.4	23.1	28.7	36.8	38.4
2001.....	13.6	14.5	15.4	14.6	17.8	22.0	28.6	36.1	39.7
2000.....	13.3	13.8	13.6	14.4	16.3	21.7	28.9	36.1	43.8
1999.....	13.2	13.1	14.5	14.5	17.3	23.2	29.4	35.1	42.2

Non-Hispanic 4/, female

2007.....	2.8	2.5	2.5	2.4	1.8	1.4	1.6	1.3	0.6
2006.....	2.9	2.8	2.4	2.5	1.9	1.6	1.2	1.3	0.7
2005.....	2.7	2.6	2.5	2.0	1.8	1.7	1.4	1.4	0.8
2004.....	3.1	2.9	2.3	2.4	1.5	1.7	1.8	1.0	*
2003.....	2.8	2.5	2.7	1.9	1.8	1.7	1.7	1.4	0.9
2002.....	2.5	2.3	2.5	2.2	1.9	1.4	1.5	1.3	0.8
2001.....	2.5	2.6	2.6	2.5	2.0	1.5	1.8	1.4	1.0
2000.....	2.3	2.6	2.3	2.2	1.9	1.6	1.4	1.5	1.1
1999.....	2.5	2.2	2.8	1.7	1.7	1.9	1.7	1.2	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic White, both sexes

2007 5/.....	9.8	10.4	10.4	10.1	10.4	10.3	14.3	13.4	12.5
2006 5/.....	10.0	10.3	10.0	9.6	10.1	11.2	13.6	13.8	12.0
2005 5/.....	10.0	10.4	9.8	9.9	9.8	11.5	13.6	15.3	12.5
2004 5/.....	10.0	9.9	9.2	9.8	9.4	11.2	13.9	13.5	12.5
2003 5/.....	9.8	10.0	10.1	9.6	10.0	11.6	13.5	14.6	13.2
2002.....	9.5	9.5	10.2	9.5	10.3	12.2	13.8	16.0	13.1
2001.....	9.1	9.5	9.9	9.3	10.4	11.5	13.9	15.3	13.4
2000.....	8.7	9.1	8.6	9.0	9.5	11.4	13.8	15.5	14.5
1999.....	8.7	8.5	9.4	8.6	9.8	12.3	14.1	14.6	13.7

Non-Hispanic White, male

2007 5/.....	16.4	17.9	18.1	17.9	19.5	20.8	30.9	32.3	37.7
2006 5/.....	16.7	17.6	17.3	16.7	19.0	22.5	30.0	33.4	36.3
2005 5/.....	16.9	17.8	16.8	18.1	18.6	23.0	29.7	37.5	37.9
2004 5/.....	16.4	16.6	16.1	17.5	17.9	22.4	30.0	33.6	39.0
2003 5/.....	16.3	17.3	17.3	17.6	19.2	23.5	29.5	36.1	41.0
2002.....	16.1	16.6	17.8	17.0	19.4	25.3	30.7	39.9	41.3
2001.....	15.6	16.3	17.1	16.2	19.7	23.7	30.8	38.5	43.0
2000.....	14.9	15.4	14.9	15.9	17.8	23.2	31.3	39.0	47.6
1999.....	14.7	14.5	15.8	15.9	18.8	24.9	31.6	37.3	45.7

Non-Hispanic White, female

2007 5/.....	3.3	3.0	2.9	2.8	2.1	1.6	1.7	1.4	0.6
2006 5/.....	3.5	3.2	2.9	2.9	2.2	1.8	1.4	1.5	0.7
2005 5/.....	3.3	3.1	2.9	2.2	2.0	1.9	1.6	1.5	0.9
2004 5/.....	3.7	3.3	2.6	2.7	1.8	1.9	2.0	1.1	*
2003 5/.....	3.3	2.9	3.1	2.1	1.9	1.9	1.9	1.4	1.0
2002.....	3.0	2.6	2.9	2.5	2.2	1.5	1.7	1.4	0.9
2001.....	2.8	2.9	3.0	2.9	2.2	1.6	1.9	1.4	1.1
2000.....	2.7	2.9	2.6	2.5	2.2	1.9	1.5	1.6	1.2
1999.....	2.8	2.6	3.2	1.9	1.8	2.2	1.8	1.3	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes

2007.....	10.0	9.3	7.7	6.0	4.1	3.5	3.9	4.3	4.6
2006.....	9.3	9.0	7.2	5.3	4.0	3.2	3.8	4.1	5.0
2005.....	8.7	7.8	6.2	5.1	3.7	3.6	3.5	5.2	5.7
2004.....	8.9	8.2	6.5	4.9	3.4	3.6	3.9	4.4	5.4
2003.....	8.2	7.5	6.2	4.5	3.4	3.5	3.7	4.6	5.1
2002.....	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 6/.....	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000.....	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
1999.....	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7

All origins 3/, male

2007.....	14.0	12.3	10.0	7.9	5.6	4.7	5.7	7.4	8.8
2006.....	12.9	12.2	9.6	6.8	5.7	4.4	5.3	6.0	10.5
2005.....	11.8	10.4	8.3	6.7	5.1	5.2	5.3	8.1	11.2
2004.....	11.8	10.8	8.6	6.8	4.7	5.4	5.7	7.6	10.7
2003.....	11.2	9.9	8.5	5.8	4.7	5.4	5.5	7.7	11.1
2002.....	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 6/.....	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000.....	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
1999.....	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5

All origins 3/, female

2007.....	6.1	6.5	5.7	4.3	2.8	2.5	2.5	2.3	2.5
2006.....	5.7	5.9	4.9	4.0	2.6	2.3	2.7	2.9	2.4
2005.....	5.7	5.3	4.3	3.6	2.4	2.2	2.2	3.4	3.1
2004.....	6.0	5.7	4.5	3.2	2.2	2.2	2.5	2.5	3.0
2003.....	5.3	5.1	4.1	3.2	2.4	2.0	2.4	2.6	2.4
2002.....	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 6/.....	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000.....	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
1999.....	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic 4/, both sexes

2007.....	10.7	9.7	8.0	6.2	4.1	3.5	3.9	4.3	4.6
2006.....	9.7	9.5	7.5	5.4	4.1	3.2	3.8	4.2	5.1
2005.....	9.2	8.1	6.4	5.1	3.7	3.6	3.4	5.3	5.8
2004.....	9.3	8.5	6.5	5.0	3.4	3.6	3.9	4.3	5.4
2003.....	8.6	7.8	6.3	4.5	3.4	3.5	3.7	4.5	5.1
2002.....	9.0	7.4	6.0	4.3	3.7	3.9	4.2	5.0	5.9
2001 6/.....	8.6	6.7	5.7	4.1	3.7	3.6	4.1	4.7	5.1
2000.....	7.7	6.3	5.3	3.8	2.9	3.5	4.1	5.1	6.5
1999.....	7.2	6.3	4.8	3.9	3.7	3.6	4.6	5.5	6.6

Non-Hispanic 4/, male

2007.....	14.8	12.6	10.1	8.0	5.5	4.6	5.7	7.3	9.0
2006.....	13.5	12.9	9.9	6.8	5.9	4.3	5.1	6.1	10.6
2005.....	12.3	10.8	8.4	6.7	5.1	5.1	5.0	8.2	11.3
2004.....	12.3	11.2	8.6	6.9	4.6	5.3	5.6	7.4	10.6
2003.....	11.6	10.3	8.4	5.8	4.5	5.3	5.4	7.5	11.2
2002.....	12.1	10.0	8.1	5.6	4.9	5.4	6.4	8.0	12.5
2001 6/.....	12.0	8.8	7.6	5.3	5.4	5.2	6.0	8.6	11.6
2000.....	10.9	8.2	6.9	4.7	3.7	4.7	6.7	8.8	14.6
1999.....	9.9	8.2	6.3	5.0	5.0	5.3	6.7	9.1	15.4

Non-Hispanic 4/, female

2007.....	6.6	6.9	6.0	4.5	2.9	2.6	2.6	2.5	2.5
2006.....	6.1	6.3	5.3	4.1	2.6	2.3	2.8	3.0	2.5
2005.....	6.2	5.6	4.4	3.8	2.5	2.3	2.2	3.4	3.2
2004.....	6.3	6.0	4.6	3.3	2.4	2.3	2.6	2.4	3.0
2003.....	5.6	5.4	4.4	3.4	2.4	2.1	2.5	2.7	2.4
2002.....	5.9	4.8	4.0	3.0	2.6	2.6	2.6	3.2	3.1
2001 6/.....	5.4	4.8	3.9	3.0	2.3	2.3	2.7	2.4	2.5
2000.....	4.6	4.5	3.8	2.9	2.2	2.5	2.4	2.9	3.2
1999.....	4.5	4.5	3.3	2.8	2.6	2.3	3.1	3.5	3.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic White, both sexes

2007 5/.....	12.2	10.9	9.0	6.8	4.5	3.7	4.0	4.4	4.8
2006 5/.....	11.0	10.7	8.3	5.9	4.4	3.2	4.0	4.3	5.4
2005 5/.....	10.4	9.1	7.0	5.6	3.9	3.8	3.5	5.4	5.7
2004 5/.....	10.3	9.5	7.1	5.2	3.6	3.6	4.1	4.3	5.4
2003 5/.....	9.6	8.7	6.9	4.8	3.5	3.6	3.8	4.8	5.2
2002.....	10.1	8.2	6.6	4.6	3.8	4.0	4.3	5.3	6.1
2001 6/.....	9.6	7.4	6.2	4.3	3.8	3.7	4.3	5.0	5.3
2000.....	8.6	6.8	5.9	4.2	3.1	3.6	4.2	5.1	6.8
1999.....	7.9	6.8	5.1	4.1	3.8	3.8	4.7	5.8	7.0

Non-Hispanic White, male

2007 5/.....	16.8	14.0	11.1	8.7	5.8	4.8	5.8	7.5	9.2
2006 5/.....	14.9	14.2	10.7	7.3	6.0	4.5	5.3	6.5	11.2
2005 5/.....	13.8	11.8	9.1	7.2	5.3	5.5	5.2	8.5	11.5
2004 5/.....	13.3	12.3	9.0	7.2	4.8	5.3	6.0	7.3	10.8
2003 5/.....	12.8	11.5	9.0	6.0	4.5	5.3	5.6	7.9	11.6
2002.....	13.3	11.2	8.7	6.1	4.9	5.7	6.7	8.3	13.0
2001 6/.....	13.3	9.5	8.1	5.3	5.4	5.1	6.2	9.0	12.3
2000.....	11.9	8.6	7.6	5.1	3.8	4.8	6.6	8.9	15.2
1999.....	10.7	8.7	6.7	5.2	5.0	5.6	6.8	9.6	16.5

Non-Hispanic White, female

2007 5/.....	7.7	7.9	6.9	4.9	3.3	2.8	2.7	2.5	2.7
2006 5/.....	7.1	7.2	6.0	4.6	2.9	2.2	3.0	3.0	2.7
2005 5/.....	7.1	6.3	4.9	4.1	2.6	2.4	2.2	3.5	3.1
2004 5/.....	7.3	6.8	5.3	3.5	2.6	2.3	2.6	2.5	3.0
2003 5/.....	6.3	6.0	4.8	3.7	2.5	2.2	2.5	2.9	2.4
2002.....	7.0	5.2	4.5	3.2	2.8	2.6	2.6	3.4	3.2
2001 6/.....	6.0	5.4	4.3	3.3	2.4	2.5	2.9	2.5	2.4
2000.....	5.3	5.0	4.2	3.3	2.4	2.6	2.5	2.8	3.4
1999.....	5.1	5.0	3.6	3.1	2.7	2.4	3.3	3.5	3.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes

2007.....	5.4	4.3	3.3	2.6	2.2	2.0	2.0	2.1	1.5
2006.....	5.6	4.5	3.4	2.9	2.0	2.2	2.0	2.3	1.9
2005.....	5.4	4.2	2.9	2.7	2.3	2.4	2.2	2.1	2.1
2004.....	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
2003.....	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 6/.....	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4

All origins 3/, male

2007.....	7.8	6.7	5.1	4.0	3.3	2.9	3.0	2.5	1.5
2006.....	8.5	6.7	5.2	4.4	2.9	3.2	2.9	2.3	2.3
2005.....	8.3	6.7	4.5	4.2	3.1	3.5	2.8	2.4	2.7
2004.....	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
2003.....	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 6/.....	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4

All origins 3/, female

2007.....	3.1	2.0	1.6	1.4	1.3	1.2	1.2	1.9	1.4
2006.....	2.8	2.5	1.7	1.6	1.3	1.4	1.4	2.3	1.8
2005.....	2.6	1.8	1.5	1.4	1.6	1.5	1.8	2.0	1.9
2004.....	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
2003.....	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 6/.....	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	5.3	4.2	3.2	2.6	2.3	1.9	2.0	2.1	1.5
2006.....	5.5	4.4	3.4	2.8	2.0	2.2	2.0	2.3	1.9
2005.....	5.3	4.0	2.9	2.7	2.3	2.4	2.2	2.0	2.2
2004.....	5.2	4.0	3.1	2.8	2.6	2.2	2.2	2.3	2.0
2003.....	5.4	4.1	3.1	2.3	2.2	2.2	2.4	2.3	2.1
2002.....	5.3	3.9	3.3	2.6	2.4	2.2	2.3	2.2	2.1
2001 6/.....	6.8	5.3	4.3	3.3	3.1	2.5	2.5	2.4	2.4
2000.....	5.0	3.8	3.2	2.5	2.4	2.3	2.3	2.6	2.3
1999.....	4.8	3.8	3.0	2.8	2.6	2.4	2.5	2.4	2.4

Non-Hispanic 4/, male

2007.....	7.5	6.6	4.9	3.9	3.2	2.7	3.0	2.5	1.4
2006.....	8.2	6.5	5.0	4.1	2.9	3.1	2.9	2.3	2.1
2005.....	8.1	6.2	4.4	4.1	3.0	3.5	2.6	2.2	2.8
2004.....	7.5	6.3	4.7	4.0	3.8	3.2	2.6	2.8	2.4
2003.....	7.9	6.1	4.5	3.4	3.2	3.0	3.4	2.9	3.0
2002.....	7.7	5.8	5.0	4.0	3.3	3.1	3.3	2.8	2.9
2001 6/.....	10.2	8.0	6.6	4.7	4.3	3.2	3.2	2.9	3.1
2000.....	7.4	5.6	4.9	3.7	3.2	3.2	2.7	3.8	3.1
1999.....	7.3	5.5	4.8	4.1	3.9	3.0	3.3	3.3	3.4

Non-Hispanic 4/, female

2007.....	3.2	2.0	1.6	1.4	1.4	1.2	1.2	1.9	1.5
2006.....	2.8	2.5	1.8	1.7	1.2	1.5	1.4	2.4	1.8
2005.....	2.6	1.8	1.5	1.4	1.7	1.5	1.8	1.9	1.9
2004.....	3.0	1.8	1.6	1.7	1.6	1.3	2.0	2.0	1.8
2003.....	2.9	2.1	1.7	1.3	1.4	1.6	1.7	2.0	1.8
2002.....	2.9	2.1	1.8	1.4	1.5	1.5	1.5	1.9	1.7
2001 6/.....	3.5	2.7	2.2	2.1	2.1	1.9	2.0	2.2	2.1
2000.....	2.6	2.0	1.6	1.4	1.6	1.5	2.0	1.9	2.0
1999.....	2.4	2.2	1.4	1.6	1.5	1.9	2.0	1.9	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic White, both sexes

2007 5/.....	3.4	3.0	2.2	1.9	1.7	1.6	1.4	1.9	1.2
2006 5/.....	3.5	3.0	2.4	2.2	1.4	1.9	1.7	2.3	1.7
2005 5/.....	3.4	2.5	2.2	2.1	1.7	1.9	1.9	1.8	1.9
2004 5/.....	3.4	2.6	2.3	2.1	2.2	1.7	1.8	2.1	1.8
2003 5/.....	3.4	2.8	2.3	1.7	1.7	1.7	2.0	2.0	2.0
2002.....	3.4	2.6	2.6	2.0	1.9	1.9	1.8	1.9	1.8
2001 6/.....	5.1	4.1	3.7	2.8	2.5	2.1	2.1	2.1	2.2
2000.....	3.2	2.6	2.4	1.8	1.7	1.8	2.0	2.0	2.1
1999.....	3.1	2.7	2.3	2.1	2.2	2.0	2.0	2.1	2.1

Non-Hispanic White, male

2007 5/.....	4.5	4.3	3.1	2.6	2.2	2.3	1.9	2.1	*
2006 5/.....	4.9	4.1	3.2	3.1	1.9	2.3	2.3	2.1	1.8
2005 5/.....	4.9	3.7	3.1	3.1	1.9	2.5	2.0	1.8	2.0
2004 5/.....	4.6	3.9	3.2	2.9	2.8	2.4	2.0	2.3	1.9
2003 5/.....	4.6	4.0	3.1	2.5	2.3	2.2	2.8	2.3	2.9
2002.....	4.3	3.5	3.6	2.9	2.5	2.4	2.4	2.1	2.7
2001 6/.....	7.4	6.0	5.3	3.8	3.2	2.5	2.6	2.4	2.8
2000.....	4.3	3.5	3.5	2.6	2.1	2.4	2.3	2.6	2.6
1999.....	4.5	3.9	3.4	3.0	3.1	2.3	2.5	2.8	2.6

Non-Hispanic White, female

2007 5/.....	2.4	1.6	1.3	1.3	1.3	1.0	1.1	1.8	1.4
2006 5/.....	2.1	1.9	1.6	1.4	1.1	1.5	1.3	2.4	1.7
2005 5/.....	1.9	1.3	1.3	1.1	1.4	1.3	1.8	1.8	1.9
2004 5/.....	2.3	1.3	1.4	1.4	1.7	1.1	1.6	1.9	1.7
2003 5/.....	2.3	1.7	1.6	1.0	1.2	1.3	1.5	1.8	1.6
2002.....	2.5	1.6	1.7	1.3	1.3	1.5	1.3	1.8	1.5
2001 6/.....	2.9	2.3	2.0	1.8	1.9	1.7	1.7	1.9	2.0
2000.....	2.0	1.7	1.4	1.2	1.4	1.3	1.8	1.6	1.9
1999.....	1.7	1.6	1.2	1.4	1.5	1.7	1.7	1.6	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic Black, both sexes

2007 5/.....	17.4	13.5	10.8	8.0	6.8	4.7	7.5	5.3	*
2006 5/.....	17.5	14.7	10.9	8.0	6.6	5.4	5.2	*	*
2005 5/.....	17.6	14.1	8.4	8.0	7.3	7.1	6.0	5.7	*
2004 5/.....	16.8	13.6	9.4	7.9	5.8	6.6	7.1	5.3	*
2003 5/.....	18.1	12.9	9.1	6.9	5.8	6.9	6.4	6.7	*
2002.....	17.9	13.2	9.0	7.7	6.0	5.5	8.2	5.9	*
2001 6/.....	18.2	13.9	10.0	7.5	7.6	6.6	7.1	6.3	*
2000.....	17.1	12.4	9.2	7.5	8.1	7.0	5.4	9.8	*
1999.....	16.4	11.2	9.0	7.6	6.7	6.1	8.4	6.2	*

Non-Hispanic Black, male

2007 5/.....	28.1	23.9	20.1	15.7	13.0	7.2	16.4	*	*
2006 5/.....	30.1	24.8	19.7	13.5	12.3	10.4	10.6	*	*
2005 5/.....	30.1	24.5	15.4	13.3	13.0	13.0	11.5	*	*
2004 5/.....	27.6	24.2	16.6	13.1	12.0	11.6	10.3	*	*
2003 5/.....	30.6	22.2	17.4	11.5	11.1	11.5	11.7	*	*
2002.....	31.4	22.8	17.2	13.8	10.2	10.9	15.0	*	*
2001 6/.....	30.1	23.6	17.6	11.7	13.0	12.1	11.9	*	*
2000.....	29.2	22.5	16.3	13.4	14.9	12.6	*	19.7	*
1999.....	27.7	18.5	17.2	13.1	13.0	10.4	14.0	*	*

Non-Hispanic Black, female

2007 5/.....	8.1	4.7	3.3	*	*	*	*	*	*
2006 5/.....	6.5	6.2	3.7	3.7	*	*	*	*	*
2005 5/.....	6.7	5.4	2.6	3.8	*	*	*	*	*
2004 5/.....	7.4	4.7	3.5	3.9	*	*	*	*	*
2003 5/.....	7.1	5.1	*	3.2	*	*	*	*	*
2002.....	6.3	5.0	*	*	*	*	*	*	*
2001 6/.....	7.8	5.7	3.8	4.1	*	*	*	*	*
2000.....	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	6.6	5.1	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes

2007.....	3.1	2.2	1.5	1.1	1.0	0.9	0.7	0.6	0.4
2006.....	3.1	2.4	1.6	1.4	0.9	0.8	0.7	0.6	0.4
2005.....	3.0	2.1	1.4	1.3	1.1	1.1	0.8	0.6	0.5
2004.....	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
2003.....	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All origins 3/, male

2007.....	4.8	3.5	2.4	1.9	1.4	1.3	1.1	*	*
2006.....	4.9	3.6	2.3	2.1	1.2	1.3	1.0	*	*
2005.....	4.7	3.4	2.2	2.0	1.4	1.7	1.0	*	*
2004.....	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
2003.....	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All origins 3/, female

2007.....	1.4	0.9	0.7	0.5	0.7	0.5	*	0.6	0.5
2006.....	1.3	1.2	0.9	0.7	0.6	0.4	*	0.7	*
2005.....	1.4	0.8	0.6	0.6	0.8	0.6	0.7	0.6	*
2004.....	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
2003.....	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic 4/, both sexes

2007	3.0	2.1	1.5	1.2	1.0	0.9	0.7	0.7	0.4
2006	3.0	2.3	1.6	1.3	0.9	0.8	0.7	0.7	0.4
2005	2.9	2.0	1.4	1.3	1.1	1.0	0.9	0.6	0.5
2004	2.9	2.1	1.7	1.4	1.2	0.9	0.9	0.7	0.5
2003	3.0	2.2	1.6	1.1	0.8	0.9	0.7	0.6	0.5
2002	2.8	2.1	1.7	1.3	1.0	0.8	1.0	0.8	*
2001	2.6	2.0	1.5	1.4	1.1	0.9	0.9	0.7	*
2000	2.7	2.0	1.7	1.2	1.0	0.9	0.7	0.6	0.7
1999	2.6	2.0	1.5	1.5	1.1	1.1	0.8	0.7	0.5

Non-Hispanic 4/, male

2007	4.6	3.4	2.3	1.9	1.4	1.3	1.1	*	*
2006	4.8	3.5	2.3	2.0	1.2	1.3	1.0	*	*
2005	4.5	3.2	2.2	2.0	1.3	1.6	1.1	*	*
2004	4.4	3.4	2.7	2.0	1.7	1.5	1.0	*	*
2003	4.7	3.5	2.5	1.7	1.3	1.3	0.8	*	*
2002	4.1	3.3	2.7	1.9	1.4	1.0	1.4	*	*
2001	4.1	3.2	2.3	2.0	1.6	1.1	1.3	*	*
2000	4.2	3.1	2.5	1.8	1.4	1.2	0.8	*	*
1999	4.1	2.9	2.5	2.5	1.5	1.5	1.0	*	*

Non-Hispanic 4/, female

2007	1.4	0.9	0.7	0.5	0.7	0.5	*	0.6	0.6
2006	1.3	1.2	0.9	0.7	0.6	0.5	*	0.8	*
2005	1.3	0.9	0.6	0.6	0.8	0.5	0.7	0.7	*
2004	1.4	0.9	0.8	0.9	0.8	*	0.7	0.8	*
2003	1.4	1.0	0.8	0.6	*	0.5	0.7	0.8	*
2002	1.5	1.0	0.9	0.7	0.6	0.6	0.7	0.7	*
2001	1.1	0.9	0.8	0.8	0.7	0.7	0.6	*	*
2000	1.3	1.0	0.9	0.6	0.7	0.7	0.7	*	*
1999	1.1	1.1	0.5	0.7	0.6	0.7	0.7	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic White, both sexes

2007 5/.....	1.8	1.4	0.9	0.9	0.8	0.8	0.6	0.6	*
2006 5/.....	1.8	1.4	1.2	1.0	0.6	0.7	0.5	0.7	*
2005 5/.....	1.8	1.3	1.0	1.0	0.8	0.8	0.8	0.5	*
2004 5/.....	1.9	1.4	1.2	1.2	1.1	0.7	0.8	0.7	*
2003 5/.....	1.8	1.5	1.3	0.9	0.6	0.7	0.6	0.6	0.5
2002.....	1.8	1.4	1.4	0.9	0.7	0.7	0.9	0.7	*
2001.....	1.7	1.4	1.2	1.2	1.0	0.7	0.7	0.6	*
2000.....	1.7	1.4	1.3	0.9	0.8	0.7	0.7	*	0.6
1999.....	1.6	1.4	1.1	1.2	0.9	0.9	0.6	0.7	0.6

Non-Hispanic White, male

2007 5/.....	2.6	2.0	1.3	1.3	0.9	1.1	*	*	*
2006 5/.....	2.4	2.0	1.5	1.4	0.6	0.9	*	*	*
2005 5/.....	2.5	2.0	1.4	1.6	0.9	1.1	0.9	*	*
2004 5/.....	2.6	2.0	1.8	1.6	1.3	1.1	0.9	*	*
2003 5/.....	2.5	2.3	1.8	1.3	1.0	1.0	*	*	*
2002.....	2.2	2.1	2.0	1.3	0.9	0.8	1.2	*	*
2001.....	2.5	2.1	1.5	1.7	1.2	0.8	1.0	*	*
2000.....	2.5	1.9	1.9	1.2	1.0	0.9	0.8	*	*
1999.....	2.4	2.1	1.7	1.7	1.3	1.2	*	*	*

Non-Hispanic White, female

2007 5/.....	1.1	0.8	0.6	0.4	0.7	*	*	*	*
2006 5/.....	1.2	0.9	0.8	0.7	0.6	*	*	0.8	*
2005 5/.....	1.1	0.7	0.6	0.5	0.8	0.5	0.8	*	*
2004 5/.....	1.1	0.7	0.7	0.8	0.9	*	0.6	0.9	*
2003 5/.....	1.2	0.8	0.8	0.5	*	*	0.6	0.8	*
2002.....	1.3	0.8	0.9	0.5	0.5	0.7	0.6	0.7	*
2001.....	0.9	0.8	0.8	0.8	0.8	0.6	0.5	*	*
2000.....	1.0	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	0.8	0.7	0.5	0.7	0.7	0.7	0.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic Black, both sexes

2007 5/.....	10.6	7.3	5.8	3.7	3.0	*	*	*	*
2006 5/.....	10.5	8.5	4.7	3.7	3.3	*	*	*	*
2005 5/.....	10.1	7.0	3.8	3.4	2.9	3.4	*	*	*
2004 5/.....	9.5	7.5	5.1	4.1	2.7	*	*	*	*
2003 5/.....	10.5	7.0	4.6	3.1	2.5	*	*	*	*
2002.....	9.7	7.2	4.2	4.1	2.8	*	*	*	*
2001.....	8.5	6.6	4.4	2.9	2.8	*	*	*	*
2000.....	9.7	6.9	4.3	2.7	2.8	2.7	*	*	*
1999.....	9.7	6.1	4.6	4.3	2.3	*	*	*	*

Non-Hispanic Black, male

2007 5/.....	18.7	13.9	10.8	7.8	5.7	*	*	*	*
2006 5/.....	20.1	15.2	8.7	6.9	6.3	*	*	*	*
2005 5/.....	18.5	12.7	7.6	5.8	5.4	7.2	*	*	*
2004 5/.....	16.6	14.2	9.7	6.9	5.7	*	*	*	*
2003 5/.....	19.7	12.3	9.1	5.2	*	*	*	*	*
2002.....	17.6	12.3	8.4	6.7	5.4	*	*	*	*
2001.....	15.6	12.1	8.7	5.0	6.0	*	*	*	*
2000.....	17.1	13.1	7.3	5.5	5.3	*	*	*	*
1999.....	17.4	9.8	9.6	8.5	*	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	3.6	1.6	*	*	*	*	*	*	*
2006 5/.....	2.2	2.9	*	*	*	*	*	*	*
2005 5/.....	2.9	2.2	*	*	*	*	*	*	*
2004 5/.....	3.3	1.8	*	*	*	*	*	*	*
2003 5/.....	2.6	2.6	*	*	*	*	*	*	*
2002.....	2.8	2.8	*	*	*	*	*	*	*
2001.....	2.4	2.0	*	*	*	*	*	*	*
2000.....	3.3	*	*	*	*	*	*	*	*
1999.....	3.1	2.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes

2007.....	2.3	2.1	1.7	1.5	1.2	1.1	1.3	1.5	1.0
2006.....	2.5	2.2	1.8	1.6	1.2	1.4	1.4	1.7	1.5
2005.....	2.4	2.1	1.6	1.4	1.2	1.4	1.4	1.5	1.7
2004.....	2.5	2.1	1.5	1.3	1.4	1.4	1.4	1.6	1.5
2003.....	2.5	1.9	1.5	1.3	1.5	1.5	1.8	1.8	1.7
2002.....	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 6/.....	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000.....	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
1999.....	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8

All origins 3/, male

2007.....	3.0	3.2	2.7	2.1	1.9	1.6	2.0	1.8	1.3
2006.....	3.6	3.1	2.9	2.3	1.7	1.9	2.0	1.8	1.8
2005.....	3.6	3.2	2.3	2.1	1.7	1.9	1.8	1.8	2.1
2004.....	3.4	3.2	2.2	2.0	2.0	1.8	1.6	2.2	1.7
2003.....	3.3	2.9	2.2	1.9	2.1	2.0	2.7	2.6	2.5
2002.....	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 6/.....	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000.....	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
1999.....	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6

All origins 3/, female

2007.....	1.7	1.1	0.8	1.0	0.6	0.7	0.8	1.3	0.9
2006.....	1.5	1.3	0.9	0.9	0.7	1.0	0.9	1.6	1.4
2005.....	1.2	1.0	0.9	0.8	0.8	1.0	1.1	1.3	1.5
2004.....	1.6	1.0	0.8	0.8	0.8	1.0	1.2	1.2	1.5
2003.....	1.6	1.0	0.9	0.7	1.0	1.0	1.1	1.3	1.4
2002.....	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 6/.....	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000.....	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999.....	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic 4/, both sexes

2007.....	2.3	2.1	1.7	1.4	1.2	1.0	1.3	1.5	1.1
2006.....	2.5	2.1	1.8	1.5	1.1	1.4	1.4	1.7	1.5
2005.....	2.4	2.0	1.5	1.4	1.2	1.4	1.3	1.4	1.7
2004.....	2.3	1.9	1.4	1.3	1.4	1.3	1.4	1.6	1.5
2003.....	2.4	1.9	1.4	1.2	1.4	1.4	1.7	1.7	1.6
2002.....	2.5	1.7	1.6	1.4	1.4	1.4	1.3	1.5	1.7
2001 6/.....	4.2	3.3	2.8	2.0	1.9	1.6	1.6	1.7	2.1
2000.....	2.3	1.8	1.5	1.3	1.4	1.3	1.5	2.1	1.6
1999.....	2.2	1.8	1.5	1.2	1.6	1.3	1.7	1.7	1.8

Non-Hispanic 4/, male

2007.....	2.9	3.2	2.6	2.0	1.8	1.4	1.9	1.8	1.3
2006.....	3.5	3.0	2.7	2.2	1.7	1.8	2.0	1.8	1.7
2005.....	3.6	3.0	2.2	2.1	1.6	1.9	1.6	1.7	2.2
2004.....	3.1	2.9	2.0	1.9	2.0	1.7	1.6	2.3	1.7
2003.....	3.2	2.7	2.1	1.7	1.9	1.7	2.6	2.5	2.3
2002.....	3.6	2.5	2.3	2.1	1.9	2.1	1.8	1.8	2.3
2001 6/.....	6.1	4.9	4.3	2.6	2.6	2.1	1.9	2.0	2.7
2000.....	3.2	2.5	2.4	1.9	1.9	2.0	1.9	3.1	2.4
1999.....	3.1	2.6	2.3	1.6	2.4	1.6	2.3	2.6	2.6

Non-Hispanic 4/, female

2007.....	1.8	1.1	0.9	0.9	0.7	0.6	0.8	1.3	0.9
2006.....	1.5	1.2	0.9	0.9	0.6	1.0	0.9	1.6	1.4
2005.....	1.3	1.0	0.9	0.8	0.8	0.9	1.1	1.3	1.4
2004.....	1.6	0.9	0.8	0.8	0.8	1.0	1.2	1.2	1.4
2003.....	1.6	1.1	0.8	0.7	1.0	1.1	1.0	1.2	1.3
2002.....	1.4	1.0	1.0	0.7	0.9	0.8	0.9	1.2	1.4
2001 6/.....	2.4	1.7	1.4	1.3	1.4	1.2	1.4	1.6	1.8
2000.....	1.4	1.0	0.7	0.8	0.9	0.8	1.2	1.5	1.3
1999.....	1.2	1.1	0.8	0.8	0.9	1.1	1.3	1.2	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic White, both sexes

2007 5/.....	1.6	1.6	1.2	1.0	0.9	0.8	0.9	1.3	0.9
2006 5/.....	1.7	1.5	1.2	1.2	0.8	1.2	1.2	1.6	1.4
2005 5/.....	1.6	1.2	1.2	1.0	0.8	1.1	1.0	1.2	1.5
2004 5/.....	1.6	1.2	1.1	1.0	1.1	1.0	1.0	1.4	1.3
2003 5/.....	1.6	1.3	1.1	0.8	1.1	1.0	1.4	1.4	1.5
2002.....	1.6	1.1	1.2	1.1	1.1	1.1	0.9	1.2	1.4
2001 6/.....	3.4	2.7	2.5	1.6	1.5	1.3	1.3	1.5	1.9
2000.....	1.5	1.2	1.1	0.9	0.9	1.0	1.3	1.6	1.5
1999.....	1.5	1.4	1.2	1.0	1.3	1.1	1.4	1.4	1.5

Non-Hispanic White, male

2007 5/.....	2.0	2.3	1.8	1.2	1.3	1.2	1.2	1.5	*
2006 5/.....	2.5	2.1	1.6	1.7	1.2	1.4	1.7	1.6	1.6
2005 5/.....	2.3	1.7	1.6	1.5	1.0	1.5	1.1	1.4	1.6
2004 5/.....	2.0	1.9	1.5	1.3	1.5	1.4	1.1	1.9	*
2003 5/.....	2.0	1.7	1.4	1.2	1.3	1.2	2.1	2.0	2.2
2002.....	2.1	1.4	1.5	1.6	1.6	1.6	1.2	1.4	2.0
2001 6/.....	4.9	3.9	3.8	2.2	2.1	1.7	1.6	1.8	2.3
2000.....	1.9	1.6	1.6	1.3	1.1	1.5	1.5	2.2	1.9
1999.....	2.1	1.8	1.7	1.3	1.8	1.2	1.8	2.1	*

Non-Hispanic White, female

2007 5/.....	1.3	0.8	0.7	0.8	0.6	*	0.6	1.2	0.8
2006 5/.....	1.0	0.9	0.7	0.7	0.5	1.1	0.8	1.5	1.3
2005 5/.....	0.9	0.6	0.7	0.6	0.7	0.7	1.0	1.1	1.4
2004 5/.....	1.2	0.6	0.6	0.6	0.8	0.7	1.0	1.1	1.3
2003 5/.....	1.1	0.9	0.8	0.5	0.9	0.9	0.9	1.0	1.2
2002.....	1.1	0.8	0.9	0.7	0.7	0.8	0.7	1.1	1.2
2001 6/.....	2.0	1.5	1.2	1.0	1.1	1.1	1.2	1.2	1.7
2000.....	1.1	0.9	0.5	0.5	0.7	0.7	1.1	1.2	1.3
1999.....	0.9	0.9	0.7	0.7	0.8	1.0	1.1	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic Black, both sexes

2007 5/.....	6.8	6.2	5.0	4.3	3.8	2.7	5.1	*	*
2006 5/.....	7.0	6.2	6.3	4.3	3.3	3.1	*	*	*
2005 5/.....	7.5	7.2	4.6	4.6	4.4	3.7	4.6	*	*
2004 5/.....	7.3	6.2	4.3	3.8	3.2	4.4	5.2	*	*
2003 5/.....	7.5	5.9	4.5	3.7	3.3	4.3	4.7	5.2	*
2002.....	8.2	6.0	4.8	3.6	3.2	3.7	5.8	*	*
2001 6/.....	9.6	7.3	5.7	4.5	4.9	4.1	4.8	*	*
2000.....	7.4	5.5	5.0	4.8	5.3	4.2	4.2	6.9	*
1999.....	6.7	5.2	4.4	3.3	4.4	3.5	5.3	*	*

Non-Hispanic Black, male

2007 5/.....	9.4	9.9	9.2	7.8	7.3	*	10.5	*	*
2006 5/.....	10.0	9.6	11.0	6.6	6.0	*	*	*	*
2005 5/.....	11.6	11.8	7.8	7.5	7.6	*	*	*	*
2004 5/.....	11.0	10.0	6.9	6.2	6.2	*	*	*	*
2003 5/.....	11.0	9.9	8.2	6.4	6.3	6.7	*	*	*
2002.....	13.7	10.5	8.7	7.1	*	7.2	10.3	*	*
2001 6/.....	14.6	11.5	8.9	6.7	7.0	7.2	*	*	*
2000.....	12.2	9.5	9.0	7.9	9.6	6.8	*	*	*
1999.....	10.3	8.8	7.7	4.6	8.5	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	4.6	3.0	*	*	*	*	*	*	*
2006 5/.....	4.3	3.3	2.3	*	*	*	*	*	*
2005 5/.....	3.8	3.2	*	*	*	*	*	*	*
2004 5/.....	4.1	2.9	*	*	*	*	*	*	*
2003 5/.....	4.6	2.6	*	*	*	*	*	*	*
2002.....	3.5	2.2	*	*	*	*	*	*	*
2001 6/.....	5.4	3.7	3.0	*	*	*	*	*	*
2000.....	3.3	2.0	*	*	*	*	*	*	*
1999.....	3.5	2.1	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	3.5	3.3	2.2	1.4	1.0	0.9	0.9	1.2	1.3
2006.....	3.5	2.8	1.8	1.2	0.9	0.8	1.0	1.1	1.4
2005.....	3.2	2.9	1.6	1.1	0.9	0.7	0.8	0.9	1.5
2004.....	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

All origins 3/, male

2007.....	4.0	3.9	2.7	1.7	1.3	1.1	1.3	1.7	1.6
2006.....	4.2	3.5	2.3	1.4	1.2	0.9	1.2	1.7	2.2
2005.....	3.8	3.5	2.1	1.1	1.1	1.0	1.0	*	1.6
2004.....	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
2003.....	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1

All origins 3/, female

2007.....	3.0	2.8	1.7	1.1	0.7	0.7	0.6	0.8	1.1
2006.....	2.7	2.0	1.5	0.9	0.7	0.7	0.8	0.8	1.0
2005.....	2.7	2.4	1.2	1.1	0.8	0.5	0.6	0.9	1.4
2004.....	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
2003.....	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	3.8	3.6	2.3	1.5	1.0	0.9	0.9	1.2	1.3
2006.....	3.7	2.9	1.9	1.2	0.9	0.9	1.0	1.2	1.3
2005.....	3.4	3.0	1.7	1.1	1.0	0.8	0.7	0.9	1.5
2004.....	3.8	2.7	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.7	2.7	1.6	1.0	0.8	0.6	0.6	0.9	1.1
2002.....	3.5	2.4	1.4	1.0	0.8	0.6	0.6	1.0	1.4
2001.....	3.1	2.0	1.1	0.6	0.6	0.5	0.7	1.1	0.9
2000.....	2.8	1.8	1.1	0.7	0.5	0.6	0.6	0.7	1.4
1999.....	2.8	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

Non-Hispanic 4/, male

2007.....	4.4	4.1	2.8	1.8	1.3	1.1	1.3	1.7	1.6
2006.....	4.5	3.7	2.3	1.4	1.2	0.9	1.3	1.8	2.2
2005.....	4.0	3.6	2.1	1.1	1.1	1.0	0.9	*	1.7
2004.....	4.8	3.4	2.0	1.1	1.2	0.9	0.8	1.3	1.7
2003.....	4.8	3.5	2.0	1.2	0.8	0.7	0.9	1.0	*
2002.....	4.2	3.0	1.8	1.2	1.0	0.9	0.8	1.1	2.1
2001.....	4.5	2.5	1.5	0.8	0.9	0.6	0.9	1.5	*
2000.....	4.0	2.4	1.6	0.9	0.8	0.8	1.0	*	1.7
1999.....	3.9	2.1	1.4	1.1	1.1	0.6	1.0	1.4	2.1

Non-Hispanic 4/, female

2007.....	3.2	3.0	1.8	1.2	0.8	0.8	0.6	0.8	1.1
2006.....	3.0	2.1	1.5	1.0	0.7	0.8	0.8	0.8	1.0
2005.....	2.9	2.5	1.2	1.2	0.8	0.5	0.6	0.8	1.4
2004.....	2.8	2.1	1.4	0.9	0.7	0.7	0.6	1.3	1.5
2003.....	2.7	1.9	1.2	0.9	0.8	0.6	*	0.9	1.0
2002.....	2.8	1.9	1.1	0.7	0.7	*	*	0.9	1.1
2001.....	1.8	1.5	0.7	0.5	*	*	0.6	0.8	0.8
2000.....	1.8	1.2	0.7	0.5	*	*	*	*	1.2
1999.....	1.8	1.1	0.7	0.6	0.5	*	0.5	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic White, both sexes

2007 5/.....	3.9	3.6	2.2	1.3	1.0	0.8	0.9	1.0	1.3
2006 5/.....	3.8	2.8	1.8	1.2	0.9	0.7	0.8	1.1	1.1
2005 5/.....	3.5	3.0	1.7	1.1	1.0	0.7	0.6	0.8	1.5
2004 5/.....	3.9	2.7	1.7	1.0	0.8	0.8	0.6	1.3	1.6
2003 5/.....	3.8	2.6	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	2.3	1.4	0.9	0.8	0.6	0.6	0.9	1.5
2001.....	3.1	1.9	1.1	0.6	0.5	0.5	0.7	1.1	1.0
2000.....	2.8	1.7	1.0	0.7	0.5	0.5	0.6	0.7	1.3
1999.....	2.6	1.4	1.0	0.8	0.8	0.4	0.7	1.0	1.4

Non-Hispanic White, male

2007 5/.....	4.4	3.9	2.5	1.6	1.2	0.9	1.2	1.4	1.7
2006 5/.....	4.3	3.5	2.0	1.3	1.1	0.7	0.8	1.7	1.9
2005 5/.....	4.0	3.4	2.1	1.0	1.1	1.0	*	*	1.8
2004 5/.....	4.8	3.3	2.0	1.0	1.0	0.9	*	1.4	1.9
2003 5/.....	4.7	3.2	1.8	1.0	0.7	*	0.9	*	*
2002.....	4.1	2.6	1.8	1.1	0.9	0.8	*	*	2.2
2001.....	4.3	2.3	1.4	0.7	0.8	*	1.0	1.5	*
2000.....	3.7	2.2	1.3	0.9	0.7	0.7	1.0	*	*
1999.....	3.4	1.8	1.2	1.0	1.1	*	1.0	1.3	2.1

Non-Hispanic White, female

2007 5/.....	3.4	3.3	1.8	1.1	0.8	0.8	0.6	0.8	1.1
2006 5/.....	3.2	2.2	1.5	1.0	0.8	0.8	0.8	0.8	0.7
2005 5/.....	3.1	2.6	1.4	1.2	0.9	0.5	0.6	0.8	1.4
2004 5/.....	3.1	2.2	1.5	1.0	0.6	0.7	0.6	1.3	1.5
2003 5/.....	2.9	2.0	1.3	0.8	0.8	0.5	*	0.8	1.1
2002.....	2.8	2.0	1.1	0.7	0.7	*	*	0.8	1.1
2001.....	1.9	1.6	0.8	0.6	*	*	0.6	0.8	0.8
2000.....	1.8	1.2	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.1	0.8	0.7	0.5	*	*	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2007.....	3.4	3.2	2.1	1.3	0.9	0.8	0.8	1.1	1.3
2006.....	3.4	2.7	1.8	1.1	0.9	0.8	0.9	1.1	1.4
2005.....	3.2	2.8	1.6	1.0	0.9	0.7	0.7	0.9	1.4
2004.....	3.5	2.5	1.6	0.9	0.8	0.7	0.6	1.2	1.6
2003.....	3.5	2.5	1.5	1.0	0.7	0.6	0.6	0.8	1.1
2002.....	3.2	2.2	1.3	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	2.9	1.9	1.0	0.6	0.5	0.4	0.6	1.0	1.0
2000.....	2.7	1.6	1.1	0.7	0.5	0.5	0.5	0.7	1.3
1999.....	2.6	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3

All origins 3/, male

2007.....	3.9	3.8	2.5	1.5	1.1	1.0	1.2	1.6	1.5
2006.....	4.2	3.5	2.2	1.3	1.1	0.8	1.2	1.7	2.2
2005.....	3.7	3.3	2.0	1.0	1.1	0.9	0.9	*	1.5
2004.....	4.5	3.1	1.9	1.0	1.0	0.8	0.7	1.1	1.8
2003.....	4.5	3.3	1.8	1.1	0.8	0.6	0.7	*	1.4
2002.....	4.0	2.8	1.7	1.3	0.9	0.8	0.7	1.0	2.2
2001.....	4.1	2.4	1.4	0.7	0.8	0.6	0.7	1.3	*
2000.....	3.7	2.1	1.5	0.9	0.7	0.7	0.8	*	*
1999.....	3.5	2.0	1.4	0.9	0.9	*	0.9	*	1.8

All origins 3/, female

2007.....	3.0	2.7	1.6	1.1	0.7	0.6	0.6	0.8	1.1
2006.....	2.7	2.0	1.4	0.9	0.6	0.7	0.8	0.8	1.0
2005.....	2.7	2.4	1.1	1.0	0.7	0.5	0.6	0.9	1.4
2004.....	2.6	2.0	1.3	0.8	0.6	0.6	0.6	1.2	1.5
2003.....	2.5	1.8	1.1	0.8	0.7	0.6	*	0.8	0.9
2002.....	2.5	1.7	0.9	0.7	0.6	0.4	0.5	0.8	1.1
2001.....	1.7	1.5	0.6	0.5	*	*	0.6	0.7	0.8
2000.....	1.6	1.1	0.7	0.4	*	*	*	*	1.2
1999.....	1.7	1.0	0.6	0.5	0.5	*	*	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 4/, both sexes

2007.....	3.7	3.4	2.1	1.4	0.9	0.8	0.8	1.1	1.3
2006.....	3.7	2.9	1.8	1.1	0.9	0.8	1.0	1.2	1.3
2005.....	3.4	2.9	1.6	1.0	0.9	0.7	0.7	0.8	1.5
2004.....	3.7	2.6	1.6	0.9	0.8	0.7	0.6	1.2	1.5
2003.....	3.7	2.6	1.5	1.0	0.8	0.6	0.6	0.8	1.1
2002.....	3.4	2.4	1.4	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	3.0	1.9	1.0	0.6	0.5	0.4	0.6	1.0	0.9
2000.....	2.8	1.7	1.0	0.6	0.5	0.5	0.5	0.7	1.3
1999.....	2.7	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3

Non-Hispanic 4/, male

2007.....	4.3	4.0	2.6	1.6	1.1	1.0	1.2	1.6	1.5
2006.....	4.4	3.7	2.2	1.3	1.1	0.9	1.2	1.8	2.2
2005.....	3.9	3.4	2.1	0.9	1.1	0.8	0.8	*	1.6
2004.....	4.7	3.3	1.9	1.0	1.0	0.8	0.7	1.0	1.7
2003.....	4.7	3.4	1.8	1.1	0.8	0.6	0.7	*	*
2002.....	4.1	2.9	1.8	1.2	0.9	0.8	0.7	1.0	2.1
2001.....	4.4	2.4	1.4	0.7	0.8	*	0.7	1.3	*
2000.....	3.9	2.2	1.4	0.9	0.7	0.7	0.9	*	*
1999.....	3.7	2.1	1.4	0.9	0.9	*	0.8	*	1.7

Non-Hispanic 4/, female

2007.....	3.2	2.9	1.7	1.2	0.8	0.7	0.5	0.8	1.1
2006.....	2.9	2.1	1.5	0.9	0.7	0.8	0.8	0.8	1.0
2005.....	2.9	2.5	1.2	1.1	0.8	0.5	0.6	0.8	1.4
2004.....	2.8	2.0	1.3	0.8	0.6	0.7	0.6	1.2	1.5
2003.....	2.7	1.9	1.2	0.8	0.7	0.6	*	0.8	1.0
2002.....	2.7	1.9	1.0	0.7	0.6	*	*	0.9	1.1
2001.....	1.8	1.5	0.7	0.5	*	*	0.6	0.8	0.8
2000.....	1.7	1.2	0.7	0.4	*	*	*	*	1.2
1999.....	1.8	1.0	0.6	0.6	0.5	*	*	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic White, both sexes

2007 5/.....	3.8	3.5	2.1	1.2	0.9	0.8	0.8	1.0	1.3
2006 5/.....	3.7	2.8	1.7	1.1	0.9	0.7	0.8	1.1	1.1
2005 5/.....	3.5	2.9	1.6	1.0	0.9	0.6	0.6	0.8	1.5
2004 5/.....	3.9	2.6	1.7	0.9	0.7	0.7	0.6	1.2	1.6
2003 5/.....	3.7	2.5	1.4	0.9	0.7	0.5	0.5	0.8	1.2
2002.....	3.4	2.2	1.4	0.9	0.7	0.5	0.5	0.9	1.5
2001.....	3.0	1.8	1.0	0.6	0.5	0.4	0.6	1.0	0.9
2000.....	2.7	1.6	0.9	0.6	0.4	0.5	0.5	0.7	1.2
1999.....	2.6	1.4	0.9	0.7	0.7	0.4	0.6	0.8	1.3

Non-Hispanic White, male

2007 5/.....	4.2	3.8	2.4	1.4	1.0	0.8	1.1	1.3	1.7
2006 5/.....	4.2	3.4	1.9	1.2	1.0	0.7	0.8	1.6	1.9
2005 5/.....	3.9	3.2	2.0	0.8	1.1	0.8	*	*	1.7
2004 5/.....	4.7	3.1	1.9	0.9	0.8	0.8	*	1.1	1.8
2003 5/.....	4.6	3.1	1.6	0.9	0.6	*	*	*	*
2002.....	3.9	2.5	1.7	1.1	0.8	0.6	*	*	2.2
2001.....	4.1	2.2	1.3	0.6	0.6	*	*	1.4	*
2000.....	3.6	2.0	1.2	0.8	0.6	0.6	0.9	*	*
1999.....	3.2	1.7	1.2	0.8	0.9	*	0.8	*	*

Non-Hispanic White, female

2007 5/.....	3.4	3.2	1.8	1.1	0.8	0.7	*	0.8	1.1
2006 5/.....	3.2	2.2	1.5	1.0	0.8	0.7	0.8	0.8	0.7
2005 5/.....	3.1	2.6	1.3	1.2	0.8	0.5	0.6	0.8	1.4
2004 5/.....	3.1	2.1	1.4	0.9	0.6	0.6	0.6	1.3	1.5
2003 5/.....	2.9	2.0	1.3	0.8	0.7	0.5	*	0.8	1.1
2002.....	2.8	2.0	1.1	0.7	0.6	*	*	0.7	1.1
2001.....	1.9	1.5	0.8	0.6	*	*	0.6	*	0.8
2000.....	1.8	1.2	0.7	*	*	*	*	*	1.2
1999.....	1.9	1.0	0.7	0.7	0.5	*	*	0.7	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/, both sexes

2007.....	0.5	0.7	1.0	1.5	2.2	2.9	5.1	6.0	7.9
2006.....	0.5	0.8	0.9	1.3	2.0	3.1	4.7	6.1	8.3
2005.....	0.5	0.8	1.0	1.4	2.0	3.3	4.7	7.2	9.1
2004.....	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
2003.....	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7

All origins 3/, male

2007.....	0.5	0.7	1.1	1.5	2.4	3.2	5.7	7.3	9.8
2006.....	0.5	0.9	1.0	1.2	2.0	2.9	5.3	7.1	8.8
2005.....	0.5	0.8	1.1	1.7	2.3	3.7	4.6	8.9	10.8
2004.....	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
2003.....	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5

All origins 3/, female

2007.....	0.4	0.7	0.9	1.5	2.0	2.7	4.6	5.2	6.9
2006.....	0.5	0.7	0.9	1.4	2.0	3.3	4.2	5.4	8.1
2005.....	0.6	0.9	1.0	1.2	1.8	3.0	4.8	6.2	8.3
2004.....	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
2003.....	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 4/, both sexes

2007.....	0.5	0.7	1.0	1.5	2.3	3.0	5.2	6.1	8.0
2006.....	0.5	0.8	1.0	1.4	2.0	3.1	4.9	6.1	8.6
2005.....	0.5	0.8	1.1	1.5	2.1	3.3	4.7	7.3	9.2
2004.....	0.6	0.8	1.3	1.6	2.6	3.6	5.0	7.6	10.8
2003.....	0.7	0.7	1.2	1.7	2.2	3.7	5.5	7.4	11.4
2002.....	0.6	0.8	1.1	1.6	2.3	3.9	5.2	7.7	11.6
2001.....	0.7	0.8	1.3	1.9	2.6	3.7	5.9	8.3	12.9
2000.....	0.7	1.0	1.2	2.1	2.7	4.4	5.8	8.1	13.7
1999.....	0.5	0.7	1.1	1.7	2.5	4.2	5.6	8.6	12.8

Non-Hispanic 4/, male

2007.....	0.5	0.7	1.2	1.5	2.5	3.3	5.9	7.4	10.1
2006.....	0.5	0.9	1.0	1.3	2.0	2.9	5.4	7.2	9.1
2005.....	0.5	0.9	1.1	1.7	2.4	3.7	4.6	9.0	11.2
2004.....	0.6	0.9	1.5	1.7	2.5	4.4	6.0	9.5	12.5
2003.....	0.7	0.7	1.4	1.7	2.3	4.2	5.7	9.0	13.2
2002.....	0.6	0.7	1.2	1.6	2.5	3.9	5.4	7.8	12.9
2001.....	0.7	0.8	1.4	2.0	2.6	4.0	6.3	9.9	14.5
2000.....	0.6	1.0	1.2	2.1	2.8	5.3	6.9	10.4	16.1
1999.....	0.4	0.7	1.2	1.7	2.4	4.8	6.5	10.3	16.8

Non-Hispanic 4/, female

2007.....	0.5	0.8	0.9	1.5	2.1	2.7	4.8	5.2	7.1
2006.....	0.6	0.7	0.9	1.5	2.0	3.3	4.4	5.4	8.3
2005.....	0.6	0.8	1.0	1.3	1.8	2.9	4.8	6.2	8.4
2004.....	0.6	0.8	1.0	1.5	2.8	2.9	4.3	6.5	10.1
2003.....	0.6	0.7	1.0	1.7	2.1	3.3	5.3	6.4	10.6
2002.....	0.5	0.9	1.0	1.7	2.2	4.0	5.0	7.7	11.1
2001.....	0.6	0.8	1.2	1.8	2.5	3.5	5.6	7.4	12.2
2000.....	0.7	0.9	1.2	2.1	2.6	3.7	5.0	6.8	12.7
1999.....	0.6	0.8	0.9	1.6	2.5	3.7	5.0	7.6	11.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic White, both sexes

2007 5/.....	0.4	0.6	1.0	1.5	2.1	2.8	4.9	6.0	7.7
2006 5/.....	0.5	0.7	0.8	1.3	1.9	3.1	4.6	5.9	8.5
2005 5/.....	0.5	0.7	0.9	1.3	1.8	2.9	4.4	7.3	9.4
2004 5/.....	0.5	0.8	1.2	1.5	2.5	3.4	4.8	7.4	10.6
2003 5/.....	0.6	0.6	1.0	1.6	2.1	3.5	5.2	7.5	11.2
2002.....	0.5	0.7	1.0	1.5	2.1	3.9	5.0	7.6	11.5
2001.....	0.6	0.7	1.2	1.7	2.4	3.3	5.8	8.2	12.2
2000.....	0.6	0.9	1.1	1.8	2.5	4.1	5.5	8.0	13.2
1999.....	0.4	0.6	0.9	1.6	2.4	3.9	5.3	8.2	12.1

Non-Hispanic White, male

2007 5/.....	0.4	0.6	1.0	1.5	2.3	3.1	5.6	7.6	9.8
2006 5/.....	0.4	0.9	0.9	1.2	1.9	2.8	5.3	7.3	9.1
2005 5/.....	0.4	0.7	1.0	1.5	2.3	3.5	4.4	9.0	11.6
2004 5/.....	0.6	0.8	1.4	1.7	2.5	4.3	6.0	9.4	12.3
2003 5/.....	0.6	0.6	1.2	1.6	2.3	4.1	5.6	9.3	13.1
2002.....	0.5	0.6	1.1	1.5	2.2	3.8	5.3	7.9	12.4
2001.....	0.6	0.7	1.2	1.7	2.5	3.3	6.2	10.1	13.5
2000.....	0.6	1.0	1.2	1.8	2.5	5.0	6.8	10.6	15.9
1999.....	0.3	0.6	1.0	1.6	2.3	4.5	6.3	9.9	15.8

Non-Hispanic White, female

2007 5/.....	0.4	0.7	0.9	1.4	1.9	2.6	4.4	5.0	6.8
2006 5/.....	0.5	0.5	0.7	1.3	1.8	3.3	4.2	5.0	8.2
2005 5/.....	0.5	0.6	0.9	1.2	1.4	2.5	4.4	6.3	8.3
2004 5/.....	0.5	0.7	1.0	1.4	2.5	2.7	4.0	6.2	9.9
2003 5/.....	0.6	0.6	0.8	1.5	2.0	3.0	4.9	6.3	10.4
2002.....	0.4	0.7	0.9	1.5	2.1	4.0	4.9	7.4	11.1
2001.....	0.5	0.7	1.2	1.6	2.3	3.2	5.5	7.1	11.6
2000.....	0.6	0.9	1.0	1.8	2.5	3.5	4.5	6.5	12.0
1999.....	0.5	0.6	0.8	1.5	2.5	3.5	4.7	7.2	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic Black, both sexes

2007 5/.....	0.9	1.6	1.6	2.8	4.2	5.3	9.4	8.4	12.5
2006 5/.....	1.0	1.7	2.4	3.2	4.0	4.3	8.4	9.7	10.4
2005 5/.....	1.0	2.2	2.3	2.9	4.6	6.6	9.4	8.4	9.3
2004 5/.....	1.1	1.5	2.7	2.5	4.7	5.6	7.6	12.2	14.1
2003 5/.....	1.3	1.6	2.5	3.0	3.7	5.7	9.7	7.8	15.2
2002.....	1.3	2.2	2.6	2.9	4.1	5.4	7.7	9.6	14.2
2001.....	1.6	1.6	2.8	4.2	4.6	8.5	8.1	11.0	23.7
2000.....	1.2	1.7	2.7	4.8	4.8	8.0	10.5	10.6	21.3
1999.....	0.9	1.8	2.7	2.6	3.5	7.3	9.4	12.3	20.4

Non-Hispanic Black, male

2007 5/.....	*	*	2.4	*	*	7.2	11.4	*	*
2006 5/.....	*	*	*	*	*	*	9.7	*	*
2005 5/.....	*	2.3	2.7	3.7	*	*	9.2	*	*
2004 5/.....	*	*	3.4	*	*	*	*	*	*
2003 5/.....	*	2.2	2.9	*	*	*	*	*	*
2002.....	*	*	*	*	5.1	*	*	*	*
2001.....	1.8	*	3.5	5.2	5.2	11.4	*	*	30.0
2000.....	*	*	*	4.7	6.4	9.6	10.1	*	*
1999.....	*	*	*	*	*	9.3	10.2	*	23.8

Non-Hispanic Black, female

2007 5/.....	*	1.9	*	2.9	4.2	*	8.2	8.4	10.9
2006 5/.....	*	1.9	2.6	3.0	4.3	*	7.7	9.6	10.1
2005 5/.....	*	2.1	*	*	4.9	6.9	9.6	7.6	9.2
2004 5/.....	*	1.7	*	*	5.5	5.2	7.5	11.2	13.7
2003 5/.....	*	*	*	3.1	3.9	6.4	10.8	7.9	15.3
2002.....	*	2.5	2.5	3.2	*	5.0	8.4	10.1	11.7
2001.....	*	*	*	3.5	4.2	6.7	8.2	11.2	21.4
2000.....	*	*	3.2	4.9	*	7.0	10.7	10.8	22.2
1999.....	*	*	*	*	*	6.0	9.0	11.0	19.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

All origins 3/, both sexes

2007.....	16.2	20.5	20.8	19.4	16.8	13.8	11.3	7.6	4.6
2006.....	15.7	19.7	19.9	18.6	16.0	13.2	11.0	8.1	4.9
2005.....	15.7	19.7	19.7	19.3	16.3	13.7	10.1	8.1	4.6
2004.....	15.5	19.5	18.9	18.5	17.0	14.1	10.7	7.3	4.3
2003.....	16.0	18.9	18.3	18.8	16.8	13.2	10.3	7.6	4.1
2002.....	15.6	18.2	18.2	18.6	16.5	14.4	10.9	7.2	4.4
2001.....	16.5	17.9	17.6	18.9	17.3	13.7	11.0	7.5	5.0
2000.....	15.9	16.7	18.2	19.3	17.2	14.3	11.2	7.9	5.4
1999.....	16.0	16.9	18.1	19.3	17.6	14.1	11.9	8.5	5.5

All origins 3/, male

2007.....	23.5	31.6	33.4	31.5	28.2	24.2	20.4	14.1	9.6
2006.....	22.8	30.6	31.8	30.1	27.1	22.2	19.1	14.1	9.5
2005.....	22.9	30.9	31.7	31.3	27.0	22.6	18.4	14.9	9.3
2004.....	22.7	30.6	30.0	30.4	28.1	24.5	19.4	13.8	9.0
2003.....	24.5	29.9	29.1	30.6	27.6	22.8	18.0	15.0	7.8
2002.....	23.7	29.2	29.2	31.0	27.9	23.8	18.7	13.4	9.7
2001.....	25.4	28.0	28.7	31.0	28.5	23.8	19.4	13.8	11.6
2000.....	25.0	26.9	28.8	31.4	28.9	25.0	21.1	14.9	11.9
1999.....	25.2	27.4	29.7	31.5	29.9	24.5	21.2	16.6	11.8

All origins 3/, female

2007.....	9.1	9.9	9.0	8.2	6.8	5.4	4.6	3.5	2.2
2006.....	8.7	9.1	8.7	8.2	6.4	5.9	5.0	4.3	2.7
2005.....	8.6	8.9	8.4	8.4	7.1	6.5	4.0	3.8	2.5
2004.....	8.4	8.8	8.3	7.8	7.3	5.8	4.4	3.3	2.3
2003.....	7.7	8.3	8.1	8.1	7.5	5.4	4.9	3.1	2.5
2002.....	7.7	7.7	7.9	7.2	6.7	6.9	5.3	3.4	2.1
2001.....	7.9	8.3	7.2	8.0	7.6	5.7	5.1	3.8	2.3
2000.....	7.1	7.0	8.3	8.3	7.2	5.9	4.3	3.8	2.7
1999.....	7.1	6.8	7.3	8.3	7.0	6.0	5.5	3.8	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Hispanic, both sexes

2007	18.1	25.2	25.2	27.4	25.0	24.0	21.3	12.1	8.9
2006	17.6	23.6	26.5	25.9	23.0	19.9	20.5	16.7	9.4
2005	17.7	23.0	24.9	25.2	26.8	24.7	16.5	11.2	*
2004	18.6	23.2	22.1	23.1	21.8	19.3	20.3	12.0	*
2003	18.3	22.8	27.7	27.4	26.1	19.1	19.7	14.8	*
2002	18.7	24.5	26.9	25.7	28.0	25.8	16.3	11.6	*
2001	21.5	23.9	27.3	29.5	27.7	24.0	22.4	16.9	*
2000	20.6	24.5	30.0	28.8	32.5	24.3	22.0	13.4	*
1999	21.3	24.2	28.0	29.6	28.5	23.2	20.7	15.3	14.0

Hispanic, male

2007	29.9	42.6	44.7	49.1	46.7	47.7	41.1	20.9	23.4
2006	28.6	40.0	46.2	45.8	42.3	40.1	39.1	27.0	*
2005	29.5	39.4	43.6	47.2	49.6	50.1	32.4	25.4	*
2004	31.9	41.5	37.4	42.1	38.9	36.0	40.9	20.8	*
2003	31.3	40.5	50.3	50.7	44.2	35.4	37.7	30.1	*
2002	31.7	42.4	49.4	47.5	54.5	50.5	28.8	22.9	*
2001	36.9	41.2	49.7	54.6	55.6	48.0	41.6	31.2	*
2000	35.9	43.4	55.9	53.3	66.0	49.1	48.0	30.9	*
1999	36.9	42.6	51.3	54.8	54.1	47.4	37.4	33.0	*

Hispanic, female

2007	5.7	7.8	6.7	7.9	6.6	*	*	*	*
2006	6.1	7.3	7.9	8.2	6.9	*	*	*	*
2005	5.4	6.8	7.3	5.8	7.9	*	*	*	*
2004	4.8	5.3	7.9	6.5	7.7	6.4	*	*	*
2003	4.9	5.6	6.8	7.1	11.2	6.6	*	*	*
2002	5.4	7.1	6.2	6.7	6.4	6.8	*	*	*
2001	6.1	7.4	7.1	7.8	*	*	*	*	*
2000	5.3	6.5	6.5	7.7	*	*	*	*	*
1999	5.8	6.7	6.9	8.0	7.8	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic 4/, both sexes

2007	15.9	19.9	20.3	18.6	16.0	13.0	10.5	7.3	4.4
2006	15.4	19.2	19.3	18.0	15.4	12.6	10.3	7.6	4.6
2005	15.4	19.2	19.1	18.7	15.4	12.8	9.7	7.9	4.4
2004	15.0	19.0	18.5	18.1	16.5	13.7	10.1	7.1	4.4
2003	15.6	18.4	17.4	18.0	16.0	12.7	9.8	7.3	4.1
2002	15.1	17.5	17.4	17.9	15.5	13.6	10.5	7.0	4.1
2001	15.9	17.2	16.6	18.0	16.4	13.0	10.4	7.2	5.0
2000	15.3	15.9	17.2	18.5	16.1	13.7	10.6	7.7	5.3
1999	15.4	16.1	17.3	18.4	16.8	13.5	11.5	8.3	5.2

Non-Hispanic 4/, male

2007	22.5	30.2	32.1	29.9	26.5	22.4	18.9	13.7	8.8
2006	21.9	29.5	30.4	28.6	25.6	20.8	17.7	13.4	8.8
2005	22.0	29.8	30.5	29.9	25.1	20.4	17.5	14.3	8.6
2004	21.4	29.3	29.2	29.3	27.2	23.6	18.1	13.5	8.9
2003	23.6	28.7	27.2	28.7	26.2	21.8	16.9	14.2	7.8
2002	22.5	27.8	27.3	29.5	25.8	22.0	18.1	12.9	9.1
2001	24.0	26.5	26.7	29.0	26.4	22.2	18.1	13.1	11.4
2000	23.7	25.2	26.6	29.6	26.2	23.6	19.6	14.3	11.7
1999	23.9	25.9	28.0	29.5	28.2	23.0	20.3	16.0	11.1

Non-Hispanic 4/, female

2007	9.5	10.0	9.1	8.2	6.8	5.4	4.4	3.3	2.3
2006	9.0	9.3	8.8	8.2	6.3	6.0	4.9	4.0	2.7
2005	9.0	9.1	8.4	8.5	7.0	6.6	4.0	3.9	2.5
2004	8.7	9.1	8.3	7.9	7.2	5.8	4.4	3.2	2.3
2003	7.9	8.5	8.2	8.2	7.2	5.3	4.8	3.1	2.5
2002	7.9	7.7	8.0	7.3	6.7	6.9	5.2	3.4	1.9
2001	8.0	8.3	7.1	8.0	7.7	5.6	4.9	3.6	2.3
2000	7.2	7.0	8.5	8.3	7.3	5.9	4.4	3.8	2.6
1999	7.2	6.7	7.3	8.3	6.9	6.0	5.4	3.7	2.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic White, both sexes

2007 5/.....	16.5	20.5	20.7	18.6	16.2	13.0	10.6	7.5	4.3
2006 5/.....	16.0	19.7	19.3	18.2	15.3	12.7	10.6	7.8	4.7
2005 5/.....	15.7	19.4	19.5	18.3	15.5	12.7	9.7	7.8	4.4
2004 5/.....	15.4	19.1	18.2	17.6	16.0	13.5	10.2	7.2	4.5
2003 5/.....	15.8	18.0	17.1	17.2	16.0	12.7	10.0	7.2	4.1
2002.....	15.2	17.2	16.6	17.1	15.4	13.3	10.5	6.8	4.0
2001.....	15.7	16.1	15.5	17.5	15.6	12.9	10.2	7.2	4.7
2000.....	14.7	15.1	16.1	17.6	15.5	13.4	10.5	7.6	5.2
1999.....	14.4	15.2	16.0	17.3	16.0	12.9	11.0	8.1	5.0

Non-Hispanic White, male

2007 5/.....	23.3	30.7	32.4	29.6	26.3	21.7	18.4	14.0	8.8
2006 5/.....	22.7	29.9	29.9	28.7	25.2	20.5	18.2	13.7	8.9
2005 5/.....	22.1	29.8	30.7	29.0	24.7	19.8	17.2	14.1	8.7
2004 5/.....	21.7	29.4	28.4	28.1	25.9	22.7	18.0	13.7	9.3
2003 5/.....	23.5	27.8	26.5	26.9	26.0	21.6	17.0	13.7	7.7
2002.....	22.5	27.3	26.1	28.0	24.9	20.9	18.0	12.5	8.7
2001.....	23.7	24.8	24.5	27.8	24.5	21.7	17.8	13.2	10.8
2000.....	22.9	23.8	24.9	28.0	24.9	22.8	19.1	14.2	11.4
1999.....	22.4	24.4	25.4	27.9	26.3	21.7	19.3	15.8	10.5

Non-Hispanic White, female

2007 5/.....	9.9	10.5	9.3	8.2	7.1	5.7	4.7	3.4	2.2
2006 5/.....	9.4	9.6	9.0	8.4	6.5	6.2	5.0	4.1	2.7
2005 5/.....	9.3	9.2	8.6	8.4	7.2	6.7	4.1	3.9	2.5
2004 5/.....	9.1	9.1	8.2	7.7	7.2	5.9	4.5	3.2	2.4
2003 5/.....	8.2	8.5	8.0	8.1	7.1	5.3	4.9	3.2	2.5
2002.....	7.9	7.4	7.6	6.9	6.9	7.1	5.2	3.3	2.1
2001.....	7.8	7.5	6.9	7.9	7.7	5.8	4.9	3.7	2.2
2000.....	6.6	6.5	7.7	8.0	7.1	5.9	4.4	3.7	2.7
1999.....	6.5	6.2	6.9	7.5	6.9	5.9	5.2	3.5	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

Non-Hispanic Black, both sexes

2007 5/.....	12.8	17.7	21.0	21.6	18.4	15.6	11.3	6.0	5.6
2006 5/.....	12.1	18.3	21.5	19.6	17.5	13.5	9.3	6.5	*
2005 5/.....	15.1	20.0	20.6	23.9	17.2	16.0	10.8	8.7	5.5
2004 5/.....	14.3	21.1	23.7	24.3	22.7	17.3	8.8	6.3	*
2003 5/.....	15.3	22.8	22.7	26.9	19.2	14.5	9.2	8.8	*
2002.....	16.3	22.2	25.6	25.3	18.7	18.0	12.3	9.1	*
2001.....	18.6	27.8	27.2	24.8	25.5	14.7	13.6	7.4	*
2000.....	21.9	24.6	27.1	26.6	24.0	17.9	13.6	8.3	6.4
1999.....	23.1	24.4	29.8	29.3	25.3	21.1	16.7	11.7	7.8

Non-Hispanic Black, male

2007 5/.....	18.7	30.1	35.9	39.2	36.4	32.7	26.9	*	*
2006 5/.....	18.2	29.6	37.7	34.5	33.6	25.6	18.0	*	*
2005 5/.....	23.8	32.4	34.0	42.0	31.6	30.2	24.3	15.8	*
2004 5/.....	22.3	33.9	41.6	43.1	43.4	35.6	17.7	*	*
2003 5/.....	25.2	39.5	37.5	49.2	33.2	28.0	17.8	21.9	*
2002.....	25.1	36.0	42.3	45.4	36.3	35.6	23.0	19.5	*
2001.....	29.5	44.5	48.6	44.5	48.7	29.6	26.5	*	*
2000.....	33.5	40.5	44.2	45.3	43.2	36.6	28.4	17.2	*
1999.....	36.1	40.5	53.7	49.0	49.1	41.2	31.9	23.0	*

Non-Hispanic Black, female

2007 5/.....	7.7	7.3	8.8	7.9	5.2	*	*	*	*
2006 5/.....	6.8	8.8	8.1	8.1	5.7	5.2	*	*	*
2005 5/.....	7.5	9.5	9.5	9.9	6.5	6.2	*	*	*
2004 5/.....	7.5	10.3	9.1	9.6	7.4	5.0	*	*	*
2003 5/.....	6.8	8.6	10.6	9.4	8.8	5.5	*	*	*
2002.....	8.7	10.4	11.8	9.5	5.7	6.2	5.9	*	*
2001.....	9.2	13.6	9.6	9.4	8.3	4.8	5.9	*	*
2000.....	11.8	11.0	13.0	11.9	9.8	5.4	*	*	*
1999.....	11.9	10.9	10.4	13.8	7.7	7.8	7.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

All origins 3/, both sexes

2007	28.0	25.7	17.0	9.3	5.3	3.9	3.8	4.0	4.8
2006	28.4	24.8	15.2	8.1	5.7	4.9	5.5	6.8	8.1
2005	25.8	21.2	12.8	7.8	5.0	4.4	4.9	6.1	7.6
2004	24.4	18.5	11.0	6.6	4.4	4.1	4.7	4.9	6.3
2003	22.7	17.0	9.8	5.7	4.5	3.7	3.9	4.6	6.0
2002	20.9	14.5	8.2	5.1	3.8	3.3	3.6	4.1	5.7
2001	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.6	5.1
2000	15.8	10.3	5.9	3.6	2.6	2.7	3.2	3.9	5.7
1999	15.5	9.3	5.8	3.8	3.2	2.9	3.6	4.2	4.8

All origins 3/, male

2007	32.9	30.4	20.5	10.5	6.0	4.4	3.4	4.4	5.5
2006	35.0	31.3	18.7	9.5	6.0	5.5	6.2	8.0	10.7
2005	31.9	26.6	16.0	9.0	5.4	5.2	6.6	6.9	9.9
2004	30.1	23.6	13.0	7.6	4.8	4.5	5.2	5.6	8.6
2003	29.2	21.7	11.5	6.6	5.2	3.9	4.7	5.3	7.8
2002	26.4	19.1	10.0	5.7	4.1	3.4	3.9	4.7	7.3
2001	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.9	5.7
2000	21.9	13.8	6.7	3.9	2.9	2.8	3.6	4.3	7.5
1999	21.9	12.2	7.1	4.7	3.7	3.1	3.5	4.4	6.2

All origins 3/, female

2007	23.2	21.1	13.7	8.3	4.6	3.5	4.0	3.8	4.4
2006	21.9	18.6	12.0	6.9	5.4	4.3	5.0	6.0	6.8
2005	19.8	16.1	9.8	6.6	4.8	3.8	3.7	5.5	6.6
2004	19.0	13.7	9.1	5.7	4.1	3.7	4.4	4.4	5.2
2003	16.5	12.4	8.2	4.9	4.0	3.4	3.3	4.2	5.2
2002	15.6	10.2	6.6	4.6	3.6	3.3	3.3	3.8	4.9
2001	11.6	8.3	5.1	3.9	2.6	2.6	3.2	3.4	4.9
2000	9.8	7.0	5.2	3.4	2.3	2.7	2.9	3.8	4.9
1999	9.2	6.6	4.6	2.9	2.7	2.7	3.7	4.0	4.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Hispanic, both sexes

2007	15.2	15.0	10.7	6.2	3.2	*	*	*	*
2006	17.2	15.2	11.1	6.0	3.2	*	*	*	*
2005	15.4	14.7	9.0	6.2	4.2	*	*	*	*
2004	14.0	11.9	10.6	5.1	3.4	*	*	*	*
2003	16.4	13.0	7.6	4.9	3.8	*	*	*	*
2002	15.9	11.1	6.5	4.4	*	*	*	*	*
2001	12.0	9.6	6.3	2.8	*	*	*	*	*
2000	12.1	8.6	6.4	*	*	*	*	*	*
1999	15.1	11.2	5.0	5.2	*	*	*	*	*

Hispanic, male

2007	22.5	22.0	14.4	9.6	5.9	*	*	*	*
2006	23.8	22.3	17.7	9.6	*	*	*	*	*
2005	22.6	21.7	14.7	10.5	*	*	*	*	*
2004	19.2	17.0	15.4	8.1	*	*	*	*	*
2003	24.3	19.1	12.3	7.9	*	*	*	*	*
2002	23.6	17.7	9.9	6.9	*	*	*	*	*
2001	18.5	15.7	10.6	*	*	*	*	*	*
2000	20.1	13.1	9.0	*	*	*	*	*	*
1999	25.9	16.5	9.2	8.6	*	*	*	*	*

Hispanic, female

2007	7.5	8.0	7.2	*	*	*	*	*	*
2006	10.2	8.0	4.9	*	*	*	*	*	*
2005	7.9	7.8	3.6	*	*	*	*	*	*
2004	8.6	6.8	6.1	*	*	*	*	*	*
2003	8.3	7.1	3.2	*	*	*	*	*	*
2002	8.1	4.6	3.4	*	*	*	*	*	*
2001	5.5	3.8	*	*	*	*	*	*	*
2000	4.2	4.3	4.0	*	*	*	*	*	*
1999	4.4	6.2	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic 4/, both sexes

2007	29.6	26.7	17.5	9.6	5.4	4.0	3.9	4.2	4.9
2006	29.6	25.7	15.6	8.2	5.9	5.0	5.6	6.9	8.2
2005	26.9	21.7	13.1	7.9	5.1	4.5	5.0	6.1	7.7
2004	25.5	19.1	11.0	6.7	4.5	4.2	4.7	4.9	6.3
2003	23.3	17.2	9.9	5.7	4.5	3.7	3.9	4.6	6.0
2002	21.3	14.8	8.3	5.1	3.8	3.3	3.5	4.2	5.8
2001	18.0	11.6	6.2	4.2	3.1	2.9	3.5	3.6	5.2
2000	15.9	10.3	5.8	3.7	2.6	2.7	3.2	4.0	5.7
1999	15.3	9.0	5.7	3.6	3.1	2.9	3.7	4.2	4.9

Non-Hispanic 4/, male

2007	34.2	31.2	20.9	10.5	6.0	4.3	3.4	4.6	5.7
2006	36.3	32.1	18.7	9.3	6.1	5.6	6.3	8.2	10.8
2005	32.9	26.8	16.0	8.8	5.4	5.2	6.7	7.0	10.0
2004	31.2	24.0	12.8	7.5	4.7	4.5	5.1	5.7	8.6
2003	29.5	21.7	11.3	6.4	5.0	3.8	4.7	5.1	7.8
2002	26.5	19.1	9.9	5.6	4.0	3.3	3.7	4.7	7.5
2001	24.1	14.8	7.3	4.2	3.6	3.3	3.8	4.0	5.8
2000	21.8	13.6	6.3	3.8	2.9	2.8	3.7	4.4	7.5
1999	21.3	11.6	6.8	4.2	3.5	3.1	3.6	4.6	6.3

Non-Hispanic 4/, female

2007	25.1	22.5	14.3	8.7	4.9	3.7	4.2	3.9	4.6
2006	23.2	19.6	12.6	7.3	5.7	4.5	5.1	6.0	7.0
2005	21.1	16.9	10.3	7.0	4.9	3.9	3.8	5.6	6.7
2004	20.0	14.4	9.4	6.0	4.2	3.9	4.4	4.5	5.3
2003	17.2	12.9	8.6	5.1	4.1	3.6	3.3	4.2	5.3
2002	16.3	10.6	6.9	4.7	3.6	3.3	3.4	3.8	5.1
2001	12.1	8.6	5.2	4.1	2.6	2.6	3.3	3.4	5.0
2000	10.2	7.1	5.2	3.6	2.4	2.7	2.9	3.8	5.0
1999	9.6	6.6	4.7	3.0	2.8	2.7	3.8	4.0	4.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic White, both sexes

2007 5/.....	31.5	27.1	17.1	9.5	5.4	4.2	4.0	4.3	5.2
2006 5/.....	30.4	25.2	14.9	7.7	5.8	4.8	5.8	7.1	8.4
2005 5/.....	27.8	21.0	12.5	7.9	5.1	4.6	5.2	6.3	8.1
2004 5/.....	26.2	18.7	10.5	6.5	4.4	4.4	4.9	5.1	6.4
2003 5/.....	23.6	16.7	9.7	5.6	4.4	3.9	4.2	4.8	6.3
2002.....	21.2	14.0	8.0	4.9	3.9	3.4	3.6	4.4	5.9
2001.....	17.3	10.8	6.0	4.0	3.1	3.0	3.6	3.8	5.2
2000.....	15.0	9.3	5.5	3.7	2.6	2.9	3.5	4.2	5.7
1999.....	14.2	8.1	5.4	3.3	3.1	2.9	3.9	4.4	4.9

Non-Hispanic White, male

2007 5/.....	35.8	30.4	19.1	9.9	5.3	4.5	3.5	4.8	6.0
2006 5/.....	36.2	30.1	16.6	7.8	5.5	4.9	6.4	8.7	11.3
2005 5/.....	33.1	24.8	14.3	8.1	4.9	5.2	6.9	7.0	10.7
2004 5/.....	31.6	22.5	11.2	6.7	4.1	4.6	5.1	5.8	8.7
2003 5/.....	29.4	20.2	10.4	5.9	4.5	3.9	5.0	5.4	7.8
2002.....	25.9	17.2	8.7	4.7	3.7	3.4	3.8	4.6	7.8
2001.....	22.4	12.8	6.5	3.4	3.5	3.2	3.6	4.2	5.7
2000.....	20.0	11.7	5.5	3.5	2.6	2.7	3.9	4.5	7.2
1999.....	19.1	9.8	5.7	3.5	3.3	3.2	3.6	4.8	6.6

Non-Hispanic White, female

2007 5/.....	27.3	23.8	15.1	9.1	5.5	3.9	4.4	4.0	4.8
2006 5/.....	24.7	20.4	13.3	7.7	6.1	4.8	5.4	6.1	7.1
2005 5/.....	22.6	17.2	10.8	7.6	5.3	4.2	3.9	5.9	7.0
2004 5/.....	20.7	15.1	9.9	6.4	4.7	4.2	4.8	4.6	5.4
2003 5/.....	17.9	13.2	9.0	5.4	4.4	3.8	3.7	4.5	5.7
2002.....	16.6	10.9	7.3	5.1	4.0	3.5	3.5	4.2	5.1
2001.....	12.2	8.8	5.6	4.5	2.7	2.8	3.6	3.5	4.9
2000.....	10.1	7.0	5.5	3.9	2.5	3.0	3.1	4.0	5.1
1999.....	9.4	6.6	5.0	3.2	3.0	2.8	4.1	4.2	4.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

Non-Hispanic Black, both sexes

2007 5/.....	28.2	34.6	26.2	13.2	8.1	3.6	3.9	*	*
2006 5/.....	34.4	37.6	25.8	15.1	7.6	6.4	4.5	*	6.6
2005 5/.....	29.9	33.4	22.7	10.5	5.6	4.6	4.3	*	*
2004 5/.....	29.7	27.4	18.4	9.7	6.1	3.1	3.5	*	*
2003 5/.....	28.9	25.8	14.8	7.6	6.5	3.0	*	*	*
2002.....	28.9	25.3	13.9	8.9	3.6	*	*	*	*
2001.....	28.6	21.9	10.1	6.9	3.4	2.9	*	*	6.2
2000.....	26.5	21.2	9.8	4.6	3.6	*	*	*	7.6
1999.....	28.1	18.4	9.6	6.4	3.8	2.9	*	*	*

Non-Hispanic Black, male

2007 5/.....	37.1	49.4	41.7	20.6	15.6	*	*	*	*
2006 5/.....	48.4	57.9	42.1	26.1	13.5	12.0	*	*	*
2005 5/.....	41.9	49.3	36.6	19.3	10.0	7.5	*	*	*
2004 5/.....	37.8	42.5	30.4	16.6	11.2	*	*	*	*
2003 5/.....	40.3	38.9	23.4	12.9	11.6	*	*	*	*
2002.....	39.7	40.5	23.3	16.1	6.7	*	*	*	*
2001.....	44.1	35.1	16.9	12.1	5.2	*	*	*	*
2000.....	41.5	33.9	16.3	7.4	5.9	*	*	*	*
1999.....	44.6	30.1	17.6	11.8	6.4	*	*	*	*

Non-Hispanic Black, female

2007 5/.....	20.5	22.0	13.4	7.5	*	*	*	*	*
2006 5/.....	22.3	20.4	12.4	6.5	*	*	*	*	7.5
2005 5/.....	19.5	20.0	11.4	3.6	*	*	*	*	*
2004 5/.....	22.7	14.8	8.5	4.3	*	*	*	*	*
2003 5/.....	19.0	14.8	7.7	3.6	*	*	*	*	*
2002.....	19.5	12.5	6.2	3.3	*	*	*	*	*
2001.....	15.1	10.7	4.5	*	*	*	*	*	*
2000.....	13.5	10.4	4.6	*	*	*	*	*	*
1999.....	13.8	8.6	3.1	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

All origins 3/, both sexes

2007.....	11.2	11.0	10.4	10.0	10.1	9.9	13.5	12.7	11.6
2006.....	11.3	11.0	9.9	9.7	9.7	10.4	12.8	13.1	11.5
2005.....	11.4	10.9	9.8	9.9	9.5	11.3	13.0	14.5	12.0
2004.....	11.3	10.7	9.7	10.0	9.5	11.0	13.3	13.4	11.9
2003.....	11.4	10.9	10.4	9.6	9.7	11.3	12.9	14.2	12.5
2002.....	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001.....	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000.....	10.7	10.3	9.6	9.1	9.7	11.5	13.4	14.7	14.2
1999.....	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5

All origins 3/, male

2007.....	18.4	18.9	18.0	17.9	18.8	19.6	29.1	29.9	33.5
2006.....	18.7	18.4	17.0	17.0	18.0	20.8	28.0	30.7	33.6
2005.....	19.1	18.7	17.0	17.9	17.4	22.6	28.1	34.6	35.4
2004.....	18.4	18.0	16.9	17.4	17.8	22.1	28.2	32.0	35.9
2003.....	18.9	18.6	17.8	17.6	18.5	22.5	27.8	33.8	37.8
2002.....	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001.....	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000.....	17.9	17.3	16.5	16.1	17.8	23.1	29.6	36.6	44.7
1999.....	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3

All origins 3/, female

2007.....	4.1	3.4	3.2	2.8	2.5	2.0	2.0	1.9	1.1
2006.....	4.1	3.9	3.2	3.1	2.4	1.9	1.6	2.1	1.1
2005.....	4.0	3.3	3.0	2.6	2.7	2.3	2.1	2.0	1.3
2004.....	4.3	3.7	3.0	3.2	2.3	2.1	2.6	1.9	1.0
2003.....	4.1	3.5	3.4	2.4	2.1	2.3	2.3	2.2	1.3
2002.....	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001.....	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000.....	3.7	3.5	3.1	2.8	2.7	2.4	2.2	1.9	1.7
1999.....	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic 4/, both sexes

2007.....	11.8	11.5	10.8	10.5	10.6	10.3	14.0	13.2	11.9
2006.....	12.0	11.6	10.5	10.1	10.2	11.0	13.3	13.5	11.8
2005.....	11.9	11.4	10.3	10.4	10.0	11.7	13.4	15.0	12.3
2004.....	11.9	11.1	10.1	10.3	10.0	11.3	13.9	13.6	12.1
2003.....	11.8	11.4	10.9	10.0	10.1	11.6	13.3	14.7	12.7
2002.....	11.5	10.8	11.1	9.9	10.4	12.2	14.1	15.8	12.7
2001.....	11.0	10.8	10.8	10.0	10.7	11.7	14.1	15.5	12.9
2000.....	10.8	10.5	9.9	9.4	9.9	11.7	13.8	15.1	14.4
1999.....	10.8	9.9	10.2	9.7	10.2	12.6	14.2	14.7	13.6

Non-Hispanic 4/, male

2007.....	19.5	19.8	18.9	18.7	19.5	20.3	30.2	31.0	34.7
2006.....	19.8	19.4	18.0	17.5	18.9	22.0	29.2	31.6	34.7
2005.....	19.9	19.7	17.7	18.8	18.3	23.3	28.9	35.9	36.3
2004.....	19.3	18.7	17.6	17.9	18.6	22.8	29.4	32.7	37.0
2003.....	19.5	19.5	18.6	18.2	19.1	23.2	28.7	35.0	38.7
2002.....	19.1	18.6	19.3	17.6	19.3	24.9	30.8	38.3	39.5
2001.....	18.5	18.2	18.5	17.2	20.0	23.5	30.9	37.9	40.8
2000.....	18.1	17.6	16.9	16.5	18.2	23.5	30.4	37.4	45.8
1999.....	18.1	16.6	17.5	17.6	19.3	25.3	31.1	36.6	43.9

Non-Hispanic 4/, female

2007.....	4.4	3.6	3.3	3.0	2.7	2.1	2.1	2.0	1.2
2006.....	4.4	4.1	3.4	3.3	2.5	2.1	1.7	2.1	1.2
2005.....	4.2	3.5	3.2	2.7	2.8	2.4	2.2	2.1	1.3
2004.....	4.6	3.9	3.1	3.4	2.5	2.1	2.7	1.9	1.0
2003.....	4.3	3.6	3.6	2.5	2.2	2.3	2.4	2.3	1.3
2002.....	4.2	3.4	3.4	2.9	2.6	2.1	2.3	2.1	1.2
2001.....	3.7	3.6	3.5	3.4	2.7	2.3	2.4	2.2	1.4
2000.....	3.7	3.6	3.3	2.9	2.8	2.4	2.2	1.9	1.7
1999.....	3.7	3.5	3.4	2.5	2.4	2.7	2.4	1.8	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic White, both sexes

2007 5/.....	12.1	12.2	11.7	11.4	11.5	11.3	15.1	14.4	13.1
2006 5/.....	12.2	12.1	11.5	10.9	11.1	12.1	14.5	14.7	12.6
2005 5/.....	12.2	12.1	11.2	11.3	11.0	12.7	14.7	16.2	13.3
2004 5/.....	12.2	11.8	10.7	11.2	10.8	12.2	14.9	14.6	13.1
2003 5/.....	12.0	12.0	11.7	10.8	11.0	12.6	14.4	15.7	13.8
2002.....	11.8	11.3	12.0	10.6	11.3	13.3	14.9	17.1	13.7
2001.....	11.4	11.4	11.5	10.9	11.7	12.5	15.1	16.4	14.0
2000.....	10.8	10.8	10.3	10.1	10.7	12.4	14.9	16.1	15.5
1999.....	10.8	10.3	10.7	10.2	11.0	13.5	15.0	15.6	14.7

Non-Hispanic White, male

2007 5/.....	19.7	20.6	20.0	19.9	21.1	22.3	32.3	33.8	38.3
2006 5/.....	19.8	20.1	19.3	18.7	20.4	23.9	31.3	34.4	37.0
2005 5/.....	20.1	20.6	19.1	20.3	19.9	24.9	31.0	38.7	39.2
2004 5/.....	19.7	19.5	18.3	19.4	19.7	24.2	31.5	34.9	40.1
2003 5/.....	19.5	20.4	19.8	19.5	20.7	24.9	30.9	37.4	42.1
2002.....	19.2	19.3	20.4	18.8	20.9	26.8	32.3	41.3	42.5
2001.....	19.0	19.0	19.3	18.5	21.4	24.8	32.9	39.9	44.0
2000.....	18.0	18.0	17.5	17.5	19.4	24.6	32.8	39.7	49.6
1999.....	17.8	17.3	18.0	18.2	20.6	26.7	32.9	38.7	47.5

Non-Hispanic White, female

2007 5/.....	4.6	3.9	3.6	3.3	3.0	2.2	2.2	2.1	1.2
2006 5/.....	4.8	4.3	3.9	3.6	2.8	2.3	1.9	2.4	1.1
2005 5/.....	4.4	3.9	3.6	2.9	2.9	2.5	2.5	2.1	1.5
2004 5/.....	4.9	4.2	3.4	3.6	2.9	2.4	2.8	2.1	1.0
2003 5/.....	4.6	3.8	3.9	2.7	2.4	2.4	2.5	2.3	1.4
2002.....	4.5	3.4	3.9	3.0	2.9	2.2	2.4	2.3	1.2
2001.....	3.9	3.8	3.9	3.8	3.0	2.4	2.5	2.2	1.5
2000.....	3.7	3.8	3.5	3.2	3.0	2.6	2.3	2.0	1.8
1999.....	3.7	3.5	3.8	2.7	2.6	2.9	2.5	1.9	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic Black, both sexes

2007 5/.....	13.6	10.6	8.5	6.6	5.8	5.0	7.2	*	*
2006 5/.....	13.5	11.5	6.8	6.9	6.4	5.4	6.0	*	6.4
2005 5/.....	13.2	10.3	6.2	6.4	5.8	7.7	5.1	6.5	*
2004 5/.....	12.3	10.2	8.8	7.0	6.2	6.7	7.3	6.1	*
2003 5/.....	12.8	10.4	7.4	6.3	6.0	6.4	5.7	6.5	*
2002.....	13.2	10.3	7.3	7.1	5.2	6.5	8.6	*	*
2001.....	11.4	9.9	8.1	5.4	5.9	7.6	6.9	8.3	*
2000.....	13.4	10.5	7.5	5.9	6.3	7.9	4.7	6.9	*
1999.....	13.7	8.8	8.7	8.0	6.6	7.0	8.5	6.2	*

Non-Hispanic Black, male

2007 5/.....	24.0	19.9	16.4	13.5	11.8	10.3	17.3	*	*
2006 5/.....	25.6	20.8	13.3	13.5	13.1	11.7	15.7	*	*
2005 5/.....	24.2	19.1	12.0	11.4	10.5	16.3	13.8	15.1	*
2004 5/.....	21.8	19.1	16.8	12.3	13.7	16.3	16.3	16.2	*
2003 5/.....	24.1	18.6	13.9	11.9	11.6	12.5	12.6	15.1	*
2002.....	24.6	18.0	14.6	12.4	10.5	14.0	20.2	*	*
2001.....	20.5	17.9	16.3	10.2	12.0	16.2	16.1	20.4	*
2000.....	23.7	19.2	13.9	11.5	13.1	18.5	9.6	17.2	*
1999.....	24.7	15.5	17.8	15.7	13.0	15.9	19.8	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

Non-Hispanic Black, female

2007 5/.....	4.5	2.6	2.1	*	*	*	*	*	*
2006 5/.....	3.0	3.7	*	*	*	*	*	*	*
2005 5/.....	3.6	2.8	*	*	*	*	*	*	*
2004 5/.....	4.1	2.6	2.4	*	*	*	*	*	*
2003 5/.....	2.9	3.5	*	*	*	*	*	*	*
2002.....	3.3	3.7	*	*	*	*	*	*	*
2001.....	3.5	3.0	*	*	*	*	*	*	*
2000.....	4.5	3.1	*	*	*	*	*	*	*
1999.....	4.3	3.2	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes races other than white and black.

5/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality."

6/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

7/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.