

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nchsed@cdc.gov](mailto:nchsed@cdc.gov), or call 301-458-4688

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,719.1	16,244.8	1,983.8	385.9	585.5	819.8	1,023.1	1,495.4	2,723.6	5,636.1	12,330.0	26,088.2
1901	1,641.5	14,139.0	1,695.0	353.8	554.6	804.6	1,029.5	1,500.0	2,776.5	5,622.8	12,462.2	26,082.7
1902	1,548.1	13,887.7	1,655.7	330.8	514.3	746.2	960.0	1,398.6	2,590.5	5,291.3	11,411.7	23,560.0
1903	1,562.8	13,261.1	1,542.1	344.1	517.4	747.1	975.7	1,430.6	2,719.8	5,504.2	12,078.4	25,369.4
1904	1,640.0	13,922.7	1,591.5	367.2	549.4	778.9	1,020.6	1,506.8	2,854.8	5,822.0	12,610.2	26,998.9
1905	1,588.9	14,120.6	1,498.9	339.9	519.5	739.2	978.1	1,465.1	2,772.2	5,618.8	12,241.1	26,151.4
1906	1,571.8	14,484.8	1,580.0	326.5	530.3	747.2	977.9	1,450.3	2,708.1	5,495.2	12,035.5	25,511.1
1907	1,592.5	13,858.0	1,468.3	317.2	530.0	752.9	1,015.6	1,505.6	2,856.7	5,876.5	12,870.7	26,911.0
1908	1,468.2	13,320.7	1,396.8	302.5	476.3	668.8	897.2	1,376.8	2,621.1	5,376.5	11,953.1	24,862.7
1909	1,424.7	12,668.9	1,348.9	281.5	438.5	634.6	872.5	1,333.5	2,561.6	5,390.1	11,836.0	24,492.2
1910	1,468.0	13,176.3	1,397.3	293.5	453.8	654.5	898.5	1,374.5	2,624.9	5,556.7	12,218.6	25,030.8
1911	1,390.5	11,404.5	1,176.0	257.2	448.0	637.6	887.9	1,349.6	2,580.3	5,499.9	12,008.4	24,641.4
1912	1,359.7	11,105.5	1,094.1	246.0	425.6	610.7	861.9	1,336.7	2,583.2	5,449.8	12,016.4	24,219.4
1913	1,380.6	11,478.7	1,193.4	267.7	437.6	618.6	873.5	1,348.2	2,548.5	5,414.7	11,792.2	23,593.3
1914	1,330.2	10,716.8	1,024.2	250.6	419.5	598.6	848.9	1,311.7	2,513.4	5,411.0	11,558.1	23,147.9
1915	1,317.6	10,237.7	924.2	229.6	407.1	582.5	833.3	1,309.0	2,554.0	5,564.0	12,007.6	24,028.6
1916	1,381.1	10,565.9	1,111.5	245.0	438.2	616.9	877.3	1,361.8	2,654.3	5,715.5	12,385.3	25,042.6
1917	1,397.1	10,457.2	1,066.0	256.0	468.9	649.1	900.8	1,385.6	2,678.6	5,728.4	12,386.2	24,593.6
1918	1,810.0	11,167.2	1,573.5	412.8	1,070.6	1,643.5	1,339.3	1,524.1	2,648.1	5,505.0	11,295.7	22,213.5
1919	1,289.4	9,995.0	928.0	269.3	531.6	752.0	861.6	1,226.9	2,310.0	4,998.3	10,782.0	22,225.0
1920	1,298.9	9,226.4	987.2	263.9	487.4	677.5	811.3	1,219.3	2,358.8	5,253.4	11,888.1	24,825.2
1921	1,149.8	8,063.2	801.2	246.9	387.3	488.5	675.7	1,119.6	2,209.2	4,897.9	11,120.5	23,914.5
1922	1,169.3	7,757.0	742.0	213.9	380.4	502.6	712.0	1,179.9	2,318.9	5,219.8	11,754.5	25,813.6
1923	1,213.0	8,107.7	806.7	214.8	387.6	504.4	730.8	1,222.1	2,385.9	5,334.0	12,346.6	27,967.3
1924	1,159.0	7,676.0	683.2	201.2	376.3	476.8	712.1	1,212.1	2,300.5	5,097.4	11,720.3	26,177.4
1925	1,168.1	7,542.7	641.0	196.6	375.9	483.7	718.7	1,216.7	2,326.1	5,173.5	11,929.0	27,226.8
1926	1,211.0	7,788.2	723.4	193.2	370.5	489.5	737.4	1,268.7	2,407.2	5,383.7	12,543.6	27,967.8
1927	1,131.5	6,883.9	591.0	188.5	347.7	469.1	705.0	1,204.6	2,291.8	5,118.5	11,586.6	25,012.1
1928	1,188.6	7,307.7	647.5	194.0	368.7	497.3	747.1	1,277.1	2,422.4	5,427.9	12,520.4	26,832.2
1929	1,187.8	7,156.7	625.5	187.1	361.0	499.0	732.9	1,266.4	2,449.9	5,398.5	12,224.1	25,431.7
1930	1,132.1	6,900.4	563.6	171.7	334.3	465.8	682.0	1,217.6	2,402.7	5,137.3	11,269.2	22,795.6
1931	1,106.5	6,444.8	526.6	166.4	318.7	449.4	673.4	1,198.4	2,355.9	4,993.2	11,053.9	22,277.8
1932	1,087.7	6,132.4	461.9	154.2	288.1	417.1	634.7	1,156.1	2,340.8	4,988.2	11,430.8	23,334.2
1933	1,068.7	6,125.7	472.6	149.8	273.9	406.1	618.4	1,143.2	2,315.7	4,904.1	11,131.8	22,228.5
1934	1,105.4	6,678.1	507.7	153.4	275.3	407.5	622.9	1,178.8	2,352.2	4,941.0	11,409.0	22,484.9
1935	1,094.5	6,093.8	440.9	152.9	270.5	402.7	622.9	1,160.4	2,315.7	4,872.4	11,312.6	22,457.9
1936	1,155.2	6,288.1	439.8	147.7	278.3	409.8	650.8	1,208.2	2,412.0	5,077.8	12,174.3	24,273.5
1937	1,125.9	6,127.2	418.7	135.4	262.8	387.5	621.7	1,180.6	2,351.8	4,899.2	11,702.1	22,720.1
1938	1,064.0	5,804.5	383.8	121.0	228.2	336.1	555.2	1,090.8	2,206.4	4,711.1	11,093.1	21,258.9
1939	1,060.4	5,367.1	318.3	109.4	212.4	317.4	530.4	1,072.8	2,214.3	4,719.4	11,253.7	22,331.3

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,791.1	17,314.4	2,054.2	384.3	594.0	824.3	1,057.7	1,569.9	2,874.1	5,929.0	12,826.4	26,871.2
1901	1,725.9	15,637.5	1,769.3	366.0	570.0	833.0	1,097.3	1,609.4	2,953.0	5,919.2	12,970.3	26,806.6
1902	1,635.3	15,336.5	1,713.8	342.2	522.9	765.9	1,032.1	1,512.3	2,798.1	5,650.9	12,050.9	24,862.0
1903	1,643.6	14,660.8	1,590.7	352.8	533.2	768.5	1,043.3	1,550.0	2,900.6	5,845.0	12,675.2	26,266.3
1904	1,727.6	15,391.1	1,659.3	369.8	549.5	799.9	1,110.8	1,636.8	3,108.6	6,173.0	13,256.6	28,074.7
1905	1,674.4	15,662.5	1,577.3	344.7	529.8	759.8	1,062.0	1,596.3	2,984.6	5,898.9	12,884.5	27,084.2
1906	1,672.6	16,017.0	1,640.7	337.7	566.1	787.5	1,092.8	1,595.7	2,944.6	5,820.5	12,651.3	26,155.4
1907	1,702.2	15,291.2	1,527.3	329.9	580.0	814.9	1,142.9	1,683.1	3,111.7	6,269.7	13,398.0	27,501.2
1908	1,552.4	14,696.5	1,456.8	312.1	504.4	700.6	981.9	1,515.5	2,844.3	5,642.4	12,590.2	25,149.6
1909	1,506.4	13,985.4	1,406.1	292.5	459.4	663.6	954.2	1,478.3	2,773.7	5,699.3	12,392.8	25,140.6
1910	1,556.4	14,533.2	1,458.0	301.7	483.6	692.9	995.1	1,523.4	2,866.6	5,873.6	12,741.6	25,579.4
1911	1,472.7	12,593.8	1,222.6	278.5	469.2	670.9	978.3	1,494.1	2,802.6	5,811.3	12,508.1	24,927.2
1912	1,448.6	12,331.5	1,147.0	257.7	451.4	647.0	950.7	1,494.9	2,817.2	5,785.9	12,521.7	24,865.0
1913	1,476.0	12,756.1	1,245.1	280.6	469.9	665.1	970.2	1,500.8	2,790.3	5,771.4	12,284.8	24,136.5
1914	1,419.0	11,894.2	1,074.8	262.7	441.8	637.6	940.4	1,446.4	2,740.5	5,776.4	12,054.7	23,686.4
1915	1,399.6	11,450.1	968.7	242.9	423.3	618.7	913.5	1,436.1	2,769.3	5,875.2	12,455.2	24,672.3
1916	1,475.4	11,820.5	1,169.2	261.3	454.9	659.5	970.5	1,509.1	2,898.3	6,063.6	12,865.5	25,546.4
1917	1,504.6	11,736.4	1,133.3	259.9	496.0	708.4	1,056.7	1,547.5	2,933.5	6,110.9	12,899.2	25,107.1
1918	1,980.5	12,453.5	1,600.6	415.3	1,215.2	1,902.4	1,528.3	1,670.5	2,870.2	5,849.5	11,811.5	22,756.1
1919	1,345.8	10,190.2	973.5	283.4	530.3	744.2	905.3	1,290.3	2,444.2	5,194.7	11,096.7	22,958.9
1920	1,338.1	10,358.0	1,027.7	280.1	477.8	643.0	821.8	1,262.7	2,463.3	5,453.3	12,214.3	25,300.9
1921	1,194.7	9,012.9	842.4	265.7	383.1	479.5	694.1	1,161.3	2,325.7	5,110.4	11,441.7	24,123.7
1922	1,226.3	8,697.8	787.6	231.0	377.6	499.1	741.0	1,249.1	2,474.8	5,508.4	12,175.5	25,784.6
1923	1,274.5	9,022.9	846.0	232.3	386.7	510.2	772.2	1,311.5	2,556.0	5,622.4	12,743.8	27,939.9
1924	1,230.4	8,623.6	723.8	219.1	376.3	483.3	758.7	1,309.2	2,488.9	5,469.4	12,284.2	26,376.0
1925	1,239.4	8,457.6	673.2	215.7	375.9	491.2	761.9	1,326.2	2,513.5	5,540.7	12,527.3	27,352.5
1926	1,285.7	8,708.8	762.4	211.7	373.2	495.4	789.3	1,387.6	2,595.5	5,764.9	13,183.5	28,129.2
1927	1,209.4	7,747.0	624.1	208.0	350.4	475.7	758.6	1,321.0	2,504.5	5,523.2	12,263.1	25,423.3
1928	1,283.4	8,232.7	683.4	212.8	376.4	510.1	801.8	1,413.0	2,646.3	5,854.4	13,226.1	27,146.5
1929	1,276.9	8,003.5	660.3	207.0	372.6	515.5	798.5	1,407.0	2,694.3	5,837.9	12,886.3	25,980.3
1930	1,225.3	7,701.3	602.6	189.8	349.6	488.9	745.9	1,359.8	2,661.0	5,582.0	11,912.2	23,671.2
1931	1,201.5	7,217.7	561.6	183.4	336.3	471.5	744.3	1,343.5	2,617.1	5,435.6	11,749.1	23,409.2
1932	1,172.9	6,854.5	487.7	169.0	301.6	431.9	692.0	1,290.8	2,605.7	5,407.2	12,107.8	24,233.7
1933	1,162.8	6,828.2	503.0	165.6	292.2	426.1	681.9	1,291.1	2,599.7	5,363.3	11,831.7	23,272.2
1934	1,211.6	7,479.3	540.6	169.6	299.0	433.9	695.0	1,349.6	2,660.3	5,434.9	12,217.5	23,512.0
1935	1,202.1	6,885.1	469.2	169.6	294.4	430.3	698.8	1,329.2	2,629.5	5,367.3	12,168.3	23,466.8
1936	1,273.0	7,066.1	471.0	165.7	304.9	440.3	739.0	1,405.6	2,767.7	5,606.6	13,057.0	25,267.1
1937	1,246.8	6,874.4	446.3	152.3	291.2	422.0	703.9	1,377.5	2,718.4	5,453.6	12,638.9	23,803.7
1938	1,170.8	6,523.9	408.3	135.1	249.4	363.2	622.1	1,264.6	2,532.1	5,245.0	11,879.5	22,215.5
1939	1,168.2	6,031.9	342.6	125.5	235.6	343.9	595.2	1,245.8	2,551.1	5,273.3	12,065.2	23,263.9

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,646.9	14,541.2	1,912.3	387.5	577.6	815.2	975.2	1,417.7	2,576.0	5,358.3	11,877.3	25,517.4
1901	1,556.9	12,610.5	1,619.6	341.4	540.0	775.7	957.5	1,385.5	2,603.5	5,342.1	12,001.2	25,557.1
1902	1,460.5	12,409.9	1,596.8	319.4	506.2	726.3	883.4	1,279.3	2,386.8	4,951.1	10,834.6	22,611.1
1903	1,481.4	11,833.1	1,492.8	335.3	502.3	725.2	903.8	1,304.9	2,541.9	5,182.0	11,542.2	24,713.2
1904	1,551.8	12,423.9	1,522.6	364.6	549.4	757.5	924.9	1,369.9	2,604.4	5,490.5	12,032.6	26,209.0
1905	1,502.5	12,546.0	1,419.2	334.9	509.7	718.0	889.0	1,326.6	2,561.9	5,354.2	11,669.3	25,486.5
1906	1,467.3	12,916.0	1,538.2	315.2	494.7	703.8	851.3	1,290.7	2,464.7	5,177.8	11,477.7	25,028.7
1907	1,478.4	12,389.0	1,408.4	304.3	480.1	686.0	875.3	1,310.8	2,592.9	5,493.1	12,395.3	26,467.2
1908	1,379.8	11,909.0	1,335.7	292.8	447.9	633.9	802.8	1,222.2	2,385.7	5,113.5	11,370.5	24,640.5
1909	1,338.8	11,317.3	1,290.4	270.4	417.4	602.7	781.9	1,172.5	2,337.0	5,083.3	11,327.7	23,986.1
1910	1,374.3	11,762.1	1,335.3	285.1	423.3	611.7	790.4	1,207.2	2,366.0	5,240.5	11,740.1	24,599.8
1911	1,303.7	10,184.1	1,128.4	255.7	425.5	600.9	787.2	1,186.6	2,339.9	5,186.3	11,548.0	24,415.1
1912	1,265.0	9,847.4	1,039.0	234.0	399.3	570.9	762.9	1,158.1	2,329.4	5,111.1	11,553.3	23,713.8
1913	1,280.1	10,168.6	1,140.5	254.6	405.1	568.1	766.2	1,175.8	2,285.5	5,054.4	11,342.5	23,170.4
1914	1,236.6	9,509.9	972.5	238.2	397.0	556.4	747.6	1,159.2	2,265.1	5,039.2	11,102.4	22,729.0
1915	1,231.3	8,995.3	878.7	216.1	390.9	543.5	744.6	1,165.0	2,318.6	5,247.2	11,599.0	23,532.4
1916	1,282.7	9,281.2	1,052.3	228.5	421.7	571.9	774.8	1,195.0	2,386.6	5,360.1	11,950.1	24,659.6
1917	1,285.4	9,147.2	1,011.5	241.8	442.4	587.4	785.1	1,202.0	2,398.6	5,336.1	11,921.1	24,205.7
1918	1,638.7	9,850.9	1,545.8	410.2	943.6	1,399.4	1,133.6	1,357.6	2,403.6	5,150.3	10,827.4	21,805.0
1919	1,231.5	7,974.8	881.4	255.0	532.9	759.9	814.1	1,154.6	2,162.1	4,795.0	10,496.2	21,676.0
1920	1,258.2	8,067.3	945.7	247.4	496.7	712.7	799.8	1,170.2	2,243.9	5,046.3	11,589.0	24,466.8
1921	1,103.4	7,082.1	758.8	227.8	391.5	497.6	655.6	1,072.7	2,082.1	4,678.2	10,825.8	23,756.1
1922	1,110.7	6,787.3	695.0	196.6	383.2	506.1	680.7	1,102.8	2,150.0	4,922.3	11,369.0	25,835.7
1923	1,149.8	7,164.7	766.1	197.0	388.6	498.7	686.1	1,123.5	2,201.4	5,035.4	11,981.4	27,988.4
1924	1,085.5	6,699.8	641.3	183.1	376.3	470.4	661.8	1,105.9	2,095.4	4,711.3	11,200.9	26,025.5
1925	1,094.7	6,597.9	607.8	172.2	376.0	476.1	672.4	1,097.2	2,121.9	4,792.6	11,377.0	27,130.3
1926	1,134.0	6,838.3	683.2	174.4	367.8	483.6	681.8	1,139.3	2,201.7	4,988.7	11,952.1	27,843.8
1927	1,051.3	5,998.9	556.9	168.6	345.0	462.4	647.8	1,078.0	2,059.3	4,699.0	10,958.7	24,696.0
1928	1,111.4	6,356.4	610.3	174.7	361.1	484.5	689.1	1,129.0	2,177.3	4,985.3	11,861.5	26,593.6
1929	1,096.4	6,286.1	589.6	166.9	349.6	482.7	663.6	1,112.9	2,183.8	4,943.1	11,602.9	25,015.9
1930	1,036.7	6,073.8	523.3	153.2	319.3	443.1	614.9	1,062.3	2,123.7	4,675.8	10,663.4	22,138.0
1931	1,009.3	5,646.5	490.6	149.0	301.4	427.6	599.2	1,040.6	2,074.7	4,536.0	10,400.3	21,428.5
1932	1,000.7	5,386.2	435.4	139.1	274.9	402.6	574.9	1,010.2	2,056.4	4,577.4	10,796.0	22,659.1
1933	972.8	5,403.0	441.3	133.5	256.0	386.5	552.4	983.3	2,010.9	4,432.3	10,773.7	21,442.3
1934	997.2	5,853.6	474.0	136.6	252.0	381.6	548.3	994.7	2,022.6	4,436.0	10,651.6	21,711.5
1935	985.1	5,280.3	411.9	135.7	247.1	375.6	544.6	979.1	1,981.0	4,369.1	10,514.3	21,698.5
1936	1,035.5	5,489.6	407.8	129.3	252.1	379.9	560.2	996.7	2,033.5	4,542.8	11,354.3	23,525.7
1937	1,003.2	5,359.9	390.3	118.2	234.9	353.7	537.5	970.2	1,962.9	4,341.3	10,835.8	21,904.6
1938	955.9	5,065.5	358.8	106.6	207.4	309.6	487.0	905.5	1,861.7	4,176.6	10,369.3	20,539.0
1939	951.4	4,684.3	293.3	92.9	189.5	291.5	464.7	888.8	1,858.8	4,168.0	10,510.5	21,629.4

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,702.4	15,942.1	1,943.5	376.9	571.1	809.7	1,012.3	1,475.6	2,700.1	5,620.0	12,333.0	26,198.2
1901	1,624.7	13,841.4	1,664.4	347.1	539.8	791.8	1,014.9	1,476.6	2,752.8	5,611.6	12,477.6	26,165.3
1902	1,530.9	13,587.0	1,620.2	324.2	498.6	733.9	946.2	1,379.6	2,571.3	5,274.8	11,418.0	23,636.7
1903	1,544.0	12,963.4	1,501.1	337.6	502.5	730.9	961.0	1,411.2	2,694.3	5,480.2	12,093.5	25,403.9
1904	1,619.5	13,611.4	1,550.8	359.6	531.9	761.9	1,003.7	1,485.5	2,821.9	5,804.0	12,609.4	27,100.4
1905	1,568.5	13,833.2	1,458.6	331.7	500.8	721.5	962.1	1,440.0	2,746.6	5,596.1	12,255.4	26,277.6
1906	1,545.8	14,149.1	1,525.7	314.5	507.5	723.5	953.2	1,423.3	2,673.6	5,476.5	12,052.5	25,690.2
1907	1,566.8	13,518.4	1,418.8	307.5	507.6	728.3	988.2	1,475.4	2,823.9	5,857.4	12,893.4	27,125.2
1908	1,445.7	13,037.5	1,356.8	293.1	457.1	646.6	873.7	1,346.0	2,591.1	5,355.8	11,978.9	25,065.3
1909	1,403.4	12,404.8	1,312.6	273.2	420.6	614.0	849.1	1,303.5	2,531.2	5,368.3	11,864.5	24,628.0
1910	1,448.8	12,228.9	1,365.0	287.1	438.0	634.9	872.7	1,345.5	2,597.1	5,540.2	12,250.6	25,246.5
1911	1,366.1	11,175.9	1,139.0	258.5	425.4	612.5	856.9	1,314.2	2,542.3	5,474.3	12,041.2	24,784.7
1912	1,336.6	10,904.7	1,065.2	238.7	403.8	586.6	829.9	1,298.9	2,543.0	5,423.4	12,032.3	24,397.0
1913	1,351.4	11,219.1	1,159.6	258.8	408.8	586.9	833.9	1,303.3	2,499.5	5,380.2	11,815.7	23,796.0
1914	1,299.6	10,454.3	988.5	241.6	390.1	565.1	807.7	1,263.9	2,462.4	5,375.4	11,576.7	23,282.1
1915	1,286.3	9,980.2	890.3	220.6	377.3	549.5	790.2	1,260.5	2,495.5	5,516.9	12,007.6	24,131.6
1916	1,344.5	10,216.7	1,062.9	234.4	395.3	571.3	824.0	1,308.0	2,593.4	5,675.2	12,397.4	25,161.3
1917	1,349.5	10,038.7	997.5	241.6	414.5	594.2	834.8	1,320.2	2,607.4	5,671.4	12,403.7	24,662.5
1918	1,751.1	10,707.3	1,474.0	387.8	992.7	1,594.3	1,269.5	1,448.9	2,574.8	5,446.0	11,296.9	22,270.3
1919	1,242.5	8,684.3	874.7	251.5	463.6	693.2	794.7	1,153.6	2,253.4	4,960.0	10,811.0	22,479.7
1920	1,256.1	8,727.0	942.9	252.0	428.3	623.1	746.7	1,145.4	2,301.3	5,208.8	11,930.5	24,980.1
1921	1,112.9	7,645.4	765.2	240.2	334.5	436.0	614.8	1,051.8	2,155.2	4,867.4	11,173.3	24,129.1
1922	1,132.5	7,345.0	706.6	207.0	325.8	445.8	642.7	1,096.8	2,260.1	5,195.2	11,843.0	26,074.5
1923	1,169.1	7,618.6	749.8	206.3	330.2	442.8	656.2	1,128.7	2,319.1	5,295.8	12,424.3	28,137.6
1924	1,104.4	7,097.7	621.2	191.0	312.6	411.2	629.9	1,105.4	2,219.9	5,029.8	11,747.1	26,135.3
1925	1,113.1	6,998.7	592.2	185.3	311.9	412.9	631.4	1,105.9	2,245.1	5,106.8	11,955.5	27,238.3
1926	1,155.9	7,253.8	667.6	182.4	308.9	413.9	648.0	1,150.4	2,318.6	5,321.8	12,595.2	28,188.2
1927	1,080.0	6,368.6	540.7	178.6	290.8	396.2	616.5	1,091.8	2,203.2	5,064.6	11,652.7	25,210.1
1928	1,142.2	6,745.5	596.5	182.9	305.1	414.5	648.0	1,144.0	2,314.1	5,363.8	12,594.7	27,160.6
1929	1,132.4	6,602.4	580.5	177.0	300.2	414.9	631.3	1,132.3	2,333.8	5,334.3	12,262.6	25,713.8
1930	1,076.8	6,392.5	516.7	160.7	275.6	384.5	585.8	1,084.2	2,281.6	5,064.3	11,321.1	23,049.3
1931	1,057.6	5,980.7	484.8	156.8	262.3	369.0	580.6	1,079.6	2,246.6	4,938.0	11,120.2	22,766.3
1932	1,048.2	5,998.8	433.0	146.5	240.1	346.9	549.6	1,046.5	2,234.6	4,963.5	11,529.5	23,893.1
1933	1,030.3	5,682.9	442.3	141.1	229.4	337.7	535.2	1,034.4	2,212.1	4,869.7	11,248.1	22,821.8
1934	1,063.5	6,163.3	467.3	145.3	230.8	336.6	534.2	1,063.0	2,246.2	4,904.0	11,523.3	23,048.6
1935	1,056.4	5,652.4	409.2	145.3	228.1	335.2	532.7	1,049.9	2,209.1	4,846.4	11,438.5	23,023.2
1936	1,111.7	5,804.0	408.7	140.2	231.5	335.5	550.6	1,087.5	2,291.6	5,027.7	12,298.2	24,901.2
1937	1,084.7	5,636.8	384.3	128.1	218.7	318.2	525.9	1,063.0	2,239.5	4,852.2	11,833.8	23,378.8
1938	1,025.5	5,340.3	349.0	112.7	186.5	273.1	467.6	977.4	2,097.3	4,673.5	11,221.4	21,910.9
1939	1,027.2	4,920.2	287.4	102.1	175.7	258.5	448.9	965.2	2,105.3	4,695.7	11,382.2	23,121.1

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

WHITE, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,774.8	17,586.5	2,016.4	3,777.6	5,798.8	813.2	1,059.3	1,550.4	2,851.9	5,913.2	12,822.9	26,917.5
1901	1,708.4	15,332.2	1,739.1	360.1	556.1	817.9	1,081.8	1,585.8	2,927.0	5,905.5	12,982.1	26,862.3
1902	1,617.5	15,003.3	1,678.7	336.1	508.3	752.8	1,018.8	1,494.3	2,776.4	5,637.0	12,039.2	24,951.6
1903	1,624.6	14,346.0	1,548.4	348.2	519.1	751.4	1,030.7	1,531.3	2,873.5	5,816.8	12,674.8	26,342.8
1904	1,706.2	15,071.1	1,616.9	363.6	532.9	780.4	1,093.6	1,615.9	3,073.1	6,154.6	13,241.8	28,124.2
1905	1,653.5	15,344.1	1,535.6	339.0	512.1	741.7	1,045.5	1,571.0	2,958.6	5,875.6	12,882.0	27,161.4
1906	1,647.3	15,666.5	1,584.8	327.6	546.9	762.9	1,068.2	1,571.3	2,910.8	5,802.2	12,665.1	26,282.4
1907	1,676.9	14,939.8	1,476.9	321.6	559.9	787.8	1,117.0	1,655.5	3,080.1	6,251.7	13,408.6	27,725.1
1908	1,530.5	14,403.8	1,416.4	303.4	487.6	678.2	957.6	1,485.8	2,816.7	5,622.7	12,613.6	25,305.3
1909	1,485.3	13,715.1	1,370.7	285.9	442.5	642.1	930.0	1,447.0	2,747.6	5,676.4	12,408.7	25,216.7
1910	1,537.2	14,296.6	1,424.1	296.3	469.4	672.1	968.7	1,495.1	2,841.1	5,857.9	12,764.1	25,794.3
1911	1,448.4	12,354.0	1,184.4	270.9	449.0	644.4	946.5	1,460.3	2,766.4	5,781.0	12,540.7	25,065.6
1912	1,425.3	12,128.7	1,118.5	251.8	431.8	621.5	916.7	1,458.8	2,777.7	5,758.2	12,517.5	24,992.0
1913	1,447.8	12,484.8	1,211.2	273.8	444.8	633.4	930.5	1,459.2	2,748.4	5,735.9	12,294.2	24,323.7
1914	1,388.8	11,614.3	1,040.7	255.3	414.5	603.3	899.1	1,403.1	2,692.8	5,736.8	12,063.6	23,828.2
1915	1,368.8	11,182.7	935.9	235.1	396.3	585.6	868.4	1,392.8	2,716.2	5,831.2	12,451.4	24,728.8
1916	1,440.3	11,433.5	1,118.6	253.8	417.9	616.9	921.5	1,462.8	2,848.3	6,028.6	12,859.3	25,593.5
1917	1,457.8	11,281.0	1,049.9	258.3	446.1	654.2	941.9	1,491.3	2,875.8	6,054.9	12,904.1	25,159.8
1918	1,925.7	11,979.0	1,501.3	394.9	1,135.4	1,860.7	1,465.9	1,604.8	2,809.9	5,800.3	11,793.5	22,776.5
1919	1,303.5	9,741.4	919.2	270.0	470.0	694.4	847.4	1,227.5	2,404.4	5,167.5	11,113.5	23,166.2
1920	1,298.1	9,810.3	984.1	271.4	424.9	594.3	765.3	1,197.9	2,420.9	5,418.5	12,250.4	25,358.2
1921	1,160.8	8,573.5	807.5	261.1	337.3	431.8	638.9	1,102.4	2,287.9	5,096.5	11,482.3	24,254.8
1922	1,191.4	8,262.5	749.7	226.0	329.4	445.2	676.6	1,172.8	2,441.1	5,503.7	12,243.6	25,791.5
1923	1,231.7	8,498.3	788.3	226.5	336.0	450.0	701.1	1,225.6	2,512.2	5,602.2	12,788.6	27,775.4
1924	1,175.6	7,991.3	660.5	211.0	319.1	417.8	680.1	1,208.8	2,420.0	5,418.2	12,274.4	25,935.8
1925	1,184.2	7,862.6	623.9	205.8	320.2	420.0	678.3	1,216.8	2,445.6	5,485.9	12,531.8	26,951.6
1926	1,229.7	8,111.8	705.0	202.6	318.6	419.3	702.9	1,271.8	2,516.5	5,716.7	13,206.2	27,983.2
1927	1,158.6	7,183.5	571.0	199.5	301.1	404.4	674.7	1,212.5	2,427.8	5,479.6	12,312.7	25,353.4
1928	1,227.3	7,615.7	630.6	203.8	320.8	428.2	707.9	1,284.2	2,552.0	5,797.5	13,269.2	27,189.4
1929	1,220.4	7,385.7	613.7	197.8	318.8	431.2	699.3	1,276.6	2,591.4	5,775.1	12,876.7	25,987.2
1930	1,169.6	7,154.0	553.7	180.5	297.7	406.5	650.9	1,231.2	2,552.2	5,512.2	11,921.4	23,762.2
1931	1,152.6	6,714.5	518.3	175.1	285.5	390.7	651.3	1,231.8	2,519.6	5,390.3	11,771.0	23,763.2
1932	1,133.0	6,387.8	458.3	162.1	258.5	360.8	606.3	1,188.1	2,513.2	5,380.6	12,171.9	24,618.5
1933	1,124.2	6,351.0	472.0	158.1	250.9	356.3	595.7	1,189.7	2,509.9	5,345.5	11,935.0	23,714.3
1934	1,168.1	6,916.5	495.8	163.1	258.4	359.8	602.8	1,235.8	2,567.7	5,407.9	12,315.7	23,939.7
1935	1,162.4	6,401.3	436.6	162.7	256.0	359.9	603.5	1,220.2	2,536.2	5,351.5	12,270.6	23,933.1
1936	1,226.3	6,532.8	438.1	158.7	262.5	361.0	632.2	1,282.2	2,659.7	5,559.2	13,162.9	25,728.1
1937	1,203.5	6,339.7	409.2	146.2	250.8	349.8	604.8	1,260.1	2,619.1	5,405.5	12,744.0	24,351.0
1938	1,132.0	6,011.6	371.6	127.7	213.1	299.6	534.4	1,153.7	2,437.7	5,213.2	12,001.5	22,747.8
1939	1,134.6	5,526.4	310.1	118.6	204.7	284.3	513.5	1,141.0	2,456.6	5,261.7	12,175.7	23,958.0

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	1,629.7	14,262.8	1,869.4	376.2	562.8	806.1	962.1	1,397.6	2,551.7	5,341.7	11,885.1	25,674.8
1901	1,540.7	12,319.7	1,588.6	334.0	524.4	765.3	943.8	1,362.5	2,582.3	5,333.2	12,018.6	25,656.3
1902	1,443.9	12,141.4	1,560.8	312.1	489.5	714.8	869.2	1,259.5	2,370.2	4,932.3	10,855.7	22,672.7
1903	1,462.9	11,552.1	1,453.0	326.9	486.8	710.1	887.0	1,285.0	2,518.4	5,162.1	11,570.2	24,713.0
1904	1,532.0	12,121.0	1,483.7	355.5	530.8	742.9	908.2	1,348.1	2,574.3	5,473.0	12,043.1	26,344.1
1905	1,482.6	12,269.6	1,380.1	324.3	490.0	700.7	873.6	1,301.9	2,536.9	5,332.4	11,897.6	25,622.8
1906	1,440.7	12,595.1	1,465.6	301.2	468.5	681.3	827.3	1,262.0	2,430.9	5,159.7	11,496.4	25,243.7
1907	1,452.7	12,061.4	1,359.6	293.2	455.4	664.2	847.2	1,279.0	2,560.7	5,474.0	12,428.1	26,671.1
1908	1,356.8	11,635.6	1,296.0	282.6	426.1	612.1	780.8	1,190.9	2,354.5	5,092.6	11,397.5	24,878.3
1909	1,317.5	11,059.5	1,253.1	260.2	398.2	583.3	759.9	1,145.0	2,303.4	5,063.7	11,367.1	24,166.0
1910	1,355.3	11,524.0	1,304.5	277.9	405.9	593.7	766.0	1,178.2	2,337.1	5,224.1	11,780.6	24,814.0
1911	1,279.3	9,966.4	1,092.4	245.9	401.4	577.6	757.6	1,150.3	2,301.2	5,166.4	11,580.5	24,561.1
1912	1,243.0	9,647.8	1,010.6	225.5	375.3	548.4	733.8	1,119.5	2,289.6	5,087.1	11,587.4	23,927.0
1913	1,250.0	9,919.3	1,106.8	243.6	372.4	536.3	727.3	1,128.2	2,230.1	5,022.1	11,378.9	23,382.8
1914	1,205.9	9,263.7	935.1	227.7	365.4	523.7	706.9	1,107.3	2,211.8	5,009.0	11,130.4	22,855.0
1915	1,199.6	8,766.3	843.5	205.8	358.2	510.5	704.1	1,111.8	2,255.4	5,198.1	11,602.9	23,669.1
1916	1,244.3	8,967.0	1,005.8	214.6	372.7	522.8	717.1	1,134.0	2,315.8	5,316.5	11,979.9	24,831.3
1917	1,236.8	8,762.4	943.6	224.5	383.5	531.5	717.9	1,128.0	2,315.0	5,280.7	11,951.5	24,285.4
1918	1,575.8	9,401.4	1,446.0	380.6	867.7	1,343.2	1,055.8	1,273.6	2,318.0	5,083.5	10,847.6	21,887.5
1919	1,179.8	7,599.0	829.1	232.7	457.6	692.0	737.2	1,070.5	2,088.5	4,746.9	10,537.6	21,964.9
1920	1,212.6	7,612.4	900.4	232.2	431.6	652.7	726.4	1,086.8	2,171.3	4,993.3	11,639.0	24,695.1
1921	1,063.2	6,682.3	721.7	219.0	331.7	440.3	588.4	995.7	2,012.0	4,632.6	10,891.4	24,033.3
1922	1,072.0	6,394.5	662.1	187.7	322.4	446.5	605.8	1,013.2	2,066.7	4,880.6	11,478.5	26,291.6
1923	1,104.5	6,707.8	709.9	185.6	324.5	435.6	607.5	1,023.1	2,112.1	4,981.8	12,091.3	28,418.8
1924	1,030.9	6,172.7	580.4	170.6	306.2	404.6	575.6	993.3	2,004.6	4,630.4	11,263.9	26,290.4
1925	1,039.8	6,102.6	559.4	164.5	303.7	405.8	580.8	986.1	2,029.2	4,716.8	11,426.1	27,461.3
1926	1,079.8	6,364.0	628.9	161.9	299.4	408.5	588.8	1,019.4	2,105.0	4,915.8	12,032.6	28,347.8
1927	999.0	5,527.6	509.2	157.3	280.4	387.9	554.1	961.5	1,960.6	4,637.8	11,042.5	25,098.8
1928	1,054.5	5,844.2	561.3	161.4	289.4	400.7	584.0	992.4	2,057.4	4,916.9	11,967.0	27,138.4
1929	1,041.8	5,791.1	546.2	155.7	281.8	398.7	558.9	975.7	2,057.1	4,880.1	11,687.7	25,506.1
1930	981.4	5,602.3	478.4	140.3	253.7	362.6	516.8	924.6	1,993.2	4,601.9	10,755.8	22,514.1
1931	960.1	5,218.4	450.2	138.0	239.3	347.5	506.1	914.7	1,956.4	4,472.9	10,508.6	22,018.1
1932	961.2	4,982.6	406.9	130.5	221.9	333.1	489.9	893.8	1,939.2	4,536.5	10,927.6	23,348.7
1933	934.2	4,991.4	411.6	123.6	208.1	319.2	471.9	867.1	1,896.3	4,383.4	10,602.8	22,149.0
1934	956.4	5,383.0	437.7	127.0	203.5	313.6	462.6	877.3	1,906.1	4,391.7	10,781.6	22,376.8
1935	948.2	4,877.7	380.9	127.3	200.5	310.7	459.0	867.4	1,863.7	4,335.5	10,662.9	22,337.2
1936	994.8	5,050.8	378.3	121.0	200.8	310.4	466.2	879.4	1,903.6	4,493.1	11,495.9	24,277.8
1937	963.7	4,909.8	358.7	109.3	186.9	287.0	444.6	852.8	1,840.2	4,298.7	10,993.2	22,645.7
1938	917.1	4,645.7	325.7	97.2	160.1	246.9	399.2	789.6	1,739.9	4,136.7	10,504.6	21,279.7
1939	918.1	4,293.0	264.0	85.0	147.0	233.0	382.9	778.4	1,737.0	4,135.9	10,657.0	22,489.6

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	2,504.0	33,390.4	4,346.4	895.5	1,148.5	1,211.9	1,483.9	2,429.7	4,212.7	6,894.2	12,091.9	21,516.4
1901	2,434.3	30,972.8	3,483.7	737.1	1,137.6	1,294.6	1,654.5	2,607.5	4,273.9	6,484.7	11,253.7	22,620.3
1902	2,356.1	30,888.0	3,735.5	714.2	1,141.0	1,212.6	1,547.0	2,301.7	3,797.5	6,550.4	10,921.4	20,306.1
1903	2,446.6	30,106.9	3,943.5	721.2	1,120.5	1,352.7	1,597.8	2,358.2	4,307.9	7,326.4	10,884.7	23,877.6
1904	2,610.5	31,565.2	3,976.4	809.0	1,275.5	1,412.5	1,737.3	2,535.3	4,899.4	7,181.1	12,676.1	22,538.1
1905	2,553.0	31,021.9	3,867.9	813.7	1,308.2	1,389.5	1,650.4	2,683.2	4,358.1	7,322.6	11,098.5	20,484.4
1906	2,417.0	27,093.8	3,639.5	779.1	1,214.5	1,375.3	1,694.8	2,331.2	4,063.3	6,405.2	11,131.5	20,173.3
1907	2,427.3	26,749.9	3,366.7	682.2	1,216.6	1,401.3	1,809.0	2,503.1	4,147.4	6,808.0	11,652.2	20,353.6
1908	2,243.1	24,754.1	3,034.4	685.3	1,113.5	1,276.5	1,607.5	2,461.0	3,864.1	6,423.2	10,506.6	18,403.8
1909	2,176.4	23,696.4	2,886.2	629.1	1,056.4	1,209.1	1,589.7	2,411.3	3,847.6	6,497.9	10,230.2	20,030.6
1910	2,172.4	23,975.1	2,834.0	567.6	1,025.9	1,216.9	1,702.8	2,460.3	3,852.1	6,429.3	10,320.1	17,636.2
1911	2,132.1	19,714.4	2,544.4	569.3	1,113.6	1,253.3	1,697.5	2,440.0	3,965.1	6,594.1	10,490.0	20,898.1
1912	2,056.1	18,272.0	2,155.9	497.3	1,061.0	1,199.0	1,678.9	2,489.6	4,064.0	6,568.4	11,279.9	19,508.0
1913	2,032.0	17,634.9	2,017.8	467.7	1,045.8	1,250.8	1,680.1	2,415.7	3,949.9	6,499.9	10,988.8	19,544.7
1914	2,016.9	16,992.0	1,914.1	457.0	1,042.6	1,270.2	1,684.5	2,443.1	3,998.9	6,536.6	10,917.8	20,452.6
1915	2,021.3	16,121.6	1,775.6	439.0	1,034.5	1,246.8	1,696.5	2,446.4	4,285.4	7,062.9	12,008.0	21,923.8
1916	1,914.7	15,252.5	1,782.5	380.6	996.4	1,288.6	1,664.5	2,277.2	3,945.5	6,600.3	12,093.9	23,377.2
1917	2,043.1	15,772.9	1,968.4	428.7	1,118.5	1,402.3	1,806.6	2,401.6	4,090.7	6,885.9	12,001.4	23,715.1
1918	2,558.6	16,771.7	2,835.2	697.8	1,910.7	2,259.3	2,237.0	2,620.1	4,062.8	6,656.2	11,270.6	21,543.8
1919	1,794.7	13,298.4	1,495.6	436.3	1,161.2	1,402.3	1,593.5	2,118.8	3,250.2	5,637.3	10,274.3	19,730.6
1920	1,767.5	14,915.7	1,458.2	379.3	1,040.9	1,283.7	1,520.3	2,147.7	3,315.6	6,024.6	11,122.5	23,229.3
1921	1,554.8	12,660.4	1,183.6	311.6	886.2	1,064.0	1,349.5	1,972.5	3,104.9	5,431.2	10,174.2	21,647.5
1922	1,524.6	11,737.4	1,068.3	271.9	832.2	1,049.7	1,411.7	2,118.8	3,172.4	5,607.2	10,357.7	23,375.7
1923	1,650.3	12,921.4	1,336.5	289.0	877.4	1,104.3	1,516.0	2,312.5	3,376.8	5,968.2	11,080.9	26,307.2
1924	1,708.1	13,407.7	1,257.5	292.1	925.9	1,114.7	1,595.6	2,476.1	3,494.3	6,257.5	11,282.2	26,590.5
1925	1,735.9	13,064.4	1,103.2	301.6	945.3	1,177.2	1,678.3	2,548.8	3,546.5	6,376.5	11,485.9	27,110.9
1926	1,777.4	12,963.4	1,247.6	293.6	917.7	1,221.0	1,720.6	2,688.6	3,742.4	6,520.4	11,677.2	25,751.6
1927	1,640.4	11,634.0	1,037.9	276.7	836.2	1,144.0	1,644.7	2,490.4	3,582.1	6,094.7	10,495.5	23,072.7
1928	1,706.6	12,117.3	1,049.1	283.5	864.9	1,205.2	1,720.0	2,649.9	3,866.4	6,505.7	11,385.5	23,924.2
1929	1,687.9	12,064.5	973.9	268.5	838.3	1,216.6	1,722.8	2,636.4	4,027.3	6,493.4	11,617.0	22,894.5
1930	1,633.0	11,000.0	932.3	260.9	800.5	1,158.9	1,612.2	2,580.2	4,084.3	6,387.2	10,421.0	20,489.4
1931	1,549.5	10,141.8	855.5	243.7	768.0	1,132.8	1,566.4	2,421.5	3,872.8	5,922.4	9,959.2	17,808.2
1932	1,445.8	9,457.3	688.8	215.9	672.4	1,014.1	1,450.1	2,292.8	3,813.2	5,569.4	9,784.7	18,193.1
1933	1,408.2	9,351.9	703.5	217.1	624.0	974.4	1,389.5	2,251.7	3,709.2	5,436.6	9,237.7	16,967.1
1934	1,475.5	10,392.1	808.2	215.5	627.0	997.2	1,440.3	2,369.0	3,778.7	5,496.3	9,533.0	17,470.5
1935	1,430.2	9,287.6	670.3	210.9	607.3	965.8	1,448.1	2,307.6	3,751.0	5,251.9	9,231.5	17,418.8
1936	1,538.1	9,830.9	659.5	205.2	651.3	1,031.6	1,560.9	2,474.2	4,033.2	5,784.7	10,110.4	18,671.1
1937	1,488.2	9,714.6	658.1	190.8	616.8	971.2	1,483.7	2,426.7	3,863.8	5,540.6	9,495.4	16,839.1
1938	1,403.0	9,208.2	625.6	182.6	564.8	870.2	1,335.8	2,306.6	3,675.4	5,205.7	8,929.2	15,441.8
1939	1,352.6	8,654.7	532.6	163.2	510.0	820.3	1,250.2	2,240.1	3,682.4	5,021.2	9,072.6	15,294.4

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	2,555.3	36,326.4	4,342.6	777.2	1,183.1	1,248.6	1,418.8	2,465.7	4,212.7	7,164.4	13,139.2	24,925.2
1901	2,559.3	33,272.2	3,597.0	720.9	1,156.7	1,404.6	1,739.1	2,704.0	4,513.6	6,975.7	11,933.0	24,079.3
1902	2,480.8	34,525.3	3,839.4	707.0	1,147.3	1,258.2	1,581.1	2,357.6	4,094.9	6,714.6	13,064.3	20,396.6
1903	2,552.3	32,752.1	4,155.6	633.8	1,150.0	1,411.4	1,566.4	2,433.6	4,526.6	7,968.7	12,710.2	22,379.6
1904	2,755.9	33,764.0	4,235.2	739.6	1,281.6	1,528.5	1,819.6	2,631.0	5,234.4	7,541.0	14,531.0	25,495.8
1905	2,681.5	33,943.4	4,111.6	689.0	1,326.9	1,430.8	1,737.7	2,814.3	4,544.2	7,617.5	13,099.9	21,348.3
1906	2,468.4	29,250.9	3,834.2	730.2	1,152.8	1,417.3	1,749.0	2,319.2	4,114.5	6,655.2	11,868.4	21,839.1
1907	2,500.9	28,652.1	3,523.9	652.8	1,203.5	1,501.5	1,834.6	2,511.7	4,213.7	7,085.0	12,796.1	19,697.0
1908	2,277.8	26,509.0	3,168.9	675.4	1,061.6	1,285.7	1,656.4	2,463.1	3,861.6	6,577.8	11,208.7	19,550.6
1909	2,226.3	25,272.3	2,957.7	572.9	1,041.5	1,238.9	1,635.3	2,504.0	3,763.1	6,778.7	11,465.8	22,362.1
1910	2,233.2	25,761.5	3,012.7	542.8	1,001.2	1,263.9	1,751.8	2,482.6	3,873.6	6,632.2	11,378.6	17,407.4
1911	2,193.6	21,398.2	2,671.1	548.2	1,068.6	1,304.7	1,753.4	2,454.1	4,014.5	7,012.1	10,962.4	20,957.7
1912	2,129.3	19,655.9	2,224.0	469.2	1,029.2	1,253.0	1,763.3	2,503.7	4,147.1	6,887.3	12,715.7	21,120.7
1913	2,099.3	19,298.2	2,094.2	437.1	1,009.0	1,294.6	1,743.9	2,419.7	3,908.9	6,825.9	11,964.9	20,130.2
1914	2,091.0	18,704.0	1,945.5	438.0	1,030.5	1,324.6	1,739.5	2,394.8	4,038.4	6,961.0	11,754.1	20,648.6
1915	2,083.3	17,923.7	1,812.1	428.7	1,001.0	1,285.3	1,774.7	2,369.7	4,229.9	7,201.5	12,586.1	23,447.7
1916	1,989.6	17,123.6	1,882.5	358.0	954.7	1,308.9	1,687.8	2,242.0	3,883.9	6,781.4	13,010.0	24,862.5
1917	2,143.4	17,646.7	2,049.5	410.9	1,114.6	1,481.1	1,894.6	2,356.3	3,993.2	7,176.3	12,795.4	24,413.0
1918	2,673.5	18,364.9	2,884.2	652.3	2,071.5	2,425.2	2,328.6	2,553.7	3,947.9	6,753.9	12,170.1	22,507.9
1919	1,805.3	14,878.7	1,562.2	410.9	1,117.7	1,317.1	1,542.7	1,990.1	3,051.1	5,618.0	10,817.1	20,897.1
1920	1,780.6	16,773.8	1,497.8	365.2	993.9	1,215.4	1,444.6	2,009.3	3,113.5	6,015.6	11,599.1	24,708.9
1921	1,569.5	13,971.4	1,219.5	310.7	831.9	1,029.9	1,307.6	1,842.4	2,900.7	5,336.1	10,752.7	22,683.1
1922	1,567.0	13,021.6	1,142.1	274.0	790.0	1,046.5	1,399.7	2,048.9	2,920.3	5,575.7	11,158.2	25,715.5
1923	1,704.7	14,319.7	1,390.7	283.5	834.2	1,126.8	1,532.1	2,249.2	3,147.9	5,930.2	12,050.3	29,690.2
1924	1,789.1	15,049.9	1,316.3	292.8	887.0	1,151.9	1,619.9	2,430.3	3,420.7	6,277.0	12,437.5	31,125.6
1925	1,816.2	14,651.6	1,444.8	310.6	887.9	1,218.2	1,698.8	2,567.9	3,453.2	6,460.6	12,454.6	31,761.4
1926	1,869.3	14,647.6	1,307.9	298.6	874.7	1,259.5	1,759.2	2,713.4	3,689.0	6,587.5	12,816.2	29,724.6
1927	1,718.9	13,121.9	1,101.3	285.9	787.7	1,161.0	1,671.3	2,504.2	3,526.5	6,261.6	11,473.1	26,151.4
1928	1,798.5	13,689.0	1,106.8	287.2	826.8	1,239.0	1,758.7	2,694.7	3,793.2	6,753.7	12,586.2	26,749.5
1929	1,795.9	13,676.7	1,028.3	282.3	813.3	1,260.2	1,803.9	2,695.7	3,966.9	6,854.0	13,036.6	25,917.5
1930	1,739.0	12,226.1	996.3	266.4	781.2	1,213.5	1,699.9	2,633.3	4,039.4	6,726.5	11,763.4	22,845.4
1931	1,653.6	11,335.7	910.4	251.5	759.7	1,180.7	1,674.3	2,462.1	3,855.1	6,165.7	11,388.7	20,172.5
1932	1,541.0	10,528.5	724.1	225.4	662.5	1,055.3	1,545.0	2,331.7	3,784.1	5,826.3	11,046.3	20,692.9
1933	1,510.3	10,394.5	744.6	224.9	633.0	1,024.9	1,512.1	2,303.4	3,717.1	5,627.8	10,159.2	19,333.0
1934	1,603.1	11,649.8	879.7	221.4	635.3	1,071.5	1,576.4	2,499.0	3,817.5	5,825.2	10,616.1	19,683.2
1935	1,559.0	10,479.0	710.0	223.3	614.3	1,037.9	1,603.3	2,445.0	3,800.1	5,588.6	10,492.0	19,277.1
1936	1,693.3	11,080.4	707.8	219.7	659.5	1,129.1	1,745.6	2,685.8	4,130.2	6,248.6	11,311.5	21,112.8
1937	1,636.0	10,904.3	710.4	198.5	630.7	1,054.4	1,629.3	2,611.1	3,977.3	6,080.5	10,900.7	18,861.2
1938	1,513.6	10,397.4	668.2	190.4	556.0	925.0	1,432.9	2,446.3	3,736.3	5,645.6	9,854.4	17,402.9
1939	1,470.2	9,863.3	573.2	177.3	498.8	874.7	1,341.8	2,377.9	3,763.4	5,413.0	10,225.0	16,983.7

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL CAUSES

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	2,443.8	29,945.4	4,350.0	1,006.0	1,119.6	1,173.3	1,558.0	2,390.1	4,212.7	6,642.1	11,315.8	19,582.7
1901	2,311.8	28,721.8	3,375.8	752.2	1,121.6	1,180.0	1,559.0	2,502.2	4,013.5	6,022.1	10,746.0	21,794.9
1902	2,234.1	27,309.5	3,636.7	721.0	1,135.7	1,165.5	1,508.7	2,240.9	3,477.7	6,394.0	9,302.3	20,255.2
1903	2,343.3	27,488.7	3,742.1	802.8	1,095.8	1,292.6	1,633.0	2,276.4	4,074.7	6,708.7	9,492.0	24,720.9
1904	2,468.4	29,376.9	3,730.6	873.8	1,270.3	1,294.6	1,645.4	2,431.8	4,544.6	6,831.3	11,246.8	20,886.1
1905	2,427.5	28,096.8	3,636.6	930.1	1,292.5	1,347.8	1,553.0	2,541.8	4,162.3	7,032.9	9,540.6	20,000.0
1906	2,360.1	24,906.9	3,453.8	825.9	1,274.1	1,326.6	1,623.3	2,347.4	3,994.6	6,126.4	10,540.0	19,165.4
1907	2,345.7	24,806.8	3,216.8	710.3	1,229.3	1,284.8	1,775.3	2,491.5	4,058.5	6,496.1	10,726.8	20,753.3
1908	2,203.9	22,951.6	2,906.2	694.8	1,165.6	1,265.5	1,542.1	2,458.1	3,867.5	6,247.9	9,925.0	17,676.4
1909	2,120.0	22,079.0	2,817.8	683.1	1,071.3	1,173.1	1,529.2	2,284.7	3,960.8	6,170.9	9,163.3	18,504.4
1910	2,102.8	22,142.8	2,662.3	591.5	1,051.0	1,159.5	1,637.3	2,429.2	3,822.9	6,189.2	9,391.2	17,787.3
1911	2,063.5	18,019.6	2,421.4	589.8	1,158.5	1,193.5	1,625.4	2,421.1	3,901.0	6,091.3	10,073.7	20,857.9
1912	1,974.4	16,883.9	2,089.7	524.6	1,092.7	1,136.7	1,570.1	2,470.4	3,955.6	6,200.8	9,994.7	18,416.7
1913	1,959.6	15,982.3	1,943.2	497.6	1,081.5	1,203.0	1,602.3	2,410.5	4,001.3	6,131.5	10,092.0	19,138.0
1914	1,937.1	15,295.0	1,883.5	475.7	1,054.4	1,211.4	1,617.4	2,507.7	3,949.1	6,054.2	10,133.3	20,315.6
1915	1,954.5	14,339.9	1,740.0	449.0	1,066.9	1,205.5	1,601.0	2,550.6	4,356.2	6,905.5	11,460.1	20,860.6
1916	1,837.5	13,410.0	1,683.7	402.8	1,035.0	1,268.4	1,638.5	2,323.5	4,024.2	6,388.3	11,203.5	22,308.1
1917	1,940.3	13,929.6	1,888.2	446.2	1,122.0	1,326.0	1,710.1	2,461.5	4,216.3	6,546.2	11,234.1	23,208.4
1918	2,441.9	15,207.4	2,786.8	742.9	1,767.9	2,103.0	2,136.7	2,709.5	4,210.9	6,542.9	10,390.9	20,856.0
1919	1,784.0	11,741.4	1,429.5	461.5	1,205.0	1,481.2	1,647.9	2,294.8	3,511.2	5,660.1	9,734.1	18,879.4
1920	1,754.2	13,107.5	1,418.7	393.3	1,083.5	1,346.8	1,603.0	2,335.3	3,579.6	6,035.3	10,639.1	22,121.6
1921	1,539.9	11,368.4	1,147.7	312.5	936.0	1,095.5	1,395.0	2,145.9	3,371.1	5,545.5	9,586.7	20,910.7
1922	1,481.8	10,478.7	996.1	269.9	870.8	1,052.6	1,424.4	2,209.3	3,503.7	5,645.6	9,545.2	21,767.5
1923	1,595.4	11,546.5	1,282.0	294.5	917.5	1,083.9	1,499.1	2,392.2	3,676.7	6,014.0	10,103.1	24,049.2
1924	1,626.7	11,792.0	1,198.0	291.5	962.1	1,080.5	1,570.4	2,532.4	3,590.6	6,233.6	10,122.7	23,598.7
1925	1,655.0	11,499.3	1,061.1	292.8	998.7	1,139.0	1,657.1	2,525.4	3,667.5	6,275.4	10,524.2	24,020.8
1926	1,685.0	11,308.5	1,186.5	288.7	958.0	1,184.8	1,680.8	2,659.0	3,811.7	6,440.1	10,550.8	23,070.6
1927	1,561.6	10,199.2	974.0	267.7	881.3	1,128.1	1,617.5	2,474.1	3,654.8	5,896.3	9,528.6	20,923.5
1928	1,615.2	10,594.3	991.2	279.8	900.1	1,174.0	1,681.7	2,597.6	3,963.2	6,215.3	10,206.0	21,897.0
1929	1,580.5	10,512.0	919.5	254.9	861.2	1,176.0	1,643.0	2,567.4	4,106.9	6,079.5	10,242.0	20,657.6
1930	1,528.0	9,787.6	868.9	255.3	818.0	1,108.1	1,526.3	2,518.5	4,142.8	6,003.1	9,144.8	18,716.1
1931	1,446.5	8,966.5	801.2	236.0	775.6	1,088.2	1,461.0	2,375.0	3,895.6	5,648.7	8,603.3	16,031.9
1932	1,351.8	8,400.7	653.9	206.6	681.3	975.7	1,357.7	2,248.6	3,850.3	5,282.1	8,590.4	16,318.8
1933	1,307.8	8,326.1	663.0	209.3	615.8	927.6	1,270.6	2,193.5	3,699.2	5,223.5	8,362.7	15,198.1
1934	1,350.2	9,157.8	737.3	209.6	619.4	928.4	1,308.7	2,224.3	3,730.5	5,131.0	8,505.6	15,820.4
1935	1,304.0	8,118.1	630.9	198.6	600.9	899.3	1,298.5	2,156.0	3,690.9	4,880.4	8,036.6	16,036.7
1936	1,386.4	8,610.5	611.6	190.8	643.8	942.0	1,383.2	2,242.6	3,915.9	5,275.3	8,972.3	16,860.5
1937	1,344.0	8,555.2	606.3	183.1	604.2	895.0	1,344.1	2,226.3	3,728.4	4,950.0	8,163.6	15,344.2
1938	1,289.5	8,051.0	583.4	174.8	572.9	820.3	1,243.0	2,155.8	3,603.8	4,726.3	8,052.2	13,996.6
1939	1,238.3	7,477.0	492.4	149.3	520.2	770.9	1,162.8	2,092.4	3,588.3	4,595.4	7,979.6	14,055.5

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ICD Revision	Years in use	ICD Nos.
Fifth	1939	13-22
Fourth	1930-1938	23-32
Third	1921-1929	31-37
Second	1910-1920	28-35
First	1900-1909	26-35

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	194.4	311.6	101.8	36.2	205.7	294.3	253.6	215.6	223.0	256.1	279.3	204.5
1901	189.9	292.7	94.3	35.1	200.8	284.0	255.5	210.1	221.3	254.1	250.3	195.7
1902	174.2	256.3	88.1	33.1	183.9	262.0	236.0	193.9	198.3	219.8	229.7	188.1
1903	177.2	255.8	92.8	34.3	189.3	266.1	234.5	195.5	205.4	220.2	223.2	159.3
1904	188.1	287.5	96.6	37.6	203.4	279.7	254.7	208.1	207.6	224.8	207.7	173.9
1905	179.9	259.4	101.2	36.7	190.6	263.7	240.1	208.3	200.6	220.9	201.5	149.4
1906	175.8	259.3	95.9	36.2	182.0	254.4	237.2	209.0	206.7	220.0	210.9	170.4
1907	174.2	262.2	93.6	34.6	175.2	246.0	239.9	213.0	208.6	229.3	230.5	149.6
1908	162.1	226.2	87.9	33.7	164.7	231.5	223.5	194.1	195.9	209.9	198.5	137.9
1909	156.3	225.3	85.2	31.7	159.2	221.4	210.9	189.1	189.2	207.9	203.1	135.7
1910	153.8	212.9	84.6	29.7	152.0	217.6	214.9	188.1	192.9	203.5	189.2	115.8
1911	155.1	206.0	83.5	31.3	157.5	216.7	215.4	185.7	198.8	218.7	194.4	149.0
1912	145.4	184.2	75.6	28.4	147.8	206.5	202.5	179.4	181.4	194.7	179.7	140.9
1913	143.5	184.9	78.2	29.0	144.1	199.3	199.8	178.6	186.7	198.4	184.8	117.0
1914	141.7	171.3	67.8	28.6	144.8	199.2	201.9	177.5	184.7	185.3	172.0	107.4
1915	140.1	168.6	68.0	26.9	146.1	198.0	196.9	176.8	177.0	187.8	167.5	117.6
1916	138.4	163.1	67.1	27.6	148.0	198.2	193.9	167.3	177.4	194.4	170.4	131.2
1917	143.5	167.7	69.2	28.3	155.9	205.9	199.9	178.4	181.3	193.7	179.7	144.8
1918	149.8	144.9	64.3	30.3	182.3	231.4	197.8	178.2	176.2	184.7	155.6	121.2
1919	125.6	121.3	50.6	24.0	147.7	187.0	170.3	155.1	154.9	170.4	152.9	113.2
1920	113.1	106.5	45.4	22.4	136.1	164.9	147.4	137.2	141.3	163.8	157.7	132.3
1921	97.6	91.3	37.1	18.9	118.4	139.7	125.3	120.6	127.8	144.5	142.6	104.3
1922	95.3	84.2	35.1	16.1	113.9	139.5	123.8	118.0	125.4	151.6	143.3	108.2
1923	91.7	78.0	35.1	15.9	109.7	131.7	119.3	114.3	120.1	144.1	142.8	117.5
1924	87.9	73.1	33.6	15.7	107.7	126.3	113.8	110.0	111.8	129.8	132.9	100.4
1925	84.8	69.7	30.3	14.8	101.7	122.5	110.6	106.4	108.2	132.7	125.7	98.2
1926	85.5	66.8	34.0	14.6	100.4	122.0	111.6	108.3	110.4	132.9	138.2	107.3
1927	79.6	58.4	29.7	13.5	93.8	116.7	103.1	100.1	100.8	121.5	124.2	97.0
1928	78.3	58.4	28.7	13.1	90.7	112.4	103.7	99.9	102.7	121.7	126.5	93.0
1929	75.3	47.5	26.9	11.9	85.5	111.7	98.5	96.9	98.9	115.7	121.6	89.1
1930	71.1	51.6	25.9	11.9	77.3	102.8	92.4	93.2	97.0	111.7	115.5	76.9
1931	67.8	45.6	24.4	10.8	71.7	96.8	89.9	90.4	93.0	106.0	110.0	83.6
1932	62.5	41.9	20.1	9.8	62.2	87.4	83.6	84.5	89.2	102.8	110.9	74.4
1933	59.6	39.8	18.9	8.8	57.5	83.8	79.1	82.3	88.7	95.4	103.9	66.5
1934	56.7	37.2	18.4	8.2	52.1	77.8	74.7	81.2	85.3	92.6	103.7	69.7
1935	55.1	35.5	16.3	7.9	48.9	74.1	75.5	77.9	83.8	90.5	101.7	64.1
1936	55.9	34.4	15.9	7.6	49.8	74.3	75.4	79.2	85.6	92.2	99.1	84.0
1937	53.8	32.1	15.9	7.0	46.9	70.4	71.8	78.3	82.9	90.0	93.4	62.3
1938	49.1	28.4	15.6	6.5	43.2	62.9	64.4	69.1	78.2	83.0	82.2	67.4
1939	47.1	26.6	13.6	6.0	39.4	58.7	60.6	69.7	75.4	82.3	85.0	67.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	201.0	338.3	107.2	29.0	187.4	291.0	275.9	255.6	267.4	282.4	273.4	269.4
1901	201.3	318.5	99.4	29.2	185.5	288.8	286.7	257.9	268.9	285.8	263.0	221.2
1902	183.5	281.5	87.5	28.1	163.1	267.3	268.1	234.3	245.5	236.0	251.3	211.9
1903	184.9	273.6	94.9	28.2	168.9	265.9	261.6	242.0	247.5	242.2	231.3	153.0
1904	197.4	317.3	97.2	29.3	177.9	283.2	295.3	254.6	249.3	248.4	215.9	197.3
1905	190.8	280.3	103.0	29.5	169.3	268.1	279.5	261.4	243.5	235.7	216.8	155.4
1906	187.7	282.3	102.8	28.8	163.3	255.8	273.3	266.1	251.7	251.5	230.6	189.8
1907	188.8	284.1	98.1	29.5	159.4	253.0	278.0	272.2	257.3	254.9	246.6	147.9
1908	177.3	241.3	92.4	28.9	153.7	240.0	262.0	243.7	246.2	239.3	226.6	145.8
1909	169.5	250.2	89.4	26.8	148.5	227.7	245.4	233.3	232.5	234.7	217.0	154.8
1910	167.1	223.8	88.1	24.4	140.7	220.3	250.7	236.9	243.1	232.1	195.0	132.8
1911	168.2	222.9	85.7	26.4	144.2	221.0	250.0	232.8	250.7	249.2	210.8	165.5
1912	159.6	195.8	80.0	23.7	138.1	213.0	239.3	228.8	228.0	221.5	181.9	156.0
1913	158.7	198.6	81.6	24.3	134.5	210.5	236.4	229.4	235.1	225.9	185.3	129.8
1914	158.5	197.2	70.7	23.9	136.0	211.9	244.1	228.7	232.3	211.4	184.2	108.1
1915	155.9	184.8	68.5	22.4	134.8	212.0	233.8	229.5	224.0	218.6	175.3	115.2
1916	152.9	181.7	70.8	23.6	132.4	211.0	230.8	215.9	226.3	225.3	166.7	113.1
1917	158.7	187.0	71.5	23.9	139.1	219.8	241.7	230.2	230.2	220.4	177.6	146.5
1918	167.7	157.1	67.0	25.1	173.6	258.3	239.2	230.0	224.8	212.9	159.6	131.2
1919	134.2	132.8	53.4	20.1	127.0	189.6	198.1	195.0	192.9	193.3	156.0	118.0
1920	116.6	115.9	48.0	18.4	111.7	161.6	162.7	165.9	172.0	188.0	165.2	139.1
1921	101.1	96.1	38.6	16.0	97.0	136.7	140.5	147.2	156.8	164.0	144.9	105.4
1922	99.5	89.6	37.4	13.3	92.6	138.2	139.5	144.6	157.9	176.4	149.6	111.6
1923	95.4	83.0	36.1	14.0	85.6	130.5	137.2	142.9	147.6	163.2	143.7	115.4
1924	91.5	78.6	35.4	13.3	83.8	125.3	130.5	136.1	140.0	151.7	137.5	108.4
1925	88.0	74.7	31.9	12.7	78.8	119.2	127.5	133.4	132.6	154.6	133.9	101.8
1926	89.1	72.7	35.1	12.8	77.5	117.9	130.4	136.7	137.3	150.6	145.5	117.2
1927	82.9	59.8	30.3	11.8	71.0	113.3	121.2	127.1	126.2	140.9	133.6	94.4
1928	82.5	62.4	28.4	11.5	70.5	110.3	121.7	128.4	129.1	138.9	137.1	79.8
1929	79.3	49.7	28.3	10.2	66.2	107.7	117.2	123.9	124.3	133.5	124.3	104.3
1930	76.2	54.9	26.7	10.4	61.3	102.0	109.6	120.5	124.0	130.9	120.9	78.6
1931	73.5	51.8	25.8	9.2	57.2	95.1	110.5	117.3	117.6	123.6	117.8	92.7
1932	68.0	44.7	20.1	8.6	48.6	86.6	101.7	110.3	115.8	120.8	118.5	67.6
1933	65.4	42.0	19.0	7.7	45.6	81.1	97.0	109.7	115.3	113.2	113.2	69.2
1934	63.1	40.1	20.0	7.0	41.6	76.6	92.0	109.6	111.7	110.9	111.0	82.2
1935	62.1	38.8	16.8	6.9	38.8	74.1	94.1	106.6	111.4	108.5	108.8	60.7
1936	63.4	34.8	16.8	6.8	39.0	73.5	94.1	110.2	115.4	113.4	112.5	89.3
1937	61.3	34.1	15.5	6.0	37.0	69.1	89.8	108.3	114.3	111.0	105.0	66.7
1938	56.0	28.7	16.4	5.5	34.2	61.2	79.5	97.4	107.5	104.2	90.5	77.9
1939	54.0	28.3	13.6	5.1	30.6	56.4	75.5	98.4	104.0	104.2	94.9	76.0

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	187.8	284.3	96.3	43.5	222.8	297.7	229.6	174.0	179.4	231.2	284.8	157.6
1901	178.5	266.2	89.2	41.0	215.1	279.0	222.3	160.2	174.6	223.9	238.7	177.1
1902	164.9	230.7	88.7	38.1	203.5	256.6	202.0	151.5	152.0	204.5	210.1	170.7
1903	169.4	237.7	90.7	40.5	208.7	266.4	205.6	146.6	164.1	199.3	216.1	163.9
1904	178.7	257.0	96.0	45.9	227.8	276.2	211.6	159.1	166.6	202.5	200.4	156.6
1905	169.0	238.1	99.3	43.9	211.1	259.1	198.1	152.3	158.1	207.0	188.0	145.0
1906	163.5	235.7	88.9	43.7	202.6	252.9	197.4	146.4	150.3	189.4	193.1	155.8
1907	159.1	239.8	89.1	39.7	190.9	238.3	198.0	148.0	158.2	204.3	216.0	150.9
1908	146.2	210.8	83.2	38.6	175.9	222.3	180.6	138.7	142.8	180.8	172.9	131.8
1909	142.3	199.7	81.0	36.6	170.0	214.6	172.6	140.0	143.3	181.3	190.5	120.8
1910	139.8	201.8	81.2	35.2	163.7	214.6	174.7	133.3	139.2	174.9	184.0	102.5
1911	141.3	188.7	81.3	36.2	171.1	211.9	176.8	132.6	142.7	188.0	179.4	136.0
1912	130.3	172.2	71.1	33.2	157.6	199.3	161.5	123.7	130.9	167.7	177.7	129.0
1913	127.4	170.9	74.7	33.8	153.7	187.3	159.1	121.2	134.1	170.6	184.3	107.1
1914	123.9	144.8	64.9	33.3	153.6	185.4	155.1	119.6	132.7	158.6	160.7	106.9
1915	123.5	151.9	67.4	31.5	157.5	182.9	156.0	117.1	125.5	156.3	160.3	119.4
1916	123.3	144.2	63.3	31.8	163.5	184.6	153.3	112.1	123.7	162.9	173.7	145.0
1917	127.7	147.8	66.9	32.7	172.4	191.3	154.2	119.7	127.6	166.3	181.6	143.5
1918	131.8	132.4	61.5	35.6	189.8	206.0	152.6	119.2	122.7	155.6	152.0	113.7
1919	116.9	109.5	47.7	27.9	167.2	184.4	140.0	109.6	113.0	146.6	150.0	109.6
1920	109.5	96.8	42.8	26.3	159.9	168.3	130.5	104.7	107.6	138.6	150.9	127.2
1921	94.1	86.4	35.4	21.8	139.3	142.8	108.7	90.8	96.1	124.4	140.4	103.4
1922	91.0	78.6	32.8	19.0	134.7	140.7	106.8	88.3	90.2	135.0	137.5	105.6
1923	87.9	72.8	34.1	17.9	133.3	133.0	99.9	82.8	90.2	124.3	142.0	119.1
1924	84.3	67.4	31.7	18.2	131.2	127.3	95.8	81.4	81.1	107.0	128.7	94.3
1925	81.4	64.5	28.7	17.0	124.3	125.7	92.5	76.9	81.6	110.0	118.1	95.3
1926	81.7	60.7	32.8	16.4	122.9	126.1	91.4	77.5	81.0	114.6	131.5	99.7
1927	76.1	56.9	29.1	15.3	116.3	120.2	83.8	70.8	72.9	101.4	115.6	99.0
1928	74.1	54.3	29.0	14.7	110.6	114.5	84.7	68.8	73.8	103.8	116.6	103.2
1929	71.3	45.2	25.5	13.7	104.4	115.6	78.7	67.5	71.2	97.4	119.1	77.6
1930	65.9	48.1	25.1	13.5	92.9	103.5	74.4	63.3	67.9	91.9	110.4	75.6
1931	62.0	39.2	23.0	12.4	85.9	98.4	68.4	61.1	66.4	87.8	102.6	76.7
1932	56.8	39.0	20.1	11.0	75.6	88.3	64.8	56.5	60.6	84.4	103.8	79.5
1933	53.7	37.6	18.8	9.9	69.3	86.5	60.6	52.8	60.1	77.1	95.3	68.4
1934	50.1	34.2	16.7	9.5	62.5	78.9	56.7	50.6	57.1	73.8	96.8	60.2
1935	47.9	32.1	15.8	8.9	58.9	74.1	56.3	47.0	54.3	72.1	95.1	66.6
1936	48.2	33.9	15.0	8.5	60.4	75.0	56.1	46.0	53.8	70.7	86.7	80.0
1937	46.2	30.0	16.2	8.1	56.5	71.7	53.3	46.3	49.5	68.8	82.7	59.0
1938	42.1	28.0	14.8	7.6	52.0	64.5	49.0	38.9	47.2	61.8	74.5	59.6
1939	40.1	24.9	13.6	7.0	48.0	60.9	45.5	39.1	45.3	60.5	76.0	60.6

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	145.9	...	...	...	...	...	...	...	...	...	...	...
1911	145.0	...	...	...	...	...	...	...	...	...	...	...
1912	136.0	...	...	...	...	...	...	...	...	...	...	...
1913	132.6	...	...	...	...	...	...	...	...	...	...	...
1914	130.3	159.9	61.1	23.1	127.1	183.8	189.2	168.2	176.8	178.6	165.7	101.0
1915	128.5	158.1	60.8	21.5	128.3	182.5	183.1	166.7	168.8	181.2	163.3	107.7
1916	125.7	154.9	60.9	21.5	126.2	179.7	178.4	157.1	170.1	186.2	164.1	122.6
1917	129.6	...	...	...	...	...	...	...	...	...	...	...
1918	134.3	136.9	56.4	23.0	151.3	209.1	181.7	166.4	167.8	176.9	147.9	110.5
1919	110.9	112.7	45.7	17.8	117.9	163.4	154.2	143.8	147.6	163.5	147.8	99.2
1920	99.5	98.1	41.2	16.9	108.1	144.1	132.6	126.0	134.4	156.7	152.8	124.2
1921	84.7	84.1	32.7	14.4	92.1	118.6	111.4	109.7	121.5	139.7	136.9	92.5
1922	82.6	77.5	30.9	12.2	88.5	118.2	108.1	107.5	118.9	145.6	138.4	99.7
1923	79.5	70.5	30.2	12.0	85.3	111.3	104.7	104.2	113.8	140.8	140.6	105.0
1924	74.9	65.3	28.6	11.2	81.9	105.4	99.4	98.4	105.0	124.9	128.1	88.9
1925	71.6	63.4	24.6	10.4	76.0	100.4	95.7	94.9	101.7	128.6	120.6	91.5
1926	72.0	59.1	27.7	10.0	75.1	98.9	96.0	97.3	103.4	128.6	135.1	97.8
1927	66.5	49.3	24.0	9.2	69.6	93.7	87.6	89.5	94.7	118.1	120.2	85.3
1928	64.9	48.7	23.3	8.6	65.1	88.8	88.2	88.4	96.8	117.7	121.6	86.9
1929	62.4	40.8	21.0	8.0	61.5	88.3	83.4	85.8	92.5	112.1	116.9	81.7
1930	57.7	42.4	19.7	7.3	53.1	78.6	76.9	82.1	91.2	106.7	111.6	73.2
1931	54.2	38.4	18.8	6.6	47.2	72.2	73.4	79.0	85.8	101.1	106.1	79.2
1932	49.0	33.0	13.7	5.4	39.5	63.5	66.4	72.7	81.8	97.6	107.4	69.8
1933	46.6	30.0	13.0	4.9	36.1	60.6	62.9	70.5	80.5	90.1	99.8	63.5
1934	44.5	26.6	12.5	4.5	31.9	55.8	59.4	70.0	78.2	88.4	101.3	63.8
1935	44.9	27.0	12.4	4.9	32.4	55.5	61.1	68.3	77.8	87.1	101.2	61.8
1936	45.0	27.6	12.4	4.6	31.8	54.4	60.1	69.0	79.0	89.2	98.4	80.2
1937	43.4	25.0	11.4	4.1	30.0	52.1	56.6	68.9	76.5	87.0	92.0	59.1
1938	39.1	21.8	11.4	4.0	26.3	45.2	50.8	59.9	71.9	79.9	80.6	64.5
1939	37.7	19.3	9.7	3.5	23.9	41.8	47.7	60.4	70.0	80.0	83.8	68.9

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	158.2	---	---	---	---	---	---	---	---	---	---	---
1911	157.5	---	---	---	---	---	---	---	---	---	---	---
1912	149.4	---	---	---	---	---	---	---	---	---	---	---
1913	147.7	---	---	---	---	---	---	---	---	---	---	---
1914	146.9	182.6	64.3	19.6	119.5	196.8	230.0	218.5	221.3	201.3	173.6	103.5
1915	144.0	171.8	61.6	17.8	119.3	196.5	218.3	217.9	212.8	209.5	168.1	99.8
1916	141.3	174.6	64.7	18.7	114.1	194.4	216.2	205.5	217.2	212.7	157.9	102.8
1917	145.8	---	---	---	---	---	---	---	---	---	---	---
1918	153.2	149.1	58.5	19.1	145.3	237.0	223.5	217.4	215.0	202.5	150.9	123.7
1919	121.1	125.2	48.5	15.2	101.1	169.1	182.9	184.5	184.5	184.1	150.0	106.0
1920	104.1	108.3	43.9	14.7	86.0	142.5	147.8	154.7	164.7	178.8	159.9	120.0
1921	89.1	88.3	34.2	12.6	74.2	116.5	126.4	136.2	149.9	158.6	136.0	85.5
1922	87.5	82.4	32.6	10.4	69.9	118.1	124.1	133.5	151.4	167.9	140.2	95.3
1923	84.4	76.0	30.8	10.9	65.7	111.2	122.5	132.5	141.6	159.2	140.5	101.6
1924	79.3	71.6	30.5	9.8	61.6	104.5	115.8	124.3	132.7	145.9	130.3	91.5
1925	75.8	67.4	26.0	9.2	57.0	97.6	112.6	121.9	126.1	149.5	126.6	90.4
1926	76.4	65.0	28.7	9.0	56.3	95.2	113.7	125.4	129.1	145.0	140.4	105.6
1927	70.7	50.8	24.5	8.2	51.0	91.0	104.8	115.5	119.4	136.3	127.4	83.5
1928	69.7	53.0	22.4	8.0	48.3	86.8	105.7	116.1	122.5	134.0	130.1	68.7
1929	67.1	42.6	22.1	7.0	45.7	85.1	101.3	112.3	117.1	128.2	117.2	93.0
1930	63.4	46.0	20.4	6.7	40.3	77.3	93.1	108.5	118.1	124.0	114.9	74.5
1931	60.1	42.8	19.5	5.9	36.1	70.3	91.1	105.0	109.3	117.2	113.1	85.7
1932	54.6	35.4	13.5	5.1	29.7	61.4	81.9	96.7	107.4	113.5	112.5	62.0
1933	52.4	30.7	13.3	4.6	27.4	57.8	77.5	96.0	105.5	106.2	107.1	65.7
1934	50.9	28.7	13.6	4.2	24.3	54.2	73.9	96.6	103.4	104.8	106.3	70.4
1935	51.7	28.7	13.0	4.3	25.2	54.8	77.2	94.8	104.1	103.7	107.2	56.9
1936	52.2	28.7	13.1	4.3	24.2	52.2	75.5	97.5	107.4	109.7	110.2	81.9
1937	50.9	25.2	11.3	3.8	23.0	50.4	71.5	97.5	106.6	106.9	102.0	60.3
1938	46.2	22.6	12.0	3.7	20.3	43.7	64.3	86.1	99.8	99.7	88.0	73.9
1939	44.7	20.3	9.7	3.2	18.1	39.5	60.0	87.0	97.7	101.3	93.0	77.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.  
 Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	132.8	...	...	...	...	...	...	...	...	...	...	...
1911	131.9	...	...	...	...	...	...	...	...	...	...	...
1912	121.8	...	...	...	...	...	...	...	...	...	...	...
1913	116.7	...	...	...	...	...	...	...	...	...	...	...
1914	112.9	136.7	57.9	26.7	134.7	169.9	144.1	111.6	128.4	155.5	158.4	98.9
1915	112.2	144.1	60.0	25.1	137.3	167.5	144.4	109.2	120.9	152.6	158.8	113.8
1916	109.5	134.6	56.9	24.3	138.2	164.0	136.9	102.7	118.8	159.3	169.7	137.7
1917	112.7	...	...	...	...	...	...	...	...	...	...	...
1918	115.4	124.3	54.3	27.0	156.6	182.8	136.1	108.9	116.2	150.7	145.3	100.5
1919	100.4	99.9	43.0	20.5	134.0	157.7	122.9	98.0	107.3	142.3	145.8	94.1
1920	94.8	87.7	38.4	19.1	129.7	145.7	115.8	94.0	101.4	133.9	146.3	127.3
1921	80.2	79.7	31.2	16.2	109.8	120.7	95.0	80.4	90.8	120.3	137.8	97.7
1922	77.4	72.4	29.1	14.0	106.8	118.2	90.8	79.0	84.3	122.8	136.7	103.0
1923	74.5	64.8	29.5	13.1	104.7	111.5	85.3	73.4	83.9	122.0	140.7	107.6
1924	70.4	58.9	26.5	12.7	102.2	106.2	81.5	70.4	75.1	103.3	126.1	86.8
1925	67.2	59.3	23.0	11.7	95.0	103.2	77.4	65.7	75.4	107.2	115.0	92.4
1926	67.5	52.9	26.6	11.0	93.8	102.5	77.0	67.0	75.7	111.7	130.2	91.8
1927	62.2	47.8	23.5	10.3	88.0	96.4	69.0	61.6	68.1	99.5	113.5	86.7
1928	59.9	44.2	24.1	9.2	81.7	90.9	69.5	58.5	69.2	100.9	113.6	100.9
1929	57.6	38.9	19.7	9.0	77.2	91.5	64.4	57.0	66.1	95.6	116.7	73.0
1930	51.9	38.6	19.0	7.8	65.7	79.9	59.8	53.4	62.6	88.8	108.6	72.2
1931	48.2	33.8	18.0	7.2	58.1	74.1	54.6	51.0	60.9	84.6	99.5	74.4
1932	43.2	30.4	13.9	5.7	49.3	65.6	50.0	46.8	54.6	81.3	102.7	75.7
1933	40.7	29.3	12.8	5.3	44.7	63.4	47.5	43.0	54.0	73.7	92.9	61.9
1934	38.0	24.5	11.4	4.8	39.3	57.5	44.3	41.5	51.6	71.7	96.6	58.8
1935	37.8	25.1	11.8	5.5	39.5	56.2	44.5	39.8	50.0	70.4	95.7	65.6
1936	37.6	26.5	11.6	4.9	39.3	56.6	44.1	38.6	49.1	68.6	87.4	79.0
1937	35.8	24.7	11.5	4.5	36.9	53.8	41.3	38.4	44.9	67.0	82.7	58.1
1938	31.9	20.9	10.7	4.4	32.3	46.6	36.9	32.0	42.5	60.2	73.8	57.4
1939	30.6	18.3	9.7	3.9	29.7	44.1	35.2	32.2	40.9	58.8	75.4	62.6

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	445.5	---	---	---	---	---	---	---	---	---	---	---
1911	461.4	---	---	---	---	---	---	---	---	---	---	---
1912	429.0	---	---	---	---	---	---	---	---	---	---	---
1913	386.5	---	---	---	---	---	---	---	---	---	---	---
1914	396.7	442.3	234.5	154.8	519.8	506.4	459.6	398.3	415.9	397.0	387.6	*
1915	401.1	417.1	248.1	153.1	522.0	508.4	472.3	413.9	420.2	394.9	312.5	*
1916	322.7	274.1	153.3	106.4	431.8	470.1	423.4	340.3	331.8	374.3	322.1	252.2
1917	332.6	---	---	---	---	---	---	---	---	---	---	---
1918	346.0	242.6	164.2	113.6	516.2	510.0	404.8	350.2	339.3	335.4	314.9	248.3
1919	284.0	209.2	102.1	81.5	422.9	447.5	346.2	292.9	276.9	285.3	241.3	250.6
1920	262.4	201.4	90.2	75.5	398.6	396.9	309.9	278.3	257.2	286.4	247.7	216.4
1921	239.3	171.5	83.2	62.7	366.3	371.9	279.1	258.1	232.3	229.3	243.9	229.0
1922	218.9	148.7	74.2	49.2	324.4	344.6	282.1	235.7	218.7	246.1	221.0	187.5
1923	213.1	151.7	81.0	49.8	317.5	330.5	273.1	232.6	213.2	198.8	178.7	239.8
1924	218.6	150.0	79.8	56.1	330.1	329.7	269.2	246.9	212.9	214.1	211.6	213.6
1925	221.3	133.0	84.5	55.8	330.3	338.8	274.9	244.7	205.6	206.7	211.1	164.4
1926	223.8	141.5	92.8	57.2	325.1	345.9	282.2	240.0	215.5	212.6	191.5	202.3
1927	208.7	141.9	80.1	51.9	301.5	329.7	268.3	221.1	189.1	182.1	191.3	211.5
1928	199.5	141.9	71.2	49.4	290.7	314.3	255.9	217.8	180.7	188.2	202.4	147.7
1929	192.0	106.7	73.0	43.7	273.5	311.2	245.2	211.0	185.6	177.4	195.7	156.5
1930	192.0	125.5	74.4	49.5	269.5	308.6	242.1	206.2	177.8	198.5	178.7	110.9
1931	191.1	103.3	68.6	44.7	267.4	305.7	249.3	207.1	191.9	188.6	174.0	123.3
1932	184.6	110.4	70.7	45.0	243.6	290.9	249.0	207.2	191.3	189.6	169.4	116.7
1933	174.2	111.3	63.4	39.1	226.5	276.4	229.5	202.9	198.4	176.9	172.1	92.3
1934	164.1	113.3	62.2	37.0	212.4	260.1	214.9	195.9	180.8	155.6	143.0	121.9
1935	145.1	97.3	44.4	30.7	180.7	228.6	207.1	177.6	164.2	139.1	109.5	84.1
1936	151.6	84.1	40.8	30.7	193.6	240.7	214.2	185.6	174.0	133.9	111.3	117.8
1937	145.0	84.1	46.7	28.8	182.4	224.2	208.3	178.6	168.6	130.8	117.3	90.8
1938	136.8	76.7	45.1	24.9	179.3	212.5	185.9	167.8	163.8	124.5	109.2	93.6
1939	129.1	80.3	40.6	24.1	164.8	202.4	174.2	169.7	149.0	111.8	105.4	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	479.3	...	...	...	...	...	...	...	...	...	...	...
1911	484.8	...	...	...	...	...	...	...	...	...	...	...
1912	459.9	...	...	...	...	...	...	...	...	...	...	...
1913	401.9	...	...	...	...	...	...	...	...	...	...	...
1914	417.8	551.5	234.2	125.5	493.6	515.2	516.4	451.9	534.0	515.0	543.4	*
1915	420.2	501.0	247.6	129.3	467.8	524.3	531.1	479.8	534.1	494.0	418.9	*
1916	322.3	278.2	156.8	86.4	378.9	463.3	445.6	381.5	405.7	481.9	372.6	*
1917	335.0	...	...	...	...	...	...	...	...	...	...	...
1918	351.0	256.6	176.9	93.7	477.0	525.0	440.4	399.0	400.6	403.3	333.8	*
1919	275.5	212.1	107.1	66.4	376.7	426.5	365.5	312.2	322.0	336.9	256.1	*
1920	255.4	204.8	92.4	55.7	362.4	384.8	326.5	295.6	285.2	336.6	256.6	337.0
1921	233.7	184.6	86.1	49.4	320.6	369.9	297.1	274.8	262.8	252.1	297.3	324.0
1922	216.6	160.9	82.6	38.1	287.5	342.2	297.3	261.2	243.4	300.2	289.6	*
1923	206.3	154.1	85.5	40.9	261.5	328.2	294.5	257.4	228.6	225.1	193.5	*
1924	215.0	149.6	80.8	45.6	282.9	336.8	291.1	268.2	238.7	243.9	250.1	290.5
1925	215.8	150.0	87.5	46.1	279.8	339.7	293.7	264.4	222.1	241.1	250.4	*
1926	221.5	149.6	95.8	49.1	272.2	345.1	317.7	265.2	251.3	246.4	229.0	*
1927	205.4	145.5	82.1	44.2	248.0	326.9	299.1	254.6	217.9	219.3	231.8	*
1928	199.4	146.1	75.8	40.7	250.7	320.2	284.5	250.9	209.3	216.5	241.0	*
1929	191.5	114.7	76.8	36.6	234.0	308.0	278.5	238.9	212.8	219.2	234.6	207.6
1930	194.3	128.4	77.0	40.3	235.8	319.1	275.0	239.6	199.2	243.3	219.1	*
1931	197.4	125.5	76.0	36.3	233.8	312.7	304.1	240.9	222.8	226.8	195.2	*
1932	191.9	117.4	73.5	37.4	206.6	307.2	298.6	248.8	222.6	235.6	219.4	*
1933	182.4	126.2	63.3	32.5	195.6	281.6	283.9	246.2	237.0	217.2	212.0	*
1934	173.4	124.2	68.9	28.9	184.3	270.1	264.8	241.0	215.3	199.0	188.2	187.7
1935	155.4	113.9	44.8	26.8	152.4	240.1	255.3	227.1	202.2	176.3	135.0	*
1936	163.9	81.2	43.2	25.9	162.7	259.0	269.2	241.5	217.0	163.3	150.1	156.1
1937	155.0	100.7	45.1	22.9	155.3	232.6	260.9	222.5	211.8	164.5	153.6	*
1938	144.0	74.8	47.8	19.2	151.5	215.3	220.2	217.6	205.7	161.6	132.2	*
1939	137.3	89.3	41.2	19.3	136.6	206.4	217.4	221.7	184.4	138.7	125.2	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

TUBERCULOSIS, ALL FORMS

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	406.8	---	---	---	---	---	---	---	---	---	---	---
1911	435.2	---	---	---	---	---	---	---	---	---	---	---
1912	394.5	---	---	---	---	---	---	---	---	---	---	---
1913	369.9	---	---	---	---	---	---	---	---	---	---	---
1914	374.0	334.1	234.7	183.4	545.3	497.0	390.4	326.7	266.6	262.8	241.5	*
1915	380.5	334.0	248.7	176.3	574.6	491.3	400.5	324.3	275.1	282.4	211.7	*
1916	323.0	270.0	149.9	126.2	480.7	476.9	398.6	286.3	237.3	248.3	273.1	*
1917	330.1	---	---	---	---	---	---	---	---	---	---	---
1918	340.9	228.8	151.7	133.3	551.0	495.9	365.8	284.4	260.3	256.6	296.4	265.9
1919	292.7	206.4	97.2	96.6	463.7	467.0	325.5	266.4	217.7	224.1	226.6	260.1
1920	269.6	198.0	88.0	95.3	431.3	408.1	291.9	254.9	220.6	226.7	238.6	*
1921	245.1	158.5	80.2	75.8	408.2	373.7	259.6	235.8	192.5	201.8	189.6	*
1922	221.2	136.7	65.8	60.1	358.3	346.8	266.2	202.7	186.3	180.0	151.4	*
1923	220.0	149.4	76.5	58.6	369.4	332.6	250.7	201.4	193.0	166.6	163.6	224.3
1924	222.3	150.3	78.7	66.3	373.9	323.2	246.4	220.7	179.1	177.5	172.9	*
1925	226.7	116.2	81.4	65.3	377.3	338.0	255.4	220.6	184.3	165.3	172.1	*
1926	226.1	133.5	89.8	65.1	374.5	346.8	245.7	210.0	168.9	172.1	154.5	*
1927	212.1	138.4	78.2	59.4	351.3	332.4	236.9	181.3	151.4	138.0	151.2	213.8
1928	199.6	137.9	66.7	57.9	327.7	308.9	227.7	179.2	142.8	155.0	164.4	*
1929	192.6	99.1	69.1	50.7	309.6	314.2	212.5	178.5	149.8	129.3	158.0	*
1930	189.8	122.6	71.9	58.7	300.0	298.8	209.8	167.5	150.1	147.8	140.3	*
1931	184.9	81.5	61.4	53.1	297.8	299.2	195.9	168.3	152.0	145.6	154.0	*
1932	177.4	103.4	67.9	52.4	277.2	275.7	200.8	160.1	151.4	138.2	122.2	*
1933	166.2	96.6	63.4	45.6	254.5	271.6	176.7	154.2	149.8	132.1	134.2	*
1934	155.0	102.7	55.6	45.1	237.8	250.8	166.7	145.7	137.9	107.5	100.2	*
1935	135.0	81.1	44.1	34.7	206.3	218.0	160.6	123.0	117.5	98.1	85.3	*
1936	139.6	86.9	38.5	35.3	221.6	223.8	161.4	124.4	122.1	101.7	74.6	*
1937	135.2	67.9	48.3	34.6	206.9	216.5	157.9	130.9	116.9	93.9	82.9	*
1938	129.8	78.5	42.5	30.6	204.5	210.1	153.1	114.0	114.4	84.1	87.4	*
1939	121.1	71.6	40.0	28.9	190.4	198.8	133.1	114.0	107.9	82.5	86.7	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: 1) Values shown differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes—	1,316	2,110	1,973
Male	729	1,067	1,055
Female	587	1,043	918

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

ICD Revision	Years in use	ICD Nos.
Fifth	1939	45-55
Fourth	1930-1938	45-53
Third	1921-1929	43-49
Second	1910-1920	39-45
First	1900-1909	39-45

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	64.0	*	2.9	1.8	3.2	14.0	52.5	139.1	260.9	421.0	544.7	623.2
1901	66.4	*	3.4	1.0	3.0	13.5	60.4	147.5	269.8	421.0	551.5	627.6
1902	66.3	5.3	3.1	1.5	3.2	13.0	54.9	142.0	274.8	430.3	572.7	703.0
1903	70.0	*	2.6	1.4	3.0	13.0	57.4	146.9	298.6	459.6	587.6	690.3
1904	71.5	4.9	2.8	1.2	3.2	14.7	56.4	146.0	301.3	486.8	608.2	675.1
1905	73.4	*	2.6	1.4	3.2	13.8	59.7	151.6	303.3	492.5	648.8	744.9
1906	69.3	*	2.7	1.2	3.4	13.4	54.7	152.8	294.0	480.5	625.4	726.8
1907	71.4	2.6	2.9	1.6	3.5	13.7	55.9	155.0	308.3	486.3	634.6	822.0
1908	71.5	2.6	3.1	1.2	3.6	12.9	55.7	150.3	309.3	497.4	687.0	769.2
1909	74.0	3.0	2.9	1.6	3.3	13.2	57.7	150.2	313.6	521.0	705.8	799.5
1910	76.2	3.1	3.3	1.5	3.5	14.1	55.6	156.7	322.4	541.7	749.9	810.9
1911	74.2	3.0	2.9	1.3	3.4	13.6	54.7	152.9	313.5	527.2	744.7	822.6
1912	77.0	3.9	3.7	1.6	3.5	14.1	56.8	154.4	324.8	547.3	761.3	847.0
1913	78.5	2.8	3.3	1.6	3.7	14.3	57.9	158.8	326.6	557.7	779.1	887.8
1914	78.7	3.4	3.3	1.7	3.6	13.8	57.8	155.1	325.2	565.6	774.0	835.8
1915	80.7	3.6	3.1	1.7	3.6	14.6	58.3	156.3	332.6	573.7	801.2	851.9
1916	81.0	2.7	3.6	1.6	4.1	14.9	58.7	158.0	334.5	582.0	819.9	895.9
1917	80.8	4.2	3.1	1.7	4.0	15.3	58.0	154.4	334.8	583.2	807.6	865.4
1918	80.8	3.4	3.8	1.6	4.2	15.4	57.5	153.6	324.8	575.0	780.8	895.3
1919	81.0	2.2	3.1	1.6	3.8	14.7	55.1	152.8	333.9	581.3	829.0	960.3
1920	83.4	3.2	3.1	1.6	3.8	14.7	56.0	155.1	341.2	607.7	900.0	1,017.5
1921	85.5	2.5	3.6	1.6	4.1	14.5	55.9	158.9	345.7	625.3	918.6	1,119.0
1922	86.2	2.5	3.1	1.7	4.0	14.7	56.1	157.6	345.7	645.0	934.2	1,164.3
1923	88.4	3.6	3.3	1.7	3.9	14.8	56.9	159.0	348.8	646.3	963.7	1,211.1
1924	90.4	2.8	3.8	2.2	4.1	15.3	58.9	163.5	348.7	656.1	973.1	1,185.7
1925	92.0	2.8	3.4	1.6	4.0	15.3	57.6	162.3	350.4	665.2	1,027.1	1,250.8
1926	94.6	2.6	3.9	2.0	4.4	15.4	58.4	164.5	353.5	679.5	1,045.0	1,275.1
1927	95.2	2.0	3.6	1.9	4.4	16.2	59.1	161.7	350.0	678.5	1,062.8	1,297.1
1928	95.7	3.7	3.6	2.1	4.6	15.7	59.1	162.4	351.6	683.1	1,071.2	1,237.6
1929	95.8	2.7	3.7	1.9	4.5	15.6	57.7	160.0	351.0	682.3	1,028.9	1,244.8
1930	97.4	3.1	4.1	2.0	4.2	16.7	58.9	159.6	355.6	677.1	1,019.7	1,196.3
1931	99.0	2.8	4.0	2.1	4.6	16.0	58.9	159.7	352.0	676.6	1,047.4	1,225.6
1932	102.3	2.8	3.9	2.2	4.6	15.8	58.6	163.6	362.1	676.8	1,081.1	1,255.7
1933	102.3	2.8	3.9	2.2	4.6	15.6	57.8	160.2	355.9	683.2	1,088.1	1,220.3
1934	106.4	2.7	4.1	2.4	4.7	16.6	59.5	166.9	366.0	683.8	1,120.4	1,199.6
1935	108.2	3.1	4.0	2.1	4.7	16.3	59.3	164.2	362.7	689.7	1,133.9	1,297.5
1936	111.4	2.7	4.1	2.5	4.7	16.7	60.9	165.6	362.5	689.7	1,185.0	1,336.5
1937	112.4	2.9	4.2	2.3	4.7	16.4	60.8	164.0	363.7	684.0	1,170.4	1,270.4
1938	114.9	3.0	4.4	2.5	4.8	17.0	60.3	166.6	364.4	685.8	1,183.5	1,267.8
1939	117.5	3.8	4.7	2.8	5.2	17.3	61.1	168.1	369.7	682.0	1,176.9	1,329.2

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	47.1	*	3.6	2.1	3.0	9.1	27.9	83.5	189.5	377.7	495.1	590.4
1901	49.2	*	3.7	1.1	2.6	9.5	32.6	90.6	205.3	368.3	499.3	703.3
1902	48.0	*	3.1	1.6	2.9	8.1	29.4	84.1	205.1	367.2	509.6	635.8
1903	51.3	*	2.5	1.6	3.4	7.9	31.0	92.3	221.5	400.5	516.3	622.9
1904	52.6	*	2.9	1.2	3.5	9.7	28.0	90.7	235.3	418.8	522.0	629.3
1905	53.8	*	*	1.6	3.4	8.1	31.6	89.9	234.9	414.3	609.3	735.7
1906	51.3	*	2.8	1.5	3.6	7.5	30.3	95.0	224.0	420.9	546.3	704.5
1907	53.5	*	3.1	1.7	3.4	8.8	30.5	97.7	242.6	424.4	573.7	707.6
1908	54.1	*	2.8	1.2	3.8	8.5	29.0	94.7	250.1	430.3	625.9	692.8
1909	56.8	*	3.1	1.7	3.5	8.3	31.1	96.9	250.7	468.4	646.0	717.1
1910	59.0	*	3.6	1.6	3.6	8.8	30.9	103.0	260.9	481.5	680.3	780.5
1911	57.7	*	3.0	1.3	3.7	8.9	28.4	100.6	252.7	472.3	693.5	758.3
1912	59.5	4.3	4.1	2.0	3.3	8.2	31.5	100.6	262.0	479.8	696.3	755.3
1913	61.4	3.1	3.4	1.9	3.7	9.0	30.8	104.4	266.7	498.9	730.9	800.1
1914	62.3	*	3.4	2.0	3.6	8.8	30.9	103.3	265.4	518.5	707.9	800.4
1915	64.9	3.8	3.6	1.9	3.7	10.0	31.7	106.5	276.1	526.1	745.8	833.6
1916	64.8	*	3.7	1.7	4.3	10.1	33.3	107.1	272.5	528.6	757.8	848.9
1917	65.3	3.8	3.2	2.0	4.0	10.3	31.8	102.3	276.7	536.4	759.8	816.3
1918	66.0	3.6	3.9	1.7	4.3	10.4	33.2	103.6	265.6	518.0	734.9	891.9
1919	66.7	*	3.2	1.7	3.8	9.3	30.4	103.2	281.4	538.5	794.0	955.1
1920	69.6	3.1	3.4	1.7	3.7	9.6	31.1	107.5	289.5	575.5	874.4	988.9
1921	71.7	3.3	3.8	1.5	4.3	9.6	32.4	108.0	296.7	596.7	897.2	1,024.7
1922	73.8	2.7	3.4	1.7	4.2	9.1	32.2	110.2	301.8	629.7	926.0	1,078.9
1923	76.3	3.6	3.5	1.8	4.1	10.4	33.5	113.2	304.9	627.9	962.5	1,170.8
1924	78.0	2.9	3.8	2.4	4.3	10.4	34.7	115.0	306.2	638.5	975.2	1,146.8
1925	79.6	2.3	3.9	1.9	4.1	10.5	33.4	114.1	305.9	647.1	1,033.8	1,254.8
1926	82.7	2.4	4.3	2.3	4.4	10.9	34.7	115.4	310.0	669.1	1,079.3	1,262.8
1927	82.5	2.6	3.5	2.1	4.7	10.4	35.1	113.3	306.8	666.1	1,067.5	1,203.9
1928	84.1	3.1	3.8	2.3	4.8	10.6	35.4	117.5	305.3	673.7	1,112.7	1,180.6
1929	84.5	2.9	4.0	2.0	4.9	10.4	35.3	116.1	306.5	679.8	1,045.7	1,239.7
1930	86.5	3.2	4.2	2.0	4.3	11.0	35.6	117.0	315.1	678.8	1,046.7	1,193.6
1931	87.9	3.1	4.2	2.2	5.1	10.8	34.9	114.5	309.8	679.3	1,086.5	1,267.3
1932	91.4	3.4	4.1	2.4	4.8	10.5	35.5	118.1	325.8	677.1	1,125.7	1,303.1
1933	92.4	2.6	4.2	2.5	5.2	10.4	35.4	119.1	316.0	694.3	1,138.5	1,280.3
1934	96.6	3.2	4.6	2.8	4.9	11.4	35.8	125.1	326.9	695.4	1,205.6	1,285.5
1935	98.1	2.8	4.5	2.3	5.0	10.2	36.5	123.8	324.3	701.7	1,202.5	1,360.3
1936	101.6	2.9	4.2	2.6	5.0	10.7	38.5	127.1	328.4	701.6	1,246.9	1,409.3
1937	103.8	3.4	4.6	2.6	5.1	10.7	37.9	127.2	335.5	702.0	1,253.4	1,379.5
1938	106.9	2.6	4.9	2.8	5.1	11.6	38.3	130.7	335.7	712.0	1,280.2	1,356.4
1939	110.0	4.5	4.9	3.0	5.8	12.1	38.7	132.4	345.3	710.6	1,277.2	1,419.3

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	80.9	*	*	1.4	3.4	18.9	78.8	196.9	330.9	462.1	589.8	647.0
1901	83.7	*	3.0	*	3.5	17.6	89.9	207.1	333.0	471.0	599.0	572.5
1902	84.7	*	3.1	1.5	3.5	18.0	81.9	202.7	343.2	490.0	629.6	751.9
1903	88.7	*	2.8	1.2	2.6	18.2	85.5	204.3	374.4	515.5	651.6	739.6
1904	90.5	*	2.8	1.3	2.9	19.8	86.5	204.4	366.5	551.1	685.2	708.8
1905	93.2	*	3.1	1.2	2.9	19.6	89.6	216.8	371.0	566.4	683.9	751.7
1906	87.9	*	2.5	1.0	3.3	19.7	81.5	216.1	366.1	538.7	697.1	745.5
1907	90.0	*	2.6	1.4	3.5	19.0	83.9	217.9	376.2	546.8	689.5	908.0
1908	89.8	*	3.3	1.2	3.3	17.8	85.3	212.3	371.8	563.6	742.8	828.3
1909	92.0	*	2.7	1.4	3.1	18.5	87.3	209.3	380.1	573.2	760.5	863.8
1910	94.5	*	2.9	1.4	3.4	20.1	83.2	217.1	388.3	601.8	813.5	834.9
1911	91.6	*	2.8	1.3	3.1	18.7	84.1	211.9	379.3	582.5	791.8	855.6
1912	95.5	3.4	3.3	1.3	3.6	20.6	85.0	215.2	393.0	615.3	820.8	919.0
1913	96.4	*	3.2	1.2	3.6	20.0	87.9	220.2	391.8	617.1	823.1	956.1
1914	96.0	4.2	3.3	1.5	3.6	19.3	87.5	213.7	390.5	613.5	834.5	863.4
1915	97.3	3.4	2.7	1.6	3.5	19.6	87.7	212.7	394.3	622.3	851.9	866.0
1916	98.0	2.8	3.5	1.6	3.8	20.1	86.5	215.8	402.5	636.5	876.1	931.7
1917	97.0	4.7	3.0	1.4	4.1	20.4	86.7	213.5	398.5	631.2	841.9	902.5
1918	95.7	3.2	3.7	1.5	4.1	20.2	83.9	210.5	390.1	633.7	822.5	897.8
1919	95.7	2.3	3.0	1.6	3.8	20.1	82.0	209.4	391.8	625.6	860.8	964.1
1920	97.7	3.4	2.7	1.4	4.0	19.9	83.4	208.9	398.0	641.0	923.4	1,039.0
1921	99.8	*	3.4	1.6	4.0	19.4	81.7	216.1	399.1	654.9	938.3	1,190.5
1922	98.9	2.2	2.8	1.7	3.8	20.2	82.0	210.4	393.2	660.8	922.6	1,229.2
1923	100.8	3.5	3.1	1.7	3.7	19.1	82.2	209.5	396.3	665.4	964.7	1,242.0
1924	103.2	2.7	3.7	1.9	3.9	20.3	85.0	216.6	395.1	674.3	971.2	1,215.5
1925	104.9	3.2	3.0	1.4	3.9	20.2	83.7	214.9	398.9	684.0	1,020.9	1,247.7
1926	106.8	2.8	3.5	1.8	4.4	19.9	83.7	217.8	401.1	690.2	1,013.3	1,284.6
1927	108.3	*	3.6	1.8	4.1	22.0	84.7	214.3	397.2	691.3	1,058.3	1,368.7
1928	107.7	4.2	3.4	1.9	4.3	20.8	84.5	211.3	402.3	692.8	1,032.4	1,281.2
1929	107.4	2.5	3.3	1.8	4.0	20.9	81.5	208.0	399.4	684.8	1,013.2	1,248.7
1930	108.6	2.9	3.9	2.0	4.0	22.4	83.3	206.1	399.3	675.4	994.2	1,198.3
1931	110.4	2.5	3.8	1.9	4.2	21.1	84.0	208.8	397.3	673.9	1,010.6	1,194.3
1932	113.4	2.1	3.7	1.9	4.5	20.9	82.7	212.9	401.1	676.6	1,039.3	1,220.1
1933	112.4	3.1	3.6	1.9	4.1	20.8	81.1	204.6	398.8	671.7	1,040.7	1,175.2
1934	116.3	2.1	3.5	2.1	4.4	21.7	84.0	212.0	407.7	672.0	1,040.5	1,135.0
1935	118.4	3.5	3.4	2.0	4.5	22.3	82.7	207.7	403.7	677.4	1,069.8	1,250.3
1936	121.3	2.4	3.9	2.4	4.4	22.5	83.8	206.9	398.8	677.7	1,127.5	1,281.7
1937	121.1	2.4	3.8	2.0	4.2	22.0	84.3	203.4	393.6	665.9	1,093.7	1,188.2
1938	123.0	3.5	4.0	2.3	4.5	22.4	82.7	204.9	394.7	659.6	1,094.6	1,201.0
1939	125.2	3.1	4.4	2.5	4.5	22.3	83.7	206.0	395.4	653.4	1,085.1	1,261.4

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	76.9	---	---	---	---	---	---	---	---	---	---	---
1911	74.8	---	---	---	---	---	---	---	---	---	---	---
1912	77.7	---	---	---	---	---	---	---	---	---	---	---
1913	79.5	---	---	---	---	---	---	---	---	---	---	---
1914	79.7	3.4	3.4	1.7	3.6	13.6	57.2	155.1	327.5	572.8	785.0	863.1
1915	81.7	3.5	3.2	1.7	3.6	14.2	58.0	156.5	334.3	580.2	811.5	871.4
1916	83.2	2.9	3.7	1.7	4.0	14.6	57.9	159.6	338.3	592.8	836.0	920.7
1917	83.1	---	---	---	---	---	---	---	---	---	---	---
1918	83.3	3.6	3.9	1.6	4.1	14.9	56.2	154.4	329.3	588.2	804.2	938.3
1919	84.1	2.3	3.1	1.7	3.7	14.2	54.4	154.5	339.7	597.6	855.7	1,024.4
1920	86.5	3.2	3.1	1.6	3.8	14.1	54.9	156.4	347.1	625.2	929.0	1,070.4
1921	88.8	2.6	3.6	1.5	3.9	13.9	55.1	160.3	352.0	641.5	948.8	1,178.4
1922	90.0	2.5	3.3	1.7	4.0	13.9	54.9	158.7	352.6	663.7	960.3	1,244.4
1923	92.1	3.7	3.4	1.8	3.9	14.0	55.5	158.7	356.7	663.0	996.9	1,274.8
1924	94.0	3.0	3.9	2.3	4.0	14.7	56.8	163.0	355.9	673.7	1,009.0	1,248.3
1925	95.5	2.8	3.5	1.7	3.9	14.4	55.6	161.2	357.1	682.0	1,060.8	1,317.8
1926	98.1	2.7	3.9	2.1	4.3	14.6	56.4	163.0	359.4	695.0	1,082.5	1,350.8
1927	98.9	2.1	3.8	2.0	4.4	15.2	57.1	160.3	355.4	693.3	1,101.0	1,374.5
1928	100.1	3.9	3.8	2.1	4.5	14.6	56.7	161.9	358.3	702.6	1,113.8	1,326.5
1929	100.2	2.8	3.9	2.0	4.4	14.7	55.5	159.6	356.8	701.4	1,068.4	1,329.7
1930	101.9	3.2	4.2	2.1	4.1	15.8	56.8	159.0	361.9	696.0	1,057.3	1,283.7
1931	103.5	2.8	4.0	2.1	4.6	14.9	56.7	159.4	357.5	694.9	1,085.5	1,314.2
1932	106.7	3.0	3.9	2.3	4.5	14.8	56.0	161.9	366.3	695.1	1,122.2	1,341.5
1933	106.6	3.1	4.0	2.2	4.6	14.6	55.3	158.0	360.0	700.7	1,124.9	1,309.3
1934	110.8	2.6	4.3	2.5	4.6	15.5	56.8	164.0	369.5	701.9	1,159.8	1,285.8
1935	113.0	3.3	4.1	2.2	4.7	15.7	57.0	163.1	366.0	709.3	1,173.5	1,388.2
1936	116.5	2.8	4.2	2.7	4.6	15.8	58.3	164.6	367.4	710.1	1,228.2	1,436.8
1937	117.4	3.0	4.5	2.4	4.6	15.5	58.1	162.2	367.9	705.6	1,212.1	1,352.6
1938	119.8	3.1	4.7	2.6	4.8	16.0	57.5	164.3	366.9	707.2	1,223.2	1,366.3
1939	122.6	4.0	4.9	2.9	5.2	16.7	58.4	165.1	372.8	704.2	1,216.4	1,426.1

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

WHITE, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	59.8	---	---	---	---	---	---	---	---	---	---	---
1911	58.5	---	---	---	---	---	---	---	---	---	---	---
1912	60.3	---	---	---	---	---	---	---	---	---	---	---
1913	62.7	---	---	---	---	---	---	---	---	---	---	---
1914	63.7	*	3.4	1.9	3.7	8.9	31.1	104.7	269.0	527.4	722.4	832.0
1915	66.1	3.9	3.6	1.9	3.7	10.0	31.6	108.1	278.9	533.3	758.8	860.8
1916	67.2	*	3.8	1.8	4.3	10.1	33.8	109.4	278.4	542.5	775.5	880.3
1917	67.9	---	---	---	---	---	---	---	---	---	---	---
1918	69.1	3.7	4.0	1.8	4.3	10.5	33.4	106.5	272.7	535.8	760.0	946.7
1919	70.3	*	3.3	1.7	3.8	9.3	30.9	106.5	291.4	558.5	826.2	1,031.7
1920	73.2	3.2	3.5	1.7	3.7	9.5	31.6	110.8	298.0	599.3	908.3	1,047.5
1921	75.4	3.4	3.8	1.5	4.2	9.7	32.8	111.2	306.0	617.9	932.8	1,084.1
1922	78.1	2.8	3.7	1.8	4.4	9.0	32.4	113.2	313.1	654.9	968.2	1,155.5
1923	80.6	3.5	3.7	1.8	4.2	10.4	33.9	115.4	316.2	652.0	1,000.8	1,243.7
1924	82.3	3.2	4.1	2.6	4.3	10.4	34.7	117.3	316.6	663.3	1,017.1	1,197.9
1925	83.6	2.4	3.8	1.9	4.0	10.5	32.9	115.3	315.8	669.2	1,077.3	1,322.4
1926	86.7	2.6	4.4	2.4	4.4	10.8	34.9	116.1	318.7	688.6	1,124.2	1,330.0
1927	86.9	2.8	3.8	2.2	4.8	10.4	35.3	114.6	316.0	686.3	1,111.6	1,273.6
1928	89.0	3.3	4.1	2.3	4.9	10.6	35.1	118.9	315.5	686.9	1,163.4	1,280.9
1929	89.5	2.9	4.1	2.1	5.0	10.2	34.7	117.9	316.0	703.5	1,090.4	1,333.2
1930	91.6	3.2	4.5	2.0	4.5	10.9	35.5	118.3	325.1	702.2	1,087.9	1,285.9
1931	93.0	3.2	4.2	2.3	5.2	10.7	34.7	116.8	318.0	701.7	1,131.3	1,359.2
1932	96.3	3.8	4.2	2.6	4.8	10.4	35.0	118.8	332.7	698.4	1,171.6	1,391.1
1933	97.4	2.7	4.4	2.5	5.2	10.2	35.1	119.5	323.0	717.3	1,180.1	1,380.8
1934	101.8	3.2	5.0	2.8	5.0	11.3	35.0	125.3	333.6	718.1	1,251.4	1,384.6
1935	103.5	2.8	4.7	2.3	5.0	10.4	36.2	125.1	330.3	726.1	1,248.6	1,454.8
1936	107.4	3.1	4.3	2.8	5.2	10.8	38.1	128.5	336.2	725.3	1,295.0	1,524.8
1937	109.5	3.7	4.9	2.7	5.2	10.9	37.2	127.0	342.8	727.3	1,299.8	1,478.9
1938	112.5	2.7	5.3	2.9	5.2	11.7	37.5	130.9	340.8	738.0	1,325.9	1,468.0
1939	115.7	4.7	5.1	3.2	6.0	12.2	38.3	131.8	350.7	737.1	1,323.5	1,531.1

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	94.9	---	---	---	---	---	---	---	---	---	---	---
1911	92.0	---	---	---	---	---	---	---	---	---	---	---
1912	96.0	---	---	---	---	---	---	---	---	---	---	---
1913	97.2	---	---	---	---	---	---	---	---	---	---	---
1914	96.7	4.0	3.4	1.4	3.6	18.6	86.1	211.8	391.0	618.8	842.4	887.5
1915	98.0	*	2.7	1.6	3.4	18.8	87.0	210.9	394.5	627.7	859.5	879.7
1916	99.9	3.0	3.5	1.6	3.8	19.3	84.3	216.1	403.4	643.8	890.7	951.5
1917	99.0	---	---	---	---	---	---	---	---	---	---	---
1918	97.5	3.5	3.8	1.5	3.9	19.2	81.0	208.3	391.1	641.8	844.1	931.9
1919	98.4	2.6	2.9	1.6	3.5	19.0	80.0	208.4	392.5	637.8	882.3	1,019.0
1920	100.4	3.2	2.7	1.4	3.9	18.8	80.5	207.4	400.4	651.8	947.9	1,087.7
1921	102.7	*	3.4	1.5	3.7	18.3	79.7	214.6	401.6	665.7	963.4	1,250.3
1922	102.2	*	3.0	1.7	3.6	18.9	79.3	208.6	395.0	672.7	953.0	1,312.6
1923	103.9	3.8	3.1	1.7	3.5	17.5	79.0	206.0	400.1	676.1	993.4	1,298.9
1924	106.1	2.8	3.7	1.9	3.6	18.9	80.7	212.5	398.2	684.4	1,001.5	1,287.5
1925	107.8	3.2	3.2	1.4	3.7	18.5	80.1	210.9	401.6	695.1	1,045.7	1,314.2
1926	109.8	2.7	3.3	1.8	4.3	18.4	79.6	213.6	403.4	701.7	1,044.2	1,367.1
1927	111.2	*	3.8	1.8	3.9	20.1	80.4	209.7	398.0	700.4	1,091.2	1,452.8
1928	111.5	4.5	3.5	1.9	4.2	18.6	79.7	208.4	404.5	708.5	1,067.7	1,361.7
1929	111.3	2.8	3.7	2.0	3.9	19.2	77.6	204.8	400.6	699.3	1,047.7	1,327.1
1930	112.5	3.3	3.9	2.1	3.8	20.6	79.4	203.2	401.2	689.6	1,028.4	1,282.1
1931	114.3	2.4	3.7	2.0	4.0	19.0	80.0	205.5	399.5	688.0	1,042.4	1,280.4
1932	117.2	*	3.7	2.0	4.2	19.0	78.1	208.4	402.0	691.8	1,075.9	1,304.2
1933	116.0	3.4	3.6	1.9	4.0	18.9	76.5	199.4	399.2	683.8	1,073.0	1,255.4
1934	120.1	*	3.6	2.2	4.2	19.7	79.6	207.5	407.5	685.5	1,074.0	1,211.3
1935	122.6	3.8	3.6	2.1	4.3	20.9	78.6	203.8	403.6	692.4	1,103.5	1,338.1
1936	125.8	2.5	4.2	2.6	4.1	20.7	79.3	203.1	400.2	694.7	1,166.1	1,370.5
1937	125.5	2.4	4.0	2.0	4.0	20.1	79.7	199.8	394.4	684.0	1,131.2	1,257.4
1938	127.2	3.5	4.2	2.3	4.4	20.3	78.0	199.9	394.3	676.6	1,128.8	1,289.6
1939	129.5	3.3	4.6	2.6	4.3	21.0	79.0	200.6	396.0	671.6	1,118.6	1,346.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	54.0	---	---	---	---	---	---	---	---	---	---	---
1911	57.3	---	---	---	---	---	---	---	---	---	---	---
1912	57.2	---	---	---	---	---	---	---	---	---	---	---
1913	55.8	---	---	---	---	---	---	---	---	---	---	---
1914	55.4	*	*	*	*	18.7	68.5	154.1	258.5	337.8	396.8	*
1915	58.3	*	*	*	4.3	22.8	64.4	151.3	281.9	368.8	448.4	453.2
1916	49.6	*	*	*	4.4	20.2	69.9	130.7	254.0	346.2	430.5	548.2
1917	49.4	---	---	---	---	---	---	---	---	---	---	---
1918	49.8	*	*	*	5.3	21.8	73.9	141.8	239.0	318.1	295.3	387.9
1919	46.8	*	*	1.2	4.8	20.6	62.6	132.4	236.9	309.6	360.2	332.1
1920	48.5	*	*	1.5	4.1	21.7	68.2	137.8	243.3	305.1	376.9	472.2
1921	49.4	*	3.5	2.0	5.8	20.1	64.8	141.9	240.5	341.7	378.0	491.7
1922	49.4	*	*	1.3	4.2	21.9	68.8	145.8	244.6	350.4	355.1	416.0
1923	51.9	*	*	1.4	4.2	22.6	72.0	162.5	231.0	353.9	421.6	590.7
1924	54.2	*	2.6	1.3	5.4	21.8	81.4	169.6	242.8	353.5	386.9	571.5
1925	56.5	*	3.1	1.3	5.0	24.1	79.8	175.0	248.9	363.4	463.3	578.4
1926	58.3	*	4.2	1.6	4.8	23.6	79.9	182.4	264.3	393.5	415.2	513.8
1927	59.0	*	*	1.1	4.8	25.1	81.1	177.1	271.1	410.3	430.7	538.8
1928	56.5	*	2.2	1.5	4.8	25.1	83.2	167.8	262.2	355.3	419.1	447.2
1929	56.0	*	*	0.9	4.5	23.7	79.4	164.5	271.5	355.1	406.8	481.5
1930	56.6	*	2.8	1.5	4.6	24.9	78.5	166.1	267.2	354.9	404.2	401.5
1931	58.2	*	4.2	1.4	4.8	25.5	80.0	162.2	275.1	368.5	418.6	414.8
1932	62.5	*	3.8	1.3	5.6	24.2	83.2	181.0	303.6	375.3	396.4	466.9
1933	64.5	*	3.3	2.1	5.0	24.2	80.6	182.9	301.4	411.2	489.6	431.7
1934	67.2	*	2.4	2.0	5.5	25.5	83.8	187.4	318.0	411.9	474.0	432.9
1935	65.8	*	2.7	1.6	5.5	22.0	79.6	175.6	319.2	403.4	478.3	489.0
1936	66.3	*	2.8	1.3	5.0	24.4	83.9	176.8	297.2	402.8	465.8	441.2
1937	68.2	*	2.2	1.6	5.0	23.6	84.9	183.3	306.2	388.6	471.0	536.2
1938	72.3	*	2.5	1.9	5.3	25.8	85.5	191.2	329.7	403.7	514.7	388.5
1939	73.5	*	3.1	1.8	5.1	22.7	84.3	200.2	327.6	399.9	506.6	465.8

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	33.9	---	---	---	---	---	---	---	---	---	---	---
1911	35.4	---	---	---	---	---	---	---	---	---	---	---
1912	35.8	---	---	---	---	---	---	---	---	---	---	---
1913	34.4	---	---	---	---	---	---	---	---	---	---	---
1914	31.7	*	*	*	*	*	27.7	73.6	167.1	251.6	219.3	*
1915	37.9	*	*	*	*	10.5	34.1	72.8	198.0	306.2	307.2	*
1916	30.4	*	*	*	*	9.9	25.9	70.4	154.6	245.6	348.2	*
1917	29.5	---	---	---	---	---	---	---	---	---	---	---
1918	27.6	*	*	*	4.3	9.2	30.0	64.0	137.5	191.8	232.8	*
1919	27.3	*	*	*	3.0	9.0	24.6	66.0	128.7	228.1	256.1	*
1920	29.0	*	*	*	3.1	10.8	25.4	69.0	158.2	193.0	295.9	383.0
1921	31.0	*	*	*	4.6	8.4	28.4	70.6	155.2	252.1	293.9	372.7
1922	31.8	*	*	*	3.1	10.1	30.1	79.0	153.6	265.2	295.4	316.4
1923	33.3	*	*	*	2.7	9.9	29.9	90.2	152.1	260.2	370.0	394.5
1924	34.3	*	*	*	4.2	9.9	34.6	89.1	165.4	246.7	322.3	595.5
1925	37.5	*	*	*	4.2	10.5	38.9	100.3	168.9	277.1	335.6	511.4
1926	40.3	*	*	*	4.0	12.4	33.0	108.4	188.5	336.4	352.9	529.1
1927	38.3	*	*	*	3.6	10.5	33.6	99.2	184.5	324.1	364.3	477.7
1928	38.9	*	*	1.8	4.0	10.6	38.7	103.8	180.6	307.5	359.3	252.4
1929	39.1	*	*	*	3.5	12.0	40.8	98.0	188.7	296.1	353.1	386.8
1930	39.4	*	*	1.9	3.1	11.8	36.5	104.6	188.2	296.7	375.6	356.1
1931	40.2	*	4.2	*	3.7	11.3	37.5	91.1	206.3	317.8	351.0	426.8
1932	45.7	*	*	*	4.6	11.1	40.5	111.1	237.9	340.2	363.5	493.7
1933	47.2	*	*	2.2	5.2	11.9	38.4	115.1	229.5	352.6	465.3	385.1
1934	50.3	*	*	2.7	4.4	11.8	43.5	122.6	243.5	367.9	459.0	397.9
1935	48.9	*	*	1.9	4.6	8.8	39.3	110.1	248.6	361.7	446.4	511.1
1936	49.6	*	*	1.5	3.5	10.4	42.7	111.8	229.8	381.5	454.0	369.0
1937	53.1	*	*	1.5	4.3	8.9	43.9	129.3	243.3	371.7	485.8	482.6
1938	56.7	*	*	2.0	4.9	11.1	46.1	128.6	269.8	384.6	521.9	347.5
1939	59.1	*	*	1.8	4.4	11.3	43.3	139.4	275.8	389.9	507.6	407.6

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MALIGNANT NEOPLASMS

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	77.0	---	---	---	---	---	---	---	---	---	---	---
1911	81.7	---	---	---	---	---	---	---	---	---	---	---
1912	81.0	---	---	---	---	---	---	---	---	---	---	---
1913	78.8	---	---	---	---	---	---	---	---	---	---	---
1914	80.9	*	*	*	*	32.5	118.3	261.7	374.0	435.8	563.5	*
1915	80.4	*	*	*	*	36.1	101.2	258.0	388.8	440.0	582.3	598.5
1916	69.5	*	*	*	*	4.9	30.4	119.3	209.8	381.1	463.9	510.6
1917	69.8	---	---	---	---	---	---	---	---	---	---	---
1918	72.3	*	*	*	6.3	33.7	121.9	246.6	369.7	464.6	356.5	505.1
1919	66.6	*	*	*	6.3	31.4	103.3	223.1	378.8	406.1	463.9	433.5
1920	68.4	*	*	*	5.0	31.7	114.9	230.9	354.5	438.4	459.1	539.0
1921	68.1	*	*	*	6.9	30.8	104.2	236.9	351.8	449.3	463.5	576.4
1922	67.2	*	*	*	5.2	32.6	109.5	232.3	364.3	454.6	415.7	484.4
1923	70.7	*	*	*	5.6	34.1	116.1	253.8	334.2	469.1	473.7	721.7
1924	74.2	*	*	*	6.4	32.7	130.3	268.6	344.0	484.5	451.8	555.7
1925	75.6	*	*	*	5.6	36.7	122.0	266.5	352.6	467.1	590.0	622.9
1926	76.3	*	5.4	*	5.5	34.2	128.1	270.7	362.7	461.9	476.8	503.5
1927	79.7	*	*	*	5.9	38.8	129.6	269.5	384.7	512.7	496.3	581.4
1928	74.1	*	*	*	5.5	38.6	127.2	242.7	370.1	411.2	477.9	586.9
1929	72.8	*	*	*	5.4	34.5	117.2	241.9	380.6	422.9	458.9	551.5
1930	73.7	*	*	*	6.0	37.2	119.5	237.5	370.0	420.7	431.4	435.6
1931	75.9	*	4.1	1.5	5.9	38.7	121.4	243.7	363.7	425.5	482.7	405.7
1932	79.1	*	4.4	*	6.5	36.5	124.7	260.2	387.3	414.7	427.6	446.7
1933	81.4	*	*	1.9	4.8	35.5	121.5	259.2	391.7	476.6	512.6	466.5
1934	83.7	*	*	*	6.5	38.1	122.9	259.5	410.5	460.8	488.2	459.0
1935	82.3	*	*	*	6.3	34.1	118.5	248.0	405.7	449.5	508.5	472.5
1936	82.6	*	*	*	6.5	37.3	123.6	248.0	378.7	426.2	477.1	494.7
1937	83.0	*	*	1.7	5.7	37.1	124.2	241.9	381.3	407.1	457.0	575.9
1938	87.5	*	*	1.8	5.6	39.3	123.2	258.7	400.2	424.6	507.9	418.6
1939	87.5	*	*	1.9	5.7	33.1	123.4	265.3	387.7	410.7	505.5	508.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes ...	282	493	456
Male....	104	193	195
Female....	178	300	261

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ICD Revision	Years in use	ICD Nos.
Fifth	1939	61
Fourth	1930-1938	59
Third	1921-1929	57
Second	1910-1920	50
First	1900-1909	50

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	11.0	*	2.0	3.3	4.2	5.0	6.8	16.3	40.3	67.0	72.0	48.1
1901	11.6	*	2.5	3.9	3.5	5.3	8.1	19.2	42.9	66.4	67.1	*
1902	11.7	*	2.3	3.7	3.6	4.3	7.2	16.7	44.7	76.6	82.7	58.8
1903	12.7	*	1.5	3.6	3.6	5.6	7.2	19.4	50.7	80.8	86.2	57.7
1904	14.2	*	2.5	4.3	3.5	5.2	8.4	23.8	55.8	91.2	99.0	56.4
1905	14.1	*	2.1	4.1	3.8	5.0	7.6	21.9	58.6	91.9	92.9	79.1
1906	13.4	*	2.2	3.4	3.9	4.2	7.8	22.6	57.3	90.7	90.7	70.3
1907	14.2	*	1.7	3.6	4.3	5.1	7.8	21.5	61.6	100.8	93.5	81.6
1908	13.8	*	2.1	3.7	4.0	4.7	7.1	22.5	57.6	98.1	94.5	71.6
1909	14.1	*	2.1	3.6	3.6	4.8	8.1	21.7	58.4	100.1	107.0	74.6
1910	15.3	*	2.4	3.8	4.6	5.1	8.3	23.7	65.0	104.5	112.0	74.1
1911	15.1	2.0	1.7	3.7	4.2	5.1	8.9	21.8	63.4	107.9	117.9	87.0
1912	15.1	*	1.7	3.8	4.4	5.1	8.3	22.9	60.6	105.7	124.3	74.9
1913	15.4	*	2.2	3.8	4.6	5.3	8.1	23.7	63.1	106.2	117.6	95.8
1914	16.2	*	2.1	3.7	4.7	5.7	8.2	25.3	64.3	115.9	125.8	95.4
1915	17.6	*	2.2	4.3	4.7	5.9	8.9	26.0	72.3	125.3	137.7	89.3
1916	16.9	*	2.3	3.1	4.2	4.9	8.9	25.5	71.8	123.7	136.5	106.4
1917	16.9	1.5	2.6	3.4	4.5	5.9	8.2	23.8	71.8	119.2	137.5	122.6
1918	16.1	1.2	2.0	3.4	5.7	7.1	9.1	22.8	61.7	108.2	114.7	94.6
1919	15.0	*	2.1	3.1	4.4	5.1	8.0	21.6	59.8	108.6	116.8	96.4
1920	16.1	1.4	2.2	3.5	4.8	5.6	8.2	23.1	64.8	113.7	135.0	110.9
1921	16.7	*	2.1	3.8	4.8	5.9	8.4	23.6	69.0	114.2	132.9	135.1
1922	18.3	1.7	2.1	3.9	5.0	6.3	9.3	26.2	73.7	130.2	150.7	145.0
1923	17.7	1.1	1.6	2.5	3.7	4.6	8.4	25.4	73.5	135.1	157.5	130.0
1924	16.4	*	1.6	2.3	3.0	3.5	6.9	22.4	68.9	126.9	153.6	148.5
1925	16.8	1.0	1.3	2.1	2.9	3.4	7.0	23.1	69.8	130.6	161.6	124.4
1926	17.9	1.0	1.6	2.4	3.6	3.4	7.0	23.7	72.9	141.3	166.0	151.8
1927	17.4	1.1	1.3	2.2	2.7	3.4	7.1	22.5	71.4	137.8	160.6	148.8
1928	19.0	*	1.4	2.6	3.4	3.9	8.0	24.4	74.1	149.0	181.1	169.4
1929	18.8	*	1.3	1.9	2.9	3.8	7.5	23.1	76.3	147.6	181.0	150.6
1930	19.1	*	1.5	2.0	2.8	3.5	6.8	23.3	76.6	151.3	181.1	133.8
1931	20.4	*	1.1	1.9	2.8	3.6	7.7	24.4	79.0	162.5	195.7	160.5
1932	22.0	*	1.1	2.1	3.0	3.7	7.6	25.4	84.7	171.2	212.9	171.7
1933	21.4	*	1.0	1.7	2.5	3.1	6.7	23.5	81.3	173.4	214.9	159.3
1934	22.2	*	1.0	1.6	2.3	3.1	6.6	24.2	84.0	174.1	223.6	171.7
1935	22.3	*	1.1	1.5	2.5	2.7	6.3	24.3	80.6	173.6	229.1	182.5
1936	23.7	1.4	1.2	1.5	2.4	2.8	6.5	25.5	83.8	182.3	244.0	193.6
1937	23.7	*	0.8	1.4	2.4	2.9	6.7	25.9	83.6	174.6	242.6	171.8
1938	23.9	1.5	0.9	1.4	2.2	2.6	6.3	24.0	79.7	179.0	251.2	191.5
1939	25.5	1.3	0.8	1.5	2.3	2.6	6.4	25.7	85.0	185.1	264.0	204.9

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	11.2	*	*	3.1	4.2	5.3	7.8	15.4	37.4	70.5	88.2	*
1901	11.4	*	2.9	3.9	4.2	6.3	8.3	18.8	37.3	65.5	65.4	*
1902	11.4	*	*	3.5	4.4	4.3	7.6	16.0	40.2	76.5	83.3	*
1903	12.2	*	*	3.6	4.1	6.0	7.5	18.6	42.8	76.2	96.8	*
1904	13.8	*	2.8	4.2	3.6	6.0	9.1	23.4	51.0	87.8	93.9	*
1905	13.3	*	2.8	3.7	4.6	6.1	8.1	19.2	51.8	82.3	97.6	*
1906	12.2	*	2.3	3.2	4.4	3.6	8.7	21.4	47.2	75.7	91.3	87.6
1907	12.9	*	2.1	3.4	4.9	5.5	7.9	18.4	50.3	87.4	96.4	123.2
1908	12.5	*	1.9	3.0	4.7	4.9	6.5	18.9	49.4	91.0	92.1	91.2
1909	13.0	*	2.3	3.6	4.0	5.1	8.1	21.0	47.0	86.6	106.6	87.7
1910	14.0	*	2.9	3.6	4.7	5.0	8.3	22.5	54.6	94.8	107.3	64.0
1911	14.1	*	1.9	3.7	4.7	5.3	9.1	19.5	56.7	93.6	112.4	98.5
1912	13.7	*	1.7	3.8	4.7	5.3	8.3	22.1	49.8	89.5	116.9	88.3
1913	14.0	*	2.4	3.9	4.9	5.5	8.7	20.2	53.8	88.3	109.7	110.4
1914	14.6	*	2.3	3.3	5.8	6.6	8.0	21.8	53.6	92.7	123.8	106.2
1915	15.5	*	2.3	3.9	5.0	6.1	8.7	22.3	58.3	104.9	125.8	97.2
1916	15.0	*	2.6	3.0	4.6	5.2	9.0	21.3	58.0	104.1	128.4	124.9
1917	15.4	*	3.0	3.0	5.2	6.4	7.8	21.1	59.8	102.2	136.5	133.6
1918	15.0	*	2.1	3.3	6.6	8.6	9.2	20.6	49.5	93.3	109.5	107.3
1919	13.4	*	2.4	3.1	5.1	5.5	7.6	17.7	48.3	92.2	104.8	119.4
1920	14.4	2.1	2.4	3.3	5.5	6.0	8.3	19.4	50.9	93.9	131.9	124.3
1921	14.5	*	2.3	3.9	5.1	6.5	8.4	18.7	51.8	93.7	125.7	133.8
1922	15.9	*	2.2	3.6	5.9	7.0	9.1	20.5	57.3	106.5	139.7	139.1
1923	15.3	*	2.0	2.3	4.1	5.0	7.5	20.0	56.9	113.2	153.6	130.5
1924	13.5	*	1.9	2.0	3.2	3.3	5.8	17.2	50.9	103.0	140.5	161.4
1925	13.8	*	1.4	2.1	3.1	3.2	5.9	18.3	52.6	105.2	142.2	124.3
1926	14.7	*	1.9	2.4	3.6	3.2	6.2	18.7	52.3	111.9	156.2	151.3
1927	14.0	*	1.1	2.0	2.8	3.3	5.5	17.0	53.1	107.0	144.5	155.6
1928	15.2	*	1.3	2.3	3.5	3.7	6.4	18.4	53.4	116.9	157.8	194.0
1929	15.1	*	1.2	2.1	3.0	3.6	6.7	16.6	55.8	114.3	159.7	162.1
1930	15.0	*	1.4	1.8	2.7	3.5	5.4	17.1	53.3	117.1	163.2	150.2
1931	16.1	*	1.2	1.7	2.9	3.6	6.3	17.7	54.9	127.5	166.6	167.4
1932	17.3	*	1.3	2.0	3.0	3.8	6.3	19.3	59.5	129.9	185.6	171.9
1933	17.0	*	1.1	1.4	2.7	3.1	5.7	16.8	59.8	135.1	188.8	164.0
1934	17.2	*	1.1	1.5	2.3	2.9	5.5	17.8	59.2	131.6	193.1	177.3
1935	17.2	*	1.1	1.3	2.3	2.7	5.5	18.8	56.1	128.0	196.1	185.8
1936	18.3	*	1.2	1.3	2.4	2.6	5.5	18.4	58.7	136.1	215.9	195.7
1937	18.0	*	0.9	1.4	2.4	2.7	5.5	18.4	57.9	128.1	207.0	180.1
1938	17.8	2.0	0.8	1.2	1.9	2.4	5.3	17.4	53.2	130.2	210.9	181.7
1939	19.1	*	0.8	1.3	2.2	2.5	4.9	18.5	57.6	134.6	227.3	207.4

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	10.7	*	*	3.4	2.9	4.4	5.7	17.3	43.0	63.3	57.2	*
1901	11.7	*	*	4.0	2.9	4.4	7.9	19.6	48.3	67.2	68.6	*
1902	12.0	*	2.5	3.9	2.9	4.2	6.7	17.4	49.2	76.6	82.2	*
1903	13.1	*	*	3.7	3.1	5.1	6.7	20.2	58.5	85.1	76.6	*
1904	14.7	*	*	4.4	3.4	4.4	7.6	24.2	60.6	94.5	103.6	*
1905	14.9	*	*	4.4	3.1	3.8	7.0	24.8	65.4	101.0	88.7	84.0
1906	14.7	*	2.0	3.5	3.5	4.9	6.8	24.0	67.7	105.2	90.1	57.4
1907	15.6	*	*	3.9	3.8	4.6	7.8	25.0	73.3	113.8	90.9	*
1908	15.1	*	2.3	4.4	3.4	4.4	7.8	26.4	66.2	105.2	96.7	56.5
1909	15.4	*	2.0	3.6	3.1	4.4	8.1	22.4	70.4	113.4	107.4	64.4
1910	16.6	*	2.0	3.9	4.5	5.3	8.3	24.9	76.1	114.2	116.4	82.0
1911	16.2	*	1.4	3.6	3.7	4.9	8.6	24.5	70.7	122.4	123.0	77.9
1912	16.5	*	1.7	3.9	4.2	4.8	8.4	23.8	72.4	122.1	131.2	64.5
1913	16.9	*	1.9	3.7	4.3	5.0	7.5	27.6	73.2	124.3	124.8	84.4
1914	17.9	*	1.9	4.1	3.6	4.7	8.5	29.2	76.0	139.6	127.6	86.9
1915	19.9	*	2.0	4.8	4.5	5.8	9.1	30.2	87.5	146.1	148.6	83.3
1916	18.9	*	2.1	3.2	3.8	4.6	8.8	30.2	87.0	143.8	143.8	92.4
1917	18.4	*	2.2	3.8	3.9	5.3	8.7	26.8	84.9	136.5	138.5	114.3
1918	17.1	*	1.8	3.5	4.9	5.7	9.0	25.2	75.2	123.6	119.4	85.0
1919	16.6	*	1.8	3.1	3.7	4.7	8.5	26.1	72.5	125.5	127.7	79.2
1920	18.0	*	2.0	3.7	4.1	5.1	8.1	27.4	80.2	134.2	137.9	100.7
1921	19.0	*	1.9	3.7	4.4	5.4	8.5	29.2	87.7	135.3	139.5	136.1
1922	20.8	*	1.9	4.1	4.2	5.6	9.5	32.4	91.5	154.6	160.9	149.4
1923	20.2	*	1.2	2.7	3.2	4.3	9.3	31.3	91.4	157.6	161.2	129.7
1924	19.4	*	1.3	2.5	2.9	3.6	8.1	28.2	88.4	151.6	165.6	138.6
1925	19.8	*	1.2	2.2	2.7	3.5	8.1	28.4	88.6	156.9	179.4	124.4
1926	21.2	*	1.3	2.4	3.5	3.6	7.7	29.1	95.3	171.8	175.1	152.2
1927	21.0	*	1.6	2.5	2.7	3.6	8.8	28.5	91.4	169.6	175.6	145.6
1928	22.8	*	1.5	2.8	3.4	4.1	9.6	30.8	96.8	182.3	202.8	150.6
1929	22.5	*	1.3	1.7	2.7	3.9	8.4	30.2	98.5	182.1	201.0	141.9
1930	23.2	*	1.6	2.2	2.9	3.5	8.2	30.1	101.8	186.8	197.9	121.4
1931	24.9	*	1.1	2.1	2.6	3.5	9.1	31.6	105.0	198.7	223.0	155.3
1932	26.8	*	0.9	2.2	3.0	3.6	8.9	32.1	111.8	213.7	238.6	171.5
1933	25.8	*	0.9	1.9	2.3	3.0	7.7	30.7	104.3	212.7	239.5	155.8
1934	27.2	*	0.9	1.8	2.3	3.3	7.7	31.1	110.4	217.5	252.2	167.6
1935	27.5	*	1.1	1.6	2.7	2.7	7.1	30.2	106.8	220.0	260.0	180.0
1936	29.3	*	1.2	1.6	2.5	2.9	7.5	33.1	110.5	229.2	270.1	192.1
1937	29.6	*	0.7	1.5	2.4	3.0	7.8	33.9	110.8	221.5	275.5	165.5
1938	30.1	*	1.0	1.6	2.4	2.8	7.2	31.0	107.7	227.9	288.3	198.8
1939	32.0	*	0.8	1.6	2.4	2.8	7.8	33.5	113.9	235.4	297.6	203.1

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	15.5	---	---	---	---	---	---	---	---	---	---	---
1911	15.4	---	---	---	---	---	---	---	---	---	---	---
1912	15.2	---	---	---	---	---	---	---	---	---	---	---
1913	15.7	---	---	---	---	---	---	---	---	---	---	---
1914	16.6	*	2.1	3.8	4.8	5.8	8.3	25.5	64.8	117.9	127.6	95.9
1915	17.9	*	2.2	4.5	4.8	5.9	8.8	26.2	72.7	127.1	139.5	92.1
1916	17.5	*	2.4	3.3	4.3	5.1	9.1	25.9	73.0	126.7	139.7	110.9
1917	17.6	---	---	---	---	---	---	---	---	---	---	---
1918	16.7	*	2.1	3.6	5.9	7.3	9.3	23.0	62.8	111.7	117.9	96.7
1919	15.8	*	2.3	3.4	4.7	5.4	8.2	21.9	61.5	112.8	120.3	100.5
1920	16.9	1.2	2.4	3.8	5.1	5.8	8.2	23.3	66.2	117.4	139.2	111.4
1921	17.5	*	2.2	4.1	5.0	6.1	8.5	23.7	70.9	117.2	137.8	140.3
1922	19.3	1.8	2.2	4.2	5.4	6.6	9.2	26.2	75.9	135.0	155.3	148.6
1923	18.6	1.1	1.8	2.7	3.9	4.7	8.1	25.2	75.5	139.5	162.2	132.0
1924	17.0	*	1.6	2.4	3.1	3.4	6.3	22.0	70.3	130.4	156.9	154.8
1925	17.4	*	1.4	2.2	3.0	3.2	6.5	22.3	71.5	134.1	166.8	125.4
1926	18.5	*	1.7	2.5	3.7	3.3	6.3	22.7	74.3	144.3	171.0	152.8
1927	18.0	*	1.4	2.3	2.8	3.3	6.5	21.6	72.3	141.4	165.1	154.2
1928	19.7	*	1.5	2.7	3.5	3.7	7.4	23.2	75.2	153.4	186.3	171.4
1929	19.5	*	1.3	2.0	2.9	3.6	6.9	22.0	77.4	152.0	186.7	157.1
1930	19.8	*	1.5	2.1	2.9	3.4	6.1	21.9	77.3	155.6	187.7	138.4
1931	21.3	*	1.2	2.0	2.8	3.3	7.0	23.2	80.2	167.4	202.0	166.6
1932	22.9	*	1.2	2.2	3.1	3.5	6.9	24.1	85.7	176.4	220.6	179.2
1933	22.2	*	1.1	1.7	2.5	2.8	5.9	22.0	81.9	178.2	222.1	166.9
1934	23.0	*	1.0	1.7	2.4	2.9	5.8	22.8	84.8	179.8	231.1	181.3
1935	23.3	*	1.1	1.5	2.5	2.6	5.6	23.1	81.3	180.1	237.2	190.0
1936	24.8	1.3	1.3	1.5	2.5	2.6	5.7	24.1	84.4	189.1	251.9	204.8
1937	24.7	*	0.8	1.4	2.3	2.6	5.9	24.3	84.7	181.2	251.9	178.2
1938	24.8	1.4	0.9	1.5	2.2	2.3	5.4	22.6	79.8	185.5	260.6	202.1
1939	26.4	1.5	0.9	1.5	2.2	2.4	5.3	23.8	84.5	192.2	273.5	218.1

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

WHITE, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	14.2	---	---	---	---	---	---	---	---	---	---	---
1911	14.3	---	---	---	---	---	---	---	---	---	---	---
1912	13.8	---	---	---	---	---	---	---	---	---	---	---
1913	14.2	---	---	---	---	---	---	---	---	---	---	---
1914	14.9	*	2.3	3.4	5.9	6.8	8.1	21.9	54.0	94.3	125.5	111.2
1915	15.7	*	2.4	4.0	5.1	6.0	8.8	22.8	58.7	107.0	127.1	97.9
1916	15.5	*	2.8	3.2	4.7	5.4	9.2	21.8	58.9	106.5	131.3	129.9
1917	16.0	---	---	---	---	---	---	---	---	---	---	---
1918	15.6	*	2.2	3.5	6.8	8.9	9.5	21.0	50.6	96.2	111.9	106.9
1919	14.1	*	2.6	3.4	5.4	5.8	7.9	18.1	49.7	95.8	106.6	122.4
1920	15.0	*	2.5	3.5	5.8	6.3	8.4	19.7	52.2	97.1	135.9	121.5
1921	15.2	*	2.5	4.2	5.5	6.7	8.6	19.0	53.3	95.8	130.0	138.6
1922	16.8	*	2.4	3.9	6.3	7.3	9.2	20.7	59.5	110.6	143.4	141.5
1923	16.0	*	2.2	2.4	4.4	5.1	7.5	19.9	58.9	117.3	157.2	131.8
1924	14.0	*	1.9	2.1	3.1	3.3	5.5	17.2	52.0	105.8	144.4	168.2
1925	14.3	*	1.5	2.2	3.1	3.2	5.7	18.0	54.3	108.4	145.3	127.8
1926	15.2	*	1.9	2.5	3.7	3.2	5.8	18.4	53.4	114.3	160.2	146.5
1927	14.4	*	1.2	2.1	2.8	3.2	5.3	16.5	53.7	109.5	147.8	155.2
1928	15.7	*	1.4	2.5	3.5	3.6	6.2	18.0	54.3	120.1	161.5	195.3
1929	15.6	*	1.3	2.1	3.0	3.5	6.4	16.1	57.2	117.5	162.5	163.3
1930	15.6	*	1.4	1.9	2.8	3.4	5.2	16.5	53.9	120.7	169.5	155.0
1931	16.7	*	1.3	1.7	2.9	3.4	6.1	17.7	56.1	130.8	170.6	170.5
1932	18.0	*	1.4	2.1	3.0	3.7	5.9	18.7	60.7	133.4	191.6	181.3
1933	17.7	*	1.2	1.5	2.6	3.0	5.1	16.0	61.0	139.6	194.7	170.3
1934	18.0	*	1.1	1.5	2.3	3.0	5.0	17.3	60.4	136.1	199.8	187.0
1935	17.9	*	1.2	1.3	2.2	2.5	5.1	18.5	57.4	132.8	202.3	193.4
1936	19.1	*	1.2	1.3	2.5	2.5	5.4	17.8	59.6	141.1	222.1	204.8
1937	18.8	*	0.9	1.3	2.3	2.4	5.3	18.0	58.9	132.7	214.4	183.2
1938	18.6	*	0.9	1.3	1.9	2.3	5.1	17.0	53.9	135.3	218.7	193.7
1939	19.9	*	0.9	1.3	2.1	2.4	4.6	17.8	58.0	140.1	235.5	220.3

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	16.8	---	---	---	---	---	---	---	---	---	---	---
1911	16.5	---	---	---	---	---	---	---	---	---	---	---
1912	16.7	---	---	---	---	---	---	---	---	---	---	---
1913	17.2	---	---	---	---	---	---	---	---	---	---	---
1914	18.3	*	1.9	4.3	3.7	4.7	8.6	29.6	76.5	141.8	129.5	84.0
1915	20.2	*	2.1	4.9	4.6	5.9	8.9	30.1	88.0	147.4	150.8	87.5
1916	19.6	*	2.1	3.4	4.0	4.7	8.9	30.6	88.4	147.1	147.2	96.4
1917	19.1	---	---	---	---	---	---	---	---	---	---	---
1918	17.8	*	1.9	3.8	5.1	5.8	9.1	25.3	76.1	127.7	123.4	88.9
1919	17.6	*	1.9	3.4	4.0	4.9	8.6	26.1	74.4	130.1	132.7	84.0
1920	18.8	*	2.2	4.0	4.4	5.2	8.0	27.4	81.4	138.1	142.3	103.9
1921	19.9	*	2.0	4.0	4.6	5.4	8.4	28.9	89.9	139.2	144.9	141.5
1922	21.9	*	2.1	4.5	4.6	5.9	9.2	32.3	93.5	159.8	166.3	154.0
1923	21.2	*	1.3	2.9	3.4	4.3	8.8	31.1	93.2	162.2	166.7	132.1
1924	20.1	*	1.3	2.7	3.0	3.5	7.3	27.3	89.9	155.6	168.5	144.3
1925	20.5	*	1.3	2.3	2.8	3.2	7.4	27.0	90.2	160.6	186.6	123.6
1926	22.0	*	1.4	2.5	3.6	3.5	6.9	27.4	96.9	175.3	181.0	157.6
1927	21.7	*	1.7	2.5	2.8	3.4	7.9	27.0	92.3	174.2	181.1	153.4
1928	23.7	*	1.7	2.9	3.6	3.8	8.7	28.8	97.7	187.7	209.3	153.0
1929	23.4	*	1.4	1.8	2.8	3.7	7.4	28.4	99.1	187.4	209.2	152.3
1930	24.0	*	1.7	2.2	2.9	3.3	7.0	27.8	102.2	191.7	204.9	125.9
1931	25.9	*	1.1	2.3	2.7	3.2	7.9	29.2	105.8	205.1	231.4	163.7
1932	27.9	*	1.0	2.3	3.2	3.3	7.9	30.0	112.1	220.3	247.8	177.7
1933	26.8	*	0.9	2.0	2.4	2.7	6.8	28.4	104.1	217.6	248.0	164.3
1934	28.2	*	0.9	1.9	2.4	2.9	6.5	28.6	110.6	224.1	260.4	177.0
1935	28.7	*	1.1	1.7	2.8	2.6	6.2	28.1	106.6	227.9	269.6	187.5
1936	30.5	*	1.3	1.7	2.5	2.7	6.1	30.7	110.5	237.4	279.5	204.9
1937	30.8	*	0.7	1.5	2.4	2.7	6.5	31.0	111.9	229.8	286.5	174.4
1938	31.2	*	1.0	1.6	2.5	2.3	5.7	28.6	107.0	235.3	299.1	208.5
1939	33.0	*	0.9	1.6	2.3	2.3	6.0	30.3	112.2	243.8	308.3	216.4

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	7.2	---	---	---	---	---	---	---	---	---	---	---
1911	7.0	---	---	---	---	---	---	---	---	---	---	---
1912	9.7	---	---	---	---	---	---	---	---	---	---	---
1913	8.7	---	---	---	---	---	---	---	---	---	---	---
1914	8.8	*	*	*	*	*	6.3	19.1	49.8	52.9	*	*
1915	10.9	*	*	*	*	*	5.6	10.4	21.6	58.6	69.2	*
1916	7.9	*	*	*	2.7	3.1	6.9	18.0	47.3	59.2	60.2	*
1917	7.8	---	---	---	---	---	---	---	---	---	---	---
1918	7.7	*	*	*	3.3	4.4	6.7	18.9	41.9	39.9	47.8	*
1919	6.4	*	*	*	1.8	2.7	6.2	18.6	31.4	38.5	55.0	*
1920	8.0	*	*	*	2.1	3.7	7.8	20.7	42.3	50.5	59.2	*
1921	8.2	*	*	*	2.5	4.6	7.5	22.3	37.0	60.5	45.3	*
1922	8.8	*	*	*	1.1	1.7	3.8	10.1	25.2	41.5	55.0	78.1
1923	9.3	*	*	1.2	1.5	4.2	10.9	26.9	43.4	61.5	81.5	*
1924	10.4	*	*	1.2	2.7	4.3	12.5	27.2	48.2	66.2	98.8	*
1925	10.6	*	*	1.2	2.7	5.1	12.0	32.1	44.2	66.5	74.0	113.4
1926	11.8	*	*	*	2.5	4.1	13.8	35.5	50.9	85.7	81.7	142.1
1927	11.7	*	*	1.5	2.5	5.0	13.3	33.0	58.0	72.5	86.1	*
1928	12.6	*	*	1.5	2.7	5.5	13.7	36.4	59.7	75.5	101.7	151.9
1929	12.2	*	*	1.3	2.6	5.3	14.1	34.3	60.6	73.6	92.4	92.3
1930	12.8	*	*	1.3	2.3	4.4	13.7	37.8	67.4	77.2	71.9	91.8
1931	13.0	*	*	1.2	2.5	5.6	14.3	36.3	63.1	79.4	91.8	104.6
1932	13.8	*	*	1.2	2.4	5.2	13.8	39.3	71.1	86.0	84.7	102.1
1933	14.3	*	*	1.2	2.6	4.8	13.7	38.6	72.5	98.9	97.5	92.3
1934	14.3	*	*	1.2	1.8	4.7	14.5	39.4	72.9	88.7	101.0	86.6
1935	13.6	*	*	1.3	2.2	3.7	12.1	36.6	71.3	79.4	96.8	115.2
1936	14.7	*	*	1.2	1.6	3.9	13.4	40.8	76.0	86.8	112.2	93.7
1937	14.9	*	*	1.6	2.9	5.3	13.2	42.4	68.2	84.4	86.9	114.3
1938	15.6	*	*	1.3	2.3	4.6	14.0	39.2	78.6	94.6	93.1	96.4
1939	17.6	*	*	1.5	2.7	4.8	15.4	46.2	92.0	94.3	102.1	87.7

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	7.3	...	...	...	...	...	...	...	...	...	...	...
1911	6.9	...	...	...	...	...	...	...	...	...	...	...
1912	9.0	...	...	...	...	...	...	...	...	...	...	...
1913	8.5	...	...	...	...	...	...	...	...	...	...	...
1914	8.7	*	*	*	*	*	*	20.1	42.5	*	*	*
1915	9.4	*	*	*	*	8.1	*	*	48.8	*	*	*
1916	7.4	*	*	*	*	*	*	14.9	40.3	54.0	*	*
1917	7.0	...	...	...	...	...	...	...	...	...	...	...
1918	7.5	*	*	*	4.5	4.4	6.0	15.2	30.9	40.6	*	*
1919	6.1	*	*	*	*	*	4.4	13.6	26.5	35.5	74.7	*
1920	7.2	*	*	*	*	*	6.9	15.8	30.1	41.4	*	*
1921	7.1	*	*	*	*	3.9	5.9	14.3	29.2	60.5	*	*
1922	7.8	*	*	*	*	4.0	7.8	18.5	27.6	47.2	84.8	*
1923	8.0	*	*	*	*	3.5	7.0	21.8	30.0	51.7	96.8	*
1924	8.6	*	*	*	3.8	3.4	8.5	17.6	36.4	58.3	80.6	*
1925	8.7	*	*	*	3.5	3.9	8.5	21.0	29.7	51.7	93.5	*
1926	9.9	*	*	*	2.8	3.8	10.8	22.7	36.8	71.8	91.6	*
1927	9.7	*	*	*	2.8	4.1	8.1	21.9	44.3	65.0	91.7	*
1928	10.2	*	*	*	3.5	4.6	9.0	23.1	41.9	66.6	102.7	*
1929	10.0	*	*	1.7	3.2	5.0	9.8	21.4	38.3	62.4	116.2	*
1930	9.4	*	*	*	1.9	4.1	7.7	23.3	45.3	57.9	60.4	*
1931	9.7	*	*	*	2.7	5.0	8.4	18.2	39.7	73.9	100.9	*
1932	10.7	*	*	*	3.0	4.4	10.1	25.7	43.0	73.3	86.0	*
1933	10.8	*	*	*	3.2	4.6	11.4	23.9	44.2	67.8	94.2	*
1934	10.2	*	*	*	2.2	2.8	10.2	22.8	44.4	67.3	84.5	*
1935	10.1	*	*	1.7	2.7	3.6	8.5	22.7	40.3	59.8	93.8	*
1936	10.5	*	*	*	*	3.1	6.5	24.4	48.1	68.2	113.5	*
1937	11.0	*	*	1.6	3.3	5.3	7.6	21.9	46.0	67.2	83.9	151.7
1938	10.2	*	*	*	2.3	3.0	7.4	22.2	45.5	66.3	81.8	*
1939	11.6	*	*	1.5	2.6	3.4	7.7	25.6	52.6	68.3	91.4	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DIABETES MELLITUS

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	7.2	---	---	---	---	---	---	---	---	---	---	---
1911	7.1	---	---	---	---	---	---	---	---	---	---	---
1912	10.4	---	---	---	---	---	---	---	---	---	---	---
1913	9.0	---	---	---	---	---	---	---	---	---	---	---
1914	8.9	*	*	*	*	*	*	*	59.0	*	*	*
1915	12.5	*	*	*	*	*	13.1	33.6	71.0	101.8	*	*
1916	8.4	*	*	*	*	*	7.7	22.0	56.3	65.3	*	*
1917	8.5	---	---	---	---	---	---	---	---	---	---	---
1918	7.9	*	*	*	*	4.5	7.4	24.0	56.0	39.0	*	*
1919	6.6	*	*	*	*	*	8.0	25.5	37.8	42.1	*	*
1920	8.8	*	*	*	*	4.1	8.8	27.3	58.2	61.5	*	*
1921	9.2	*	*	*	3.0	5.2	9.3	32.9	47.2	60.5	*	*
1922	9.8	*	*	*	*	3.6	12.5	33.9	59.9	64.6	71.3	*
1923	10.7	*	*	*	*	4.8	14.9	33.3	60.9	73.7	66.0	*
1924	12.1	*	*	*	*	5.2	16.8	39.1	63.5	75.9	117.1	*
1925	12.5	*	*	*	*	6.3	15.8	45.6	63.0	84.3	54.6	*
1926	13.7	*	*	*	2.3	4.3	16.9	50.8	69.1	102.4	71.9	*
1927	13.8	*	*	1.8	2.1	5.9	18.6	46.1	76.0	81.3	80.6	*
1928	14.9	*	*	1.7	1.9	6.3	18.2	51.9	83.1	86.0	100.8	*
1929	14.4	*	*	*	2.1	5.5	18.3	49.2	90.0	86.5	69.3	*
1930	16.1	*	*	*	2.7	4.7	19.7	54.5	96.3	99.1	82.9	*
1931	16.2	*	*	*	2.3	6.1	20.0	57.0	93.2	85.6	83.2	*
1932	16.8	*	*	*	1.9	6.0	17.4	54.7	106.9	100.2	83.5	*
1933	17.6	*	*	1.4	2.0	5.0	15.8	55.2	107.9	134.1	100.7	*
1934	18.3	*	*	*	1.5	6.5	18.7	57.9	108.4	112.5	116.6	*
1935	17.0	*	*	*	1.7	3.8	15.5	51.9	109.3	101.1	99.6	114.0
1936	18.8	*	*	*	1.7	4.7	20.1	58.8	109.8	107.2	111.0	*
1937	18.7	*	*	1.5	2.5	5.4	18.5	64.6	94.8	103.4	89.7	*
1938	20.9	*	*	1.6	2.3	6.2	20.3	57.6	117.5	125.5	103.9	113.3
1939	23.4	*	*	1.5	2.8	6.1	22.7	68.4	137.8	122.6	112.3	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both Sexes...	39	62	47
Male...	14	21	18
Female	25	41	29

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ICD Revision	Years in use	ICD Nos.
Fifth	1939-58	83, 90-103
Fourth	1930-1938	56, 82, 90-95, 97-103
Third	1921-1929	51, 74, 75, 83, 87-90, 91b,c, 92-96, 151
Second	1910-1920	47, 64-66, 77-85, 142
First	1900-1909	64-66, 77-86, 142

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	264.3	329.8	394.4	33.7	42.0	56.3	132.4	311.3	776.6	1,899.2	3,912.0	5,825.5
1901	261.5	291.3	30.5	25.2	35.8	63.5	128.2	309.4	795.1	1,880.2	4,069.9	5,800.7
1902	264.9	278.9	28.9	26.1	35.0	64.1	130.9	311.2	803.6	1,940.8	4,050.4	6,058.7
1903	273.5	240.3	24.8	23.3	34.9	61.5	130.4	314.4	834.4	2,016.3	4,411.9	6,602.6
1904	290.6	236.1	27.1	27.6	35.4	61.6	133.3	321.2	896.1	2,166.6	4,702.8	7,494.3
1905	287.6	244.7	23.0	27.0	35.1	60.9	135.2	328.1	872.3	2,139.0	4,642.2	7,427.3
1906	272.4	212.9	21.8	24.7	33.8	59.8	130.5	311.1	853.6	2,115.1	4,648.0	7,881.1
1907	293.7	199.4	21.5	24.8	35.1	61.1	136.1	333.5	917.9	2,297.5	5,138.8	8,638.4
1908	270.4	189.4	20.4	24.5	31.4	54.2	118.7	300.5	843.9	2,140.5	4,798.0	8,203.6
1909	274.3	191.6	18.6	23.8	29.2	51.4	118.2	295.1	829.8	2,166.9	4,932.1	8,470.2
1910	287.2	182.1	22.2	31.5	35.5	55.6	125.9	310.3	852.7	2,243.4	5,175.5	8,819.2
1911	281.9	156.7	20.0	30.0	33.9	55.5	124.6	303.0	833.4	2,234.5	5,105.5	8,932.9
1912	286.0	132.6	18.4	26.0	33.5	54.2	126.0	311.4	849.3	2,254.9	5,204.7	9,114.6
1913	280.8	120.3	18.8	28.6	32.2	54.1	123.1	305.5	814.0	2,207.6	5,172.1	8,944.1
1914	283.8	102.8	17.7	29.3	32.5	52.7	120.8	299.1	817.2	2,268.4	5,127.2	9,205.9
1915	290.5	104.5	16.4	27.4	32.2	50.9	120.3	299.7	830.3	2,304.0	5,318.6	9,762.2
1916	294.3	94.0	17.1	26.1	32.6	55.2	124.6	313.4	852.3	2,325.7	5,452.8	9,925.1
1917	299.4	94.3	16.5	25.4	33.8	56.7	127.7	313.6	860.2	2,351.5	5,559.5	10,133.2
1918	296.8	93.2	16.9	25.7	42.5	68.7	131.9	311.5	839.6	2,255.8	5,203.2	9,521.4
1919	266.8	79.4	14.0	24.0	29.1	45.8	105.3	272.4	758.0	2,079.8	5,091.1	9,774.0
1920	282.5	87.1	15.7	23.5	31.2	48.4	104.8	282.1	788.7	2,202.4	5,610.8	11,089.8
1921	273.1	63.0	15.1	24.8	30.5	42.7	98.5	276.1	759.4	2,127.6	5,391.6	10,928.8
1922	284.9	65.5	14.3	21.1	28.8	43.0	104.6	292.2	794.0	2,247.4	5,832.7	11,656.4
1923	297.9	68.0	13.7	20.9	28.6	43.5	109.5	309.0	826.6	2,269.5	5,818.9	12,379.7
1924	301.5	62.9	12.8	22.2	30.2	44.5	113.8	319.7	820.0	2,265.4	5,838.6	12,236.8
1925	301.8	60.7	13.7	21.5	28.7	45.0	112.1	313.6	812.3	2,232.3	5,783.0	12,795.5
1926	318.3	56.1	12.9	19.4	28.9	45.8	116.9	329.4	848.5	2,340.1	6,075.8	13,231.4
1927	311.3	53.4	12.2	19.6	28.3	45.1	116.1	323.6	828.5	2,284.9	5,801.6	12,276.8
1928	328.9	51.9	12.2	18.0	27.2	48.7	124.6	344.5	878.5	2,407.1	6,122.9	12,723.6
1929	332.4	49.8	12.4	18.2	27.9	47.9	122.7	348.3	898.5	2,411.6	5,987.1	12,065.5
1930	327.8	47.3	10.5	16.1	26.3	47.3	119.0	343.9	896.5	2,332.4	5,701.6	11,359.5
1931	323.5	36.8	10.2	15.1	24.1	44.6	116.9	336.6	877.9	2,249.1	5,549.3	11,090.3
1932	334.4	28.8	9.3	14.2	22.7	43.3	114.9	333.4	887.9	2,288.8	5,769.9	11,578.9
1933	334.3	28.9	9.9	13.5	21.4	42.3	112.4	337.8	895.5	2,261.0	5,701.7	11,192.7
1934	349.2	28.5	9.0	12.6	21.4	42.6	118.2	352.7	912.4	2,294.8	5,896.6	11,574.9
1935	353.4	43.3	9.1	14.3	20.7	41.3	116.5	347.2	905.6	2,274.6	5,877.3	11,571.6
1936	380.9	45.2	8.8	14.2	21.1	41.6	123.9	362.8	949.7	2,378.7	6,364.6	12,511.2
1937	377.9	34.8	8.2	12.4	20.0	40.6	120.8	362.4	934.5	2,319.8	6,147.1	11,923.5
1938	377.1	36.1	7.8	12.6	19.5	38.6	118.3	358.2	910.5	2,300.3	5,990.3	11,481.8
1939	386.0	30.4	7.3	11.9	18.6	37.3	116.3	357.0	919.4	2,301.2	6,041.1	11,986.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	259.3	376.3	43.4	29.9	40.2	62.2	125.4	306.6	817.9	2,056.2	4,211.4	6,138.9
1901	271.2	335.5	32.9	23.7	35.7	62.9	127.7	314.8	853.6	2,051.7	4,372.5	6,375.5
1902	274.9	321.0	28.1	24.1	34.1	59.8	128.7	315.7	876.6	2,135.4	4,415.6	6,480.4
1903	283.7	270.2	26.0	21.8	36.8	60.0	126.8	320.8	894.3	2,210.6	4,821.0	7,152.6
1904	299.0	270.9	26.9	24.4	35.2	57.4	132.1	325.5	977.6	2,335.1	5,118.4	7,925.3
1905	293.6	274.1	24.9	22.1	34.1	58.5	133.5	332.7	928.2	2,291.2	4,998.2	8,061.3
1906	280.5	242.3	21.5	22.3	33.7	58.4	134.8	314.3	915.7	2,279.9	5,051.2	8,209.8
1907	304.8	237.1	21.3	23.6	36.4	58.9	139.9	350.2	992.6	2,507.4	5,497.0	9,276.2
1908	276.5	205.0	18.9	21.8	30.5	50.4	118.0	309.2	901.1	2,289.4	5,197.6	8,602.0
1909	281.4	224.4	18.3	20.7	27.6	49.1	117.5	304.0	894.0	2,319.7	5,322.6	8,982.1
1910	294.1	205.3	22.5	28.9	34.2	54.5	127.3	316.2	909.1	2,418.8	5,523.9	9,377.3
1911	288.8	178.9	19.5	27.5	33.0	54.2	125.6	313.5	886.3	2,380.3	5,458.0	9,275.4
1912	295.3	151.6	18.7	22.8	32.4	51.6	126.5	328.8	917.5	2,433.1	5,537.4	9,627.9
1913	288.3	134.4	19.1	25.9	31.5	52.5	123.4	315.3	870.1	2,381.9	5,451.3	9,417.1
1914	291.9	121.0	19.0	26.9	31.8	50.2	122.8	303.4	877.5	2,438.5	5,441.9	9,547.4
1915	297.1	120.0	16.9	25.2	30.4	48.6	120.7	304.3	892.2	2,447.6	5,626.9	10,246.7
1916	302.7	103.4	18.1	24.4	31.2	54.6	125.8	326.1	917.4	2,478.7	5,732.3	10,315.1
1917	309.7	100.5	16.5	23.8	33.0	56.5	129.1	324.9	922.6	2,522.0	5,913.1	10,483.7
1918	305.0	104.3	17.5	24.0	40.5	64.8	126.9	307.9	890.2	2,377.9	5,427.1	9,820.8
1919	267.3	90.4	14.3	19.0	27.9	42.4	100.5	266.3	791.4	2,138.4	5,192.0	10,143.8
1920	280.3	102.3	16.3	21.7	29.8	42.0	98.3	275.4	813.7	2,262.6	5,743.7	11,204.3
1921	272.4	68.1	15.3	23.5	28.7	38.9	94.2	271.7	790.3	2,197.3	5,513.8	11,009.4
1922	289.0	74.2	15.1	20.0	27.6	39.9	101.6	297.0	844.1	2,357.6	5,800.8	11,634.2
1923	304.9	78.2	14.2	19.5	27.4	41.5	108.2	316.4	890.0	2,395.7	6,032.9	12,415.8
1924	311.8	68.6	13.5	21.3	29.2	42.4	115.0	333.5	888.5	2,425.3	6,086.5	12,280.1
1925	313.5	70.7	13.8	20.6	27.4	42.9	113.5	334.5	889.1	2,388.3	6,060.9	12,723.3
1926	332.5	65.5	12.8	18.5	27.4	44.5	119.2	356.0	923.8	2,535.3	6,352.8	13,393.9
1927	327.6	60.5	12.2	18.7	26.9	42.8	121.4	349.3	916.8	2,472.8	6,119.6	12,490.6
1928	347.3	59.1	12.7	17.3	26.3	47.1	130.8	375.2	972.1	2,609.0	6,454.8	12,832.8
1929	353.6	58.2	12.7	16.6	26.5	47.3	131.7	385.0	1,005.3	2,626.0	6,289.7	12,329.3
1930	349.9	53.8	10.7	15.5	24.4	46.1	127.7	382.4	1,007.2	2,543.8	5,997.3	11,703.1
1931	347.9	40.5	10.5	14.4	22.8	43.7	127.5	378.2	996.0	2,464.0	5,862.9	11,643.6
1932	358.5	29.5	9.6	13.8	22.2	41.8	126.1	376.8	1,014.2	2,494.5	6,080.6	11,874.9
1933	362.1	31.3	10.7	12.6	20.9	42.2	124.0	386.8	1,035.5	2,490.0	6,023.1	11,536.6
1934	380.7	33.6	8.8	12.0	21.1	43.5	132.0	411.1	1,062.4	2,545.6	6,258.7	11,929.9
1935	385.9	49.7	8.7	13.5	21.0	42.0	130.5	404.2	1,061.4	2,528.7	6,272.4	11,941.8
1936	416.9	54.0	9.3	14.1	21.6	42.2	139.9	428.1	1,125.6	2,648.0	6,763.2	12,952.6
1937	416.2	38.8	7.9	11.6	19.4	42.2	137.9	434.4	1,111.6	2,608.6	6,581.8	12,273.3
1938	414.8	39.3	8.6	12.1	19.2	40.5	136.9	429.7	1,086.2	2,588.0	6,361.0	11,798.3
1939	425.7	34.6	7.5	12.0	18.8	39.6	135.5	433.9	1,103.8	2,600.4	6,403.7	12,326.8

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	259.2	282.4	35.3	37.5	43.7	69.5	139.9	316.1	736.0	1,750.2	3,639.1	5,598.6
1901	251.8	246.2	28.0	26.7	35.8	64.0	128.8	303.7	737.8	1,717.8	3,795.3	5,383.3
1902	254.8	236.0	29.7	28.1	35.8	68.6	133.2	306.5	731.9	1,756.6	3,720.7	5,571.3
1903	263.3	209.9	23.6	24.8	33.2	63.0	134.3	307.7	775.4	1,832.6	4,044.4	6,200.3
1904	282.2	200.6	27.2	30.9	35.7	65.9	134.5	316.7	815.7	2,007.5	4,331.3	7,177.8
1905	281.5	214.6	21.0	32.0	36.0	63.3	137.0	323.2	817.0	1,975.9	4,325.8	6,960.2
1906	264.1	182.8	22.1	27.1	34.0	61.3	125.7	307.6	789.7	1,954.5	4,282.8	7,634.9
1907	282.2	160.8	21.6	25.9	33.9	63.4	132.0	315.2	840.5	2,092.8	4,815.9	8,158.8
1908	264.0	173.4	21.9	27.2	32.3	58.3	119.6	290.8	783.6	1,993.3	4,432.6	7,895.0
1909	266.7	158.0	19.0	26.9	30.8	53.9	119.0	285.3	761.8	2,015.3	4,575.6	8,070.6
1910	280.0	158.3	21.9	34.3	36.9	56.9	124.3	303.6	792.2	2,068.3	4,856.9	8,380.7
1911	274.5	133.8	20.5	32.6	34.7	56.8	123.6	291.1	776.2	2,087.7	4,780.8	8,661.8
1912	276.3	113.1	18.2	29.2	34.5	57.1	125.5	291.7	775.4	2,075.4	4,899.8	8,711.3
1913	273.0	105.9	18.5	31.3	32.9	55.8	122.7	294.5	753.0	2,031.5	4,917.3	8,576.0
1914	275.4	84.2	16.5	31.8	33.2	55.3	118.6	294.1	751.4	2,095.4	4,838.4	8,940.3
1915	283.5	88.7	15.9	29.6	34.0	53.5	119.8	294.5	762.5	2,157.8	5,037.1	9,388.8
1916	285.4	84.4	16.1	27.8	33.9	56.0	123.3	299.0	780.8	2,169.5	5,199.5	9,628.5
1917	288.6	87.9	16.5	26.9	34.6	56.9	126.1	300.8	791.7	2,176.7	5,238.9	9,868.4
1918	288.6	81.9	16.3	27.4	44.3	72.3	137.5	315.6	783.9	2,130.0	5,000.0	9,296.0
1919	266.3	68.1	13.7	21.8	30.1	49.3	110.5	279.3	721.2	2,019.2	4,999.6	9,497.3
1920	284.8	71.5	15.1	25.3	32.5	54.9	112.1	289.6	761.2	2,140.1	5,488.9	11,003.6
1921	273.8	57.8	14.9	26.1	32.3	46.5	103.1	281.0	725.6	2,055.6	5,279.3	10,867.7
1922	280.6	56.6	13.6	22.3	30.0	46.1	107.8	286.9	739.8	2,133.8	5,478.7	11,673.2
1923	290.6	57.5	13.2	22.5	29.8	45.6	111.0	300.8	757.7	2,138.9	5,622.2	12,352.1
1924	290.9	56.9	12.1	23.1	31.1	46.7	112.5	304.5	745.4	2,099.4	5,610.3	12,203.7
1925	289.7	50.3	13.6	22.3	30.0	47.1	110.5	290.7	728.6	2,070.5	5,526.5	12,851.0
1926	303.6	46.5	13.1	20.3	30.3	47.1	114.5	300.4	766.4	2,137.9	5,819.8	13,106.4
1927	294.6	46.1	12.2	20.5	29.7	47.5	110.4	295.6	731.9	2,090.2	5,506.4	12,112.4
1928	310.1	44.5	11.7	18.8	28.2	50.3	118.0	311.2	776.1	2,197.7	5,812.9	12,640.0
1929	310.7	41.1	12.0	19.8	29.2	48.4	113.3	308.3	782.3	2,189.3	5,703.2	11,865.6
1930	305.1	40.6	10.3	16.8	28.0	48.4	109.9	301.9	777.0	2,113.0	5,423.1	11,101.5
1931	298.6	33.0	9.8	15.8	25.3	45.6	105.8	291.3	750.9	2,027.1	5,254.6	10,674.9
1932	309.7	28.1	9.0	14.7	23.2	44.7	103.1	286.4	752.3	2,077.2	5,478.6	11,356.8
1933	306.0	26.4	9.0	14.5	21.8	42.5	103.3	284.7	745.3	2,025.8	5,399.4	10,933.7
1934	317.2	23.2	9.1	13.1	21.8	41.7	103.9	289.7	752.0	2,038.4	5,557.3	11,307.7
1935	320.3	36.6	9.5	15.0	20.5	40.7	102.1	285.9	739.3	2,016.2	5,508.7	11,293.0
1936	344.4	36.1	8.4	14.3	20.6	41.0	107.5	292.7	762.6	2,106.3	5,994.2	12,179.0
1937	339.0	30.7	8.4	13.2	20.6	39.1	103.3	285.4	746.5	2,029.1	5,745.1	11,660.2
1938	338.8	32.9	6.9	13.2	19.8	36.7	99.3	282.1	724.5	2,012.3	5,649.2	11,243.7
1939	345.8	26.1	7.1	11.8	18.3	35.0	96.9	275.2	724.8	2,003.3	5,709.1	11,729.9

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	285.7	---	---	---	---	---	---	---	---	---	---	---
1911	279.5	---	---	---	---	---	---	---	---	---	---	---
1912	283.4	---	---	---	---	---	---	---	---	---	---	---
1913	278.2	---	---	---	---	---	---	---	---	---	---	---
1914	280.6	101.3	17.3	29.4	31.5	48.8	110.9	280.4	796.7	2,257.2	5,149.2	9,297.7
1915	287.2	103.7	16.3	27.4	31.3	47.1	110.3	282.6	805.8	2,290.6	5,333.3	9,881.9
1916	293.2	94.4	17.1	26.6	31.5	49.6	112.4	293.7	825.6	2,317.3	5,497.4	10,126.7
1917	297.1	---	---	---	---	---	---	---	---	---	---	---
1918	295.2	93.7	16.6	26.1	40.8	62.8	116.9	288.5	811.3	2,241.5	5,250.8	9,759.6
1919	267.5	81.5	13.8	20.9	27.1	40.7	91.8	248.2	737.1	2,078.8	5,165.7	10,190.5
1920	283.4	88.7	15.8	24.1	29.6	42.5	91.1	258.2	766.7	2,197.2	5,687.3	11,499.9
1921	273.8	63.2	15.1	25.3	28.8	37.2	84.4	251.6	736.7	2,125.6	5,466.8	11,325.8
1922	286.7	66.9	14.3	21.6	27.4	37.0	88.9	261.8	770.3	2,252.1	5,738.0	12,159.0
1923	298.5	68.7	13.7	21.3	27.2	37.5	92.7	274.4	798.3	2,263.4	5,917.9	12,817.4
1924	299.6	62.4	12.7	22.5	28.1	37.4	94.3	280.4	785.1	2,248.4	5,918.7	12,602.7
1925	299.6	60.1	13.5	21.7	26.4	37.9	92.5	273.2	777.7	2,212.8	5,863.7	13,186.9
1926	316.3	55.5	12.6	19.5	26.7	38.0	96.2	285.2	812.5	2,323.9	6,164.0	13,716.0
1927	310.3	53.9	11.9	19.8	26.6	37.6	95.6	282.1	791.5	2,272.1	5,892.0	12,759.5
1928	327.6	52.9	12.2	18.0	25.0	39.4	100.5	295.3	833.1	2,394.8	6,236.0	13,301.4
1929	330.7	50.0	12.1	18.1	25.4	38.2	98.5	299.0	851.6	2,395.3	6,079.1	12,595.7
1930	324.8	46.5	10.1	16.0	23.7	37.6	94.7	291.9	843.2	2,310.4	5,790.9	11,825.6
1931	322.8	36.1	10.1	15.1	22.1	36.0	94.8	291.9	832.2	2,235.1	5,641.5	11,599.2
1932	335.0	28.6	9.1	14.1	21.0	35.1	93.7	290.5	841.6	2,283.0	5,870.3	12,137.1
1933	335.1	27.4	9.5	13.2	19.9	34.6	92.4	294.8	848.9	2,254.1	5,801.6	11,741.9
1934	349.1	27.5	8.6	12.2	19.6	34.5	95.9	306.8	864.2	2,280.8	5,993.3	12,114.0
1935	355.1	43.5	9.0	14.1	19.2	33.9	94.8	305.0	859.2	2,273.6	5,992.2	12,127.3
1936	382.2	45.2	8.4	13.9	19.3	33.9	100.6	318.9	898.6	2,368.5	6,475.6	13,140.7
1937	379.4	34.5	8.0	12.0	18.5	33.4	99.2	317.9	885.8	2,306.1	6,264.6	12,536.1
1938	378.3	35.8	7.4	12.1	17.7	31.7	96.7	313.4	861.4	2,292.4	6,098.9	12,061.6
1939	389.3	29.7	6.8	11.5	17.2	31.0	96.5	317.1	873.2	2,303.3	6,152.8	12,652.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	293.1	---	---	---	---	---	---	---	---	---	---	---
1911	286.8	---	---	---	---	---	---	---	---	---	---	---
1912	293.0	---	---	---	---	---	---	---	---	---	---	---
1913	286.2	---	---	---	---	---	---	---	---	---	---	---
1914	289.2	119.8	18.4	27.0	31.1	46.3	113.0	288.0	860.6	2,431.8	5,465.2	9,667.4
1915	294.5	118.3	16.8	25.3	29.8	44.7	111.3	290.3	872.8	2,441.2	5,644.4	10,350.0
1916	302.1	103.3	17.6	24.8	30.4	48.3	114.8	310.0	897.6	2,480.4	5,782.9	10,495.0
1917	308.2	---	---	---	---	---	---	---	---	---	---	---
1918	304.3	105.4	17.3	24.3	39.3	58.8	112.8	289.3	870.2	2,374.4	5,481.8	10,017.9
1919	269.0	92.4	14.0	19.5	26.5	38.1	88.9	246.1	780.8	2,151.9	5,278.2	10,554.4
1920	281.7	105.1	16.3	22.3	28.6	36.5	86.3	255.2	800.3	2,269.1	5,830.6	11,599.0
1921	274.4	68.4	15.3	24.0	27.7	34.5	81.9	251.9	781.0	2,210.2	5,597.2	11,393.6
1922	292.0	75.9	14.8	20.4	26.6	34.6	87.5	271.9	833.8	2,381.0	5,922.0	12,047.2
1923	306.6	78.3	14.2	19.8	26.5	35.8	93.2	287.1	875.1	2,405.0	6,140.8	12,732.5
1924	310.9	68.2	13.2	21.6	27.7	35.9	97.5	299.5	862.8	2,425.3	6,171.8	12,492.1
1925	312.2	70.7	13.7	20.7	26.0	36.5	95.7	296.6	864.9	2,381.3	6,155.3	12,947.6
1926	331.4	64.6	12.1	18.6	25.8	37.2	100.7	314.8	897.5	2,533.5	6,446.8	13,744.5
1927	327.8	61.1	12.0	18.9	25.8	36.2	103.3	312.7	889.2	2,474.8	6,219.4	12,839.6
1928	347.1	59.7	12.5	17.2	24.6	38.2	109.3	330.0	937.5	2,613.7	6,570.7	13,338.9
1929	352.7	58.6	12.3	16.6	24.3	38.0	109.3	340.5	969.4	2,620.3	6,379.0	12,774.5
1930	348.0	54.1	10.4	15.2	22.6	37.3	105.3	334.5	964.2	2,531.8	6,081.3	12,122.4
1931	348.9	40.1	10.3	14.4	21.4	35.9	107.1	340.9	961.5	2,465.0	5,948.3	12,111.1
1932	360.9	29.6	9.3	13.7	20.9	34.7	106.7	340.8	979.8	2,503.4	6,175.0	12,392.0
1933	364.8	30.3	10.4	12.3	20.0	34.9	105.6	350.4	1,002.8	2,500.7	6,127.3	12,027.5
1934	382.5	32.6	8.3	11.8	19.9	35.7	111.7	371.3	1,026.2	2,546.7	6,358.1	12,430.1
1935	389.4	49.8	8.7	13.5	19.7	35.1	110.7	367.5	1,026.8	2,543.4	6,390.4	12,467.2
1936	420.1	54.4	8.8	13.7	20.0	35.1	118.5	389.2	1,088.1	2,652.3	6,882.6	13,558.3
1937	419.7	39.5	7.8	11.3	18.5	35.5	118.4	395.5	1,077.3	2,608.3	6,700.8	12,834.8
1938	418.2	39.3	8.4	11.7	18.0	34.2	117.5	390.9	1,052.3	2,592.9	6,474.6	12,345.5
1939	431.5	33.3	7.1	11.6	18.0	33.6	118.6	401.0	1,071.6	2,621.2	6,522.3	12,955.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	277.9	---	---	---	---	---	---	---	---	---	---	---
1911	271.8	---	---	---	---	---	---	---	---	---	---	---
1912	273.2	---	---	---	---	---	---	---	---	---	---	---
1913	269.8	---	---	---	---	---	---	---	---	---	---	---
1914	271.6	82.2	16.2	31.8	31.9	51.5	108.5	271.9	727.2	2,080.2	4,859.6	9,008.6
1915	279.5	88.6	15.7	29.7	32.7	49.8	109.3	274.0	733.0	2,137.8	5,049.6	9,519.3
1916	283.9	85.3	16.6	28.4	32.5	51.0	109.8	275.4	747.2	2,151.8	5,239.4	9,845.5
1917	285.5	---	---	---	---	---	---	---	---	---	---	---
1918	286.0	81.6	15.9	27.9	42.2	66.5	121.4	287.6	747.1	2,105.6	5,041.7	9,564.3
1919	266.0	70.3	13.6	22.2	27.8	43.4	94.9	250.7	689.4	2,003.8	5,064.1	9,917.6
1920	285.1	71.7	15.3	25.9	30.5	48.6	96.5	261.4	730.3	2,123.3	5,556.7	11,425.2
1921	273.1	57.9	14.9	26.6	29.9	40.0	87.1	251.2	689.0	2,039.0	5,347.9	11,274.1
1922	281.3	57.5	13.8	22.8	28.1	39.4	90.3	250.8	702.5	2,120.6	5,570.6	12,244.7
1923	290.2	58.8	13.2	22.8	27.8	39.2	92.2	260.6	716.1	2,118.3	5,714.1	12,883.3
1924	288.1	56.4	12.1	23.4	28.4	38.9	90.8	259.8	701.5	2,066.4	5,686.8	12,688.7
1925	286.6	49.0	13.3	22.7	26.8	39.2	88.9	248.0	683.8	2,039.4	5,595.8	13,373.0
1926	300.8	46.0	13.1	20.4	27.6	38.8	91.5	253.3	720.7	2,108.5	5,903.7	13,693.9
1927	292.2	46.5	11.8	20.7	27.5	39.0	87.4	249.2	686.0	2,063.6	5,589.3	12,697.4
1928	307.6	45.9	11.8	18.9	25.4	40.7	91.1	257.8	720.4	2,169.3	5,924.4	13,272.5
1929	308.0	41.0	12.0	19.7	26.4	38.4	87.1	254.1	725.1	2,163.5	5,798.4	12,459.8
1930	301.0	38.6	9.8	16.9	24.7	37.9	83.4	245.6	714.2	2,081.8	5,517.4	11,602.8
1931	296.1	32.0	10.0	15.7	22.8	36.2	81.9	239.0	694.6	1,998.7	5,353.1	11,214.9
1932	308.5	27.5	8.9	14.4	21.0	35.5	80.1	236.3	695.0	2,057.4	5,584.8	11,945.7
1933	304.6	24.4	8.6	14.2	19.7	34.3	78.6	235.0	685.6	2,002.1	5,495.6	11,526.5
1934	315.0	22.2	8.8	12.7	19.3	33.4	79.4	237.5	692.8	2,010.5	5,651.8	11,869.0
1935	319.9	37.0	9.3	14.6	18.7	32.8	78.3	238.0	682.1	2,000.8	5,621.0	11,871.0
1936	343.6	35.7	8.1	14.1	18.5	32.6	82.1	243.7	698.9	2,082.9	6,098.0	12,825.9
1937	338.2	29.3	8.3	12.7	18.4	31.4	79.4	235.2	684.4	2,003.7	5,861.8	12,310.8
1938	337.6	32.2	6.4	12.6	17.3	29.1	75.3	231.0	661.0	1,993.4	5,753.6	11,847.5
1939	346.4	26.1	6.5	11.4	16.4	28.5	73.8	227.8	665.2	1,989.0	5,815.2	12,423.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	343.5	---	---	---	---	---	---	---	---	---	---	---
1911	353.4	---	---	---	---	---	---	---	---	---	---	---
1912	365.9	---	---	---	---	---	---	---	---	---	---	---
1913	339.6	---	---	---	---	---	---	---	---	---	---	---
1914	355.9	139.2	28.3	28.1	53.9	129.8	322.3	739.8	1,415.2	2,623.1	4,369.9	7,363.6
1915	363.3	124.5	19.6	26.4	51.6	127.2	319.6	700.7	1,554.2	2,730.8	4,810.4	7,319.1
1916	309.3	88.0	17.0	19.5	46.5	138.1	304.9	648.6	1,418.2	2,510.5	4,377.5	7,094.3
1917	330.3	---	---	---	---	---	---	---	---	---	---	---
1918	317.7	88.1	20.5	20.9	60.9	142.6	325.6	646.9	1,385.2	2,534.3	4,215.1	6,710.6
1919	258.7	57.5	15.7	15.8	47.0	101.8	253.4	566.2	1,104.5	2,097.3	3,783.2	5,695.5
1920	273.2	68.9	14.8	17.8	46.1	114.7	255.5	582.5	1,154.7	2,292.5	4,230.4	6,866.5
1921	265.9	60.7	15.6	20.0	46.2	102.2	254.1	584.4	1,134.7	2,161.7	4,043.6	6,735.4
1922	257.1	52.6	14.6	17.6	40.9	100.9	253.0	635.1	1,138.1	2,173.3	3,969.2	6,960.0
1923	291.6	61.9	13.5	17.8	41.0	102.6	286.4	712.5	1,245.0	2,371.8	4,206.3	8,112.1
1924	319.9	67.3	14.4	19.7	48.3	114.0	323.8	784.2	1,336.7	2,557.7	4,529.0	8,648.0
1925	324.8	67.0	15.3	19.3	49.3	114.4	327.6	798.6	1,333.2	2,585.2	4,434.0	8,868.7
1926	338.6	62.6	16.3	18.7	48.2	120.9	344.6	859.5	1,392.0	2,637.5	4,593.7	8,357.9
1927	321.7	48.4	15.1	17.8	42.6	115.2	332.9	795.8	1,366.7	2,518.0	4,307.9	7,548.2
1928	340.7	43.2	12.8	17.8	44.3	127.6	361.6	852.7	1,484.7	2,613.7	4,394.5	7,589.4
1929	347.9	48.1	14.2	18.8	47.5	130.2	358.5	851.7	1,535.8	2,688.6	4,534.2	7,298.5
1930	354.8	53.8	13.8	17.0	46.8	129.9	354.5	875.6	1,636.8	2,709.9	4,241.4	7,122.9
1931	329.8	42.4	10.5	15.4	39.9	117.6	329.2	796.2	1,513.4	2,485.1	4,027.7	6,434.3
1932	328.5	30.7	11.0	15.4	36.5	112.8	317.8	778.0	1,530.4	2,383.9	4,096.9	6,444.9
1933	327.8	39.9	12.6	16.0	33.3	106.6	297.5	775.2	1,522.5	2,367.9	4,075.3	6,323.5
1934	350.1	35.4	12.1	15.0	35.9	109.8	323.5	824.6	1,561.5	2,504.9	4,309.8	6,779.1
1935	338.3	41.4	9.9	15.9	32.7	103.2	315.5	785.2	1,530.1	2,289.1	3,977.7	6,618.1
1936	369.4	44.8	11.6	16.7	35.6	106.5	335.5	823.1	1,638.2	2,523.1	4,515.0	6,892.5
1937	364.6	37.0	9.0	15.0	32.0	101.4	314.8	833.4	1,589.6	2,506.7	4,177.6	6,454.9
1938	366.5	38.1	10.3	16.5	34.4	97.0	311.4	838.5	1,571.3	2,404.5	4,159.0	6,308.8
1939	356.5	35.4	10.7	14.8	29.7	90.5	291.6	790.7	1,542.2	2,273.8	4,145.5	6,049.8

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	327.3	---	---	---	---	---	---	---	---	---	---	---
1911	347.9	---	---	---	---	---	---	---	---	---	---	---
1912	361.8	---	---	---	---	---	---	---	---	---	---	---
1913	333.8	---	---	---	---	---	---	---	---	---	---	---
1914	350.6	149.4	33.8	25.3	46.9	128.4	311.8	641.4	1,335.7	2,639.3	4,652.0	6,978.7
1915	353.6	159.4	*	23.4	42.6	127.0	301.7	606.1	1,428.1	2,640.2	5,036.3	8,006.5
1916	311.5	104.8	24.8	18.7	42.0	150.6	286.8	581.1	1,309.0	2,444.7	4,556.6	7,702.4
1917	331.2	---	---	---	---	---	---	---	---	---	---	---
1918	313.6	91.3	20.7	20.4	53.5	140.4	308.2	558.4	1,249.5	2,443.6	4,334.8	7,415.7
1919	248.5	69.0	17.6	14.3	42.2	92.1	228.8	491.9	953.8	1,928.8	3,752.7	6,059.7
1920	264.7	69.2	15.9	15.4	41.1	107.0	230.6	507.6	1,020.0	2,157.6	4,259.6	7,123.2
1921	250.6	64.9	15.7	18.8	37.6	89.4	230.8	500.8	932.7	1,987.9	4,100.5	6,788.7
1922	259.7	56.9	17.8	16.5	36.0	93.1	245.5	560.2	980.1	2,019.0	3,989.5	7,521.9
1923	288.2	78.1	13.7	16.4	35.6	99.8	267.7	635.7	1,092.2	2,253.8	4,360.2	9,044.4
1924	320.9	72.7	16.5	18.7	42.5	108.6	306.6	713.0	1,235.4	2,424.0	4,754.4	9,992.7
1925	327.8	70.7	14.8	19.4	40.6	107.3	313.3	765.5	1,224.2	2,505.7	4,545.0	10,255.7
1926	344.2	74.8	18.8	17.6	41.9	116.8	327.4	827.5	1,288.0	2,565.9	4,829.6	9,564.5
1927	325.6	54.8	14.5	16.8	37.0	105.4	318.2	748.6	1,285.2	2,438.8	4,526.6	8,856.0
1928	349.3	53.6	14.6	17.6	39.6	126.1	350.9	824.8	1,392.6	2,534.5	4,731.0	8,149.0
1929	351.6	54.9	15.9	16.6	44.6	129.4	358.4	824.9	1,449.0	2,717.6	4,904.9	8,264.9
1930	367.6	51.0	13.5	17.6	39.5	124.3	352.0	856.7	1,551.9	2,741.2	4,629.8	7,897.1
1931	338.6	43.6	12.3	14.3	35.0	111.8	331.7	752.4	1,433.1	2,447.2	4,457.2	7,368.7
1932	336.7	28.7	12.4	14.0	32.4	104.0	319.9	741.8	1,452.9	2,354.0	4,516.6	7,116.1
1933	337.1	38.9	12.6	15.4	35.5	104.8	301.4	750.7	1,441.3	2,330.8	4,336.6	7,163.2
1934	364.7	40.8	12.7	14.1	31.0	110.9	326.3	813.7	1,514.6	2,529.9	4,639.7	7,371.8
1935	354.0	48.9	8.5	13.8	31.5	102.3	319.6	780.3	1,495.0	2,323.2	4,338.2	7,221.6
1936	388.0	51.0	12.9	17.0	34.5	104.4	341.6	832.1	1,599.0	2,589.2	4,796.3	7,494.1
1937	384.2	33.6	8.8	13.4	26.3	101.0	319.2	842.8	1,547.3	2,612.2	4,613.6	7,203.9
1938	384.2	38.7	10.0	15.1	29.7	95.3	316.8	842.9	1,519.3	2,525.8	4,474.1	6,850.2
1939	372.9	45.1	10.2	14.9	25.7	93.2	289.4	789.7	1,516.9	2,347.8	4,429.8	6,638.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE CARDIOVASCULAR SYSTEM

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	....	....	....	....	....	....	....	....	....	....	....	....
1901	....	....	....	....	....	....	....	....	....	....	....	....
1902	....	....	....	....	....	....	....	....	....	....	....	....
1903	....	....	....	....	....	....	....	....	....	....	....	....
1904	....	....	....	....	....	....	....	....	....	....	....	....
1905	....	....	....	....	....	....	....	....	....	....	....	....
1906	....	....	....	....	....	....	....	....	....	....	....	....
1907	....	....	....	....	....	....	....	....	....	....	....	....
1908	....	....	....	....	....	....	....	....	....	....	....	....
1909	....	....	....	....	....	....	....	....	....	....	....	....
1910	362.0	....	....	....	....	....	....	....	....	....	....	....
1911	359.6	....	....	....	....	....	....	....	....	....	....	....
1912	370.4	....	....	....	....	....	....	....	....	....	....	....
1913	345.9	....	....	....	....	....	....	....	....	....	....	....
1914	361.5	129.1	23.0	30.9	60.8	131.3	335.2	871.5	1,515.6	2,604.6	4,105.2	7,632.7
1915	373.8	90.1	19.8	29.4	60.3	127.4	341.4	829.3	1,714.8	2,833.8	4,596.4	6,839.6
1916	306.9	71.5	*	20.3	50.6	125.7	325.2	737.2	1,557.9	2,587.6	4,203.5	6,656.6
1917	329.5	....	....	....	....	....	....	....	....	....	....	....
1918	322.0	85.0	20.4	21.4	67.5	144.8	344.7	766.1	1,560.0	2,639.6	4,097.9	6,207.6
1919	269.1	46.2	13.9	17.3	51.3	110.9	279.7	667.8	1,301.9	2,296.9	3,813.6	5,429.7
1920	281.8	68.6	13.7	20.1	50.6	121.7	282.8	684.0	1,330.8	2,452.9	4,200.7	6,674.3
1921	281.5	56.5	15.4	21.2	54.1	113.9	279.3	695.8	1,398.2	2,370.5	3,985.7	6,697.4
1922	274.7	48.4	11.4	18.6	45.3	107.9	281.4	732.1	1,345.7	2,361.8	3,948.7	6,573.7
1923	295.1	45.9	13.3	19.1	46.1	105.2	306.0	809.2	1,445.2	2,516.8	4,051.0	7,489.8
1924	319.0	62.0	12.3	20.7	53.7	118.9	341.8	871.7	1,469.1	2,721.6	4,302.8	7,760.9
1925	321.8	63.4	15.7	19.1	57.4	121.1	342.4	839.1	1,474.7	2,680.6	4,323.8	7,947.1
1926	332.9	50.7	13.7	19.7	54.0	124.7	362.3	897.7	1,527.1	2,723.3	4,360.4	7,543.7
1927	317.8	42.3	15.6	18.8	47.9	124.4	348.0	851.7	1,473.4	2,612.0	4,091.6	6,635.3
1928	332.1	33.1	10.9	18.1	48.7	128.9	372.3	885.4	1,606.4	2,706.3	4,054.0	7,187.9
1929	334.2	41.5	12.5	20.9	50.1	130.9	358.7	883.0	1,650.3	2,655.3	4,175.1	6,583.4
1930	342.0	56.5	14.2	16.5	53.4	135.1	356.8	897.5	1,747.2	2,674.5	3,872.1	6,540.2
1931	321.1	41.2	8.7	16.6	44.4	122.9	326.8	846.3	1,616.8	2,527.6	3,620.2	5,732.2
1932	320.4	32.7	9.7	16.8	40.1	120.9	315.7	819.1	1,629.2	2,417.4	3,699.6	5,941.7
1933	318.6	40.8	12.6	16.6	37.6	108.4	293.7	802.7	1,624.5	2,409.3	3,827.2	5,695.7
1934	335.8	30.1	11.4	16.0	40.4	108.8	320.8	836.6	1,619.8	2,477.1	3,996.9	6,337.1
1935	323.0	33.9	11.2	17.9	33.7	104.1	311.5	790.5	1,573.1	2,251.5	3,636.0	6,169.2
1936	351.2	38.8	10.3	16.3	36.6	108.4	329.7	813.2	1,685.6	2,450.5	4,248.5	6,446.4
1937	345.5	40.3	9.3	16.6	37.1	101.9	310.5	823.3	1,640.1	2,391.3	3,764.5	5,901.2
1938	349.3	37.6	10.5	17.9	38.8	98.5	306.2	833.6	1,632.5	2,272.4	3,860.4	5,909.9
1939	340.6	26.0	11.2	14.7	33.2	88.1	293.6	791.8	1,571.5	2,193.3	3,875.8	5,617.6

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Cause of death:

DISEASES OF CARDIOVASCULAR SYSTEM

NOTES:

For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes .....	724	1,112	1,194
Male.....	397	592	653
Female....	327	520	541

For 1901-09 deaths from rheumatic fever are included with total deaths, but are not distributed by age.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ICD Revision	Years in use	ICD Nos
Fifth	1939	90-95
Fourth	1930-1938	90-95
Third	1921-1929	87-90
Second	1910-1920	77-80
First	1900-1909	77-80

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	137.4	147.8	15.0	23.3	28.8	43.4	80.8	173.0	414.1	957.3	1,751.9	2,249.8
1901	140.0	122.1	13.7	20.3	27.4	44.8	82.0	175.4	437.8	966.6	1,868.6	2,228.6
1902	145.4	122.8	13.8	21.5	27.9	46.2	85.5	181.1	449.4	1,013.7	1,937.3	2,461.6
1903	151.8	103.7	10.5	19.1	27.2	45.2	88.2	185.0	472.6	1,080.3	2,141.2	2,705.7
1904	163.7	109.6	12.7	23.6	28.9	45.6	91.1	195.0	510.3	1,179.5	2,294.4	3,177.0
1905	161.9	119.1	10.9	23.4	28.4	46.1	92.6	195.0	501.1	1,139.9	2,280.1	3,184.1
1906	154.2	99.7	10.8	21.4	27.6	45.0	90.7	189.7	494.4	1,146.3	2,244.0	3,268.4
1907	166.6	90.3	12.2	21.6	29.1	45.7	94.5	209.3	530.3	1,236.1	2,511.4	3,682.3
1908	152.0	92.2	11.8	21.8	25.8	40.5	83.1	185.4	475.8	1,150.0	2,328.4	3,487.8
1909	153.0	88.9	10.7	20.9	25.0	39.7	83.2	181.8	470.4	1,155.1	2,353.9	3,517.0
1910	158.9	99.2	12.8	22.3	26.7	40.8	85.7	189.5	480.7	1,191.7	2,481.6	3,811.6
1911	156.4	82.5	10.7	21.0	25.6	40.7	85.4	183.5	477.0	1,202.6	2,445.8	3,777.9
1912	158.7	74.4	9.9	18.7	25.6	40.1	88.2	190.7	478.3	1,207.2	2,524.4	3,738.3
1913	154.6	73.7	11.1	20.0	24.3	40.6	86.7	184.3	458.9	1,160.7	2,454.3	3,792.9
1914	158.2	60.2	9.8	21.5	25.0	39.6	83.9	182.9	462.7	1,215.8	2,507.5	3,977.9
1915	163.9	60.5	9.6	19.8	25.5	39.5	84.8	184.7	479.2	1,258.7	2,617.7	4,344.2
1916	167.2	55.6	10.2	19.0	25.3	42.6	88.3	193.8	492.0	1,273.3	2,734.4	4,498.5
1917	169.9	56.0	9.6	18.2	26.8	44.1	89.6	195.4	498.9	1,279.5	2,784.5	4,542.1
1918	171.6	55.3	10.3	19.3	35.0	55.8	96.3	195.4	489.6	1,236.5	2,611.0	4,269.1
1919	147.9	49.8	8.4	14.9	22.9	35.1	74.1	165.0	425.4	1,113.9	2,506.7	4,323.5
1920	159.6	52.0	9.7	17.4	24.5	37.5	73.1	171.6	452.0	1,200.9	2,857.4	5,214.4
1921	156.2	43.6	9.9	18.6	24.3	33.4	69.8	174.1	440.6	1,166.7	2,762.9	5,172.8
1922	165.0	45.6	9.5	16.0	23.4	33.8	74.7	185.7	469.5	1,249.6	2,925.5	5,735.0
1923	174.0	46.5	9.5	16.0	23.4	34.3	78.7	198.3	493.2	1,273.0	3,069.3	6,103.6
1924	175.7	43.1	8.8	16.8	24.6	35.2	81.5	203.1	486.9	1,261.9	3,079.0	6,076.5
1925	184.8	42.8	9.3	16.5	23.7	36.5	83.8	209.1	508.8	1,319.8	3,213.0	6,745.7
1926	198.6	39.8	8.7	14.8	23.7	36.5	86.5	221.3	539.9	1,419.6	3,508.1	7,238.9
1927	195.3	37.9	8.5	14.9	23.6	36.5	86.0	218.1	530.9	1,390.2	3,347.0	6,825.0
1928	207.7	34.9	8.1	13.6	22.2	38.5	91.7	231.4	560.6	1,484.5	3,612.0	7,267.1
1929	211.2	35.1	8.1	13.6	22.6	38.4	89.3	232.5	576.6	1,495.6	3,563.5	7,105.9
1930	214.2	33.1	6.9	12.1	21.3	38.2	89.4	238.9	598.6	1,493.9	3,476.7	6,680.9
1931	213.4	24.0	6.6	11.3	19.7	36.1	88.0	236.8	592.3	1,453.5	3,436.7	6,554.6
1932	224.1	17.5	5.9	10.2	18.4	34.9	87.2	237.3	609.3	1,510.4	3,651.2	7,136.2
1933	228.0	17.4	6.1	9.9	17.3	34.1	86.2	244.7	628.3	1,523.0	3,677.7	6,926.9
1934	240.3	16.5	5.4	9.3	17.1	34.4	90.5	258.8	648.0	1,562.6	3,825.4	7,291.0
1935	245.4	24.6	5.7	10.1	16.5	33.1	89.5	254.8	646.9	1,570.6	3,882.6	7,399.2
1936	266.6	25.9	5.0	10.2	16.6	33.4	94.5	268.3	684.8	1,655.3	4,255.2	8,112.7
1937	268.9	19.4	4.9	8.8	15.8	32.4	92.6	272.0	685.5	1,643.2	4,190.0	7,894.0
1938	269.7	20.3	4.4	8.8	15.5	30.6	91.8	270.6	671.1	1,639.1	4,105.1	7,640.0
1939	275.5	18.2	4.4	8.3	14.9	29.4	89.2	268.2	680.5	1,638.2	4,132.9	7,945.7

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	141.0	160.1	16.3	19.4	27.0	40.8	73.3	174.0	440.6	1,063.3	1,950.3	2,504.9
1901	147.4	138.8	13.6	18.8	26.6	44.5	78.8	182.1	474.3	1,092.2	2,081.5	2,580.8
1902	153.7	138.3	12.8	19.4	26.9	41.7	82.0	187.6	514.0	1,141.8	2,179.4	2,799.6
1903	159.3	115.8	11.6	17.1	29.4	43.0	84.0	191.8	515.4	1,214.0	2,377.8	2,928.8
1904	170.4	128.5	12.9	20.2	28.8	41.4	87.1	198.2	565.0	1,295.4	2,606.2	3,424.0
1905	167.4	127.8	11.6	18.9	27.2	43.5	90.4	198.3	543.9	1,263.2	2,511.0	3,574.8
1906	160.8	105.9	9.9	18.5	27.0	43.7	91.8	194.5	539.9	1,257.3	2,497.8	3,464.3
1907	175.6	107.3	12.2	20.5	29.9	43.2	95.4	225.2	586.6	1,372.2	2,740.5	4,076.6
1908	156.5	97.9	10.8	18.8	24.7	37.1	80.9	192.3	520.5	1,232.9	2,580.2	3,713.0
1909	158.0	103.5	10.1	17.7	23.0	37.7	82.0	189.4	515.1	1,254.9	2,542.0	3,822.9
1910	164.1	112.3	12.5	19.6	25.6	39.2	86.2	195.1	515.9	1,292.0	2,711.0	4,182.2
1911	161.3	92.5	10.0	18.5	24.8	39.1	84.6	194.5	511.8	1,288.9	2,635.9	3,992.5
1912	164.9	84.1	9.8	16.1	24.7	37.8	87.6	204.3	524.6	1,314.2	2,695.2	3,858.5
1913	159.5	81.1	11.3	17.7	23.3	38.4	85.5	194.1	496.8	1,262.4	2,574.7	4,089.6
1914	163.7	69.7	10.4	19.0	24.1	37.5	84.8	189.1	502.1	1,317.5	2,664.8	4,095.2
1915	168.6	66.8	9.9	17.5	23.7	37.6	84.8	190.8	519.7	1,345.7	2,786.2	4,544.5
1916	173.5	60.9	10.8	17.3	23.9	41.9	89.4	207.0	538.5	1,360.7	2,880.5	4,636.1
1917	178.1	60.0	9.5	16.6	25.8	43.8	90.5	208.4	548.2	1,389.6	2,970.9	4,695.2
1918	177.8	61.1	10.0	17.3	33.1	50.9	92.2	198.8	526.1	1,329.0	2,746.2	4,405.9
1919	150.0	56.3	8.6	13.6	21.9	31.6	70.2	165.3	455.8	1,159.8	2,578.9	4,449.9
1920	160.1	61.3	9.8	15.8	23.2	32.2	69.2	171.6	475.0	1,247.8	2,957.5	5,234.7
1921	158.5	46.2	10.0	17.4	22.8	30.0	67.8	179.6	471.6	1,219.9	2,849.0	5,195.1
1922	170.7	51.2	9.7	14.7	22.2	31.4	74.0	198.2	512.9	1,329.7	3,044.0	5,784.3
1923	181.3	51.8	9.7	14.8	22.5	33.0	79.4	214.5	546.8	1,352.6	3,203.9	6,028.4
1924	185.4	45.8	9.1	15.8	23.5	34.1	84.9	222.9	542.1	1,365.7	3,245.6	6,137.0
1925	195.9	50.4	8.9	15.7	22.4	34.7	86.9	234.3	573.1	1,435.2	3,384.1	6,722.7
1926	210.5	45.8	8.5	14.0	22.6	35.6	89.8	248.3	604.6	1,549.2	3,659.8	7,324.4
1927	208.6	42.2	8.4	14.0	22.1	34.5	91.7	245.2	602.3	1,522.9	3,525.2	6,951.9
1928	222.6	38.4	8.7	12.7	21.4	37.3	97.5	261.8	637.8	1,624.0	3,802.2	7,342.5
1929	228.0	41.3	8.2	12.4	21.3	37.7	97.6	268.2	658.7	1,652.4	3,736.6	7,233.5
1930	234.6	37.7	7.2	11.4	19.5	37.3	99.2	280.6	697.1	1,659.3	3,679.9	6,977.1
1931	235.3	26.5	6.7	10.4	18.7	35.3	99.9	282.3	697.9	1,617.3	3,640.6	6,941.0
1932	246.7	17.2	6.1	9.8	17.8	33.7	100.0	284.9	722.2	1,679.8	3,847.2	7,339.0
1933	254.5	19.2	6.6	9.1	16.8	34.0	99.1	298.5	758.4	1,711.0	3,925.5	7,105.7
1934	270.6	18.7	5.1	8.8	16.6	35.4	105.7	322.0	789.8	1,773.4	4,095.9	7,509.9
1935	277.0	28.4	5.5	9.4	16.4	34.0	105.4	316.8	791.2	1,789.0	4,189.5	7,709.6
1936	302.1	30.6	5.4	9.7	17.0	34.3	112.2	338.0	850.1	1,887.1	4,601.3	8,440.3
1937	306.8	20.8	4.6	8.0	15.2	33.7	111.2	348.1	855.6	1,896.1	4,540.5	8,237.9
1938	306.7	21.6	5.0	8.3	15.3	32.6	111.9	347.6	835.7	1,889.1	4,393.1	7,865.5
1939	314.6	19.9	4.6	8.4	14.9	31.5	109.8	349.4	857.7	1,897.6	4,409.9	8,174.5

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	133.7	135.2	13.7	27.3	30.5	46.0	89.0	172.0	388.1	856.7	1,571.0	2,065.3
1901	132.6	105.1	13.8	21.7	28.2	45.2	85.4	168.4	402.1	847.6	1,675.4	1,972.9
1902	137.1	107.0	14.8	23.6	28.8	50.7	89.1	174.2	385.9	892.6	1,718.7	2,215.2
1903	144.3	91.3	9.5	21.1	25.2	47.4	92.7	177.9	430.6	953.9	1,928.7	2,542.5
1904	157.0	90.3	12.5	27.0	29.1	49.8	95.2	191.6	456.4	1,070.1	2,015.7	2,995.7
1905	156.5	110.1	10.2	28.0	29.5	48.8	95.0	191.4	458.7	1,023.5	2,074.9	2,896.3
1906	147.4	93.3	11.8	24.2	28.3	46.4	89.4	184.5	447.7	1,028.3	2,014.1	3,121.8
1907	157.2	72.9	12.2	22.8	28.4	48.4	93.5	191.7	472.1	1,103.4	2,304.9	3,385.8
1908	147.3	86.3	12.9	25.0	26.9	44.2	85.5	177.8	428.7	1,068.0	2,098.1	3,313.3
1909	147.7	73.8	11.3	24.1	27.1	42.0	84.5	173.4	423.1	1,056.1	2,182.1	3,278.2
1910	153.4	85.9	13.1	25.0	27.9	42.6	85.2	183.1	443.0	1,091.7	2,271.8	3,520.4
1911	151.1	72.3	11.4	23.5	26.5	42.4	86.3	171.0	439.4	1,115.7	2,270.6	3,608.0
1912	152.1	64.5	10.1	21.3	26.6	42.7	88.9	175.3	428.1	1,099.3	2,367.8	3,643.8
1913	149.3	66.1	10.9	22.3	25.4	42.9	88.0	173.3	417.8	1,058.1	2,344.3	3,561.9
1914	152.4	50.5	9.2	23.9	25.9	41.8	83.0	175.8	419.6	1,112.2	2,363.2	3,886.7
1915	158.9	54.0	9.3	22.2	27.3	41.5	84.8	177.7	435.1	1,170.1	2,463.9	4,189.9
1916	160.5	50.3	9.7	20.7	26.8	43.2	87.1	178.8	441.0	1,184.1	2,601.9	4,394.0
1917	161.3	51.9	9.7	19.9	27.8	44.5	88.6	180.7	444.7	1,166.6	2,615.5	4,426.4
1918	165.4	49.3	10.6	21.3	36.6	60.4	100.7	191.6	449.3	1,141.3	2,488.3	4,166.1
1919	145.8	43.3	8.2	16.2	23.9	38.5	78.3	164.6	391.8	1,066.3	2,441.0	4,229.0
1920	159.1	42.5	9.7	19.1	25.7	42.9	77.4	171.5	426.7	1,152.3	2,765.6	5,199.0
1921	153.8	40.9	9.7	19.9	25.8	36.8	72.1	167.9	406.8	1,111.6	2,683.8	5,155.9
1922	159.2	39.7	9.3	17.4	24.6	36.2	75.4	171.9	422.5	1,166.9	2,817.0	5,697.7
1923	166.5	41.0	9.3	17.3	24.2	35.6	77.9	180.4	435.1	1,190.7	2,945.5	6,161.1
1924	165.6	40.4	8.4	17.9	25.7	36.4	77.8	181.3	426.9	1,154.0	2,925.6	6,030.2
1925	173.4	35.0	9.7	17.3	25.0	38.3	80.4	181.6	438.6	1,200.1	3,055.3	6,763.3
1926	186.3	33.7	8.9	15.6	24.8	37.5	83.1	191.8	469.3	1,285.3	3,367.8	7,173.2
1927	181.5	33.6	8.6	15.8	25.0	38.5	79.9	188.7	452.8	1,252.6	3,181.6	6,727.4
1928	192.5	31.2	7.6	14.5	23.0	39.7	85.4	198.3	476.1	1,339.8	3,334.3	7,209.5
1929	193.9	28.8	8.0	14.9	23.8	39.1	80.6	193.6	487.2	1,333.2	3,401.1	7,009.2
1930	193.4	28.4	6.7	12.8	23.1	39.1	79.1	193.4	492.3	1,322.3	3,285.3	6,458.4
1931	190.9	21.5	6.4	12.2	20.7	36.8	75.5	187.4	478.8	1,284.2	3,245.1	6,264.6
1932	201.2	17.9	5.7	10.6	19.1	36.0	73.9	185.8	488.2	1,336.0	3,467.5	6,984.0
1933	201.1	15.5	5.7	10.8	17.8	34.1	72.8	186.5	488.7	1,329.8	3,444.7	6,792.2
1934	209.5	14.3	5.7	9.8	17.5	33.4	74.6	190.6	496.3	1,347.0	3,571.9	7,126.2
1935	213.3	20.7	5.9	10.8	16.6	32.2	73.1	188.2	492.8	1,348.5	3,596.2	7,165.5
1936	230.5	21.1	4.7	10.6	16.2	32.5	76.4	193.5	508.9	1,420.7	3,933.6	7,866.1
1937	230.5	17.9	5.2	9.6	16.3	31.2	73.6	190.8	505.1	1,388.7	3,865.9	7,635.2
1938	232.2	19.1	3.9	9.3	15.7	28.8	71.3	188.4	497.0	1,388.8	3,840.0	7,471.7
1939	236.1	16.5	4.3	8.3	14.8	27.3	68.3	181.7	493.5	1,380.0	3,879.2	7,773.6

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	157.6	---	---	---	---	---	---	---	---	---	---	---
1911	154.3	---	---	---	---	---	---	---	---	---	---	---
1912	156.4	---	---	---	---	---	---	---	---	---	---	---
1913	152.5	---	---	---	---	---	---	---	---	---	---	---
1914	155.7	59.7	9.6	21.4	24.3	36.7	76.9	170.7	451.1	1,210.8	2,517.3	4,013.7
1915	161.5	60.3	9.4	20.0	24.9	36.8	77.8	173.9	466.0	1,251.2	2,618.6	4,400.8
1916	166.1	56.3	10.2	19.4	24.5	38.6	80.0	182.6	477.0	1,269.2	2,758.1	4,583.9
1917	157.9	---	---	---	---	---	---	---	---	---	---	---
1918	169.8	56.8	10.1	19.6	34.0	51.5	86.2	180.9	472.4	1,228.4	2,632.7	4,356.7
1919	147.5	51.9	8.3	15.3	21.5	31.4	64.9	150.8	413.8	1,113.4	2,542.3	4,484.2
1920	159.5	53.6	9.9	17.9	23.4	33.3	64.1	157.9	441.2	1,197.9	2,891.9	5,381.9
1921	155.8	43.8	9.8	19.1	23.2	29.2	60.1	158.9	428.4	1,166.4	2,799.6	5,359.7
1922	165.7	47.6	9.6	16.4	22.5	29.3	63.9	168.4	457.8	1,254.8	2,984.0	5,995.2
1923	173.8	47.7	9.4	16.4	22.4	29.8	66.7	176.9	478.9	1,272.9	3,124.2	6,325.3
1924	174.0	43.8	8.9	17.0	23.1	30.0	68.6	179.1	467.4	1,254.9	3,124.3	6,264.6
1925	182.9	42.7	9.2	16.7	21.9	31.1	69.6	183.8	490.2	1,309.6	3,260.5	6,967.0
1926	197.3	39.8	8.4	15.0	22.1	30.7	72.5	194.5	520.2	1,413.2	3,563.3	7,514.7
1927	194.5	38.3	8.1	15.2	22.3	30.7	71.9	192.7	510.4	1,387.0	3,401.6	7,115.6
1928	207.1	35.8	8.1	13.6	20.7	31.9	75.3	202.1	536.2	1,482.2	3,684.1	7,612.1
1929	210.3	36.1	7.9	13.7	20.8	31.3	73.0	203.4	551.2	1,489.1	3,623.3	7,443.7
1930	213.1	32.7	6.6	12.0	19.4	30.9	73.5	208.1	570.0	1,486.9	3,537.8	6,981.8
1931	213.8	24.1	6.3	11.3	18.3	29.8	73.5	210.9	568.3	1,451.0	3,505.3	6,862.6
1932	226.4	17.4	5.8	10.1	17.2	28.8	73.1	213.5	585.6	1,515.8	3,726.5	7,508.3
1933	230.6	16.8	5.7	9.7	16.1	28.4	72.8	219.4	604.2	1,526.8	3,755.3	7,308.6
1934	242.6	16.1	5.1	9.0	15.8	28.3	75.8	231.6	623.7	1,563.4	3,895.3	7,681.7
1935	248.7	25.1	5.5	9.9	15.4	27.9	75.4	230.6	624.3	1,581.1	3,971.9	7,796.6
1936	269.6	26.2	4.7	9.9	15.2	27.8	79.6	242.6	658.6	1,660.1	4,342.2	8,545.6
1937	272.2	19.0	4.7	8.5	14.7	27.3	78.6	246.0	660.3	1,646.8	4,280.1	8,323.0
1938	273.1	20.7	4.2	8.5	14.1	25.7	77.8	244.7	645.9	1,646.2	4,194.4	8,057.7
1939	281.0	17.9	4.0	8.1	13.8	25.2	76.6	246.5	659.6	1,656.4	4,225.9	8,435.0

--- Data not available.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	162.9	---	---	---	---	---	---	---	---	---	---	---
1911	159.4	---	---	---	---	---	---	---	---	---	---	---
1912	162.7	---	---	---	---	---	---	---	---	---	---	---
1913	157.6	---	---	---	---	---	---	---	---	---	---	---
1914	161.2	69.1	10.2	19.0	23.7	34.5	77.5	178.6	492.2	1,311.1	2,668.6	4,142.6
1915	166.4	66.6	9.7	17.7	23.4	34.8	78.2	181.7	507.9	1,338.9	2,783.5	4,578.8
1916	172.4	60.9	10.5	17.6	23.4	37.5	82.0	197.6	525.9	1,359.5	2,901.7	4,708.1
1917	176.2	---	---	---	---	---	---	---	---	---	---	---
1918	176.3	63.0	9.9	17.5	32.5	46.5	82.2	186.2	512.3	1,324.1	2,767.4	4,473.8
1919	149.9	58.3	8.4	14.0	20.9	28.7	62.2	152.8	447.7	1,166.1	2,614.7	4,597.1
1920	160.2	63.7	9.9	16.2	22.6	28.3	61.0	160.3	468.1	1,251.2	2,992.3	5,358.8
1921	158.7	46.6	9.9	17.8	22.4	26.6	58.9	166.7	466.5	1,225.5	2,885.3	5,365.7
1922	171.9	53.9	9.8	15.1	21.7	27.5	64.4	182.9	507.9	1,342.4	3,106.6	5,985.3
1923	181.5	52.3	9.6	15.1	21.9	28.8	68.2	195.2	539.7	1,359.1	3,265.4	6,163.8
1924	184.0	47.1	9.2	15.9	22.6	29.2	72.6	201.6	527.1	1,366.4	3,285.1	6,218.5
1925	194.1	51.3	8.8	15.9	21.4	29.8	73.5	209.2	559.0	1,429.3	3,433.2	6,817.3
1926	209.3	46.0	8.0	14.2	21.4	30.1	77.2	222.2	589.7	1,547.4	3,711.5	7,505.6
1927	208.3	42.6	8.1	14.3	21.4	29.5	78.8	222.1	586.3	1,525.5	3,578.8	7,147.7
1928	222.4	38.9	8.5	12.6	20.4	30.8	82.8	234.4	619.3	1,629.7	3,871.6	7,635.0
1929	227.5	42.6	8.0	12.5	20.0	30.9	82.3	241.1	639.1	1,650.4	3,789.0	7,507.0
1930	234.1	37.8	6.9	11.2	18.4	30.7	84.4	252.1	675.2	1,656.9	3,732.9	7,244.6
1931	236.8	26.5	6.3	10.5	17.7	29.6	86.5	260.9	680.5	1,622.5	3,701.0	7,215.2
1932	250.3	17.1	5.9	9.7	17.1	28.5	86.6	265.7	706.8	1,695.0	3,915.2	7,691.6
1933	258.7	18.8	6.2	8.9	16.1	28.8	86.7	277.5	743.9	1,726.2	4,006.4	7,457.2
1934	274.4	18.4	4.7	8.6	15.9	29.5	92.3	298.6	773.2	1,784.8	4,164.8	7,874.2
1935	281.7	28.5	5.4	9.3	15.5	29.0	92.3	296.4	777.5	1,810.1	4,279.7	8,075.2
1936	306.7	31.3	5.0	9.5	15.9	29.1	97.9	315.8	833.7	1,903.5	4,692.5	8,861.1
1937	311.7	20.7	4.4	7.9	14.7	29.1	98.7	325.9	840.7	1,908.6	4,628.8	8,624.2
1938	312.0	22.0	4.7	8.1	14.5	28.2	99.3	325.6	822.3	1,907.4	4,481.5	8,268.3
1939	322.2	19.3	4.3	8.1	14.3	27.4	99.3	333.5	846.7	1,929.4	4,508.3	8,633.9

--- Data not available.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	152.1	---	---	---	---	---	---	---	---	---	---	---
1911	149.0	---	---	---	---	---	---	---	---	---	---	---
1912	149.7	---	---	---	---	---	---	---	---	---	---	---
1913	147.2	---	---	---	---	---	---	---	---	---	---	---
1914	150.0	50.1	9.0	23.9	24.9	39.1	76.3	161.7	406.4	1,109.1	2,378.7	3,912.9
1915	156.4	53.9	9.1	22.2	26.5	39.0	77.4	165.3	420.5	1,162.3	2,468.3	4,262.9
1916	159.5	51.5	10.0	21.2	25.6	39.8	77.8	165.8	423.7	1,177.5	2,628.3	4,489.0
1917	159.2	---	---	---	---	---	---	---	---	---	---	---
1918	163.3	50.3	10.2	21.7	35.3	56.1	90.5	174.9	428.9	1,130.5	2,510.8	4,268.2
1919	145.1	45.4	8.3	16.6	22.2	34.2	67.8	148.6	376.8	1,059.3	2,476.9	4,399.5
1920	158.7	43.2	9.9	19.6	24.1	38.6	67.5	155.3	412.0	1,143.1	2,800.4	5,399.3
1921	152.8	40.9	9.6	20.4	24.1	31.9	61.4	150.2	387.2	1,105.8	2,721.5	5,355.1
1922	159.3	41.2	9.4	17.7	23.4	31.2	63.5	152.5	404.3	1,165.4	2,872.4	6,002.7
1923	165.8	43.0	9.1	17.6	22.9	30.9	65.2	157.0	413.7	1,184.6	2,995.1	6,450.7
1924	163.7	40.5	8.6	18.1	23.5	30.9	64.2	154.6	403.3	1,140.3	2,977.0	6,300.4
1925	171.3	33.7	9.5	17.5	22.3	32.4	65.5	156.4	416.1	1,186.5	3,101.8	7,083.4
1926	184.9	33.4	8.8	15.8	22.7	31.3	67.5	164.7	445.3	1,275.3	3,426.8	7,521.8
1927	180.2	33.8	8.1	16.1	23.3	32.0	64.4	161.1	428.4	1,244.6	3,237.8	7,090.6
1928	191.4	32.6	7.7	14.7	21.0	33.0	67.3	167.2	446.4	1,330.3	3,509.7	7,594.5
1929	192.7	29.3	7.8	14.9	21.6	31.6	63.2	162.6	456.8	1,322.9	3,468.2	7,395.6
1930	191.5	27.4	6.3	12.9	20.4	31.2	61.9	160.4	458.0	1,311.4	3,354.2	6,784.5
1931	190.2	21.6	6.3	12.1	18.8	29.9	59.9	156.8	449.0	1,274.7	3,321.5	6,598.0
1932	202.0	17.6	5.7	10.5	17.3	29.1	59.0	157.1	457.0	1,334.4	3,549.6	7,370.8
1933	201.8	14.8	5.2	10.6	16.1	28.0	58.3	156.7	456.1	1,322.9	3,519.4	7,196.6
1934	210.1	13.8	5.4	9.5	15.7	27.1	58.7	159.5	465.5	1,338.3	3,643.0	7,536.5
1935	214.9	21.5	5.7	10.5	15.3	26.7	57.9	160.1	462.4	1,349.5	3,685.1	7,586.6
1936	231.7	20.9	4.4	10.3	14.5	26.5	60.5	164.4	474.2	1,415.2	4,017.1	8,307.8
1937	231.9	17.2	5.0	9.3	14.6	25.6	57.9	160.8	470.5	1,384.9	3,958.1	8,095.9
1938	233.5	19.4	3.7	8.9	13.7	23.3	55.8	158.6	460.7	1,386.5	3,930.6	7,898.9
1939	239.2	16.5	3.7	8.0	13.3	22.9	53.5	154.1	463.4	1,386.5	3,967.9	8,284.9

--- Data not available.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	204.8	---	---	---	---	---	---	---	---	---	---	---
1911	217.6	---	---	---	---	---	---	---	---	---	---	---
1912	227.2	---	---	---	---	---	---	---	---	---	---	---
1913	199.8	---	---	---	---	---	---	---	---	---	---	---
1914	213.1	72.4	14.7	21.9	39.9	96.9	225.9	471.4	801.7	1,373.0	2,168.8	3,259.6
1915	217.4	64.2	14.0	17.3	38.1	92.7	223.9	435.8	870.1	1,495.3	2,586.4	3,189.5
1916	182.8	47.0	10.0	13.4	35.9	100.6	210.9	383.7	810.9	1,364.6	2,161.7	3,300.4
1917	197.1	---	---	---	---	---	---	---	---	---	---	---
1918	194.5	36.7	13.2	15.4	45.4	110.5	226.3	407.0	820.8	1,394.3	2,160.7	3,235.1
1919	152.1	28.4	9.2	10.9	35.5	75.1	174.3	337.1	617.2	1,120.7	1,881.0	2,750.6
1920	160.7	34.1	8.1	12.4	34.9	83.8	172.2	343.5	632.1	1,252.3	2,234.5	3,489.0
1921	160.7	40.9	10.8	14.2	34.6	79.3	177.9	365.1	643.6	1,170.7	2,102.3	3,199.3
1922	158.5	25.5	8.7	13.3	31.0	76.8	183.5	381.3	638.5	1,167.6	2,002.3	3,304.2
1923	176.2	34.4	10.9	13.3	31.6	78.4	204.0	448.4	705.4	1,275.4	2,175.3	3,942.0
1924	192.5	36.0	7.5	15.3	37.9	85.8	220.0	487.0	775.9	1,380.9	2,338.3	4,231.6
1925	204.7	44.7	10.6	14.7	39.7	89.4	238.9	513.1	788.6	1,503.5	2,420.5	4,525.1
1926	212.2	40.2	11.2	13.0	38.0	92.4	240.5	542.2	836.3	1,536.8	2,580.8	4,465.9
1927	203.1	34.7	11.9	12.5	33.9	89.9	235.9	507.1	828.8	1,447.0	2,444.6	3,978.0
1928	213.1	26.8	8.5	13.3	33.7	95.3	252.2	533.4	886.4	1,523.5	2,509.1	4,201.8
1929	218.7	26.9	9.5	13.4	36.3	99.3	248.0	529.7	921.4	1,606.9	2,620.0	4,068.5
1930	224.7	36.7	9.8	12.2	36.3	100.6	243.3	553.7	996.0	1,614.5	2,476.4	3,945.7
1931	209.2	23.4	8.3	11.3	31.2	89.2	226.9	503.8	926.2	1,495.9	2,303.2	3,736.5
1932	203.4	18.7	7.0	11.2	28.6	86.6	222.1	484.8	939.3	1,420.3	2,397.3	3,713.0
1933	205.3	21.4	9.2	11.4	35.5	81.1	210.7	503.1	952.3	1,464.3	2,414.4	3,542.4
1934	220.1	19.4	7.5	11.1	27.0	85.0	225.2	538.2	975.3	1,550.3	2,677.5	3,816.1
1935	217.0	21.3	7.1	11.5	24.9	76.8	218.0	505.9	951.1	1,417.5	2,405.1	3,855.6
1936	240.3	23.9	7.2	12.2	27.8	79.8	230.6	537.2	1,037.2	1,587.8	2,805.7	4,248.5
1937	239.9	22.1	5.8	10.7	24.6	75.5	218.8	547.6	1,024.9	1,594.6	2,679.3	4,064.0
1938	240.1	17.4	6.0	11.3	26.6	72.3	216.0	547.9	1,010.4	1,545.1	2,598.9	3,912.9
1939	227.3	20.6	7.1	10.3	23.2	65.4	199.6	502.8	962.5	1,407.3	2,354.2	3,586.6

--- Data not available.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	205.0	...	...	...	...	...	...	...	...	...	...	...
1911	218.0	...	...	...	...	...	...	...	...	...	...	...
1912	228.9	...	...	...	...	...	...	...	...	...	...	...
1913	202.9	...	...	...	...	...	...	...	...	...	...	...
1914	217.8	84.3	*	19.4	32.1	96.4	225.9	417.9	773.4	1,509.8	2,535.7	3,078.8
1915	219.1	*	*	13.3	31.9	94.1	210.6	386.9	843.7	1,551.2	2,876.6	3,799.0
1916	189.4	60.5	15.0	13.2	30.9	109.8	198.4	355.9	787.1	1,386.0	2,388.2	3,589.2
1917	204.4	...	...	...	...	...	...	...	...	...	...	...
1918	197.5	36.2	11.9	14.3	39.3	106.1	220.5	368.0	773.9	1,418.5	2,323.3	3,577.4
1919	150.3	35.1	10.4	9.5	30.9	65.0	158.2	305.0	579.3	1,061.3	1,981.3	2,985.3
1920	158.8	33.9	8.3	11.2	29.0	78.2	160.3	302.5	582.1	1,192.5	2,363.3	3,952.2
1921	156.3	41.2	10.7	13.3	27.6	69.8	166.9	328.9	548.2	1,128.4	2,233.5	3,321.5
1922	159.2	24.7	8.8	11.7	26.6	70.3	173.2	357.6	578.1	1,146.9	2,108.8	3,782.5
1923	179.7	46.7	11.0	12.5	27.8	76.5	199.3	425.7	642.4	1,253.8	2,251.3	4,588.0
1924	199.7	32.1	8.5	14.6	32.1	83.1	219.2	460.5	745.0	1,354.4	2,628.7	5,257.8
1925	214.9	41.8	9.9	13.9	31.4	84.4	237.4	519.6	769.3	1,534.1	2,594.4	5,681.8
1926	223.1	44.2	12.5	12.5	33.1	90.0	230.9	547.7	811.3	1,581.2	2,822.9	5,345.3
1927	212.3	37.9	10.9	12.2	28.5	82.5	231.6	496.5	815.5	1,477.7	2,669.6	4,911.8
1928	224.2	34.2	9.9	13.6	29.5	96.0	247.3	534.0	862.5	1,533.9	2,771.6	4,635.0
1929	233.2	29.4	9.5	12.0	32.1	98.1	252.9	535.8	900.6	1,683.8	2,925.1	4,736.3
1930	239.3	36.9	9.8	12.8	29.0	95.8	247.6	563.3	975.4	1,699.6	2,816.8	4,549.5
1931	221.8	25.8	9.8	10.0	27.3	85.3	234.7	496.9	917.8	1,535.0	2,647.6	4,433.4
1932	213.1	17.9	7.9	10.7	24.1	79.7	233.3	479.4	918.5	1,440.1	2,718.6	4,094.3
1933	216.1	22.0	9.2	10.7	22.1	79.1	219.1	508.0	939.1	1,484.9	2,614.9	3,974.4
1934	236.5	21.3	7.8	9.9	22.3	86.4	234.5	557.9	997.0	1,609.4	2,972.8	4,248.9
1935	234.9	27.8	6.1	9.8	23.9	77.0	229.7	525.2	963.3	1,494.7	2,712.1	4,425.0
1936	260.6	25.1	8.0	11.6	26.0	78.9	246.5	569.1	1,057.6	1,664.9	3,097.4	4,648.4
1937	262.0	21.5	5.5	9.3	19.2	73.9	228.4	581.1	1,044.1	1,732.5	3,079.3	4,749.8
1938	259.5	18.5	6.6	10.1	22.1	71.5	228.0	582.6	1,006.4	1,658.3	2,925.9	4,203.7
1939	245.8	25.0	6.2	10.4	19.6	67.4	204.9	521.6	999.1	1,512.0	2,771.6	4,017.9

... Data not available.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

DISEASES OF THE HEART

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	204.5	...	...	...	...	...	...	...	...	...	...	...
1911	217.2	...	...	...	...	...	...	...	...	...	...	...
1912	225.3	...	...	...	...	...	...	...	...	...	...	...
1913	196.4	...	...	...	...	...	...	...	...	...	...	...
1914	208.1	*	*	24.3	47.5	97.5	225.9	542.9	837.4	1,217.5	1,824.5	3,385.9
1915	215.4	*	*	21.2	44.1	91.3	240.2	502.3	903.5	1,431.7	2,311.4	2,764.3
1916	175.9	*	*	13.6	40.6	91.4	225.0	420.3	841.3	1,339.5	1,941.5	3,092.6
1917	189.5	...	...	...	...	...	...	...	...	...	...	...
1918	191.5	37.1	14.5	16.4	50.8	114.6	232.8	459.5	881.2	1,366.1	2,001.7	2,990.8
1919	153.9	*	8.1	12.4	39.5	84.3	191.5	381.0	666.8	1,191.3	1,781.1	2,579.4
1920	162.7	34.3	7.9	13.6	40.2	89.0	185.2	398.9	697.4	1,323.4	2,104.0	3,142.2
1921	165.1	40.6	11.0	15.0	41.0	88.1	189.8	413.3	767.9	1,221.6	1,973.5	3,112.4
1922	157.7	26.3	8.6	14.9	34.9	82.6	194.3	411.9	717.9	1,193.0	1,894.2	2,975.5
1923	172.8	22.4	10.8	14.1	35.2	80.0	208.9	477.1	788.1	1,302.0	2,098.7	3,510.8
1924	185.2	40.0	6.4	16.1	43.3	88.3	220.8	519.5	816.2	1,413.3	2,046.8	3,554.7
1925	194.5	47.6	11.3	15.5	47.5	94.1	240.5	505.2	813.7	1,466.8	2,248.0	3,756.5
1926	201.1	36.2	9.8	13.5	42.6	94.7	250.3	535.7	868.8	1,483.7	2,341.4	3,872.6
1927	193.9	31.7	12.8	12.9	38.9	96.8	240.4	519.7	846.3	1,410.5	2,222.2	3,326.2
1928	202.1	19.7	7.0	13.0	37.7	94.7	257.1	532.7	917.9	1,511.3	2,251.2	3,891.0
1929	204.4	24.5	9.5	14.8	40.1	100.5	243.2	522.7	948.9	1,518.6	2,324.6	3,574.4
1930	210.2	36.5	9.7	11.6	42.8	105.0	239.1	542.6	1,022.9	1,518.2	2,152.8	3,491.3
1931	196.9	21.0	6.8	12.7	34.7	92.9	219.3	511.6	937.1	1,451.8	1,976.6	3,212.9
1932	193.7	19.5	6.1	11.6	32.7	93.1	211.2	491.0	965.8	1,398.2	2,093.1	3,427.1
1933	194.6	20.8	9.3	12.2	30.5	83.0	202.6	497.5	968.9	1,441.3	2,224.0	3,219.3
1934	204.1	17.5	7.3	12.3	31.2	83.7	216.3	516.3	948.3	1,484.8	2,397.4	3,493.3
1935	199.4	*	8.0	13.2	25.9	76.6	206.7	484.6	936.1	1,332.3	2,114.0	3,432.2
1936	220.4	22.8	6.4	12.9	29.4	80.7	215.3	502.3	1,012.5	1,503.2	2,529.3	3,952.0
1937	218.4	22.6	6.2	12.0	29.6	76.9	209.7	511.3	1,002.0	1,443.8	2,300.3	3,557.1
1938	221.2	16.4	5.5	12.5	30.8	73.0	204.6	510.4	1,015.1	1,421.8	2,288.8	3,698.6
1939	209.4	16.3	8.0	10.2	26.4	63.5	194.7	482.7	920.0	1,293.5	2,348.0	3,269.8

... Data not available.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

NOTES:  
For 1932-34, deaths of Mexicans are included with total deaths for the white population; but by age, are included with deaths for the ALL OTHER population.  
Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes ....	536	785	825
Male.....	301	443	478
Female.....	235	342	347

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ICD Revision	Years in use	ICD Nos.
Fifth	1939	83
Fourth	1930-1938	82
Third	1921-1929	74, 75, 83
Second	1910-1920	64-66, 82
First	1900-1909	64-66, 82

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	106.9	117.2	19.2	4.4	8.0	15.4	40.1	117.7	319.6	828.2	1,913.0	3,024.6
1901	106.9	118.7	14.1	4.0	6.2	14.9	40.5	121.7	326.4	822.5	1,933.4	2,918.2
1902	103.9	96.7	12.1	3.1	5.2	14.7	39.2	117.8	321.2	828.9	1,823.3	2,957.6
1903	105.2	84.0	11.3	3.0	6.2	13.7	36.9	116.7	327.1	826.7	1,938.7	3,088.9
1904	108.6	80.6	10.7	3.0	5.5	13.3	35.6	115.4	347.4	859.6	2,018.7	3,323.8
1905	105.9	77.3	8.7	2.6	5.4	11.7	37.2	118.6	331.5	851.2	1,942.6	3,144.5
1906	98.6	70.2	7.9	2.3	4.9	12.2	33.9	108.3	318.2	823.2	1,943.9	3,379.4
1907	104.5	59.8	6.5	2.2	4.8	12.7	35.7	110.6	340.5	885.9	2,070.9	3,617.4
1908	95.6	53.6	5.8	1.8	4.4	11.6	30.9	101.1	319.0	814.5	1,895.1	3,238.5
1909	95.5	58.0	4.8	2.0	3.4	9.8	29.8	99.0	307.0	813.1	1,919.5	3,200.3
1910	95.8	53.2	4.6	1.7	3.9	10.2	30.5	99.3	306.1	822.1	1,943.4	3,148.8
1911	91.8	42.9	4.3	1.4	3.6	9.8	29.9	97.9	290.9	798.0	1,846.6	3,123.6
1912	91.9	33.8	3.8	1.3	3.2	9.2	29.1	98.1	300.6	796.2	1,812.0	3,101.7
1913	91.1	23.8	2.9	1.5	3.1	8.7	27.3	99.4	289.9	796.8	1,856.6	2,937.3
1914	93.6	20.7	3.0	1.2	3.3	8.6	27.7	98.2	296.8	834.4	1,839.8	3,082.0
1915	94.5	17.9	2.6	1.3	2.6	7.3	28.1	97.3	296.2	827.2	1,906.5	3,153.8
1916	94.7	14.5	2.5	1.2	3.1	8.6	28.5	101.6	302.5	830.4	1,903.5	3,140.4
1917	95.9	15.7	2.4	1.3	3.2	8.1	29.9	101.1	304.2	837.6	1,925.4	3,190.2
1918	94.0	16.1	2.8	1.6	3.8	9.0	27.9	98.5	295.9	808.6	1,823.1	3,022.3
1919	89.9	13.5	2.4	1.3	3.1	7.4	24.9	91.9	283.7	772.3	1,839.1	3,160.1
1920	93.0	13.2	2.3	1.2	3.0	7.5	25.2	94.4	286.9	809.1	1,964.0	3,429.0
1921	89.2	9.7	1.8	0.9	2.4	6.0	23.3	89.0	274.4	778.7	1,895.6	3,382.1
1922	92.1	9.1	1.8	1.0	2.1	6.3	24.2	93.3	277.3	807.5	1,966.9	3,554.7
1923	95.7	11.4	1.7	0.9	2.2	6.3	25.5	96.5	287.2	813.7	2,004.8	3,752.0
1924	97.2	10.9	1.7	0.8	2.1	6.1	26.1	101.3	285.7	818.0	2,025.6	3,690.6
1925	89.5	9.8	1.7	1.0	2.4	5.8	22.6	89.9	258.2	736.3	1,866.8	3,594.6
1926	91.3	9.8	1.6	0.9	2.3	6.2	24.5	91.8	260.6	740.1	1,867.9	3,543.6
1927	88.1	8.7	1.4	0.9	2.0	5.9	23.7	88.2	248.4	715.8	1,781.0	3,245.7
1928	92.0	9.0	1.8	1.0	2.5	6.9	25.9	94.3	261.4	735.4	1,830.5	3,310.3
1929	90.8	8.1	1.8	1.0	2.4	6.2	25.8	94.1	259.5	719.3	1,756.2	2,964.7
1930	89.0	8.3	1.4	0.8	2.4	6.4	25.1	92.4	256.6	687.4	1,651.0	2,837.8
1931	86.8	8.3	1.3	0.8	2.0	6.2	24.6	88.5	249.5	656.8	1,563.1	2,734.3
1932	87.5	6.6	1.4	0.9	2.0	6.0	23.6	85.5	244.0	648.3	1,585.4	2,738.5
1933	94.1	7.3	1.7	0.8	1.9	6.0	22.4	82.9	233.7	610.5	1,508.5	2,609.0
1934	85.5	7.6	1.6	0.7	2.2	5.9	23.8	83.5	229.8	605.4	1,530.0	2,537.0
1935	85.7	13.1	1.6	1.1	2.1	6.0	23.1	83.0	228.3	587.8	1,487.6	2,533.8
1936	91.0	14.2	1.8	1.3	2.3	6.0	25.3	84.9	234.2	607.0	1,575.6	2,673.6
1937	86.7	11.2	1.8	0.9	2.3	6.1	23.9	81.0	218.5	567.7	1,466.2	2,450.7
1938	85.9	12.1	1.7	1.1	2.0	5.9	23.0	79.0	211.6	557.8	1,414.9	2,373.2
1939	87.8	8.8	1.3	0.7	1.8	5.8	22.9	78.7	211.5	559.0	1,428.6	2,456.0

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	106.2	142.0	21.5	4.3	7.6	15.0	40.8	108.7	328.1	855.6	1,980.5	2,963.4
1901	107.1	141.9	16.3	4.0	6.5	14.9	42.4	117.2	341.6	848.3	1,975.2	2,960.9
1902	102.7	115.5	11.7	2.8	5.3	14.5	39.3	112.8	324.1	866.9	1,864.3	2,861.0
1903	104.8	99.8	11.6	3.3	5.5	14.3	36.4	113.7	334.6	864.7	2,007.5	3,163.8
1904	107.1	89.7	9.8	3.1	5.3	13.4	36.8	113.7	363.2	884.8	2,028.9	3,264.0
1905	103.5	90.7	8.8	2.2	5.3	11.9	37.0	116.3	335.3	872.5	1,944.9	3,144.8
1906	97.2	85.3	8.1	2.5	5.2	12.2	35.6	102.9	327.3	838.1	2,000.5	3,263.5
1907	103.1	69.8	6.2	1.9	5.0	12.8	36.8	106.6	348.2	924.0	2,102.4	3,587.3
1908	94.0	58.8	5.3	2.1	4.5	11.0	31.4	98.7	321.9	844.6	1,918.9	3,254.2
1909	94.2	69.4	5.3	2.1	3.6	9.4	28.8	95.9	314.1	832.2	1,979.7	3,201.3
1910	94.1	61.2	5.0	1.9	3.7	10.6	30.6	95.3	315.9	849.2	1,944.3	3,128.9
1911	90.4	49.5	4.4	1.6	3.7	9.8	30.2	94.1	298.7	819.0	1,852.9	2,978.7
1912	91.2	39.7	4.0	1.1	3.1	9.1	28.8	97.0	309.6	825.5	1,827.8	3,121.7
1913	89.9	26.7	2.8	1.6	3.1	9.1	27.4	95.8	294.3	827.4	1,874.6	2,812.9
1914	92.6	23.1	3.2	1.4	3.3	8.5	27.5	93.9	304.9	860.9	1,870.2	3,009.1
1915	93.1	21.2	2.6	1.5	2.5	6.9	27.2	92.9	308.6	847.0	1,902.2	3,174.3
1916	93.4	14.9	2.6	1.1	3.1	8.6	27.7	98.3	309.8	859.5	1,887.2	3,132.3
1917	94.3	14.3	2.5	1.4	3.3	7.9	29.0	96.7	306.6	854.9	1,936.3	3,086.7
1918	92.8	18.6	3.1	1.6	3.7	9.8	25.8	89.3	300.6	809.4	1,810.9	2,943.8
1919	86.2	15.1	2.3	1.3	3.0	7.2	23.1	83.5	279.1	761.9	1,780.5	3,143.4
1920	87.9	15.5	2.5	1.2	3.0	6.7	22.1	85.1	279.3	798.7	1,885.5	3,281.3
1921	85.1	11.2	2.0	0.8	2.2	6.1	21.0	78.7	269.9	773.1	1,846.4	3,283.6
1922	88.8	9.8	1.9	1.2	2.1	6.0	21.9	84.6	278.7	810.8	1,925.1	3,308.9
1923	93.4	14.4	1.9	0.8	2.1	6.0	23.3	87.0	290.5	831.2	1,989.1	3,582.7
1924	95.3	12.0	1.9	0.8	2.2	5.7	24.0	93.5	291.6	838.7	2,010.0	3,461.4
1925	87.6	10.1	2.0	1.2	2.3	5.8	20.7	83.8	261.6	745.9	1,875.0	3,361.9
1926	91.0	12.1	1.5	0.8	2.3	6.3	22.9	88.4	262.5	774.1	1,897.2	3,415.3
1927	87.9	10.3	1.7	0.9	2.0	5.8	22.3	83.4	254.8	737.5	1,819.5	3,190.0
1928	92.1	10.8	1.8	1.0	2.2	6.9	24.8	90.0	266.1	760.5	1,880.6	3,223.7
1929	91.3	9.2	1.9	0.9	2.5	6.3	25.1	89.5	270.0	739.4	1,798.8	2,919.1
1930	88.8	10.4	1.5	0.8	2.3	6.5	24.1	88.0	263.3	708.9	1,662.7	2,733.6
1931	87.3	8.8	1.4	0.8	2.0	6.2	23.6	83.9	255.9	685.8	1,601.2	2,719.2
1932	87.4	7.3	1.4	0.8	2.1	5.9	21.9	80.5	252.7	664.3	1,622.0	2,750.6
1933	83.5	7.4	1.9	0.8	1.9	6.0	21.2	77.5	238.5	631.4	1,508.6	2,627.4
1934	84.6	9.1	1.5	0.8	2.4	5.8	22.1	77.8	233.2	621.9	1,538.7	2,494.0
1935	85.1	14.9	1.3	1.2	2.2	6.0	21.3	77.2	235.0	605.5	1,506.2	2,490.9
1936	89.9	17.2	2.0	1.5	2.6	6.0	23.5	79.5	239.2	627.7	1,561.9	2,636.4
1937	85.7	12.9	1.8	1.0	2.3	6.7	22.4	76.3	221.1	587.2	1,480.3	2,360.4
1938	85.1	13.9	1.9	1.0	2.1	5.9	21.3	72.6	218.2	579.9	1,434.3	2,338.8
1939	86.9	10.0	1.5	0.7	2.0	5.9	21.2	73.3	214.1	583.2	1,450.4	2,436.1

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	107.7	91.8	16.9	4.4	8.4	15.8	39.3	127.1	311.2	802.2	1,851.4	3,068.9
1901	106.8	95.1	11.8	4.0	5.9	14.8	38.5	126.4	311.4	798.1	1,895.5	2,887.3
1902	105.1	77.5	12.4	3.4	5.0	15.0	39.0	123.1	318.4	793.0	1,786.2	3,028.1
1903	105.6	67.9	11.0	2.7	6.8	13.2	37.4	119.8	319.7	790.7	1,876.9	3,034.2
1904	110.1	71.2	11.7	2.9	5.7	13.2	34.4	117.2	331.9	835.7	2,009.6	3,367.7
1905	108.4	63.6	8.5	3.1	5.4	11.4	37.3	120.9	327.8	831.1	1,940.5	3,144.3
1906	100.1	54.7	7.8	2.1	4.5	12.2	31.9	114.3	308.8	808.8	1,892.6	3,466.2
1907	106.0	49.5	6.8	2.5	4.6	12.6	34.4	115.0	332.5	848.8	2,042.5	3,640.0
1908	97.2	48.2	6.2	1.5	4.3	12.3	30.4	103.8	316.0	784.8	1,873.4	3,226.3
1909	97.0	46.2	4.4	2.0	3.1	10.3	30.9	102.5	299.5	794.2	1,864.4	3,199.6
1910	97.5	44.9	4.3	1.6	4.1	9.8	30.4	103.8	295.6	795.1	1,942.6	3,164.4
1911	93.3	36.1	4.1	1.3	3.5	9.8	29.5	102.2	282.4	776.8	1,831.6	3,238.2
1912	92.6	27.7	3.6	1.4	3.3	9.4	29.3	99.3	290.8	766.6	1,797.5	3,086.0
1913	92.4	20.8	3.0	1.4	3.1	8.3	27.3	103.6	285.1	765.9	1,840.1	3,034.1
1914	94.5	18.1	2.7	1.1	3.3	8.7	27.9	103.0	288.1	807.4	1,812.0	3,138.7
1915	95.9	14.5	2.5	1.2	2.8	7.8	29.1	102.2	282.7	807.0	1,910.4	3,137.9
1916	95.9	14.0	2.3	1.3	3.2	8.7	29.4	105.4	294.5	800.6	1,918.3	3,146.6
1917	97.6	17.1	2.4	1.2	3.0	8.2	31.0	106.0	301.6	819.9	1,915.6	3,268.4
1918	95.2	13.5	2.5	1.5	4.0	8.3	30.2	108.9	290.7	807.8	1,834.1	3,081.3
1919	93.7	11.9	2.6	1.2	3.1	7.5	26.9	101.5	288.9	783.1	1,892.3	3,172.5
1920	98.2	10.9	2.1	1.2	3.0	8.3	28.7	104.8	295.2	819.8	2,035.9	3,540.3
1921	93.5	8.3	1.6	0.9	2.5	6.0	25.8	100.6	279.3	784.6	1,940.7	3,456.7
1922	95.4	8.4	1.7	0.8	2.1	6.7	26.6	102.9	275.8	804.0	2,005.2	3,741.3
1923	98.1	8.3	1.5	1.0	2.2	6.6	27.9	106.9	283.8	795.5	2,019.2	3,881.7
1924	99.1	9.8	1.4	0.7	2.0	6.4	28.4	109.8	279.4	796.5	2,039.8	3,866.1
1925	91.4	9.4	1.4	0.8	2.5	5.7	24.7	96.6	254.6	726.3	1,859.3	3,773.0
1926	91.6	7.3	1.7	1.0	2.3	6.1	26.2	95.4	258.5	705.0	1,840.7	3,642.1
1927	88.3	7.1	1.2	0.9	2.0	5.9	25.1	93.5	241.3	693.3	1,745.2	3,288.6
1928	91.9	7.1	1.9	0.9	2.7	7.0	27.0	98.9	256.2	709.3	1,783.7	3,376.5
1929	90.3	7.0	1.7	1.1	2.4	6.2	26.7	99.1	248.0	698.4	1,716.2	2,999.2
1930	89.2	6.2	1.3	0.7	2.4	6.3	26.1	97.2	249.4	665.2	1,640.0	2,916.0
1931	86.2	7.7	1.3	0.8	2.0	6.2	25.6	93.4	242.5	626.9	1,527.2	2,745.6
1932	87.5	6.0	1.3	0.9	2.0	6.0	25.4	91.0	234.7	631.8	1,551.2	2,729.5
1933	84.6	7.3	1.5	0.8	1.9	5.9	23.7	88.7	228.5	589.0	1,508.4	2,595.1
1934	86.5	6.1	1.7	0.6	2.0	6.0	25.4	89.6	226.2	588.5	1,521.9	2,569.3
1935	86.4	11.3	1.8	1.1	2.0	5.9	24.9	89.2	221.2	569.8	1,470.3	2,566.1
1936	92.2	11.0	1.5	1.1	2.1	5.9	27.0	90.6	229.0	586.1	1,588.3	2,701.5
1937	87.8	9.5	1.9	0.9	2.2	5.6	25.5	86.0	215.7	548.0	1,453.1	2,518.7
1938	86.7	10.2	1.5	1.2	2.0	5.8	24.7	85.9	204.7	535.7	1,397.0	2,399.1
1939	88.8	7.6	1.2	0.7	1.7	5.6	24.6	84.5	208.7	535.0	1,408.6	2,471.0

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	95.6	---	---	---	---	---	---	---	---	---	---	---
1911	91.5	---	---	---	---	---	---	---	---	---	---	---
1912	91.6	---	---	---	---	---	---	---	---	---	---	---
1913	90.5	---	---	---	---	---	---	---	---	---	---	---
1914	93.0	20.4	2.9	1.3	3.1	8.1	25.8	93.1	289.7	829.1	1,847.6	3,106.0
1915	93.7	17.5	2.5	1.3	2.4	6.6	25.9	92.2	286.8	822.8	1,914.5	3,184.1
1916	94.4	14.2	2.3	1.1	2.9	7.3	25.6	94.7	292.7	826.7	1,914.8	3,186.8
1917	95.4	---	---	---	---	---	---	---	---	---	---	---
1918	93.9	15.6	2.7	1.5	3.3	8.0	24.4	91.6	286.0	803.8	1,841.2	3,093.6
1919	90.5	13.2	2.2	1.2	2.6	6.5	21.6	83.7	275.9	770.5	1,862.2	3,285.1
1920	93.4	13.2	2.3	1.1	2.6	6.3	21.4	86.1	277.7	807.0	1,992.1	3,540.4
1921	89.5	9.8	1.7	0.7	1.9	5.0	19.5	80.8	265.2	777.1	1,916.8	3,468.3
1922	92.3	8.8	1.6	0.9	1.7	5.0	19.7	81.4	266.4	805.8	1,996.7	3,658.0
1923	95.8	11.1	1.7	0.8	1.8	5.0	21.1	84.7	274.5	808.1	2,034.2	3,841.7
1924	96.4	10.3	1.4	0.7	1.7	4.5	20.2	87.5	272.2	808.5	2,046.6	3,768.5
1925	88.5	9.3	1.6	0.9	2.0	4.3	17.6	76.4	243.6	727.2	1,885.9	3,643.7
1926	90.0	9.0	1.5	0.8	1.9	4.6	18.3	76.0	245.2	731.1	1,890.2	3,632.3
1927	87.1	8.7	1.4	0.8	1.7	4.4	18.0	73.4	233.3	707.0	1,804.6	3,336.5
1928	90.5	9.2	1.7	0.9	2.0	4.8	18.7	75.9	242.4	726.1	1,854.8	3,419.3
1929	89.1	7.6	1.7	0.9	2.0	4.2	18.5	75.6	240.1	710.6	1,772.9	3,042.7
1930	86.9	8.5	1.4	0.7	1.8	4.3	17.5	73.4	235.4	675.1	1,669.8	2,907.7
1931	85.3	7.9	1.3	0.8	1.6	4.3	17.8	71.7	231.0	647.5	1,577.0	2,822.0
1932	85.9	6.6	1.3	0.8	1.7	4.3	17.1	68.4	224.5	639.4	1,601.9	2,825.7
1933	82.6	6.8	1.7	0.7	1.7	4.2	16.5	67.0	214.2	602.3	1,521.5	2,685.8
1934	83.6	7.7	1.5	0.7	1.7	4.2	16.7	66.7	209.2	593.8	1,548.6	2,592.9
1935	83.8	13.1	1.6	1.1	1.8	4.1	16.1	66.6	207.2	577.3	1,505.0	2,607.4
1936	88.9	14.2	1.8	1.2	2.1	4.1	17.7	67.9	212.2	593.7	1,590.5	2,763.9
1937	84.5	11.4	1.8	0.9	2.0	4.3	17.0	64.0	197.9	552.3	1,483.7	2,542.1
1938	83.5	11.6	1.7	1.0	1.7	4.2	16.0	61.7	190.3	545.3	1,427.6	2,450.7
1939	85.4	8.4	1.3	0.7	1.6	4.1	16.3	62.0	188.9	545.9	1,440.4	2,539.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	94.4	---	---	---	---	---	---	---	---	---	---	---
1911	90.4	---	---	---	---	---	---	---	---	---	---	---
1912	91.3	---	---	---	---	---	---	---	---	---	---	---
1913	89.7	---	---	---	---	---	---	---	---	---	---	---
1914	92.5	23.6	3.1	1.4	3.1	8.1	26.1	90.3	299.7	860.0	1,884.0	3,040.4
1915	92.9	20.5	2.6	1.4	2.3	6.4	25.4	89.4	302.9	848.3	1,916.9	3,205.8
1916	93.7	14.9	2.4	1.0	2.8	7.3	25.4	93.2	304.6	861.6	1,903.5	3,182.0
1917	94.4	---	---	---	---	---	---	---	---	---	---	---
1918	93.4	18.3	3.0	1.5	3.2	8.7	23.5	84.8	295.2	811.2	1,834.1	2,986.6
1919	87.4	14.8	2.1	1.2	2.6	6.4	20.6	77.4	277.5	766.0	1,813.6	3,262.3
1920	88.7	15.7	2.5	1.2	2.6	5.7	19.2	78.3	274.1	801.0	1,920.8	3,397.2
1921	86.0	10.9	1.9	0.7	1.8	5.2	18.1	72.7	266.1	778.1	1,873.1	3,370.2
1922	89.6	9.6	1.6	1.0	1.7	4.8	18.1	75.7	274.0	818.8	1,962.7	3,389.9
1923	94.0	14.2	1.9	0.7	1.8	4.9	19.9	78.2	283.0	832.2	2,019.9	3,637.9
1924	95.1	11.0	1.7	0.7	1.8	4.4	19.3	82.0	282.5	836.9	2,038.6	3,498.2
1925	87.2	9.6	1.9	1.0	2.0	4.6	16.7	72.1	252.3	743.3	1,901.1	3,389.9
1926	90.3	11.0	1.4	0.7	2.0	4.8	17.3	74.7	251.6	772.7	1,927.5	3,472.1
1927	87.5	10.3	1.7	0.8	1.7	4.6	17.5	70.6	244.1	736.1	1,850.2	3,245.8
1928	91.0	11.1	1.7	1.0	1.8	5.0	18.5	73.2	251.5	758.8	1,906.9	3,310.2
1929	90.0	8.5	1.7	0.9	2.0	4.3	18.5	73.3	254.9	734.9	1,819.3	2,978.4
1930	86.9	10.7	1.5	0.8	1.7	4.5	17.1	70.6	245.2	700.8	1,681.4	2,780.7
1931	86.3	8.7	1.4	0.8	1.7	4.5	17.2	69.8	241.8	682.5	1,613.6	2,794.7
1932	86.2	7.4	1.3	0.8	1.7	4.4	16.4	65.4	236.6	659.8	1,637.6	2,814.5
1933	82.5	7.1	1.9	0.7	1.7	4.3	15.8	63.7	223.7	629.0	1,522.9	2,677.7
1934	83.0	9.4	1.4	0.7	2.0	4.2	15.9	62.0	216.8	614.5	1,558.9	2,539.5
1935	83.4	15.3	1.3	1.1	1.9	4.2	15.2	62.3	216.9	599.2	1,524.1	2,569.8
1936	88.0	17.1	2.1	1.4	2.3	4.3	17.1	64.1	221.2	617.2	1,578.7	2,715.6
1937	83.8	13.6	1.8	0.9	2.0	4.7	16.0	61.0	204.3	575.8	1,499.7	2,444.6
1938	83.0	13.9	1.8	1.0	1.8	4.4	15.2	57.4	199.8	568.3	1,452.2	2,401.5
1939	84.8	9.1	1.4	0.7	1.8	4.2	15.4	58.1	195.4	574.8	1,462.1	2,508.7

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	96.8	---	---	---	---	---	---	---	---	---	---	---
1911	92.7	---	---	---	---	---	---	---	---	---	---	---
1912	92.0	---	---	---	---	---	---	---	---	---	---	---
1913	91.3	---	---	---	---	---	---	---	---	---	---	---
1914	93.4	17.1	2.7	1.1	3.1	8.0	25.4	96.3	278.8	797.8	1,814.2	3,157.3
1915	94.5	14.5	2.5	1.1	2.6	7.0	26.4	95.3	269.4	796.9	1,912.3	3,167.3
1916	95.2	13.6	2.2	1.2	2.9	7.4	25.8	96.4	279.8	791.3	1,925.0	3,190.5
1917	96.5	---	---	---	---	---	---	---	---	---	---	---
1918	94.5	12.7	2.5	1.4	3.4	7.3	25.4	99.2	275.9	796.1	1,847.6	3,174.6
1919	93.7	11.6	2.4	1.1	2.6	6.6	22.7	90.7	274.2	775.2	1,906.1	3,302.1
1920	98.4	10.5	2.1	1.1	2.6	6.8	23.9	94.7	281.5	813.3	2,057.0	3,648.3
1921	93.2	8.7	1.4	0.8	2.0	4.7	21.0	89.8	264.3	776.1	1,956.6	3,543.1
1922	95.0	8.1	1.7	0.7	1.6	5.3	21.5	87.7	258.3	792.6	2,027.6	3,863.7
1923	97.7	7.9	1.5	0.9	1.8	5.2	22.3	91.8	265.5	783.4	2,047.3	4,000.0
1924	97.8	9.5	1.1	0.7	1.6	4.7	21.1	93.4	261.1	779.3	2,053.9	3,978.5
1925	89.8	9.0	1.3	0.8	2.0	4.1	18.7	80.9	234.2	710.7	1,872.0	3,841.1
1926	89.7	6.9	1.6	0.9	1.9	4.3	19.4	77.4	238.3	688.4	1,855.9	3,757.0
1927	86.7	7.0	1.1	0.8	1.7	4.2	18.4	76.5	221.7	677.1	1,762.5	3,407.0
1928	89.9	7.3	1.8	0.9	2.1	4.6	18.9	78.7	232.5	692.4	1,806.3	3,503.5
1929	88.3	6.6	1.6	1.0	1.9	4.2	18.6	78.1	224.3	685.6	1,729.4	3,091.6
1930	86.9	6.2	1.2	0.7	1.8	4.1	17.8	76.5	225.0	648.5	1,658.9	3,003.1
1931	84.2	7.1	1.3	0.8	1.6	4.1	18.4	73.7	219.4	611.5	1,542.5	2,842.5
1932	85.7	5.8	1.2	0.8	1.7	4.2	17.8	71.6	211.6	618.6	1,568.4	2,834.1
1933	82.6	6.5	1.4	0.7	1.7	4.1	17.2	70.5	204.2	575.0	1,520.1	2,692.0
1934	84.2	5.9	1.7	0.6	1.5	4.2	17.5	70.8	201.3	572.7	1,538.9	2,633.1
1935	84.1	10.8	1.8	1.1	1.6	3.9	17.1	71.2	197.0	555.0	1,487.1	2,635.7
1936	89.8	11.3	1.5	1.0	1.8	4.0	18.4	72.1	202.6	570.0	1,601.5	2,800.3
1937	85.3	9.2	1.9	0.8	1.9	3.9	18.1	67.2	191.3	528.8	1,469.0	2,615.5
1938	84.0	9.3	1.5	1.1	1.7	4.1	16.9	66.3	180.2	522.4	1,405.1	2,487.8
1939	86.0	7.6	1.1	0.7	1.5	3.9	17.2	66.2	182.0	517.4	1,420.6	2,562.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	102.3	---	---	---	---	---	---	---	---	---	---	---
1911	98.8	---	---	---	---	---	---	---	---	---	---	---
1912	99.1	---	---	---	---	---	---	---	---	---	---	---
1913	104.7	---	---	---	---	---	---	---	---	---	---	---
1914	106.8	*	*	*	7.9	19.5	66.2	217.6	506.0	999.4	1,573.5	2,600.9
1915	111.1	*	*	*	6.7	21.2	72.8	216.4	574.0	966.6	1,630.7	2,534.8
1916	98.3	*	*	2.5	6.4	27.6	72.0	219.1	510.6	911.1	1,631.8	2,489.0
1917	102.7	---	---	---	---	---	---	---	---	---	---	---
1918	94.6	22.6	4.1	2.4	9.6	22.3	72.8	199.0	487.0	903.9	1,446.3	2,180.0
1919	84.0	16.1	4.7	2.3	7.4	16.8	61.1	192.6	413.3	801.9	1,433.8	1,936.1
1920	87.7	14.0	*	1.9	6.2	21.2	66.9	198.8	440.2	844.3	1,457.4	2,282.3
1921	85.7	*	*	2.0	6.5	17.8	65.1	192.2	426.0	807.6	1,515.7	2,471.9
1922	89.7	11.7	3.5	2.1	5.8	19.1	68.7	227.0	436.2	833.8	1,496.9	2,505.5
1923	95.0	14.0	*	1.9	5.3	18.5	72.4	233.8	475.7	906.2	1,525.0	2,877.5
1924	104.6	17.0	4.2	1.4	5.8	20.8	90.0	265.3	486.5	980.2	1,681.3	2,926.9
1925	99.8	14.9	2.6	1.7	6.3	19.8	77.8	252.9	479.5	899.7	1,547.4	3,101.8
1926	104.3	17.2	*	1.7	5.7	22.1	92.4	280.7	492.0	905.4	1,492.0	2,651.1
1927	97.3	*	*	1.8	4.7	19.5	84.5	256.9	467.4	874.8	1,390.8	2,356.6
1928	105.6	*	2.4	1.4	6.2	25.2	96.1	284.2	514.6	891.7	1,459.7	2,341.4
1929	105.8	13.0	2.9	1.6	6.2	23.4	96.8	282.8	522.9	866.9	1,493.1	2,263.0
1930	108.2	*	*	1.4	6.9	24.5	98.5	286.3	550.8	899.2	1,344.4	2,202.3
1931	100.6	11.0	*	1.1	5.0	22.5	90.0	261.4	506.3	814.6	1,333.7	1,931.8
1932	101.1	*	2.1	1.3	4.7	20.3	85.7	263.1	514.5	793.7	1,311.6	1,936.8
1933	97.3	11.3	2.0	1.2	3.9	20.6	77.8	244.6	496.1	737.5	1,296.6	1,927.7
1934	102.7	*	2.5	0.9	5.7	20.1	88.7	256.1	506.9	779.5	1,225.6	2,039.5
1935	102.9	13.4	*	1.6	4.9	21.9	86.7	253.5	512.3	742.1	1,201.2	1,878.0
1936	110.0	13.7	*	1.8	4.6	21.4	93.7	262.2	531.4	795.2	1,326.3	1,867.4
1937	106.2	9.8	*	1.4	4.6	21.2	86.1	261.1	494.7	776.8	1,171.5	1,635.0
1938	107.1	15.3	2.1	1.7	4.4	19.6	84.7	264.6	499.2	722.9	1,199.5	1,681.4
1939	109.2	11.9	1.9	1.1	3.4	20.2	81.4	260.0	516.0	725.5	1,228.1	1,715.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	83.9	---	---	---	---	---	---	---	---	---	---	---
1911	89.4	---	---	---	---	---	---	---	---	---	---	---
1912	88.0	---	---	---	---	---	---	---	---	---	---	---
1913	92.8	---	---	---	---	---	---	---	---	---	---	---
1914	94.5	*	*	*	8.0	15.6	55.0	172.8	446.2	886.6	1,401.3	2,339.9
1915	95.4	*	*	*	*	17.8	61.5	168.4	465.8	808.8	1,405.7	2,491.8
1916	89.8	*	*	*	6.3	28.4	61.9	178.6	413.2	816.7	1,508.7	2,410.3
1917	92.3	---	---	---	---	---	---	---	---	---	---	---
1918	84.7	*	*	*	8.9	23.8	55.4	149.5	396.4	775.8	1,348.3	2,422.2
1919	74.0	*	*	2.5	7.1	16.2	50.4	151.7	302.3	698.0	1,227.2	1,960.5
1920	79.4	*	*	*	6.1	18.6	53.7	163.5	358.2	762.9	1,283.2	2,083.3
1921	75.3	*	*	*	6.2	15.8	52.8	147.7	327.1	692.5	1,393.4	2,333.1
1922	80.3	*	5.4	2.2	5.4	18.1	61.3	178.1	341.9	695.7	1,363.0	2,502.5
1923	87.7	*	*	*	5.1	17.6	59.7	182.9	391.6	816.5	1,511.3	2,995.3
1924	97.5	21.4	*	*	6.1	18.7	74.9	222.5	415.4	866.5	1,564.4	3,064.6
1925	91.8	*	*	2.5	5.6	18.9	65.6	216.4	390.2	788.8	1,455.4	3,054.0
1926	98.3	23.2	*	*	5.6	21.8	85.0	244.9	412.6	797.4	1,406.1	2,794.7
1927	91.1	*	*	*	4.7	17.6	74.5	222.5	398.3	760.9	1,329.7	2,609.0
1928	101.6	*	*	*	5.7	24.0	88.7	256.8	443.0	787.2	1,490.7	2,423.5
1929	103.1	*	*	1.6	7.0	23.8	91.2	250.0	457.5	812.2	1,481.5	2,377.6
1930	106.4	*	*	1.6	7.2	24.0	93.7	260.2	492.7	840.5	1,359.2	2,305.9
1931	96.5	*	*	*	4.7	21.8	87.1	225.1	435.4	738.7	1,397.3	2,029.4
1932	97.9	*	*	*	5.4	19.7	75.8	233.0	457.4	734.7	1,363.6	2,162.1
1933	92.7	*	*	*	4.0	20.6	73.2	214.3	423.4	667.8	1,276.0	2,179.8
1934	98.6	*	*	1.5	5.7	20.0	81.7	228.2	438.5	727.7	1,209.9	2,086.9
1935	99.6	*	*	1.7	4.6	21.7	79.1	230.2	461.7	693.2	1,213.5	1,781.7
1936	106.5	18.1	*	2.3	5.2	21.0	84.1	239.7	465.7	769.6	1,285.1	1,923.2
1937	102.9	*	*	*	4.7	23.5	82.2	237.0	433.8	735.8	1,159.2	1,599.3
1938	104.3	*	*	1.6	4.6	19.0	77.5	234.5	452.0	726.2	1,137.6	1,771.0
1939	105.8	16.7	*	*	3.9	20.9	74.3	237.5	454.2	685.2	1,255.5	1,779.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

VASCULAR LESIONS AFFECTING CENTRAL NERVOUS SYSTEM

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	123.3	---	---	---	---	---	---	---	---	---	---	---
1911	109.3	---	---	---	---	---	---	---	---	---	---	---
1912	111.4	---	---	---	---	---	---	---	---	---	---	---
1913	117.5	---	---	---	---	---	---	---	---	---	---	---
1914	120.0	*	*	*	7.8	23.6	80.0	277.5	581.5	1,127.7	1,735.1	2,783.4
1915	128.0	*	*	*	*	24.9	86.5	281.6	711.8	1,146.0	1,843.8	2,564.8
1916	107.1	*	*	*	6.5	26.8	83.3	272.2	635.1	1,021.5	1,751.5	2,545.7
1917	113.5	---	---	---	---	---	---	---	---	---	---	---
1918	104.7	*	*	*	10.3	21.0	91.8	265.7	603.8	1,052.6	1,542.1	2,007.2
1919	94.2	*	*	*	7.6	17.4	72.6	248.5	558.7	925.0	1,639.5	1,918.3
1920	96.1	*	*	*	6.2	23.5	81.2	246.5	547.5	941.0	1,634.0	2,431.2
1921	96.2	*	*	*	6.9	19.7	78.6	251.4	554.9	945.8	1,639.9	2,570.6
1922	99.2	*	*	*	6.2	20.0	76.6	290.2	560.1	1,002.7	1,632.9	2,649.3
1923	102.2	*	*	2.2	5.4	19.4	85.7	297.9	585.9	1,016.5	1,538.9	2,798.9
1924	111.8	*	*	*	5.4	22.6	105.7	317.9	579.5	1,119.7	1,798.7	2,836.1
1925	107.9	*	*	*	7.0	20.6	90.3	297.5	595.2	1,033.0	1,638.9	3,133.6
1926	110.3	*	*	2.0	5.9	22.3	100.1	323.3	594.9	1,034.9	1,576.9	2,554.2
1927	103.5	*	*	1.9	4.8	21.3	94.7	297.7	558.0	1,010.0	1,451.2	2,180.4
1928	109.5	*	*	*	6.6	26.3	103.3	316.2	609.2	1,014.1	1,429.2	2,282.4
1929	108.5	*	*	1.7	5.4	23.0	102.4	321.0	609.0	929.7	1,504.3	2,178.2
1930	110.0	*	*	*	6.6	25.0	103.2	316.5	626.4	965.7	1,330.4	2,124.2
1931	104.6	*	*	*	5.2	23.2	92.8	303.0	597.5	899.9	1,273.3	1,858.4
1932	104.2	*	*	1.8	4.1	20.9	95.2	297.3	587.2	859.7	1,262.4	1,767.8
1933	101.8	*	*	1.4	3.9	20.6	82.2	278.7	587.4	815.2	1,316.1	1,739.2
1934	106.8	*	*	*	5.6	20.1	95.5	287.1	591.9	836.9	1,240.6	2,004.1
1935	106.1	*	*	*	5.1	22.2	94.1	279.2	574.3	796.1	1,189.5	1,949.6
1936	113.4	*	*	*	4.1	21.8	103.0	286.9	610.9	823.4	1,365.3	1,826.0
1937	109.4	*	*	1.5	4.4	19.2	89.8	287.2	567.4	821.6	1,183.2	1,661.3
1938	109.9	16.4	*	1.9	4.2	20.2	91.6	297.1	554.8	719.3	1,258.2	1,615.4
1939	112.6	*	*	3.0	19.5	88.2	284.1	587.7	769.4	1,202.1	1,668.2	

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Notes: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes ...	133	235	280
Male....	65	104	127
Female....	68	131	153

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

INFLUENZA AND PNEUMONIA

ICD Revision	Years in use	ICD Nos.
Fifth	1939-33	107-109
Fourth	1930-1938	11, 107-109
Third	1921-1929	11, 100, 101
Second	1910-1920	10, 91, 92
First	1900-1909	10, 92, 93

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	202.2	1,682.1	386.6	38.2	48.1	76.2	114.1	175.2	229.4	692.4	1,531.3	2,899.5
1901	197.2	1,520.3	315.3	34.4	49.3	76.9	112.2	168.9	332.7	724.6	1,746.8	3,610.2
1902	161.3	1,650.8	356.3	31.5	37.4	56.4	84.1	127.2	232.2	477.5	1,049.0	1,916.1
1903	169.3	1,540.0	308.8	30.6	37.8	61.6	94.4	143.2	269.0	567.3	1,327.9	2,551.0
1904	192.1	1,645.7	332.0	32.1	42.3	72.2	117.4	171.3	322.1	656.0	1,471.7	3,235.5
1905	169.3	1,598.6	298.4	27.5	35.0	57.0	94.6	142.0	271.9	583.0	1,349.8	2,922.6
1906	156.3	1,649.9	324.9	26.4	33.2	51.4	82.9	126.0	239.5	492.5	1,089.2	2,168.0
1907	180.0	1,620.7	310.3	28.7	37.0	62.7	100.7	152.7	297.5	663.5	1,570.0	3,278.9
1908	150.9	1,518.6	275.9	26.4	29.4	45.0	75.5	124.7	238.0	536.0	1,305.6	2,613.9
1909	148.1	1,619.8	289.1	24.6	26.8	42.7	73.6	113.0	228.8	509.3	1,166.8	2,360.1
1910	155.9	1,629.0	285.6	25.4	29.2	46.9	78.8	125.7	250.2	556.6	1,292.8	2,555.0
1911	145.4	1,409.3	244.2	23.3	28.6	43.0	73.9	120.6	235.4	551.7	1,325.1	2,792.2
1912	138.4	1,491.0	238.2	20.2	26.1	41.6	69.4	110.8	219.9	493.3	1,172.4	2,396.5
1913	140.8	1,502.5	250.3	23.2	27.2	42.4	73.2	112.9	214.8	483.7	1,160.2	2,415.0
1914	132.4	1,439.8	214.8	21.9	25.5	38.7	67.5	110.2	211.5	460.1	1,096.3	2,244.6
1915	145.9	1,489.4	205.5	21.8	28.0	43.0	71.0	121.1	240.0	567.9	1,371.5	2,930.4
1916	163.3	1,414.0	221.7	23.0	35.0	51.4	87.7	140.4	277.3	661.3	1,631.5	3,698.8
1917	164.5	1,474.5	211.5	24.0	38.9	59.3	98.1	148.8	281.4	614.6	1,503.0	3,187.4
1918	588.5	2,273.3	718.0	176.2	580.5	992.6	554.8	347.8	381.9	646.3	1,179.0	2,230.6
1919	223.0	1,594.2	293.9	63.3	141.4	235.9	181.0	163.9	233.2	459.6	913.9	1,842.2
1920	207.3	1,495.0	283.7	45.1	101.3	180.6	164.0	164.9	255.8	545.9	1,194.3	2,379.1
1921	98.7	979.5	146.8	22.0	26.3	32.7	51.6	78.2	145.5	335.4	842.6	1,841.9
1922	132.3	1,365.0	190.0	24.0	37.0	51.2	77.7	108.1	193.4	457.1	1,126.0	2,530.3
1923	151.7	1,356.6	200.7	26.0	43.5	56.0	87.9	126.9	228.7	549.8	1,431.9	3,474.0
1924	115.2	1,184.8	151.8	20.0	32.2	41.0	67.9	103.1	170.4	379.9	953.0	2,361.3
1925	121.7	1,175.1	150.3	23.3	36.1	44.3	74.1	104.3	177.8	404.9	1,071.6	2,758.5
1926	141.7	1,400.5	198.1	23.0	35.4	50.4	83.9	122.7	204.5	470.0	1,296.9	3,245.4
1927	102.2	1,016.0	129.1	18.8	27.3	37.4	62.4	92.2	153.0	340.0	873.6	2,168.8
1928	142.5	1,321.5	176.6	27.5	42.8	56.6	91.6	128.4	203.7	469.3	1,305.5	3,303.9
1929	146.5	1,321.9	179.1	25.8	41.0	58.3	92.6	132.2	217.7	496.7	1,407.1	3,492.3
1930	102.5	1,090.1	123.1	18.8	27.8	38.9	61.1	95.7	164.6	337.0	834.3	1,884.4
1931	107.5	1,105.4	125.3	19.6	30.4	44.6	67.8	101.0	164.9	340.4	884.0	2,174.7
1932	107.3	1,030.5	115.8	20.4	30.5	42.5	65.2	95.5	162.9	348.4	1,006.5	2,616.2
1933	95.7	990.0	109.0	17.5	25.0	38.0	59.7	88.4	149.3	300.5	836.2	2,168.8
1934	96.9	1,055.3	115.4	17.8	24.6	36.3	57.6	92.3	157.1	302.5	802.9	1,900.2
1935	104.2	1,002.9	111.6	19.9	28.6	43.6	66.6	101.6	166.9	325.9	853.5	2,071.8
1936	119.6	1,090.4	119.3	21.8	35.6	50.1	77.8	118.9	194.1	372.2	977.6	2,330.2
1937	114.9	1,036.4	112.4	17.9	29.4	45.8	72.1	116.9	189.8	357.6	960.3	2,425.9
1938	80.4	901.9	90.2	13.7	18.4	27.0	42.3	70.9	123.3	248.2	677.5	1,592.1
1939	75.7	793.7	70.3	11.4	15.2	22.2	37.1	63.5	113.7	245.6	717.0	1,930.1

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	205.8	1,869.3	396.3	34.2	58.2	87.2	130.1	188.4	312.8	630.9	1,345.5	2,495.1
1901	201.7	1,671.7	328.5	34.6	57.9	88.3	128.0	185.7	321.7	659.0	1,609.0	3,312.5
1902	170.8	1,839.0	369.3	30.6	43.3	66.6	102.0	148.8	240.0	454.5	929.0	1,756.7
1903	174.1	1,685.3	318.1	28.3	42.5	71.7	111.5	164.1	266.5	518.3	1,153.2	2,469.8
1904	206.6	1,829.8	354.6	30.2	51.0	90.8	148.9	204.7	343.5	621.8	1,372.0	2,874.7
1905	176.8	1,785.2	307.9	26.4	39.4	68.4	117.5	164.7	282.4	524.0	1,208.2	2,675.3
1906	188.0	1,845.0	344.1	25.7	41.4	64.1	104.4	152.8	256.3	453.8	984.9	2,040.6
1907	189.1	1,796.7	322.6	28.1	46.7	76.2	125.5	179.1	304.5	623.9	1,395.0	3,006.4
1908	157.0	1,702.8	286.5	26.6	34.7	50.4	89.2	144.8	247.2	493.0	1,216.0	2,367.0
1909	154.5	1,779.0	304.1	24.5	31.6	48.8	89.6	134.3	239.4	473.0	1,044.5	2,177.2
1910	164.0	1,812.4	296.4	25.6	35.7	54.3	98.5	151.7	267.7	522.1	1,157.1	2,301.0
1911	152.4	1,564.7	255.4	23.0	34.8	51.4	91.1	142.4	250.0	519.2	1,182.2	2,597.5
1912	147.2	1,663.7	251.2	20.2	31.4	50.0	86.2	133.8	238.5	470.9	1,065.0	2,259.7
1913	150.6	1,670.5	263.0	22.4	32.8	51.6	93.0	138.8	233.7	460.7	1,069.8	2,346.0
1914	141.0	1,602.0	224.2	22.1	30.3	46.9	84.7	134.3	227.8	446.7	985.7	1,996.3
1915	152.7	1,658.2	214.9	22.4	32.8	50.0	85.9	140.9	253.9	541.9	1,244.2	2,731.4
1916	173.1	1,590.2	236.3	23.2	42.1	59.7	107.6	170.2	296.8	632.8	1,507.6	3,449.6
1917	182.4	1,665.0	224.1	24.6	51.0	79.1	130.8	188.3	313.5	608.4	1,369.4	3,028.7
1918	669.0	2,520.5	712.0	162.5	700.6	1,216.6	691.1	411.8	420.8	655.8	1,112.9	2,111.2
1919	230.0	1,785.1	301.4	59.0	142.5	232.8	204.3	181.9	244.0	444.7	836.4	1,717.5
1920	213.8	1,691.6	295.4	45.4	105.2	172.2	178.8	180.9	267.2	534.0	1,100.0	2,269.5
1921	104.9	1,094.0	156.6	22.8	30.6	35.8	59.8	92.2	156.8	323.9	790.2	1,706.0
1922	139.0	1,436.2	201.4	24.7	41.6	51.7	88.7	123.2	200.0	441.5	1,031.4	2,387.5
1923	158.1	1,504.5	210.2	26.6	49.1	59.4	100.0	147.0	241.2	527.9	1,309.5	3,142.2
1924	126.5	1,325.9	162.2	20.6	37.7	47.3	84.9	127.0	191.1	389.0	916.0	2,208.7
1925	132.2	1,327.0	159.1	23.6	41.5	50.5	88.8	128.1	197.2	410.8	1,028.0	2,639.8
1926	151.7	1,577.1	213.0	24.4	41.2	53.7	98.5	146.8	221.3	465.5	1,229.9	2,956.8
1927	112.5	1,140.4	136.8	19.5	31.5	43.8	78.6	113.3	174.0	351.1	850.9	2,107.3
1928	153.8	1,483.1	188.8	28.1	49.0	62.1	106.4	155.2	226.3	473.7	1,266.3	3,245.4
1929	157.2	1,485.6	187.8	26.3	46.0	63.9	108.7	157.8	240.0	503.8	1,368.1	3,391.3
1930	113.5	1,211.1	129.7	19.3	32.6	45.9	77.7	118.4	186.2	349.1	816.8	1,916.3
1931	118.4	1,232.8	134.0	20.3	35.1	51.2	83.1	124.2	189.5	348.1	865.6	2,171.4
1932	113.8	1,148.9	122.2	20.5	33.4	43.2	72.1	113.0	176.5	357.5	994.9	2,657.1
1933	103.8	1,102.6	116.6	18.0	27.8	40.9	70.7	106.1	167.8	313.2	825.7	2,218.5
1934	108.4	1,183.5	124.3	18.8	28.7	41.8	73.2	115.2	183.4	324.4	800.2	1,887.5
1935	117.0	1,136.5	120.4	21.3	33.9	50.4	82.4	127.2	197.9	347.4	859.2	2,093.8
1936	134.5	1,212.5	126.6	22.7	42.7	57.5	96.6	150.0	229.7	405.5	1,005.4	2,298.3
1937	129.0	1,161.3	121.2	19.0	34.3	51.8	88.0	146.4	227.0	387.4	995.5	2,471.0
1938	89.7	1,012.9	96.0	14.0	21.1	30.8	51.0	89.1	146.4	271.5	701.3	1,630.6
1939	83.1	885.6	74.9	12.0	16.4	24.3	43.5	76.7	136.2	270.7	743.3	1,996.3

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	198.5	1,491.1	376.8	42.1	38.6	65.1	96.9	161.4	345.7	750.8	1,700.7	3,189.2
1901	192.7	1,365.8	302.0	34.3	41.1	65.3	95.3	151.3	343.5	786.7	1,871.9	3,826.4
1902	151.9	1,458.8	343.1	32.4	31.8	46.1	65.1	104.5	224.6	499.3	1,157.3	2,032.3
1903	164.4	1,391.8	299.3	33.2	33.2	51.3	76.1	121.2	271.5	613.7	1,484.8	2,610.4
1904	177.5	1,457.8	309.1	34.0	34.1	53.1	84.0	136.0	300.9	688.4	1,560.7	3,496.9
1905	161.8	1,408.0	288.8	28.6	30.9	45.3	70.3	118.0	261.6	638.8	1,475.7	3,106.2
1906	144.1	1,450.1	305.4	27.0	25.0	37.7	59.3	96.6	222.1	530.3	1,183.5	2,263.4
1907	170.6	1,440.3	297.8	29.3	27.4	48.1	73.4	123.7	290.3	702.1	1,727.9	3,483.8
1908	144.4	1,329.6	265.1	26.3	24.1	39.0	60.2	102.2	228.2	578.6	1,387.7	2,805.0
1909	141.4	1,456.4	273.7	24.7	22.0	36.0	55.9	89.3	217.5	545.2	1,278.4	2,502.9
1910	147.2	1,440.7	274.6	25.1	22.5	38.8	56.9	96.5	231.4	591.0	1,417.0	2,754.4
1911	138.0	1,249.8	232.7	23.6	22.2	33.7	54.7	95.9	219.6	584.5	1,456.8	2,946.4
1912	129.1	1,313.9	225.0	20.2	20.8	32.3	50.7	84.0	199.6	516.0	1,270.9	2,503.9
1913	130.4	1,330.3	237.3	24.0	21.6	32.5	51.1	83.6	194.3	507.0	1,242.7	2,468.6
1914	123.3	1,273.5	205.1	21.8	20.7	29.9	48.4	82.9	193.7	473.7	1,197.9	2,437.7
1915	138.9	1,316.5	196.0	21.2	23.2	35.5	54.6	98.7	224.8	594.4	1,487.7	3,083.8
1916	153.0	1,233.7	206.8	22.8	28.0	42.6	65.8	106.7	255.9	690.4	1,743.7	3,888.4
1917	145.9	1,279.5	198.6	23.3	27.0	38.6	62.4	104.0	246.2	620.9	1,624.1	3,307.3
1918	507.5	2,020.4	724.2	190.5	475.1	781.4	406.5	275.0	339.2	636.5	1,239.0	2,320.5
1919	215.8	1,399.0	286.2	67.7	140.3	238.9	155.5	143.4	221.4	475.1	984.2	1,935.4
1920	200.6	1,293.6	271.7	44.9	97.5	189.1	147.8	146.8	243.4	558.2	1,280.7	2,461.6
1921	92.2	861.3	136.7	21.3	22.0	29.7	42.5	62.6	133.2	347.2	890.8	1,944.9
1922	125.4	1,088.6	178.2	23.2	32.6	50.7	65.8	91.2	186.3	473.2	1,212.8	2,638.7
1923	145.2	1,204.2	190.9	25.5	38.0	52.7	74.9	104.9	215.3	572.5	1,544.4	3,728.0
1924	103.6	1,039.5	141.0	19.4	26.7	34.7	49.4	76.9	147.9	370.5	987.0	2,478.1
1925	110.8	1,018.2	141.1	22.9	30.7	38.2	58.3	78.2	156.8	398.8	1,111.9	2,849.5
1926	131.5	1,218.2	182.6	21.6	29.6	47.0	68.2	96.5	186.2	474.6	1,358.8	3,467.2
1927	91.6	888.4	121.2	18.2	23.2	31.1	45.2	69.3	130.1	328.5	894.6	2,216.1
1928	130.9	1,155.3	163.9	26.8	36.7	51.1	75.8	99.3	179.0	464.7	1,342.2	3,348.7
1929	135.6	1,153.6	170.2	25.4	36.2	52.7	75.6	104.3	193.4	489.3	1,443.8	3,568.9
1930	91.2	965.2	116.4	18.3	23.0	32.0	43.7	70.8	141.2	324.4	850.9	1,860.4
1931	96.4	973.8	116.4	18.8	25.8	38.1	51.7	75.7	138.4	332.4	901.3	2,177.1
1932	100.7	908.2	109.3	20.3	27.6	41.8	57.9	76.5	148.4	339.1	1,017.4	2,585.5
1933	87.4	874.0	101.2	17.0	22.3	35.2	48.4	69.4	129.5	287.5	846.1	2,131.3
1934	85.2	923.4	106.1	16.7	20.5	31.0	41.5	67.6	128.9	280.2	805.3	1,909.8
1935	91.2	865.6	102.5	18.5	23.4	36.9	50.4	74.1	133.7	304.0	848.1	2,055.2
1936	104.5	965.0	111.8	21.0	28.6	42.8	58.5	85.5	156.3	338.5	951.9	2,354.2
1937	100.6	908.1	103.4	16.8	24.7	39.9	55.9	85.3	150.5	327.5	927.6	2,391.9
1938	71.1	788.0	84.1	13.3	15.8	23.2	33.4	51.4	99.0	224.9	655.5	1,563.2
1939	68.2	699.3	65.6	10.8	14.0	20.1	30.6	49.5	89.9	220.6	692.9	1,880.3

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	152.6	---	---	---	---	---	---	---	---	---	---	---
1911	141.9	---	---	---	---	---	---	---	---	---	---	---
1912	135.0	---	---	---	---	---	---	---	---	---	---	---
1913	137.3	---	---	---	---	---	---	---	---	---	---	---
1914	128.4	1,386.5	203.9	20.8	23.3	35.5	63.5	105.7	206.8	455.4	1,100.3	2,286.6
1915	141.5	1,439.1	194.5	20.8	25.6	39.1	66.2	115.6	233.3	560.3	1,372.0	2,990.9
1916	158.3	1,356.2	209.4	21.7	30.0	46.1	82.0	133.8	270.4	657.0	1,639.3	3,778.0
1917	157.1	---	---	---	---	---	---	---	---	---	---	---
1918	572.3	2,152.6	679.1	164.9	557.2	992.6	542.8	334.1	372.1	639.0	1,178.0	2,261.2
1919	211.1	1,491.0	271.4	55.9	127.3	228.4	169.8	152.0	224.6	453.0	912.1	1,892.7
1920	198.4	1,367.2	266.7	41.8	92.9	174.5	154.6	153.6	247.6	542.2	1,204.1	2,429.2
1921	94.7	908.0	136.8	21.2	22.2	29.3	47.2	73.3	141.5	334.5	855.6	1,906.0
1922	127.3	1,180.0	180.7	22.7	31.3	45.9	71.3	99.4	188.4	456.9	1,146.2	2,624.3
1923	142.6	1,229.8	181.8	23.9	35.0	47.3	77.3	114.1	218.5	544.6	1,447.3	3,580.8
1924	105.6	1,056.8	134.9	18.1	23.3	32.0	57.6	91.0	162.0	370.6	959.5	2,425.4
1925	112.2	1,055.3	134.0	21.0	27.0	35.2	63.0	91.8	169.6	399.6	1,080.2	2,818.1
1926	131.2	1,260.2	177.6	20.9	27.2	40.0	72.2	108.4	192.7	462.2	1,305.7	3,320.3
1927	94.5	903.8	114.0	17.2	21.3	30.0	53.6	81.4	144.5	337.1	883.4	2,235.0
1928	133.1	1,186.2	160.5	25.5	34.9	46.7	79.9	113.2	192.2	461.7	1,322.7	3,409.9
1929	135.7	1,184.7	163.5	23.6	32.2	46.3	78.1	114.6	201.6	486.0	1,411.4	3,561.2
1930	94.4	983.6	110.2	17.1	20.9	29.9	51.4	82.8	153.7	331.1	842.0	1,958.4
1931	98.9	999.8	112.9	17.8	23.6	34.2	56.8	86.5	152.9	332.8	891.7	2,270.7
1932	99.4	907.1	102.5	18.9	24.7	34.0	55.6	82.6	150.5	340.5	1,014.7	2,720.2
1933	87.6	850.3	96.0	15.6	19.6	29.8	50.6	76.7	137.8	292.8	842.6	2,262.5
1934	87.7	889.7	100.1	15.9	19.0	27.9	47.1	78.7	144.7	295.6	807.0	1,958.1
1935	96.6	894.1	101.6	18.6	23.0	34.8	55.7	88.3	154.6	320.6	863.9	2,160.6
1936	109.2	970.0	108.5	20.3	28.0	38.7	62.8	100.9	176.4	358.9	980.6	2,390.5
1937	105.1	915.3	100.5	16.1	22.4	34.8	57.8	99.7	174.8	345.7	967.1	2,518.8
1938	73.1	790.9	80.0	12.1	13.5	20.1	33.3	58.6	112.2	241.4	680.5	1,654.9
1939	69.3	689.4	60.0	10.1	11.6	16.4	29.1	52.6	101.8	238.9	721.5	2,020.3

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	160.4	---	---	---	---	---	---	---	---	---	---	---
1911	148.5	---	---	---	---	---	---	---	---	---	---	---
1912	143.5	---	---	---	---	---	---	---	---	---	---	---
1913	146.7	---	---	---	---	---	---	---	---	---	---	---
1914	136.7	1,548.0	213.2	21.0	27.6	43.1	80.7	129.5	223.2	441.3	986.6	2,039.7
1915	147.7	1,604.1	203.8	21.4	29.7	45.4	79.8	135.4	248.5	534.2	1,243.3	2,770.7
1916	166.9	1,526.5	222.3	22.1	35.9	53.4	100.7	162.8	290.8	626.8	1,510.5	3,515.8
1917	172.8	---	---	---	---	---	---	---	---	---	---	---
1918	652.0	2,387.5	673.6	152.8	668.7	1,218.9	680.1	397.8	411.8	648.2	1,111.4	2,145.3
1919	218.2	1,671.3	279.3	53.3	128.6	226.0	193.1	170.0	236.0	438.0	827.3	1,752.8
1920	204.2	1,550.6	278.0	42.2	96.5	166.2	168.3	168.9	261.1	531.2	1,104.1	2,287.5
1921	100.3	1,021.4	146.2	21.9	25.8	31.9	54.9	87.0	153.4	323.3	799.0	1,741.9
1922	133.0	1,345.8	191.1	23.7	35.2	45.5	81.0	114.0	195.7	441.3	1,045.6	2,455.4
1923	147.5	1,362.0	191.0	24.7	39.3	48.9	87.7	133.3	232.8	523.6	1,317.4	3,208.2
1924	115.1	1,185.5	144.8	18.7	27.0	35.9	72.0	113.8	182.8	380.4	918.5	2,254.5
1925	120.9	1,191.4	141.6	21.6	31.1	38.5	75.2	114.0	189.8	406.6	1,029.8	2,656.4
1926	139.6	1,417.9	190.6	22.3	32.1	41.7	84.8	132.0	210.9	458.1	1,232.6	2,995.3
1927	103.6	1,012.8	120.8	17.9	25.1	34.7	68.3	100.9	165.9	347.6	861.5	2,166.2
1928	143.1	1,333.2	172.4	26.2	40.3	50.6	93.0	139.1	215.9	465.4	1,280.6	3,314.5
1929	144.5	1,327.9	171.3	24.2	36.6	49.7	91.8	138.8	223.2	492.8	1,360.5	3,428.0
1930	103.8	1,089.7	115.6	17.8	24.7	34.9	65.4	104.4	175.4	342.4	819.4	1,980.3
1931	108.4	1,114.7	121.0	18.5	27.1	38.6	70.6	108.8	177.4	340.6	867.4	2,265.1
1932	104.7	1,011.6	107.6	19.1	27.4	33.5	61.2	99.1	164.4	349.0	1,001.1	2,748.8
1933	94.4	949.5	102.8	16.2	21.9	31.6	59.9	93.2	155.2	305.3	826.4	2,297.3
1934	97.7	999.7	107.6	17.1	22.8	31.8	60.2	100.0	171.0	316.2	800.5	1,933.1
1935	108.1	1,013.4	110.0	20.0	27.9	40.2	69.2	112.0	185.5	342.0	867.9	2,165.1
1936	121.9	1,076.4	114.4	21.3	33.8	43.4	77.9	129.2	211.7	389.4	1,002.3	2,345.4
1937	117.2	1,023.6	107.7	17.1	26.2	38.5	71.7	126.7	210.8	373.1	997.1	2,553.1
1938	81.0	888.5	84.5	12.6	15.7	22.3	40.5	74.8	134.4	263.9	704.0	1,675.7
1939	75.8	766.1	63.1	10.6	13.0	17.9	34.2	65.1	123.9	264.2	746.0	2,072.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	144.4	---	---	---	---	---	---	---	---	---	---	---
1911	134.9	---	---	---	---	---	---	---	---	---	---	---
1912	126.1	---	---	---	---	---	---	---	---	---	---	---
1913	127.4	---	---	---	---	---	---	---	---	---	---	---
1914	119.7	1,220.7	194.4	20.6	19.0	27.3	44.6	78.9	189.1	469.6	1,204.6	2,479.6
1915	135.1	1,269.8	185.0	20.2	21.4	32.4	51.2	93.3	216.8	586.8	1,489.4	3,161.4
1916	149.3	1,181.3	196.1	21.2	24.1	38.3	61.6	101.2	248.3	687.6	1,755.7	3,978.1
1917	140.8	---	---	---	---	---	---	---	---	---	---	---
1918	492.2	1,911.4	684.8	177.2	459.5	779.3	393.4	262.4	328.7	629.5	1,238.3	2,348.9
1919	203.8	1,305.8	263.4	58.4	126.1	230.9	144.5	131.8	212.1	468.4	988.7	1,997.6
1920	192.5	1,178.5	255.2	41.4	89.4	183.0	139.5	136.5	233.0	553.6	1,295.2	2,536.1
1921	88.8	790.4	127.2	20.5	18.6	26.7	38.9	58.1	128.6	345.9	907.3	2,031.0
1922	121.4	1,008.3	169.9	21.8	27.4	46.3	60.8	83.3	180.5	472.8	1,237.8	2,753.8
1923	137.6	1,092.9	172.3	23.0	30.8	45.6	66.0	93.1	203.1	566.2	1,566.0	3,870.0
1924	95.7	923.5	124.6	17.4	19.6	28.0	41.9	66.4	139.7	360.6	997.1	2,558.2
1925	103.3	914.1	126.1	20.4	22.9	32.0	49.9	67.7	147.8	392.3	1,126.4	2,943.9
1926	122.6	1,096.6	164.2	19.5	22.3	38.3	58.6	82.9	173.1	466.4	1,372.9	3,573.3
1927	85.1	791.3	107.0	16.5	17.5	25.4	37.8	60.3	121.4	326.4	903.7	2,288.4
1928	122.9	1,034.0	148.2	24.7	29.6	42.7	65.9	85.3	166.6	457.8	1,361.8	3,483.3
1929	126.7	1,036.4	155.4	23.0	27.8	42.9	63.4	88.4	178.4	479.0	1,459.0	3,662.5
1930	84.7	873.5	104.7	16.4	17.1	25.0	36.6	59.4	130.5	319.4	863.4	1,941.9
1931	89.2	880.4	104.5	17.0	20.0	29.9	42.3	62.4	126.9	324.8	914.6	2,274.9
1932	94.0	798.6	97.2	18.6	22.0	34.5	49.7	64.8	135.8	331.8	1,027.5	2,698.7
1933	80.6	747.7	88.9	15.0	17.4	28.0	40.8	58.9	119.2	280.0	857.7	2,236.3
1934	77.5	775.8	92.3	14.7	15.2	24.1	33.5	55.7	116.8	274.7	813.0	1,976.9
1935	84.9	770.6	92.9	17.2	18.1	29.4	41.7	62.8	121.9	299.0	860.1	2,157.2
1936	96.3	860.0	102.4	19.3	22.3	34.0	47.1	70.7	139.2	328.3	960.4	2,424.5
1937	92.8	803.4	93.1	15.1	18.7	31.2	43.4	70.9	136.9	318.3	939.4	2,492.9
1938	65.1	689.9	75.5	11.5	11.4	17.8	26.0	41.3	89.0	218.9	659.0	1,639.2
1939	62.7	610.0	56.9	9.5	10.3	14.8	23.8	39.2	78.7	213.8	699.2	1,981.0

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	273.6	---	---	---	---	---	---	---	---	---	---	---
1911	251.4	---	---	---	---	---	---	---	---	---	---	---
1912	239.0	---	---	---	---	---	---	---	---	---	---	---
1913	219.5	---	---	---	---	---	---	---	---	---	---	---
1914	222.1	2,712.9	485.1	47.9	72.5	102.8	148.0	217.6	346.3	608.7	959.8	1,401.8
1915	245.0	2,685.4	481.8	45.6	79.1	121.4	168.2	251.4	436.6	809.9	1,354.4	1,695.5
1916	235.4	2,190.5	392.1	40.4	100.8	129.0	170.8	253.8	422.3	755.6	1,442.1	2,587.7
1917	254.3	---	---	---	---	---	---	---	---	---	---	---
1918	794.7	3,743.9	1,211.6	305.5	831.3	992.4	709.2	548.1	571.9	788.1	1,198.7	1,869.7
1919	351.2	2,650.8	533.3	133.3	271.6	317.9	302.4	308.7	376.3	570.0	945.8	1,347.1
1920	304.4	2,950.2	463.4	77.5	179.5	249.3	267.6	306.4	392.9	609.0	1,017.7	1,862.5
1921	143.0	1,766.0	252.8	30.0	64.8	70.3	99.3	140.6	212.4	350.9	609.8	1,165.2
1922	180.7	2,086.2	276.1	34.5	84.8	102.0	142.2	206.3	266.7	451.0	807.4	1,652.1
1923	242.6	2,604.8	376.7	44.8	116.0	141.5	199.9	277.1	381.1	635.7	1,180.6	2,433.0
1924	212.6	2,454.2	307.9	37.3	109.1	129.0	178.5	245.9	294.8	539.5	846.2	1,731.9
1925	219.2	2,391.0	304.2	44.1	116.7	133.7	195.9	254.5	302.4	500.5	929.3	2,160.5
1926	249.7	2,759.1	389.7	42.8	108.1	150.6	212.0	295.1	382.5	612.5	1,149.1	2,492.6
1927	178.3	2,050.0	265.4	33.3	78.6	106.0	156.6	215.9	276.8	391.5	710.6	1,520.7
1928	226.5	2,478.8	303.2	43.5	104.6	141.4	205.9	285.3	356.9	597.5	1,043.9	2,362.5
1929	244.3	2,536.5	300.5	43.9	110.4	160.8	233.8	311.9	436.6	679.0	1,340.3	2,872.8
1930	175.9	1,949.5	224.9	32.5	82.7	115.6	155.2	227.2	315.8	437.6	708.2	1,212.0
1931	185.6	1,946.4	223.1	34.1	84.5	133.0	173.0	250.3	330.8	468.3	756.0	1,296.6
1932	179.1	1,976.7	220.9	33.0	76.6	114.4	156.7	228.8	334.7	478.6	870.2	1,659.6
1933	166.9	2,007.4	208.7	32.7	67.4	106.1	145.0	207.9	305.1	420.4	733.4	1,337.9
1934	177.9	2,250.0	228.8	32.2	68.6	106.4	153.9	232.3	323.5	406.5	735.8	1,385.3
1935	171.0	1,790.2	183.8	29.6	73.6	117.4	166.6	239.8	332.3	403.1	681.8	1,280.0
1936	211.4	1,971.6	195.1	33.2	95.9	145.3	213.9	307.2	433.1	559.6	929.0	1,791.9
1937	201.3	1,921.9	195.5	31.3	85.9	138.3	201.0	299.4	393.0	519.4	845.6	1,596.9
1938	145.1	1,716.3	160.4	25.3	57.8	85.5	122.4	202.7	272.7	338.3	625.6	1,032.1
1939	131.9	1,561.2	141.4	21.0	44.0	71.8	107.6	182.3	273.2	331.2	640.0	1,126.1

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	290.6	---	---	---	---	---	---	---	---	---	---	---
1911	267.6	---	---	---	---	---	---	---	---	---	---	---
1912	256.1	---	---	---	---	---	---	---	---	---	---	---
1913	238.4	---	---	---	---	---	---	---	---	---	---	---
1914	237.4	2,914.4	504.3	46.7	89.4	122.9	162.6	238.8	352.7	608.6	953.3	1,067.3
1915	262.8	2,968.3	499.2	46.7	98.0	142.9	202.9	260.6	401.7	774.2	1,275.4	1,879.1
1916	263.7	2,463.5	432.7	38.0	126.9	155.1	209.2	287.8	414.2	755.3	1,441.5	2,488.9
1917	313.9	---	---	---	---	---	---	---	---	---	---	---
1918	884.6	4,176.8	1,208.4	274.8	1,041.9	1,187.9	831.5	600.3	581.6	794.0	1,141.9	1,695.5
1919	357.8	2,973.1	541.0	112.4	276.6	310.9	327.8	314.6	365.2	549.0	988.9	1,366.4
1920	319.6	3,342.3	482.6	76.7	189.4	243.6	293.7	318.6	361.0	579.0	1,030.2	2,083.3
1921	156.4	1,913.1	269.4	30.9	77.6	80.4	114.7	152.0	207.6	333.9	639.6	1,312.4
1922	198.1	2,334.2	298.0	33.6	96.8	114.5	167.1	219.3	256.5	445.5	819.0	1,711.5
1923	265.1	2,943.4	391.5	42.8	135.4	166.8	231.7	296.1	354.6	592.3	1,186.8	2,440.1
1924	242.7	2,753.7	325.3	37.8	133.2	164.3	226.4	274.5	304.5	526.1	878.1	1,713.9
1925	250.5	2,738.9	326.4	42.7	137.0	173.6	241.9	288.1	299.1	480.3	998.7	2,457.4
1926	277.7	3,160.6	425.0	44.8	124.8	174.8	251.7	316.7	365.8	591.0	1,185.2	2,537.0
1927	202.0	2,357.6	281.2	33.7	87.5	131.6	191.1	248.2	281.9	410.8	682.7	1,494.4
1928	251.9	2,809.2	320.6	43.7	119.6	164.5	243.0	315.3	352.4	605.1	1,053.3	2,605.3
1929	273.9	2,933.7	317.8	43.5	122.4	189.5	279.3	345.2	447.9	681.3	1,486.0	3,056.9
1930	203.2	2,214.7	243.6	31.6	98.6	142.8	201.6	257.7	322.3	458.5	773.5	1,335.5
1931	211.1	2,199.0	238.7	34.5	101.0	161.6	208.2	279.1	342.4	469.3	835.7	1,315.2
1932	197.5	2,230.3	239.4	32.5	83.8	128.3	180.1	253.5	330.2	491.6	892.6	1,813.1
1933	187.9	2,247.1	224.5	32.8	76.1	120.8	174.5	234.4	324.3	430.2	814.7	1,517.4
1934	204.5	2,546.0	250.9	32.4	77.2	127.7	197.5	268.7	337.3	442.0	795.1	1,478.9
1935	197.3	2,050.2	197.7	31.0	84.3	139.1	207.1	282.1	353.9	423.7	716.5	1,453.1
1936	248.4	2,237.5	214.0	33.2	116.7	179.7	271.9	365.8	456.7	623.6	1,056.3	1,873.5
1937	235.2	2,199.3	217.1	33.3	102.5	168.4	239.6	353.9	432.4	574.1	969.9	1,730.3
1938	168.2	1,953.9	178.3	24.7	66.9	105.7	148.7	241.0	299.5	366.5	657.5	1,223.0
1939	148.9	1,791.7	158.5	22.3	45.1	81.7	128.3	201.3	293.8	350.0	698.8	1,306.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Influenza and Pneumonia, except Pneumonia of Newborn

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	254.2	...	...	...	...	...	...	...	...	...	...	...
1911	233.3	...	...	...	...	...	...	...	...	...	...	...
1912	219.9	...	...	...	...	...	...	...	...	...	...	...
1913	199.1	...	...	...	...	...	...	...	...	...	...	...
1914	205.5	2,513.1	466.4	49.1	56.1	81.1	130.1	189.3	338.2	608.7	965.9	1,635.6
1915	225.8	2,405.6	464.7	44.5	60.7	98.4	125.9	238.9	481.1	850.5	1,429.2	1,567.4
1916	206.2	1,921.7	351.9	42.7	76.7	103.1	127.7	209.2	432.6	755.8	1,442.7	2,658.9
1917	213.5	...	...	...	...	...	...	...	...	...	...	...
1918	703.2	3,318.8	1,214.8	335.7	644.3	808.2	575.3	477.7	559.5	781.3	1,254.3	1,993.9
1919	344.4	2,333.2	525.7	154.1	267.2	324.4	275.1	300.6	390.9	594.9	902.9	1,333.0
1920	288.9	2,568.7	444.2	78.3	170.6	254.5	239.0	289.8	434.6	644.6	1,005.0	1,697.2
1921	129.4	1,621.1	236.1	29.0	53.0	60.9	82.5	125.5	218.6	371.3	579.4	1,060.5
1922	163.2	1,843.2	254.0	35.3	73.8	90.6	116.1	189.5	280.1	480.0	795.7	1,611.3
1923	219.9	2,272.0	361.8	46.8	98.1	118.5	166.6	253.2	415.7	689.0	1,174.2	2,428.3
1924	182.3	2,159.5	290.3	36.8	86.7	96.7	128.7	210.7	282.1	555.9	814.3	1,743.8
1925	187.7	2,047.9	281.6	45.4	97.7	96.5	148.3	213.3	306.7	524.8	860.4	1,963.2
1926	221.6	2,364.6	354.0	40.9	92.5	128.0	171.1	269.3	404.2	638.3	1,113.4	2,462.7
1927	154.5	1,753.4	245.5	33.0	70.3	82.1	121.5	177.6	270.2	368.6	738.2	1,539.1
1928	201.4	2,158.6	285.8	43.3	90.7	120.0	169.3	250.4	362.8	588.6	1,034.6	2,188.2
1929	214.9	2,153.9	283.1	44.3	99.4	134.1	189.1	273.2	421.8	676.3	1,199.1	2,736.7
1930	148.9	1,687.3	206.5	33.4	68.3	90.2	109.7	191.8	307.3	414.0	646.1	1,119.1
1931	160.4	1,697.8	207.6	33.7	69.5	106.3	138.5	217.4	315.9	467.1	680.4	1,282.6
1932	160.9	1,726.7	202.6	33.6	70.1	101.3	134.0	200.8	340.5	464.0	649.1	1,544.5
1933	146.3	1,771.7	193.0	32.7	59.6	92.5	116.3	178.1	280.9	409.6	656.2	1,203.6
1934	151.7	1,959.5	207.0	32.0	60.9	86.7	111.7	191.8	306.3	367.2	679.5	1,315.6
1935	145.3	1,534.9	170.0	28.3	64.0	97.4	127.5	193.1	305.9	380.5	649.0	1,151.3
1936	175.2	1,712.0	176.3	33.2	77.1	113.7	158.2	243.0	404.5	489.3	808.4	1,731.3
1937	168.2	1,651.7	174.2	29.3	70.8	110.6	163.9	240.1	346.0	459.6	727.8	1,498.2
1938	122.6	1,485.1	142.8	25.8	49.5	67.2	97.3	161.4	241.2	307.5	595.3	891.4
1939	115.4	1,336.6	124.5	19.7	43.1	62.8	87.9	162.1	249.3	310.7	584.2	993.3

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Notes: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total, separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes.....	1,271	1,943	1,950
Male	683	1,053	1,068
Female	588	890	882

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ICD Revision	Years in use	ICD Nos.
Fifth	1939	117
Fourth	1930-1938	117
Third	1921-1929	111
Second	1910-1920	102
First	1900-1909	103

ALL RACES, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	2.7	*	*	*	1.0	2.5	2.8	4.1	7.0	13.8	16.7	*
1901	2.8	*	*	*	1.3	1.6	2.9	4.3	8.5	12.9	22.9	*
1902	3.0	*	*	*	1.2	2.3	3.8	5.1	7.7	13.0	19.6	*
1903	2.7	*	*	*	1.0	2.0	2.9	4.4	7.7	11.6	20.7	*
1904	3.0	*	*	*	1.3	2.1	3.9	4.7	9.1	13.5	16.1	*
1905	3.0	*	*	*	1.1	2.0	4.0	5.0	7.2	13.9	23.1	*
1906	3.3	*	*	*	1.1	2.4	4.1	5.9	10.1	14.8	18.7	*
1907	3.2	*	*	*	1.1	2.3	3.8	6.5	9.5	15.7	18.9	*
1908	3.2	*	*	*	1.0	2.4	4.0	6.3	9.5	13.3	19.9	27.8
1909	3.4	*	*	*	1.1	2.4	3.8	6.3	10.3	14.9	25.1	22.6
1910	4.0	*	*	0.3	1.5	2.7	4.8	7.9	12.1	18.0	23.4	27.1
1911	3.5	*	*	*	1.1	2.5	4.6	6.1	10.6	17.2	21.8	25.0
1912	3.8	*	*	0.2	1.3	2.7	4.7	7.7	10.5	16.0	26.9	26.2
1913	3.8	*	*	0.2	1.1	2.8	5.2	6.7	11.8	17.5	24.2	26.3
1914	4.0	*	0.4	0.2	1.2	3.3	5.1	7.3	11.7	17.1	25.4	23.2
1915	4.2	*	*	0.2	1.2	2.9	5.6	8.1	12.4	18.3	27.8	23.5
1916	4.4	1.5	*	0.2	1.4	2.9	6.1	8.7	13.2	18.6	26.5	38.6
1917	4.3	*	*	0.2	1.3	3.1	6.1	8.0	12.8	19.3	25.6	34.6
1918	3.9	1.3	*	0.2	1.1	2.8	5.1	7.3	11.8	16.0	22.2	27.9
1919	3.5	*	*	*	0.8	2.2	4.7	7.3	10.5	15.4	21.3	24.4
1920	3.6	*	*	0.1	0.8	2.4	4.5	7.2	11.0	16.2	25.3	33.1
1921	4.8	1.1	*	0.2	1.2	3.5	6.8	10.1	14.1	19.5	25.5	33.2
1922	5.0	*	*	0.2	1.2	3.6	6.8	10.5	14.6	22.4	28.9	34.5
1923	5.2	1.0	*	0.2	1.3	3.7	6.7	11.9	15.4	20.2	27.6	37.0
1924	5.5	1.5	*	0.1	1.3	3.5	7.9	11.4	16.5	21.8	31.8	33.6
1925	5.9	1.0	*	*	1.3	4.0	8.3	12.6	16.9	24.1	33.3	36.0
1926	6.0	*	*	0.1	1.3	4.2	8.3	13.0	17.4	24.7	32.6	40.5
1927	6.3	*	*	0.1	1.2	3.9	9.0	14.0	19.0	24.0	32.1	37.3
1928	6.4	*	*	*	1.3	4.2	8.9	14.4	18.0	24.9	31.8	45.2
1929	6.4	*	*	0.1	1.3	4.3	8.8	14.1	18.3	24.2	32.4	36.1
1930	6.2	*	*	0.1	1.1	3.8	8.4	13.6	18.9	23.4	30.8	28.8
1931	6.1	*	*	0.1	1.1	3.7	7.7	13.4	17.5	24.1	31.0	31.3
1932	6.0	*	*	0.1	1.0	3.5	7.2	13.1	17.8	23.4	32.0	32.5
1933	6.0	*	*	0.1	1.0	3.1	7.4	13.4	18.3	22.5	30.9	34.4
1934	6.1	1.1	*	*	1.0	3.2	7.4	13.4	18.6	22.8	28.9	27.5
1935	6.6	*	*	0.1	0.9	3.5	7.8	14.8	20.2	23.7	33.0	32.7
1936	6.7	1.4	*	0.1	1.0	3.2	8.3	14.5	19.3	24.6	31.7	39.3
1937	6.8	*	*	0.1	0.9	3.3	7.8	14.6	20.9	23.9	33.6	32.5
1938	6.5	1.1	*	*	0.8	3.0	7.1	13.4	19.7	23.7	32.8	34.3
1939	6.8	1.0	*	*	0.8	2.7	7.0	14.1	21.1	25.3	34.8	35.2

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ALL RACES, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	2.7	*	*	*	*	1.9	2.3	3.8	8.1	16.6	23.8	*
1901	2.7	*	*	*	*	*	3.0	4.2	9.4	14.7	22.8	*
1902	2.9	*	*	*	*	1.9	3.3	5.8	8.5	15.5	21.0	*
1903	2.8	*	*	*	*	1.5	2.7	4.7	9.0	15.0	20.0	*
1904	3.0	*	*	*	*	1.7	3.6	5.1	10.5	16.0	15.9	*
1905	3.2	*	*	*	*	2.0	4.1	6.6	8.3	14.1	29.8	*
1906	3.3	*	*	*	*	2.1	4.3	6.0	11.7	15.8	19.9	*
1907	3.5	*	*	*	0.7	2.2	4.4	6.8	11.4	19.0	22.5	*
1908	3.5	*	*	*	0.7	2.3	4.2	7.3	11.6	16.1	25.2	*
1909	3.8	*	*	*	0.8	2.5	4.3	7.4	12.7	17.1	27.8	*
1910	4.5	*	*	*	1.2	2.9	5.7	9.7	14.9	18.7	25.2	*
1911	3.9	*	*	*	0.7	2.3	5.2	7.5	13.9	19.1	23.6	*
1912	4.4	*	*	*	1.2	2.8	5.7	9.9	12.7	18.6	30.5	*
1913	4.4	*	*	*	0.9	3.2	5.9	8.1	14.4	19.0	27.3	*
1914	4.5	*	*	*	1.1	3.5	6.1	8.6	14.6	18.0	27.9	*
1915	5.1	*	*	*	1.2	3.6	7.0	10.2	15.7	21.1	28.6	*
1916	5.4	*	*	*	1.5	3.5	7.6	10.8	17.0	21.4	28.5	47.3
1917	5.4	*	*	*	1.3	3.6	8.1	10.1	15.8	24.4	28.4	33.8
1918	5.1	*	*	*	1.1	3.9	6.7	9.4	15.5	19.4	29.1	38.1
1919	4.3	*	*	*	0.9	2.5	6.1	9.2	12.8	18.8	24.3	28.5
1920	4.4	*	*	*	0.8	2.9	5.5	9.3	13.6	19.8	26.2	29.7
1921	6.5	*	*	*	1.3	4.5	9.7	13.8	19.4	24.5	30.2	50.0
1922	6.7	*	*	*	1.4	5.1	9.6	14.3	19.8	28.8	31.6	39.4
1923	7.2	*	*	*	1.5	5.2	10.0	16.8	21.2	26.7	35.2	37.6
1924	7.8	2.0	*	*	1.6	5.4	11.9	16.3	23.3	29.5	39.6	30.8
1925	8.5	*	*	*	1.9	6.0	12.8	18.7	23.6	32.3	42.6	35.5
1926	8.7	*	*	*	1.9	6.2	12.9	19.0	24.6	33.0	41.2	50.1
1927	9.2	*	*	*	1.7	6.0	14.0	20.5	27.7	33.1	37.4	48.3
1928	9.4	*	*	*	1.6	6.8	14.0	21.6	26.1	35.0	37.4	47.3
1929	9.6	*	*	*	1.8	7.1	14.0	21.5	27.2	33.8	43.7	41.9
1930	9.4	*	*	*	1.7	5.8	13.6	21.1	28.5	32.4	40.4	24.7
1931	9.2	*	*	*	1.5	5.9	12.4	21.1	25.9	34.1	40.8	42.9
1932	9.3	*	*	0.2	1.5	5.6	11.6	20.8	27.5	34.5	42.6	37.5
1933	9.3	*	*	*	1.4	5.1	12.2	21.2	28.3	31.9	41.7	37.3
1934	9.5	*	*	*	1.5	5.2	12.2	21.2	29.3	33.0	38.9	35.5
1935	10.5	*	*	0.2	1.5	5.7	13.2	23.9	31.8	34.9	44.0	43.1
1936	10.7	*	*	*	1.6	5.4	13.9	23.5	30.8	36.8	44.8	46.8
1937	11.0	*	*	*	1.4	5.6	13.1	24.2	34.2	35.9	48.6	42.6
1938	10.5	*	*	*	1.4	4.9	12.0	22.2	32.2	36.2	48.8	43.3
1939	10.9	*	*	*	1.2	4.6	11.7	23.7	34.3	38.9	50.3	44.4

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ALL RACES, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	2.7	*	*	*	1.3	3.1	3.4	4.3	6.0	11.1	*	*
1901	2.9	*	*	*	1.8	2.3	2.9	4.4	7.7	11.2	22.9	*
1902	3.0	*	*	*	1.5	2.8	4.3	4.2	6.9	10.7	18.3	*
1903	2.6	*	*	*	1.1	2.5	3.1	4.0	6.4	8.4	21.4	*
1904	3.0	*	*	*	1.8	2.5	4.3	4.2	7.8	11.0	16.3	*
1905	2.7	*	*	*	1.3	1.9	3.9	3.3	6.2	13.8	17.2	*
1906	3.2	*	*	*	1.7	2.6	3.9	5.8	8.5	13.7	17.6	*
1907	3.0	*	*	*	1.4	2.4	3.1	6.2	7.5	12.5	15.8	*
1908	2.8	*	*	*	1.3	2.4	3.8	5.2	7.2	10.5	15.0	*
1909	3.0	*	*	*	1.4	2.3	3.3	5.1	7.8	12.8	22.6	*
1910	3.5	*	*	*	1.9	2.4	3.9	5.9	9.1	17.3	21.8	*
1911	3.1	*	*	*	1.6	2.7	3.9	4.6	7.1	15.3	20.3	*
1912	3.1	*	*	*	1.3	2.6	3.6	5.2	8.2	13.4	23.7	*
1913	3.3	*	*	*	1.3	2.5	4.4	5.1	8.8	15.9	21.2	*
1914	3.4	*	*	*	1.3	3.1	4.0	5.9	8.5	16.2	23.1	*
1915	3.3	*	*	*	1.2	2.2	4.2	5.8	8.9	15.6	27.1	*
1916	3.4	*	*	*	1.3	2.3	4.5	6.4	9.0	15.8	24.7	32.1
1917	3.3	*	*	*	1.2	2.5	3.9	5.7	9.6	14.2	23.0	35.3
1918	2.7	*	*	*	1.1	1.8	3.4	4.9	7.7	12.6	16.0	*
1919	2.6	*	*	*	0.8	1.8	3.3	5.1	8.0	11.9	18.6	21.3
1920	2.8	*	*	*	0.8	1.9	3.4	4.7	8.1	12.5	24.4	35.6
1921	3.1	*	*	*	1.1	2.5	3.6	6.0	8.3	14.4	21.2	20.5
1922	3.3	*	*	*	0.9	2.2	3.7	6.3	9.0	15.8	26.4	30.9
1923	3.1	*	*	*	1.1	2.1	3.2	6.5	9.1	13.4	20.6	36.5
1924	3.1	*	*	*	0.9	1.7	3.6	6.1	9.1	13.7	24.5	35.8
1925	3.2	*	*	*	0.8	2.0	3.5	6.0	9.6	15.7	24.7	36.3
1926	3.3	*	*	*	0.6	2.1	3.4	6.5	9.7	16.1	24.6	33.2
1927	3.3	*	*	*	0.7	1.8	3.7	7.0	9.5	14.5	27.2	28.9
1928	3.3	*	*	*	0.9	1.7	3.5	6.7	9.2	14.5	26.5	43.7
1929	3.0	*	*	*	0.7	1.6	3.4	6.1	8.6	14.3	21.7	31.8
1930	3.0	*	*	*	0.6	1.9	3.0	5.5	8.6	14.0	21.8	31.8
1931	2.9	*	*	*	0.7	1.6	2.8	5.2	8.4	13.7	21.8	22.6
1932	2.6	*	*	*	0.5	1.4	2.5	4.8	7.5	12.0	22.1	28.8
1933	2.7	*	*	*	0.6	1.2	2.4	5.0	7.5	12.8	20.7	32.2
1934	2.6	*	*	*	0.4	1.1	2.5	4.9	7.1	12.4	19.6	21.6
1935	2.7	*	*	*	0.4	1.2	2.1	4.9	7.9	12.3	22.7	24.8
1936	2.7	*	*	*	0.4	1.0	2.5	4.8	7.1	12.4	19.6	33.6
1937	2.5	*	*	*	0.4	1.2	2.3	4.3	6.7	11.8	19.8	24.8
1938	2.4	*	*	*	0.3	1.1	2.2	4.0	6.5	11.2	18.1	27.5
1939	2.6	*	*	*	0.3	0.9	2.3	3.9	7.2	11.7	20.7	24.7

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

WHITE, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	4.0	---	---	---	---	---	---	---	---	---	---	---
1911	3.5	---	---	---	---	---	---	---	---	---	---	---
1912	3.7	---	---	---	---	---	---	---	---	---	---	---
1913	3.8	---	---	---	---	---	---	---	---	---	---	---
1914	4.0	*	0.4	*	1.2	3.3	5.0	7.4	11.7	17.2	25.6	24.4
1915	4.2	*	*	0.2	1.2	2.9	5.5	8.0	12.4	18.3	28.4	23.0
1916	4.4	1.6	*	0.2	1.3	2.8	6.0	8.6	13.2	18.7	27.2	40.6
1917	4.3	---	---	---	---	---	---	---	---	---	---	---
1918	3.9	1.4	*	0.2	1.1	2.7	5.1	7.3	11.9	16.2	22.6	26.3
1919	3.5	*	*	*	0.8	2.1	4.6	7.0	10.4	15.4	21.5	24.3
1920	3.6	*	*	0.2	0.7	2.3	4.3	7.0	10.9	16.3	25.4	35.0
1921	4.8	1.2	*	0.1	1.1	3.4	6.6	10.0	14.0	19.4	26.3	34.4
1922	5.1	*	*	0.2	1.1	3.5	6.7	10.4	14.7	22.4	29.3	35.1
1923	5.2	*	*	0.1	1.2	3.5	6.7	11.8	15.3	20.8	28.4	37.2
1924	5.6	1.5	*	0.1	1.2	3.6	7.8	11.3	16.7	22.2	32.0	34.1
1925	6.0	1.0	*	*	1.3	3.9	8.2	12.5	17.1	24.1	34.0	37.3
1926	6.1	*	*	0.1	1.2	4.0	8.2	13.0	17.5	25.1	33.5	43.5
1927	6.3	*	*	0.1	1.1	3.6	8.7	13.8	19.2	24.2	33.1	38.0
1928	6.5	*	*	*	1.1	4.1	8.9	14.3	18.1	25.4	32.6	47.5
1929	6.4	*	*	*	1.1	4.0	8.6	14.0	18.5	24.5	33.6	39.3
1930	6.2	*	*	0.1	1.0	3.6	8.2	13.1	18.8	23.7	31.6	30.7
1931	6.1	*	*	0.1	0.9	3.5	7.6	13.2	17.4	24.4	31.3	32.3
1932	6.0	*	*	0.1	0.9	3.2	6.9	12.8	17.6	23.7	32.9	34.5
1933	6.0	*	*	0.1	0.8	2.9	7.0	13.3	18.0	22.8	31.5	36.4
1934	6.1	*	*	*	0.8	2.9	7.1	13.0	18.8	23.3	29.5	29.2
1935	6.7	*	*	0.1	0.9	3.2	7.4	14.6	20.2	24.2	33.9	34.9
1936	6.8	1.4	*	0.1	1.0	3.0	8.0	14.3	19.4	25.0	32.7	42.0
1937	6.9	*	*	0.1	0.8	3.1	7.6	14.5	20.8	24.2	33.9	34.8
1938	6.5	1.1	*	*	0.7	2.7	6.7	13.0	19.7	24.1	34.2	37.2
1939	6.8	*	*	*	0.7	2.4	6.7	13.8	21.1	25.7	36.0	34.1

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

WHITE, MALE												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	4.5	---	---	---	---	---	---	---	---	---	---	---
1911	3.9	---	---	---	---	---	---	---	---	---	---	---
1912	4.4	---	---	---	---	---	---	---	---	---	---	---
1913	4.4	---	---	---	---	---	---	---	---	---	---	---
1914	4.5	*	*	*	1.1	3.4	5.9	8.7	14.8	18.0	27.9	*
1915	5.1	*	*	*	1.2	3.7	6.8	10.1	15.8	21.0	29.2	*
1916	5.4	*	*	*	1.3	3.5	7.5	10.8	17.3	21.6	29.2	48.7
1917	5.4	---	---	---	---	---	---	---	---	---	---	---
1918	5.2	*	*	*	1.1	4.0	6.7	9.5	15.8	19.6	29.7	36.6
1919	4.4	*	*	*	0.9	2.5	6.0	9.0	12.9	19.0	24.5	*
1920	4.4	*	*	*	0.7	2.8	5.4	9.3	13.5	20.3	26.7	31.1
1921	6.6	*	*	*	1.3	4.4	9.6	13.9	19.6	24.6	31.0	53.1
1922	6.9	*	*	*	1.4	5.1	9.6	14.5	20.4	29.0	33.1	39.0
1923	7.4	*	*	*	1.5	5.2	10.0	16.9	21.7	27.6	36.0	38.4
1924	8.0	2.2	*	*	1.5	5.5	12.1	16.3	23.8	30.5	40.6	31.0
1925	8.7	*	*	*	1.9	5.9	12.9	18.7	24.3	32.8	43.5	34.9
1926	9.0	*	*	*	1.9	6.2	12.9	19.3	25.1	33.8	42.4	52.2
1927	9.4	*	*	*	1.7	5.7	13.8	20.7	28.3	33.9	38.8	48.2
1928	9.7	*	*	*	1.6	6.8	14.2	21.8	26.8	36.1	38.6	49.1
1929	9.9	*	*	*	1.6	6.8	14.0	21.6	27.9	34.3	45.7	45.5
1930	9.5	*	*	*	1.6	5.6	13.4	21.0	28.9	33.1	41.3	27.5
1931	9.3	*	*	*	1.3	5.7	12.3	21.0	26.2	34.9	41.1	44.8
1932	9.4	*	*	*	1.3	5.2	11.5	20.5	27.4	35.2	43.7	40.7
1933	9.5	*	*	*	1.2	4.9	11.9	21.5	28.5	32.8	42.4	39.8
1934	9.7	*	*	*	1.4	5.0	11.9	21.1	30.1	34.0	39.6	38.6
1935	10.7	*	*	*	1.4	5.4	12.8	24.1	32.2	36.1	45.7	45.5
1936	11.0	*	*	*	1.6	5.2	13.8	23.7	31.4	37.8	46.3	49.6
1937	11.2	*	*	*	1.2	5.2	13.1	24.2	34.8	36.6	49.3	44.3
1938	10.6	*	*	*	1.3	4.5	11.5	22.1	32.6	37.2	50.9	48.0
1939	11.2	*	*	*	1.1	4.3	11.6	23.6	34.9	39.9	52.2	46.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

WHITE, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	3.5	---	---	---	---	---	---	---	---	---	---	---
1911	3.0	---	---	---	---	---	---	---	---	---	---	---
1912	3.0	---	---	---	---	---	---	---	---	---	---	---
1913	3.2	---	---	---	---	---	---	---	---	---	---	---
1914	3.4	*	*	*	1.3	3.1	3.9	5.8	8.3	16.4	23.5	*
1915	3.3	*	*	*	1.1	2.0	4.1	5.7	8.7	15.6	27.6	*
1916	3.4	*	*	*	1.3	2.1	4.3	6.0	8.8	15.7	25.4	34.4
1917	3.2	---	---	---	---	---	---	---	---	---	---	---
1918	2.6	*	*	*	1.0	1.6	3.2	4.8	7.6	12.7	16.1	*
1919	2.6	*	*	*	0.7	1.7	3.1	4.7	7.8	11.8	18.9	*
1920	2.7	*	*	*	0.7	1.9	3.1	4.4	8.0	12.2	24.2	38.0
1921	3.0	*	*	*	1.0	2.3	3.4	5.8	8.0	14.1	21.9	*
1922	3.2	*	*	*	0.8	1.9	3.5	6.0	8.6	15.6	25.8	32.1
1923	3.0	*	*	*	1.0	1.8	3.1	6.2	8.5	13.8	21.5	36.2
1924	3.0	*	*	*	0.8	1.7	3.1	5.9	9.2	13.7	24.1	36.6
1925	3.2	*	*	*	0.7	1.8	3.2	5.8	9.4	15.2	25.3	39.2
1926	3.2	*	*	*	0.6	1.8	3.0	6.2	9.3	16.2	25.3	36.7
1927	3.2	*	*	*	0.6	1.6	3.1	6.4	9.4	14.3	27.8	30.1
1928	3.1	*	*	*	0.7	1.4	3.2	6.1	8.6	14.5	27.1	46.2
1929	2.9	*	*	*	0.7	1.2	2.8	5.8	8.3	14.4	22.3	34.6
1930	2.8	*	*	*	0.4	1.6	2.7	4.6	8.1	13.9	22.5	33.1
1931	2.8	*	*	*	0.6	1.4	2.6	4.8	8.0	13.7	22.1	22.9
1932	2.5	*	*	*	0.4	1.2	2.0	4.4	7.2	11.8	22.9	29.8
1933	2.5	*	*	*	0.4	0.8	2.0	4.5	6.9	12.6	21.2	33.9
1934	2.5	*	*	*	0.3	0.9	2.1	4.3	6.9	12.4	20.0	22.1
1935	2.6	*	*	*	0.4	1.0	1.8	4.4	7.4	12.1	23.0	27.0
1936	2.5	*	*	*	0.3	0.8	2.0	4.3	6.8	12.1	20.0	36.2
1937	2.5	*	*	*	0.4	1.0	2.0	4.1	6.0	11.7	19.5	27.6
1938	2.3	*	*	*	0.2	0.8	1.7	3.4	6.2	11.0	18.8	29.0
1939	2.4	*	*	*	0.3	0.5	1.8	3.4	6.6	11.7	21.2	24.8

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ALL OTHER, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	4.8	---	---	---	---	---	---	---	---	---	---	---
1911	4.6	---	---	---	---	---	---	---	---	---	---	---
1912	5.0	---	---	---	---	---	---	---	---	---	---	---
1913	4.7	---	---	---	---	---	---	---	---	---	---	---
1914	3.9	*	*	*	*	4.5	7.1	*	*	*	*	*
1915	4.9	*	*	*	*	4.2	8.4	10.8	*	*	*	*
1916	4.5	*	*	*	2.4	4.5	8.0	11.6	13.1	*	*	*
1917	4.6	---	---	---	---	---	---	---	---	---	---	---
1918	3.6	*	*	*	1.8	3.8	6.3	7.6	10.3	*	*	*
1919	3.7	*	*	*	*	2.9	6.3	10.5	11.8	14.9	*	*
1920	3.8	*	*	*	*	3.4	6.6	9.4	13.3	14.4	*	*
1921	4.7	*	*	*	2.2	4.6	8.2	11.2	15.9	21.4	*	*
1922	4.5	*	*	*	1.7	4.6	8.1	11.5	13.1	22.3	*	*
1923	4.7	*	*	*	1.9	5.1	7.6	13.4	16.3	10.2	*	*
1924	4.5	*	*	*	2.0	2.8	9.4	12.7	12.8	14.0	27.8	*
1925	5.0	*	*	*	1.7	5.2	8.8	14.0	13.5	24.2	*	*
1926	5.2	*	*	*	1.7	5.7	9.9	13.6	16.1	17.3	*	*
1927	6.0	*	*	*	1.6	6.7	12.6	16.0	15.6	19.3	*	*
1928	5.5	*	*	*	2.3	5.3	9.5	16.5	17.3	16.1	*	*
1929	5.8	*	*	*	2.1	7.0	11.2	14.8	15.4	19.1	*	*
1930	6.3	*	*	*	2.2	6.0	11.0	18.6	20.3	19.3	*	*
1931	5.8	*	*	*	2.2	5.6	8.7	16.2	19.3	18.4	25.6	*
1932	5.9	*	*	*	2.0	5.8	9.6	16.3	21.7	19.4	*	*
1933	5.7	*	*	*	1.9	5.3	10.5	14.5	21.9	17.4	21.0	*
1934	5.7	*	*	*	2.3	5.2	10.6	17.0	15.3	15.3	19.6	*
1935	6.1	*	*	*	1.5	5.8	10.8	16.5	21.1	17.5	*	*
1936	5.9	*	*	*	1.5	4.7	11.2	16.2	18.0	20.2	*	*
1937	6.2	*	*	*	1.8	5.4	9.2	15.8	22.1	20.0	29.5	*
1938	6.2	*	*	*	1.6	5.5	10.9	17.1	19.9	18.8	*	*
1939	6.3	*	*	*	1.5	5.5	9.5	17.7	21.1	19.9	*	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ALL OTHER, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	4.8	---	---	---	---	---	---	---	---	---	---	---
1911	3.6	---	---	---	---	---	---	---	---	---	---	---
1912	5.8	---	---	---	---	---	---	---	---	---	---	---
1913	4.6	---	---	---	---	---	---	---	---	---	---	---
1914	4.2	*	*	*	*	*	*	*	*	*	*	*
1915	5.4	*	*	*	*	*	10.3	*	*	*	*	*
1916	4.7	*	*	*	*	*	8.2	9.8	*	*	*	*
1917	4.9	---	---	---	---	---	---	---	---	---	---	---
1918	3.7	*	*	*	*	*	6.5	8.6	*	*	*	*
1919	3.9	*	*	*	*	*	7.1	10.4	*	*	*	*
1920	4.3	*	*	*	*	4.6	7.1	9.9	15.6	*	*	*
1921	5.6	*	*	*	*	5.0	10.5	12.7	17.8	23.5	*	*
1922	4.9	*	*	*	*	4.8	9.7	12.3	11.7	25.5	*	*
1923	5.7	*	*	*	*	5.7	10.7	16.0	15.2	*	*	*
1924	5.2	*	*	*	*	3.6	9.8	15.5	15.9	*	*	*
1925	6.5	*	*	*	*	7.2	11.5	18.3	14.0	23.1	*	*
1926	6.3	*	*	*	2.5	6.4	12.3	15.5	16.8	19.1	*	*
1927	7.6	*	*	*	2.2	9.2	15.6	18.4	19.3	19.9	*	*
1928	6.6	*	*	*	2.4	6.7	12.4	19.4	16.8	16.8	*	*
1929	7.7	*	*	*	3.1	9.5	13.3	20.0	17.7	25.1	*	*
1930	8.0	*	*	*	2.4	7.3	16.3	21.6	23.0	22.3	*	*
1931	8.1	*	*	*	2.9	8.1	13.0	22.2	22.8	22.0	*	*
1932	8.3	*	*	*	2.6	9.0	12.5	23.0	29.0	22.4	*	*
1933	7.3	*	*	*	2.3	6.3	14.7	18.3	26.1	19.3	*	*
1934	7.7	*	*	*	2.9	7.0	15.1	21.9	19.8	17.7	*	*
1935	8.6	*	*	*	2.1	8.4	16.8	22.2	25.8	18.9	*	*
1936	8.0	*	*	*	1.7	7.1	15.1	21.4	23.2	23.4	*	*
1937	9.1	*	*	*	3.0	8.8	13.3	24.4	25.8	26.4	35.7	*
1938	8.9	*	*	*	2.2	8.4	15.7	23.2	27.5	23.5	*	*
1939	8.6	*	*	*	2.3	7.3	12.4	24.7	25.8	26.7	*	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ULCER OF STOMACH AND DUODENUM

ALL OTHER, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	4.8	...	...	...	...	...	...	...	...	...	...	...
1911	5.7	...	...	...	...	...	...	...	...	...	...	...
1912	4.2	...	...	...	...	...	...	...	...	...	...	...
1913	4.7	...	...	...	...	...	...	...	...	...	...	...
1914	3.5	*	*	*	*	*	*	*	*	*	*	*
1915	4.4	*	*	*	*	*	*	*	*	*	*	*
1916	4.3	*	*	*	*	*	*	*	*	*	*	*
1917	4.2	...	...	...	...	...	7.7	14.1	*	*	*	*
1918	3.5	*	*	*	*	4.1	6.1	*	*	*	*	*
1919	3.4	*	*	*	*	3.4	5.4	10.6	*	*	*	*
1920	3.3	*	*	*	*	*	6.1	8.7	*	*	*	*
1921	3.9	*	*	*	*	4.3	5.8	9.2	*	*	*	*
1922	4.1	*	*	*	*	4.4	6.4	10.5	14.8	*	*	*
1923	3.6	*	*	*	*	4.5	4.4	10.2	17.7	*	*	*
1924	3.8	*	*	*	2.2	*	9.0	9.1	*	*	*	*
1925	3.5	*	*	*	*	3.4	6.0	8.7	12.8	25.5	*	*
1926	4.0	*	*	*	*	5.1	7.4	11.3	15.1	*	*	*
1927	4.4	*	*	*	*	4.3	9.6	13.2	10.7	*	*	*
1928	4.4	*	*	*	2.2	4.0	6.6	13.1	17.9	*	*	*
1929	4.0	*	*	*	*	4.7	9.1	8.8	12.3	*	*	*
1930	4.5	*	*	*	2.0	4.8	5.7	15.0	16.8	*	*	*
1931	3.5	*	*	*	*	3.3	4.5	9.3	14.7	*	*	*
1932	3.5	*	*	*	*	2.8	6.8	8.6	12.3	15.9	*	*
1933	4.1	*	*	*	1.5	4.4	6.4	10.2	16.6	15.4	*	*
1934	3.7	*	*	*	1.7	3.4	6.2	11.6	9.8	12.6	*	*
1935	3.7	*	*	*	*	3.3	5.0	10.2	15.3	16.1	*	*
1936	3.8	*	*	*	*	2.5	7.4	10.5	11.6	16.8	*	*
1937	3.3	*	*	*	*	2.3	5.3	6.4	17.7	13.1	*	*
1938	3.6	*	*	*	*	3.0	6.4	10.4	10.9	13.8	*	*
1939	4.1	*	*	*	*	3.8	6.7	10.3	15.7	12.4	*	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population; but by age, are included with deaths for the nonwhite population. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes.....	12	21	20
Male.....	7	17	14
Female.....	5	4	6

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ICD Revision	Years in use	ICD Nos.
Fifth	1939-119, 120	
Fourth	1930-1938 119, 120	
Third	1921-1929 15, 113, 114	
Second	1910-1920 13, 104, 105	
First	1900-1909 13, 105, 106	

ALL RACES, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	142.7	4,429.8	303.0	8.8	5.7	7.5	10.5	18.6	49.3	130.4	316.9	676.1
1901	118.5	3,572.4	261.4	8.8	4.7	6.6	10.0	17.2	49.7	116.3	270.5	558.4
1902	104.9	3,167.6	242.6	6.9	3.5	5.4	8.3	15.0	42.2	97.6	242.2	425.5
1903	100.3	3,113.2	210.4	6.4	3.1	4.8	7.6	14.1	35.1	94.4	245.1	551.8
1904	111.5	3,537.8	236.0	7.8	3.3	4.3	7.0	14.2	34.5	97.0	242.1	553.2
1905	118.4	3,796.9	261.9	6.9	3.3	4.4	6.9	12.4	34.3	88.4	233.3	617.5
1906	123.6	3,883.8	274.8	6.0	2.7	4.6	6.1	13.0	32.7	94.7	287.3	692.4
1907	115.0	3,693.8	251.8	5.8	2.5	3.6	6.1	11.3	31.2	78.6	234.6	536.4
1908	112.5	3,630.1	252.2	5.8	2.2	3.9	6.0	9.8	26.3	74.7	223.9	586.2
1909	101.8	3,254.1	243.0	5.6	1.8	3.1	5.1	10.3	24.3	68.7	215.5	546.2
1910	115.4	3,778.8	271.8	6.1	1.9	3.2	5.0	9.3	23.9	74.3	230.7	584.5
1911	86.8	2,782.9	197.4	4.3	2.2	3.2	5.5	9.0	21.0	63.1	181.3	451.6
1912	79.6	2,515.6	191.3	4.1	1.9	3.1	4.8	8.7	21.2	56.1	172.6	409.9
1913	86.7	2,690.5	221.7	5.2	2.0	3.3	5.5	9.7	20.1	58.5	175.7	449.4
1914	75.1	2,344.2	186.5	5.1	2.0	3.5	5.3	8.4	18.2	51.2	156.5	385.5
1915	67.5	2,134.4	168.2	4.5	1.7	2.8	4.9	7.5	16.9	45.9	128.9	330.0
1916	75.5	2,205.1	213.2	4.9	2.0	3.2	5.7	7.9	16.8	44.6	133.2	340.4
1917	75.2	2,191.9	217.8	6.0	2.2	3.7	5.9	9.5	19.4	49.3	139.1	340.9
1918	72.2	2,113.0	208.6	5.7	2.2	3.7	5.1	8.8	16.0	43.7	133.8	327.3
1919	55.2	1,683.2	147.3	4.6	1.8	3.1	4.5	7.5	13.2	37.6	106.9	290.9
1920	53.7	1,625.9	141.3	4.1	1.7	3.0	4.3	7.4	12.7	32.7	107.5	258.3
1921	50.7	1,466.5	146.1	4.5	1.4	2.0	2.8	4.6	10.1	29.6	115.7	302.9
1922	38.9	1,169.7	105.0	2.9	1.2	1.7	2.6	4.5	8.5	22.9	77.0	222.0
1923	39.1	1,180.3	118.3	3.0	1.3	1.6	2.2	4.1	7.4	22.6	74.3	220.4
1924	33.7	1,009.5	100.9	2.7	1.2	1.6	2.3	3.7	7.6	21.7	71.0	193.3
1925	38.6	1,176.4	115.4	3.1	1.2	1.6	2.5	4.1	8.1	22.6	79.0	249.2
1926	32.9	1,039.0	100.1	2.7	1.1	1.5	2.2	4.4	6.9	20.4	64.2	195.8
1927	27.1	835.7	85.8	2.7	1.0	1.5	2.3	3.6	6.4	16.0	58.2	164.2
1928	26.4	821.6	92.9	2.5	1.1	1.4	2.1	3.7	6.2	17.6	57.5	166.4
1929	23.3	754.9	80.8	2.4	1.1	1.4	2.3	3.6	6.2	14.8	49.3	131.7
1930	26.0	836.9	95.6	3.0	1.1	1.5	2.2	3.6	7.2	16.7	59.5	173.5
1931	20.5	693.7	70.6	2.4	1.0	1.3	1.9	3.6	6.1	13.6	46.9	134.0
1932	16.1	554.1	56.8	1.8	0.9	1.2	1.8	3.2	5.4	11.6	41.4	110.9
1933	17.3	590.5	66.9	1.9	1.0	1.5	1.8	3.4	6.2	13.2	42.2	130.9
1934	18.4	675.3	65.9	2.1	1.0	1.5	2.1	3.5	6.0	13.5	44.6	134.8
1935	14.1	539.6	43.6	1.4	1.0	1.5	1.8	3.0	5.0	11.2	32.8	101.8
1936	16.4	629.9	57.9	1.6	0.9	1.4	1.7	3.1	5.2	10.2	40.7	111.4
1937	14.7	596.3	47.2	1.3	0.9	1.2	1.9	2.6	4.3	9.3	30.1	86.8
1938	14.3	567.8	45.2	1.3	0.9	1.2	1.6	2.4	4.5	8.8	30.2	84.3
1939	11.6	448.5	35.3	1.0	0.8	1.1	1.6	2.4	3.5	8.0	25.1	88.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ALL RACES, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	150.5	4,748.6	317.4	8.7	5.8	6.8	9.7	17.6	49.1	115.7	313.9	651.4
1901	124.9	3,850.7	272.4	9.2	4.4	5.4	9.3	15.4	46.9	113.2	258.3	470.8
1902	110.3	3,407.4	253.6	7.1	3.4	4.0	7.0	12.2	40.0	90.0	236.5	446.2
1903	106.1	3,368.5	225.9	6.3	3.0	3.9	7.3	12.4	31.8	86.8	232.8	475.4
1904	117.2	3,815.2	234.4	8.0	3.1	3.5	6.3	12.7	35.0	95.7	238.6	576.0
1905	126.3	4,116.5	284.8	7.4	3.3	4.3	6.3	10.8	33.4	81.8	210.8	564.7
1906	129.6	4,226.5	285.5	5.3	2.7	4.4	5.8	12.3	27.3	82.4	276.5	693.6
1907	120.7	4,012.0	262.0	5.8	2.3	2.9	5.9	10.6	30.8	69.2	207.2	521.0
1908	116.9	3,920.6	265.4	5.7	1.6	3.3	5.5	9.8	22.2	64.6	196.1	507.4
1909	106.3	3,524.1	254.2	5.6	1.5	2.7	4.4	10.0	22.3	66.4	201.7	505.6
1910	120.7	4,120.3	288.2	6.1	1.6	2.6	4.3	8.0	21.1	61.6	213.9	529.0
1911	90.6	3,037.7	204.4	4.2	1.6	3.5	5.0	8.6	20.2	55.1	160.8	402.2
1912	83.0	2,736.4	200.0	4.0	1.8	3.2	5.0	8.6	18.9	48.9	150.5	390.0
1913	91.1	2,936.8	231.7	5.7	2.0	3.2	5.7	9.7	19.0	54.3	156.8	406.8
1914	79.6	2,577.2	199.8	5.3	1.6	3.4	5.8	8.6	16.8	46.0	139.4	355.3
1915	71.5	2,349.1	178.9	4.7	1.7	2.9	5.2	7.7	15.8	40.0	109.8	304.3
1916	80.2	2,498.5	221.3	5.1	2.0	3.4	6.4	8.1	16.6	43.3	114.6	317.3
1917	80.8	2,440.3	231.7	5.9	2.2	3.5	6.1	10.3	19.4	44.7	125.9	330.1
1918	78.2	2,331.7	219.9	6.2	2.3	4.0	5.4	8.6	15.9	38.3	116.7	286.5
1919	59.4	1,860.2	158.2	4.9	1.6	3.2	4.9	7.0	13.3	34.0	93.9	267.3
1920	58.0	1,813.3	151.4	4.2	1.6	2.7	4.7	7.5	12.1	30.3	102.1	221.5
1921	54.0	1,620.6	153.5	4.8	1.2	1.4	2.5	4.0	8.8	24.9	100.5	286.6
1922	42.1	1,299.7	113.1	3.0	1.3	1.4	2.4	4.3	7.4	20.1	70.7	196.9
1923	41.9	1,307.8	125.2	3.0	1.1	1.4	2.2	4.0	7.0	18.0	64.9	198.3
1924	36.4	1,137.0	107.0	2.7	1.0	1.3	2.2	3.6	6.8	18.7	60.2	189.7
1925	41.5	1,303.1	121.4	3.2	1.2	1.3	2.5	4.2	7.7	21.1	75.0	230.8
1926	35.2	1,150.7	104.6	2.8	0.9	1.4	2.2	4.2	6.3	18.3	58.6	179.8
1927	29.1	938.3	90.6	2.8	0.9	1.3	2.3	3.5	5.7	13.7	50.8	158.8
1928	28.5	937.8	100.2	2.7	1.0	1.3	2.0	3.5	6.1	16.1	49.5	144.7
1929	25.0	836.3	86.9	2.6	1.0	1.1	2.2	3.4	5.8	13.0	46.0	119.2
1930	28.3	938.2	104.5	3.2	1.0	1.2	2.1	3.5	7.1	15.8	54.5	163.5
1931	21.8	764.6	73.3	2.5	0.9	1.2	1.8	3.5	6.1	13.2	42.5	124.5
1932	17.1	614.8	59.8	1.7	0.8	1.1	1.7	3.3	4.9	11.0	36.7	105.1
1933	18.5	657.0	72.2	2.0	0.9	1.3	1.7	3.2	6.2	12.6	39.9	104.2
1934	19.7	748.6	70.6	2.1	0.9	1.3	1.9	3.6	6.0	13.0	41.3	132.0
1935	15.1	599.2	47.4	1.5	0.9	1.4	1.7	3.0	4.7	10.2	29.9	90.0
1936	17.4	689.7	62.2	1.7	0.7	1.2	1.8	2.9	5.3	9.1	37.5	87.9
1937	15.9	667.0	50.6	1.4	0.7	1.0	1.9	2.7	3.8	8.2	28.1	81.8
1938	15.1	621.6	47.9	1.3	0.8	1.0	1.4	2.5	4.2	8.3	26.5	84.5
1939	12.4	496.2	37.5	1.0	0.8	1.0	1.4	2.6	3.4	7.2	22.4	88.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ALL RACES, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	134.9	4,104.5	288.5	8.8	5.6	8.2	11.4	19.6	49.5	144.4	219.6	692.6
1901	112.1	3,288.6	250.2	8.3	4.9	7.8	10.8	19.1	52.4	119.2	281.6	621.9
1902	99.5	2,923.0	231.4	6.7	3.7	6.7	9.6	17.8	44.2	104.7	247.3	410.5
1903	94.4	2,852.8	194.7	6.5	3.1	5.7	7.9	16.0	38.3	101.6	256.1	607.6
1904	105.8	3,254.6	237.6	7.7	3.5	5.0	7.8	15.7	34.0	98.3	245.1	536.5
1905	110.4	3,470.5	238.7	6.5	3.2	4.6	7.5	14.1	35.3	94.7	272.1	656.3
1906	117.3	3,532.8	263.9	6.7	2.7	4.8	6.4	13.7	38.2	106.6	297.0	691.6
1907	109.1	3,367.7	241.3	5.9	2.6	4.4	6.3	12.0	31.7	87.8	259.2	548.0
1908	107.8	3,332.0	238.8	5.9	2.7	4.5	6.5	9.8	30.6	84.5	249.4	647.1
1909	97.0	2,977.0	231.6	5.7	2.1	3.5	5.9	10.8	26.3	70.9	228.0	577.9
1910	109.9	3,428.0	255.1	6.0	2.2	3.9	5.8	10.9	27.0	87.0	246.0	628.0
1911	82.8	2,521.5	190.2	4.3	2.7	2.9	6.0	9.4	21.9	71.2	200.3	490.8
1912	75.0	2,289.0	182.4	4.3	2.0	3.0	4.6	8.9	23.8	63.4	192.9	425.6
1913	82.1	2,438.0	211.5	4.7	2.0	3.3	5.3	9.7	21.3	62.7	193.0	482.6
1914	70.5	2,105.3	173.0	5.0	2.4	3.7	4.8	8.2	19.8	56.5	172.3	408.9
1915	63.2	1,914.4	157.2	4.4	1.6	2.7	4.6	7.4	18.1	52.0	146.4	349.7
1916	70.6	2,025.0	204.9	4.7	2.1	3.1	4.9	7.7	17.1	46.0	150.0	358.0
1917	69.4	1,937.5	203.6	6.0	2.1	3.8	5.8	8.5	19.4	54.0	151.1	349.1
1918	66.2	1,889.3	197.0	5.2	2.0	3.3	4.7	9.0	16.1	49.1	149.3	358.1
1919	50.9	1,502.1	136.1	4.4	2.0	2.9	4.1	8.0	13.0	41.5	118.7	308.5
1920	49.3	1,433.9	131.1	4.0	1.7	3.4	3.9	7.4	13.4	35.2	112.4	286.0
1921	47.4	1,307.4	138.5	4.3	1.5	2.5	3.2	5.3	11.4	34.5	129.7	315.3
1922	35.7	1,035.6	96.5	2.7	1.2	2.0	2.9	4.7	9.6	25.7	82.7	241.1
1923	36.1	1,048.8	111.2	2.9	1.4	1.7	2.2	4.2	8.0	26.4	83.0	237.3
1924	30.9	878.2	94.5	2.8	1.4	1.8	2.4	3.8	8.4	24.8	80.9	196.1
1925	35.6	1,045.4	109.2	3.0	1.1	1.8	2.5	4.1	8.5	24.3	82.7	263.3
1926	30.6	923.7	95.5	2.7	1.2	1.7	2.2	4.7	7.5	22.6	69.4	208.1
1927	25.0	730.6	80.7	2.6	1.1	1.8	2.3	3.7	7.2	18.4	65.1	203.8
1928	24.3	722.7	85.4	2.4	1.1	1.5	2.2	4.0	6.3	19.2	65.0	183.0
1929	21.5	671.2	74.4	2.2	1.2	1.7	2.3	3.8	6.6	16.7	52.3	141.2
1930	23.7	732.3	86.4	2.8	1.2	1.7	2.4	3.7	7.4	17.7	64.2	181.1
1931	19.1	620.4	67.8	2.2	1.2	1.4	2.0	3.7	6.1	14.0	51.1	141.1
1932	15.0	491.3	53.7	1.8	1.0	1.3	1.9	3.1	5.9	12.3	45.9	115.2
1933	16.0	522.0	61.4	1.9	1.1	1.8	1.9	3.7	6.1	13.9	44.4	151.1
1934	17.0	599.9	61.1	2.1	1.1	1.7	2.2	3.5	6.1	14.0	47.6	136.9
1935	13.1	478.3	39.7	1.4	1.0	1.6	2.0	3.0	5.4	12.2	35.5	110.7
1936	15.3	568.5	53.5	1.4	1.1	1.6	1.7	3.4	5.1	11.2	43.6	129.1
1937	13.5	523.7	43.8	1.3	1.1	1.3	1.9	2.5	4.8	10.4	31.9	90.5
1938	13.4	512.4	42.5	1.2	1.0	1.4	1.8	2.4	4.8	9.3	33.7	84.1
1939	10.7	399.5	33.1	0.9	0.8	1.2	1.7	2.1	3.7	8.8	27.6	88.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

WHITE, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	114.6	---	---	---	---	---	---	---	---	---	---	---
1911	86.1	---	---	---	---	---	---	---	---	---	---	---
1912	79.0	---	---	---	---	---	---	---	---	---	---	---
1913	86.0	---	---	---	---	---	---	---	---	---	---	---
1914	74.3	2,312.3	183.9	5.0	1.8	3.4	5.0	8.0	17.7	50.6	157.3	390.3
1915	66.5	2,103.8	164.8	4.3	1.6	2.6	4.6	7.3	16.2	45.3	127.8	336.2
1916	74.0	2,244.5	206.6	4.7	1.9	3.1	5.2	7.5	16.2	43.8	132.1	347.5
1917	72.6	---	---	---	---	---	---	---	---	---	---	---
1918	69.7	2,075.6	194.7	5.3	1.9	3.3	4.5	8.1	15.3	42.7	133.4	333.4
1919	53.8	1,663.7	141.0	4.4	1.5	2.7	4.2	6.7	12.7	36.6	106.2	303.3
1920	52.1	1,584.4	134.9	3.9	1.4	2.6	3.8	6.8	12.1	31.8	108.2	261.7
1921	49.1	1,425.3	140.7	4.3	1.1	1.6	2.4	4.0	9.2	28.5	117.4	311.2
1922	37.4	1,134.0	99.4	2.8	1.0	1.4	2.2	3.7	7.6	22.1	77.2	221.3
1923	37.1	1,134.7	109.8	2.8	1.0	1.3	1.8	3.4	6.6	21.9	75.3	219.6
1924	30.9	939.2	89.9	2.5	0.9	1.2	2.0	2.9	6.7	20.4	69.8	188.9
1925	36.1	1,104.2	108.3	2.9	0.9	1.4	1.9	3.5	7.2	21.6	79.1	250.9
1926	30.5	976.8	91.4	2.5	0.8	1.3	1.8	3.6	6.0	19.3	64.4	196.8
1927	24.8	774.3	77.9	2.6	0.8	1.3	1.8	2.9	5.5	15.2	57.9	181.4
1928	24.2	760.9	84.9	2.4	0.8	1.0	1.6	2.9	5.3	16.9	57.2	165.7
1929	21.3	697.7	75.1	2.2	0.9	1.1	1.9	2.9	5.0	13.8	48.5	127.6
1930	24.0	784.1	89.1	2.9	0.8	1.1	1.9	2.8	6.3	15.9	59.7	169.9
1931	18.7	646.3	64.1	2.2	0.9	1.0	1.6	2.7	5.2	12.3	45.3	135.6
1932	14.8	494.5	52.2	1.6	0.8	1.0	1.4	2.6	4.4	10.8	39.9	110.2
1933	15.8	501.0	57.3	1.7	0.8	1.2	1.4	2.6	4.9	12.1	41.0	130.4
1934	16.5	575.8	55.6	1.9	0.8	1.2	1.5	2.6	5.0	12.3	43.4	131.6
1935	12.7	503.1	38.9	1.3	0.8	1.2	1.5	2.4	4.2	10.3	32.6	103.8
1936	14.8	581.9	53.0	1.4	0.8	1.0	1.2	2.4	4.5	9.5	40.4	112.4
1937	13.2	550.9	41.7	1.2	0.8	1.0	1.5	2.0	3.5	8.9	29.9	88.6
1938	12.5	513.4	39.6	1.1	0.8	1.0	1.3	1.8	3.6	8.1	29.6	79.8
1939	10.3	410.0	31.2	0.9	0.8	0.9	1.2	2.0	3.0	7.5	24.5	88.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

WHITE, MALE												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	119.8	---	---	---	---	---	---	---	---	---	---	---
1911	90.0	---	---	---	---	---	---	---	---	---	---	---
1912	82.6	---	---	---	---	---	---	---	---	---	---	---
1913	90.6	---	---	---	---	---	---	---	---	---	---	---
1914	78.9	2,543.8	197.3	5.1	1.5	3.2	5.5	8.4	16.6	45.1	140.7	366.1
1915	70.6	2,313.7	174.9	4.6	1.7	2.8	5.0	7.6	15.1	39.8	108.0	307.0
1916	78.7	2,474.8	214.4	4.9	1.9	3.3	5.9	7.8	16.3	42.7	112.2	317.5
1917	78.3	---	---	---	---	---	---	---	---	---	---	---
1918	76.0	2,296.3	205.9	5.8	2.1	4.0	5.0	8.1	15.4	37.2	116.1	300.8
1919	58.1	1,842.3	151.2	4.7	1.3	3.0	4.7	6.8	12.9	33.2	93.1	279.2
1920	56.5	1,770.1	144.7	4.1	1.5	2.5	4.4	7.2	11.7	29.6	102.3	225.2
1921	52.6	1,580.6	148.7	4.7	1.1	1.2	2.2	3.6	8.4	24.1	100.9	297.9
1922	40.6	1,294.5	106.8	2.8	1.1	1.3	2.0	3.6	6.9	19.6	70.7	206.5
1923	40.0	1,261.6	116.2	2.9	1.1	1.3	1.9	3.5	6.3	18.3	65.8	200.4
1924	33.5	1,060.8	94.7	2.5	0.8	1.1	2.0	3.1	6.3	17.6	58.9	181.7
1925	39.1	1,228.9	114.6	3.0	1.0	1.1	2.0	3.7	7.1	20.5	75.0	224.7
1926	32.7	1,082.9	95.7	2.7	0.7	1.2	1.9	3.5	5.7	17.1	58.2	177.6
1927	26.7	869.4	82.6	2.7	0.8	1.1	2.0	2.8	5.0	13.1	50.9	149.4
1928	26.1	852.8	91.2	2.5	0.8	1.0	1.5	2.7	5.1	15.2	49.1	144.0
1929	22.8	767.2	80.3	2.3	0.8	0.9	1.9	2.9	4.9	12.2	45.0	117.8
1930	26.4	885.5	98.0	3.1	0.7	1.0	1.8	2.9	6.3	14.9	54.4	153.0
1931	19.9	716.7	65.9	2.3	0.7	0.9	1.5	2.9	5.3	12.0	40.6	131.4
1932	15.9	555.3	54.7	1.6	0.7	1.0	1.4	2.7	4.1	10.0	35.5	105.4
1933	17.0	558.3	62.4	1.7	0.7	1.1	1.5	2.5	5.2	11.5	38.8	101.2
1934	17.7	640.3	58.7	1.9	0.8	1.0	1.5	2.9	5.1	12.0	39.8	131.7
1935	13.7	556.0	43.0	1.4	0.8	1.1	1.4	2.6	4.0	9.6	29.8	88.6
1936	15.7	637.8	56.7	1.5	0.7	1.0	1.3	2.3	4.7	8.5	37.5	85.1
1937	14.3	617.8	44.3	1.2	0.6	0.9	1.7	2.3	3.1	7.8	27.9	84.7
1938	13.4	560.2	42.0	1.2	0.7	0.9	1.2	1.9	3.6	7.5	26.2	79.1
1939	11.1	449.1	33.5	1.0	0.8	0.9	1.1	2.3	3.0	7.0	22.2	85.8

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.  
 Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

WHITE, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	109.0	---	---	---	---	---	---	---	---	---	---	---
1911	81.9	---	---	---	---	---	---	---	---	---	---	---
1912	75.3	---	---	---	---	---	---	---	---	---	---	---
1913	81.2	---	---	---	---	---	---	---	---	---	---	---
1914	69.5	2,074.7	170.2	5.0	2.2	3.6	4.5	7.6	19.0	56.2	172.6	409.3
1915	62.3	1,888.3	154.4	4.1	1.5	2.4	4.1	6.9	17.3	50.9	145.8	358.9
1916	69.2	2,008.1	198.6	4.5	1.9	2.8	4.4	7.2	16.2	45.0	150.0	370.4
1917	66.7	---	---	---	---	---	---	---	---	---	---	---
1918	63.5	1,849.0	183.3	4.7	1.6	2.7	3.9	8.1	15.2	48.4	149.0	358.0
1919	49.4	1,480.3	130.6	4.0	1.7	2.4	3.6	6.5	12.4	40.1	118.1	321.4
1920	47.6	1,393.3	124.8	3.7	1.4	2.8	3.1	6.3	12.5	34.1	113.6	289.2
1921	45.6	1,264.1	132.4	4.0	1.1	2.1	2.6	4.5	10.1	32.9	132.4	321.3
1922	34.1	998.8	91.6	2.7	1.0	1.6	2.3	3.8	8.5	24.7	83.0	232.6
1923	34.1	1,003.4	103.1	2.7	1.0	1.3	1.7	3.2	6.8	25.7	84.0	234.4
1924	28.2	813.4	84.9	2.5	1.0	1.3	1.9	2.8	7.0	23.2	79.8	194.5
1925	33.1	974.8	101.7	2.8	0.8	1.6	1.8	3.4	7.3	22.7	82.8	271.3
1926	28.2	866.8	87.0	2.4	0.8	1.4	1.8	3.8	6.3	21.6	70.1	211.7
1927	22.8	676.2	72.9	2.4	0.8	1.4	1.7	2.9	6.1	17.3	64.4	206.4
1928	22.1	655.7	78.3	2.2	0.8	1.0	1.6	3.1	5.5	18.7	64.7	182.4
1929	19.8	625.8	69.7	2.1	1.0	1.4	1.8	2.8	5.1	15.6	51.8	135.1
1930	21.6	679.0	79.8	2.6	1.0	1.2	2.0	2.6	6.2	17.0	64.6	182.6
1931	17.4	573.1	62.1	2.1	1.0	1.1	1.7	2.6	5.2	12.7	49.8	138.7
1932	13.7	431.3	49.6	1.6	0.8	1.1	1.5	2.4	4.6	11.5	44.0	113.8
1933	14.6	441.6	52.0	1.6	0.9	1.4	1.4	2.6	4.7	12.7	43.1	152.5
1934	15.3	509.1	52.4	1.8	0.9	1.3	1.6	2.4	4.9	12.6	46.8	131.5
1935	11.8	448.3	34.7	1.3	0.8	1.3	1.5	2.2	4.3	11.1	35.2	115.2
1936	13.8	524.2	49.2	1.3	0.9	1.1	1.2	2.4	4.4	10.5	43.1	133.0
1937	12.1	481.7	39.0	1.1	1.0	1.1	1.3	1.6	3.9	10.0	31.8	91.5
1938	11.7	465.0	37.0	1.0	0.9	1.1	1.4	1.6	3.6	8.7	32.7	80.3
1939	9.6	369.6	28.9	0.8	0.8	1.0	1.3	1.7	2.9	8.0	26.6	91.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ALL OTHER, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	147.1	---	---	---	---	---	---	---	---	---	---	---
1911	109.3	---	---	---	---	---	---	---	---	---	---	---
1912	95.9	---	---	---	---	---	---	---	---	---	---	---
1913	102.4	---	---	---	---	---	---	---	---	---	---	---
1914	94.2	3,105.6	250.6	7.3	4.9	6.7	11.4	17.9	32.4	70.1	129.2	*
1915	89.0	2,862.8	254.2	10.1	*	6.6	10.6	13.9	38.3	66.1	167.6	*
1916	97.5	2,540.6	304.6	7.8	4.1	5.7	12.3	14.0	30.0	62.2	159.6	241.2
1917	109.8	---	---	---	---	---	---	---	---	---	---	---
1918	103.6	2,569.2	383.7	10.5	5.6	7.8	12.8	18.9	28.8	62.4	141.2	256.0
1919	69.9	1,882.3	214.1	7.2	4.4	7.0	7.7	17.5	21.6	55.3	118.9	169.2
1920	71.4	2,098.6	210.1	5.6	4.0	7.3	10.2	15.9	23.7	48.0	95.1	223.0
1921	68.5	1,920.1	203.7	6.5	3.6	5.6	7.4	12.0	24.5	49.5	85.4	215.5
1922	53.8	1,514.4	156.5	3.7	2.7	4.5	7.4	13.5	20.3	34.8	73.7	228.5
1923	58.9	1,628.2	197.8	4.2	3.4	4.4	6.2	12.7	20.5	33.6	58.6	228.1
1924	62.1	1,706.4	202.8	4.6	3.9	5.3	6.1	12.5	21.4	44.6	90.5	236.7
1925	64.2	1,909.0	182.9	4.8	3.4	3.8	9.3	11.4	21.4	41.8	78.1	232.5
1926	57.6	1,641.0	181.3	4.6	3.5	3.7	6.5	14.2	19.9	40.6	60.3	185.9
1927	49.8	1,401.6	155.9	3.9	2.9	4.3	7.3	12.0	19.4	31.5	63.3	211.5
1928	46.8	1,340.6	156.2	3.9	3.0	4.8	7.2	12.2	18.5	29.3	63.0	175.0
1929	40.5	1,261.3	125.0	3.9	3.1	3.9	6.1	10.7	21.8	31.3	61.6	168.5
1930	44.0	1,262.4	146.8	4.4	3.3	4.4	5.9	11.8	20.8	29.5	56.7	206.5
1931	37.1	1,071.0	121.7	3.8	2.3	3.5	5.6	12.6	18.4	34.3	73.7	119.6
1932	27.5	1,011.3	93.0	3.2	1.9	2.9	5.5	9.5	19.7	26.2	66.9	116.7
1933	30.0	1,242.8	139.9	3.8	2.5	4.3	5.6	12.4	22.9	31.1	61.2	135.1
1934	34.7	1,393.0	142.5	3.9	2.2	4.3	6.7	12.6	20.2	32.1	63.6	163.5
1935	26.2	803.8	77.6	2.3	1.9	4.3	5.2	9.5	16.4	23.7	36.5	84.1
1936	30.4	981.3	91.9	2.6	2.1	4.4	6.3	11.2	14.4	19.2	45.4	102.7
1937	28.1	928.7	85.9	2.5	1.8	2.8	5.4	9.2	14.8	14.8	33.0	70.3
1938	29.3	966.5	84.6	2.6	1.7	2.8	4.4	9.5	16.2	18.6	40.6	124.8
1939	22.4	731.8	63.5	1.6	1.3	2.8	4.7	6.5	11.1	15.0	36.2	82.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ALL OTHER, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	150.4	...	...	...	...	...	...	...	...	...	...	...
1911	109.4	...	...	...	...	...	...	...	...	...	...	...
1912	95.6	...	...	...	...	...	...	...	...	...	...	...
1913	102.9	...	...	...	...	...	...	...	...	...	...	...
1914	96.0	3,389.2	262.9	9.3	*	*	11.5	*	*	73.2	*	*
1915	92.6	3,207.4	283.1	8.3	*	*	*	*	*	36.3	*	*
1916	103.1	2,838.5	319.4	7.9	*	*	12.4	12.6	23.4	55.8	171.0	*
1917	114.8	...	...	...	...	...	...	...	...	...	...	...
1918	106.5	2,772.4	400.7	10.8	5.0	4.6	10.6	15.5	23.2	60.2	127.4	*
1919	73.0	2,046.7	234.4	6.5	4.4	5.0	7.1	10.1	19.0	45.8	106.7	*
1920	74.0	2,319.5	223.8	4.6	*	5.0	7.5	11.3	17.8	42.5	99.8	*
1921	69.3	2,071.5	205.6	5.8	*	4.1	5.1	8.8	16.2	37.0	93.4	*
1922	56.6	1,649.8	172.1	4.3	2.6	*	6.3	11.6	14.0	27.4	70.2	*
1923	61.3	1,774.5	209.4	3.7	*	3.1	5.1	9.5	15.6	28.6	*	*
1924	66.5	1,911.7	222.6	4.3	2.8	4.1	4.6	9.7	13.4	36.4	80.6	*
1925	67.1	2,075.6	186.2	4.9	3.0	3.6	8.6	10.3	16.5	30.5	74.3	298.3
1926	61.0	1,825.2	188.6	4.0	2.9	2.6	5.8	12.8	14.0	38.2	64.6	*
1927	52.9	1,595.0	162.5	3.9	2.2	3.3	5.8	11.0	14.5	23.4	*	257.2
1928	50.5	1,491.9	171.9	4.2	3.1	3.4	6.5	10.7	18.7	29.8	55.8	*
1929	44.4	1,471.3	139.6	4.4	2.8	3.0	5.3	7.7	15.8	26.6	62.6	*
1930	46.3	1,374.2	157.2	3.9	3.3	2.7	5.0	9.0	17.1	28.9	55.9	258.2
1931	39.2	1,156.0	132.2	4.0	2.3	3.4	5.7	10.4	16.5	33.4	74.6	*
1932	28.6	1,083.8	100.9	2.7	1.8	2.5	4.8	8.7	15.0	26.5	55.9	*
1933	32.2	1,394.5	148.4	4.1	1.9	3.5	4.5	9.8	19.2	28.9	56.9	*
1934	37.3	1,551.6	160.7	3.7	1.8	3.6	5.2	10.4	17.0	27.9	65.3	*
1935	28.0	920.1	80.4	2.3	1.6	4.3	4.5	7.5	13.1	19.4	*	*
1936	32.4	1,080.7	101.2	3.0	*	3.3	6.2	9.3	13.7	16.8	38.4	*
1937	30.3	1,038.1	95.2	2.8	1.8	2.6	4.3	7.3	12.4	13.4	*	*
1938	31.0	1,086.6	89.6	2.0	*	*	3.2	8.1	11.7	18.5	*	133.7
1939	24.2	853.3	66.0	1.5	*	2.2	4.2	6.3	7.3	10.2	*	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

GASTRITIS, DUODENITIS, ENTERITIS AND COLITIS, EXCEPT DIARRHEA OF NEWBORN

ALL OTHER, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	143.4	...	...	...	...	...	...	...	...	...	...	...
1911	109.2	...	...	...	...	...	...	...	...	...	...	...
1912	96.3	...	...	...	...	...	...	...	...	...	...	...
1913	101.9	...	...	...	...	...	...	...	...	...	...	...
1914	92.3	2,824.4	238.7	*	*	*	11.3	24.1	44.7	66.5	*	*
1915	85.3	2,522.0	225.9	11.8	*	*	13.7	19.1	40.8	88.7	*	*
1916	91.7	2,247.2	290.0	7.7	4.7	7.2	12.1	15.9	38.3	69.7	148.4	*
1917	104.7	...	...	...	...	...	...	...	...	...	...	...
1918	100.7	2,369.7	367.0	10.1	6.1	10.9	15.1	23.6	36.2	65.0	154.6	358.9
1919	66.9	1,720.4	194.1	7.9	4.4	8.9	8.4	27.7	25.0	66.6	131.0	*
1920	68.8	1,883.6	196.6	6.7	5.2	9.5	13.0	22.2	31.3	54.6	90.4	252.3
1921	67.6	1,771.0	201.9	7.2	4.6	6.9	9.9	16.3	35.3	64.6	77.3	253.6
1922	51.0	1,381.6	140.8	3.2	2.8	6.0	8.6	15.9	28.5	43.8	77.2	316.3
1923	56.5	1,484.4	186.1	4.6	4.7	5.6	7.3	16.7	26.8	39.7	66.0	263.3
1924	57.7	1,504.5	182.8	4.9	4.9	6.4	7.6	16.0	31.8	54.7	100.4	210.8
1925	61.3	1,744.7	179.5	4.6	3.7	4.0	10.0	12.9	27.8	55.5	81.9	188.8
1926	54.3	1,460.1	173.8	5.1	4.1	4.7	7.2	15.8	27.5	43.6	55.9	*
1927	46.7	1,215.1	149.2	3.9	3.5	5.2	8.8	13.2	25.8	41.2	78.1	179.6
1928	43.1	1,194.0	140.5	3.6	2.9	6.1	8.0	13.9	18.3	28.7	70.1	188.4
1929	36.6	1,059.0	110.4	3.4	3.3	4.8	6.9	14.1	29.7	36.7	60.6	195.5
1930	41.8	1,151.8	136.4	4.8	3.4	6.0	6.8	15.0	25.6	30.2	57.4	167.5
1931	34.9	987.3	111.4	3.6	2.4	3.6	5.5	15.2	20.8	35.2	72.8	163.6
1932	26.4	939.8	85.2	3.7	2.0	3.3	6.2	10.5	25.8	25.8	77.4	127.6
1933	28.0	1,093.7	131.6	3.5	2.9	4.9	6.7	15.3	27.4	33.5	65.2	138.2
1934	32.2	1,237.3	124.4	4.2	2.6	5.0	8.2	15.1	24.2	36.7	61.9	184.7
1935	24.4	689.6	74.9	2.4	2.3	4.4	5.9	11.8	20.4	28.5	40.9	*
1936	28.4	884.3	82.6	2.2	2.9	5.4	6.3	13.3	15.2	21.8	52.0	*
1937	25.9	822.1	76.7	2.2	1.8	3.1	6.6	11.3	17.7	16.3	33.9	*
1938	27.6	849.7	79.7	3.2	2.0	4.1	5.5	10.9	21.5	18.7	49.5	118.2
1939	20.5	613.4	61.1	1.7	1.5	3.4	5.1	6.8	15.4	20.3	44.9	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

NOTES:  
For 1932-34 deaths of Mexicans are included with total deaths for the white population; but by age, are included with deaths for the nonwhite population. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes-----	569	1,401	1,177
Male-----	298	740	643
Female-----	271	661	534

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ICD Revision	Years in use	ICD Nos.
Fifth	1939	124
Fourth	1930-1938	124
Third	1921-1929	122
Second	1910-1920	113
First	1900-1909	112

ALL RACES, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	12.5	*	*	*	0.8	5.7	16.1	29.0	47.9	66.1	60.3	*
1901	13.1	*	*	*	0.9	5.0	16.6	31.8	47.7	72.7	67.4	*
1902	13.0	*	*	*	0.9	5.9	14.7	31.2	49.1	71.8	66.8	*
1903	13.5	*	*	*	0.6	5.3	17.2	28.6	53.5	74.5	82.5	50.8
1904	13.9	*	*	*	0.7	4.8	16.9	33.4	52.6	76.6	74.7	*
1905	14.0	*	*	*	0.8	5.6	15.5	33.9	53.4	76.2	77.5	59.3
1906	14.1	*	*	0.4	0.9	4.9	17.1	34.5	56.9	75.0	79.0	76.6
1907	14.8	*	*	0.4	0.7	5.4	17.4	36.2	58.1	80.9	95.2	80.1
1908	13.5	*	*	0.4	0.6	4.1	15.9	32.9	54.9	77.2	83.3	72.9
1909	13.4	*	*	0.3	0.6	4.3	14.2	31.5	54.6	78.7	91.7	75.8
1910	13.2	*	*	*	0.6	4.3	14.7	31.4	54.2	77.0	86.6	77.2
1911	13.6	*	0.6	0.4	0.7	4.3	14.4	32.5	52.6	82.8	100.0	75.9
1912	13.1	*	*	0.2	0.5	3.9	14.0	29.7	53.2	78.8	94.0	75.8
1913	12.9	*	*	0.3	0.6	3.8	14.3	30.0	49.5	80.1	86.5	77.2
1914	12.5	*	0.5	0.3	0.5	3.6	13.3	29.5	45.4	76.9	95.2	76.1
1915	12.1	1.7	0.5	0.3	0.6	3.5	12.2	26.7	46.7	73.1	90.6	86.2
1916	11.8	*	0.5	0.3	0.6	3.7	11.9	25.8	48.1	70.3	89.2	78.7
1917	10.9	*	*	0.2	0.5	2.7	10.3	24.6	43.2	66.9	90.6	81.1
1918	9.6	*	0.5	0.3	0.5	2.4	8.5	19.2	38.4	60.9	80.9	74.0
1919	7.9	*	0.3	0.2	0.3	1.4	5.8	15.4	31.4	55.3	83.7	73.2
1920	7.1	*	0.3	0.2	0.5	1.3	4.5	12.9	28.1	53.2	78.2	76.6
1921	7.3	1.1	0.3	0.2	0.4	1.4	4.4	13.2	29.9	54.4	78.8	84.5
1922	7.4	1.1	0.3	0.3	0.4	1.4	5.1	13.7	28.9	52.8	78.8	99.7
1923	7.1	*	0.3	0.2	0.4	1.2	4.9	14.0	27.5	49.7	73.1	96.8
1924	7.3	*	0.3	0.3	0.4	1.5	5.2	15.3	27.1	50.2	64.8	84.9
1925	7.2	*	0.3	0.2	0.4	1.4	5.8	14.1	27.5	46.1	71.7	83.8
1926	7.2	*	*	0.1	0.3	1.6	5.6	13.7	27.8	44.0	71.2	77.6
1927	7.4	*	0.3	0.2	0.5	1.5	5.9	14.4	26.7	48.8	66.6	76.0
1928	7.5	*	0.3	0.2	0.4	1.6	6.4	15.0	27.7	46.3	63.6	68.3
1929	7.2	*	*	0.1	0.4	1.5	6.3	14.5	26.1	43.8	57.1	69.5
1930	7.2	*	*	0.2	0.4	1.5	6.1	14.0	26.8	43.0	57.5	64.4
1931	7.4	*	0.3	0.2	0.4	1.7	5.9	14.8	26.8	42.0	59.4	68.1
1932	7.2	*	*	0.2	0.3	1.4	5.7	14.0	26.1	41.2	58.4	57.6
1933	7.4	*	*	0.2	0.3	1.6	5.9	14.3	26.9	41.9	59.9	58.1
1934	7.7	*	0.3	0.1	0.4	1.8	6.4	15.1	27.2	40.4	60.1	52.8
1935	7.9	1.2	*	0.1	0.3	1.6	6.4	15.7	27.3	41.5	60.2	60.3
1936	8.3	1.2	0.3	0.2	0.3	1.7	7.1	15.8	28.8	41.6	57.7	66.7
1937	8.5	*	0.3	0.2	0.4	1.7	7.2	17.0	28.3	41.3	59.4	57.3
1938	8.3	1.3	*	0.2	0.4	1.7	6.9	16.2	27.1	40.7	58.3	54.9
1939	8.3	1.0	*	0.2	0.4	1.7	6.6	15.5	27.8	40.3	55.6	58.6

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ALL RACES, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	15.5	*	*	*	*	7.5	18.9	35.7	63.6	82.2	96.0	*
1901	16.6	*	*	*	1.2	6.9	20.7	41.0	61.4	90.8	81.1	*
1902	16.8	*	*	*	*	7.1	20.0	40.0	67.8	90.0	82.5	*
1903	17.0	*	*	*	*	6.6	20.8	37.2	71.1	92.8	96.8	*
1904	17.2	*	*	*	*	5.7	21.4	42.8	69.2	90.5	82.6	*
1905	18.0	*	*	*	*	6.9	19.6	45.6	72.9	94.8	94.6	*
1906	17.9	*	*	*	1.2	5.5	21.9	44.2	76.9	95.1	91.3	91.3
1907	19.1	*	*	*	0.9	6.8	22.3	47.5	78.2	102.7	116.8	95.1
1908	17.3	*	*	0.6	0.7	4.6	20.1	42.7	72.3	98.8	99.6	103.3
1909	17.5	*	*	*	0.8	5.3	18.6	43.0	72.9	98.4	114.6	85.1
1910	17.2	*	*	*	0.6	4.9	19.2	41.6	73.1	101.0	103.7	80.7
1911	17.9	*	*	*	0.8	5.5	17.8	44.8	72.8	108.3	115.0	85.9
1912	17.3	*	*	*	0.6	5.1	18.0	39.8	73.3	103.6	116.3	104.7
1913	17.1	*	*	*	0.8	4.7	19.1	39.8	68.1	105.9	103.8	91.1
1914	16.7	*	*	0.4	0.5	4.7	17.7	40.7	63.7	98.9	116.7	71.4
1915	15.9	*	*	*	0.7	4.2	15.5	36.2	64.2	94.3	112.7	99.0
1916	15.7	*	*	*	0.8	4.5	15.4	35.3	65.3	94.1	105.8	96.2
1917	14.3	*	*	*	0.6	3.4	13.3	32.2	60.3	86.9	106.1	98.2
1918	12.9	*	*	0.3	0.6	3.0	11.3	26.5	52.1	80.4	96.3	74.8
1919	10.3	*	*	*	0.4	1.6	7.8	20.1	43.3	70.3	99.3	85.5
1920	9.0	*	*	*	0.5	1.4	5.8	16.7	36.7	67.6	90.5	94.6
1921	9.2	*	*	*	0.5	1.7	5.6	16.7	39.0	67.5	90.5	98.7
1922	9.4	*	*	0.3	0.4	1.6	6.4	17.7	38.7	66.3	94.5	115.5
1923	9.3	*	*	0.2	0.4	1.3	6.3	19.1	37.5	62.4	88.7	106.7
1924	9.5	*	*	0.3	0.3	1.8	6.7	20.2	37.1	64.7	75.4	98.5
1925	9.2	*	*	0.2	0.3	1.5	7.3	18.8	36.6	57.2	79.8	95.9
1926	9.2	*	*	*	0.4	1.6	6.8	18.3	36.7	56.7	84.4	83.1
1927	9.5	*	*	0.3	0.5	1.8	7.3	19.1	34.7	63.3	79.8	81.5
1928	9.7	*	*	0.3	0.4	1.9	7.6	19.7	37.3	60.8	77.8	73.8
1929	9.1	*	*	0.2	0.4	1.6	7.7	18.7	35.3	55.6	64.4	75.4
1930	9.2	*	*	0.2	0.4	1.6	7.3	17.9	36.4	55.8	65.7	66.3
1931	9.5	*	*	*	0.4	1.9	7.2	19.5	36.2	54.2	70.3	82.4
1932	9.3	*	*	*	0.3	1.6	7.1	18.2	35.2	53.5	70.5	60.9
1933	9.5	*	*	0.2	0.3	1.8	7.3	18.4	36.9	53.0	69.3	55.2
1934	10.1	*	*	*	0.3	2.0	8.2	20.1	37.5	53.1	71.5	59.6
1935	10.5	*	*	0.2	0.2	1.9	8.4	20.4	37.9	55.4	75.5	76.1
1936	11.0	*	*	0.2	0.3	1.9	9.2	21.0	39.9	55.4	73.4	73.7
1937	11.2	*	*	0.2	0.4	1.9	9.4	22.8	38.8	55.6	71.1	59.8
1938	11.0	*	*	0.2	0.4	2.0	8.5	21.6	38.1	54.5	68.6	66.6
1939	11.1	*	*	*	0.3	1.8	8.6	21.4	38.9	54.6	67.6	67.6

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ALL RACES, FEMALE

Years												
1900	9.5	*	*	*	*	3.9	13.1	21.9	32.4	51.2	55.1	*
1901	9.5	*	*	*	*	3.0	12.1	22.2	34.2	55.5	55.0	*
1902	9.2	*	*	*	*	4.6	9.1	21.9	30.7	54.5	52.7	*
1903	10.0	*	*	*	*	3.9	13.3	19.6	36.3	57.1	69.7	*
1904	10.6	*	*	*	*	3.9	12.2	23.4	36.2	63.4	67.7	*
1905	9.9	*	*	*	*	4.3	11.1	21.5	34.1	58.6	62.2	*
1906	10.1	*	*	*	0.7	4.2	11.8	23.8	36.2	55.4	67.9	*
1907	10.4	*	*	*	0.6	3.8	12.0	23.7	37.3	59.7	75.6	68.8
1908	9.6	*	*	*	*	3.6	11.1	21.9	36.6	55.8	68.4	49.4
1909	9.2	*	*	*	*	3.3	9.4	18.8	35.3	59.1	70.9	68.5
1910	9.2	*	*	*	0.6	3.6	9.7	20.0	34.0	53.2	70.9	74.5
1911	9.1	*	*	*	0.4	0.6	2.9	10.5	18.7	30.8	57.1	86.2
1912	8.6	*	*	*	0.4	2.6	9.5	18.4	31.4	53.9	73.6	53.2
1913	8.5	*	*	*	0.4	2.8	9.0	18.9	29.3	54.1	70.6	66.4
1914	8.0	*	*	*	0.5	2.4	8.4	16.8	25.5	54.6	75.5	79.8
1915	8.1	*	*	*	0.5	2.7	8.7	15.9	27.6	51.6	70.4	76.3
1916	7.7	*	*	*	0.4	2.7	7.9	15.1	29.2	46.0	74.2	65.4
1917	7.3	*	*	*	0.4	1.9	7.0	16.0	24.4	46.3	76.5	68.1
1918	6.2	*	0.6	*	0.4	1.8	5.5	10.8	23.3	40.8	66.9	73.3
1919	5.4	*	*	*	*	1.3	3.6	10.0	18.2	39.8	69.4	63.9
1920	5.1	*	*	0.3	0.5	1.1	3.0	8.6	18.6	38.2	66.9	63.1
1921	5.4	*	*	0.3	0.4	1.0	3.2	9.3	19.9	40.9	68.2	73.7
1922	5.3	*	*	0.2	0.4	1.3	3.6	9.3	18.4	38.8	64.4	87.7
1923	5.0	*	*	0.2	0.3	1.1	3.5	8.3	16.7	36.4	58.9	89.3
1924	5.0	*	*	*	0.4	1.2	3.5	9.9	16.1	35.1	55.0	74.5
1925	5.3	*	*	0.2	0.4	1.3	4.2	9.1	17.5	34.6	64.3	74.5
1926	5.1	*	*	*	0.3	1.5	4.2	8.7	18.1	30.8	58.9	73.5
1927	5.3	*	*	0.2	0.4	1.3	4.3	9.4	18.0	33.7	54.4	71.8
1928	5.2	*	*	*	0.5	1.4	5.0	9.8	17.3	31.2	50.4	64.0
1929	5.2	*	*	*	0.4	1.4	4.8	9.9	16.1	31.6	50.2	64.9
1930	5.2	*	*	*	0.5	1.3	4.8	9.8	16.5	29.8	49.8	63.0
1931	5.2	*	*	0.2	0.4	1.5	4.6	9.8	16.6	29.4	49.2	57.4
1932	5.1	*	*	0.2	0.3	1.3	4.3	9.5	16.4	28.6	47.1	55.1
1933	5.4	*	*	0.2	0.4	1.4	4.3	9.9	16.3	30.5	51.0	60.3
1934	5.3	*	*	*	0.5	1.5	4.5	9.8	16.2	27.3	49.4	47.7
1935	5.3	*	*	*	0.4	1.2	4.4	10.6	16.0	27.4	45.9	48.4
1936	5.5	*	*	0.2	0.3	1.4	5.0	10.2	17.0	27.7	43.0	61.4
1937	5.7	*	*	0.2	0.4	1.5	4.9	10.8	17.0	27.0	48.6	55.3
1938	5.7	*	*	0.2	0.4	1.4	5.2	10.4	15.4	26.9	48.8	46.1
1939	5.5	*	*	0.2	0.5	1.6	4.6	9.2	16.1	26.1	44.5	51.9

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .  
 Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

WHITE, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	13.4	---	---	---	---	---	---	---	---	---	---	---
1911	13.7	---	---	---	---	---	---	---	---	---	---	---
1912	13.1	---	---	---	---	---	---	---	---	---	---	---
1913	12.9	---	---	---	---	---	---	---	---	---	---	---
1914	12.5	*	0.5	0.3	0.4	3.4	13.0	29.6	45.5	77.6	95.8	77.4
1915	12.1	1.7	0.5	0.2	0.5	3.4	11.9	26.5	46.6	73.8	91.1	86.3
1916	12.0	*	0.4	0.2	0.5	3.6	11.7	26.2	48.4	71.3	90.9	81.2
1917	11.1	---	---	---	---	---	---	---	---	---	---	---
1918	9.7	*	0.5	0.3	0.5	2.3	8.4	19.0	38.0	61.1	82.4	76.9
1919	8.0	*	0.4	0.1	0.3	1.2	5.6	15.2	31.1	55.3	84.6	78.1
1920	7.1	*	0.3	0.2	0.3	1.1	4.2	12.2	28.3	53.7	80.6	78.3
1921	7.3	1.1	0.3	0.2	0.4	1.3	4.0	12.7	29.3	54.6	80.1	86.7
1922	7.5	1.2	0.3	0.2	0.3	1.3	4.6	13.2	29.0	53.9	81.3	104.7
1923	7.2	*	0.3	0.2	0.3	1.0	4.6	13.5	27.4	50.2	74.5	97.2
1924	7.3	*	*	0.2	0.3	1.3	5.0	14.7	26.9	50.7	66.2	86.5
1925	7.3	*	0.3	0.2	0.3	1.2	5.3	13.7	27.4	46.2	72.9	87.0
1926	7.2	*	*	0.1	0.3	1.4	5.3	13.2	27.9	44.1	71.9	78.3
1927	7.5	*	0.3	0.2	0.4	1.4	5.6	13.8	26.7	49.4	67.6	76.6
1928	7.6	*	0.3	0.2	0.3	1.4	6.0	14.6	27.9	47.1	65.7	70.3
1929	7.2	*	*	0.1	0.3	1.3	5.9	13.8	26.3	44.2	58.2	73.2
1930	7.3	*	*	0.2	0.3	1.3	5.7	13.6	26.6	43.3	58.2	65.6
1931	7.5	*	0.3	0.2	0.3	1.4	5.5	14.6	26.8	42.3	61.1	72.7
1932	7.4	*	*	0.2	0.2	1.3	5.4	13.9	26.1	41.8	59.5	60.7
1933	7.6	*	*	0.2	0.2	1.5	5.4	13.9	26.8	42.4	60.7	60.9
1934	7.9	*	*	0.1	0.3	1.6	6.0	15.0	27.2	40.9	61.4	55.2
1935	8.2	1.2	*	0.2	0.3	1.4	6.2	15.7	27.4	42.5	61.3	62.9
1936	8.5	1.3	*	0.2	0.3	1.5	6.8	15.5	28.9	42.1	59.1	70.8
1937	8.8	*	0.3	0.2	0.3	1.6	7.1	17.0	28.4	42.1	61.1	58.8
1938	8.6	1.2	*	0.2	0.4	1.6	6.8	16.2	27.4	41.7	59.4	57.8
1939	8.6	*	*	0.2	0.4	1.6	6.5	15.5	28.1	41.3	56.7	62.4

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

WHITE, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	17.3	---	---	---	---	---	---	---	---	---	---	---
1911	18.0	---	---	---	---	---	---	---	---	---	---	---
1912	17.3	---	---	---	---	---	---	---	---	---	---	---
1913	17.2	---	---	---	---	---	---	---	---	---	---	---
1914	16.7	*	*	0.4	0.5	4.5	17.3	40.8	63.8	99.5	116.8	74.8
1915	15.9	*	*	*	0.6	4.1	14.8	36.2	64.3	95.6	113.0	97.9
1916	16.0	*	*	*	0.6	4.4	15.2	36.0	65.9	96.4	107.4	101.0
1917	14.6	---	---	---	---	---	---	---	---	---	---	---
1918	13.1	*	*	0.3	0.5	2.9	11.1	26.6	51.8	81.2	97.6	76.3
1919	10.4	*	*	*	0.4	1.3	7.5	20.2	43.3	70.0	100.6	89.6
1920	9.1	*	*	*	0.3	1.3	5.5	16.1	37.1	68.3	93.7	97.8
1921	9.2	*	*	*	0.4	1.6	5.1	16.3	38.4	67.7	92.7	103.2
1922	9.6	*	*	0.3	0.3	1.4	5.9	17.2	39.4	67.9	96.4	118.4
1923	9.4	*	*	*	0.4	1.1	6.0	18.5	37.7	63.4	90.5	109.8
1924	9.5	*	*	0.3	0.3	1.5	6.4	19.5	37.3	65.8	76.2	99.6
1925	9.2	*	*	*	0.3	1.3	6.6	18.4	36.8	58.0	81.9	96.9
1926	9.3	*	*	*	0.4	1.4	6.6	17.8	37.1	56.8	84.3	84.4
1927	9.6	*	*	0.2	0.4	1.6	7.0	18.4	34.8	64.1	81.1	84.7
1928	9.9	*	*	0.3	0.4	1.7	7.3	19.4	38.0	62.4	80.3	75.1
1929	9.2	*	*	*	0.3	1.4	7.2	17.7	35.7	56.3	65.0	78.5
1930	9.3	*	*	0.2	0.3	1.5	6.8	17.5	36.5	56.5	66.2	66.7
1931	9.7	*	*	*	0.3	1.6	6.9	19.3	36.6	54.5	72.3	86.7
1932	9.6	*	*	*	0.3	1.4	6.8	18.3	35.6	54.6	72.1	62.9
1933	9.7	*	*	*	0.2	1.7	6.8	18.2	37.3	53.9	70.5	57.9
1934	10.4	*	*	*	0.3	1.8	7.9	20.1	37.7	54.2	72.7	61.2
1935	10.8	*	*	*	0.2	1.7	8.2	20.5	38.3	56.8	76.8	78.8
1936	11.2	*	*	0.2	0.2	1.8	8.9	20.7	40.2	56.2	74.7	78.8
1937	11.6	*	*	0.2	0.3	1.8	9.5	22.7	39.3	56.7	73.0	59.6
1938	11.4	*	*	*	0.3	2.0	8.3	22.0	38.9	56.0	69.5	68.7
1939	11.5	*	*	*	0.3	1.7	8.6	21.6	39.6	56.0	68.7	71.5

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

WHITE, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	9.2	---	---	---	---	---	---	---	---	---	---	---
1911	9.2	---	---	---	---	---	---	---	---	---	---	---
1912	8.6	---	---	---	---	---	---	---	---	---	---	---
1913	8.4	---	---	---	---	---	---	---	---	---	---	---
1914	8.1	*	*	*	0.4	2.1	8.3	17.0	25.7	55.4	76.6	79.5
1915	8.1	*	*	*	0.5	2.6	8.7	15.5	27.3	51.7	71.1	77.3
1916	7.9	*	*	*	0.4	2.7	8.0	15.1	29.4	45.9	76.0	66.1
1917	7.5	---	---	---	---	---	---	---	---	---	---	---
1918	6.3	*	*	*	0.4	1.7	5.5	10.5	22.9	40.7	68.7	77.4
1919	5.4	*	*	*	*	1.0	3.4	9.6	17.8	40.3	70.1	69.4
1920	5.1	*	*	*	0.3	0.8	2.7	7.8	18.7	38.7	68.7	63.7
1921	5.4	*	*	0.3	0.3	0.9	2.9	8.6	19.4	41.1	68.6	74.1
1922	5.4	*	*	*	*	1.1	3.3	8.9	17.9	39.5	67.5	94.2
1923	5.0	*	*	0.2	0.3	1.0	3.1	8.0	16.3	36.6	59.8	87.4
1924	5.0	*	*	*	0.3	1.0	3.4	9.4	15.8	35.2	57.1	76.3
1925	5.2	*	*	*	0.3	1.1	3.9	8.5	17.3	34.1	64.7	79.4
1926	5.2	*	*	*	0.2	1.4	3.9	8.3	17.8	31.0	60.5	73.5
1927	5.3	*	*	*	0.3	1.2	4.0	8.9	17.9	34.3	55.2	70.3
1928	5.3	*	*	*	0.3	1.2	4.7	9.4	17.1	31.5	52.1	68.1
1929	5.2	*	*	*	0.3	1.3	4.4	9.6	16.2	31.7	51.9	69.1
1930	5.2	*	*	*	0.3	1.2	4.5	9.5	15.9	29.8	50.7	64.8
1931	5.3	*	*	*	0.3	1.2	4.2	9.4	16.3	29.8	50.6	62.2
1932	5.2	*	*	0.2	*	1.1	4.0	9.2	16.2	28.7	47.8	59.0
1933	5.4	*	*	0.2	0.3	1.3	4.0	9.3	15.7	30.6	51.6	63.2
1934	5.4	*	*	*	0.3	1.4	4.1	9.5	16.1	27.4	50.9	50.6
1935	5.4	*	*	*	0.4	1.1	4.2	10.5	16.0	28.1	46.8	50.9
1936	5.7	*	*	0.2	0.3	1.3	4.6	10.0	17.1	28.0	44.6	64.7
1937	5.9	*	*	0.2	0.3	1.3	4.7	10.8	17.1	27.5	50.1	58.1
1938	5.8	*	*	*	0.4	1.2	5.2	10.2	15.5	27.5	50.1	49.6
1939	5.7	*	*	*	0.4	1.6	4.4	9.0	16.0	26.8	45.7	55.6

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ALL OTHER, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	11.0	...	...	...	...	...	...	...	...	...	...	...
1911	12.0	...	...	...	...	...	...	...	...	...	...	...
1912	13.6	...	...	...	...	...	...	...	...	...	...	...
1913	12.0	...	...	...	...	...	...	...	...	...	...	...
1914	11.8	*	*	*	*	7.9	18.8	27.0	42.7	54.5	*	*
1915	12.4	*	*	*	*	5.6	19.0	31.2	50.8	50.7	*	*
1916	8.4	*	*	*	*	5.2	13.5	19.8	41.0	48.2	*	*
1917	8.1	...	...	...	...	...	...	...	...	...	...	...
1918	8.2	*	*	*	*	3.7	9.9	21.4	45.8	56.4	49.9	*
1919	7.2	*	*	*	*	4.3	8.0	18.0	35.7	54.7	67.4	*
1920	6.8	*	*	*	1.9	3.5	7.8	22.1	24.3	44.3	*	*
1921	7.4	*	*	*	*	2.7	8.8	20.0	39.8	51.4	55.7	*
1922	6.3	*	*	*	*	2.8	9.3	19.3	28.2	35.8	39.8	*
1923	6.4	*	*	*	*	2.4	8.3	19.8	29.8	41.2	51.5	*
1924	6.8	*	*	*	1.4	3.8	7.4	22.4	29.0	41.1	41.8	*
1925	7.1	*	*	*	*	3.6	11.0	19.9	28.8	43.9	52.1	*
1926	6.5	*	*	*	*	3.2	8.4	18.8	27.6	41.6	58.9	*
1927	6.6	*	*	*	1.3	2.8	8.9	21.2	27.0	37.6	49.4	*
1928	6.2	*	*	*	1.1	3.6	9.7	19.2	25.2	31.8	32.1	*
1929	6.5	*	*	*	1.1	2.8	10.2	21.1	24.2	37.4	38.5	*
1930	6.7	*	*	*	1.3	2.9	10.1	18.3	30.5	37.4	46.8	*
1931	6.4	*	*	*	0.9	4.0	9.6	17.7	26.6	36.9	31.0	*
1932	5.5	*	*	*	1.0	2.6	8.3	15.3	25.9	31.5	39.7	*
1933	6.2	*	*	*	0.9	2.5	9.8	18.2	28.2	34.9	45.9	*
1934	6.0	*	*	*	1.2	3.3	9.3	16.5	27.9	32.4	38.3	*
1935	5.9	*	*	*	*	2.8	8.6	15.9	25.4	27.4	42.0	*
1936	6.6	*	*	*	*	3.0	10.0	18.4	27.4	34.9	33.8	*
1937	6.2	*	*	*	0.9	2.9	7.5	17.6	25.9	30.3	31.3	*
1938	5.6	*	*	*	*	2.5	7.7	15.5	21.8	27.1	39.8	*
1939	5.7	*	*	*	*	2.3	7.6	15.4	24.3	27.8	36.2	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ALL OTHER, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	13.4	...	...	...	...	...	...	...	...	...	...	...
1911	16.2	...	...	...	...	...	...	...	...	...	...	...
1912	17.9	...	...	...	...	...	...	...	...	...	...	...
1913	14.9	...	...	...	...	...	...	...	...	...	...	...
1914	16.6	*	*	*	*	8.6	25.4	37.5	59.5	79.0	*	*
1915	16.3	*	*	*	*	*	27.8	36.1	61.4	*	*	*
1916	11.2	*	*	*	*	7.3	19.0	24.3	53.4	48.4	*	*
1917	11.5	...	...	...	...	...	...	...	...	...	...	...
1918	10.8	*	*	*	*	4.8	14.0	25.1	56.8	67.2	*	*
1919	9.3	*	*	*	*	4.4	10.1	19.7	43.3	74.5	78.3	*
1920	8.0	*	*	*	*	3.5	9.1	23.6	29.5	56.3	*	*
1921	9.1	*	*	*	*	*	11.2	20.7	48.1	63.9	*	*
1922	8.1	*	*	*	*	3.4	11.6	23.4	29.8	42.5	67.3	*
1923	8.1	*	*	*	*	3.1	9.0	25.3	34.8	48.0	59.8	*
1924	8.8	*	*	*	*	4.8	9.8	27.8	35.2	47.3	63.9	*
1925	8.5	*	*	*	*	3.9	14.7	22.4	33.8	43.4	*	*
1926	8.2	*	*	*	*	3.8	9.5	23.2	31.2	54.6	86.2	*
1927	8.4	*	*	*	*	3.3	10.2	26.2	32.7	49.4	58.6	*
1928	7.6	*	*	*	*	4.1	11.5	23.3	28.7	36.7	*	*
1929	8.6	*	*	*	*	3.4	12.8	27.9	30.6	44.2	55.9	*
1930	8.3	*	*	*	*	3.2	12.4	22.3	34.9	44.5	58.1	*
1931	7.7	*	*	*	*	3.8	11.1	21.2	31.1	49.1	*	*
1932	6.5	*	*	*	*	3.0	9.7	18.1	30.9	34.6	43.0	*
1933	7.3	*	*	*	*	2.8	12.0	20.1	30.7	39.7	51.0	*
1934	7.4	*	*	*	*	3.9	11.3	19.6	35.6	37.6	51.9	*
1935	7.6	*	*	*	*	3.6	10.3	19.3	32.5	36.7	54.4	*
1936	8.8	*	*	*	*	3.3	11.9	24.0	36.3	45.3	53.1	*
1937	8.1	*	*	*	*	2.9	8.8	23.6	33.4	40.8	39.3	*
1938	7.2	*	*	*	*	2.2	9.9	18.0	27.8	34.8	53.9	*
1939	7.3	*	*	*	*	2.6	8.7	19.6	30.0	38.2	49.1	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CIRRHOSIS OF LIVER

ALL OTHER, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	8.3	...	...	...	...	...	...	...	...	...	...	...
1911	7.4	...	...	...	...	...	...	...	...	...	...	...
1912	8.8	...	...	...	...	...	...	...	...	...	...	...
1913	8.8	...	...	...	...	...	...	...	...	...	...	...
1914	6.6	*	*	*	*	*	*	*	*	*	*	*
1915	8.2	*	*	*	*	*	*	24.5	37.3	*	*	*
1916	5.6	*	*	*	*	*	7.3	14.1	25.2	47.9	*	*
1917	4.7	...	...	...	...	...	...	...	...	...	...	...
1918	5.5	*	*	*	*	*	5.3	16.4	31.6	43.9	*	*
1919	5.1	*	*	*	*	4.3	5.8	15.6	25.7	31.2	*	*
1920	5.5	*	*	*	*	3.4	6.5	20.0	17.5	30.0	*	*
1921	5.6	*	*	*	*	*	6.2	19.1	28.9	36.3	*	*
1922	4.6	*	*	*	*	*	7.0	14.1	26.1	27.7	*	*
1923	4.7	*	*	*	*	*	7.6	12.9	23.3	32.9	*	*
1924	4.7	*	*	*	*	2.8	4.9	15.7	20.8	33.5	*	*
1925	5.7	*	*	*	*	3.4	7.2	16.8	22.4	44.4	57.4	*
1926	4.7	*	*	*	*	2.6	7.4	13.7	22.9	26.1	*	*
1927	4.8	*	*	*	*	2.3	7.6	15.2	19.5	23.7	*	*
1928	4.8	*	*	*	1.7	3.1	7.8	14.5	20.5	26.0	*	*
1929	4.3	*	*	*	*	2.2	7.7	13.2	15.7	29.7	*	*
1930	5.1	*	*	*	*	2.7	7.8	13.7	24.8	29.4	*	*
1931	5.0	*	*	*	*	4.1	8.2	13.6	20.8	23.2	*	*
1932	4.4	*	*	*	*	2.3	6.9	12.1	19.5	28.1	*	*
1933	5.0	*	*	*	*	2.3	7.8	15.0	24.9	29.5	41.0	*
1934	4.5	*	*	*	1.8	2.7	7.3	13.1	18.2	26.5	*	*
1935	4.2	*	*	*	*	2.1	6.9	12.1	16.7	17.2	*	*
1936	4.5	*	*	*	*	2.7	8.1	12.3	16.5	23.5	*	*
1937	4.3	*	*	*	*	2.9	6.3	11.0	17.0	18.9	*	*
1938	4.1	*	*	*	*	2.7	5.5	12.9	14.6	18.7	*	*
1939	4.2	*	*	*	*	2.0	6.6	11.0	17.8	16.6	*	*

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

NOTES:

For 1932-34, deaths of Mexicans are included with total deaths for the white population; but by age, are included with deaths for the nonwhite population. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes.....	30	41	55
Male.....	19	21	30
Female.....	11	20	25

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ICD Revision	Years in use	ICD Nos.
Fifth	1939	130
Fourth	1930-1938	130
Third	1921-1929	128
Second	1910-1920	119
First	1900-1909	119

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	7.7	20.3	7.7	3.1	4.7	6.6	7.9	10.0	13.1	18.9	28.0	*
1901	8.6	21.2	9.6	4.2	5.0	6.6	8.8	11.7	17.3	17.2	28.5	*
1902	9.9	25.7	10.1	4.5	5.6	8.9	9.6	12.1	17.0	27.0	36.9	*
1903	10.1	23.6	8.3	4.3	5.5	8.8	10.9	12.5	19.3	27.1	38.5	60.0
1904	10.7	24.3	9.7	4.6	5.3	8.0	11.2	14.0	20.6	31.3	48.6	70.0
1905	10.6	25.1	7.8	3.8	5.5	9.3	10.5	14.8	21.9	31.7	40.3	43.9
1906	9.7	28.6	8.0	4.0	4.7	7.7	9.8	12.2	19.4	30.1	44.1	59.4
1907	10.3	32.2	9.6	3.5	4.9	7.7	10.9	14.1	23.0	27.8	45.2	62.0
1908	9.9	27.7	9.6	4.2	4.4	8.0	10.2	13.2	19.6	27.8	43.1	74.3
1909	9.6	29.6	10.1	3.4	4.0	6.8	8.9	12.5	19.7	32.4	47.8	78.0
1910	10.2	31.3	9.3	4.3	4.2	7.0	10.1	13.6	21.4	32.9	48.4	88.7
1911	9.6	33.0	8.2	3.3	4.1	6.5	9.6	13.4	19.5	30.5	51.8	79.6
1912	10.0	28.5	8.3	3.2	4.2	6.6	10.4	14.3	19.8	35.0	58.4	99.3
1913	9.9	26.3	8.7	3.6	4.2	7.1	10.5	13.2	20.1	30.0	52.6	92.4
1914	8.5	24.5	7.5	3.1	3.8	5.8	8.9	11.4	16.3	26.1	48.5	86.5
1915	8.5	25.6	8.3	3.0	3.7	5.8	8.7	11.3	15.1	27.3	47.3	85.4
1916	8.0	24.7	6.8	2.8	3.9	5.6	7.7	11.2	16.6	23.9	42.6	69.3
1917	7.9	25.8	8.8	3.2	3.6	5.2	7.5	10.4	16.1	23.7	43.2	71.4
1918	7.2	22.1	7.2	2.8	3.5	5.2	6.8	8.8	13.0	21.7	38.1	66.1
1919	6.4	21.2	5.4	2.8	2.9	4.0	5.9	8.3	11.9	19.5	38.2	75.5
1920	6.3	22.5	5.4	2.6	3.1	4.0	5.7	7.6	10.8	20.3	42.0	81.8
1921	6.3	22.8	6.8	3.1	2.9	3.7	5.3	7.9	10.3	19.1	36.4	65.2
1922	6.0	22.9	6.3	2.5	2.7	3.2	5.6	7.8	11.0	17.9	35.0	76.5
1923	6.0	18.2	6.0	2.5	2.8	3.7	5.6	8.1	10.3	19.6	33.8	75.6
1924	5.9	20.5	5.5	2.5	2.6	3.4	5.6	8.0	10.7	16.5	36.1	83.3
1925	5.4	18.4	5.6	2.1	2.5	3.1	5.0	7.1	9.4	15.7	31.8	65.8
1926	5.0	16.5	5.1	2.1	2.3	3.0	5.2	6.6	8.7	12.7	26.7	62.8
1927	4.7	15.3	4.4	2.0	2.0	3.0	4.9	6.3	8.4	12.4	23.7	56.0
1928	4.8	15.3	4.3	1.9	2.1	3.1	4.6	6.9	8.3	12.5	27.0	61.9
1929	4.6	13.1	3.9	1.9	2.1	2.9	4.8	6.4	7.5	12.4	26.3	54.6
1930	4.3	12.4	4.0	1.7	2.0	2.7	4.4	5.8	8.1	11.7	25.4	50.8
1931	3.8	11.1	3.7	1.5	1.8	2.3	3.8	5.4	7.0	9.2	20.8	38.7
1932	3.6	9.8	3.2	1.4	1.5	2.3	3.7	4.9	6.6	9.4	17.0	49.7
1933	3.8	10.8	3.6	1.5	1.8	2.3	3.6	5.0	6.7	10.7	20.3	40.1
1934	3.6	9.1	3.1	1.5	1.5	2.1	3.4	4.8	6.3	9.8	20.7	45.0
1935	3.5	9.2	3.0	1.4	1.6	2.3	3.4	4.5	5.9	9.9	19.5	36.4
1936	3.2	8.8	2.9	1.2	1.5	2.3	3.0	4.0	5.8	8.4	17.6	36.2
1937	3.1	9.3	3.0	1.2	1.4	2.0	2.9	4.0	5.5	7.7	17.1	29.5
1938	2.8	7.6	2.8	1.2	1.3	1.7	2.6	3.5	4.4	6.6	14.3	28.9
1939	2.5	7.2	2.4	1.0	1.2	1.6	2.4	2.9	4.3	6.5	12.7	32.1

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .  
 Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	7.4	19.0	6.7	2.8	3.8	5.6	7.3	9.5	14.4	24.5	35.8	*
1901	9.0	26.3	10.0	4.6	4.4	6.4	9.0	12.8	18.7	19.5	30.7	*
1902	10.4	32.9	12.2	4.9	5.6	7.8	9.3	13.2	18.7	28.2	47.5	*
1903	10.4	27.7	8.9	4.8	5.2	7.0	11.4	12.1	20.7	30.8	52.2	*
1904	10.9	25.5	9.8	4.0	4.4	7.0	12.1	14.9	22.3	38.0	52.3	*
1905	11.0	32.2	7.7	3.5	5.2	8.2	9.7	16.7	24.9	39.0	43.2	*
1906	10.0	32.4	7.5	4.2	4.5	7.3	10.3	12.2	21.6	33.0	53.0	*
1907	11.1	38.0	11.3	3.5	4.3	6.6	11.8	14.8	25.9	33.9	60.4	84.5
1908	10.5	32.3	11.3	4.1	3.7	7.0	10.8	14.3	22.7	33.9	50.0	79.0
1909	10.1	37.4	11.0	3.6	3.5	6.2	9.1	13.6	21.8	33.9	59.4	95.4
1910	10.5	38.1	8.8	4.2	3.9	6.7	9.7	14.9	23.4	34.4	57.2	104.4
1911	9.9	36.9	8.5	3.2	3.6	6.2	9.7	13.9	20.5	33.4	65.6	83.8
1912	10.7	31.5	9.2	3.2	3.8	6.8	10.7	16.3	21.6	36.2	70.2	123.1
1913	10.3	28.6	9.8	3.5	4.0	6.9	10.5	14.0	21.9	33.0	62.8	114.3
1914	9.0	29.9	8.3	2.9	3.2	5.5	9.1	12.8	17.2	30.0	55.8	111.7
1915	9.0	30.6	9.1	2.9	3.4	5.9	8.5	11.4	16.5	30.9	60.8	111.6
1916	8.6	28.2	7.4	2.9	3.6	5.6	8.5	12.1	17.9	26.2	50.6	81.0
1917	8.6	30.4	7.3	3.3	3.8	5.4	8.1	10.1	17.8	27.4	53.1	93.4
1918	7.7	26.2	7.8	3.1	3.6	5.2	6.8	9.2	13.9	23.1	46.1	74.8
1919	6.8	23.0	5.5	3.0	2.9	3.9	5.8	8.6	13.4	21.8	45.4	96.3
1920	6.6	28.4	5.5	2.7	2.9	3.7	5.3	7.6	11.0	23.0	50.5	108.1
1921	6.6	27.0	7.2	3.2	2.9	3.5	5.2	8.0	10.6	20.5	43.1	100.0
1922	6.5	28.3	7.1	2.5	2.4	3.1	5.4	8.7	11.6	21.1	39.1	111.6
1923	6.3	20.5	6.1	2.7	2.5	3.5	5.3	8.6	11.3	21.6	37.6	100.4
1924	6.3	25.2	6.0	2.3	2.4	3.4	5.4	8.9	11.2	19.2	42.3	113.3
1925	5.8	21.5	6.0	1.9	2.4	3.0	5.0	7.9	10.9	18.6	36.1	74.6
1926	5.3	19.9	4.9	2.1	2.3	2.8	5.3	6.9	10.3	13.7	31.6	80.8
1927	5.0	18.0	4.8	2.0	1.9	2.6	5.2	6.9	9.9	14.4	24.9	67.6
1928	5.3	19.8	5.0	1.9	2.3	3.1	4.8	8.1	9.5	14.2	33.8	69.9
1929	4.9	14.9	4.1	2.0	2.0	2.8	4.9	6.9	8.7	15.0	29.1	65.2
1930	4.7	13.6	4.7	1.7	1.9	2.7	4.3	6.0	9.7	14.1	30.1	56.5
1931	4.1	13.1	4.2	1.5	1.7	2.2	4.0	6.1	7.3	10.6	23.5	51.5
1932	3.9	11.4	3.4	1.5	1.7	2.5	3.8	5.2	7.3	10.4	21.5	61.7
1933	4.1	13.6	3.9	1.5	1.9	2.3	3.6	5.5	7.7	12.3	23.2	55.2
1934	3.8	10.8	3.0	1.6	1.5	2.1	3.5	5.2	7.3	11.0	24.9	43.0
1935	3.9	10.3	3.5	1.4	1.8	2.2	3.6	4.9	6.7	11.5	24.8	43.1
1936	3.5	10.9	3.4	1.3	1.5	2.1	3.0	4.6	6.4	9.6	21.1	46.1
1937	3.5	11.3	3.3	1.2	1.5	2.2	2.9	4.6	6.5	9.5	20.4	35.1
1938	3.0	9.3	3.0	1.2	1.3	1.8	2.6	3.5	5.1	7.6	14.7	37.9
1939	2.7	8.7	2.9	1.1	1.0	1.7	2.4	3.2	4.7	6.5	14.6	44.4

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	8.0	21.7	8.8	3.4	5.5	7.6	8.5	10.6	12.3	13.6	21.0	*
1901	8.2	15.9	9.1	3.9	5.5	6.9	8.7	10.6	15.9	15.0	26.4	*
1902	9.4	18.4	8.1	4.0	5.6	10.0	9.9	11.0	15.3	25.9	27.4	*
1903	9.8	19.4	7.7	3.9	5.8	10.6	10.4	12.9	17.9	23.6	26.2	*
1904	10.5	23.0	9.6	5.2	6.2	8.9	10.1	13.1	18.9	24.9	45.4	*
1905	10.2	17.9	7.8	4.2	5.7	10.5	11.3	12.9	18.8	24.8	37.7	*
1906	9.4	24.8	8.6	3.8	5.0	8.1	9.2	12.3	17.2	27.2	36.0	57.4
1907	9.6	26.2	7.8	3.4	5.4	8.9	9.9	13.3	20.1	21.9	31.5	*
1908	9.3	22.9	8.0	4.4	5.1	9.0	9.4	12.1	16.2	21.8	36.8	70.6
1909	9.0	21.5	9.3	3.2	4.5	7.3	8.8	11.3	17.4	30.9	37.2	64.4
1910	9.9	24.3	9.8	4.4	4.5	7.4	10.5	12.2	19.2	31.3	40.3	76.4
1911	9.2	29.0	7.8	3.4	4.5	6.8	9.5	12.7	18.6	27.6	39.1	76.3
1912	9.4	25.3	7.3	3.2	4.6	6.4	10.0	12.1	17.9	33.7	47.7	80.6
1913	9.4	24.0	7.5	3.8	4.3	7.3	10.6	12.4	18.0	26.9	43.3	75.4
1914	8.1	18.9	6.7	3.2	4.4	6.2	8.7	9.9	15.4	22.2	41.9	67.0
1915	8.0	20.5	7.5	3.1	4.1	5.7	8.9	11.3	13.5	23.7	34.9	65.2
1916	7.4	21.1	6.1	2.7	4.2	5.5	6.9	10.1	15.1	21.5	35.4	60.3
1917	7.2	21.0	6.3	3.2	3.4	4.9	6.8	10.8	14.2	19.8	34.2	54.7
1918	6.7	18.0	6.6	2.5	3.4	5.2	6.9	8.4	12.1	20.3	30.8	59.5
1919	6.0	19.4	5.2	2.5	2.9	4.2	6.0	7.9	10.3	17.0	31.6	59.9
1920	6.1	16.5	5.4	2.6	3.3	4.3	6.1	7.7	10.6	17.5	34.2	62.1
1921	6.0	18.6	6.4	3.1	3.0	3.9	5.5	7.9	9.9	17.6	30.2	38.9
1922	5.6	17.4	5.5	2.5	2.9	3.3	5.9	6.9	10.5	14.7	31.3	49.8
1923	5.8	15.7	5.9	2.3	3.0	3.9	5.9	7.6	9.2	17.5	30.3	56.7
1924	5.5	15.8	4.9	2.6	2.9	3.4	5.8	7.1	10.2	13.7	30.4	60.3
1925	5.0	15.1	5.3	2.2	2.7	3.2	5.1	6.2	7.7	12.7	27.8	59.0
1926	4.7	12.9	5.2	2.1	2.4	3.1	5.1	6.3	6.8	11.6	22.2	49.0
1927	4.4	12.5	4.1	2.0	2.1	3.4	4.6	5.7	6.9	10.2	22.6	47.0
1928	4.2	10.6	3.6	1.9	2.0	3.2	4.3	5.7	6.9	10.6	20.6	55.7
1929	4.2	11.3	3.8	1.8	2.1	3.0	4.6	5.9	6.2	9.8	23.6	46.6
1930	4.0	11.3	3.3	1.6	2.1	2.7	4.4	5.6	6.4	9.1	21.0	46.4
1931	3.5	9.1	3.1	1.5	1.8	2.5	3.6	4.6	6.6	7.9	18.2	29.0
1932	3.2	8.0	3.0	1.3	1.4	2.1	3.7	4.6	5.8	8.4	12.7	40.7
1933	3.4	8.0	3.3	1.6	1.6	2.3	3.5	4.4	5.7	9.1	17.6	28.7
1934	3.3	7.4	3.2	1.4	1.6	2.2	3.3	4.3	5.2	8.7	16.8	46.6
1935	3.1	8.0	2.5	1.5	1.3	2.3	3.2	4.1	4.9	8.2	14.6	31.4
1936	3.0	6.7	2.5	1.0	1.6	2.5	3.0	3.4	5.2	7.2	14.4	28.8
1937	2.8	7.2	2.6	1.1	1.4	1.8	3.0	3.4	4.5	6.0	14.1	25.3
1938	2.5	5.7	2.6	1.2	1.3	1.6	2.5	3.5	3.6	5.7	13.8	22.0
1939	2.4	5.7	1.9	0.9	1.3	1.5	2.4	2.5	3.8	6.4	11.0	22.8

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator. .

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	9.9	---	---	---	---	---	---	---	---	---	---	---
1911	9.3	---	---	---	---	---	---	---	---	---	---	---
1912	9.7	---	---	---	---	---	---	---	---	---	---	---
1913	9.5	---	---	---	---	---	---	---	---	---	---	---
1914	8.1	23.9	7.2	3.0	3.6	5.5	8.2	10.6	15.0	25.3	48.7	85.8
1915	8.0	25.1	7.6	2.9	3.5	5.4	7.8	10.3	13.8	25.5	46.8	83.0
1916	7.4	24.4	5.9	2.7	3.6	5.1	6.7	10.0	15.0	22.6	42.6	70.3
1917	7.2	---	---	---	---	---	---	---	---	---	---	---
1918	6.4	20.9	6.1	2.6	3.1	4.3	5.8	7.5	11.7	19.7	37.4	67.1
1919	5.7	20.1	4.9	2.6	2.6	3.4	5.2	7.1	10.3	17.2	36.5	71.7
1920	5.6	20.9	5.2	2.5	2.6	3.3	4.5	6.2	9.1	18.5	40.1	80.2
1921	5.5	21.2	6.0	2.9	2.4	2.9	4.3	6.5	9.1	17.9	35.4	63.8
1922	5.1	21.3	5.4	2.4	2.2	2.5	4.4	5.8	9.2	16.2	31.6	69.6
1923	5.2	16.6	5.2	2.4	2.2	3.0	4.4	6.1	8.4	17.6	31.4	72.6
1924	4.9	18.6	4.7	2.3	2.1	2.6	4.3	5.6	8.6	14.9	33.7	72.4
1925	4.5	17.2	4.7	1.9	1.9	2.4	3.8	5.3	7.5	13.7	30.5	58.2
1926	4.1	15.5	4.5	1.9	1.9	2.1	3.8	4.8	7.1	11.1	25.6	57.1
1927	3.9	14.2	3.8	1.8	1.6	2.2	3.7	4.5	6.9	11.1	22.2	51.4
1928	3.8	13.6	3.5	1.6	1.6	2.2	3.3	4.9	6.3	11.0	25.3	59.3
1929	3.6	11.2	3.2	1.7	1.5	1.9	3.2	4.5	5.6	10.8	25.2	51.3
1930	3.4	10.8	3.3	1.5	1.4	1.9	3.0	3.8	6.0	9.9	23.9	44.2
1931	3.0	10.1	2.9	1.3	1.4	1.6	2.8	3.8	5.4	8.0	18.7	36.3
1932	3.0	8.8	2.8	1.3	1.2	1.8	2.7	3.7	5.2	8.5	16.1	47.6
1933	3.1	9.9	3.1	1.3	1.4	1.6	2.6	3.6	5.1	9.1	19.3	38.7
1934	2.8	7.2	2.7	1.3	1.2	1.4	2.2	3.3	4.9	8.2	19.2	45.1
1935	2.8	8.8	2.7	1.3	1.3	1.6	2.3	3.3	4.4	8.4	18.1	33.9
1936	2.6	7.9	2.5	1.0	1.1	1.6	2.0	2.8	4.5	6.8	16.3	36.6
1937	2.5	8.2	2.4	1.1	1.1	1.4	2.0	2.8	4.4	6.6	15.8	28.6
1938	2.1	6.7	2.3	1.0	1.0	1.1	1.6	2.4	3.2	5.5	13.3	28.0
1939	2.0	6.7	2.0	0.9	0.9	1.2	1.6	2.0	3.1	5.3	12.0	31.4

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	10.2	---	---	---	---	---	---	---	---	---	---	---
1911	9.5	---	---	---	---	---	---	---	---	---	---	---
1912	10.3	---	---	---	---	---	---	---	---	---	---	---
1913	10.0	---	---	---	---	---	---	---	---	---	---	---
1914	8.6	29.4	8.0	2.9	3.0	5.1	8.5	12.0	16.0	28.7	55.7	109.3
1915	8.4	30.9	8.2	2.8	3.2	5.5	7.6	10.6	15.2	28.7	59.8	107.4
1916	8.0	27.4	6.2	2.9	3.5	5.1	7.4	10.9	16.3	25.1	50.2	81.2
1917	7.9	---	---	---	---	---	---	---	---	---	---	---
1918	7.0	24.8	6.7	2.9	3.2	4.5	6.0	7.9	12.7	20.8	45.3	76.3
1919	6.2	22.9	5.0	2.8	2.6	3.6	5.2	7.4	11.7	19.0	43.0	97.0
1920	5.8	26.5	5.4	2.5	2.4	3.1	4.3	6.2	9.2	20.3	48.0	106.7
1921	5.8	25.4	6.5	3.0	2.3	2.8	4.3	6.6	9.3	19.2	41.8	94.4
1922	5.5	26.2	6.0	2.3	2.1	2.5	4.4	6.5	9.5	19.2	35.1	99.7
1923	5.5	18.3	5.3	2.6	2.1	3.0	4.4	7.0	9.4	19.1	34.5	92.0
1924	5.3	22.5	5.1	2.1	1.9	2.8	4.3	6.6	9.1	17.1	38.1	92.9
1925	4.8	20.3	4.8	1.7	1.9	2.3	3.7	5.9	8.8	16.2	34.6	65.9
1926	4.5	19.3	4.4	2.0	2.0	2.2	4.1	5.1	8.5	12.0	29.8	68.3
1927	4.3	16.6	4.0	1.8	1.6	2.1	4.2	5.2	8.2	13.0	23.8	63.5
1928	4.3	17.9	4.0	1.7	1.9	2.3	3.7	5.8	7.2	12.3	31.2	66.6
1929	3.9	13.0	3.2	1.8	1.5	2.0	3.5	5.0	6.6	12.7	28.1	63.0
1930	3.7	12.2	3.7	1.4	1.4	2.0	3.0	3.9	7.3	11.7	28.1	47.1
1931	3.3	11.5	3.4	1.3	1.4	1.7	3.1	4.5	5.7	9.1	20.3	45.7
1932	3.4	10.1	3.1	1.4	1.4	2.0	3.2	4.1	5.8	9.5	20.1	60.1
1933	3.4	12.8	3.3	1.3	1.5	1.8	2.7	4.1	5.9	10.3	22.2	52.7
1934	3.1	8.5	2.6	1.4	1.2	1.5	2.5	3.8	5.9	9.0	23.0	41.9
1935	3.2	9.4	3.1	1.2	1.6	1.6	2.5	3.7	5.2	9.9	23.0	39.8
1936	2.8	9.5	2.9	1.1	1.2	1.5	2.2	3.4	5.0	7.9	19.0	45.7
1937	2.8	10.1	2.6	1.1	1.2	1.6	2.0	3.3	5.4	8.0	18.4	32.8
1938	2.3	8.7	2.5	1.0	1.2	1.3	1.8	2.3	3.8	6.1	13.5	37.7
1939	2.2	8.3	2.4	1.0	0.9	1.2	1.8	2.3	3.5	5.0	13.5	42.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.  
 Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).  
 Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	9.5	---	---	---	---	---	---	---	---	---	---	---
1911	8.9	---	---	---	---	---	---	---	---	---	---	---
1912	9.0	---	---	---	---	---	---	---	---	---	---	---
1913	8.9	---	---	---	---	---	---	---	---	---	---	---
1914	7.6	18.3	6.4	3.1	4.3	5.9	7.9	9.1	13.9	21.9	42.3	67.5
1915	7.5	19.1	7.0	3.0	3.8	5.4	8.1	10.0	12.3	22.4	34.9	64.2
1916	6.9	21.3	5.7	2.5	3.6	5.1	6.0	9.0	13.5	20.1	35.8	62.0
1917	6.6	---	---	---	---	---	---	---	---	---	---	---
1918	5.8	16.9	5.4	2.2	2.9	4.1	5.6	7.0	10.6	18.6	30.2	60.1
1919	5.2	17.1	4.8	2.3	2.5	3.1	5.2	6.7	8.8	15.3	30.6	52.6
1920	5.4	15.1	5.0	2.5	2.8	3.6	4.8	6.2	9.0	16.6	32.9	60.3
1921	5.2	16.9	5.5	2.9	2.5	2.9	4.2	6.4	8.9	16.6	29.6	40.4
1922	4.7	16.1	4.8	2.4	2.3	2.6	4.4	5.0	8.9	13.2	28.5	46.5
1923	4.8	14.8	5.0	2.3	2.3	2.9	4.4	5.2	7.4	16.1	28.5	57.5
1924	4.5	14.6	4.3	2.5	2.2	2.4	4.2	4.6	8.2	12.6	29.7	56.5
1925	4.1	14.1	4.5	2.1	2.0	2.5	3.9	4.6	6.2	11.1	26.8	52.2
1926	3.7	11.6	4.5	1.9	1.9	2.0	3.5	4.5	5.5	10.1	21.7	48.3
1927	3.4	11.6	3.6	1.8	1.5	2.3	3.1	3.7	5.5	9.0	20.8	42.0
1928	3.3	9.2	3.0	1.6	1.4	2.1	2.9	3.9	5.2	9.7	19.8	53.8
1929	3.2	9.4	3.3	1.7	1.5	1.9	2.8	3.9	4.6	8.9	22.4	42.4
1930	3.1	9.4	2.9	1.5	1.4	1.8	2.9	3.6	4.5	8.0	19.9	42.0
1931	2.8	8.6	2.4	1.4	1.5	1.5	2.5	3.1	5.2	6.8	17.2	29.3
1932	2.6	7.3	2.4	1.2	1.1	1.6	2.3	3.4	4.6	7.6	12.4	38.2
1933	2.7	6.9	2.9	1.3	1.3	1.5	2.4	3.1	4.3	8.0	16.5	28.0
1934	2.5	5.9	2.9	1.3	1.1	1.3	1.9	2.8	3.8	7.4	15.7	47.4
1935	2.4	8.1	2.4	1.3	1.0	1.5	2.1	2.9	3.6	6.8	13.5	29.4
1936	2.2	6.2	2.0	0.9	1.1	1.7	1.8	2.2	3.9	5.8	13.7	29.7
1937	2.1	6.3	2.2	1.1	0.9	1.2	1.9	2.2	3.3	5.1	13.5	25.3
1938	1.9	4.7	2.1	1.0	0.8	0.9	1.4	2.4	2.6	5.0	13.1	20.6
1939	1.8	5.1	1.6	0.7	1.0	1.2	1.5	1.6	2.7	5.6	10.6	22.7

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	22.0	---	---	---	---	---	---	---	---	---	---	---
1911	19.5	---	---	---	---	---	---	---	---	---	---	---
1912	21.0	---	---	---	---	---	---	---	---	---	---	---
1913	18.9	---	---	---	---	---	---	---	---	---	---	---
1914	17.9	38.1	15.7	4.8	7.9	13.6	22.8	30.2	53.8	51.4	*	*
1915	20.9	37.7	26.6	5.2	8.8	13.0	26.4	34.7	53.1	84.5	*	*
1916	16.6	29.0	18.8	3.8	8.6	12.9	22.3	30.4	50.5	52.2	*	*
1917	16.8	---	---	---	---	---	---	---	---	---	---	---
1918	17.4	37.4	21.5	5.2	8.4	16.2	20.1	28.8	38.7	60.9	52.1	*
1919	13.5	32.9	10.4	4.3	6.1	11.3	13.7	23.2	39.3	57.2	67.4	*
1920	14.5	40.8	7.8	4.0	7.8	11.5	18.3	25.9	39.8	52.4	75.4	*
1921	14.5	40.9	15.4	5.1	7.8	12.7	17.2	26.1	29.4	38.5	54.0	*
1922	14.7	38.8	15.1	3.3	6.6	9.6	18.3	30.3	38.0	44.6	88.4	140.6
1923	14.7	33.9	14.2	3.2	7.5	10.3	18.0	31.3	37.5	52.4	72.9	*
1924	15.9	39.7	12.5	4.0	7.6	11.3	19.7	36.2	41.3	44.1	75.2	190.5
1925	14.8	29.8	14.9	3.9	7.8	10.5	18.8	28.7	36.5	51.9	53.5	141.8
1926	13.8	25.6	10.7	3.7	5.8	11.1	20.6	28.5	32.3	42.1	45.5	120.3
1927	12.8	25.4	9.9	3.5	5.7	10.6	18.2	26.9	30.6	35.8	48.1	100.7
1928	13.4	29.6	10.7	4.1	6.2	11.0	16.8	27.8	35.6	37.1	52.0	84.4
1929	13.6	29.8	9.4	3.8	6.6	11.0	20.4	26.6	33.3	39.9	44.0	84.3
1930	13.1	25.4	9.4	3.0	6.4	9.4	17.9	26.2	37.5	41.8	50.1	110.9
1931	10.7	19.4	9.6	2.9	4.3	8.8	13.9	21.6	28.2	30.9	55.5	*
1932	8.9	17.4	6.7	2.4	4.0	6.8	13.2	16.7	25.3	23.7	31.4	*
1933	10.1	17.6	7.2	3.2	4.7	7.8	12.5	18.8	28.2	35.5	37.3	*
1934	10.2	22.8	5.7	3.0	4.3	8.1	14.3	19.4	25.2	34.5	45.8	*
1935	9.5	12.1	5.3	2.7	3.7	8.0	13.3	17.7	25.1	31.1	42.9	*
1936	9.4	15.8	6.2	2.3	4.7	7.8	12.4	16.7	24.4	31.1	40.1	*
1937	8.9	17.4	7.1	2.0	4.4	7.1	11.7	17.1	20.9	23.6	39.1	*
1938	8.3	13.7	6.3	2.6	3.9	6.8	10.8	15.6	19.9	21.2	30.5	*
1939	7.2	10.7	4.8	2.1	3.2	5.5	9.1	12.9	19.7	21.7	24.7	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	21.1	...	...	...	...	...	...	...	...	...	...	...
1911	20.8	...	...	...	...	...	...	...	...	...	...	...
1912	21.4	...	...	...	...	...	...	...	...	...	...	...
1913	17.5	...	...	...	...	...	...	...	...	...	...	...
1914	18.3	*	*	*	8.0	13.7	20.8	29.2	48.2	67.3	*	*
1915	21.7	*	34.5	*	*	13.6	26.9	28.7	53.0	98.2	*	*
1916	17.5	40.3	24.8	*	5.1	13.5	24.2	30.8	48.7	50.2	*	*
1917	17.8	...	...	...	...	...	...	...	...	...	...	...
1918	17.1	44.1	22.3	5.0	8.0	14.0	17.4	26.7	35.8	65.8	*	*
1919	13.2	*	11.1	4.5	5.5	7.3	12.7	22.4	39.2	65.3	85.4	*
1920	15.1	50.2	6.7	4.7	7.9	10.6	16.4	23.6	39.0	66.6	92.7	*
1921	14.5	44.9	15.1	5.0	7.9	11.1	15.4	23.9	29.7	40.3	*	*
1922	15.7	48.3	18.1	4.0	5.5	9.0	16.5	31.7	39.3	48.1	99.4	*
1923	14.3	43.4	14.0	3.9	5.5	8.1	14.8	26.0	36.9	60.0	85.4	*
1924	16.6	52.4	14.6	4.7	7.0	9.4	16.8	34.3	39.4	51.9	108.4	334.1
1925	15.9	34.3	17.0	4.1	6.4	10.3	19.1	29.9	39.6	59.1	60.5	*
1926	13.3	26.3	9.4	3.2	4.5	8.8	18.6	27.2	35.2	42.7	62.0	*
1927	12.3	30.9	11.4	3.5	4.6	7.7	15.9	25.4	32.4	38.1	*	*
1928	14.4	37.0	13.0	3.4	5.7	10.1	15.7	30.4	38.0	45.1	71.4	*
1929	14.3	32.3	11.1	4.3	6.6	10.0	19.1	26.5	35.4	51.8	44.7	*
1930	14.0	24.6	12.1	3.6	5.7	8.7	17.9	26.1	39.4	52.7	62.6	*
1931	11.4	25.8	11.1	3.3	4.3	7.1	13.8	22.5	27.9	34.8	76.8	*
1932	8.9	21.5	6.0	2.5	4.0	6.8	10.1	16.3	25.6	24.4	45.2	*
1933	10.7	19.5	8.0	3.1	5.0	6.7	11.9	19.8	29.9	42.7	39.3	*
1934	10.4	28.1	6.2	3.5	3.4	7.1	13.1	19.3	24.9	39.9	55.7	*
1935	10.1	16.9	6.7	2.4	3.7	7.6	14.0	18.0	25.0	32.9	54.4	*
1936	9.7	21.6	6.9	2.6	4.0	7.1	10.8	17.4	24.6	33.1	54.9	*
1937	9.5	20.7	8.4	2.2	3.6	7.5	11.2	18.3	20.2	27.8	53.6	*
1938	8.6	*	6.7	2.8	3.0	6.7	10.0	16.4	21.7	26.2	34.8	*
1939	7.5	*	5.9	2.1	2.6	6.4	7.8	13.5	19.8	25.0	*	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACUTE NEPHRITIS AND NEPHRITIS WITH EDEMA, INCLUDING NEPHROSIS

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	23.0	---	---	---	---	---	---	---	---	---	---	---
1911	18.1	---	---	---	---	---	---	---	---	---	---	---
1912	20.6	---	---	---	---	---	---	---	---	---	---	---
1913	20.4	---	---	---	---	---	---	---	---	---	---	---
1914	17.5	*	*	*	7.8	13.5	25.4	31.6	60.8	*	*	*
1915	20.1	*	*	*	11.2	12.4	25.7	42.7	53.3	69.0	*	*
1916	15.7	*	12.8	4.8	11.8	12.3	20.2	30.0	52.7	54.5	*	*
1917	15.7	---	---	---	---	---	---	---	---	---	---	---
1918	17.7	30.9	20.8	5.4	8.8	18.3	23.0	31.6	42.5	55.2	*	*
1919	13.8	42.3	9.7	4.1	6.7	15.0	14.7	24.4	39.3	47.5	*	*
1920	13.9	31.7	8.9	3.3	7.7	12.3	20.4	29.1	40.8	35.5	*	*
1921	14.5	36.9	15.7	5.1	7.8	14.2	19.0	29.0	28.9	36.3	**	*
1922	13.7	29.4	12.2	2.7	7.5	10.1	20.2	29.7	36.2	40.4	77.2	*
1923	15.2	24.5	14.4	2.5	9.3	12.4	21.3	38.0	38.2	43.1	60.3	*
1924	15.2	27.3	10.3	3.3	8.1	13.0	22.8	38.5	43.8	34.6	*	*
1925	13.7	25.4	12.8	3.6	9.1	10.7	18.4	27.1	32.6	43.3	*	*
1926	14.2	24.8	12.0	4.1	7.1	13.3	22.8	30.1	28.6	41.4	*	*
1927	13.2	20.2	8.5	3.6	6.7	13.3	20.5	28.7	28.2	32.9	52.9	*
1928	12.5	22.4	8.4	4.7	6.5	11.9	17.8	24.7	32.4	27.8	*	*
1929	12.9	27.4	7.8	3.3	6.6	11.9	21.7	26.7	30.6	26.2	43.3	*
1930	12.1	26.1	6.7	2.4	7.0	10.0	18.0	26.2	34.9	29.4	*	*
1931	10.0	*	8.1	2.6	4.3	10.4	14.0	20.7	28.6	26.4	*	*
1932	9.0	*	7.4	2.2	4.0	6.9	16.2	17.2	25.0	22.8	*	*
1933	9.5	*	6.3	3.4	4.5	8.8	13.1	17.8	26.0	27.5	*	*
1934	10.1	17.5	5.3	2.6	5.0	9.0	15.5	19.6	25.6	28.4	36.4	*
1935	8.9	*	3.9	3.1	3.6	8.3	12.5	17.3	25.2	29.1	*	*
1936	9.1	*	5.4	1.9	5.4	8.4	13.9	15.8	24.1	29.0	*	*
1937	8.3	*	5.8	1.8	5.0	6.7	12.2	15.8	21.8	18.9	*	*
1938	8.0	*	5.9	2.5	4.7	7.0	11.5	14.7	17.8	15.7	*	*
1939	6.8	*	3.7	2.0	3.7	4.6	10.3	12.3	19.6	18.0	*	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

NOTES:  
Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ICD Revision	Years in use	ICD Nos.
Fifth	1939	158-161
Fourth	1930-1938	158-161
Third	1921-1929	160-162
Second	1910-1920	151, 152
First	1900-1909	151, 152

ALL RACES, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	62.6	2,856.9	..	..	..	..	..	..	..	..	..	..
1901	58.5	2,666.8	..	..	..	..	..	..	..	..	..	..
1902	61.4	2,801.3	..	..	..	..	..	..	..	..	..	..
1903	62.5	2,855.8	..	..	..	..	..	..	..	..	..	..
1904	65.4	2,995.3	..	..	..	..	..	..	..	..	..	..
1905	72.9	3,348.5	..	..	..	..	..	..	..	..	..	..
1906	77.4	3,482.5	..	..	..	..	..	..	..	..	..	..
1907	78.1	3,530.7	..	..	..	..	..	..	..	..	..	..
1908	75.9	3,450.9	..	..	..	..	..	..	..	..	..	..
1909	72.5	3,335.4	..	..	..	..	..	..	..	..	..	..
1910	73.0	3,380.4	..	..	..	..	..	..	..	..	..	..
1911	78.3	3,651.9	..	..	..	..	..	..	..	..	..	..
1912	80.6	3,734.6	..	..	..	..	..	..	..	..	..	..
1913	83.2	3,849.8	..	..	..	..	..	..	..	..	..	..
1914	79.0	3,684.3	..	..	..	..	..	..	..	..	..	..
1915	76.6	3,612.3	..	..	..	..	..	..	..	..	..	..
1916	77.5	3,598.5	..	..	..	..	..	..	..	..	..	..
1917	75.5	3,519.3	..	..	..	..	..	..	..	..	..	..
1918	76.4	3,569.5	..	..	..	..	..	..	..	..	..	..
1919	65.5	3,132.1	..	..	..	..	..	..	..	..	..	..
1920	69.2	3,217.6	..	..	..	..	..	..	..	..	..	..
1921	67.7	3,066.0	..	..	..	..	..	..	..	..	..	..
1922	62.8	2,901.6	..	..	..	..	..	..	..	..	..	..
1923	62.4	2,996.3	..	..	..	..	..	..	..	..	..	..
1924	62.6	3,018.7	..	..	..	..	..	..	..	..	..	..
1925	59.2	2,882.0	..	..	..	..	..	..	..	..	..	..
1926	57.5	2,918.5	..	..	..	..	..	..	..	..	..	..
1927	54.5	2,794.8	..	..	..	..	..	..	..	..	..	..
1928	53.4	2,891.1	..	..	..	..	..	..	..	..	..	..
1929	50.6	2,848.8	..	..	..	..	..	..	..	..	..	..
1930	49.6	2,803.6	..	..	..	..	..	..	..	..	..	..
1931	45.4	2,638.5	..	..	..	..	..	..	..	..	..	..
1932	42.8	2,614.3	..	..	..	..	..	..	..	..	..	..
1933	41.0	2,607.3	..	..	..	..	..	..	..	..	..	..
1934	43.0	2,787.9	..	..	..	..	..	..	..	..	..	..
1935	40.2	2,597.7	..	..	..	..	..	..	..	..	..	..
1936	40.4	2,656.2	..	..	..	..	..	..	..	..	..	..
1937	40.0	2,631.5	..	..	..	..	..	..	..	..	..	..
1938	39.4	2,541.9	..	..	..	..	..	..	..	..	..	..
1939	38.9	2,509.9	..	..	..	..	..	..	..	..	..	..

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ALL RACES, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	72.2	3,263.3	..	..	..	..	..	..	..	..	..	..
1901	67.7	3,060.6	..	..	..	..	..	..	..	..	..	..
1902	70.8	3,206.8	..	..	..	..	..	..	..	..	..	..
1903	71.2	3,233.6	..	..	..	..	..	..	..	..	..	..
1904	74.1	3,375.0	..	..	..	..	..	..	..	..	..	..
1905	83.0	3,791.9	..	..	..	..	..	..	..	..	..	..
1906	87.0	3,940.0	..	..	..	..	..	..	..	..	..	..
1907	86.9	3,957.2	..	..	..	..	..	..	..	..	..	..
1908	84.9	3,908.2	..	..	..	..	..	..	..	..	..	..
1909	80.7	3,761.2	..	..	..	..	..	..	..	..	..	..
1910	81.3	3,821.6	..	..	..	..	..	..	..	..	..	..
1911	86.3	4,031.9	..	..	..	..	..	..	..	..	..	..
1912	89.6	4,208.9	..	..	..	..	..	..	..	..	..	..
1913	92.7	4,349.4	..	..	..	..	..	..	..	..	..	..
1914	87.8	4,148.7	..	..	..	..	..	..	..	..	..	..
1915	85.6	4,087.9	..	..	..	..	..	..	..	..	..	..
1916	87.7	4,115.1	..	..	..	..	..	..	..	..	..	..
1917	85.3	4,004.6	..	..	..	..	..	..	..	..	..	..
1918	87.8	4,067.4	..	..	..	..	..	..	..	..	..	..
1919	74.1	3,551.0	..	..	..	..	..	..	..	..	..	..
1920	78.1	3,651.1	..	..	..	..	..	..	..	..	..	..
1921	77.0	3,486.8	..	..	..	..	..	..	..	..	..	..
1922	71.3	3,287.0	..	..	..	..	..	..	..	..	..	..
1923	70.5	3,384.5	..	..	..	..	..	..	..	..	..	..
1924	70.9	3,420.8	..	..	..	..	..	..	..	..	..	..
1925	67.3	3,273.0	..	..	..	..	..	..	..	..	..	..
1926	65.5	3,316.4	..	..	..	..	..	..	..	..	..	..
1927	62.2	3,196.3	..	..	..	..	..	..	..	..	..	..
1928	61.2	3,308.9	..	..	..	..	..	..	..	..	..	..
1929	57.3	3,219.2	..	..	..	..	..	..	..	..	..	..
1930	56.3	3,166.3	..	..	..	..	..	..	..	..	..	..
1931	52.2	3,020.0	..	..	..	..	..	..	..	..	..	..
1932	49.1	2,978.7	..	..	..	..	..	..	..	..	..	..
1933	46.7	2,958.2	..	..	..	..	..	..	..	..	..	..
1934	49.1	3,168.6	..	..	..	..	..	..	..	..	..	..
1935	46.4	2,978.6	..	..	..	..	..	..	..	..	..	..
1936	46.4	3,036.1	..	..	..	..	..	..	..	..	..	..
1937	45.8	2,995.5	..	..	..	..	..	..	..	..	..	..
1938	45.4	2,913.6	..	..	..	..	..	..	..	..	..	..
1939	44.7	2,861.5	..	..	..	..	..	..	..	..	..	..

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ALL RACES, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	53.1	2,442.2	..	..	..	..	..	..	..	..	..	..
1901	49.2	2,265.1	..	..	..	..	..	..	..	..	..	..
1902	51.9	2,387.7	..	..	..	..	..	..	..	..	..	..
1903	53.7	2,470.4	..	..	..	..	..	..	..	..	..	..
1904	56.5	2,607.7	..	..	..	..	..	..	..	..	..	..
1905	62.6	2,895.7	..	..	..	..	..	..	..	..	..	..
1906	67.4	3,034.0	..	..	..	..	..	..	..	..	..	..
1907	68.9	3,093.7	..	..	..	..	..	..	..	..	..	..
1908	66.4	2,981.7	..	..	..	..	..	..	..	..	..	..
1909	63.8	2,898.3	..	..	..	..	..	..	..	..	..	..
1910	64.2	2,927.3	..	..	..	..	..	..	..	..	..	..
1911	69.7	3,150.6	..	..	..	..	..	..	..	..	..	..
1912	71.1	3,247.8	..	..	..	..	..	..	..	..	..	..
1913	73.1	3,337.3	..	..	..	..	..	..	..	..	..	..
1914	69.7	3,208.3	..	..	..	..	..	..	..	..	..	..
1915	67.1	3,125.0	..	..	..	..	..	..	..	..	..	..
1916	66.8	3,069.5	..	..	..	..	..	..	..	..	..	..
1917	65.3	3,022.4	..	..	..	..	..	..	..	..	..	..
1918	64.9	3,059.9	..	..	..	..	..	..	..	..	..	..
1919	56.7	2,703.7	..	..	..	..	..	..	..	..	..	..
1920	60.0	2,773.5	..	..	..	..	..	..	..	..	..	..
1921	58.1	2,631.2	..	..	..	..	..	..	..	..	..	..
1922	54.1	2,504.3	..	..	..	..	..	..	..	..	..	..
1923	54.0	2,506.3	..	..	..	..	..	..	..	..	..	..
1924	54.0	2,604.4	..	..	..	..	..	..	..	..	..	..
1925	50.8	2,478.1	..	..	..	..	..	..	..	..	..	..
1926	49.4	2,508.0	..	..	..	..	..	..	..	..	..	..
1927	46.6	2,383.2	..	..	..	..	..	..	..	..	..	..
1928	45.5	2,451.4	..	..	..	..	..	..	..	..	..	..
1929	43.8	2,467.9	..	..	..	..	..	..	..	..	..	..
1930	42.8	2,429.1	..	..	..	..	..	..	..	..	..	..
1931	38.4	2,244.6	..	..	..	..	..	..	..	..	..	..
1932	36.4	2,237.8	..	..	..	..	..	..	..	..	..	..
1933	35.1	2,246.3	..	..	..	..	..	..	..	..	..	..
1934	36.8	2,306.1	..	..	..	..	..	..	..	..	..	..
1935	34.0	2,206.2	..	..	..	..	..	..	..	..	..	..
1936	34.3	2,266.3	..	..	..	..	..	..	..	..	..	..
1937	34.1	2,257.6	..	..	..	..	..	..	..	..	..	..
1938	33.2	2,160.1	..	..	..	..	..	..	..	..	..	..
1939	33.0	2,148.8	..	..	..	..	..	..	..	..	..	..

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

WHITE, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	72.0	3,321.0	*	...	...	...	...	...	...	...	...	...
1911	77.3	3,539.6	*	...	...	...	...	...	...	...	...	...
1912	79.9	3,683.0	*	...	...	...	...	...	...	...	...	...
1913	82.3	3,802.1	*	...	...	...	...	...	...	...	...	...
1914	78.0	3,628.6	*	...	...	...	...	...	...	...	...	...
1915	75.6	3,555.5	*	...	...	...	...	...	...	...	...	...
1916	75.4	3,521.9	*	...	...	...	...	...	...	...	...	...
1917	73.5	3,443.2	*	...	...	...	...	...	...	...	...	...
1918	74.7	3,502.3	*	...	...	...	...	...	...	...	...	...
1919	63.9	3,069.5	*	...	...	...	...	...	...	...	...	...
1920	67.7	3,136.6	*	...	...	...	...	...	...	...	...	...
1921	66.1	2,995.3	*	...	...	...	...	...	...	...	...	...
1922	61.6	2,845.3	*	...	...	...	...	...	...	...	...	...
1923	61.0	2,934.4	*	...	...	...	...	...	...	...	...	...
1924	60.5	2,921.0	*	...	...	...	...	...	...	...	...	...
1925	57.1	2,786.0	*	...	...	...	...	...	...	...	...	...
1926	55.6	2,837.6	*	...	...	...	...	...	...	...	...	...
1927	52.7	2,721.3	*	...	...	...	...	...	...	...	...	...
1928	51.7	2,811.5	*	...	...	...	...	...	...	...	...	...
1929	49.0	2,760.4	*	...	...	...	...	...	...	...	...	...
1930	47.8	2,731.3	*	...	...	...	...	...	...	...	...	...
1931	43.7	2,574.6	*	...	...	...	...	...	...	...	...	...
1932	40.9	2,540.8	*	...	...	...	...	...	...	...	...	...
1933	38.9	2,530.4	*	...	...	...	...	...	...	...	...	...
1934	40.9	2,708.5	*	...	...	...	...	...	...	...	...	...
1935	38.4	2,531.7	*	...	...	...	...	...	...	...	...	...
1936	38.4	2,578.6	*	...	...	...	...	...	...	...	...	...
1937	37.8	2,541.5	*	...	...	...	...	...	...	...	...	...
1938	37.4	2,466.4	*	...	...	...	...	...	...	...	...	...
1939	36.7	2,413.0	*	...	...	...	...	...	...	...	...	...

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

WHITE, MALE												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	80.3	3,755.1	...	...	...	...	...	...	...	...	...	...
1911	85.6	3,971.7	...	...	...	...	...	...	...	...	...	...
1912	89.0	4,158.2	...	...	...	...	...	...	...	...	...	...
1913	92.1	4,303.9	...	...	...	...	...	...	...	...	...	...
1914	86.8	4,089.4	...	...	...	...	...	...	...	...	...	...
1915	84.7	4,031.0	...	...	...	...	...	...	...	...	...	...
1916	85.5	4,029.8	...	...	...	...	...	...	...	...	...	...
1917	83.3	3,929.1	...	...	...	...	...	...	...	...	...	...
1918	86.3	4,001.2	...	...	...	...	...	...	...	...	...	...
1919	72.4	3,479.9	...	...	...	...	...	...	...	...	...	...
1920	76.5	3,556.9	...	...	...	...	...	...	...	...	...	...
1921	75.5	3,414.1	...	...	...	...	...	...	...	...	...	...
1922	70.3	3,233.1	...	...	...	...	...	...	...	...	...	...
1923	69.2	3,319.6	...	...	...	...	...	...	...	...	...	...
1924	68.7	3,113.0	...	...	...	...	...	...	...	...	...	...
1925	65.2	3,170.3	...	...	...	...	...	...	...	...	...	...
1926	63.4	3,227.3	...	...	...	...	...	...	...	...	...	...
1927	60.4	3,118.8	...	...	...	...	...	...	...	...	...	...
1928	59.4	3,224.4	...	...	...	...	...	...	...	...	...	...
1929	55.5	3,120.6	...	...	...	...	...	...	...	...	...	...
1930	54.3	3,089.3	...	...	...	...	...	...	...	...	...	...
1931	50.5	2,958.9	...	...	...	...	...	...	...	...	...	...
1932	47.0	2,900.7	...	...	...	...	...	...	...	...	...	...
1933	44.6	2,882.9	...	...	...	...	...	...	...	...	...	...
1934	46.8	3,086.4	...	...	...	...	...	...	...	...	...	...
1935	44.5	2,915.4	...	...	...	...	...	...	...	...	...	...
1936	44.3	2,951.1	...	...	...	...	...	...	...	...	...	...
1937	43.5	2,902.5	...	...	...	...	...	...	...	...	...	...
1938	43.4	2,836.1	...	...	...	...	...	...	...	...	...	...
1939	42.2	2,753.1	...	...	...	...	...	...	...	...	...	...

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1938 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

WHITE, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	63.3	2,875.2	---	---	---	---	---	---	---	---	---	---
1911	68.6	3,095.9	---	---	---	---	---	---	---	---	---	---
1912	70.2	3,195.1	---	---	---	---	---	---	---	---	---	---
1913	72.1	3,286.9	---	---	---	---	---	---	---	---	---	---
1914	68.6	3,155.6	---	---	---	---	---	---	---	---	---	---
1915	66.0	3,067.6	---	---	---	---	---	---	---	---	---	---
1916	64.8	3,000.2	---	---	---	---	---	---	---	---	---	---
1917	63.2	2,944.0	---	---	---	---	---	---	---	---	---	---
1918	63.1	2,989.9	---	---	---	---	---	---	---	---	---	---
1919	55.2	2,648.2	---	---	---	---	---	---	---	---	---	---
1920	58.6	2,704.2	---	---	---	---	---	---	---	---	---	---
1921	56.5	2,560.8	---	---	---	---	---	---	---	---	---	---
1922	52.7	2,443.5	---	---	---	---	---	---	---	---	---	---
1923	52.6	2,535.6	---	---	---	---	---	---	---	---	---	---
1924	52.0	2,515.3	---	---	---	---	---	---	---	---	---	---
1925	48.8	2,387.3	---	---	---	---	---	---	---	---	---	---
1926	47.6	2,433.6	---	---	---	---	---	---	---	---	---	---
1927	44.7	2,311.0	---	---	---	---	---	---	---	---	---	---
1928	43.7	2,383.8	---	---	---	---	---	---	---	---	---	---
1929	42.2	2,387.3	---	---	---	---	---	---	---	---	---	---
1930	41.1	2,359.8	---	---	---	---	---	---	---	---	---	---
1931	36.7	2,175.3	---	---	---	---	---	---	---	---	---	---
1932	34.6	2,166.8	---	---	---	---	---	---	---	---	---	---
1933	33.1	2,155.6	---	---	---	---	---	---	---	---	---	---
1934	34.7	2,317.0	---	---	---	---	---	---	---	---	---	---
1935	32.1	2,134.8	---	---	---	---	---	---	---	---	---	---
1936	32.5	2,193.7	---	---	---	---	---	---	---	---	---	---
1937	32.0	2,168.0	---	---	---	---	---	---	---	---	---	---
1938	31.4	2,083.8	---	---	---	---	---	---	---	---	---	---
1939	31.0	2,061.0	---	---	---	---	---	---	---	---	---	---

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	108.8	5,973.1	---	---	---	---	---	---	---	---	---	---
1911	107.0	5,805.9	---	---	---	---	---	---	---	---	---	---
1912	102.1	5,575.2	---	---	---	---	---	---	---	---	---	---
1913	101.6	4,979.2	---	---	---	---	---	---	---	---	---	---
1914	101.4	5,015.8	---	---	---	---	---	---	---	---	---	---
1915	99.9	4,963.1	---	---	---	---	---	---	---	---	---	---
1916	107.5	4,627.1	---	---	---	---	---	---	---	---	---	---
1917	102.3	4,486.4	---	---	---	---	---	---	---	---	---	---
1918	97.6	4,388.9	---	---	---	---	---	---	---	---	---	---
1919	82.7	3,773.1	---	---	---	---	---	---	---	---	---	---
1920	86.0	4,139.7	---	---	---	---	---	---	---	---	---	---
1921	84.6	3,843.4	---	---	---	---	---	---	---	---	---	---
1922	74.7	3,445.6	---	---	---	---	---	---	---	---	---	---
1923	75.7	3,605.1	---	---	---	---	---	---	---	---	---	---
1924	83.7	3,986.3	---	---	---	---	---	---	---	---	---	---
1925	80.5	3,856.3	---	---	---	---	---	---	---	---	---	---
1926	77.1	3,701.7	---	---	---	---	---	---	---	---	---	---
1927	72.2	3,473.1	---	---	---	---	---	---	---	---	---	---
1928	69.2	3,572.4	---	---	---	---	---	---	---	---	---	---
1929	65.6	3,631.4	---	---	---	---	---	---	---	---	---	---
1930	66.5	3,387.2	---	---	---	---	---	---	---	---	---	---
1931	60.7	3,148.1	---	---	---	---	---	---	---	---	---	---
1932	60.4	3,177.6	---	---	---	---	---	---	---	---	---	---
1933	59.1	3,157.1	---	---	---	---	---	---	---	---	---	---
1934	62.0	3,360.7	---	---	---	---	---	---	---	---	---	---
1935	56.8	3,075.2	---	---	---	---	---	---	---	---	---	---
1936	57.8	3,224.0	---	---	---	---	---	---	---	---	---	---
1937	58.9	3,289.9	---	---	---	---	---	---	---	---	---	---
1938	56.4	3,095.7	---	---	---	---	---	---	---	---	---	---
1939	58.5	3,222.9	---	---	---	---	---	---	---	---	---	---

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1938 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	116.2	6,728.6	---	---	---	---	---	---	---	---	---	---
1911	108.3	6,240.9	---	---	---	---	---	---	---	---	---	---
1912	105.0	6,040.4	---	---	---	---	---	---	---	---	---	---
1913	106.8	5,447.2	---	---	---	---	---	---	---	---	---	---
1914	108.5	5,591.3	---	---	---	---	---	---	---	---	---	---
1915	105.4	5,465.9	---	---	---	---	---	---	---	---	---	---
1916	120.0	5,283.8	---	---	---	---	---	---	---	---	---	---
1917	111.3	4,985.0	---	---	---	---	---	---	---	---	---	---
1918	106.9	4,893.1	---	---	---	---	---	---	---	---	---	---
1919	92.8	4,293.8	---	---	---	---	---	---	---	---	---	---
1920	96.5	4,754.3	---	---	---	---	---	---	---	---	---	---
1921	93.3	4,307.5	---	---	---	---	---	---	---	---	---	---
1922	81.7	3,823.1	---	---	---	---	---	---	---	---	---	---
1923	83.0	4,039.5	---	---	---	---	---	---	---	---	---	---
1924	93.8	4,515.8	---	---	---	---	---	---	---	---	---	---
1925	89.7	4,342.0	---	---	---	---	---	---	---	---	---	---
1926	86.6	4,203.2	---	---	---	---	---	---	---	---	---	---
1927	80.2	3,935.7	---	---	---	---	---	---	---	---	---	---
1928	77.5	4,056.1	---	---	---	---	---	---	---	---	---	---
1929	73.4	4,124.7	---	---	---	---	---	---	---	---	---	---
1930	74.6	3,803.4	---	---	---	---	---	---	---	---	---	---
1931	67.8	3,520.5	---	---	---	---	---	---	---	---	---	---
1932	68.3	3,592.9	---	---	---	---	---	---	---	---	---	---
1933	65.7	3,520.4	---	---	---	---	---	---	---	---	---	---
1934	69.7	3,777.8	---	---	---	---	---	---	---	---	---	---
1935	63.7	3,447.7	---	---	---	---	---	---	---	---	---	---
1936	65.9	3,675.9	---	---	---	---	---	---	---	---	---	---
1937	66.1	3,696.2	---	---	---	---	---	---	---	---	---	---
1938	63.7	3,499.4	---	---	---	---	---	---	---	---	---	---
1939	67.0	3,682.7	---	---	---	---	---	---	---	---	---	---

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1938 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

CERTAIN DISEASES OF EARLY INFANCY

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	100.3	5,198.1	---	---	---	---	---	---	---	---	---	---
1911	105.5	5,488.4	---	---	---	---	---	---	---	---	---	---
1912	98.8	5,108.6	---	---	---	---	---	---	---	---	---	---
1913	95.9	4,514.3	---	---	---	---	---	---	---	---	---	---
1914	93.8	4,445.4	---	---	---	---	---	---	---	---	---	---
1915	93.9	4,466.0	---	---	---	---	---	---	---	---	---	---
1916	94.7	3,980.3	---	---	---	---	---	---	---	---	---	---
1917	93.0	3,995.9	---	---	---	---	---	---	---	---	---	---
1918	88.1	3,893.8	---	---	---	---	---	---	---	---	---	---
1919	72.5	3,260.0	---	---	---	---	---	---	---	---	---	---
1920	75.3	3,541.5	---	---	---	---	---	---	---	---	---	---
1921	75.8	3,385.9	---	---	---	---	---	---	---	---	---	---
1922	67.6	3,075.5	---	---	---	---	---	---	---	---	---	---
1923	67.6	3,178.0	---	---	---	---	---	---	---	---	---	---
1924	73.6	3,465.3	---	---	---	---	---	---	---	---	---	---
1925	71.2	3,377.3	---	---	---	---	---	---	---	---	---	---
1926	67.7	3,209.0	---	---	---	---	---	---	---	---	---	---
1927	64.2	3,027.1	---	---	---	---	---	---	---	---	---	---
1928	60.9	3,103.6	---	---	---	---	---	---	---	---	---	---
1929	57.9	3,156.3	---	---	---	---	---	---	---	---	---	---
1930	58.5	2,975.7	---	---	---	---	---	---	---	---	---	---
1931	53.8	2,781.5	---	---	---	---	---	---	---	---	---	---
1932	52.7	2,768.1	---	---	---	---	---	---	---	---	---	---
1933	52.6	2,819.5	---	---	---	---	---	---	---	---	---	---
1934	54.5	2,951.3	---	---	---	---	---	---	---	---	---	---
1935	50.0	2,709.6	---	---	---	---	---	---	---	---	---	---
1936	50.0	2,782.7	---	---	---	---	---	---	---	---	---	---
1937	51.8	2,894.0	---	---	---	---	---	---	---	---	---	---
1938	49.2	2,702.7	---	---	---	---	---	---	---	---	---	---
1939	50.3	2,774.9	---	---	---	---	---	---	---	---	---	---

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1938 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ICD Revision	Years in use	ICD Nos.
Fifth	1939	163, 164
Fourth	1930-1938	163-171
Third	1921-1929	165-174
Second	1910-1920	155-163
First	1900-1909	155-163

ALL RACES, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	10.2	..	..	*	6.8	11.2	15.8	19.9	23.5	26.1	29.2	*
1901	10.4	..	..	*	6.8	11.4	15.4	20.5	25.6	25.6	29.2	*
1902	10.3	..	..	*	7.4	10.3	16.1	20.0	24.3	25.5	30.3	*
1903	11.3	..	..	*	6.6	13.3	18.9	22.9	23.0	27.9	24.4	*
1904	12.2	..	..	*	8.5	14.0	17.2	25.3	27.9	27.7	32.2	*
1905	13.5	..	..	*	9.4	14.9	19.6	26.5	30.3	33.0	37.9	57.1
1906	12.8	..	..	*	8.0	14.1	19.8	26.5	30.5	31.4	31.0	*
1907	14.5	..	..	*	10.8	17.0	21.1	28.4	33.6	35.5	32.7	*
1908	16.8	..	..	*	11.3	19.4	25.2	36.0	40.2	35.7	34.3	33.2
1909	16.0	..	..	0.3	10.5	18.1	23.8	33.7	37.5	35.9	34.0	30.5
1910	15.3	..	..	0.3	10.7	17.7	22.4	29.9	37.0	35.3	38.7	33.4
1911	16.0	..	..	0.2	10.9	18.8	24.5	32.1	37.3	35.6	34.8	27.8
1912	15.6	..	..	0.3	9.8	18.4	23.6	31.1	38.7	34.2	38.1	28.9
1913	15.4	..	..	0.2	10.4	18.9	22.9	30.1	36.7	35.1	31.6	27.1
1914	16.1	..	..	0.3	10.8	18.9	23.6	32.5	40.0	35.7	32.7	31.3
1915	16.2	..	..	0.3	9.8	18.6	23.9	32.2	41.3	39.7	35.1	36.8
1916	13.7	..	..	0.2	8.7	15.5	20.1	26.9	35.9	36.3	34.4	29.9
1917	13.0	..	..	0.3	8.3	16.7	18.9	25.0	30.1	34.7	32.5	27.0
1918	12.3	..	..	0.2	7.3	15.8	19.2	22.0	29.0	30.0	30.4	31.5
1919	11.5	..	..	0.2	6.6	13.7	18.1	21.6	27.7	29.4	32.2	39.5
1920	10.2	..	..	0.2	6.2	11.5	15.7	20.4	24.2	27.4	28.2	32.5
1921	12.4	..	..	0.3	6.9	13.3	19.0	26.2	31.3	32.5	32.3	35.0
1922	11.7	..	..	0.2	6.4	12.1	18.0	24.8	29.7	33.8	32.8	35.1
1923	11.5	..	..	0.2	5.7	11.9	17.8	24.0	29.5	31.2	35.0	35.4
1924	11.9	..	..	0.2	6.3	11.9	18.6	24.9	31.3	33.4	30.8	31.0
1925	12.0	..	..	0.2	5.9	12.2	18.0	25.2	30.3	35.8	31.7	32.4
1926	12.6	..	..	0.2	6.5	12.2	19.2	26.4	31.9	36.6	36.1	32.1
1927	13.2	..	..	0.3	6.4	13.1	19.6	27.7	33.7	37.4	35.9	34.0
1928	13.5	..	..	0.2	6.7	13.4	19.4	29.0	34.3	40.4	39.4	29.9
1929	13.9	..	..	0.2	7.1	13.6	20.3	28.7	35.0	41.0	40.8	25.3
1930	15.6	..	..	0.2	7.5	14.9	22.7	33.0	41.2	42.8	40.1	29.2
1931	16.8	..	..	0.2	7.2	15.1	22.9	36.0	46.9	48.6	42.1	33.9
1932	17.4	..	..	0.2	7.3	15.5	22.8	36.4	49.1	51.3	45.5	27.5
1933	15.9	..	..	0.2	7.2	13.9	21.3	33.2	43.0	46.1	45.1	31.7
1934	14.9	..	..	0.2	7.4	14.5	19.9	29.6	37.2	40.9	41.2	25.6
1935	14.3	..	..	0.2	7.1	14.3	19.0	28.6	34.9	35.8	38.7	27.3
1936	14.3	..	..	0.3	6.8	14.5	19.7	27.6	34.1	34.7	38.0	26.8
1937	15.0	..	..	0.3	6.9	14.4	20.7	30.1	35.5	34.5	38.6	27.7
1938	15.3	..	..	0.2	6.7	14.5	21.0	30.7	37.5	34.4	36.1	27.7
1939	14.1	..	..	0.3	5.9	12.9	18.9	28.2	34.4	34.0	36.0	24.1

... Category not applicable

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ALL RACES, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	15.7	...	...	*	7.7	15.2	24.5	32.9	40.0	44.6	56.4	*
1901	15.5	...	...	*	8.0	15.6	20.7	32.1	43.9	42.5	58.3	*
1902	15.4	...	...	*	8.4	13.9	24.7	30.4	39.4	42.1	59.1	*
1903	17.2	...	...	...	8.1	18.0	28.7	37.2	36.9	47.7	43.8	*
1904	18.5	...	...	*	9.7	18.9	26.3	40.1	45.8	48.9	57.6	*
1905	20.2	...	...	*	10.6	20.1	29.5	41.3	51.4	58.1	67.0	108.8
1906	19.6	...	...	*	9.8	20.0	30.1	41.3	50.0	55.0	57.6	*
1907	21.9	...	...	*	13.5	24.0	30.8	44.1	54.4	61.5	59.4	*
1908	25.5	...	...	*	13.8	27.0	37.9	56.2	66.8	59.1	59.8	66.8
1909	24.2	...	...	*	12.3	25.0	36.7	52.3	60.9	62.2	62.4	64.5
1910	23.0	...	...	*	12.9	24.9	32.6	46.4	61.2	58.2	68.9	61.7
1911	23.9	...	...	*	13.0	26.1	36.8	49.4	60.7	58.6	61.9	54.5
1912	23.4	...	...	*	11.5	25.7	34.9	48.1	62.2	58.8	68.3	57.5
1913	23.2	...	...	*	12.0	26.7	33.9	47.4	59.9	59.7	55.8	44.6
1914	24.1	...	...	*	12.5	27.0	34.8	50.4	62.9	59.5	58.2	62.3
1915	24.3	...	...	*	11.5	26.1	35.0	49.2	66.5	65.4	61.8	75.6
1916	20.7	...	...	*	10.4	21.7	30.1	40.1	57.3	61.0	63.2	64.1
1917	19.2	...	...	0.3	9.9	23.4	26.8	36.5	47.8	57.0	58.1	48.3
1918	18.2	...	...	*	9.2	23.4	28.0	31.6	43.7	48.5	52.1	60.7
1919	16.5	...	...	0.3	7.5	19.3	25.7	29.8	41.2	46.9	56.5	80.0
1920	14.5	...	...	0.2	7.1	15.1	21.5	28.7	36.6	44.6	49.9	63.5
1921	18.9	...	...	0.4	8.3	18.7	28.5	39.9	49.6	54.4	57.9	73.0
1922	17.6	...	...	0.2	7.1	17.0	26.2	37.4	46.6	56.7	57.6	70.9
1923	17.0	...	...	0.3	6.0	16.1	26.0	36.3	45.4	52.7	61.6	67.8
1924	18.1	...	...	0.2	7.4	16.4	27.5	38.0	49.7	56.1	56.0	61.6
1925	18.0	...	...	0.3	7.2	16.3	26.1	37.7	48.3	60.4	59.0	63.9
1926	18.8	...	...	0.3	7.4	16.1	28.0	39.6	50.3	61.9	64.4	62.6
1927	19.9	...	...	0.3	7.9	17.9	28.6	42.4	53.7	63.6	64.9	67.6
1928	20.7	...	...	0.2	7.8	18.4	28.6	45.1	55.4	68.4	73.1	57.1
1929	21.1	...	...	0.3	8.6	18.4	30.0	43.9	54.8	69.4	73.5	51.2
1930	24.1	...	...	0.3	9.1	20.6	34.2	51.9	66.6	72.3	72.6	58.3
1931	26.2	...	...	0.3	8.7	20.9	34.4	56.9	77.0	84.3	76.1	70.4
1932	27.4	...	...	0.3	9.0	21.8	34.7	58.0	81.4	87.6	84.9	53.4
1933	24.9	...	...	0.2	8.8	19.4	31.6	52.8	70.2	79.9	83.8	64.5
1934	22.8	...	...	0.3	9.2	20.3	29.2	45.0	60.1	70.6	75.1	51.3
1935	21.7	...	...	0.3	9.0	19.7	27.5	44.1	55.5	60.1	71.9	57.0
1936	21.7	...	...	0.4	8.8	19.7	29.0	41.9	54.4	58.2	69.3	53.9
1937	22.8	...	...	0.5	9.0	20.4	30.4	45.5	56.1	58.1	70.3	57.1
1938	23.5	...	...	0.3	9.0	20.9	31.1	47.0	60.2	58.0	66.0	54.6
1939	21.7	...	...	0.4	8.0	18.8	27.5	42.1	54.8	56.3	67.3	47.7

... Category not applicable

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ALL RACES, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	4.7	...	...	*	6.0	7.1	8.5	6.3	7.3	8.6	*	*
1901	5.3	...	...	*	5.6	7.1	9.7	8.5	7.7	9.5	*	*
1902	5.2	...	...	*	6.3	6.6	6.9	9.0	9.6	9.9	*	*
1903	5.4	...	...	*	5.1	8.5	8.4	7.8	9.3	9.2	*	*
1904	6.0	...	...	*	7.3	8.9	7.5	9.7	10.2	7.7	*	*
1905	6.8	...	...	*	8.3	9.6	9.2	10.8	9.4	9.3	*	*
1906	5.8	...	...	*	6.2	7.8	8.4	10.3	10.4	8.4	*	*
1907	6.9	...	...	*	8.0	9.5	10.4	11.0	12.1	10.2	*	*
1908	7.8	...	...	*	8.8	11.1	11.1	13.5	12.1	12.4	10.9	*
1909	7.3	...	...	*	8.8	10.6	9.5	13.0	12.7	9.8	8.0	*
1910	7.2	...	...	*	8.4	9.6	11.0	11.4	11.1	12.5	11.1	*
1911	7.5	...	...	*	8.8	10.8	10.8	12.7	12.0	12.3	9.8	*
1912	7.3	...	...	*	8.1	10.4	11.0	12.0	13.2	9.5	10.4	*
1913	7.1	...	...	*	8.7	10.3	10.7	10.5	11.6	10.3	9.6	*
1914	7.7	...	...	*	9.2	10.2	11.2	12.2	15.0	11.6	9.3	*
1915	7.6	...	...	*	8.0	10.4	11.5	13.0	13.7	13.6	10.8	*
1916	6.4	...	...	*	6.9	8.9	9.2	11.9	12.5	11.1	8.2	*
1917	6.6	...	...	*	6.7	9.7	10.3	11.9	10.6	11.8	9.3	*
1918	6.2	...	...	*	5.7	8.7	9.6	11.0	12.9	11.1	10.6	*
1919	6.3	...	...	*	5.7	8.2	9.8	12.2	12.7	11.4	10.0	*
1920	5.7	...	...	*	5.3	7.7	9.3	11.1	10.6	9.5	8.3	*
1921	5.7	...	...	*	5.5	7.8	8.7	10.8	11.3	9.8	8.8	*
1922	5.7	...	...	*	5.7	7.2	9.0	10.8	11.4	10.1	10.1	*
1923	5.8	...	...	*	5.4	7.7	8.9	10.6	12.2	9.0	10.5	*
1924	5.6	...	...	*	5.2	7.4	8.9	10.5	11.4	9.7	7.5	*
1925	5.8	...	...	*	4.7	8.0	9.3	11.6	10.7	10.3	6.5	*
1926	6.2	...	...	*	5.7	8.2	9.7	12.0	11.8	10.3	10.0	*
1927	6.2	...	...	0.2	5.1	8.4	10.1	11.7	11.8	10.4	8.9	*
1928	6.2	...	...	*	5.7	8.4	9.6	11.6	11.3	11.4	7.9	*
1929	6.6	...	...	*	5.7	8.7	10.1	12.0	13.4	11.7	10.2	*
1930	6.9	...	...	*	6.1	9.1	10.7	12.3	13.8	12.2	9.5	*
1931	7.1	...	...	*	5.6	9.4	10.9	13.2	14.5	11.8	10.2	*
1932	7.1	...	...	*	5.7	9.3	10.5	13.0	14.3	14.0	8.5	*
1933	6.8	...	...	*	5.7	8.4	10.5	12.0	13.9	11.4	8.7	*
1934	6.8	...	...	*	5.5	8.7	10.4	12.9	12.7	10.4	9.6	*
1935	6.8	...	...	*	5.2	9.1	10.2	12.0	13.1	11.2	7.7	*
1936	6.8	...	...	0.2	4.8	9.3	10.1	12.4	12.6	11.0	9.1	*
1937	7.0	...	...	*	4.9	8.5	10.7	13.7	13.6	10.8	9.3	*
1938	6.9	...	...	*	4.4	8.2	10.8	13.3	13.4	10.8	8.5	*
1939	6.5	...	...	*	3.8	7.2	10.1	13.4	12.9	11.7	7.4	*

... Category not applicable

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

WHITE, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	15.4	...	...	...	...	...	...	...	...	...	...	...
1911	16.1	...	...	...	...	...	...	...	...	...	...	...
1912	15.8	...	...	...	...	...	...	...	...	...	...	...
1913	15.6	...	...	...	...	...	...	...	...	...	...	...
1914	16.4	...	0.3	11.0	19.0	24.0	33.5	40.8	36.3	33.4	32.8	...
1915	16.5	...	0.3	9.9	18.6	24.3	33.0	42.2	40.6	35.8	38.6	...
1916	14.3	...	0.2	9.0	15.9	20.9	27.9	37.3	37.6	35.5	32.0	...
1917	13.6	...	...	...	...	...	...	...	...	...	...	...
1918	12.8	...	0.3	7.5	16.4	19.9	23.0	30.0	30.7	31.3	34.2	...
1919	12.2	...	0.2	6.9	14.2	19.1	22.9	29.0	30.9	33.9	43.5	...
1920	10.8	...	0.2	6.5	12.0	16.7	21.5	25.2	28.5	29.4	35.7	...
1921	13.1	...	0.3	7.1	13.8	20.0	27.8	32.8	33.7	33.8	37.6	...
1922	12.5	...	0.2	6.7	12.7	18.9	26.4	31.3	35.5	34.2	38.9	...
1923	12.2	...	0.2	6.0	12.5	18.7	25.4	31.1	32.7	36.8	38.4	...
1924	12.7	...	0.2	6.6	12.4	19.5	26.4	32.9	34.9	32.3	33.5	...
1925	12.7	...	0.2	6.2	12.7	19.0	26.8	31.9	37.2	33.1	35.0	...
1926	13.4	...	0.2	6.9	12.7	20.3	27.8	33.4	38.1	37.8	34.8	...
1927	14.0	...	0.3	6.8	13.6	20.7	29.4	35.5	39.0	37.8	37.5	...
1928	14.5	...	0.2	7.1	14.0	20.5	31.0	36.1	42.4	41.4	32.3	...
1929	14.9	...	0.2	7.5	14.2	21.6	30.5	36.9	42.9	42.6	27.2	...
1930	16.8	...	0.2	8.0	15.7	24.2	35.3	43.4	44.5	42.0	31.5	...
1931	18.0	...	0.2	7.6	15.8	24.4	38.6	49.3	51.0	44.2	36.3	...
1932	18.7	...	0.2	7.6	16.4	24.2	38.9	51.5	53.9	47.4	30.1	...
1933	17.2	...	0.2	7.5	14.6	22.5	35.4	45.5	48.5	47.4	34.9	...
1934	16.1	...	0.2	7.6	15.2	21.3	31.6	39.2	43.1	42.9	27.4	...
1935	15.4	...	0.2	7.4	15.0	20.1	30.6	36.7	37.9	40.6	30.0	...
1936	15.4	...	0.3	7.1	15.4	21.0	29.6	35.9	36.7	39.8	29.8	...
1937	16.1	...	0.3	7.3	15.2	22.1	32.1	37.4	36.5	40.5	30.5	...
1938	16.4	...	0.2	7.0	15.3	22.5	32.7	39.4	36.6	37.5	30.2	...
1939	15.3	...	0.3	6.2	13.7	20.2	30.0	36.4	36.2	37.5	26.8	...

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

WHITE, MALE												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	23.2	...	...	...	...	...	...	...	...	...	...	...
1911	24.1	...	...	...	...	...	...	...	...	...	...	...
1912	23.8	...	...	...	...	...	...	...	...	...	...	...
1913	23.6	...	...	...	...	...	...	...	...	...	...	...
1914	24.5	...	...	*	12.6	27.0	35.3	52.1	64.1	60.6	59.4	65.2
1915	24.8	...	...	*	11.7	26.2	35.8	50.6	68.2	66.8	63.1	79.1
1916	21.5	...	...	*	10.8	22.1	31.1	41.8	59.6	63.2	65.4	50.5
1917	20.1	...	...	...	...	...	...	...	...	...	...	...
1918	19.0	...	...	*	9.5	24.1	29.0	33.3	45.2	49.8	53.9	65.7
1919	17.5	...	...	0.3	7.7	19.9	27.1	31.7	43.3	49.4	59.7	88.1
1920	15.4	...	...	*	7.5	15.7	22.7	30.2	38.3	46.6	52.2	69.6
1921	20.0	...	...	0.4	8.5	19.4	29.9	42.5	52.2	56.6	60.7	78.2
1922	18.7	...	...	0.3	7.4	17.8	27.4	40.0	49.3	59.7	60.1	78.0
1923	18.1	...	...	0.3	6.3	16.9	27.3	38.4	48.0	55.3	65.2	74.1
1924	19.2	...	...	*	7.7	17.1	28.8	40.5	52.3	59.0	58.9	67.3
1925	19.1	...	...	0.3	7.4	17.0	27.3	40.3	51.1	63.0	61.5	68.4
1926	20.0	...	...	0.3	7.7	16.8	29.6	42.0	53.0	64.7	67.7	67.1
1927	21.2	...	...	0.3	8.3	18.4	30.0	45.2	56.8	66.3	68.5	74.1
1928	22.1	...	...	0.2	8.1	19.3	30.1	48.2	58.6	72.1	77.0	61.1
1929	22.5	...	...	0.3	8.9	19.2	31.8	46.7	58.1	72.6	76.8	54.8
1930	25.9	...	...	0.3	9.6	21.6	36.2	55.7	70.6	75.3	76.1	63.8
1931	28.2	...	...	0.3	9.2	21.8	36.4	61.2	81.4	88.6	80.1	75.2
1932	29.4	...	...	0.3	9.4	23.0	36.5	62.1	85.9	92.2	88.5	58.3
1933	26.8	...	...	0.2	9.1	20.4	33.2	56.3	74.7	84.1	88.2	70.9
1934	24.6	...	...	0.3	9.6	21.3	31.0	48.1	63.6	74.8	78.1	56.2
1935	23.3	...	...	0.3	9.5	20.7	29.0	47.0	58.5	63.6	75.8	62.6
1936	23.3	...	...	0.4	9.1	21.0	30.7	44.9	57.3	61.6	73.0	59.9
1937	24.5	...	...	0.5	9.5	21.5	32.2	48.5	59.2	61.6	73.8	62.6
1938	25.3	...	...	0.3	9.4	22.2	33.1	50.1	63.6	61.9	68.5	59.9
1939	23.4	...	...	0.5	8.5	19.8	29.3	44.8	58.2	60.2	70.4	52.9

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

... Category not applicable

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

WHITE, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	7.2	...	...	...	...	...	...	...	...	...	...	...
1911	7.6	...	...	...	...	...	...	...	...	...	...	...
1912	7.4	...	...	...	...	...	...	...	...	...	...	...
1913	7.2	...	...	...	...	...	...	...	...	...	...	...
1914	7.8	...	...	*	9.3	10.3	11.5	12.6	15.5	11.7	9.5	*
1915	7.7	...	...	*	8.1	10.4	11.6	13.3	14.0	14.0	10.8	*
1916	6.8	...	...	*	7.2	9.3	9.7	12.4	12.9	11.5	8.5	*
1917	6.9	...	...	...	...	...	...	...	...	...	...	...
1918	6.5	...	...	*	5.8	9.1	10.1	11.5	13.4	11.2	10.9	*
1919	6.7	...	...	*	6.1	8.5	10.5	12.9	13.4	11.9	10.6	*
1920	6.1	...	...	*	5.5	8.2	10.0	11.7	11.1	9.9	8.6	*
1921	6.1	...	...	*	5.7	8.0	9.2	11.5	11.8	10.2	9.3	*
1922	6.1	...	...	*	6.0	7.6	9.6	11.4	11.9	10.7	10.7	*
1923	6.2	...	...	*	5.7	8.1	9.4	11.3	12.9	9.5	10.9	*
1924	5.9	...	...	*	5.4	7.6	9.5	11.1	11.9	10.1	8.0	*
1925	6.2	...	...	*	5.0	8.4	9.9	12.3	11.3	10.7	6.9	*
1926	6.6	...	...	*	6.1	8.6	10.2	12.6	12.3	10.7	10.3	*
1927	6.6	...	...	0.2	5.3	8.8	10.6	12.4	12.4	10.9	9.4	*
1928	6.6	...	...	*	6.1	8.8	10.3	12.5	11.9	11.8	8.2	*
1929	7.1	...	...	*	6.0	9.3	10.8	12.9	14.1	12.3	10.7	*
1930	7.4	...	...	*	6.4	9.8	11.4	13.2	14.5	12.7	10.0	*
1931	7.6	...	...	*	5.9	9.9	11.8	14.2	15.2	12.3	10.5	*
1932	7.7	...	...	*	5.9	9.9	11.1	13.9	15.0	14.7	9.0	*
1933	7.3	...	...	*	5.9	9.0	11.3	12.9	14.6	12.1	9.1	*
1934	7.3	...	...	*	5.7	9.2	11.1	13.8	13.3	10.9	9.9	*
1935	7.2	...	...	*	5.4	9.4	10.9	13.0	13.8	11.9	7.9	*
1936	7.3	...	...	0.2	5.0	9.8	10.9	13.3	13.4	11.6	9.0	*
1937	7.6	...	...	*	5.1	9.0	11.6	14.6	14.4	11.5	9.8	*
1938	7.4	...	...	*	4.6	8.5	11.7	14.3	14.0	11.4	9.0	*
1939	7.1	...	...	*	3.9	7.6	11.0	14.3	13.7	12.4	7.5	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ALL OTHER, BOTH SEXES

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	11.8	...	...	...	...	...	...	...	...	...	...	...
1911	12.7	...	...	...	...	...	...	...	...	...	...	...
1912	9.7	...	...	...	...	...	...	...	...	...	...	...
1913	8.9	...	...	...	...	...	...	...	...	...	...	...
1914	9.8	...	...	*	8.1	18.3	15.5	9.1	17.4	*	*	*
1915	9.3	...	...	*	6.3	17.4	14.8	13.1	*	*	*	*
1916	5.0	...	...	*	4.3	9.5	9.2	8.5	*	*	*	*
1917	4.9	...	...	...	...	...	...	...	...	...	...	...
1918	5.5	...	...	*	5.2	9.3	9.9	6.1	10.3	16.5	*	*
1919	4.0	...	...	*	3.8	8.0	7.2	5.8	*	*	*	*
1920	3.6	...	...	*	3.7	5.6	5.4	7.6	6.3	*	*	*
1921	4.6	...	...	*	5.0	7.7	8.0	6.7	6.7	*	*	*
1922	4.2	...	...	*	3.8	6.1	8.9	7.6	6.7	*	*	*
1923	3.7	...	...	*	3.0	5.5	7.8	8.0	6.4	*	*	*
1924	4.3	...	...	*	3.6	7.2	8.4	6.9	8.8	*	*	*
1925	4.0	...	...	*	3.5	6.7	7.7	6.0	5.8	10.1	*	*
1926	4.4	...	...	*	3.3	7.3	7.2	8.6	9.3	*	*	*
1927	4.7	...	...	*	3.5	8.6	8.6	8.2	8.2	*	*	*
1928	4.8	...	...	*	4.0	7.9	8.6	8.7	10.5	*	*	*
1929	4.9	...	...	*	4.4	7.8	7.7	10.1	9.3	10.2	*	*
1930	5.0	...	...	*	4.0	7.7	8.5	9.0	10.6	13.8	*	*
1931	5.2	...	...	*	4.0	8.8	8.5	8.7	13.2	9.0	*	*
1932	5.5	...	...	*	4.8	8.1	10.3	10.1	15.7	9.3	*	*
1933	4.9	...	...	*	4.9	7.4	9.8	10.9	9.0	9.8	*	*
1934	4.7	...	...	*	5.2	8.4	7.7	8.8	10.5	6.9	*	*
1935	5.1	...	...	*	4.3	8.4	8.4	8.3	10.7	6.2	*	*
1936	4.7	...	...	*	4.2	7.0	7.9	6.9	10.8	6.9	*	*
1937	4.9	...	...	*	3.9	7.6	8.3	8.7	10.0	7.5	*	*
1938	4.9	...	...	*	4.1	7.5	7.8	8.7	11.2	6.1	*	*
1939	4.2	...	...	*	3.3	6.5	7.0	8.6	7.3	6.0	*	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ALL OTHER, MALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	16.9	...	...	...	...	...	...	...	...	...	...	...
1911	19.3	...	...	...	...	...	...	...	...	...	...	...
1912	13.8	...	...	...	...	...	...	...	...	...	...	...
1913	12.7	...	...	...	...	...	...	...	...	...	...	...
1914	14.6	...	...	*	8.4	26.9	23.6	13.9	29.7	*	*	*
1915	13.3	...	...	*	*	24.0	20.2	18.7	*	*	*	*
1916	7.9	...	...	*	4.8	15.5	15.4	12.6	*	*	*	*
1917	7.2	...	...	...	...	...	...	...	...	...	...	...
1918	8.4	...	...	*	6.7	14.8	15.7	8.6	16.1	*	*	*
1919	6.0	...	...	*	5.2	11.8	10.9	8.8	*	*	*	*
1920	5.2	...	...	*	3.7	8.5	8.3	10.7	*	*	*	*
1921	6.9	...	...	*	6.3	10.8	12.8	10.3	*	*	*	*
1922	6.3	...	...	*	4.6	9.2	14.1	10.7	9.9	*	*	*
1923	5.6	...	...	*	3.5	7.5	11.9	13.2	10.4	*	*	*
1924	6.3	...	...	*	4.1	9.6	13.6	10.2	13.8	*	*	*
1925	6.1	...	...	*	4.8	9.7	12.0	8.2	9.1	*	*	*
1926	6.2	...	...	*	3.8	9.7	10.3	12.1	14.0	*	*	*
1927	7.0	...	...	*	4.3	12.8	12.8	12.4	12.6	*	*	*
1928	7.2	...	...	*	5.2	10.8	14.0	14.2	15.8	*	*	*
1929	7.6	...	...	*	5.7	11.6	11.9	16.6	13.8	17.5	*	*
1930	7.6	...	...	*	4.4	11.9	13.4	14.3	16.5	23.0	*	*
1931	8.0	...	...	*	4.6	12.7	14.2	14.6	20.0	14.9	*	*
1932	8.6	...	...	*	5.9	11.9	16.1	16.8	24.9	16.3	*	*
1933	7.7	...	...	*	5.9	11.2	16.3	17.5	13.8	17.4	*	*
1934	6.9	...	...	*	6.3	11.9	11.3	13.8	16.7	*	*	*
1935	7.4	...	...	*	5.1	10.7	13.0	14.8	17.8	10.8	*	*
1936	7.0	...	...	*	5.5	9.2	13.2	10.6	17.8	11.2	*	*
1937	7.4	...	...	*	4.4	10.7	13.7	13.2	15.9	13.0	*	*
1938	7.4	...	...	*	5.5	9.8	12.7	14.0	17.2	9.0	*	*
1939	6.5	...	...	*	3.9	10.1	11.8	13.1	12.0	9.3	*	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

... Category not applicable

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

SUICIDE

ALL OTHER, FEMALE

Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	6.0	...	...	...	...	...	...	...	...	...	...	...
1911	5.3	...	...	...	...	...	...	...	...	...	...	...
1912	5.2	...	...	...	...	...	...	...	...	...	...	...
1913	4.7	...	...	...	...	...	...	...	...	...	...	...
1914	4.6	...	...	*	7.8	8.9	*	*	*	*	*	*
1915	5.1	...	...	*	*	10.4	*	*	*	*	*	*
1916	2.0	...	...	*	*	*	*	*	*	*	*	*
1917	2.5	...	...	...	...	...	...	...	...	...	...	...
1918	2.5	...	...	*	3.8	4.1	*	*	*	*	*	*
1919	2.0	...	...	*	*	4.6	*	*	*	*	*	*
1920	2.0	...	...	*	3.7	*	*	*	*	*	*	*
1921	2.3	...	...	*	3.7	4.9	*	*	*	*	*	*
1922	2.1	...	...	*	3.1	3.3	*	*	*	*	*	*
1923	1.8	...	...	*	2.5	3.6	3.6	*	*	*	*	*
1924	2.3	...	...	*	3.2	4.9	*	*	*	*	*	*
1925	2.0	...	...	*	2.3	3.9	*	*	*	*	*	*
1926	2.6	...	...	*	2.8	5.1	3.9	*	*	*	*	*
1927	2.4	...	...	*	2.8	4.8	4.3	*	*	*	*	*
1928	2.4	...	...	*	2.9	5.3	3.2	*	*	*	*	*
1929	2.3	...	...	*	3.3	4.2	3.6	*	*	*	*	*
1930	2.4	...	...	*	3.6	3.8	3.7	*	*	*	*	*
1931	2.5	...	...	*	3.5	5.2	3.0	*	*	*	*	*
1932	2.5	...	...	*	3.7	4.5	4.7	*	*	*	*	*
1933	2.2	...	...	*	4.0	3.9	3.5	3.4	*	*	*	*
1934	2.6	...	...	*	4.3	5.1	4.2	*	*	*	*	*
1935	2.7	...	...	*	3.6	6.3	4.0	*	*	*	*	*
1936	2.4	...	...	*	3.0	4.9	2.7	*	*	*	*	*
1937	2.5	...	...	*	3.4	4.7	3.0	3.8	*	*	*	*
1938	2.5	...	...	*	2.9	5.3	3.1	*	*	*	*	*
1939	2.0	...	...	*	2.7	3.2	2.4	3.7	*	*	*	*

NOTES:  
 For 1932-34, deaths of Mexicans are included with total deaths for the white population; but by age, are included with deaths for the nonwhite population. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes.....	40	60	64
Male.....	28	41	45
Female.....	12	19	19

... Category not applicable

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ICD Revision	Years in use	ICD Nos.
Fifth	1939-1969	176, 178-195
Fourth	1930-1938	176, 178-195, 201-214
Third	1921-1929	176-189, 192-196, 201, 202pt., 203
Second	1910-1920	165-181, 185, 186
First	1900-1909	164-175, 176pt.

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	72.3	128.8	75.3	38.3	53.4	62.6	78.3	80.6	90.9	131.5	274.8	658.9
1901	83.8	163.6	82.5	40.4	53.9	69.4	83.8	103.9	128.4	172.6	359.7	863.8
1902	72.5	100.0	68.4	37.5	52.7	65.4	77.9	84.1	100.9	139.4	275.4	799.4
1903	81.4	118.1	78.3	41.7	57.7	72.1	85.8	92.2	108.5	159.2	358.5	944.2
1904	85.4	122.2	86.2	48.7	60.9	74.8	86.2	94.4	112.3	158.5	393.3	1,034.2
1905	81.3	133.6	79.9	43.5	59.5	68.5	84.3	87.8	114.4	160.0	344.9	1,083.3
1906	94.4	131.1	87.4	45.6	74.9	91.0	99.1	104.8	120.8	170.9	382.3	1,061.3
1907	94.8	131.8	92.5	44.0	76.4	91.0	104.9	107.7	125.2	164.7	375.2	1,071.3
1908	82.9	128.0	87.6	41.1	64.5	74.5	84.4	95.1	111.8	154.5	367.3	981.4
1909	79.9	117.8	84.7	38.3	57.7	72.0	83.3	91.9	105.0	153.7	385.7	1,109.4
1910	84.2	114.1	82.6	38.0	64.1	80.2	90.3	95.8	109.3	167.2	376.3	1,081.3
1911	83.6	134.0	85.9	37.3	60.5	73.2	86.5	98.1	116.9	170.8	401.2	1,213.3
1912	80.8	112.7	85.6	38.6	60.6	72.1	81.1	92.5	112.2	159.0	386.0	1,090.8
1913	83.8	118.6	84.2	40.2	65.7	76.4	84.5	94.7	114.4	164.1	384.1	1,050.6
1914	76.7	118.8	81.1	37.2	57.1	67.0	75.7	84.5	104.2	159.6	376.5	1,049.0
1915	73.5	102.5	79.2	38.1	53.7	62.0	71.7	81.0	99.3	155.4	376.7	1,038.5
1916	81.6	122.5	83.1	39.5	59.5	70.1	83.2	94.9	116.0	169.6	391.9	1,112.4
1917	86.0	134.8	84.8	43.7	63.2	71.9	84.5	100.7	123.4	181.5	418.4	1,056.7
1918	81.4	105.2	83.4	46.8	61.8	66.4	76.3	88.9	113.8	177.4	388.0	1,054.7
1919	71.2	96.8	81.5	45.4	54.8	55.7	64.3	72.1	94.3	147.2	359.1	1,003.8
1920	70.0	100.2	80.2	44.3	55.1	53.9	58.9	69.1	89.7	156.2	391.0	1,115.0
1921	66.7	90.8	74.0	41.8	51.0	50.9	58.0	66.5	87.6	150.8	378.0	1,151.7
1922	68.4	91.1	73.3	41.0	51.1	51.8	59.7	72.0	93.4	156.4	397.2	1,207.4
1923	74.5	93.4	72.6	42.2	56.2	57.3	68.3	81.7	104.1	171.5	425.5	1,278.1
1924	74.0	101.0	75.1	42.1	56.6	54.1	67.2	80.8	102.1	170.6	415.4	1,293.6
1925	76.7	98.2	70.5	41.9	59.3	56.9	70.0	86.3	108.2	179.1	443.1	1,349.4
1926	77.3	106.3	68.4	41.0	58.7	57.6	71.4	88.8	109.8	178.0	453.7	1,327.0
1927	77.2	95.0	68.6	40.6	59.4	57.9	71.4	86.1	112.1	181.0	447.0	1,274.2
1928	78.2	100.0	65.7	40.1	59.6	59.6	70.7	89.7	112.4	186.0	472.9	1,265.3
1929	79.9	98.9	64.5	39.1	62.8	62.8	71.8	90.1	118.2	192.4	463.2	1,228.8
1930	79.8	103.4	61.2	36.1	61.8	62.3	70.1	90.4	120.1	198.1	477.8	1,302.0
1931	77.8	93.9	62.2	34.9	59.2	57.6	66.9	88.5	120.8	195.2	479.2	1,253.2
1932	70.8	90.2	57.5	32.5	52.4	51.7	58.6	75.9	107.8	180.4	454.9	1,263.2
1933	71.9	95.6	56.5	33.3	52.7	53.2	58.8	78.5	109.0	179.1	465.9	1,228.3
1934	79.3	106.9	56.2	33.9	56.9	58.3	64.7	90.0	123.6	202.9	516.1	1,317.9
1935	77.8	105.1	56.7	32.2	56.0	58.0	64.6	85.9	117.2	192.6	501.4	1,373.9
1936	85.4	124.9	58.8	32.8	58.0	61.0	68.9	93.8	132.4	220.5	583.9	1,543.8
1937	81.1	112.6	56.0	32.6	60.4	58.6	64.8	86.0	120.8	198.9	537.0	1,371.6
1938	71.8	111.1	51.5	28.3	50.9	49.7	54.7	74.3	103.3	179.2	500.9	1,332.3
1939	70.4	109.8	50.7	27.8	50.5	49.4	52.5	69.3	98.5	172.3	495.8	1,314.8

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	110.3	138.4	89.8	56.5	92.8	108.5	132.6	129.2	140.9	181.3	283.7	515.9
1901	124.0	192.3	89.4	59.6	93.9	118.0	138.2	163.3	187.6	218.9	378.8	686.3
1902	110.2	103.2	78.1	56.0	90.0	112.2	130.0	140.4	152.3	194.2	301.9	613.5
1903	123.0	133.6	86.4	63.3	100.3	124.1	143.5	151.2	164.5	215.1	347.3	748.6
1904	125.1	137.0	99.3	67.7	99.7	122.2	139.4	151.6	169.2	214.1	414.4	917.3
1905	122.8	158.3	89.6	63.7	104.3	118.1	142.6	147.2	175.9	208.9	344.1	870.4
1906	145.8	147.7	94.5	68.5	132.6	157.3	168.7	170.6	192.1	237.1	381.5	905.3
1907	146.7	142.7	99.8	65.2	136.9	158.3	179.5	180.2	195.8	215.3	372.0	925.9
1908	125.8	146.2	98.1	60.8	112.5	127.7	143.2	156.2	173.7	194.4	379.3	823.4
1909	120.3	126.1	93.1	55.6	101.6	124.8	141.3	151.1	162.2	198.1	368.8	941.5
1910	129.1	126.5	91.9	55.4	113.6	140.9	154.8	158.1	171.2	216.5	363.8	884.8
1911	124.6	144.4	95.9	54.6	103.8	125.8	147.3	158.4	178.8	218.1	377.9	932.2
1912	121.3	122.3	96.3	56.2	105.4	125.3	137.0	153.4	172.4	205.0	340.0	833.3
1913	126.6	127.3	93.2	58.1	115.7	132.2	144.7	155.9	178.9	212.0	356.3	796.2
1914	114.2	127.0	91.4	54.7	99.2	116.3	128.2	137.7	159.2	204.8	340.6	802.2
1915	109.1	110.9	90.2	56.3	93.0	107.6	121.8	131.0	149.0	196.4	348.5	828.2
1916	122.1	130.9	91.0	57.3	104.8	121.6	141.7	154.6	176.6	219.3	378.3	955.2
1917	126.9	145.2	90.7	61.0	109.9	125.3	143.2	162.1	186.1	228.6	387.4	798.6
1918	121.9	114.7	92.5	67.3	112.1	119.8	127.6	141.7	171.9	231.7	360.4	879.2
1919	103.4	109.8	88.4	65.2	95.2	96.0	107.3	113.4	138.6	188.4	327.7	800.4
1920	99.6	111.0	85.6	62.9	95.1	92.8	96.4	107.7	128.6	191.3	349.4	905.1
1921	95.3	98.4	82.8	60.1	88.0	87.1	95.5	103.8	125.1	184.4	341.3	893.6
1922	97.7	99.8	79.6	58.2	86.6	89.2	99.3	113.8	137.1	191.7	365.1	907.0
1923	107.1	102.9	79.7	59.4	95.4	98.1	112.7	129.9	154.3	215.4	382.7	957.5
1924	105.3	110.8	82.8	58.6	94.9	92.3	110.5	128.0	148.5	207.4	385.1	968.2
1925	109.3	104.3	76.4	59.2	98.5	96.5	114.7	136.3	156.4	226.2	405.4	1,102.1
1926	110.5	114.8	74.8	57.6	98.6	97.7	118.1	140.0	161.1	221.4	416.2	1,005.7
1927	111.4	103.6	77.5	57.2	100.4	98.9	117.5	137.2	164.5	225.9	423.2	1,004.3
1928	112.1	107.9	73.3	56.2	100.1	102.2	117.5	140.9	163.3	230.1	438.7	960.0
1929	114.9	108.4	70.6	55.1	104.8	106.8	119.0	143.0	172.7	238.5	444.6	892.3
1930	114.4	113.3	68.3	50.2	103.3	106.5	117.2	142.6	175.0	246.3	439.0	1,026.6
1931	110.7	101.3	68.9	48.7	97.8	97.7	110.4	139.4	175.7	240.4	453.2	1,001.1
1932	99.3	98.2	64.9	45.1	86.1	87.1	96.5	118.7	156.4	216.4	424.1	968.6
1933	100.8	103.1	62.6	47.1	85.0	89.5	95.6	123.2	157.6	219.4	425.1	977.1
1934	112.5	116.2	63.8	47.4	93.4	97.3	106.6	143.9	182.3	249.9	493.0	1,085.6
1935	109.4	114.5	62.5	44.8	90.7	97.1	106.5	135.4	171.6	238.4	474.9	1,096.3
1936	119.3	135.7	64.7	45.4	94.6	101.7	113.4	149.3	195.8	269.9	547.2	1,227.8
1937	114.8	125.3	62.9	45.5	98.9	98.7	106.9	137.1	181.7	245.5	515.5	1,147.9
1938	99.5	123.3	58.3	39.7	84.6	83.8	89.8	117.6	151.7	216.3	451.1	1,080.2
1939	97.7	122.6	57.9	40.0	84.2	83.6	87.3	110.3	144.2	209.3	452.6	999.4

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	34.1	119.1	60.6	20.0	16.3	16.1	20.9	29.9	41.9	84.3	266.7	775.7
1901	43.5	134.3	75.5	21.1	16.1	20.1	26.1	41.7	70.4	128.7	342.4	992.6
1902	34.5	96.7	58.6	18.8	17.3	17.9	22.6	25.0	50.4	87.6	251.6	934.8
1903	39.5	102.3	70.0	19.8	17.2	19.1	24.6	30.0	53.3	106.3	368.6	1,087.3
1904	45.5	107.2	72.9	29.5	23.7	26.2	29.6	34.1	56.1	106.0	374.4	1,119.9
1905	39.4	108.4	70.1	23.1	16.5	17.5	22.3	25.0	53.5	113.8	345.6	1,240.2
1906	41.1	114.1	80.1	22.5	17.8	19.8	22.5	32.8	47.5	106.4	383.0	1,178.2
1907	40.9	120.6	85.1	22.6	16.1	18.4	22.7	28.1	52.2	115.3	378.1	1,180.7
1908	37.8	109.2	76.9	21.1	16.0	16.1	18.9	26.9	46.4	115.0	356.2	1,103.7
1909	37.5	109.3	76.1	20.7	13.2	14.0	18.9	26.1	44.5	109.6	401.1	1,240.4
1910	36.7	101.3	73.1	20.3	13.6	12.8	18.2	25.7	43.0	118.0	387.7	1,235.6
1911	40.3	123.3	75.7	19.8	16.4	15.3	18.8	30.2	50.0	123.1	422.8	1,435.9
1912	38.1	102.9	74.6	20.8	15.2	13.8	18.9	23.7	46.8	112.6	428.2	1,293.1
1913	38.6	109.6	75.0	21.9	15.4	15.8	17.8	25.6	44.3	115.7	409.4	1,248.6
1914	37.2	110.3	70.6	19.5	14.7	13.8	17.4	24.4	44.1	113.6	409.5	1,241.0
1915	36.2	93.8	68.0	19.6	14.3	12.9	16.2	24.4	45.0	113.5	402.5	1,200.5
1916	39.2	113.8	75.0	21.3	14.6	15.5	18.9	27.1	49.5	118.9	404.2	1,232.0
1917	43.5	124.2	78.7	26.1	17.6	16.4	20.4	31.1	54.6	133.2	446.4	1,251.6
1918	40.8	95.5	74.1	25.9	17.6	16.1	20.3	28.8	49.8	121.4	413.1	1,186.8
1919	38.0	83.5	74.4	25.4	16.5	15.3	17.5	24.9	45.4	104.6	387.6	1,156.0
1920	39.3	89.2	74.7	25.3	16.1	14.4	17.7	25.5	47.0	119.9	429.2	1,273.0
1921	37.2	83.1	65.0	23.2	14.7	14.4	16.9	24.7	46.7	116.2	411.7	1,347.1
1922	38.2	82.1	66.7	23.6	16.5	14.7	16.8	25.5	46.0	119.9	426.7	1,435.4
1923	41.0	83.6	65.2	24.8	17.7	16.7	20.4	28.5	49.6	126.0	464.8	1,523.5
1924	41.7	90.9	67.1	25.3	18.8	15.9	20.5	29.3	51.5	132.4	443.3	1,542.7
1925	43.1	91.9	64.3	24.3	20.7	17.3	21.9	31.8	55.6	130.2	477.9	1,539.2
1926	43.1	97.6	61.9	24.1	19.3	17.5	21.4	33.1	53.8	133.0	488.4	1,574.0
1927	42.0	86.1	59.5	23.7	18.9	16.8	22.2	30.4	54.9	134.5	469.1	1,481.8
1928	43.3	91.8	57.9	23.7	19.8	17.3	21.0	33.9	56.6	140.3	504.8	1,498.8
1929	43.9	89.1	58.2	22.7	21.4	19.2	22.0	32.4	58.9	144.8	480.7	1,483.7
1930	44.3	93.2	53.9	21.6	21.2	18.7	20.7	33.4	60.9	148.1	514.5	1,508.8
1931	44.1	86.3	55.2	20.7	21.4	18.0	21.3	33.1	61.7	148.4	503.6	1,442.4
1932	41.7	81.8	50.0	19.6	19.3	16.8	19.2	29.5	55.7	143.4	483.8	1,484.3
1933	42.4	87.9	50.4	19.2	20.1	17.5	20.5	30.2	56.9	137.6	504.3	1,417.6
1934	45.5	97.3	48.4	20.1	21.0	20.0	21.4	31.9	60.9	154.8	537.7	1,492.8
1935	45.7	95.5	50.9	19.2	22.0	19.6	21.3	32.7	59.2	146.0	526.2	1,582.8
1936	51.0	113.8	52.8	19.9	22.2	21.1	23.3	34.3	65.0	170.5	618.1	1,781.6
1937	47.0	99.5	48.9	19.4	22.5	19.2	21.7	31.3	56.2	152.0	556.8	1,540.0
1938	43.7	98.5	44.5	16.5	17.7	16.4	19.0	28.1	52.0	142.1	546.9	1,522.1
1939	42.7	96.6	43.3	15.3	17.4	15.9	17.1	25.6	50.3	135.5	535.3	1,552.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	84.0	---	---	---	---	---	---	---	---	---	---	---
1911	83.1	---	---	---	---	---	---	---	---	---	---	---
1912	80.3	---	---	---	---	---	---	---	---	---	---	---
1913	83.1	---	---	---	---	---	---	---	---	---	---	---
1914	76.1	109.2	79.6	36.6	56.4	66.3	75.1	84.0	104.3	160.1	378.3	1,080.2
1915	73.0	96.6	77.5	37.4	53.0	61.7	71.1	80.7	98.7	155.3	380.2	1,060.5
1916	81.3	113.6	79.7	39.0	58.7	69.8	82.7	95.5	116.3	171.0	397.3	1,159.6
1917	85.2	---	---	---	---	---	---	---	---	---	---	---
1918	80.9	96.4	80.7	46.5	60.7	66.1	74.9	88.6	113.7	177.1	392.9	1,090.2
1919	70.7	86.9	79.1	45.1	53.5	54.9	63.9	71.6	94.6	147.0	364.4	1,070.5
1920	69.4	89.2	77.1	43.8	53.2	53.2	58.8	68.8	89.5	156.5	399.3	1,166.5
1921	66.5	81.1	72.7	41.7	49.3	50.5	57.5	66.2	88.0	151.6	387.1	1,218.6
1922	68.2	78.6	70.4	41.2	49.7	51.3	59.4	71.3	93.3	158.0	409.8	1,283.3
1923	73.9	82.7	69.3	41.9	54.2	55.8	67.7	81.1	104.5	172.5	434.9	1,342.6
1924	72.8	87.4	71.0	41.6	54.2	52.3	65.7	79.3	101.4	171.2	422.6	1,350.1
1925	75.5	87.4	67.3	41.3	56.7	54.5	67.8	84.8	107.9	180.7	451.8	1,417.2
1926	75.9	97.5	64.9	40.2	56.3	54.6	68.9	87.1	109.0	178.6	464.6	1,392.2
1927	75.9	83.0	66.3	39.9	57.2	54.9	68.5	84.0	111.7	182.0	455.9	1,344.7
1928	77.3	87.7	63.5	39.7	57.6	57.1	68.2	87.7	112.7	187.2	485.2	1,344.1
1929	79.1	87.4	62.4	38.5	61.3	59.8	69.3	88.2	117.3	194.2	473.8	1,307.8
1930	79.3	91.3	58.9	35.5	60.7	60.0	67.6	88.9	119.3	199.4	489.1	1,389.3
1931	77.7	83.7	61.0	34.2	58.7	55.3	65.2	87.1	120.3	196.7	493.3	1,346.9
1932	70.1	79.1	55.5	31.6	50.9	48.9	56.1	74.3	106.9	181.7	466.3	1,347.4
1933	71.0	82.2	54.8	32.2	51.0	50.4	56.0	76.3	107.3	179.9	479.5	1,316.7
1934	78.4	92.1	53.2	32.9	55.0	55.0	61.7	87.3	121.9	204.6	530.2	1,402.3
1935	77.7	93.1	55.3	31.6	55.0	55.7	61.9	84.5	116.4	194.2	515.7	1,463.3
1936	84.9	110.6	57.3	32.2	56.7	57.9	65.3	91.5	130.9	221.7	600.5	1,644.0
1937	80.9	98.8	53.8	32.0	59.4	56.3	61.8	83.8	119.6	200.8	552.6	1,471.1
1938	71.6	100.3	49.9	27.3	49.4	47.4	52.0	72.7	102.1	182.2	517.7	1,431.2
1939	70.2	98.5	49.1	26.8	49.2	47.4	50.2	67.1	97.5	174.6	511.7	1,412.6

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

WHITE, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	128.8	---	---	---	---	---	---	---	---	---	---	---
1911	124.0	---	---	---	---	---	---	---	---	---	---	---
1912	120.6	---	---	---	---	---	---	---	---	---	---	---
1913	125.9	---	---	---	---	---	---	---	---	---	---	---
1914	113.6	116.8	90.3	54.5	98.0	115.1	127.7	138.0	160.0	205.9	340.8	818.6
1915	108.5	104.4	88.6	55.8	92.0	107.5	121.5	131.0	148.8	196.2	350.9	849.5
1916	122.0	120.3	88.0	57.5	103.3	121.2	141.3	156.4	177.9	220.7	382.0	986.7
1917	126.2	---	---	---	---	---	---	---	---	---	---	---
1918	121.4	105.8	90.2	67.6	110.0	119.6	126.0	142.5	173.1	232.2	360.9	899.3
1919	102.9	100.4	86.5	65.2	92.7	95.2	107.0	113.7	140.2	187.8	332.0	849.5
1920	98.9	99.7	83.2	63.4	91.4	91.6	96.4	108.1	128.8	191.9	355.6	930.4
1921	94.9	88.6	82.3	60.4	84.5	86.1	95.1	103.9	126.9	186.7	347.9	933.6
1922	97.6	86.7	78.1	59.1	84.1	88.1	99.6	113.5	138.1	194.3	374.4	954.7
1923	106.1	91.6	77.2	59.3	91.8	95.5	111.4	129.7	156.4	217.1	388.8	987.0
1924	103.7	96.2	80.0	58.6	90.9	89.5	108.3	127.0	148.6	209.3	388.9	988.0
1925	107.4	96.1	73.7	58.6	94.1	92.5	111.1	134.5	157.3	228.2	409.3	1,135.1
1926	108.2	105.6	71.7	56.9	94.3	92.5	114.5	137.9	160.8	222.6	421.5	1,025.7
1927	108.8	90.4	75.3	56.5	95.9	93.5	112.7	134.7	165.1	227.2	425.1	1,038.4
1928	110.5	95.3	72.3	56.2	96.4	97.8	113.6	138.8	165.4	231.3	446.1	990.6
1929	113.3	97.8	69.3	54.7	102.0	101.8	114.8	140.9	173.2	240.2	447.3	931.1
1930	113.1	100.7	66.5	49.9	100.8	102.4	113.0	141.1	175.9	247.1	442.4	1,081.9
1931	110.3	91.7	68.9	47.9	96.5	94.2	107.8	138.0	176.7	242.4	465.6	1,061.1
1932	97.6	87.1	63.3	44.1	82.8	82.3	92.3	117.0	156.0	217.8	431.8	1,011.9
1933	98.7	89.4	61.6	45.6	82.5	84.6	91.2	120.3	156.3	219.9	430.3	1,029.7
1934	110.4	102.6	61.1	46.1	89.8	91.7	101.8	140.4	181.3	251.0	502.3	1,143.1
1935	108.4	102.3	62.1	44.2	88.1	93.1	101.9	134.2	171.4	239.8	484.4	1,153.3
1936	117.7	120.0	64.7	44.7	91.8	96.6	107.3	146.3	195.0	271.5	558.1	1,286.9
1937	113.6	112.6	60.7	45.1	96.5	94.8	102.0	134.3	181.2	247.2	526.1	1,222.3
1938	98.4	112.9	57.8	38.6	81.9	79.6	85.4	115.9	151.2	219.3	463.6	1,141.3
1939	96.5	109.8	56.8	38.5	81.9	80.3	83.4	106.8	143.7	211.6	464.0	1,062.0

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	36.6	---	---	---	---	---	---	---	---	---	---	---
1911	39.9	---	---	---	---	---	---	---	---	---	---	---
1912	37.9	---	---	---	---	---	---	---	---	---	---	---
1913	38.1	---	---	---	---	---	---	---	---	---	---	---
1914	36.6	101.3	68.6	18.5	14.5	13.5	17.0	23.3	43.7	113.7	412.8	1,284.8
1915	35.7	88.6	66.2	18.8	13.9	12.5	15.7	24.1	44.2	113.8	406.9	1,224.0
1916	38.8	106.6	71.2	20.0	14.2	15.2	18.4	27.1	49.2	120.5	411.2	1,291.6
1917	42.6	---	---	---	---	---	---	---	---	---	---	---
1918	40.3	86.8	71.0	25.0	17.5	15.6	19.3	27.9	48.9	120.7	421.9	1,234.5
1919	37.6	73.0	71.5	24.5	16.3	14.4	17.0	24.2	44.8	105.0	393.7	1,236.2
1920	38.8	78.4	70.7	23.8	15.6	13.9	17.3	24.9	46.8	120.2	439.0	1,344.5
1921	37.2	73.3	62.8	22.7	14.6	14.3	16.3	24.4	46.1	115.6	422.8	1,435.7
1922	38.0	70.2	62.6	22.9	15.9	14.5	15.9	25.0	45.5	120.9	441.9	1,535.3
1923	40.8	73.6	61.0	24.2	17.1	15.9	20.4	28.2	48.9	126.9	477.1	1,618.5
1924	40.8	78.3	61.6	24.3	17.7	14.8	19.5	27.7	50.7	132.1	453.6	1,631.6
1925	42.6	78.3	60.7	23.7	19.6	16.2	21.1	31.0	54.7	131.9	490.8	1,636.7
1926	42.7	89.0	57.9	23.1	18.5	16.3	19.9	32.3	53.2	133.4	504.2	1,677.4
1927	41.9	75.3	56.9	22.9	18.6	16.1	21.2	29.3	54.0	135.4	484.3	1,582.5
1928	43.1	79.8	54.3	22.8	19.1	16.3	19.8	32.4	55.8	141.8	521.5	1,616.4
1929	43.8	76.7	55.2	21.9	21.0	17.9	20.9	31.0	57.2	146.9	498.6	1,594.1
1930	44.5	81.5	51.1	20.7	21.1	17.8	19.6	32.2	59.0	150.1	533.0	1,620.1
1931	44.4	75.5	52.8	20.1	21.3	16.8	20.2	32.0	60.4	149.8	519.3	1,561.4
1932	41.9	70.7	47.3	18.8	19.3	15.7	18.1	28.2	54.8	144.8	498.7	1,599.2
1933	42.6	74.7	47.8	18.3	19.8	16.6	19.1	28.9	55.2	139.1	525.8	1,533.0
1934	45.6	81.1	45.1	19.2	20.5	18.8	19.9	30.2	59.1	157.4	556.4	1,696.4
1935	46.3	83.5	48.3	18.5	22.2	18.8	20.3	31.2	58.4	148.0	544.9	1,697.1
1936	51.5	100.8	49.7	19.3	21.9	19.7	21.9	32.9	63.3	171.7	639.8	1,913.2
1937	47.6	84.5	46.7	18.5	22.5	18.4	20.4	30.0	54.8	154.4	577.1	1,658.7
1938	44.3	87.3	41.8	15.7	17.2	15.6	17.7	26.7	50.6	145.3	567.5	1,649.8
1939	43.4	86.7	41.3	14.8	16.6	14.9	16.3	24.8	49.1	138.1	555.4	1,677.2

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	92.8	---	---	---	---	---	---	---	---	---	---	---
1911	100.8	---	---	---	---	---	---	---	---	---	---	---
1912	96.3	---	---	---	---	---	---	---	---	---	---	---
1913	97.9	---	---	---	---	---	---	---	---	---	---	---
1914	90.7	348.9	119.8	50.2	70.3	82.1	88.1	96.9	101.2	143.2	313.8	422.2
1915	85.8	241.6	121.8	53.6	68.5	66.5	82.6	89.3	118.7	158.3	258.2	587.5
1916	85.3	242.1	129.9	45.8	70.5	73.9	90.3	83.9	109.4	139.5	261.9	449.6
1917	96.4	---	---	---	---	---	---	---	---	---	---	---
1918	87.9	212.9	117.8	50.1	73.5	70.7	93.7	93.6	115.3	182.7	286.7	636.2
1919	75.8	198.2	106.8	49.1	66.2	63.9	68.4	77.9	88.7	151.7	266.2	350.9
1920	76.4	225.4	113.7	49.4	72.8	61.9	60.5	73.7	92.8	151.0	242.3	583.7
1921	69.0	198.1	88.2	42.6	66.9	55.4	63.8	70.9	80.5	137.6	216.0	444.5
1922	69.9	211.9	99.3	39.9	62.3	56.9	61.9	79.7	94.1	131.4	198.9	498.0
1923	79.9	198.6	103.2	45.2	73.1	71.9	74.6	88.3	97.9	154.1	271.6	649.2
1924	85.9	235.8	112.9	46.1	77.7	71.5	83.6	98.8	111.8	159.9	296.5	738.9
1925	88.7	208.0	100.3	47.2	82.6	80.4	94.0	104.7	112.6	149.7	298.8	669.1
1926	91.0	192.1	101.4	48.7	80.0	87.0	98.7	109.2	121.4	166.5	271.9	672.4
1927	89.9	205.5	89.5	46.9	78.4	85.0	101.7	109.5	118.3	163.8	300.2	584.1
1928	85.7	205.2	83.6	43.2	75.1	81.0	94.7	110.2	108.3	166.3	285.3	565.3
1929	86.9	200.0	81.0	43.7	74.2	87.8	96.4	109.5	131.3	161.9	295.8	517.6
1930	84.2	201.1	79.1	40.4	70.3	82.5	94.3	106.2	131.2	176.9	294.2	508.5
1931	77.9	174.8	71.5	40.3	63.2	76.9	83.5	102.8	127.5	168.6	245.6	396.1
1932	76.9	175.4	73.8	39.4	64.3	75.7	83.0	92.0	121.1	158.8	264.6	488.7
1933	79.6	193.6	69.7	41.9	65.8	76.2	84.7	101.4	132.1	165.5	244.8	444.9
1934	87.8	214.0	78.3	42.4	71.7	85.5	92.8	118.0	146.5	177.5	284.2	522.7
1935	79.5	192.2	67.3	36.9	64.3	77.1	89.4	100.4	127.5	169.1	264.7	576.2
1936	89.6	229.2	69.0	38.0	68.7	86.8	101.5	118.1	153.5	203.1	308.2	649.7
1937	83.2	213.7	71.3	37.1	68.3	77.3	91.8	108.9	136.8	173.4	274.6	483.5
1938	73.6	189.8	62.7	35.2	62.7	69.6	79.4	91.1	118.5	140.3	217.6	450.8
1939	72.1	192.8	61.5	34.8	61.4	66.4	72.3	93.4	112.6	142.9	224.9	443.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	137.7	...	...	...	...	...	...	...	...	...	...	...
1911	143.1	...	...	...	...	...	...	...	...	...	...	...
1912	141.8	...	...	...	...	...	...	...	...	...	...	...
1913	142.2	...	...	...	...	...	...	...	...	...	...	...
1914	129.1	375.3	120.7	59.4	123.1	140.5	138.1	131.2	136.0	172.6	333.7	*
1915	122.4	269.5	129.9	68.4	115.2	110.0	128.4	131.6	156.2	205.1	270.0	*
1916	124.4	276.2	134.1	54.1	124.5	128.9	148.0	127.3	151.8	189.8	293.2	*
1917	136.8	...	...	...	...	...	...	...	...	...	...	...
1918	127.1	226.7	122.4	64.2	134.5	121.8	147.9	131.7	150.1	222.7	351.4	633.5
1919	108.4	208.2	109.4	64.7	120.1	105.7	111.3	109.4	114.2	197.1	256.1	311.9
1920	107.4	242.8	111.2	58.7	130.6	107.4	96.1	102.5	125.9	181.5	242.4	643.4
1921	100.0	208.3	87.7	57.6	122.3	98.4	100.1	103.2	98.4	146.8	228.2	453.7
1922	99.1	229.6	94.4	50.6	107.3	100.6	96.6	117.4	123.7	154.8	225.2	431.5
1923	116.8	217.1	102.7	60.6	127.1	124.0	125.6	132.2	125.6	190.0	287.5	642.9
1924	121.3	259.7	108.5	58.1	131.5	121.1	134.8	139.2	148.3	178.4	325.1	755.3
1925	129.6	188.6	102.3	64.8	138.6	137.2	155.1	156.5	145.0	193.0	343.9	738.6
1926	134.3	206.4	103.7	64.1	137.9	149.1	158.7	163.7	166.5	200.9	331.3	786.9
1927	137.0	229.3	97.1	63.7	140.2	150.6	169.1	165.3	155.8	203.7	392.3	649.2
1928	126.5	219.1	82.0	55.6	129.7	141.5	157.6	161.7	138.6	211.9	328.0	676.6
1929	129.7	205.8	81.5	58.2	128.5	150.4	161.7	163.2	167.1	210.1	402.2	537.8
1930	126.3	217.2	82.7	53.0	124.1	142.4	159.6	158.1	162.9	233.7	382.3	525.3
1931	114.7	179.8	69.1	55.6	108.6	129.2	136.6	153.5	163.1	208.3	247.9	452.9
1932	114.8	185.6	77.4	53.3	113.7	128.7	138.1	135.5	161.5	194.9	296.8	570.3
1933	119.4	205.7	69.7	58.4	114.3	131.6	137.6	152.4	173.2	212.4	341.6	508.4
1934	131.5	216.9	84.6	57.6	123.4	145.7	152.9	179.6	194.0	234.4	341.8	570.5
1935	118.6	204.9	65.4	49.6	112.2	132.1	150.2	147.1	173.3	218.3	318.8	584.2
1936	133.1	253.1	64.3	51.2	117.8	146.2	170.5	180.6	205.5	249.3	368.0	695.5
1937	125.2	221.2	79.0	48.5	119.0	132.9	153.3	166.6	187.6	224.0	341.2	475.7
1938	109.4	201.7	61.9	48.4	107.8	121.2	130.2	135.4	158.3	178.3	243.5	528.0
1939	108.8	219.6	66.0	50.9	103.4	113.6	122.8	148.2	151.1	181.6	262.3	433.5

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	41.4	---	---	---	---	---	---	---	---	---	---	---
1911	53.5	---	---	---	---	---	---	---	---	---	---	---
1912	45.6	---	---	---	---	---	---	---	---	---	---	---
1913	50.3	---	---	---	---	---	---	---	---	---	---	---
1914	49.3	322.7	118.8	41.2	19.1	19.0	27.0	51.0	57.3	109.8	295.1	*
1915	46.4	213.9	114.0	39.2	23.2	19.9	26.8	31.8	71.0	105.1	247.0	741.0
1916	45.0	208.4	125.8	37.7	20.5	19.5	25.7	26.9	55.1	80.6	231.6	414.9
1917	54.9	---	---	---	---	---	---	---	---	---	---	---
1918	48.1	199.4	113.2	36.1	19.4	22.6	34.4	42.2	70.5	136.4	223.4	638.0
1919	42.7	188.4	104.3	33.7	18.6	25.2	22.5	34.9	55.2	97.8	276.2	379.3
1920	44.9	208.6	116.2	40.2	20.5	19.9	21.7	34.5	49.5	114.7	242.2	539.0
1921	37.5	188.0	88.7	27.8	16.1	15.8	24.4	27.9	57.1	126.5	203.7	438.0
1922	40.4	194.5	104.2	29.3	21.0	17.3	25.4	30.9	55.2	102.7	172.2	543.7
1923	42.8	180.3	103.7	30.2	23.0	24.5	21.2	33.0	61.5	109.9	255.5	653.4
1924	50.3	212.4	117.3	34.3	27.8	25.9	30.2	49.0	64.1	137.3	267.7	728.2
1925	47.6	227.2	98.1	30.1	30.4	27.6	30.8	41.4	70.5	97.6	254.0	622.9
1926	47.3	178.0	99.1	33.8	25.9	28.6	36.8	44.0	62.9	125.3	213.1	595.1
1927	42.6	182.6	81.8	30.4	20.7	23.5	33.1	43.6	69.1	116.3	209.1	538.7
1928	45.0	191.7	85.2	31.0	24.7	25.0	32.5	50.0	68.2	112.9	243.3	485.5
1929	44.3	194.4	80.6	29.4	24.5	29.6	32.2	47.0	84.0	106.6	192.6	502.7
1930	42.4	185.2	75.5	27.9	21.7	26.6	30.3	46.1	90.0	112.5	210.4	495.9
1931	41.6	170.0	74.0	25.3	22.0	28.1	31.6	44.7	81.7	124.0	243.4	353.4
1932	39.5	165.3	70.2	25.6	19.5	26.4	29.3	42.7	69.5	118.5	234.2	427.6
1933	40.4	181.6	69.7	25.5	22.0	24.8	33.3	44.1	80.5	113.3	152.9	397.4
1934	45.0	211.2	72.0	27.3	24.9	29.8	34.7	49.4	86.3	114.4	229.5	487.0
1935	41.1	179.6	69.2	24.3	20.8	26.3	30.8	48.9	71.2	114.7	213.4	570.2
1936	47.1	205.9	73.7	25.0	24.2	32.2	35.2	49.7	90.6	152.5	251.5	615.7
1937	42.3	206.4	63.7	25.8	22.4	26.4	32.9	46.2	76.1	118.0	211.6	489.2
1938	38.8	178.3	63.5	22.2	21.7	22.5	30.8	43.3	71.7	98.9	192.9	394.0
1939	36.5	166.8	57.0	18.9	23.3	23.6	24.1	34.7	68.0	101.0	189.4	451.5

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Notes: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) 1922-1929 total excludes deaths from legal executions.

3) 1909 total includes auto collisions with motor cycle.

4) 1926 total excludes food poisoning.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

MOTOR-VEHICLE ACCIDENTS

ICD Revision	Years in use	ICD Nos.
Fifth	1939-170	
Fourth	1930-1938 206, 208, 210, 211	
Third	1921-1929 188pt	
Second	1910-1920 175pt	
First	1900-1909 166pt	

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	0.4	*	*	0.5	*	0.4	*	0.8	*	*	*	*
1907	0.7	*	*	0.9	0.4	0.8	0.7	0.9	1.2	*	*	*
1908	0.8	*	*	1.0	0.5	0.6	0.8	0.9	1.7	1.6	*	*
1909	1.2	*	0.7	1.3	0.7	1.2	1.2	1.6	1.8	3.2	*	*
1910	1.8	*	0.6	2.4	1.1	1.3	1.9	1.9	2.6	3.7	4.8	*
1911	2.1	*	1.0	2.6	1.3	1.7	2.0	2.8	3.8	4.3	3.8	*
1912	2.8	*	2.2	3.6	2.0	1.9	3.0	3.3	4.6	5.2	5.2	*
1913	3.8	*	2.5	4.9	2.7	3.3	3.6	4.1	5.8	7.1	8.6	*
1914	4.2	*	2.7	5.0	3.1	3.2	4.1	5.0	6.4	8.4	7.8	*
1915	5.8	*	3.7	6.5	4.5	4.5	5.4	7.1	9.0	11.2	13.4	*
1916	7.1	*	5.0	7.7	5.4	5.9	6.7	8.3	11.5	13.5	14.7	16.8
1917	8.6	1.7	5.9	9.4	6.7	7.2	8.7	10.3	12.9	15.7	18.0	16.6
1918	9.3	2.0	7.1	10.9	6.7	6.8	8.1	10.5	14.7	17.9	21.0	19.4
1919	9.3	1.3	7.9	12.2	6.6	6.6	7.7	9.0	14.1	19.4	24.6	20.9
1920	10.3	1.8	9.1	13.0	7.8	7.9	7.7	10.2	14.8	22.1	28.4	25.5
1921	11.3	2.1	9.8	13.2	7.5	8.4	9.8	12.2	16.6	26.6	29.5	35.0
1922	12.4	2.5	9.9	13.6	9.1	9.6	10.4	13.3	18.9	28.9	37.6	33.4
1923	14.6	2.8	10.4	14.3	11.6	11.2	13.7	16.8	22.8	33.9	43.9	43.5
1924	15.3	3.9	11.7	14.7	12.5	11.5	13.5	17.5	23.5	36.6	47.7	39.5
1925	16.8	3.9	12.0	15.0	14.3	12.4	14.7	19.1	27.5	42.3	56.0	58.1
1926	19.7	5.6	12.3	15.9	18.1	15.8	18.2	23.8	29.6	46.1	62.7	56.4
1927	21.6	6.0	14.5	16.0	19.9	17.7	20.0	24.8	35.1	51.5	68.6	70.0
1928	23.2	5.8	14.3	15.5	21.9	19.1	21.5	28.0	38.9	55.5	75.8	68.7
1929	25.5	5.8	15.2	15.6	25.6	21.8	22.9	31.1	42.4	61.8	84.3	69.5
1930	26.7	6.6	14.5	14.7	27.4	24.0	23.7	32.2	44.3	66.3	85.0	83.4
1931	27.1	6.4	14.9	14.5	27.9	24.3	25.3	32.0	45.6	63.7	85.9	75.8
1932	23.6	4.5	12.9	12.0	22.6	21.6	22.4	28.1	40.2	57.0	77.4	75.5
1933	25.0	5.2	13.5	12.8	24.5	23.5	23.2	30.3	41.7	57.4	77.4	63.8
1934	28.6	5.5	13.0	13.2	28.2	27.3	26.9	35.9	48.7	66.0	84.2	71.6
1935	28.6	5.5	13.7	12.3	28.9	27.8	27.2	35.6	47.5	62.3	83.4	87.3
1936	29.7	6.4	14.5	12.7	30.5	28.3	27.9	36.2	50.1	64.6	85.4	69.4
1937	30.8	8.5	13.8	12.7	33.0	27.8	28.3	36.9	51.1	68.0	92.4	81.5
1938	25.1	6.2	12.0	10.8	25.3	22.8	22.0	29.9	41.9	57.9	79.4	71.7
1939	24.7	6.8	12.5	10.3	26.5	23.2	21.9	27.6	39.5	55.1	74.6	63.6

--- Data not available.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

Motor-vehicle Accidents

[Rates per 100,000 population for specified group.]

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	0.7	*	*	1.0	*	*	*	1.3	*	*	*	*
1907	1.1	*	*	1.2	0.7	1.1	1.2	1.5	2.0	*	*	*
1908	1.3	*	*	1.4	0.8	0.8	1.3	1.6	2.3	*	*	*
1909	1.8	*	*	2.1	1.0	1.8	1.7	2.4	2.8	4.5	*	*
1910	2.7	*	*	3.6	1.7	2.2	2.8	3.2	3.6	6.3	8.5	*
1911	3.3	*	1.5	4.2	2.1	2.7	3.1	4.0	5.8	6.9	*	*
1912	4.3	*	2.9	5.2	3.0	3.2	4.7	5.2	7.0	8.0	8.0	*
1913	5.9	*	3.1	7.7	4.3	5.1	5.8	6.3	8.5	10.7	13.1	*
1914	6.5	*	3.1	8.0	4.7	5.2	6.4	7.9	9.5	12.8	11.2	*
1915	8.8	*	4.3	10.0	7.3	7.2	8.4	10.5	12.9	17.3	21.4	*
1916	11.0	*	6.2	12.1	8.7	9.7	10.5	12.3	17.3	20.6	22.7	*
1917	13.2	*	7.1	14.0	10.8	11.6	13.8	15.9	18.8	21.9	28.0	*
1918	14.2	*	9.1	16.7	11.1	11.5	13.0	15.5	21.1	26.9	30.2	36.7
1919	13.8	*	10.0	18.5	10.5	10.6	11.7	12.5	19.3	28.3	36.2	38.0
1920	15.2	*	11.5	19.8	12.1	12.9	11.9	14.2	19.6	30.3	42.2	48.6
1921	16.6	*	12.6	19.5	12.0	13.6	15.4	17.4	22.0	35.9	43.7	60.8
1922	18.4	2.1	12.6	20.0	14.0	15.8	16.2	19.8	25.6	40.3	58.2	60.4
1923	21.5	3.1	13.0	20.5	17.9	17.9	20.9	25.0	31.7	46.5	66.1	67.8
1924	22.3	4.9	14.8	21.2	19.2	18.3	20.3	25.8	31.1	49.9	70.9	62.8
1925	24.3	4.0	14.9	21.4	21.6	19.7	22.1	27.6	36.3	58.3	80.8	94.7
1926	28.6	5.9	14.8	22.2	27.6	24.8	27.6	34.8	40.4	64.3	92.6	99.0
1927	31.6	6.6	18.5	22.2	31.2	28.3	30.4	36.0	49.1	70.1	99.0	115.9
1928	34.1	5.3	17.5	21.3	34.7	30.9	33.0	40.2	54.5	77.7	111.0	106.3
1929	37.9	6.0	18.4	21.7	38.8	35.1	35.4	46.3	59.9	87.2	132.0	104.3
1930	39.8	8.1	17.5	20.2	43.4	39.0	37.4	47.4	63.0	91.9	122.2	149.3
1931	40.4	6.0	18.1	19.9	43.5	39.3	39.4	47.1	65.7	89.0	129.9	128.8
1932	35.2	5.3	16.3	16.1	35.0	35.0	35.3	41.8	58.3	80.8	115.6	119.3
1933	37.7	5.3	17.0	17.7	38.0	38.6	36.6	45.8	61.0	82.6	114.1	114.3
1934	43.7	6.5	16.1	17.9	44.7	44.4	43.1	56.1	72.7	96.0	127.8	116.9
1935	43.6	5.0	16.8	17.0	44.8	45.5	43.2	54.6	70.9	92.7	125.5	152.8
1936	45.4	5.9	17.5	17.6	47.5	45.9	44.2	56.0	75.7	94.8	131.1	121.2
1937	47.0	8.7	16.1	17.7	51.9	45.1	44.5	57.0	78.1	100.5	140.4	137.5
1938	38.4	6.9	14.6	15.1	40.4	37.1	34.8	45.8	63.5	85.2	121.7	126.5
1939	37.9	6.4	14.0	14.4	42.5	38.1	35.0	42.4	59.6	80.3	116.3	106.3

... Data not available.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	0.1	*	*	*	*	*	*	*	*	*	*	*
1907	0.3	*	*	*	*	*	*	*	*	*	*	*
1908	0.4	*	*	*	*	*	*	*	*	*	*	*
1909	0.6	*	*	0.6	*	0.6	*	*	*	*	*	*
1910	0.8	*	*	1.3	*	*	0.8	*	1.4	*	*	*
1911	0.9	*	*	1.1	0.5	0.6	0.8	1.4	1.7	*	*	*
1912	1.3	*	1.5	2.0	1.0	0.6	1.0	1.3	2.0	2.5	*	*
1913	1.7	*	1.8	2.1	1.1	1.3	1.2	1.6	3.0	3.3	*	*
1914	1.8	*	2.2	2.0	1.4	1.1	1.7	1.6	3.1	3.9	*	*
1915	2.6	*	3.1	2.9	1.7	1.6	2.0	3.3	4.8	5.0	6.1	*
1916	3.0	*	3.7	3.2	2.1	1.9	2.5	3.7	5.2	6.2	7.5	*
1917	3.9	*	4.6	4.7	2.6	2.6	3.1	3.9	6.3	9.4	8.9	*
1918	4.3	*	5.0	5.1	2.8	2.5	2.8	4.8	7.8	8.7	12.7	*
1919	4.7	*	5.8	5.9	2.9	2.7	3.4	5.1	8.4	10.2	14.1	*
1920	5.2	*	6.7	6.1	3.5	2.9	3.2	5.8	9.6	13.6	15.7	*
1921	5.8	2.3	7.0	6.8	3.2	3.2	3.8	6.4	10.7	17.0	16.4	*
1922	6.2	2.9	7.2	7.2	4.3	3.4	4.0	6.1	11.6	17.2	18.8	*
1923	7.6	2.5	7.8	8.1	5.4	4.5	5.9	7.9	13.2	20.8	23.5	25.0
1924	8.2	2.9	8.5	8.2	6.0	4.8	6.2	8.5	15.2	22.8	26.3	21.7
1925	9.1	3.8	9.0	8.5	7.1	5.1	6.6	9.7	17.9	25.7	33.0	30.0
1926	10.4	5.3	9.7	9.4	8.6	6.7	8.0	11.9	17.8	27.1	35.0	23.6
1927	11.3	5.4	10.4	9.6	8.8	7.1	8.9	12.5	19.9	32.3	40.4	34.7
1928	12.0	6.4	10.9	9.7	9.3	7.3	9.3	14.7	21.8	32.5	44.9	39.9
1929	12.8	5.6	12.0	9.4	11.5	8.6	9.7	14.6	23.3	35.6	39.6	43.1
1930	13.3	5.2	11.5	9.0	11.7	9.1	9.3	15.7	24.0	39.6	50.0	33.8
1931	13.6	6.7	11.6	9.0	12.5	9.4	10.5	15.6	24.1	37.6	44.5	36.1
1932	11.7	3.7	9.4	7.8	10.4	8.4	8.8	13.2	20.8	32.4	41.5	42.6
1933	12.0	5.1	9.8	7.8	11.2	8.6	9.4	13.6	21.1	31.5	42.8	25.8
1934	13.1	4.6	9.7	8.2	12.1	10.5	10.2	14.2	23.0	35.4	43.4	37.5
1935	13.3	6.1	10.6	7.4	13.2	10.5	10.6	15.2	22.5	31.4	44.2	38.0
1936	13.9	6.9	11.5	7.8	13.8	11.1	11.2	15.1	22.9	34.1	42.9	30.4
1937	14.3	8.3	11.5	7.6	14.4	10.9	11.8	15.4	22.5	35.3	47.9	39.3
1938	11.6	5.5	9.4	6.5	10.6	8.8	9.0	12.9	19.1	30.5	40.5	30.6
1939	11.4	7.1	11.0	6.0	10.7	8.6	8.7	11.9	18.3	29.9	36.4	31.5

--- Data not available.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	1.8	---	---	---	---	---	---	---	---	---	---	---
1911	2.1	---	---	---	---	---	---	---	---	---	---	---
1912	2.9	---	---	---	---	---	---	---	---	---	---	---
1913	3.9	---	---	---	---	---	---	---	---	---	---	---
1914	4.3	*	2.7	5.1	3.1	3.3	4.2	5.2	6.5	8.5	7.8	*
1915	5.9	*	3.8	6.6	4.6	4.6	5.4	7.2	9.1	11.3	13.5	*
1916	7.3	*	5.2	8.0	5.6	6.1	6.9	8.5	11.8	13.6	15.2	17.2
1917	8.9	---	---	---	---	---	---	---	---	---	---	---
1918	9.5	2.1	7.3	11.3	7.0	7.1	8.3	10.8	15.0	18.2	21.2	18.4
1919	9.8	1.4	8.4	13.0	7.0	6.8	7.9	9.4	14.3	19.8	25.2	23.0
1920	10.8	1.9	9.6	13.7	8.1	8.1	8.0	10.7	15.0	22.8	29.5	26.7
1921	11.8	2.1	10.4	14.0	7.9	8.7	10.2	12.5	17.0	27.3	30.1	38.3
1922	12.9	2.7	10.5	14.6	9.7	9.8	10.5	13.7	19.4	29.6	39.3	35.7
1923	15.1	2.8	11.1	15.2	11.9	11.3	13.8	17.2	23.2	35.0	45.0	43.8
1924	15.6	4.0	12.2	15.5	12.7	11.4	13.4	17.3	23.8	37.2	48.3	38.8
1925	17.1	4.0	12.6	15.6	14.5	12.2	14.4	19.1	28.0	42.9	57.2	59.9
1926	20.0	6.0	12.7	16.5	18.4	15.5	17.9	23.9	29.9	46.5	64.5	58.7
1927	22.0	6.1	15.1	16.6	20.7	17.6	19.7	24.3	35.7	52.0	69.3	69.9
1928	23.7	6.0	15.0	16.2	22.8	19.1	21.3	27.8	39.3	56.4	78.9	66.9
1929	26.0	6.0	16.0	16.3	26.4	21.4	22.6	30.8	42.5	63.0	86.1	70.9
1930	27.2	7.2	15.2	15.2	28.1	23.8	23.4	32.0	44.4	67.3	87.1	87.9
1931	27.6	6.6	15.8	15.0	28.9	23.9	24.9	31.8	45.7	64.2	88.5	80.9
1932	23.9	4.5	13.4	12.2	23.4	21.3	21.8	27.6	40.3	57.5	79.4	76.1
1933	25.2	5.5	14.0	13.1	25.0	23.2	22.6	29.9	41.5	57.7	78.3	65.0
1934	28.8	5.8	13.3	13.3	28.9	26.7	26.4	35.5	48.3	66.5	85.9	71.4
1935	29.0	5.6	14.1	12.8	29.8	27.6	26.4	35.5	47.4	62.9	84.8	87.0
1936	30.0	6.2	15.3	13.2	31.2	27.9	26.6	35.8	49.7	65.2	86.9	70.4
1937	31.2	8.7	14.6	13.2	33.9	27.6	27.5	36.4	50.9	68.7	93.7	84.4
1938	25.3	6.4	12.5	11.1	25.7	22.3	21.0	29.6	41.7	59.0	81.2	75.6
1939	24.9	7.2	12.9	10.4	27.1	22.9	21.2	27.0	39.3	55.8	76.6	64.9

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	2.7	---	---	---	---	---	---	---	---	---	---	---
1911	3.3	---	---	---	---	---	---	---	---	---	---	---
1912	4.3	---	---	---	---	---	---	---	---	---	---	---
1913	6.0	---	---	---	---	---	---	---	---	---	---	---
1914	6.6	*	3.2	8.2	4.8	5.3	6.5	8.3	9.6	12.9	11.0	*
1915	9.0	*	4.5	10.3	7.4	7.4	8.6	10.5	12.9	17.5	21.5	*
1916	11.3	*	6.4	12.6	9.1	10.0	10.7	12.8	17.7	20.7	23.4	*
1917	13.6	---	---	---	---	---	---	---	---	---	---	---
1918	14.6	*	9.3	17.2	11.5	11.8	13.3	16.0	21.6	27.1	30.1	35.1
1919	14.4	*	10.7	19.6	11.0	10.9	11.9	13.0	19.5	28.8	37.7	41.8
1920	15.8	*	12.2	20.8	12.5	13.2	12.3	14.9	19.9	31.4	43.8	50.4
1921	17.3	*	13.4	20.6	12.6	14.1	15.8	17.8	22.6	37.2	44.6	66.4
1922	19.2	2.2	13.3	21.4	14.8	16.1	16.5	20.3	26.4	41.5	60.7	65.0
1923	22.1	3.1	13.7	21.6	18.3	17.9	21.0	25.5	32.3	48.4	68.3	65.9
1924	22.6	5.2	15.3	22.2	19.3	18.2	20.1	25.5	31.6	50.8	71.4	64.6
1925	24.7	4.0	15.5	22.1	21.9	19.6	21.6	27.6	37.1	59.2	82.9	99.4
1926	29.0	6.2	15.3	23.0	27.9	24.4	27.2	35.0	40.9	65.1	95.5	103.1
1927	32.0	6.7	19.1	22.9	32.2	28.0	30.0	35.4	50.0	70.6	98.8	114.1
1928	34.8	5.3	18.5	22.2	35.9	31.0	32.6	39.8	55.4	78.6	114.4	99.3
1929	38.4	6.2	19.3	22.6	40.9	34.4	34.7	45.6	59.9	88.9	134.0	103.3
1930	40.3	8.6	18.5	20.9	44.1	38.8	36.7	46.9	63.4	93.4	124.4	155.0
1931	41.0	6.4	19.3	20.4	44.9	38.7	38.8	46.7	66.2	89.7	133.8	135.3
1932	35.5	5.0	17.0	16.3	35.8	34.6	34.3	41.2	58.6	81.7	119.1	117.5
1933	37.7	5.4	17.7	18.0	38.4	37.8	35.4	45.1	60.7	82.8	114.5	114.1
1934	43.8	6.7	16.7	18.1	45.1	43.2	42.1	55.4	72.3	96.2	129.7	114.9
1935	43.9	5.2	17.3	17.7	45.6	44.9	41.7	54.3	70.8	93.4	127.0	153.6
1936	45.4	5.5	18.4	18.2	48.2	45.2	41.7	55.0	75.0	95.7	133.0	122.9
1937	47.3	8.8	16.7	18.1	52.9	44.5	43.0	56.1	77.9	101.3	142.3	142.0
1938	38.5	7.0	15.1	15.4	40.6	36.1	33.0	45.5	63.3	86.5	124.4	132.3
1939	38.0	6.8	14.4	14.4	43.4	37.4	33.6	41.2	59.4	81.3	119.3	106.0

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	0.8	---	---	---	---	---	---	---	---	---	---	---
1911	0.9	---	---	---	---	---	---	---	---	---	---	---
1912	1.3	---	---	---	---	---	---	---	---	---	---	---
1913	1.7	---	---	---	---	---	---	---	---	---	---	---
1914	1.9	*	2.3	2.1	1.5	1.1	1.7	1.7	3.1	4.1	*	*
1915	2.6	*	3.0	2.9	1.8	1.6	2.0	3.3	4.9	5.0	6.3	*
1916	3.2	*	3.8	3.3	2.2	1.9	2.6	3.8	5.3	6.4	7.8	*
1917	4.1	---	---	---	---	---	---	---	---	---	---	---
1918	4.4	*	5.3	5.3	3.0	2.6	2.8	4.9	7.8	9.1	13.1	*
1919	5.0	*	6.1	6.2	3.2	2.8	3.5	5.4	8.6	10.6	13.9	*
1920	5.5	*	6.9	6.5	3.8	2.9	3.3	6.0	9.8	14.1	16.4	*
1921	6.1	2.4	7.4	7.3	3.4	3.3	3.9	6.6	11.0	17.2	16.9	*
1922	6.5	3.2	7.5	7.8	4.6	3.5	4.0	6.4	11.9	17.4	19.8	*
1923	7.9	2.4	8.4	8.6	5.5	4.5	6.0	8.2	13.4	21.3	23.7	26.6
1924	8.4	2.8	9.0	8.7	6.2	4.7	6.2	8.5	15.5	23.3	27.1	*
1925	9.4	4.0	9.5	9.0	7.2	4.8	6.6	9.9	18.2	26.2	33.5	29.1
1926	10.7	5.7	10.1	9.8	8.8	6.6	7.9	12.1	18.0	27.4	36.0	24.2
1927	11.6	5.5	10.9	10.1	9.3	7.0	8.7	12.3	20.3	32.8	42.1	35.6
1928	12.3	6.7	11.4	10.1	9.9	7.2	9.3	14.7	22.0	33.4	45.8	42.0
1929	13.2	5.7	12.5	9.8	12.0	8.4	9.6	14.6	23.8	36.4	41.1	46.3
1930	13.7	5.7	11.8	9.4	12.3	9.0	9.3	15.9	24.1	40.2	52.0	37.6
1931	13.9	6.9	12.3	9.4	13.1	9.1	10.3	15.8	23.9	38.0	45.9	40.0
1932	11.9	4.0	9.6	8.0	11.1	8.2	8.6	13.0	20.9	32.9	42.3	45.1
1933	12.3	5.5	10.1	8.1	11.7	8.6	9.2	13.4	21.1	32.0	44.2	28.0
1934	13.4	4.9	9.9	8.3	12.7	10.4	10.0	14.0	22.9	36.1	44.9	38.6
1935	13.7	6.0	10.8	7.8	14.1	10.5	10.6	15.3	22.7	32.1	45.5	36.8
1936	14.2	7.0	12.2	8.2	14.3	10.8	11.0	15.2	22.9	34.4	44.0	30.9
1937	14.7	8.5	12.4	8.0	14.9	11.0	11.6	15.4	22.4	36.2	48.9	40.9
1938	11.8	5.8	9.8	6.6	10.9	8.6	8.7	12.7	18.9	31.5	41.6	32.9
1939	11.6	7.6	11.4	6.3	10.9	8.5	8.6	11.9	18.1	30.6	37.6	32.4

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	*	---	---	---	---	---	---	---	---	---	---	---
1911	1.6	---	---	---	---	---	---	---	---	---	---	---
1912	2.4	---	---	---	---	---	---	---	---	---	---	---
1913	2.4	---	---	---	---	---	---	---	---	---	---	---
1914	2.0	*	*	*	*	*	*	*	*	*	*	*
1915	3.8	*	*	*	*	*	*	*	*	*	*	*
1916	3.5	*	*	3.6	*	3.5	4.2	*	*	*	*	*
1917	4.4	---	---	---	---	---	---	---	---	---	---	---
1918	5.7	*	4.1	6.5	4.1	4.0	6.2	6.6	9.5	*	*	*
1919	4.7	*	*	5.0	2.8	4.4	5.6	4.8	10.8	12.4	*	*
1920	5.3	*	3.8	5.8	4.5	5.4	4.8	4.6	11.0	*	*	*
1921	5.7	*	3.5	5.6	3.8	4.7	6.4	8.6	9.2	14.7	*	*
1922	6.8	*	4.8	5.3	4.5	7.5	9.0	9.5	10.8	18.2	*	*
1923	9.7	*	4.1	7.0	9.2	10.1	12.5	12.7	17.0	14.7	*	*
1924	12.3	*	7.0	8.1	10.7	12.3	15.0	19.7	18.7	26.1	39.0	*
1925	13.5	*	7.2	8.9	12.2	13.9	17.7	19.1	19.5	30.7	35.6	*
1926	16.4	*	7.9	10.1	15.3	18.6	21.3	22.6	25.3	37.7	32.1	*
1927	17.8	*	9.4	10.8	13.2	18.8	23.3	29.6	27.0	43.8	57.0	*
1928	18.1	*	8.3	10.2	14.9	18.5	23.3	30.4	32.5	41.7	45.3	84.4
1929	21.5	*	9.1	10.2	19.4	25.3	25.9	35.0	41.0	41.5	57.2	*
1930	22.1	*	9.2	10.1	21.5	25.2	26.6	34.4	42.8	49.6	51.2	*
1931	22.5	*	7.6	10.7	19.5	27.6	28.5	33.9	44.3	55.0	42.7	*
1932	20.9	*	9.7	10.2	16.3	24.1	28.0	32.7	39.1	47.3	42.9	*
1933	23.3	*	9.9	10.7	20.3	26.1	29.3	34.8	44.8	53.3	63.1	*
1934	26.8	*	10.3	12.0	23.3	32.1	31.9	40.7	53.8	59.4	56.1	73.8
1935	24.9	*	10.5	8.3	21.7	29.3	34.0	37.1	48.5	53.1	60.2	90.3
1936	27.8	*	8.9	8.9	25.3	31.4	40.2	40.6	56.0	56.7	60.6	60.4
1937	27.4	*	8.5	9.7	26.2	29.7	35.7	41.9	54.6	57.6	69.5	*
1938	23.5	*	9.0	9.0	22.5	27.1	31.4	33.4	45.1	43.8	48.3	*
1939	23.1	*	9.3	9.3	22.0	25.9	28.3	34.8	43.1	46.2	41.2	*

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	4	...	...	...	...	...	...	...	...	...	...	...
1911	2.4	...	...	...	...	...	...	...	...	...	...	...
1912	4.0	...	...	...	...	...	...	...	...	...	...	...
1913	3.6	...	...	...	...	...	...	...	...	...	...	...
1914	3.5	*	*	*	*	*	*	*	*	*	*	*
1915	5.6	*	*	*	*	*	*	*	*	*	*	*
1916	5.7	*	*	5.7	*	*	7.5	*	*	*	*	*
1917	7.2	...	...	...	...	...	...	...	...	...	...	...
1918	9.0	*	10.2	7.3	7.0	9.4	8.9	*	*	*	*	*
1919	7.4	*	7.2	5.3	7.5	9.3	6.9	15.6	*	*	*	*
1920	8.4	*	9.6	8.1	9.0	7.5	6.2	15.6	*	*	*	*
1921	8.9	*	9.2	6.3	8.1	10.3	12.5	13.0	*	*	*	*
1922	10.5	*	5.4	8.3	7.0	12.9	13.1	14.6	14.9	22.7	*	*
1923	15.1	*	6.2	10.5	14.3	17.0	20.2	19.5	23.0	*	*	*
1924	19.0	*	10.0	12.4	18.0	19.3	23.1	28.8	24.7	35.5	63.9	*
1925	20.1	*	9.2	14.0	18.4	20.5	27.6	28.6	25.1	43.4	*	*
1926	25.1	*	9.4	15.1	24.8	29.7	32.7	33.5	33.6	51.8	*	*
1927	27.6	*	12.7	16.3	22.6	30.3	35.5	42.4	37.2	61.5	101.9	*
1928	27.6	*	9.1	14.1	25.0	29.9	37.2	44.1	43.5	63.5	60.3	*
1929	33.6	*	10.9	14.4	31.6	41.3	41.9	52.6	59.9	60.1	100.6	*
1930	34.9	*	9.4	14.6	37.3	41.2	44.7	52.0	58.2	67.5	87.2	*
1931	34.7	*	8.6	16.1	31.6	44.6	45.2	51.3	58.4	76.8	65.8	*
1932	32.6	*	11.1	14.2	28.3	38.9	45.5	48.4	54.6	67.2	58.1	*
1933	37.4	*	11.8	15.8	34.9	45.4	48.2	52.6	64.3	80.6	108.0	*
1934	43.0	*	12.1	16.7	40.6	54.5	53.1	63.1	77.1	91.8	96.0	*
1935	40.5	*	12.5	11.8	38.0	50.2	57.8	57.2	71.9	82.5	101.3	146.0
1936	44.7	*	10.6	13.0	42.0	51.2	68.0	65.4	83.9	81.9	98.9	*
1937	44.1	*	11.9	14.4	43.6	50.3	58.4	66.3	80.2	90.2	109.0	*
1938	37.5	*	11.4	12.8	38.4	45.5	51.6	49.6	65.7	69.1	76.5	*
1939	37.1	*	11.0	14.4	35.6	43.8	47.7	56.1	62.5	69.2	67.7	*

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

Motor-vehicle Accidents

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	*	...	...	...	...	...	...	...	...	...	...	...
1911	*	...	...	...	...	...	...	...	...	...	...	...
1912	*	...	...	...	...	...	...	...	...	...	...	...
1913	*	...	...	...	...	...	...	...	...	...	...	...
1914	*	*	*	*	*	*	*	*	*	*	*	*
1915	2.0	*	*	*	*	*	*	*	*	*	*	*
1916	1.1	*	*	*	*	*	*	*	*	*	*	*
1917	1.6	...	...	...	...	...	...	...	...	...	...	...
1918	2.3	*	*	*	*	*	*	*	*	*	*	*
1919	2.0	*	*	2.7	*	*	*	*	*	*	*	*
1920	2.2	*	*	*	*	*	*	*	*	*	*	*
1921	2.4	*	*	*	*	*	*	*	*	*	*	*
1922	3.2	*	*	2.4	2.2	*	4.7	*	*	*	*	*
1923	4.3	*	*	3.6	4.5	3.7	4.4	*	*	*	*	*
1924	5.6	*	*	3.8	3.9	5.9	6.5	8.6	11.0	*	*	*
1925	6.8	*	5.2	3.9	6.4	7.8	7.5	7.5	12.3	*	*	*
1926	7.8	*	6.3	5.2	6.5	8.1	9.6	9.7	14.5	*	*	*
1927	8.0	*	6.0	5.4	4.5	8.0	11.0	14.5	13.6	22.7	*	*
1928	8.5	*	7.4	6.4	5.5	7.9	9.5	14.3	17.9	*	*	*
1929	9.5	*	7.4	6.1	8.2	10.5	10.2	14.5	16.1	20.1	*	*
1930	9.4	*	8.9	5.7	7.1	10.3	8.8	13.9	22.7	29.4	*	*
1931	10.4	*	6.6	5.4	8.5	11.7	12.2	14.0	26.2	30.4	*	*
1932	9.4	*	8.2	6.2	5.4	10.2	11.0	14.9	19.5	25.1	*	*
1933	9.3	*	8.1	5.6	7.1	8.3	11.1	14.8	20.2	22.8	*	*
1934	10.8	*	8.5	7.4	7.7	11.4	11.5	15.8	24.9	23.4	*	*
1935	9.6	*	8.6	4.7	7.0	9.9	11.0	14.8	19.8	20.8	*	*
1936	11.3	*	7.2	4.7	10.1	13.2	13.4	13.5	22.2	29.0	*	*
1937	11.1	*	5.2	5.0	10.6	10.8	14.0	15.3	24.1	22.0	*	*
1938	10.0	*	6.7	5.1	8.1	10.3	12.1	15.8	20.9	16.2	*	*
1939	9.5	*	7.7	4.3	9.6	9.6	9.7	11.9	20.5	21.2	*	*

... Data not available. No original tables for the time period 1906-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Notes: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices, but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes ...	196	266	309
Male....	164	226	259
Female....	32	40	50

3) For 1926, deaths from automobile collisions with motorcycles included in total but not distributed by age.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ICD Revision	Years in use	ICD Nos.
Fifth	1939-1969	171-176, 178-195
Fourth	1930-1938	176, 178-195, 201-205, 207, 209, 212-214
Third	1921-1929	176-187, 188pt, 189, 192-196, 201, 202pt, 203
Second	1910-1920	165-174, 175pt, 176-181, 185, 186
First	1900-1909	164, 165, 166pt, 167-175, 176pt

ALL RACES, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	72.3	128.8	75.3	38.3	53.4	62.6	78.3	80.6	90.9	131.5	274.8	668.9
1901	83.8	163.6	82.5	40.4	53.9	69.4	83.8	103.9	128.4	172.6	359.7	863.8
1902	72.5	100.0	68.4	37.5	52.7	65.4	77.9	84.1	100.9	139.4	275.4	799.4
1903	81.4	118.1	78.3	41.7	57.7	72.1	85.8	92.2	108.5	159.2	358.5	944.2
1904	85.4	122.2	86.2	48.7	60.9	74.8	86.2	94.4	112.3	158.5	393.3	1,034.2
1905	81.3	133.6	79.9	43.5	59.5	68.5	84.3	87.8	114.4	160.0	344.9	1,083.3
1906	94.0	131.1	87.2	45.1	74.8	90.6	98.8	104.1	120.1	169.5	381.8	1,058.2
1907	94.1	131.8	92.3	43.2	76.0	90.2	104.2	106.8	124.0	163.6	373.3	1,069.8
1908	82.1	127.7	87.2	40.1	64.0	73.9	83.6	94.1	110.1	152.9	364.5	980.0
1909	78.7	117.8	84.0	36.9	57.0	70.8	82.1	90.3	103.2	150.5	383.0	1,108.2
1910	82.4	113.8	82.0	35.6	63.1	78.9	88.4	93.9	106.7	163.4	371.5	1,077.1
1911	81.5	133.7	84.9	34.7	59.1	71.5	84.5	95.3	113.1	166.5	397.5	1,207.8
1912	78.0	112.4	83.4	35.0	58.6	70.2	78.2	89.1	107.6	153.7	380.8	1,087.2
1913	79.9	118.0	81.7	35.2	63.0	73.2	80.9	90.6	108.6	157.1	375.5	1,047.2
1914	72.5	118.0	78.5	32.2	54.0	63.8	71.5	79.5	97.8	151.2	368.7	1,042.6
1915	67.7	101.8	75.5	31.6	49.2	57.5	66.3	73.9	90.4	144.2	363.3	1,027.5
1916	74.5	121.4	78.1	31.8	54.1	64.2	76.5	86.6	104.4	156.1	377.2	1,095.7
1917	77.4	133.1	78.9	34.3	56.6	64.8	75.8	90.4	110.6	165.8	400.4	1,040.1
1918	72.2	103.2	76.3	35.9	55.1	59.6	68.1	78.4	99.0	159.4	367.0	1,035.3
1919	61.8	95.5	73.6	33.2	48.2	49.0	56.6	63.0	80.2	127.8	334.5	982.9
1920	59.7	98.4	71.1	31.3	47.3	46.0	51.2	58.9	74.9	134.2	362.6	1,089.4
1921	55.5	88.8	64.2	28.6	43.4	42.5	48.2	54.3	71.0	124.2	348.5	1,116.7
1922	56.0	88.6	63.4	27.4	42.0	42.2	49.3	58.7	74.5	127.5	359.6	1,174.0
1923	59.8	90.6	62.1	27.9	44.6	46.1	54.6	64.9	81.3	137.6	381.6	1,234.5
1924	58.6	97.1	63.4	27.3	44.1	42.6	53.7	63.3	78.6	134.0	367.7	1,254.1
1925	59.9	94.3	58.4	26.9	45.0	44.5	55.3	67.2	80.7	136.8	387.2	1,291.4
1926	57.6	100.7	56.2	25.1	40.7	41.8	53.2	65.0	80.2	131.9	391.1	1,270.7
1927	55.6	88.9	54.1	24.6	39.5	40.2	51.4	61.3	77.0	129.5	378.4	1,204.3
1928	55.0	94.2	51.5	24.5	37.7	40.6	49.1	61.7	73.5	130.5	396.1	1,196.6
1929	54.3	93.1	49.3	23.5	37.2	41.0	48.9	59.0	75.8	130.6	378.9	1,159.3
1930	53.1	96.8	46.7	21.4	34.4	38.4	46.4	58.2	75.9	131.8	392.8	1,218.6
1931	50.7	87.5	47.3	20.4	31.4	33.3	41.6	56.5	75.2	131.4	393.3	1,177.4
1932	47.2	85.7	44.6	20.5	29.8	30.1	36.3	47.8	67.6	123.4	377.6	1,187.8
1933	46.9	90.4	43.0	20.5	28.2	29.7	35.6	48.2	67.3	121.7	388.6	1,164.6
1934	50.8	101.4	43.3	20.8	28.6	31.0	37.8	54.1	75.0	136.9	431.9	1,246.3
1935	49.3	99.6	43.0	19.9	27.2	30.2	37.4	50.3	69.7	130.2	417.9	1,286.5
1936	55.7	118.5	44.2	20.1	27.5	32.7	41.0	57.6	82.3	155.9	498.6	1,474.4
1937	50.4	104.1	42.2	19.9	27.3	30.7	36.5	49.1	69.7	130.9	444.6	1,290.1
1938	46.7	104.8	39.5	17.4	25.5	27.0	32.7	44.4	61.3	121.3	421.5	1,260.6
1939	45.6	103.0	38.2	17.5	24.0	26.2	30.5	41.6	59.0	117.3	421.1	1,251.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ALL RACES, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	110.2	138.4	89.8	56.5	92.8	108.5	122.6	129.2	140.9	181.5	283.7	515.9
1901	124.0	102.3	89.4	59.6	93.9	118.0	138.2	163.3	187.6	218.9	378.8	686.3
1902	110.2	103.2	78.1	56.0	90.0	112.2	130.0	140.4	152.3	194.2	301.9	613.5
1903	123.0	133.6	86.4	63.3	100.3	124.1	143.5	151.2	164.5	215.1	347.3	748.6
1904	125.1	137.0	99.3	67.7	99.7	122.2	139.4	151.6	169.2	214.1	414.4	917.3
1905	122.8	158.3	89.6	63.7	104.3	118.1	142.6	147.2	175.9	208.9	344.1	870.4
1906	145.0	147.7	94.3	67.4	132.4	156.7	168.2	169.3	190.5	234.6	380.5	901.7
1907	145.6	142.7	99.5	64.0	136.3	157.1	178.3	178.7	193.8	213.7	368.5	922.3
1908	124.5	146.0	97.4	59.3	111.6	126.8	141.9	154.6	171.4	191.8	374.5	820.4
1909	118.5	126.1	92.3	53.5	100.6	123.0	139.6	148.7	159.4	193.6	364.3	939.0
1910	126.4	126.5	91.2	51.9	111.9	138.7	151.9	154.9	167.6	210.2	355.3	877.7
1911	121.3	144.2	94.4	50.4	101.7	123.1	144.3	154.4	173.0	211.2	372.2	923.8
1912	117.0	122.1	93.4	50.9	102.4	122.1	132.3	148.2	165.4	197.0	332.0	829.2
1913	120.7	126.6	90.1	50.4	111.4	127.1	138.9	149.5	170.5	201.3	343.2	792.3
1914	107.8	125.8	88.4	46.7	94.4	111.1	121.9	129.7	149.7	192.0	329.4	789.4
1915	100.2	110.0	85.9	46.3	85.7	100.4	113.4	120.5	136.2	179.2	327.1	808.4
1916	111.2	130.0	84.8	45.2	96.1	112.0	131.2	142.3	159.3	198.7	355.6	923.2
1917	113.7	143.1	83.6	47.1	99.0	113.8	129.4	146.2	167.3	206.8	359.4	768.0
1918	107.6	112.6	83.4	50.7	101.0	108.3	114.6	126.2	150.8	204.9	330.3	842.5
1919	89.5	108.3	78.4	46.7	84.8	85.4	95.6	100.9	119.3	160.1	291.5	762.4
1920	84.4	109.4	74.1	43.2	83.0	79.9	84.5	93.5	109.0	161.0	307.2	856.5
1921	78.7	96.5	70.1	40.6	76.0	73.5	80.1	86.5	103.1	148.5	297.6	832.7
1922	79.4	97.7	67.1	38.2	72.5	73.3	83.1	94.0	111.5	151.5	306.9	846.6
1923	85.6	99.8	66.7	38.9	77.5	80.2	91.7	105.0	122.6	168.9	316.6	889.7
1924	83.0	106.0	68.0	37.4	75.8	74.0	90.2	102.2	117.5	157.6	314.2	905.4
1925	85.1	100.2	61.5	37.8	76.9	76.8	92.6	108.6	120.2	167.9	324.6	1,007.4
1926	81.9	108.9	60.0	35.3	71.0	72.8	90.4	105.2	120.7	157.1	323.6	906.7
1927	79.8	97.0	59.0	35.0	69.2	70.6	87.1	101.2	115.4	155.8	324.2	888.4
1928	78.1	102.6	55.8	34.9	65.4	71.3	84.5	100.7	108.9	152.4	327.7	853.7
1929	77.0	102.4	52.2	33.4	65.0	71.7	83.6	96.7	112.8	151.2	312.6	788.0
1930	74.6	105.2	50.8	30.0	59.9	67.5	79.8	95.3	112.0	154.4	316.7	877.3
1931	70.3	95.3	50.9	28.8	54.3	58.4	71.0	92.3	110.1	151.4	323.2	872.4
1932	64.0	93.0	48.6	29.0	51.1	52.1	61.1	76.9	98.1	135.6	308.5	849.3
1933	63.1	97.8	45.5	29.3	48.0	50.9	59.0	77.5	96.6	136.7	311.0	862.9
1934	68.7	109.7	47.7	29.5	48.8	52.9	63.5	87.8	109.6	154.0	365.2	968.6
1935	65.8	109.5	45.7	27.8	45.9	51.6	63.3	80.8	100.7	145.7	349.3	943.4
1936	73.9	129.8	47.2	27.8	47.0	55.8	69.2	93.4	120.1	175.2	416.1	1,106.6
1937	67.8	116.7	46.8	27.7	47.0	53.6	62.4	80.2	103.6	145.1	375.1	1,010.4
1938	61.1	116.4	43.7	24.6	44.3	46.7	55.0	71.8	88.2	131.0	329.4	953.7
1939	59.8	116.2	43.9	25.6	41.6	45.6	52.3	67.9	84.6	129.0	336.3	899.2

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ALL RACES, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	34.1	119.1	60.6	20.0	16.3	16.1	20.0	29.9	41.9	84.5	266.7	775.7
1901	43.5	134.3	75.5	21.1	16.1	20.1	26.1	41.7	70.4	128.7	342.4	992.6
1902	34.5	96.7	58.6	18.8	17.3	17.9	22.6	25.0	50.4	87.6	251.6	934.8
1903	39.5	102.3	70.0	19.8	17.2	19.1	24.6	30.0	53.3	106.3	368.6	1,087.3
1904	45.5	107.2	72.9	29.5	23.7	26.2	29.6	34.1	56.1	106.0	374.4	1,119.9
1905	39.4	108.4	70.1	23.1	16.5	17.5	22.3	25.0	53.5	113.8	345.6	1,240.2
1906	41.0	114.1	79.9	22.5	17.7	19.6	22.4	32.5	47.5	105.9	383.0	1,175.4
1907	40.6	120.6	85.0	22.1	15.9	18.0	22.6	27.9	51.8	114.7	377.6	1,180.7
1908	37.5	109.0	76.7	20.6	15.9	15.7	18.6	26.7	45.4	114.4	355.4	1,103.7
1909	36.8	109.3	75.5	20.1	12.8	13.4	18.3	25.4	43.7	107.7	400.1	1,240.4
1910	35.9	100.7	72.5	19.0	13.2	12.4	17.4	25.3	41.5	116.7	386.4	1,233.7
1911	39.4	122.9	75.1	18.7	15.9	14.7	18.0	28.7	48.3	121.4	420.8	1,432.6
1912	36.8	102.4	73.1	18.8	14.2	13.3	17.9	22.4	44.9	110.1	425.6	1,289.8
1913	36.9	109.3	73.2	19.8	14.2	14.5	16.6	24.0	41.3	112.3	404.9	1,245.6
1914	35.4	110.0	68.4	17.5	13.3	12.7	15.8	22.7	41.1	109.6	404.7	1,239.5
1915	33.6	93.3	64.9	16.8	12.6	11.3	14.2	21.2	40.3	108.5	396.4	1,196.3
1916	36.2	112.7	71.3	18.1	12.5	13.6	16.4	23.4	44.3	112.7	396.7	1,226.8
1917	39.6	122.8	74.1	21.4	15.0	13.8	17.3	27.2	48.3	123.8	437.5	1,245.6
1918	36.6	93.6	69.1	20.8	14.8	13.6	17.5	23.9	42.1	112.7	400.3	1,180.5
1919	33.3	82.3	68.6	19.5	13.6	12.7	14.1	19.8	37.0	94.4	373.6	1,147.8
1920	34.0	87.1	68.0	19.2	12.6	11.6	14.5	19.7	37.4	106.4	413.5	1,264.9
1921	31.4	80.7	58.0	16.4	11.5	11.2	13.1	18.2	36.0	99.2	395.3	1,331.7
1922	32.0	79.2	59.5	16.4	12.2	11.3	12.8	19.3	34.5	102.7	407.8	1,422.4
1923	33.4	81.1	57.4	16.7	12.3	12.3	14.6	20.6	36.4	105.2	441.3	1,498.5
1924	33.5	88.1	58.6	17.1	12.8	11.1	14.3	20.8	36.2	109.5	416.9	1,521.0
1925	33.9	88.2	55.2	15.9	13.6	12.2	15.3	22.0	37.7	104.6	444.9	1,509.2
1926	32.7	92.3	52.2	14.8	10.7	10.8	13.3	21.2	36.0	105.9	453.4	1,550.4
1927	30.8	80.7	49.0	14.1	10.1	9.7	13.3	17.9	35.0	102.2	428.7	1,447.1
1928	31.4	85.5	46.9	14.0	10.4	10.0	11.7	19.2	34.8	107.8	459.9	1,458.9
1929	31.1	83.4	46.2	13.4	9.9	10.6	12.3	17.8	35.6	109.2	441.0	1,440.7
1930	31.0	88.0	42.5	12.6	9.4	9.6	11.4	17.7	36.9	108.4	464.5	1,474.9
1931	30.6	79.6	43.6	11.8	8.8	8.6	10.8	17.5	37.6	110.8	459.1	1,406.3
1932	30.0	78.2	40.5	11.8	8.9	8.4	10.4	16.3	34.9	110.9	442.3	1,441.7
1933	30.4	82.8	40.5	11.4	8.9	8.9	11.1	16.6	35.8	106.2	461.5	1,391.8
1934	32.4	92.7	38.7	11.9	8.9	9.5	11.2	17.7	37.9	119.4	494.4	1,455.4
1935	32.4	89.4	40.3	11.7	8.8	9.1	10.7	17.5	36.7	114.6	482.0	1,544.9
1936	37.2	106.9	41.2	12.2	8.3	10.0	12.0	19.3	42.2	136.4	575.1	1,751.2
1937	32.7	91.2	37.4	11.7	8.0	8.3	9.9	15.9	33.7	116.7	508.9	1,500.7
1938	32.1	93.0	35.2	10.1	7.1	7.6	10.0	15.1	32.9	111.6	506.4	1,491.5
1939	31.3	89.5	32.3	9.3	6.6	7.2	8.5	13.7	32.0	105.6	498.9	1,520.7

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

WHITE, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	82.2	---	---	---	---	---	---	---	---	---	---	---
1911	80.9	---	---	---	---	---	---	---	---	---	---	---
1912	77.5	---	---	---	---	---	---	---	---	---	---	---
1913	79.2	---	---	---	---	---	---	---	---	---	---	---
1914	71.8	108.4	76.8	31.5	53.3	63.0	70.8	78.8	97.8	151.6	370.5	1,073.5
1915	67.1	95.9	73.8	30.8	48.4	57.2	65.7	73.5	89.6	144.0	366.6	1,049.0
1916	74.0	112.5	74.6	31.0	53.0	63.8	75.8	87.0	104.5	157.4	382.1	1,142.5
1917	76.3	---	---	---	---	---	---	---	---	---	---	---
1918	71.4	94.3	73.4	35.2	53.7	59.0	66.6	77.8	98.7	158.9	371.7	1,071.8
1919	61.0	85.5	70.7	32.1	46.5	48.1	56.0	62.2	80.3	127.2	339.2	1,047.5
1920	58.6	87.3	67.4	30.0	45.1	45.1	50.8	58.1	74.5	133.7	369.8	1,139.8
1921	54.7	79.0	62.2	27.7	41.3	41.8	47.3	53.7	71.0	124.3	356.9	1,180.3
1922	55.3	75.9	60.0	26.5	40.0	41.5	48.9	57.7	73.9	128.4	370.5	1,247.6
1923	58.8	79.9	58.2	26.7	42.3	44.5	53.9	63.9	81.3	137.5	389.9	1,298.8
1924	57.1	83.4	58.8	26.1	41.4	40.9	52.3	62.0	77.6	134.0	374.4	1,311.3
1925	58.4	83.4	54.8	25.7	42.2	42.3	53.4	65.7	79.9	137.8	394.6	1,357.3
1926	56.0	91.5	52.2	23.7	38.0	39.1	51.0	63.2	79.2	132.1	400.1	1,333.4
1927	53.9	76.9	51.2	23.3	36.5	37.4	48.8	59.7	76.0	130.0	386.6	1,274.8
1928	53.6	81.7	48.4	23.5	34.8	38.0	46.9	59.9	73.3	130.9	406.3	1,277.2
1929	53.1	81.5	46.4	22.2	34.9	38.5	46.7	57.4	74.8	131.2	387.8	1,236.9
1930	52.1	84.1	43.7	20.3	32.6	36.2	44.2	56.9	75.0	132.1	402.0	1,301.4
1931	50.1	77.1	45.1	19.2	29.8	31.4	40.2	55.3	74.6	132.5	404.8	1,266.0
1932	46.2	74.5	42.1	19.4	27.5	27.5	34.3	46.7	66.6	124.2	386.9	1,271.3
1933	45.8	76.7	40.9	19.1	26.0	27.3	33.4	46.4	65.8	122.3	401.3	1,251.7
1934	49.6	86.3	39.9	19.6	26.1	28.3	35.3	51.8	73.6	138.1	444.3	1,335.9
1935	48.7	87.5	41.1	18.7	25.2	28.1	35.4	49.0	69.0	131.2	430.9	1,376.4
1936	55.0	104.4	42.0	18.9	25.5	30.0	38.8	55.7	81.2	156.6	513.6	1,573.6
1937	49.8	90.1	39.2	18.9	25.5	28.7	34.3	47.4	68.8	132.0	458.9	1,386.7
1938	46.3	93.9	37.4	16.2	23.7	25.1	31.0	43.1	60.4	123.2	436.5	1,355.5
1939	45.2	91.2	36.2	16.5	22.1	24.5	29.0	40.1	58.2	118.9	435.1	1,347.7

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

WHITE, MALE												
Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	126.1	---	---	---	---	---	---	---	---	---	---	---
1911	120.7	---	---	---	---	---	---	---	---	---	---	---
1912	116.2	---	---	---	---	---	---	---	---	---	---	---
1913	119.9	---	---	---	---	---	---	---	---	---	---	---
1914	107.0	115.5	87.1	46.3	93.2	109.8	121.2	129.7	150.4	193.0	329.8	805.1
1915	99.5	103.5	84.2	45.5	84.6	100.1	112.9	120.4	135.9	178.6	329.4	828.8
1916	110.7	119.3	81.6	45.0	94.2	111.2	130.6	143.6	160.1	200.0	358.6	954.3
1917	112.6	---	---	---	---	---	---	---	---	---	---	---
1918	106.8	103.5	80.9	50.4	98.5	107.8	112.7	126.5	151.5	205.1	330.7	864.2
1919	88.5	98.8	75.8	45.6	81.6	84.3	95.1	100.8	120.7	159.1	294.3	807.7
1920	83.1	98.2	71.0	42.6	78.9	78.3	84.1	93.2	109.0	160.5	311.8	880.0
1921	77.6	86.8	68.9	39.8	71.9	72.0	79.3	86.1	104.2	149.5	303.3	867.2
1922	78.4	84.6	64.7	37.7	69.3	71.9	83.0	93.2	111.7	152.8	313.7	889.7
1923	84.0	88.5	63.6	37.7	73.5	77.6	90.5	104.3	124.1	168.7	320.5	921.1
1924	81.2	91.0	64.7	36.4	71.6	71.3	88.3	101.4	117.0	158.5	317.5	923.3
1925	82.7	92.1	58.2	36.4	72.2	72.9	89.5	106.9	120.2	169.0	326.4	1,035.7
1926	79.2	99.3	56.4	33.9	66.4	68.2	87.3	103.0	119.8	157.5	326.0	922.7
1927	76.8	83.7	56.2	33.7	63.7	65.5	82.8	99.2	115.1	156.6	326.3	924.3
1928	75.8	90.0	53.7	34.1	60.5	66.8	80.9	99.0	110.0	152.7	331.7	891.4
1929	74.9	91.6	49.9	32.1	61.1	67.5	80.0	95.3	113.3	151.3	313.3	827.8
1930	72.8	92.1	48.0	29.0	56.7	63.7	76.3	94.2	112.6	153.7	318.1	926.9
1931	69.3	85.3	49.7	27.5	51.6	55.4	69.0	91.3	110.5	152.7	331.8	925.8
1932	62.0	82.1	46.3	27.8	47.0	47.8	57.9	75.9	97.4	136.1	312.7	894.4
1933	61.0	84.0	43.9	27.7	44.2	46.8	55.9	75.2	95.6	137.1	315.8	915.6
1934	66.6	96.0	44.4	28.0	44.7	48.4	59.7	85.0	109.0	154.8	372.5	1,028.2
1935	64.5	97.1	44.8	26.5	42.5	48.1	60.2	79.9	100.6	146.4	357.4	999.7
1936	72.3	114.5	46.3	26.5	43.6	51.3	65.6	91.2	120.0	175.8	425.1	1,164.1
1937	66.3	103.8	44.0	26.9	43.6	50.3	59.0	78.2	103.3	145.9	383.8	1,080.3
1938	59.9	105.9	42.7	23.1	41.3	43.5	52.5	70.5	87.9	132.8	339.2	1,009.0
1939	58.4	103.0	42.4	24.1	38.6	42.9	49.8	65.7	84.3	130.4	344.8	954.0

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

WHITE, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	35.8	---	---	---	---	---	---	---	---	---	---	---
1911	39.0	---	---	---	---	---	---	---	---	---	---	---
1912	36.6	---	---	---	---	---	---	---	---	---	---	---
1913	36.4	---	---	---	---	---	---	---	---	---	---	---
1914	34.8	101.0	66.3	16.4	13.0	12.4	15.3	21.6	40.6	109.6	407.9	1,283.3
1915	33.1	88.2	63.1	15.9	12.1	11.0	13.7	20.8	39.3	108.8	400.6	1,219.6
1916	35.6	105.4	67.4	16.7	12.0	13.3	15.8	23.3	43.9	114.1	403.3	1,286.1
1917	38.5	---	---	---	---	---	---	---	---	---	---	---
1918	35.9	84.9	65.7	19.7	14.5	13.0	16.4	23.0	41.0	111.6	408.7	1,228.7
1919	32.6	71.9	65.4	18.3	13.1	11.6	13.5	18.8	36.2	94.4	379.8	1,227.2
1920	33.2	76.1	63.8	17.3	11.9	11.0	14.0	18.9	37.0	106.1	422.7	1,335.6
1921	31.1	70.9	55.4	15.4	11.2	10.9	12.3	17.8	35.1	88.5	405.8	1,418.8
1922	31.5	66.9	55.1	15.1	11.3	11.0	12.0	18.7	33.5	103.5	422.2	1,522.0
1923	32.9	71.1	52.6	15.6	11.6	11.3	14.3	20.0	35.5	105.5	453.4	1,591.9
1924	32.4	75.5	52.6	15.6	11.5	10.2	13.4	19.2	35.2	108.8	426.5	1,612.8
1925	33.3	74.3	51.2	14.7	12.4	11.4	14.5	21.1	36.5	105.7	457.3	1,607.6
1926	32.0	83.3	47.8	13.2	9.7	9.7	12.0	20.2	35.3	106.0	468.3	1,653.2
1927	30.4	69.8	46.0	12.8	9.4	9.1	12.5	17.0	33.7	102.6	442.3	1,546.9
1928	30.8	73.1	43.0	12.7	9.3	9.1	10.5	17.7	33.7	108.4	475.7	1,574.4
1929	30.6	71.0	42.7	12.1	9.0	9.5	11.3	16.4	33.4	110.5	457.5	1,547.8
1930	30.8	75.8	39.3	11.3	8.8	8.8	10.3	16.3	34.9	109.8	481.0	1,582.5
1931	30.5	68.6	40.5	10.7	8.2	7.6	9.9	16.2	36.5	111.7	473.4	1,521.3
1932	30.0	66.6	37.7	10.9	8.2	7.5	9.5	15.2	33.9	112.0	456.4	1,554.1
1933	30.3	69.2	37.7	10.3	8.1	8.0	9.9	15.4	34.1	107.2	481.6	1,505.0
1934	32.2	76.2	35.3	10.8	7.8	8.4	9.9	16.2	36.3	121.2	511.5	1,567.9
1935	32.5	77.5	37.4	10.7	8.1	8.2	9.7	15.9	35.7	115.9	499.4	1,660.3
1936	37.3	93.8	37.6	11.1	7.6	8.9	10.9	17.7	40.3	137.2	595.8	1,882.3
1937	32.9	76.0	34.3	10.5	7.6	7.4	8.9	14.5	32.4	118.1	528.3	1,617.8
1938	32.5	81.5	32.0	9.1	6.3	7.0	9.0	14.0	31.7	113.7	525.9	1,616.9
1939	31.8	79.1	29.8	8.5	5.7	6.4	7.8	12.9	30.9	107.5	517.7	1,644.8

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ALL OTHER, BOTH SEXES

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	91.9	---	---	---	---	---	---	---	---	---	---	---
1911	99.2	---	---	---	---	---	---	---	---	---	---	---
1912	94.0	---	---	---	---	---	---	---	---	---	---	---
1913	95.6	---	---	---	---	---	---	---	---	---	---	---
1914	88.7	348.9	119.3	47.7	68.8	79.7	86.3	96.1	97.2	138.5	304.6	422.2
1915	82.0	241.6	118.3	50.3	65.5	63.1	78.7	83.5	112.5	150.6	249.1	587.5
1916	81.8	242.1	127.6	42.3	68.3	70.4	86.2	80.5	103.6	128.4	258.9	438.6
1917	91.9	---	---	---	---	---	---	---	---	---	---	---
1918	82.2	212.2	113.6	43.6	69.4	66.7	87.5	86.9	105.9	170.7	269.3	605.1
1919	71.0	197.6	104.4	44.2	63.4	59.5	62.8	73.2	77.9	139.2	252.0	350.9
1920	71.1	224.1	109.9	43.6	68.3	56.5	55.7	69.0	81.7	142.3	233.3	570.6
1921	63.3	196.2	84.7	37.0	63.1	50.7	57.3	62.3	71.3	122.9	198.6	444.5
1922	63.0	211.3	94.5	34.6	57.8	49.4	52.9	70.2	83.3	113.2	187.1	486.3
1923	70.2	195.3	99.1	38.2	63.8	61.9	62.1	75.6	80.8	139.3	245.8	608.3
1924	73.5	233.2	105.9	38.0	67.0	59.2	68.6	79.1	93.0	133.9	257.5	692.8
1925	75.3	204.8	93.0	38.4	70.4	66.5	76.3	85.6	93.1	119.0	263.2	629.4
1926	74.5	190.5	93.6	38.6	64.7	68.4	77.3	86.5	96.1	128.9	239.7	639.6
1927	72.1	200.2	80.1	36.1	65.2	66.2	78.4	80.0	91.3	120.0	243.2	513.6
1928	67.6	201.1	75.3	33.0	60.2	62.5	71.4	79.9	75.8	124.6	240.0	489.9
1929	65.4	195.7	71.9	33.5	54.8	62.5	70.5	74.5	90.3	120.4	238.6	461.4
1930	62.1	198.9	69.9	30.3	48.9	57.3	67.7	71.9	88.5	127.2	243.0	466.5
1931	55.4	170.8	63.9	29.6	43.7	49.3	54.9	68.9	83.2	113.7	202.9	366.2
1932	56.0	171.4	64.1	29.2	48.0	51.6	54.9	59.3	81.9	111.5	221.7	419.4
1933	56.3	190.2	59.8	31.2	45.5	50.1	55.3	66.6	97.3	112.2	181.7	392.1
1934	61.1	210.3	68.0	30.3	48.4	53.4	60.9	77.2	92.6	118.1	228.1	448.9
1935	54.6	187.2	56.8	28.6	42.5	47.8	55.4	63.3	79.0	115.9	204.5	485.8
1936	61.8	222.0	60.1	29.2	43.5	55.4	61.3	77.5	97.6	146.4	247.6	589.2
1937	55.8	206.5	62.8	27.4	42.1	47.6	56.1	67.1	82.2	115.8	205.1	427.8
1938	50.1	184.9	53.7	26.3	40.1	42.5	48.0	57.7	73.4	96.5	169.3	414.0
1939	49.0	189.5	52.2	25.5	39.4	40.5	44.0	58.6	69.5	96.8	183.7	391.8

--- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.



HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ALL OTHER, MALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	...	...	...	...	...	...	...	...	...	...	...	...
1901	...	...	...	...	...	...	...	...	...	...	...	...
1902	...	...	...	...	...	...	...	...	...	...	...	...
1903	...	...	...	...	...	...	...	...	...	...	...	...
1904	...	...	...	...	...	...	...	...	...	...	...	...
1905	...	...	...	...	...	...	...	...	...	...	...	...
1906	...	...	...	...	...	...	...	...	...	...	...	...
1907	...	...	...	...	...	...	...	...	...	...	...	...
1908	...	...	...	...	...	...	...	...	...	...	...	...
1909	...	...	...	...	...	...	...	...	...	...	...	...
1910	136.2	...	...	...	...	...	...	...	...	...	...	...
1911	140.7	...	...	...	...	...	...	...	...	...	...	...
1912	137.8	...	...	...	...	...	...	...	...	...	...	...
1913	138.6	...	...	...	...	...	...	...	...	...	...	...
1914	125.6	375.3	119.7	55.2	119.9	135.8	135.3	130.5	130.3	163.9	314.6	*
1915	116.8	269.5	128.9	64.2	109.6	105.7	122.6	122.3	145.0	196.4	251.3	*
1916	118.6	276.2	131.0	48.4	120.7	123.6	140.5	122.2	142.4	171.2	287.1	*
1917	129.6	...	...	...	...	...	...	...	...	...	...	...
1918	118.1	226.7	115.7	54.0	127.2	114.8	138.5	122.8	138.2	200.2	320.6	577.6
1919	101.0	208.2	106.8	57.5	114.8	98.1	102.0	102.5	98.7	176.5	245.4	311.9
1920	99.0	240.1	107.4	49.1	122.5	98.4	88.6	96.4	110.3	168.9	228.1	612.7
1921	91.0	205.8	83.6	48.4	116.0	90.4	89.8	90.7	85.4	131.1	200.5	453.7
1922	88.6	228.5	89.0	42.4	100.3	87.6	83.5	102.8	108.8	132.1	204.7	417.1
1923	101.6	213.8	96.4	50.1	112.8	107.0	105.4	112.7	102.6	172.5	256.1	555.2
1924	102.3	257.5	98.5	45.7	113.4	101.9	111.7	110.4	123.5	142.9	261.2	711.7
1925	109.5	184.3	93.1	50.8	120.2	116.6	127.6	128.0	119.9	149.6	297.1	696.0
1926	109.3	204.3	94.3	49.0	113.1	119.4	126.0	130.2	132.9	149.1	285.5	732.6
1927	109.4	223.3	84.3	47.5	117.7	120.3	133.6	122.9	118.6	142.1	290.4	514.5
1928	98.9	214.5	72.9	41.5	104.7	111.5	120.4	117.5	95.1	148.4	257.8	508.9
1929	96.1	201.9	70.6	43.9	97.0	109.1	119.9	110.6	107.2	150.0	301.7	424.6
1930	91.4	213.6	73.3	38.3	86.8	101.2	114.9	106.1	104.8	166.2	295.1	427.4
1931	79.9	177.1	60.5	39.5	77.0	84.6	91.4	102.2	104.7	131.5	182.1	383.2
1932	82.2	178.4	66.3	39.1	85.4	89.7	92.7	87.1	106.9	127.6	238.7	434.1
1933	82.0	201.5	57.9	42.6	79.4	86.2	89.4	99.7	108.9	131.8	233.6	392.8
1934	88.5	211.8	72.5	40.9	82.8	91.2	99.7	116.5	117.8	142.6	245.8	435.4
1935	78.1	201.6	52.9	37.8	74.2	81.9	92.4	89.8	101.4	135.8	217.6	438.1
1936	88.4	244.5	53.8	38.1	75.8	94.9	102.5	115.2	121.6	167.4	269.1	589.0
1937	81.0	213.5	67.1	34.1	75.5	82.6	94.9	100.4	107.4	133.8	232.2	379.2
1938	72.0	195.8	50.5	35.6	69.4	75.8	78.6	85.7	92.7	109.2	167.0	454.5
1939	71.7	216.2	55.0	36.5	67.8	69.8	75.1	92.1	88.6	112.4	194.6	342.9

... Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

HIST290. DEATH RATES FROM SELECTED CAUSES, BY 10-YEAR GROUPS, RACE, AND SEX: DEATH-REGISTRATION STATES, 1900-1932, AND UNITED STATES, 1933-1939

[Rates per 100,000 population for specified group.]

ACCIDENTS, EXCEPT MOTOR-VEHICLE

ALL OTHER, FEMALE

Years	All Ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years & over
1900	---	---	---	---	---	---	---	---	---	---	---	---
1901	---	---	---	---	---	---	---	---	---	---	---	---
1902	---	---	---	---	---	---	---	---	---	---	---	---
1903	---	---	---	---	---	---	---	---	---	---	---	---
1904	---	---	---	---	---	---	---	---	---	---	---	---
1905	---	---	---	---	---	---	---	---	---	---	---	---
1906	---	---	---	---	---	---	---	---	---	---	---	---
1907	---	---	---	---	---	---	---	---	---	---	---	---
1908	---	---	---	---	---	---	---	---	---	---	---	---
1909	---	---	---	---	---	---	---	---	---	---	---	---
1910	41.0	---	---	---	---	---	---	---	---	---	---	---
1911	52.9	---	---	---	---	---	---	---	---	---	---	---
1912	45.0	---	---	---	---	---	---	---	---	---	---	---
1913	49.2	---	---	---	---	---	---	---	---	---	---	---
1914	48.9	322.7	118.8	40.4	19.1	19.0	26.5	50.1	55.5	109.8	295.1	*
1915	44.4	213.9	108.0	36.7	22.8	17.4	25.2	30.9	71.0	98.5	247.0	741.0
1916	43.9	208.4	124.3	36.2	19.9	17.8	25.3	25.7	53.9	78.4	231.6	414.9
1917	53.4	---	---	---	---	---	---	---	---	---	---	---
1918	45.8	197.9	111.6	33.4	18.0	21.4	31.7	38.7	64.2	136.4	219.1	624.8
1919	40.7	187.2	102.0	30.9	18.0	23.7	20.7	33.1	50.7	95.1	258.5	379.3
1920	42.7	208.6	112.4	38.2	19.2	17.8	19.8	32.0	44.4	110.6	238.6	539.0
1921	35.1	186.8	85.8	25.8	14.6	14.2	22.1	24.4	52.9	113.0	196.6	438.0
1922	37.2	194.5	100.1	26.9	18.8	14.8	20.7	27.9	49.8	90.0	169.2	533.8
1923	38.5	177.1	101.7	26.6	18.5	20.8	16.7	28.9	52.4	98.6	235.4	643.7
1924	44.7	209.2	113.4	30.4	23.9	20.0	23.7	40.5	53.1	122.8	253.8	680.3
1925	40.8	225.1	93.0	26.2	24.0	19.8	23.3	33.8	58.2	82.1	229.4	585.2
1926	39.5	177.0	92.8	28.6	19.4	20.5	27.2	34.4	48.3	104.6	194.4	576.8
1927	34.6	177.8	75.9	25.0	16.2	15.5	22.1	29.2	55.5	93.7	196.5	513.0
1928	36.5	188.1	77.8	24.6	19.2	17.1	23.0	35.7	50.3	96.7	212.6	465.7
1929	34.9	189.7	73.2	23.3	16.3	19.1	22.0	32.5	67.9	86.5	177.5	488.7
1930	33.1	184.3	66.6	22.3	14.5	16.3	21.5	32.2	67.3	83.1	193.4	495.9
1931	31.2	164.7	67.4	19.8	13.5	16.5	19.4	30.7	55.6	93.6	222.6	353.4
1932	30.1	164.4	62.0	19.4	14.1	16.2	18.2	27.8	50.1	93.4	205.6	408.4
1933	31.1	179.1	61.6	19.9	14.9	16.5	22.2	29.3	60.3	90.5	132.4	391.6
1934	34.2	208.7	63.5	19.9	17.2	18.4	23.3	33.5	61.4	91.0	211.3	459.0
1935	31.6	173.0	60.6	19.6	13.8	16.4	19.8	34.1	51.4	93.9	192.0	521.3
1936	35.8	200.0	66.5	20.3	14.1	19.0	21.7	36.2	68.5	123.4	227.3	589.4
1937	31.2	199.6	58.5	20.8	11.8	15.6	18.9	30.9	52.0	96.0	179.4	463.7
1938	28.8	174.2	56.8	17.1	13.6	12.2	18.7	27.5	50.8	82.6	171.5	384.1
1939	27.0	163.5	49.3	14.6	13.6	13.9	14.4	22.7	47.4	79.7	173.3	427.7

-- Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

\* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Notes: 1) Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

2) For 1932-34, deaths of Mexicans are included with total deaths for the white population in other publications reflecting later practices; but by age, are included with deaths for the ALL OTHER population. These tables reflected the original practice of including Mexican totals in the ALL OTHER total. Figures for deaths of Mexicans are not available separately by cause and age; total deaths of Mexicans from this cause are:

	1932	1933	1934
Both sexes ....	288	462	482
Male.....	201	327	345
Female....	87	135	137

3) For 1909 total excludes deaths from food poisoning, they are included in age distribution

4) For 1922-29 total excludes deaths from legal executions, they are included in age distribution

5) Not tabulated separately by age

6) Year in which each state was admitted to the death-registration area:

Year	Year	Year
1880 Massachusetts	1908 Washington	1919 Florida
New Jersey	Wisconsin	Mississippi
District of Columbia /1	1909 Ohio	1920 Nebraska
1890 Connecticut	1910 Minnesota	1922 Georgia /4
Delaware /2	Montana	Idaho
New Hampshire	Utah	Wyoming
New York	1911 Kentucky	1923 Iowa
Rhode Island	Missouri	1924 North Dakota
Vermont	1913 Virginia	1925 Alabama
1900 Maine	1914 Kansas	West Virginia
Michigan	1916 South Carolina	1926 Arizona
Indiana	North Carolina	1927 Arkansas
1906 California	1917 Tennessee	1928 Oklahoma
Colorado	1918 Illinois	1929 Nevada
Maryland	Louisiana	New Mexico
Pennsylvania	Oregon	1933 Texas
South Dakota /3		

/1 Included in States.

/2 Dropped from the registration area in 1900; readmitted in 1919.

/3 Dropped from the registration area in 1910; readmitted in 1930.

/4 Dropped from the registration area in 1925; readmitted in 1928.

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.