Vital and Health Statistics

Series 3, Number 50 June 2025



Anthropometric Reference Data for Children and Adults: United States, August 2021–August 2023

Analytical and Epidemiological Studies



Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested citation

Fryar CD, Gu Q, Afful J, Carroll MD, Ogden CL. Anthropometric reference data for children and adults: United States, August 2021–August 2023. Vital Health Stat 3. 2025 Jun;(50): 1–28. DOI: https://dx.doi.org/10.15620/cdc/174595.

For sale by the U.S. Government Publishing Office Superintendent of Documents Mail Stop: SSOP Washington, DC 20401–0001 Printed on acid-free paper.

NATIONAL CENTER FOR HEALTH STATISTICS

Vital and Health Statistics

Series 3, Number 50 June 2025

Anthropometric Reference Data for Children and Adults: United States, August 2021–August 2023

Analytical and Epidemiological Studies

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES Centers for Disease Control and Prevention National Center for Health Statistics

Hyattsville, Maryland June 2025

National Center for Health Statistics

Brian C. Moyer, Ph.D., *Director*Amy M. Branum, Ph.D., *Associate Director for Science*

Division of Health and Nutrition Examination Surveys

Alan E. Simon, M.D., *Director*Lara J. Akinbami, M.D., *Associate Director for Science*

Contents

| Abs | tract |
|--------|--|
| Intr | oduction |
| 9 E | thods |
| Res | ults |
| Disc | cussion |
| Refe | erences |
| App | pendix. Supporting Tables |
| Тех | rt Table |
| | ailed tables with body measurements or indices from the National Health and Nutrition Examination Survey, rust 2021–August 2023 |
| De | tailed Tables |
| 1. | Weight in kilograms for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 2. | Weight in pounds for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 6 |
| 3. | Weight in kilograms for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 4. | Weight in pounds for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 5. | Height in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 6. | Height in inches for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 8 |
| 7. | Height in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021—August 2023 |
| 8. | Height in inches for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 9. | Body mass index values for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 10. | Body mass index values for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021—August 2023 |

Contents—Con.

| 11. | Waist circumference in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
|-----|---|
| 12. | Waist circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021—August 2023 |
| 13. | Hip circumference in centimeters for children and adolescents ages 12–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 14. | Hip circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 15. | Mid-upper arm circumference in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 16. | Mid-upper arm circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 17. | Upper arm length in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 18. | Upper arm length in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021—August 2023 |
| 19. | Upper leg length in centimeters for children and adolescents ages 8–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| 20. | Upper leg length in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023 |
| Аp | pendix Tables |
| l. | Body measurements or indices from the National Health and Nutrition Examination Survey, by age and survey period, 1999–2000 through August 2021–August 2023 |
| II. | Historical summary of published anthropometry reference data |

Anthropometric Reference Data for Children and Adults: United States, August 2021–August 2023

by Cheryl D. Fryar, M.S.P.H., Qiuping Gu, M.D., Joseph Afful, M.S., Margaret D. Carroll, M.S.P.H., and Cynthia L. Ogden, Ph.D.

Abstract

Objective

This report presents anthropometric reference data for the U.S. population age 2 years and older during August 2021–August 2023.

Methods

Body measurements were obtained from 8,545 National Health and Nutrition Examination Survey (NHANES) participants age 2 years and older during August 2021—August 2023. Measurements included weight, height, circumferences, and limb lengths.

Results

Weighted population means, standard errors of the means, and selected percentiles of body measurement values are reported for participants age 2 years and older. Body measurement results are shown by sex and age.

Conclusions

These nationally representative NHANES data show the distribution of measured weight, height, body mass index, circumferences, and lengths in the U.S. population during August 2021–August 2023.

Keywords: anthropometry • body measures • nutrition surveys • National Health and Nutrition Examination Survey (NHANES)

Introduction

Anthropometry, the science of measuring human body dimensions, serves as a key indicator of nutritional status, overall health, dietary sufficiency, and growth patterns. The National Center for Health Statistics (NCHS) provides nationally representative, in-person body measurement data collected through the National Health and Nutrition Examination Survey (NHANES), the primary source of body measurement information for the U.S. population. These measurements offer insights into the average weight, height, length, and circumferences of U.S. children, adolescents, and adults. Previous reports by NCHS have detailed anthropometric reference tables that describe the distribution of various body measurements within the U.S. population during 1988-2018 (1-6). This report presents updated anthropometric reference data collected through NHANES from August 2021 to August 2023 for age 2 years and older by sex and age.

Methods

NHANES, conducted by NCHS, collects data in household interviews followed by health examinations in mobile examination centers (MECs). The MECs are staffed by full-time personnel, including health technicians who perform body measurements on survey participants using a standard protocol.

Surveys were conducted on a periodic basis from 1960 to 1994. NHANES became a continuous survey in 1999 and continued through March 2020, when data collection was paused due to the COVID-19 pandemic. NHANES data collection resumed in August 2021 using a new sample design to reduce contact between interviewers and survey participants during the continuing COVID-19 pandemic. The 2-year public-use data files for August 2021—August 2023 were used for this report (7–9). During this survey period, the overall examination response rate was 25.6%.

Sample Description

NHANES is a complex, multistage probability sample of the civilian noninstitutionalized U.S. population. The NHANES August 2021—August 2023 sample included participants of all ages (9). Due to small sample sizes for some body measurements, only those age 2 years and older are included in this report. The analytic sample for this report was based on the August 2021—August 2023 examined sample of 8,545 people age 2 years and older. Pregnant females were excluded from the tabulations of weight, body mass index (BMI), and circumference measurements.

Age for adults was categorized into 10-year age groups and age 80 and older. For participants ages 2–19 years, results were reported by 2-year age groups. This is a change from previous anthropometric reports that were based on 4 years of NHANES data and a corresponding larger sample size, with results for children and adolescents shown by single year of age.

Body Measures

The NHANES Anthropometry Procedures Manual describes the protocol, equipment, quality control, and measurement procedures in detail (10). All NHANES participants were eligible for the anthropometry examination component, although not all age groups were eligible for all measurements. A list of body measures by age eligibility, collected by NHANES from 1999–2000 through August 2021–August 2023, is included in Appendix Table I.

Weight was measured to the nearest 0.1 kilogram using a digital floor scale.

Upper arm length was measured with a tape measure from the posterior border of the acromion process to the tip of the olecranon process; during the measurement, the upper arm length midpoint was marked. The mid-arm circumference was measured with a tape measure at the marked point. Upper arm length and mid-arm circumference were measured to the nearest 0.1 centimeter.

Standing height or stature and waist circumference were measured to the nearest 0.1 centimeter on participants. Standing height was measured with a wall-mounted stadiometer. Waist circumference was measured with a tape measure at the uppermost lateral border of the hip crest (ilium).

Upper leg length was measured on participants age 8 and older. Upper leg measurements to the nearest 0.1 centimeter were taken on seated participants; the distance from the inguinal crease to the distal end of the femur was measured. Hip circumference was measured to the nearest 0.1 centimeter on participants age 12 years and older.

Weight and standing height values were recorded automatically. The other body measurement data were

recorded using computer-assisted data entry. All results were based on a single body measurement examination.

One index is reported, BMI, calculated as weight in kilograms divided by height in meters squared (kg/m²). BMI is often used to define obesity.

Statistical Analysis

Unweighted sample sizes and weighted means, standard errors, and selected percentiles were estimated. Data were weighted using examination sample weights to produce national estimates. The NHANES examination sample weights incorporate the differential probabilities of selection and include adjustments for noncoverage and nonresponse. Standard errors were estimated using SUDAAN by Taylor series linearization because of the complex sample design (11).

All estimates of means and percentiles were considered statistically reliable based on not having less than 8 degrees of freedom or a relative standard error greater than 30%. Although NCHS now uses new criteria for presentation of proportions (12), similar criteria for means and percentiles have not been established. Estimates with standard errors for the percentile that could not be calculated using SUDAAN were omitted from the tables and are identified with a dagger (†).

Results

A list of detailed tables with distributions for the various body measurements and BMI is found in the text Table. Tables 1–20 contain the sample size, mean, standard error, and selected percentiles for the different measures. Results are reported by sex and age group. For example, during August 2021–August 2023, the average man age 20 and older weighed 199 pounds (Table 4) and was 68.9 inches (just under 5 feet 9 inches) tall (Table 8). The average woman age 20 and older weighed about 172 pounds (171.8) (Table 4) and was 63.5 inches (5 feet 3.5 inches) tall (Table 8). Results from previous surveys conducted from 1960 to 2018 for most measurements have been reported (1–6,13–24). A historical summary of published body measurement reference data is available in Appendix Table II.

Discussion

This report presents the latest anthropometric reference data for U.S. children and adults. This is the first time since NHANES III (1988–1994) that hip circumference measurements have been included in the report.

Reference data derived from anthropometric measurements are widely used. The anthropometric data collected through NHANES are used for various applications, including public health research, health and safety policy formulation,

Table. Detailed tables with body measurements or indices from the National Health and Nutrition Examination Survey, August 2021–August 2023

| Table number | Measurement unit or index | Age group (years) |
|--------------|-------------------------------------|-------------------|
| 1 Weight | (kilograms) | 2–19 |
| 2 Weight | | 2–19 |
| 3 Weight | (kilograms) | 20 and older |
| 4 Weight | (pounds) | 20 and older |
| 5 Standin | g height (centimeters) | 2–19 |
| 6 Standin | g height (inches) | 2–19 |
| 7 Standin | g height (centimeters) | 20 and older |
| 8 Standin | g height (inches) | 20 and older |
| 9 Body m | ass index | 2–19 |
| 10 Body m | ass index | 20 and older |
| 11 Waist ci | rcumference (centimeters) | 2–19 |
| 12 Waist ci | rcumference (centimeters) | 20 and older |
| 13 Hip circ | umference (centimeters) | 12–19 |
| 14 Hip circ | umference (centimeters) | 20 and older |
| 15 Mid-upp | per arm circumference (centimeters) | 2–19 |
| 16 Mid-upp | per arm circumference (centimeters) | 20 and older |
| 17 Upper a | rm length (centimeters) | 2–19 |
| 18 Upper a | rm length (centimeters) | 20 and older |
| 19 Upper le | eg length (centimeters) | 8–19 |
| 20 Upper le | eg length (centimeters) | 20 and older |

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

clinical practices, and product development (25–32). These measurements serve as critical references for understanding population health trends and informing interventions aimed at improving health outcomes in adults and youth.

References

- McDowell MA, Fryar CD, Ogden CL. Anthropometric reference data for children and adults: United States, 1988–1994. Vital Health Stat 11. 2009;(249):1–68.
- McDowell MA, Fryar CD, Hirsch R, Ogden CL. Anthropometric reference data for children and adults: U.S. population, 1999–2002. Adv Data. 2005;(361): 1–5.
- 3. McDowell MA, Fryar CD, Ogden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2003–2006. National health statistics reports. 2008;(10):1–48.
- 4. Fryar CD, Gu Q, Ogden CL. Anthropometric reference data for children and adults: United States, 2007–2010. Vital Health Stat 11. 2012;(252):1–48.
- 5. Fryar CD, Gu Q, Ogden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2011–2014. Vital Health Stat 3. 2016;(39):1–46.
- Fryar CD, Carroll MD, Gu Q, Afful J, Ogden CL. Anthropometric reference data for children and adults: United States, 2015–2018. Vital Health Stat 3. 2021;(46):1–35.
- National Center for Health Statistics. National Health and Nutrition Examination Survey, August 2021–August 2023 data documentation, codebook, and frequencies: Body measures. 2024. Available from: https:// wwwn.cdc.gov/Nchs/Data/Nhanes/Public/2021/DataFiles/BMX L.htm.
- 8. National Center for Health Statistics. National Health and Nutrition Examination Survey: Brief overview and analytic guidance: NHANES

- August 2021—August 2023 public use file release. 2024. Available from: https://wwwn.cdc.gov/nchs/nhanes/continuousnhanes/overviewbrief.aspx?Cycle=2021-2023.
- Terry AL, Chiappa MM, McAllister J, Woodwell DA, Graber JE. Plan and operations of the National Health and Nutrition Examination Survey, August 2021–August 2023. National Center for Health Statistics. Vital Health Stat 1. 2024 May; (66):1–21. DOI: https://dx.doi.org/10.15620/ cdc/151927.
- 10. National Center for Health Statistics. National Health and Nutrition Examination Survey (NHANES) anthropometry procedures manual. 2021 May. Available from: https://wwwn.cdc.gov/nchs/data/nhanes/public/2021/manuals/2021-Anthropometry-Procedures-Manual-508.pdf.
- Wolter KM. Introduction to variance estimation. 2nd ed. New York: Springer; 2007.
- Parker JD, Talih M, Malec DJ, Beresovsky V, Carroll M, Gonzalez JF Jr, et al. National Center for Health Statistics data presentation standards for proportions. Vital Health Stat 2. 2017 Aug;(175):1–22.
- 13. Stoudt HW, Damon A,
 McFarland R, Roberts J. Weight,
 height, and selected body
 dimensions of adults, United
 States—1960–1962. Vital Health
 Stat 11. 1965 Jun;(8):1–44.
- 14. Stoudt HW. Skinfolds, body girths, biacromial diameter, and selected anthropometric indices of adults, United States, 1960–1962. Vital Health Stat 11. 1970 Feb;(35):1–63.
- 15. Roberts J. Weight by height and age of adults, United States—1960–1962. Vital Health Stat 11. 1966 May;(14):1–38.

- Malina RM, Hamill PVV, Lemeshow S. Selected body measurements of children 6–11 years, United States. Rockville, MD: Vital Health Stat 11. 1973 Jan;(123): 1–48.
- 17. Hamill PV, Johnston FE, Lemeshow S. Height and weight of youths 12–17 years, United States. Vital Health Stat 11. 1973(124):1–81.
- 18. Abraham S, Johnson CL, Najjar MF. Weight and height of adults 18–74 years of age: United States, 1971–74. Vital Health Stat 11. 1979(211):1–49.
- Johnson CL, Fulwood R, Abraham S, Bryner JD. Basic data on anthropometric measurements and angular measurements of the hip and knee joints for selected age groups 1–74 years of age. Vital Health Stat 11. 1981(219):1–68.
- 20. Najjar MF, Rowland M. Anthropometric reference data and prevalence of overweight, United States, 1976–80. Vital Health Stat 11. 1987(238):1–73.
- 21. Najjar MF, Kuczmarski RJ. Anthropometric data and prevalence of overweight for Hispanics: 1982–84. Vital Health Stat 11. 1989(239):1–106.
- 22. Ogden CL, Fryar CD, Carroll MD, Flegal KM. Mean body weight, height, and body mass index, United States 1960–2002. Adv Data. 2004(347):1–17.
- 23. Fryar CD, Kruszon-Moran D, Gu Q, Ogden CL. Mean body weight, height, waist circumference, and body mass index among adults: United States, 1999–2000 through 2015–2016. Natl Health Stat Report. 2018 Dec;(122):1–16.
- 24. Fryar CD, Kruszon-Moran D, Gu Q, Carroll M, Ogden CL. Mean body weight, height, waist circumference, and body mass index among children and adolescents: United States, 1999–2018. National health statistics reports. 2012(160):1–23.
- Ogden CL, Fryar CD, Martin CB, Freedman DS, Carroll MD, Gu Q, Hales CM. Trends in obesity prevalence by race and Hispanic origin—1999–2000 to 2017–2018.
 JAMA. 2020 Sep 22;324(12):1208–10. DOI: 10.1001/jama.2020.14590. PMID: 32857101.
- Emmerich SD, Fryar CD, Stierman B, Gu Q, Afful J, Ogden CL. Trends in obesity-related measures among US children, adolescents, and adults. JAMA. 2025 Mar 25;333(12):1082–4. DOI: 10.1001/ jama.2024.27676. PMID: 39946125.
- 27. Executive Office of the President. White House Task Force on Childhood Obesity: Report to the President. Solving the problem of childhood obesity. 2010.
- 28. Ostchega Y, Hughes JP, Prineas RJ, Zhang G, Nwankwo T, Chiappa MM. Mid-arm circumference and recommended blood pressure cuffs for children and adolescents ages between 3 and 19 years: Data from

- the National Health and Nutrition Examination Survey, 1999–2010. Blood Press Monit. 2014;19(1):26–31.
- Ostchega Y, Hughes JP, Zhang G, Nwankwo T, Chiappa MM. Mean mid-arm circumference and blood pressure cuff sizes for U.S. adults: National Health and Nutrition Examination Survey, 1999–2010. Blood Press Monit. 2013;18(3):138–43.
- Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, et al. 2000 CDC growth charts for the United States: Methods and development. Vital Health Stat 11. 2002 May;(246):1–190. PMID: 12043359. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_246.pdf.
- U.S. Department of Transportation. Advisory circular: Aircraft weight and balance control. Federal Aviation Administration. 2019 May 6. Available from: https:// www.faa.gov/documentLibrary/media/Advisory_ Circular/AC_120-27F.pdf.
- 32. Department of Homeland Security. Passenger weight and inspected vessel stability requirements; Final rule. 2010 Dec 14;75(239). Available from: https://www.dco.uscg.mil/Portals/9/DCO%20Documents/5p/CG-5PC/CG-CVC/CVC1/policy/pwivsr/Passenger_Weight_and_Inspected_Vessel_Stability_Final_Rule.pdf.

Table 1. Weight in kilograms for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023

| 01 | Number of | | Standard | | | | | Percentile | | | | |
|--|--------------------|------|------------------------|------|------|------|-----------|------------|------|------|-------|-------|
| Sex and age group (years) ¹ | examined people | Mean | error of – the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Kilograms | | | | | |
| 2–3 | 125 | 15.0 | 0.3 | 11.5 | 11.9 | 12.5 | 13.2 | 14.6 | 16.4 | 16.9 | 17.8 | 19.2 |
| 4–5 | 155 | 20.3 | 0.4 | 15.0 | 15.5 | 16.3 | 17.3 | 19.5 | 21.8 | 23.6 | 25.1 | 28.0 |
| 6–7 | 122 | 25.5 | 0.6 | 18.0 | 19.4 | 20.4 | 21.6 | 24.1 | 28.4 | 32.1 | 34.0 | 36.2 |
| 8–9 | 171 | 33.5 | 0.8 | 23.0 | 24.5 | 25.2 | 27.4 | 30.5 | 37.4 | 41.3 | 43.5 | 53.8 |
| 10–11 | 140 | 46.7 | 1.6 | 27.2 | 29.6 | 31.3 | 35.7 | 40.5 | 55.8 | 62.5 | 68.5 | † |
| 12–13 | 143 | 58.3 | 1.6 | 37.3 | 39.6 | 42.7 | 47.1 | 55.2 | 63.0 | 75.4 | 81.1 | † |
| 14–15 | 145 | 71.2 | 1.9 | 43.4 | 48.4 | 51.6 | 55.0 | 66.2 | 82.8 | 95.2 | 101.7 | 112.3 |
| 16–17 | 136 | 76.3 | 2.7 | 50.6 | 53.1 | 55.7 | 60.0 | 74.0 | 85.7 | 96.3 | 100.4 | † |
| 18–19 | 138 | 77.0 | 2.0 | 51.8 | 53.9 | 56.3 | 62.6 | 72.3 | 86.3 | 95.3 | 103.1 | 110.3 |
| Female | | | | | | | | | | | | |
| 2–3 | 100 | 14.5 | 0.3 | 10.8 | 11.3 | 11.9 | 12.5 | 14.5 | 15.9 | 16.4 | 17.2 | 19.2 |
| 4–5 | 151 | 19.5 | 0.3 | 14.9 | 15.5 | 15.9 | 16.5 | 18.3 | 21.2 | 23.2 | 25.9 | 28.7 |
| 6–7 | 149 | 26.0 | 0.6 | 18.0 | 18.7 | 19.7 | 21.2 | 24.5 | 28.2 | 32.1 | 35.7 | 40.2 |
| 8–9 | 135 | 34.3 | 1.3 | 23.0 | 24.6 | 25.5 | 28.0 | 31.4 | 37.8 | 44.0 | 45.3 | 57.3 |
| 10–11 | 151 | 46.0 | 1.2 | 30.2 | 32.3 | 33.4 | 35.3 | 43.2 | 52.5 | 59.7 | 64.4 | 73.6 |
| 12–13 | 161 | 55.4 | 1.4 | 36.6 | 41.3 | 41.9 | 46.5 | 52.3 | 60.9 | 65.3 | 73.1 | 81.4 |
| 14–15 | 126 | 61.9 | 1.8 | 39.7 | 45.1 | 47.3 | 50.6 | 57.1 | 68.7 | 80.4 | 86.4 | 97.4 |
| 16–17 | 147 | 64.4 | 2.0 | 45.2 | 48.1 | 48.8 | 51.6 | 59.7 | 71.6 | 78.6 | 86.6 | 103.2 |
| 18–19 | 128 | 69.1 | 2.1 | 43.2 | 45.4 | 48.9 | 52.7 | 63.7 | 82.6 | 93.0 | 99.3 | 110.7 |

 $[\]dagger$ Estimate not shown because the standard error of the 95th percentile could not be computed. $^1\!Age$ is at time of examination.

Table 2. Weight in pounds for children and adolescents ages 2-19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| | Number of | | Standard | | | | | Percentile | | | | |
|--------|--------------------|-------|----------------------|-------|-------|-------|--------|------------|-------|-------|-------|-------|
| 0 0 ,1 | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Pounds | | | | | |
| 2–3 | 125 | 33.1 | 0.8 | 25.4 | 26.3 | 27.6 | 29.2 | 32.2 | 36.2 | 37.3 | 39.1 | 42.4 |
| 4–5 | 155 | 44.8 | 0.9 | 33.2 | 34.3 | 35.9 | 38.1 | 43.1 | 48.0 | 52.0 | 55.4 | 61.7 |
| 6–7 | 122 | 56.2 | 1.3 | 39.7 | 42.8 | 45.0 | 47.7 | 53.1 | 62.5 | 70.7 | 74.8 | 79.8 |
| 8–9 | 171 | 73.9 | 1.7 | 50.6 | 53.9 | 55.5 | 60.4 | 67.2 | 82.4 | 91.0 | 95.9 | 118.5 |
| 10–11 | 140 | 103.0 | 3.6 | 60.0 | 65.2 | 69.0 | 78.8 | 89.2 | 123.1 | 137.8 | 151.0 | † |
| 12–13 | 143 | 128.5 | 3.5 | 82.1 | 87.3 | 94.1 | 103.9 | 121.7 | 138.9 | 166.3 | 178.8 | † |
| 14–15 | 145 | 156.9 | 4.2 | 95.7 | 106.7 | 113.7 | 121.3 | 145.9 | 182.5 | 210.0 | 224.2 | 247.6 |
| 16–17 | 136 | 168.3 | 5.9 | 111.6 | 117.1 | 122.8 | 132.3 | 163.1 | 189.0 | 212.2 | 221.4 | † |
| 18–19 | 138 | 169.8 | 4.4 | 114.1 | 118.8 | 124.2 | 138.0 | 159.4 | 190.2 | 210.1 | 227.4 | 243.2 |
| Female | | | | | | | | | | | | |
| 2–3 | 100 | 32.0 | 0.6 | 23.8 | 25.0 | 26.3 | 27.5 | 32.1 | 35.0 | 36.2 | 38.0 | 42.3 |
| 4–5 | 151 | 43.0 | 0.8 | 32.7 | 34.1 | 35.1 | 36.4 | 40.2 | 46.7 | 51.0 | 57.2 | 63.2 |
| 6–7 | 149 | 57.4 | 1.4 | 39.8 | 41.2 | 43.5 | 46.7 | 53.9 | 62.1 | 70.8 | 78.6 | 88.7 |
| 8–9 | 135 | 75.7 | 2.8 | 50.8 | 54.2 | 56.1 | 61.7 | 69.2 | 83.4 | 97.1 | 99.9 | 126.3 |
| 10–11 | 151 | 101.4 | 2.6 | 66.6 | 71.2 | 73.6 | 77.8 | 95.2 | 115.8 | 131.5 | 142.1 | 162.2 |
| 12–13 | 161 | 122.2 | 3.1 | 80.8 | 90.9 | 92.5 | 102.6 | 115.3 | 134.3 | 143.9 | 161.2 | 179.4 |
| 14–15 | 126 | 136.5 | 3.9 | 87.5 | 99.3 | 104.2 | 111.6 | 125.8 | 151.6 | 177.2 | 190.5 | 214.7 |
| 16–17 | 147 | 141.9 | 4.4 | 99.6 | 106.0 | 107.6 | 113.7 | 131.7 | 157.8 | 173.4 | 190.8 | 227.4 |
| 18–19 | 128 | 152.3 | 4.6 | 95.1 | 100.1 | 107.9 | 116.1 | 140.5 | 182.1 | 205.1 | 218.9 | 244.0 |

[†] Estimate not shown because the standard error of the 95th percentile could not be computed. ¹Age is at time of examination.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 3. Weight in kilograms for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Carrand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|----------------------|------|------|------|----------|------------|-------|-------|-------|-------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | K | ilograms | | | | | |
| 20 and older | 2,688 | 90.3 | 0.9 | 60.9 | 66.6 | 70.0 | 75.4 | 86.6 | 100.6 | 111.0 | 118.8 | 131.8 |
| 20-29 | | 86.8 | 1.8 | 57.0 | 60.5 | 64.2 | 72.2 | 83.7 | 98.2 | 108.1 | 114.7 | 128.3 |
| 30–39 | | 91.4 | 1.1 | 63.6 | 69.8 | 71.1 | 77.1 | 87.1 | 101.5 | 110.0 | 121.0 | 132.5 |
| 40–49 | 347 | 95.3 | 2.0 | 66.0 | 69.7 | 73.2 | 78.5 | 88.4 | 107.7 | 119.7 | 125.1 | 141.9 |
| 50–59 | 415 | 92.1 | 1.5 | 61.0 | 68.1 | 71.2 | 76.8 | 87.8 | 102.0 | 113.6 | 119.8 | 138.7 |
| 60–69 | 651 | 89.2 | 1.1 | 62.6 | 67.1 | 69.9 | 76.0 | 86.9 | 99.0 | 108.3 | 114.0 | 123.9 |
| 70–79 | 446 | 87.7 | 1.5 | 60.8 | 66.2 | 68.8 | 74.5 | 86.6 | 96.5 | 106.8 | 111.7 | 121.3 |
| 80 and older | 157 | 81.7 | 0.9 | 58.0 | 63.3 | 66.7 | 70.8 | 79.9 | 90.0 | 96.5 | 99.6 | 108.5 |
| Female | | | | | | | | | | | | |
| 20 and older | 3,254 | 77.9 | 0.8 | 50.7 | 54.7 | 57.8 | 62.6 | 73.5 | 89.2 | 99.9 | 107.2 | 119.2 |
| 20–29 | , | 74.8 | 2.1 | 47.6 | 52.9 | 54.9 | 59.4 | 68.9 | 82.9 | 98.2 | 107.3 | 121.3 |
| 30–39 | | 80.9 | 0.9 | 51.4 | 55.4 | 58.9 | 63.1 | 75.0 | 93.3 | 107.9 | 114.6 | 125.9 |
| 40–49 | 442 | 80.4 | 1.4 | 52.8 | 56.0 | 58.7 | 63.9 | 74.8 | 90.6 | 101.1 | 111.5 | 123.7 |
| 50–59 | | 81.6 | 0.9 | 53.1 | 57.9 | 60.2 | 65.2 | 78.5 | 95.5 | 105.0 | 111.5 | 121.4 |
| 60–69 | | 77.5 | 1.0 | 51.6 | 55.5 | 58.9 | 63.3 | 74.3 | 89.9 | 98.4 | 104.5 | 109.3 |
| 70–79 | | 73.8 | 0.8 | 49.8 | 54.6 | 56.8 | 62.0 | 71.7 | 82.4 | 91.4 | 96.3 | 103.5 |
| 80 and older | | 67.9 | 0.8 | 46.8 | 50.6 | 52.9 | 57.9 | 65.7 | 75.2 | 83.0 | 87.3 | 94.2 |

NOTES: Data exclude pregnant females. Sample size is unweighted; estimates are weighted.

Table 4. Weight in pounds for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Courand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|-------|----------------------|-------|-------|-------|--------|------------|-------|-------|-------|-------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Pounds | | | | | |
| 20 and older | 2,688 | 199.0 | 2.0 | 134.2 | 146.8 | 154.3 | 166.3 | 191.0 | 221.9 | 244.8 | 261.8 | 290.6 |
| 20–29 | 287 | 191.3 | 4.0 | 125.6 | 133.4 | 141.6 | 159.1 | 184.5 | 216.6 | 238.3 | 252.9 | 282.9 |
| 30-39 | 385 | 201.5 | 2.5 | 140.3 | 153.8 | 156.7 | 169.9 | 192.0 | 223.8 | 242.5 | 266.8 | 292.0 |
| 40–49 | 347 | 210.1 | 4.4 | 145.4 | 153.7 | 161.3 | 173.1 | 195.0 | 237.5 | 263.8 | 275.7 | 312.8 |
| 50–59 | 415 | 203.0 | 3.3 | 134.4 | 150.1 | 157.0 | 169.3 | 193.5 | 225.0 | 250.5 | 264.1 | 305.9 |
| 60–69 | 651 | 196.7 | 2.5 | 138.0 | 148.0 | 154.2 | 167.5 | 191.7 | 218.2 | 238.7 | 251.3 | 273.2 |
| 70–79 | 446 | 193.4 | 3.2 | 134.0 | 145.8 | 151.7 | 164.3 | 190.9 | 212.8 | 235.5 | 246.3 | 267.5 |
| 80 and older | 157 | 180.1 | 2.0 | 127.9 | 139.5 | 147.0 | 156.1 | 176.2 | 198.4 | 212.7 | 219.6 | 239.2 |
| Female | | | | | | | | | | | | |
| 20 and older | 3,254 | 171.8 | 1.7 | 111.8 | 120.6 | 127.4 | 138.0 | 162.0 | 196.7 | 220.1 | 236.3 | 262.7 |
| 20–29 | 337 | 164.9 | 4.6 | 105.0 | 116.5 | 120.9 | 130.9 | 151.9 | 182.7 | 216.5 | 236.5 | 267.4 |
| 30–39 | 491 | 178.3 | 2.0 | 113.4 | 122.1 | 129.9 | 139.2 | 165.3 | 205.7 | 237.9 | 252.7 | 277.5 |
| 40–49 | 442 | 177.3 | 3.0 | 116.4 | 123.4 | 129.5 | 140.9 | 164.8 | 199.8 | 222.9 | 245.8 | 272.7 |
| 50–59 | | 180.0 | 2.0 | 117.1 | 127.7 | 132.8 | 143.7 | 173.0 | 210.6 | 231.5 | 245.8 | 267.5 |
| 60–69 | 810 | 171.0 | 2.1 | 113.7 | 122.3 | 129.9 | 139.5 | 163.8 | 198.1 | 216.8 | 230.3 | 241.0 |
| 70–79 | | 162.8 | 1.8 | 109.9 | 120.3 | 125.1 | 136.6 | 158.0 | 181.6 | 201.4 | 212.2 | 228.2 |
| 80 and older | | 149.7 | 1.8 | 103.1 | 111.6 | 116.6 | 127.6 | 144.8 | 165.7 | 182.9 | 192.4 | 207.8 |

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 5. Height in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|-------|----------------------|-------|-------|-------|------------|------------|-------|-------|-------|-------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | Ce | entimeters | | | | | |
| 2–3 | 107 | 95.2 | 0.8 | 84.7 | 86.1 | 88.3 | 90.3 | 95.3 | 99.1 | 100.3 | 102.5 | 107.5 |
| 4–5 | 154 | 110.2 | 0.8 | 100.0 | 101.8 | 102.9 | 105.4 | 109.5 | 114.1 | 117.3 | 119.7 | 122.1 |
| 6–7 | 121 | 122.6 | 0.6 | 110.8 | 112.7 | 113.8 | 118.2 | 122.7 | 127.2 | 129.6 | 130.4 | 132.1 |
| 8–9 | 171 | 134.6 | 0.3 | 123.9 | 126.0 | 127.3 | 129.8 | 134.1 | 139.3 | 142.6 | 143.9 | 146.4 |
| 10–11 | 140 | 146.8 | 0.8 | † | 134.3 | 136.5 | 140.1 | 147.5 | 153.1 | 154.8 | 156.4 | † |
| 12–13 | 142 | 162.0 | 1.1 | 148.3 | 150.2 | 151.7 | 154.5 | 163.0 | 168.2 | 170.8 | 172.8 | 176.6 |
| 14–15 | 144 | 171.1 | 0.7 | 157.3 | 160.9 | 162.4 | 166.0 | 171.7 | 176.4 | 178.4 | 179.7 | 182.2 |
| 16–17 | 136 | 175.0 | 0.6 | 163.2 | 166.8 | 167.9 | 169.6 | 174.4 | 179.7 | 182.6 | 186.2 | 189.6 |
| 18–19 | 138 | 175.1 | 0.7 | 161.4 | 164.6 | 165.5 | 169.0 | 174.4 | 181.2 | 184.1 | 186.0 | 187.3 |
| Female | | | | | | | | | | | | |
| 2–3 | 93 | 94.4 | 0.8 | 82.9 | 85.0 | 87.1 | 89.2 | 94.3 | 99.0 | 101.6 | 102.7 | 105.5 |
| 4–5 | 149 | 108.9 | 0.6 | 98.8 | 100.4 | 101.8 | 103.7 | 108.5 | 112.7 | 116.6 | 119.1 | 121.8 |
| 6–7 | 149 | 122.7 | 0.6 | 113.1 | 114.1 | 114.8 | 117.0 | 122.2 | 127.2 | 129.9 | 131.3 | 134.3 |
| 8–9 | 135 | 134.8 | 0.8 | 123.3 | 126.4 | 127.8 | 129.6 | 133.8 | 139.1 | 142.7 | 144.4 | 146.0 |
| 10–11 | 151 | 148.7 | 0.8 | 133.9 | 137.3 | 140.2 | 142.9 | 147.4 | 153.5 | 158.4 | 160.6 | 163.4 |
| 12–13 | 162 | 156.4 | 0.4 | 142.2 | 145.4 | 149.2 | 151.7 | 157.2 | 161.3 | 164.3 | 164.8 | 166.6 |
| 14–15 | 126 | 159.6 | 0.7 | 150.2 | 152.0 | 153.1 | 155.9 | 159.3 | 163.9 | 165.3 | 166.6 | 169.4 |
| 16–17 | 147 | 161.5 | 0.6 | † | 152.5 | 153.3 | 156.0 | 161.1 | 166.2 | 169.1 | 170.3 | 171.7 |
| 18–19 | 128 | 161.5 | 0.7 | 150.4 | 151.5 | 153.2 | 156.2 | 162.1 | 167.0 | 169.2 | 170.0 | 171.6 |

[†] Estimate not shown because the standard error of the 5th and 95th percentiles could not be computed.

Age is at time of examination.

NOTE: Sample size is unweighted; estimates are weighted.

Table 6. Height in inches for children and adolescents ages 2-19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|------|----------------------|------|------|------|--------|------------|------|------|------|------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Inches | | | | | |
| 2–3 | 107 | 37.5 | 0.3 | 33.3 | 33.9 | 34.8 | 35.5 | 37.5 | 39.0 | 39.5 | 40.4 | 42.3 |
| 4–5 | 154 | 43.4 | 0.3 | 39.4 | 40.1 | 40.5 | 41.5 | 43.1 | 44.9 | 46.2 | 47.1 | 48.1 |
| 6–7 | 121 | 48.3 | 0.3 | 43.6 | 44.4 | 44.8 | 46.5 | 48.3 | 50.1 | 51.0 | 51.3 | 52.0 |
| 8–9 | 171 | 53.0 | 0.1 | 48.8 | 49.6 | 50.1 | 51.1 | 52.8 | 54.9 | 56.1 | 56.7 | 57.6 |
| 10–11 | 140 | 57.8 | 0.3 | † | 52.9 | 53.7 | 55.1 | 58.1 | 60.3 | 60.9 | 61.6 | † |
| 12–13 | 142 | 63.8 | 0.4 | 58.4 | 59.1 | 59.7 | 60.8 | 64.2 | 66.2 | 67.2 | 68.0 | 69.5 |
| 14–15 | 144 | 67.4 | 0.3 | 61.9 | 63.4 | 63.9 | 65.4 | 67.6 | 69.5 | 70.2 | 70.7 | 71.7 |
| 16–17 | 136 | 68.9 | 0.2 | 64.2 | 65.7 | 66.1 | 66.8 | 68.7 | 70.8 | 71.9 | 73.3 | 74.7 |
| 18–19 | 138 | 68.9 | 0.3 | 63.6 | 64.8 | 65.1 | 66.5 | 68.7 | 71.4 | 72.5 | 73.2 | 73.7 |
| Female | | | | | | | | | | | | |
| 2–3 | 93 | 37.2 | 0.3 | 32.6 | 33.5 | 34.3 | 35.1 | 37.1 | 39.0 | 40.0 | 40.5 | 41.5 |
| 4–5 | 149 | 42.9 | 0.2 | 38.9 | 39.5 | 40.1 | 40.8 | 42.7 | 44.4 | 45.9 | 46.9 | 48.0 |
| 6–7 | 149 | 48.3 | 0.2 | 44.5 | 44.9 | 45.2 | 46.1 | 48.1 | 50.1 | 51.2 | 51.7 | 52.9 |
| 8–9 | 135 | 53.1 | 0.3 | 48.5 | 49.8 | 50.3 | 51.0 | 52.7 | 54.8 | 56.2 | 56.9 | 57.5 |
| 10–11 | 151 | 58.5 | 0.3 | 52.7 | 54.1 | 55.2 | 56.2 | 58.0 | 60.4 | 62.4 | 63.2 | 64.3 |
| 12–13 | 162 | 61.6 | 0.2 | 56.0 | 57.2 | 58.7 | 59.7 | 61.9 | 63.5 | 64.7 | 64.9 | 65.6 |
| 14–15 | 126 | 62.8 | 0.3 | 59.1 | 59.8 | 60.3 | 61.4 | 62.7 | 64.5 | 65.1 | 65.6 | 66.7 |
| 16–17 | | 63.6 | 0.2 | † | 60.0 | 60.3 | 61.4 | 63.4 | 65.4 | 66.6 | 67.1 | 67.6 |
| 18–19 | 128 | 63.6 | 0.3 | 59.2 | 59.6 | 60.3 | 61.5 | 63.8 | 65.7 | 66.6 | 66.9 | 67.6 |

[†] Estimate not shown because the standard error of the 5th and 95th percentiles could not be computed. ¹Age is at time of examination.

NOTE: Sample size is unweighted; estimates are weighted.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 7. Height in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|-------|----------------------|-------|-------|-------|------------|------------|-------|-------|-------|-------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | С | entimeters | | | | | |
| 20 and older | 2,690 | 175.1 | 0.3 | 162.2 | 165.2 | 167.5 | 170.0 | 175.2 | 180.1 | 183.0 | 184.8 | 187.2 |
| 20-29 | 287 | 176.3 | 0.6 | 162.0 | 167.7 | 169.3 | 171.3 | 176.1 | 181.0 | 184.7 | 185.9 | 188.1 |
| 30-39 | 385 | 176.1 | 0.4 | 163.6 | 166.6 | 168.0 | 171.3 | 176.4 | 180.9 | 183.4 | 185.1 | 187.1 |
| 40–49 | 348 | 176.4 | 0.6 | 162.3 | 166.9 | 169.5 | 171.1 | 176.2 | 182.6 | 185.0 | 186.8 | 189.9 |
| 50–59 | 417 | 174.3 | 0.4 | 163.0 | 164.7 | 167.2 | 169.0 | 174.5 | 179.0 | 181.3 | 183.6 | 185.6 |
| 60–69 | 652 | 174.7 | 0.5 | 161.8 | 165.2 | 167.0 | 169.7 | 175.0 | 179.8 | 182.5 | 184.1 | 186.6 |
| 70–79 | 446 | 172.4 | 0.4 | 159.5 | 162.6 | 164.6 | 167.5 | 172.7 | 176.9 | 179.8 | 181.6 | 184.1 |
| 80 and older | 155 | 170.5 | 0.7 | 158.3 | 160.1 | 162.8 | 166.1 | 169.9 | 175.0 | 177.9 | 179.3 | † |
| Female | | | | | | | | | | | | |
| 20 and older | 3.307 | 161.2 | 0.2 | 149.6 | 152.0 | 153.9 | 156.4 | 161.2 | 166.1 | 168.6 | 170.4 | 172.5 |
| 20–29 | 357 | 163.5 | 0.4 | 153.2 | 155.1 | 156.1 | 158.4 | 163.3 | 168.2 | 170.8 | 171.9 | 174.6 |
| 30–39 | | 162.5 | 0.4 | 150.1 | 152.8 | 154.8 | 158.2 | 162.8 | 167.4 | 169.5 | 170.8 | 172.9 |
| 40–49 | 444 | 161.7 | 0.3 | 149.8 | 152.3 | 153.7 | 157.2 | 162.0 | 166.3 | 169.0 | 170.4 | 172.1 |
| 50–59 | | 161.2 | 0.3 | 150.4 | 152.6 | 154.4 | 156.4 | 160.8 | 166.2 | 168.4 | 170.4 | 172.4 |
| 60–69 | | 160.5 | 0.3 | 149.6 | 151.6 | 153.4 | 155.6 | 160.4 | 165.1 | 167.6 | 169.4 | 171.2 |
| 70–79 | | 158.3 | 0.4 | 148.6 | 150.7 | 152.2 | 154.4 | 157.9 | 162.3 | 164.4 | 166.2 | 168.2 |
| 80 and older | | 156.8 | 0.4 | 143.7 | 145.9 | 149.6 | 151.2 | 156.9 | 161.9 | 164.6 | 165.5 | † |

[†] Estimate not shown because the standard error of the 95th percentile could not be computed.

NOTE: Sample size is unweighted; estimates are weighted.

Table 8. Height in inches for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023

| Courand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|----------------------|------|------|------|--------|------------|------|------|------|------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Inches | | | | | |
| 20 and older | 2,690 | 68.9 | 0.1 | 63.9 | 65.0 | 65.9 | 66.9 | 69.0 | 70.9 | 72.1 | 72.7 | 73.7 |
| 20–29 | 287 | 69.4 | 0.2 | 63.8 | 66.0 | 66.6 | 67.5 | 69.3 | 71.2 | 72.7 | 73.2 | 74.1 |
| 30–39 | 385 | 69.3 | 0.2 | 64.4 | 65.6 | 66.1 | 67.4 | 69.5 | 71.2 | 72.2 | 72.9 | 73.7 |
| 40–49 | 348 | 69.4 | 0.3 | 63.9 | 65.7 | 66.7 | 67.4 | 69.4 | 71.9 | 72.9 | 73.5 | 74.8 |
| 50–59 | 417 | 68.6 | 0.2 | 64.2 | 64.9 | 65.8 | 66.6 | 68.7 | 70.5 | 71.4 | 72.3 | 73.1 |
| 60–69 | 652 | 68.8 | 0.2 | 63.7 | 65.0 | 65.8 | 66.8 | 68.9 | 70.8 | 71.8 | 72.5 | 73.5 |
| 70–79 | 446 | 67.9 | 0.2 | 62.8 | 64.0 | 64.8 | 66.0 | 68.0 | 69.6 | 70.8 | 71.5 | 72.5 |
| 80 and older | 155 | 67.1 | 0.3 | 62.3 | 63.0 | 64.1 | 65.4 | 66.9 | 68.9 | 70.0 | 70.6 | † |
| Female | | | | | | | | | | | | |
| 20 and older | 3,307 | 63.5 | 0.1 | 58.9 | 59.8 | 60.6 | 61.6 | 63.5 | 65.4 | 66.4 | 67.1 | 67.9 |
| 20–29 | 357 | 64.4 | 0.2 | 60.3 | 61.1 | 61.4 | 62.4 | 64.3 | 66.2 | 67.2 | 67.7 | 68.7 |
| 30–39 | 511 | 64.0 | 0.1 | 59.1 | 60.2 | 60.9 | 62.3 | 64.1 | 65.9 | 66.7 | 67.2 | 68.1 |
| 40–49 | 444 | 63.7 | 0.1 | 59.0 | 60.0 | 60.5 | 61.9 | 63.8 | 65.5 | 66.5 | 67.1 | 67.7 |
| 50–59 | | 63.5 | 0.1 | 59.2 | 60.1 | 60.8 | 61.6 | 63.3 | 65.4 | 66.3 | 67.1 | 67.9 |
| 60–69 | 816 | 63.2 | 0.1 | 58.9 | 59.7 | 60.4 | 61.3 | 63.1 | 65.0 | 66.0 | 66.7 | 67.4 |
| 70–79 | | 62.3 | 0.1 | 58.5 | 59.3 | 59.9 | 60.8 | 62.2 | 63.9 | 64.7 | 65.4 | 66.2 |
| 80 and older | 176 | 61.7 | 0.2 | 56.6 | 57.5 | 58.9 | 59.5 | 61.8 | 63.8 | 64.8 | 65.2 | † |

[†] Estimate not shown because the standard error of the 95th percentile could not be computed.

NOTE: Sample size is unweighted; estimates are weighted.

Table 9. Body mass index values for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|------|----------------------|------|------|------|-----------|------------|------|------|------|------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | Body | mass inde | х | | | | |
| 2–3 | 107 | 16.8 | 0.2 | † | 14.6 | 15.0 | 15.5 | 16.6 | 17.5 | 18.2 | 18.6 | 19.2 |
| 4–5 | | 16.6 | 0.2 | 14.0 | 14.5 | 14.7 | 15.2 | 16.0 | 17.4 | 17.9 | 18.5 | 21.8 |
| 6–7 | | 16.9 | 0.3 | 13.5 | 14.1 | 14.6 | 14.8 | 16.1 | 17.7 | 20.0 | 21.0 | 22.2 |
| 8–9 | 171 | 18.3 | 0.4 | 14.0 | 14.3 | 14.8 | 15.5 | 17.2 | 19.6 | 21.9 | 23.2 | 26.8 |
| 10–11 | 140 | 21.3 | 0.7 | 15.2 | 15.4 | 16.0 | 16.7 | 19.0 | 25.0 | 26.6 | 29.6 | 33.3 |
| 12–13 | 142 | 22.1 | 0.5 | 16.1 | 16.7 | 17.3 | 18.1 | 20.3 | 24.8 | 27.5 | 28.4 | 32.8 |
| 14–15 | 144 | 24.2 | 0.6 | 16.2 | 17.0 | 18.5 | 19.5 | 21.6 | 27.6 | 32.3 | 35.0 | 38.1 |
| 16–17 | 136 | 24.9 | 0.7 | 16.9 | 17.7 | 18.1 | 20.1 | 23.7 | 27.9 | 30.5 | 33.2 | 36.0 |
| 18–19 | 138 | 25.0 | 0.7 | 17.4 | 18.7 | 19.5 | 20.4 | 23.6 | 27.9 | 32.3 | 32.7 | 36.9 |
| Female | | | | | | | | | | | | |
| 2–3 | 93 | 16.4 | 0.3 | † | 14.2 | 14.4 | 15.1 | 16.1 | 17.4 | 18.1 | 18.4 | 19.1 |
| 4–5 | 149 | 16.3 | 0.2 | 13.6 | 14.1 | 14.3 | 14.8 | 15.6 | 16.8 | 18.5 | 19.9 | 21.3 |
| 6–7 | 149 | 17.1 | 0.4 | 13.5 | 14.0 | 14.2 | 14.9 | 15.9 | 17.9 | 20.7 | 21.9 | 23.0 |
| 8–9 | | 18.7 | 0.5 | 13.8 | 14.5 | 14.9 | 15.4 | 17.7 | 20.7 | 22.5 | 25.0 | 26.8 |
| 10–11 | 151 | 20.6 | 0.4 | 15.0 | 15.9 | 16.3 | 16.8 | 19.4 | 24.0 | 25.1 | 27.4 | 29.2 |
| 12–13 | | 22.6 | 0.6 | 16.6 | 17.2 | 17.9 | 18.8 | 21.4 | 24.9 | 26.6 | 28.3 | 32.3 |
| 14–15 | 126 | 24.2 | 0.6 | † | 17.9 | 18.8 | 19.7 | 22.2 | 27.3 | 30.2 | 33.4 | 37.4 |
| 16–17 | 147 | 24.6 | 0.7 | 17.9 | 18.1 | 19.0 | 20.0 | 22.6 | 27.6 | 30.3 | 33.3 | 36.4 |
| 18–19 | | 26.4 | 0.8 | 17.3 | 18.1 | 18.9 | 20.8 | 24.7 | 30.1 | 33.0 | 34.9 | 41.4 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed.

NOTES: Data exclude pregnant females. Body mass index (BMI) is calculated as: BMI equals weight in kilograms divided by height in meters squared. Sample size is unweighted; estimates are weighted.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 10. Body mass index values for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| • | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|----------------------|------|------|------|------------|------------|------|------|------|------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | Bod | y mass Ind | ex | | | | |
| 20 and older | 2,680 | 29.4 | 0.3 | 20.9 | 22.3 | 23.4 | 24.9 | 28.2 | 32.6 | 35.1 | 37.8 | 42.1 |
| 20–29 | 287 | 27.8 | 0.5 | 18.9 | 19.9 | 21.6 | 23.3 | 27.0 | 31.1 | 34.2 | 36.1 | 39.6 |
| 30–39 | 385 | 29.5 | 0.4 | 21.2 | 22.7 | 23.3 | 24.9 | 27.8 | 33.0 | 35.6 | 38.0 | 42.5 |
| 40–49 | 347 | 30.5 | 0.6 | 22.2 | 23.3 | 24.1 | 25.7 | 28.9 | 33.5 | 35.8 | 39.2 | 45.0 |
| 50-59 | 415 | 30.2 | 0.4 | 21.6 | 23.0 | 24.1 | 25.5 | 29.0 | 33.3 | 36.6 | 38.9 | 44.6 |
| 60-69 | 650 | 29.2 | 0.3 | 21.2 | 22.7 | 23.9 | 25.2 | 28.3 | 32.1 | 34.1 | 36.0 | 39.9 |
| 70–79 | 444 | 29.4 | 0.4 | 21.6 | 22.9 | 23.9 | 25.6 | 28.8 | 32.3 | 34.6 | 36.5 | 39.2 |
| 80 and older | 152 | 27.8 | 0.4 | 21.1 | 22.8 | 24.2 | 24.6 | 27.2 | 30.2 | 32.1 | 32.6 | 34.8 |
| Female | | | | | | | | | | | | |
| 20 and older | 3.249 | 30.0 | 0.3 | 19.7 | 21.5 | 22.4 | 24.1 | 28.3 | 34.4 | 38.4 | 40.9 | 44.9 |
| 20–29 | 336 | 28.0 | 0.7 | 18.6 | 19.9 | 21.0 | 22.2 | 25.5 | 31.6 | 36.6 | 39.7 | 45.6 |
| 30–39 | | 30.5 | 0.4 | 19.7 | 21.3 | 22.3 | 24.2 | 28.6 | 35.2 | 40.4 | 42.1 | 47.4 |
| 40–49 | 441 | 30.7 | 0.6 | 19.7 | 21.9 | 23.0 | 24.6 | 28.5 | 35.1 | 38.8 | 41.0 | 45.6 |
| 50–59 | 506 | 31.4 | 0.3 | 20.8 | 22.3 | 23.4 | 24.9 | 30.0 | 36.5 | 40.2 | 42.4 | 45.9 |
| 60-69 | 810 | 30.1 | 0.4 | 20.0 | 22.0 | 23.1 | 24.8 | 28.8 | 34.5 | 37.9 | 40.0 | 42.5 |
| 70–79 | 493 | 29.5 | 0.4 | 20.7 | 22.0 | 23.2 | 24.7 | 28.0 | 33.4 | 36.5 | 38.3 | 42.2 |
| 80 and older | 173 | 27.7 | 0.4 | 19.2 | 21.1 | 22.0 | 23.6 | 26.9 | 31.3 | 33.3 | 34.7 | 36.3 |

NOTES: Data exclude pregnant females. Body mass index (BMI) is calculated as: BMI equals weight in kilograms divided by height in meters squared. Sample size is unweighted; estimates are weighted.

¹Age is at time of examination.

Table 11. Waist circumference in centimeters for children and adolescents ages 2-19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|------|-------------------|------|------|------|------------|------------|------|-------|-------|-------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | C | entimeters | | | | | |
| 2–3 | 106 | 49.8 | 0.5 | 43.7 | 45.5 | 45.9 | 47.3 | 49.3 | 51.4 | 52.7 | 53.3 | 56.6 |
| 4–5 | 150 | 54.5 | 0.6 | 46.0 | 48.2 | 48.8 | 50.7 | 53.4 | 56.0 | 58.4 | 62.8 | 68.3 |
| 6–7 | 116 | 57.9 | 1.0 | † | 51.1 | 51.7 | 52.8 | 55.6 | 61.2 | 66.4 | 69.0 | 73.0 |
| 8–9 | 167 | 63.9 | 0.9 | 53.2 | 54.3 | 55.4 | 56.7 | 60.4 | 68.3 | 73.1 | 75.9 | 84.6 |
| 10–11 | 139 | 73.8 | 1.7 | † | 57.1 | 59.0 | 61.3 | 68.2 | 83.4 | 92.4 | 97.5 | 105.8 |
| 12–13 | 139 | 78.3 | 1.3 | 61.2 | 62.7 | 64.0 | 67.2 | 73.2 | 87.4 | 91.9 | 96.1 | † |
| 14–15 | 139 | 83.1 | 1.5 | 63.6 | 66.9 | 68.5 | 69.6 | 77.1 | 91.8 | 105.5 | 109.9 | 116.7 |
| 16–17 | 133 | 85.3 | 2.0 | † | 67.8 | 68.8 | 73.4 | 81.3 | 93.9 | 102.2 | 109.1 | 116.7 |
| 18–19 | 134 | 87.0 | 1.9 | 68.4 | 70.4 | 71.9 | 75.5 | 81.9 | 97.3 | 103.5 | 107.3 | 115.3 |
| Female | | | | | | | | | | | | |
| 2–3 | 88 | 49.2 | 0.5 | 41.3 | 43.3 | 44.8 | 46.1 | 49.4 | 51.2 | 53.3 | 53.7 | 57.6 |
| 4–5 | 141 | 54.1 | 0.6 | 46.7 | 48.5 | 49.1 | 50.2 | 52.0 | 55.5 | 60.5 | 64.5 | 67.3 |
| 6–7 | 146 | 59.4 | 0.9 | 49.5 | 50.7 | 51.4 | 53.4 | 57.3 | 61.2 | 69.0 | 73.0 | 78.4 |
| 8–9 | 133 | 65.0 | 1.2 | 52.7 | 54.2 | 55.2 | 56.5 | 62.6 | 69.0 | 76.4 | 80.8 | 87.7 |
| 10–11 | 150 | 72.1 | 0.8 | 57.3 | 60.1 | 60.9 | 62.9 | 69.3 | 79.9 | 86.7 | 89.5 | 93.4 |
| 12–13 | 155 | 76.7 | 1.1 | 62.5 | 64.8 | 66.4 | 67.5 | 74.9 | 81.1 | 86.2 | 92.6 | 97.3 |
| 14–15 | 118 | 80.2 | 1.5 | † | 66.3 | 67.3 | 70.0 | 77.2 | 87.0 | 97.3 | 99.3 | 104.2 |
| 16–17 | 141 | 82.2 | 1.5 | 65.2 | 67.1 | 68.6 | 71.1 | 78.6 | 89.8 | 97.9 | 104.6 | 109.0 |
| 18–19 | 124 | 85.0 | 1.7 | 64.5 | 67.0 | 68.0 | 71.4 | 80.7 | 92.4 | 100.5 | 108.0 | 119.4 |

 $^{\ \ ^1}$ Estimate not shown because the standard error of the 5th and 95th percentiles could not be computed. $\ ^1$ Age is at time of examination.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 12. Waist circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Courand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|-------|-------------------|------|------|------|-------------|------------|-------|-------|-------|-------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | (| Centimeters | 3 | | | | |
| 20 and older | 2,605 | 103.0 | 0.7 | 79.3 | 83.5 | 86.8 | 91.4 | 100.9 | 112.3 | 119.8 | 124.3 | 133.2 |
| 20-29 | 282 | 95.1 | 1.3 | 71.3 | 75.7 | 77.9 | 82.6 | 93.4 | 106.1 | 112.6 | 117.8 | 124.4 |
| 30-39 | 378 | 101.3 | 0.9 | 80.3 | 82.6 | 85.9 | 90.0 | 98.6 | 109.9 | 118.9 | 123.1 | 130.5 |
| 40–49 | 340 | 104.7 | 1.4 | 83.7 | 86.6 | 88.3 | 92.2 | 101.5 | 114.0 | 120.7 | 127.7 | 139.6 |
| 50-59 | 407 | 106.4 | 1.0 | 83.2 | 88.2 | 90.5 | 94.2 | 104.3 | 114.8 | 123.4 | 128.4 | 137.0 |
| 60-69 | 631 | 105.4 | 0.9 | 84.2 | 87.4 | 90.2 | 95.5 | 104.5 | 113.1 | 119.5 | 123.4 | 132.2 |
| 70–79 | 429 | 107.5 | 1.1 | 84.6 | 88.6 | 92.8 | 97.3 | 106.6 | 117.1 | 122.6 | 126.5 | 131.3 |
| 80 and older | 138 | 104.1 | 1.1 | 84.5 | 89.6 | 92.8 | 95.3 | 102.6 | 111.4 | 117.0 | 120.5 | 126.0 |
| Female | | | | | | | | | | | | |
| 20 and older | 3,121 | 97.9 | 0.7 | 73.8 | 77.0 | 80.0 | 84.9 | 95.7 | 109.2 | 116.9 | 121.5 | 129.2 |
| 20–29 | 329 | 90.5 | 1.7 | 68.5 | 72.3 | 74.5 | 77.2 | 85.3 | 98.5 | 112.3 | 118.6 | 127.3 |
| 30–39 | 472 | 97.8 | 0.8 | 72.6 | 75.6 | 78.2 | 83.9 | 93.1 | 108.8 | 121.6 | 126.6 | 133.6 |
| 40–49 | 420 | 98.2 | 1.0 | 75.3 | 77.0 | 81.3 | 85.2 | 95.7 | 109.2 | 115.9 | 121.6 | 130.8 |
| 50-59 | 491 | 101.5 | 0.6 | 77.7 | 81.5 | 84.0 | 88.4 | 100.8 | 112.6 | 119.3 | 122.0 | 130.8 |
| 60-69 | 782 | 100.3 | 0.7 | 77.1 | 81.7 | 83.6 | 88.8 | 99.6 | 110.4 | 117.1 | 120.7 | 125.5 |
| 70–79 | 465 | 99.6 | 0.8 | 76.7 | 81.2 | 84.6 | 89.1 | 98.9 | 108.4 | 115.3 | 119.3 | 123.7 |
| 80 and older | 162 | 96.5 | 1.1 | 76.4 | 79.3 | 84.1 | 88.8 | 94.4 | 107.2 | 110.5 | 111.9 | 119.1 |

NOTES: Data exclude pregnant females. Sample size is unweighted; estimates are weighted.

Table 13. Hip circumference in centimeters for children and adolescents ages 12-19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|-------|----------------------|------|------|------|------------|------------|-------|-------|-------|-------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | Ce | entimeters | | | | | |
| 12–13 | 131 | 90.5 | 1.0 | 74.2 | 77.4 | 80.4 | 82.8 | 89.2 | 95.3 | 101.9 | 103.9 | 112.1 |
| 14–15 | 136 | 97.0 | 1.3 | 76.9 | 81.6 | 85.1 | 87.2 | 93.6 | 104.5 | 110.9 | 118.3 | 124.5 |
| 16–17 | 132 | 99.9 | 1.5 | † | 85.5 | 87.5 | 90.6 | 99.2 | 105.5 | 109.6 | 116.9 | 121.8 |
| 18–19 | 134 | 100.1 | 1.4 | 84.1 | 85.1 | 87.9 | 90.6 | 98.2 | 107.1 | 111.8 | 114.3 | 123.5 |
| Female | | | | | | | | | | | | |
| 12–13 | 153 | 91.9 | 1.1 | 76.8 | 79.6 | 82.6 | 84.6 | 90.5 | 96.2 | 101.7 | 105.1 | 112.6 |
| 14–15 | 117 | 96.9 | 1.3 | † | 85.7 | 87.1 | 88.9 | 94.1 | 102.8 | 108.2 | 111.3 | 121.5 |
| 16–17 | 140 | 99.1 | 1.3 | 85.5 | 86.6 | 87.0 | 89.5 | 96.6 | 104.4 | 109.1 | 114.3 | 122.4 |
| 18–19 | 123 | 101.8 | 1.5 | 83.9 | 87.2 | 88.4 | 91.1 | 99.2 | 110.0 | 113.2 | 119.4 | 130.3 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed.

Age is at time of examination.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 14. Hip circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| 0 | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|-------|------------------------|------|------|------|------------|------------|-------|-------|-------|-------|
| Sex and age group | examined people | Mean | error of - the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Centimeter | S | | | | |
| 20 and older | 2,569 | 105.6 | 0.5 | 89.5 | 92.6 | 94.5 | 97.7 | 103.6 | 110.8 | 115.8 | 120.7 | 129.1 |
| 20-29 | 278 | 104.5 | 0.9 | 87.0 | 89.6 | 92.5 | 96.2 | 103.0 | 110.1 | 115.4 | 119.4 | 127.8 |
| 30–39 | 375 | 106.3 | 0.7 | 91.0 | 93.7 | 95.6 | 98.5 | 104.7 | 111.2 | 115.6 | 121.0 | 127.6 |
| 40–49 | 334 | 107.4 | 1.1 | 92.1 | 94.2 | 96.3 | 98.8 | 104.3 | 113.3 | 117.9 | 122.6 | 133.8 |
| 50-59 | 402 | 106.2 | 0.8 | 89.7 | 92.3 | 94.4 | 97.4 | 103.7 | 110.6 | 119.3 | 122.7 | 134.8 |
| 60–69 | 622 | 104.2 | 0.6 | 89.5 | 91.9 | 93.8 | 96.9 | 103.0 | 109.3 | 113.2 | 118.2 | 123.6 |
| 70–79 | 422 | 105.0 | 0.8 | 90.0 | 92.3 | 94.0 | 96.8 | 103.2 | 110.5 | 115.3 | 121.1 | 126.1 |
| 80 and older | 136 | 102.5 | 0.8 | 90.3 | 92.0 | 94.0 | 96.1 | 101.4 | 107.1 | 109.0 | 112.7 | 123.0 |
| Female | | | | | | | | | | | | |
| 20 and older | 3.098 | 109.8 | 0.5 | 89.7 | 92.9 | 95.5 | 98.6 | 106.8 | 118.1 | 126.0 | 131.5 | 138.4 |
| 20–29 | 328 | 107.0 | 1.2 | 87.1 | 91.7 | 93.4 | 96.6 | 104.4 | 113.7 | 121.1 | 127.8 | 135.7 |
| 30–39 | 466 | 111.6 | 0.7 | 91.0 | 94.0 | 96.5 | 99.9 | 108.1 | 120.0 | 129.9 | 135.2 | 142.0 |
| 40–49 | 420 | 111.1 | 0.9 | 90.3 | 94.7 | 96.5 | 99.5 | 107.4 | 120.9 | 127.2 | 130.2 | 139.7 |
| 50–59 | 490 | 111.5 | 0.8 | 90.1 | 93.5 | 96.3 | 99.6 | 108.3 | 121.3 | 128.8 | 133.6 | 139.2 |
| 60–69 | 780 | 110.1 | 0.8 | 88.9 | 92.7 | 95.4 | 99.2 | 108.0 | 119.5 | 126.2 | 131.1 | 137.3 |
| 70–79 | 460 | 108.6 | 0.7 | 89.4 | 93.1 | 95.6 | 98.7 | 106.4 | 116.1 | 123.3 | 128.3 | 134.4 |
| 80 and older | 154 | 104.7 | 0.7 | 87.7 | 91.1 | 92.8 | 96.8 | 103.1 | 112.2 | 117.0 | 119.8 | † |

[†] Estimate not shown because the standard error of the 95th percentile could not be computed.

NOTES: Data exclude pregnant females. Sample size is unweighted; estimates are weighted.

NOTES: Data exclude pregnant females. Sample size is unweighted; estimates are weighted.

Table 15. Mid-upper arm circumference in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|------|----------------------|------|------|------|------------|------------|------|------|------|------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | Ce | entimeters | | | | | |
| 2–3 | 112 | 16.6 | 0.2 | 14.2 | 15.0 | 15.1 | 15.6 | 16.4 | 17.1 | 18.0 | 18.4 | 19.1 |
| 4–5 | 151 | 18.1 | 0.2 | 15.1 | 15.7 | 16.3 | 16.8 | 17.8 | 18.8 | 19.5 | 20.2 | 21.9 |
| 6–7 | 118 | 19.3 | 0.3 | † | 16.6 | 16.9 | 17.5 | 18.7 | 20.1 | 22.9 | 23.5 | 24.1 |
| 8–9 | 169 | 21.5 | 0.3 | 17.2 | 17.7 | 18.3 | 19.0 | 20.3 | 23.4 | 24.6 | 25.9 | 29.5 |
| 10–11 | 139 | 25.1 | 0.6 | 18.5 | 19.4 | 19.8 | 21.5 | 23.5 | 28.0 | 30.8 | 32.8 | 34.5 |
| 12-13 | 140 | 27.1 | 0.4 | 21.0 | 21.8 | 22.6 | 23.9 | 26.1 | 29.6 | 31.3 | 34.1 | 36.0 |
| 14–15 | 141 | 29.7 | 0.6 | 21.7 | 22.8 | 24.4 | 26.0 | 28.7 | 33.4 | 36.2 | 37.8 | 40.4 |
| 16–17 | 133 | 31.4 | 0.7 | † | 24.6 | 25.6 | 27.1 | 31.4 | 34.5 | 37.5 | 38.6 | 40.1 |
| 18–19 | 136 | 31.4 | 0.6 | 23.7 | 25.6 | 26.9 | 27.9 | 30.6 | 34.0 | 37.0 | 38.4 | 40.7 |
| Female | | | | | | | | | | | | |
| 2–3 | 91 | 16.6 | 0.2 | 14.2 | 14.7 | 15.0 | 15.4 | 16.3 | 17.6 | 18.0 | 18.2 | 19.0 |
| 4–5 | 147 | 17.9 | 0.2 | 15.2 | 15.7 | 16.0 | 16.5 | 17.4 | 18.3 | 19.5 | 21.8 | 22.7 |
| 6–7 | 147 | 19.8 | 0.3 | 16.3 | 16.4 | 17.0 | 17.8 | 19.1 | 21.0 | 23.0 | 23.7 | 25.7 |
| 8–9 | | 22.1 | 0.5 | † | 18.2 | 18.5 | 19.4 | 21.3 | 24.0 | 25.5 | 27.5 | 29.3 |
| 10–11 | 151 | 24.7 | 0.3 | 19.1 | 20.0 | 20.5 | 21.2 | 23.8 | 27.4 | 29.1 | 30.2 | 32.4 |
| 12–13 | | 26.7 | 0.5 | 20.7 | 21.5 | 22.0 | 23.2 | 25.9 | 28.7 | 31.1 | 33.2 | 35.4 |
| 14–15 | | 28.3 | 0.4 | 19.7 | 22.9 | 23.5 | 25.4 | 26.8 | 30.7 | 32.7 | 35.0 | 38.5 |
| 16–17 | | 29.1 | 0.5 | 23.3 | 23.7 | 24.3 | 25.5 | 27.8 | 31.6 | 33.4 | 35.8 | 39.5 |
| 18–19 | | 30.4 | 0.6 | 22.3 | 23.2 | 23.9 | 25.6 | 29.8 | 33.8 | 37.5 | 38.4 | 40.6 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed. ¹Age is at time of examination.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 16. Mid-upper arm circumference in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Courand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|----------------------|------|------|------|------------|------------|------|------|------|------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | С | entimeters | | | | | |
| 20 and older | 2,652 | 34.7 | 0.2 | 27.7 | 29.0 | 30.0 | 31.5 | 34.4 | 37.5 | 39.2 | 40.4 | 42.9 |
| 20–29 | 285 | 34.3 | 0.5 | 25.9 | 27.9 | 29.1 | 31.1 | 34.4 | 37.3 | 38.7 | 40.1 | 42.3 |
| 30-39 | 381 | 35.4 | 0.3 | 28.1 | 29.8 | 30.9 | 32.2 | 34.9 | 38.0 | 39.7 | 40.8 | 43.2 |
| 40–49 | 343 | 36.0 | 0.4 | 28.9 | 30.2 | 31.2 | 32.9 | 35.6 | 38.6 | 40.3 | 42.5 | 44.2 |
| 50–59 | 415 | 35.2 | 0.3 | 28.4 | 29.7 | 30.2 | 31.7 | 34.5 | 38.1 | 39.7 | 41.4 | 44.4 |
| 60–69 | 641 | 34.0 | 0.2 | 27.5 | 29.0 | 29.9 | 31.2 | 33.6 | 36.4 | 37.7 | 39.0 | 40.8 |
| 70–79 | 436 | 33.4 | 0.4 | 27.4 | 28.5 | 29.2 | 30.4 | 33.3 | 35.9 | 37.7 | 39.0 | 40.1 |
| 80 and older | 151 | 31.7 | 0.3 | 26.1 | 27.5 | 27.9 | 29.1 | 31.2 | 34.1 | 35.9 | 37.0 | 38.5 |
| Female | | | | | | | | | | | | |
| 20 and older | 3.198 | 32.9 | 0.2 | 25.1 | 26.4 | 27.2 | 28.7 | 32.1 | 36.1 | 38.9 | 40.5 | 43.6 |
| 20–29 | 333 | 31.9 | 0.5 | 24.3 | 25.3 | 26.0 | 27.6 | 30.9 | 35.1 | 37.6 | 39.8 | 43.3 |
| 30–39 | 480 | 33.9 | 0.3 | 25.6 | 26.5 | 27.2 | 29.7 | 32.6 | 37.5 | 40.9 | 43.2 | 45.9 |
| 40–49 | 436 | 33.6 | 0.5 | 25.5 | 26.8 | 27.8 | 29.2 | 32.5 | 36.6 | 39.3 | 41.2 | 43.9 |
| 50–59 | 500 | 33.8 | 0.2 | 26.4 | 27.5 | 28.5 | 29.8 | 33.1 | 37.4 | 39.4 | 41.0 | 43.7 |
| 60–69 | 801 | 32.8 | 0.3 | 25.3 | 26.7 | 27.8 | 29.0 | 32.2 | 35.9 | 38.5 | 39.9 | 42.2 |
| 70–79 | 481 | 32.1 | 0.2 | 25.1 | 26.3 | 27.0 | 28.6 | 31.6 | 35.2 | 37.3 | 38.6 | 40.6 |
| 80 and older | 167 | 29.8 | 0.3 | 22.6 | 23.7 | 25.4 | 26.6 | 29.3 | 32.8 | 34.2 | 35.2 | 37.4 |

NOTES: Data exclude pregnant females. Sample size is unweighted; estimates are weighted.

Table 17. Upper arm length in centimeters for children and adolescents ages 2–19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021–August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|------|----------------------|------|------|------|------------|------------|------|------|------|------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | - | | | | | Ce | entimeters | | | | | |
| 2–3 | 112 | 19.4 | 0.2 | 16.3 | 17.1 | 17.6 | 18.0 | 19.2 | 20.2 | 20.9 | 21.6 | 22.0 |
| 4–5 | 151 | 22.6 | 0.2 | 20.0 | 20.5 | 20.9 | 21.4 | 22.5 | 23.7 | 24.1 | 24.6 | 25.9 |
| 6–7 | 118 | 25.7 | 0.2 | 22.1 | 22.9 | 23.5 | 24.3 | 26.0 | 27.0 | 27.7 | 28.0 | 28.5 |
| 8–9 | 169 | 28.7 | 0.1 | 25.3 | 25.9 | 26.2 | 27.2 | 28.7 | 29.9 | 30.5 | 31.3 | 32.5 |
| 10–11 | 139 | 31.8 | 0.2 | 27.0 | 28.4 | 28.9 | 29.7 | 31.8 | 33.6 | 34.5 | 35.2 | 36.4 |
| 12-13 | 140 | 35.5 | 0.3 | 31.7 | 32.3 | 33.1 | 33.8 | 35.5 | 37.3 | 38.0 | 38.4 | 39.0 |
| 14–15 | 141 | 37.8 | 0.2 | 33.4 | 34.3 | 35.0 | 35.9 | 37.9 | 39.1 | 40.4 | 41.1 | 42.3 |
| 16–17 | 133 | 39.1 | 0.2 | 35.9 | 36.5 | 37.0 | 37.4 | 38.9 | 40.2 | 41.3 | 41.9 | 42.6 |
| 18–19 | 136 | 38.6 | 0.2 | 34.7 | 35.4 | 35.9 | 36.8 | 38.5 | 40.1 | 41.4 | 41.9 | 42.2 |
| Female | | | | | | | | | | | | |
| 2–3 | 91 | 19.3 | 0.2 | 16.5 | 16.9 | 17.0 | 18.1 | 19.2 | 20.5 | 21.1 | 21.4 | 21.8 |
| 4–5 | 147 | 22.3 | 0.2 | 19.5 | 20.1 | 20.2 | 20.8 | 22.3 | 23.7 | 24.1 | 24.6 | 25.4 |
| 6–7 | 147 | 25.7 | 0.1 | 22.9 | 23.5 | 23.9 | 24.4 | 25.5 | 26.9 | 27.7 | 27.9 | 28.4 |
| 8–9 | 135 | 28.4 | 0.3 | 25.1 | 26.0 | 26.3 | 26.9 | 28.1 | 29.6 | 30.4 | 31.3 | 32.2 |
| 10–11 | 151 | 32.3 | 0.2 | 27.8 | 29.3 | 30.1 | 30.5 | 32.0 | 33.7 | 35.1 | 35.8 | 36.0 |
| 12–13 | 158 | 34.2 | 0.2 | 30.5 | 31.3 | 31.8 | 32.6 | 34.2 | 35.6 | 36.8 | 37.1 | 37.9 |
| 14–15 | 123 | 35.0 | 0.2 | 31.5 | 32.0 | 32.4 | 34.2 | 35.0 | 36.4 | 37.1 | 37.5 | 38.0 |
| 16–17 | 143 | 35.5 | 0.3 | † | 32.7 | 33.2 | 34.0 | 35.4 | 37.0 | 37.8 | 38.3 | 39.0 |
| 18–19 | 128 | 35.4 | 0.3 | 31.6 | 32.1 | 33.2 | 33.9 | 35.4 | 36.9 | 37.3 | 37.7 | 38.6 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed.

NOTE: Sample size is unweighted; estimates are weighted.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 18. Upper arm length in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Courand | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|----------------------|------|------|------|------------|------------|------|------|------|------|
| Sex and age group | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | C | entimeters | | | | | |
| 20 and older | 2,654 | 39.6 | 0.1 | 35.8 | 36.6 | 37.1 | 38.0 | 39.5 | 41.0 | 42.0 | 42.5 | 43.5 |
| 20–29 | 285 | 39.4 | 0.2 | 35.7 | 36.5 | 37.1 | 37.8 | 39.4 | 40.7 | 41.3 | 42.0 | 43.2 |
| 30–39 | 381 | 39.3 | 0.1 | 36.0 | 36.9 | 37.1 | 38.0 | 39.1 | 40.5 | 41.6 | 42.0 | 42.8 |
| 40–49 | 343 | 39.6 | 0.2 | 35.6 | 36.2 | 36.9 | 38.0 | 39.7 | 41.1 | 42.2 | 42.8 | 43.9 |
| 50-59 | 416 | 39.5 | 0.2 | 36.0 | 36.9 | 37.2 | 37.9 | 39.5 | 40.9 | 41.7 | 42.2 | 42.9 |
| 60-69 | 641 | 39.9 | 0.2 | 35.9 | 36.8 | 37.4 | 38.3 | 39.9 | 41.7 | 42.5 | 42.9 | 43.8 |
| 70–79 | 436 | 39.6 | 0.2 | 35.4 | 36.1 | 36.9 | 37.9 | 39.6 | 41.2 | 42.3 | 42.9 | 43.9 |
| 80 and older | 152 | 39.9 | 0.5 | 35.5 | 36.5 | 36.9 | 38.0 | 39.9 | 41.5 | 42.9 | 43.7 | 44.3 |
| Female | | | | | | | | | | | | |
| 20 and older | 3,239 | 36.2 | 0.1 | 32.4 | 33.2 | 33.8 | 34.6 | 36.1 | 37.6 | 38.4 | 39.0 | 39.8 |
| 20–29 | 352 | 36.0 | 0.2 | 32.5 | 33.3 | 33.8 | 34.6 | 35.9 | 37.3 | 38.2 | 38.9 | 39.5 |
| 30-39 | 500 | 36.3 | 0.1 | 32.3 | 33.1 | 33.6 | 34.5 | 36.4 | 37.9 | 38.6 | 39.0 | 39.9 |
| 40–49 | 437 | 36.1 | 0.2 | 32.2 | 32.9 | 33.5 | 34.6 | 36.0 | 37.6 | 38.5 | 39.1 | 39.6 |
| 50-59 | 500 | 36.2 | 0.2 | 32.2 | 33.4 | 33.9 | 34.6 | 36.0 | 37.9 | 38.5 | 39.1 | 39.9 |
| 60–69 | 800 | 36.3 | 0.1 | 32.8 | 33.3 | 33.9 | 34.6 | 36.3 | 37.7 | 38.5 | 39.0 | 39.9 |
| 70–79 | 482 | 36.1 | 0.1 | 32.5 | 33.4 | 33.8 | 34.5 | 36.1 | 37.3 | 38.0 | 38.7 | 39.8 |
| 80 and older | 168 | 36.0 | 0.2 | 32.1 | 32.7 | 33.5 | 34.5 | 36.1 | 37.2 | 37.9 | 38.4 | † |

[†] Estimate not shown because the standard error of the 95th percentile could not be computed.

NOTE: Sample size is unweighted; estimates are weighted.

¹Age is at time of examination.

Table 19. Upper leg length in centimeters for children and adolescents ages 8-19 years and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| Sex and | Number of | | Standard | | | | | Percentile | | | | |
|-----------------------------------|--------------------|--------------|----------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| age group (years) ¹ | examined people | Mean | error of the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | _ | | | | | Ce | entimeters | | | | | |
| 8–9 | 163 137 | 31.4 | 0.2 | 27.3 | 28.0 | 28.5 | 29.5 | 31.1 | 32.7 | 34.0 | 35.2 | 36.5 |
| 10–11 | | 35.2 39.1 | 0.3 0.3 | 29.9 34.0 | 31.1 34.9 | 31.9 36.2 | 32.9 36.8 | 35.2 38.9 | 37.4 41.1 | 37.9 42.5 | 38.7 42.9 | 39.6 43.5 |
| 14–15 | | 41.6 | 0.2 | 36.7 | 37.5 | 38.8 | 39.9 | 41.3 | 43.5 | 44.3 | 45.0 | 46.7 |
| 16–17 | | 42.3 41.9 | 0.3 0.2 | 38.0 37.2 | 38.8 38.3 | 39.5 38.9 | 40.3 39.5 | 42.1 41.9 | 44.5 44.0 | 45.3 45.2 | 46.0 45.5 | 46.5 46.4 |
| Female | | | | | | | | | | | | |
| 8–9 | | 31.5 35.2 | 0.3 0.2 | 27.2 30.9 | 28.3 32.1 | 28.6 32.7 | 29.4 33.2 | 31.6 35.0 | 33.0 36.8 | 33.5 38.0 | 34.7 38.8 | 35.8 39.3 |
| 12–13 | | 37.3 37.9 | 0.2 0.3 | 32.5 34.4 | 34.6 35.0 | 35.0 35.7 | 35.5 36.2 | 37.1 37.2 | 39.0 39.4 | 40.0 40.3 | 40.7 40.8 | 41.2 42.5 |
| 16–17 | | 37.8 38.3 | 0.3 0.4 | † 34.6 | 34.8 35.2 | 35.2 35.4 | 35.9 36.5 | 37.7 38.0 | 39.9 39.9 | 40.3 40.7 | 41.0 41.5 | 42.1 43.0 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed. ¹Age is at time of examination.

NOTE: Sample size is unweighted; estimates are weighted.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey.

Table 20. Upper leg length in centimeters for adults age 20 and older and number of examined people, mean, standard error of the mean, and selected percentiles, by sex and age: United States, August 2021-August 2023

| | Number of | | Standard | | | | | Percentile | | | | |
|-------------------|--------------------|------|------------------------|------|------|------|-------------|------------|------|------|------|------|
| Sex and age group | examined people | Mean | error of - the mean | 5th | 10th | 15th | 25th | 50th | 75th | 85th | 90th | 95th |
| Male | | | | | | | Centimeters | 3 | | | | |
| 20 and older | 2,559 | 41.0 | 0.2 | 35.8 | 37.1 | 38.0 | 39.1 | 41.0 | 43.1 | 44.1 | 44.8 | 45.9 |
| 20-29 | 277 | 42.3 | 0.3 | 38.1 | 39.2 | 39.6 | 40.3 | 42.0 | 44.4 | 45.1 | 45.8 | 46.4 |
| 30–39 | 374 | 41.9 | 0.2 | 37.5 | 38.5 | 39.0 | 40.0 | 41.7 | 43.8 | 44.6 | 45.0 | 46.2 |
| 40–49 | 331 | 41.3 | 0.2 | 36.0 | 37.5 | 38.5 | 39.3 | 41.4 | 43.2 | 44.0 | 44.8 | 45.9 |
| 50–59 | 397 | 40.4 | 0.2 | 35.4 | 36.5 | 37.5 | 38.4 | 40.3 | 42.2 | 43.4 | 44.1 | 45.1 |
| 60–69 | 618 | 40.4 | 0.2 | 35.0 | 36.4 | 37.2 | 38.4 | 40.5 | 42.5 | 43.3 | 44.1 | 45.1 |
| 70–79 | 425 | 39.4 | 0.1 | 34.4 | 35.5 | 36.2 | 37.3 | 39.4 | 41.2 | 42.2 | 42.9 | 43.7 |
| 80 and older | 137 | 39.4 | 0.5 | † | 35.9 | 36.2 | 37.2 | 39.7 | 41.3 | 42.1 | 43.0 | 43.7 |
| Female | | | | | | | | | | | | |
| 20 and older | 3,116 | 36.8 | 0.1 | 31.9 | 33.2 | 33.9 | 34.9 | 36.9 | 38.7 | 39.7 | 40.4 | 41.4 |
| 20–29 | 343 | 38.4 | 0.2 | 34.3 | 35.1 | 35.7 | 36.6 | 38.2 | 40.1 | 40.9 | 41.4 | 42.4 |
| 30–39 | 483 | 37.6 | 0.2 | 32.9 | 34.0 | 34.6 | 35.8 | 37.6 | 39.4 | 40.3 | 41.1 | 42.2 |
| 40–49 | 421 | 37.1 | 0.2 | 32.7 | 33.6 | 34.1 | 35.2 | 37.2 | 38.9 | 39.6 | 40.0 | 41.2 |
| 50–59 | 485 | 36.4 | 0.2 | 31.6 | 32.8 | 33.4 | 34.6 | 36.5 | 38.2 | 39.3 | 40.0 | 40.9 |
| 60–69 | 770 | 36.0 | 0.1 | 31.2 | 32.5 | 33.3 | 34.2 | 36.2 | 37.9 | 38.7 | 39.4 | 40.2 |
| 70–79 | 461 | 35.4 | 0.2 | 31.2 | 32.0 | 32.7 | 33.5 | 35.5 | 37.4 | 38.1 | 38.6 | 39.5 |
| 80 and older | 153 | 35.3 | 0.3 | 30.2 | 31.5 | 32.9 | 33.7 | 35.1 | 37.0 | 37.6 | 38.3 | 40.0 |

[†] Estimate not shown because the standard error of the 5th percentile could not be computed.

NOTE: Sample size is unweighted; estimates are weighted.

Appendix. Supporting Tables

Table I. Body measurements or indices from the National Health and Nutrition Examination Survey, by age and survey period, 1999–2000 through August 2021–August 2023

| | | | | | | | Surv | ey period | | | | | |
|------------------------------|--------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------------|-----------------------------|
| Measurement | Age | 1999– 2000 | 2001– 2002 | 2003– 2004 | 2005– 2006 | 2007– 2008 | 2009– 2010 | 2011– 2012 | 2013– 2014 | 2015– 2016 | 2017– 2018 | 2019– March 2020 | August 2021– August 2023 |
| Weight, height, and length | | | | | | | | | | | | | |
| Weight | All ages | Х | Х | Х | Χ | Χ | Х | Χ | Χ | Х | Х | X | X |
| Standing height | 2 years and older | Χ | Χ | Χ | Χ | Х | Χ | Χ | Χ | Χ | Χ | X | Х |
| Recumbent length | 0-47 months | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | X | X |
| Upper arm length | 2 months and older | X | X | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | X | X |
| Upper leg length | 8 years and older | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | X | Χ |
| Skinfold | | | | | | | | | | | | | |
| Subscapular skinfold | 2 months and older | Х | Χ | Х | Χ | Χ | Х | | | | | | |
| Triceps skinfold | 2 months and older | Х | Х | Х | Х | Х | Х | | | | | | |
| Circumference and breadth | | | | | | | | | | | | | |
| Head circumference | 0-6 months | Χ | Χ | Χ | Χ | Х | Χ | Χ | Χ | Χ | Χ | Х | Х |
| Waist circumference | 2 years and older | Χ | Χ | Χ | Х | Х | Χ | Χ | Χ | Χ | Х | X | Х |
| Sagittal abdominal diameter | 8 years and older | | | | | | | Χ | Χ | Χ | | | |
| Mid-arm circumference | 2 months and older | X | Х | Χ | Χ | Χ | Χ | Χ | Χ | Χ | Χ | X | X |
| Mid-thigh circumference | 8 years and older | X | X | X | X | | | | | | | | |
| Maximal calf circumference | 8 years and older | X | X | Χ | Χ | | | | | | | | |
| Hip (buttocks) circumference | 12 years and older | ••• | | | | | | | | | X | Х | Χ |
| Index | | | | | | | | | | | | | |
| Body mass index | 2 years and older | Χ | Х | Х | Х | Х | Χ | Χ | Χ | Х | Х | X | X |

^{...} Category not applicable.

NOTE: Data for 2017-2018 and 2019-March 2020 were also released as a combined 2017-March 2020 data file.

Table II. Historical summary of published anthropometry reference data

| Measurement or index | Survey period | | | | | | | | | | | |
|---|--------------------------|------------------------|------------------------|--------------------------|------------------------|------------------------|-------------------------|-------------|-------------------------|-------------------------|-------------------------|-----------------------------|
| | 1960–1962 ^{1–3} | 1963–1965 ⁴ | 1966–1970 ⁵ | 1971–1974 ^{6,7} | 1976–1980 ⁸ | 1988–1994 ⁹ | 1999–2002 ¹⁰ | 2003–200611 | 2007–2010 ¹² | 2011–2014 ¹³ | 2015–2018 ¹⁴ | August 2021– August 2023 |
| Weight, height, and length | | | | | | | | | | | | |
| Weight | Х | Х | Х | X | X | Х | Χ | Χ | Χ | X | Х | X |
| Standing height | Χ | X | Х | X | Χ | Х | X | X | X | X | X | X |
| Recumbent length | | | | | Х | Х | X | X | X | X | X | |
| Sitting height | Х | X | | X | Х | Х | | | | | | |
| Knee height | Х | X | | | | Х | | | | | | |
| Elbow rest height | Х | | | | | | | | | | | |
| Popliteal height | Х | Х | | | | | | | | | | |
| Thigh clearance height | X | X | ••• | | | | | | | ••• | | |
| Buttock-knee length | X | X | | | | | | | | | | |
| Buttock-popliteal length | X | X | | | ••• | | ••• | ••• | ••• | | ••• | ••• |
| Elbow-wrist length | | X | ••• | ••• | ••• | ••• | ••• | | | ••• | ••• | |
| Hand length | ••• | X | ••• | | | ••• | ••• | | | ••• | ••• | ••• |
| • | ••• | | ••• | | | | | | | | | |
| Upper arm length | ••• | Х | ••• | | | X | X | X | X | X | X | X |
| Upper leg length | ••• | ••• | • • • • | ••• | ••• | Χ | Х | X | Х | Х | Х | Х |
| Foot length | ••• | X | ••• | ••• | ••• | ••• | ••• | ••• | • • • • | ••• | ••• | |
| Crown-rump length | | ••• | ••• | ••• | Χ | ••• | | ••• | ••• | | ••• | ••• |
| Index | | | | | | | | | | | | |
| Body mass index ¹⁵ Skinfold | | ••• | ••• | ••• | Х | Х | Х | Х | X | Х | Х | Х |
| Subscapular skinfold | Х | X | Х | X | X | Х | X | X | Χ | | | |
| Triceps skinfold | Х | Х | X | X | x | Х | Х | Х | X | | | |
| Thigh skinfold | | | | | | Х | | | | | | |
| Suprailiac skinfold | | | | | | Χ | | | | | | |
| Circumference and breadth | | | | | | | | | | | | |
| Head circumference | | | | | Х | Х | Χ | Χ | Χ | Χ | Х | |
| Chest circumference | х | X | ••• | X | X | | | | | | | |
| Waist circumference | | | ••• | | | | | | | | | |
| Sagittal abdominal diameter | Х | Х | ••• | | | Х | Χ | Χ | Χ | X | Х | Χ |
| Mid-arm circumference | v | | | | | | | | | X | | |
| | Х | Х | Χ | Х | Х | X | X | X | Х | Χ | X | Х |
| Mid-thigh circumference | | | • • • • | | | Х | Х | X | | ••• | ••• | |
| Hip (buttocks) circumference | ••• | Х | ••• | ••• | ••• | Χ | ••• | ••• | ••• | ••• | ••• | Х |
| Maximal calf circumference | ••• | ••• | ••• | ••• | ••• | • • • • | Х | Х | ••• | ••• | ••• | ••• |
| Chest breadth | | Χ | • • • • | | | | ••• | | | ••• | • • • • | ••• |
| Hand breadth | | X | • • • • | | | | ••• | | | ••• | • • • • | ••• |
| Wrist breadth | | | | | | Х | | | | ••• | | |
| Biacromial breadth | X | | | | | Х | | | | | | |
| Bi-iliac breadth | | | | | | Х | | | | | | |
| Bitrochanteric breadth | | | | X | X | | | | | | | |
| Elbow breadth | | | | Х | Х | Х | | | | | | |
| Elbow-elbow breadth | Χ | Χ | | | | | | | | | | |
| Seat breadth | X | X | | | | | | | | | | |
| Foot breadth | | Х | | | | | | | | | | |

See footnotes at end of table.

.. Category not applicable.

1Stoudt HW, Damon A, McFarland R, Roberts J. Weight, height, and selected body dimensions of adults, United States, 1960–1962. Vital Health Stat 11. 1965 Jun;(8):1–44. PMID: 14338862. Available from: https://www.cdc.gov/nchs/data/series/sr 11/sr11 008.pdf.

²Stoudt HW, Damon A, McFarland RA, Roberts J. Skinfolds, body girths, biacromial diameter, and selected anthropometric indices of adults, United States, 1960–1962. Vital Health Stat 11. 1970 Feb;(35):1–63. 1970. PMID: 5309272. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_035acc.pdf.

³Roberts J. Weight by height and age of adults, United States, 1960–1962. Vital Health Stat 11. 1966 May;(14):1–38. PMID: 5296365. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_014acc.pdf.

⁴Malina RM, Hamill PVV, Lemeshow S. Selected body measurements of children 6–11 years, United States. Vital Health Stat 11. 1973 Jan;(123):1–48. PMID: 4540661. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_123acc.pdf.

⁵Hamill PVV, Johnston FE, Lemeshow S. Height and weight of youths 12–17 years, United States. Vital Health Stat 11. 1973 Jan;(124):1–81. PMID: 4539878s. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_124.pdf.

⁶Abraham S, Johnson CL, Najjar MF. Weight and height of adults 18–74 years of age: United States, 1971–74. Vital Health Stat 11. 1979 May;(211):1–49. PMID: 473616. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_211.pdf.

⁷Johnson CL, Fulwood R, Abraham S, Bryner JD. Basic data on anthropometric measurements and angular measurements of the hip and knee joints for selected age groups 1–74 years of age: United States, 1971–1975. Vital Health Stat 11. 1981 Apr;(219):1–68. PMID: 15806754. Available from: https://www.cdc.gov/nchs/data/series/sr 11/sr11 219.pdf.

⁸Najjar MF, Rowland M. Anthropometric reference data and prevalence of overweight: United States, 1976–80. Vital Health Stat 11. 1987 Oct;(238):1–73.PMID: 3424692. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_238.pdf.

9McDowell MA, Fryar CD, Ogden CL. Anthropometric reference data for children and adults: United States, 1988–1994. Vital Health Stat 11. 2009 Apr;(249):1–68. PMID: 19642512. Available from: https://www.cdc.gov/nchs/data/series/sr 11/sr11 249.pdf.

¹⁰McDowell MA, Fryar CD, Hirsch R, Ogden CL. Anthropometric reference data for children and adults: U.S. population, 1999–2002. Adv data. 2005 July 7;(361):1–5. PMID: 16018338. Available from: https://www.cdc.gov/nchs/data/ad/ad361.pdf.

¹¹McDowell MA, Fryar CD, Ogden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2003–2006. Natl Health Stat Report. 2008 Oct 22;(10):1–48. PMID: 25585443. Available from: https://www.cdc.gov/nchs/data/nhsr/nhsr010.pdf.

¹²Fryar CD, Gu Q, Ogden CL. Anthropometric reference data for children and adults: United States, 2007–2010. Vital Health Stat 11. 2012 Oct;(252):1–48. PMID: 25204692. Available from: https://www.cdc.gov/nchs/data/series/sr_11/sr11_252.pdf.

¹³Fryar CD, Gu Q, Ögden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2011–2014. Vital Health Stat 3 Anal Stud. 2016 Aug;(39):1–46. PMID: 28437242. Available from: https://www.cdc.gov/nchs/data/series/sr 03/sr03 039.pdf.

¹⁴Fryar CD, Carroll MD, Gu Q, Afful J, Ogden CL. Anthropometric reference data for children and adults: United States, 2015–2018.Vital Health Stat 3. 2021 Jan;(46):1–44. PMID: 33541517. Available from: https://www.cdc.gov/nchs/data/series/sr 03/sr03-046-508.pdf.

¹⁵Body mass index (BMI) is calculated as: BMI equals weight in kilograms divided by height in meters squared.

NOTE: Terminology for a particular measure may have changed across survey years.

Vital and Health Statistics Series Descriptions

Active Series

Series 1. Programs and Collection Procedures

Reports describe the programs and data systems of the National Center for Health Statistics, and the data collection and survey methods used. Series 1 reports also include definitions, survey design, estimation, and other material necessary for understanding and analyzing the data.

Series 2. Data Evaluation and Methods Research

Reports present new statistical methodology including experimental tests of new survey methods, studies of vital and health statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. Reports also include comparison of U.S. methodology with those of other countries.

Series 3. Analytical and Epidemiological Studies

Reports present data analyses, epidemiological studies, and descriptive statistics based on national surveys and data systems. As of 2015, Series 3 includes reports that would have previously been published in Series 5, 10–15, and 20–23.

Discontinued Series

Series 4. Documents and Committee Reports

Reports contain findings of major committees concerned with vital and health statistics and documents. The last Series 4 report was published in 2002; these are now included in Series 2 or another appropriate series.

Series 5. International Vital and Health Statistics Reports

Reports present analytical and descriptive comparisons of U.S. vital and health statistics with those of other countries. The last Series 5 report was published in 2003; these are now included in Series 3 or another appropriate series.

Series 6. Cognition and Survey Measurement

Reports use methods of cognitive science to design, evaluate, and test survey instruments. The last Series 6 report was published in 1999; these are now included in Series 2.

Series 10. Data From the National Health Interview Survey

Reports present statistics on illness; accidental injuries; disability; use of hospital, medical, dental, and other services; and other health-related topics. As of 2015, these are included in Series 3.

Series 11. Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey

Reports present 1) estimates of the medically defined prevalence of specific diseases in the United States and the distribution of the population with respect to physical, physiological, and psychological characteristics and 2) analysis of relationships among the various measurements. As of 2015, these are included in Series 3.

Series 12. Data From the Institutionalized Population Surveys

The last Series 12 report was published in 1974; these reports were included in Series 13, and as of 2015 are in Series 3.

Series 13. Data From the National Health Care Survey

Reports present statistics on health resources and use of health care resources based on data collected from health care providers and provider records. As of 2015, these reports are included in Series 3.

Series 14. Data on Health Resources: Manpower and Facilities

The last Series 14 report was published in 1989; these reports were included in Series 13, and are now included in Series 3.

Series 15. Data From Special Surveys

Reports contain statistics on health and health-related topics from surveys that are not a part of the continuing data systems of the National Center for Health Statistics. The last Series 15 report was published in 2002; these reports are now included in Series 3.

Series 16. Compilations of Advance Data From Vital and Health Statistics

The last Series 16 report was published in 1996. All reports are available online; compilations are no longer needed.

Series 20. Data on Mortality

Reports include analyses by cause of death and demographic variables, and geographic and trend analyses. The last Series 20 report was published in 2007; these reports are now included in Series 3.

Series 21. Data on Natality, Marriage, and Divorce

Reports include analyses by health and demographic variables, and geographic and trend analyses. The last Series 21 report was published in 2006; these reports are now included in Series 3.

Series 22. Data From the National Mortality and Natality Surveys

The last Series 22 report was published in 1973. Reports from sample surveys of vital records were included in Series 20 or 21, and are now included in Series 3.

Series 23. Data From the National Survey of Family Growth

Reports contain statistics on factors that affect birth rates, factors affecting the formation and dissolution of families, and behavior related to the risk of HIV and other sexually transmitted diseases. The last Series 23 report was published in 2011; these reports are now included in Series 3.

Series 24. Compilations of Data on Natality, Mortality, Marriage, and Divorce

The last Series 24 report was published in 1996. All reports are available online; compilations are no longer needed.

For answers to questions about this report or for a list of reports published in these series, contact:

Information Dissemination Staff
National Center for Health Statistics
Centers for Disease Control and Prevention
3311 Toledo Road, Room 4551, MS P08
Hyattsville, MD 20782

Tel: 1-800-CDC-INFO (1-800-232-4636)

TTY: 1-888-232-6348

Internet: https://www.cdc.gov/nchs

Online request form: https://www.cdc.gov/info

For e-mail updates on NCHS publication releases, subscribe online at: https://www.cdc.gov/nchs/updates/.

U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES

Centers for Disease Control and Prevention National Center for Health Statistics 3311 Toledo Road, Room 4551, MS P08 Hyattsville, MD 20782–2064

OFFICIAL BUSINESS PENALTY FOR PRIVATE USE, \$300 FIRST CLASS MAIL POSTAGE & FEES PAID CDC/NCHS PERMIT NO. G-284



For more NCHS Series Reports, visit: https://www.cdc.gov/nchs/products/series.htm