

Menveo New Ready-to-Use Single Vial Presentation

Menveo is a MenACWY vaccine product approved to prevent invasive meningococcal disease caused by *Neisseria meningitidis* serogroups A, C, Y, and W.

Menveo is now available in two presentations:





- **Two-vial** (requires reconstitution)
Recommended for ages **2 months and older***
- **Single-vial** (ready-to-use)
Recommended for ages **10 years and older**
NOT for use in individuals before their 10th birthday

• To extend existing stocks of MenACWY vaccine products, CDC recommends that health care providers administer specific presentations of age-appropriate vaccines.

Ages 2 months through 23 months	Ages 2 years through 9 years	Ages 10 years and older
Menveo Two-vial (requires reconstitution)	Menveo Two-vial (requires reconstitution) or MenQuadfi (single-dose vial)	Menveo Single-vial (ready-to-use) or MenQuadfi (single-dose vial)

- Menactra is no longer distributed in the United States; available doses will expire in October 2023.
- Develop a plan to add MenQuadfi to your inventory if vaccinating children ages 2 years through 9 years.

Important Differences Between Menveo Presentations

Age Indications	Two-vial (vaccine + vaccine)	Single-vial (ready-to-use)
		
	Ages 2 months and older	Ages 10 years and older
Presentation	Supplied as two vials Requires reconstitution <ul style="list-style-type: none"> • Orange cap vial contains MenA lyophilized conjugate • Gray cap vial contains MenCYW-135 liquid conjugate 	Supplied as one ready-to-use vial Reconstitution not required <ul style="list-style-type: none"> • Pink cap vial
Packaging	Each carton contains 10 vials for 5 doses  Five vials of MenA lyophilized conjugate component AND Five vials of MenCYW135 liquid conjugate component	Each carton contains 10 vials for 10 doses  Ten single vials of ready-to-use Menveo
Storage	Store refrigerated between 36°F and 46°F (2°C and 8°C) Protect from light Do not freeze	
Preparation	<ul style="list-style-type: none"> • Reconstitution required • Only use vaccine vials supplied by manufacturer • Mix vaccines vials before administering 	<ul style="list-style-type: none"> • Ready-to-use • Reconstitution not required
	See Menveo package insert for preparation guidance	
Dosage/Route	<ul style="list-style-type: none"> • 0.5 mL/intramuscular (IM) injection • Use immediately after reconstituted 	<ul style="list-style-type: none"> • 0.5 mL/intramuscular (IM) injection



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

* Off-label recommendations based on Advisory Committee on Immunization Practices- Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020. MMWR Recomm Rep 2020;69 (No. RR-9):1-41. DOI: <http://dx.doi.org/10.15585/mmwr.rr6909a1external icon>